

Dear Reviewer,

Thank you for your feedback and input. Overall, the boundary conditions imposed by proxy information collected over decades of work basically amount to one undeniable fact: we are unable to fully reconcile and align glacial-interglacial CO₂ variability with changes in the biosphere and gas exchange with the ocean. We therefore need a new hypothesis.

On the most basic level, the constraints imposed by radiocarbon clearly indicate we are missing one fundamental parameter: a source of carbon depleted or devoid of radiocarbon (Broecker and Clark 2010; Zhao et al., 2018). When asking ourselves what this source may be, the transition during times of highest CO₂ increase between glacial and interglacial clearly reveals a negative stable carbon isotopic shift of atmospheric CO₂ (Smith et al., 1999; Schmitt et al., 2012), which is a strong indicator of respired organic carbon acting as source to the atmosphere (Bauska et al., 2016).

With this, exogenous kerogen becomes an obvious candidate as the missing player in the global carbon cycle. It fulfills the basic requirements: it is abundant, it is isotopically light, and it is radiocarbon-dead. For an additional host of reasons, it appears plausible, as elaborated in the manuscript, with kerogen reburial efficiency higher during glacial episodes in Earth's history as a straightforward supporting observation (see references in Blattmann et al., 2018).

As pointed out by yourself (the reviewer), viewing the glacial and interglacial episodes as individual time intervals, one may draw the conclusion that there is biospheric uptake of isotopically light carbon on land (Shackleton 1977; Shackleton et al., 1983), which I agree, is most likely true. However, this view regards the two climate states as steady state conditions and does not consider transitional dynamics. Also, stable carbon isotopic changes are trickier to interpret as carbon isotopes fractionate as they rotate between different carbon pools changing in size; in contrast, radiocarbon isotope composition of atmospheric CO₂, which is fractionation corrected, is more straightforward in its interpretation. Ultimately, biospheric activity and atmosphere-ocean exchange cannot account for the radiocarbon budget and its evolution from the Last Glacial Maximum to the present. However, yes, the kerogen cycle is one "cogwheel operating under manifold feedbacks in the greater Earth system" (original manuscript) and clearly this point needs to be expanded on to present a mediated view.

How important is this cogwheel? And you (the reviewer) rightfully points out that a back-of-the-envelope calculation is lacking. The "Glacial Burial Hypothesis" (Zeng, 2003), which contains parallels to the hypothesis presented here, takes care of this in a detailed manner. Zeng (2003) proposes the oxidation of ice sheet-covered soil organic carbon during glacial retreat. Overall, the modeling results of Zeng (2003, 2007), if soil organic carbon is reinterpreted as kerogen, lend further support to the idea that kerogen oxidation drives a substantial part of atmospheric CO₂ increase in the wake of glacial episodes. In quantitative terms, Zeng (2003) attributes the oxidation of 547 PgC soil organic carbon to a 60 PgC increase in atmospheric CO₂ (increase of 30 ppm). I view these estimates as conservative, because unlike Ning Zeng's assumption of soil organic carbon oxidation, the supply of kerogen shed and subject to temporary storage on land over millennia of glaciation is much vaster!

Overall, the dynamism of the exogenous kerogen cycle over the Ice Ages may seem further strengthened by increased bedrock exhumation (Herman et al., 2013; Herman et al., 2015), thereby leading to greater detrital fluxes of kerogen reburied into ocean sediments and also enhanced supply of ground kerogen exposed to the elements in the wake of glacial episodes.

I thank the reviewer for his/her time and effort and look forward to future discussions and strengthening the manuscript based on this input and adding the references cited herein.

Sincerely,

Thomas Blattmann

01.10.2019 Yokosuka

References

- Bauska, T. K., Baggenstos, D., Brook, E. J., Mix, A. C., Marcott, S. A., Petrenko, V. V., Schaefer, H., Severinghaus, J. P., and Lee, J. E., 2016, Carbon isotopes characterize rapid changes in atmospheric carbon dioxide during the last deglaciation: *Proceedings of the National Academy of Sciences*, v. 113, no. 13, p. 3465-3470.
- Blattmann, T. M., Letsch, D., and Eglinton, T. I., 2018, On the geological and scientific legacy of petrogenic organic carbon: *American Journal of Science*, v. 318, no. 8, p. 861-881.
- Broecker, W., and Clark, E., 2010, Search for a glacial-age ¹⁴C-depleted ocean reservoir: *Geophysical Research Letters*, v. 37, no. 13.
- Herman, F., Seward, D., Valla, P. G., Carter, A., Kohn, B., Willett, S. D., and Ehlers, T. A., 2013, Worldwide acceleration of mountain erosion under a cooling climate: *Nature*, v. 504, p. 423.
- Herman, F., Beyssac, O., Brughelli, M., Lane, S. N., Leprince, S., Adatte, T., Lin, J. Y. Y., Avouac, J.-P., and Cox, S. C., 2015, Erosion by an Alpine glacier: *Science*, v. 350, no. 6257, p. 193.
- Schmitt, J., Schneider, R., Elsig, J., Leuenberger, D., Lourantou, A., Chappellaz, J., Köhler, P., Joos, F., Stocker, T. F., Leuenberger, M., and Fischer, H., 2012, Carbon isotope constraints on the deglacial CO₂ rise from ice cores: *Science*, v. 336, no. 6082, p. 711-714.
- Shackleton, N. J., 1977, Carbon-13 in Uvigerina: Tropical rainforest history and the equatorial Pacific carbonate dissolution cycles, *in* Andersen, N. R., and Malahoff, A., eds., *The Fate of Fossil Fuel CO₂ in the Oceans*, Plenum Press, p. 401-427.
- Shackleton, N. J., Hall, M. A., Line, J., and Shuxi, C., 1983, Carbon isotope data in core V19-30 confirm reduced carbon dioxide concentration in the ice age atmosphere: *Nature*, v. 306, no. 5941, p. 319-322.
- Smith, H. J., Fischer, H., Wahlen, M., Mastroianni, D., and Deck, B., 1999, Dual modes of the carbon cycle since the Last Glacial Maximum: *Nature*, v. 400, no. 6741, p. 248-250.
- Zhao, N., Marchal, O., Keigwin, L., Amrhein, D., and Gebbie, G., 2018, A synthesis of deglacial deep-sea radiocarbon records and their (in)consistency with modern ocean ventilation: *Paleoceanography and Paleoclimatology*, v. 33, no. 2, p. 128-151.
- Zeng, N., 2003, Glacial-interglacial atmospheric CO₂ change — The glacial burial hypothesis: *Advances in Atmospheric Sciences*, v. 20, no. 5, p. 677-693.
- Zeng, N., 2007, Quasi-100 ky glacial-interglacial cycles triggered by subglacial burial carbon release: *Climate of the Past*, v. 3, no. 1, p. 135-153.