Biogeosciences Discuss., https://doi.org/10.5194/bg-2019-289-AC1, 2019 © Author(s) 2019. This work is distributed under the Creative Commons Attribution 4.0 License.



Interactive comment on "On modeling the Southern Ocean Phytoplankton Functional Types" by Svetlana N. Losa et al.

Svetlana N. Losa et al.

svetlana.losa@awi.de

Received and published: 29 August 2019

Dear readers, we would like to draw your attention and remind that, when submitted, the discussed manuscript was supplemented with three videos on the model simulated chlorophyll "a" distribution for diatoms, haptophytes and prokaryotes in the Souther Ocean for the time period from August 2002 to April 2012. These videos also depict comparisons with available in situ observations and clearly illustrate phenology/bloom development for these three phytoplankton types over the aforementioned time period.

Video supplement

Simulated distribution of diatom chlorophyll concentration in the Southern Ocean https://doi.org/10.5446/42871

C1

Simulated distribution of haptophytes chlorophyll concentration in the Southern Ocean https://doi.org/10.5446/42873

Simulated distribution of prokaryotes chlorophyll concentration in the Southern Ocean https://doi.org/10.5446/42872

Interactive comment on Biogeosciences Discuss., https://doi.org/10.5194/bg-2019-289, 2019.