

Interactive comment on “Effect of legume intercropping on N₂O emission and CH₄ uptake during maize production in the Ethiopian Rift valley” by Shimelis G. Raji and Peter Dörsch

Shimelis G. Raji and Peter Dörsch

peter.doersch@umb.no

Received and published: 26 September 2019

General comments: This paper reports results from an experiment conducted in Ethiopia measuring yields and GHG fluxes from maize cultivated as monocrop and intercropped with 2 legumes. There is an urgent need to increase the empirical base quantifying GHG fluxes from agricultural systems in Sub-Saharan Africa and therefore this study could be a valuable contribution to the literature. Understanding the interactions between cereal and legume crops and quantifying C footprints are also commendable scientific goals, and requirements to design future climate-smart farming. However, this study seems to have a number of experimental shortcomings that

C1

require at least clarification to be able to assess its suitability for publication in Biogeosciences.

Response: We thank the reviewer for constructive comments and criticism. The reviewer's main points of critique can be summarized as i) lack of ancillary data (e.g. soil mineral N) and N-fluxes (e.g. quantification of BNF) and ii) too much speculation about underlying processes. We agree with the reviewer that our study has experimental shortcomings, but we believe that our research has some salient points worth communicating to a broader audience: i) Intercropping and mulching legumes to maize under Rift Valley conditions did not cause major N₂O emissions, nor inhibit CH₄ uptake during a dry and a wet year ii) Legume intercropping therefore appears as a viable option for climate-smart intensification which is urgently needed in the region iii) Even though being highly insecure, numbers of leguminous N input, N₂O-EFs, etc. presented in our paper can be used as first estimates in the absence of better data We understand the reviewer's frustration about the lack of ancillary data (soil moisture in 2015, mineral N content, below ground legume biomass, etc.) but as in any empirical study, there are limitations to the number of variables which can be measured, particularly so when relying on local research infrastructure.

1. The introduction doesn't follow a logical flow. It includes interesting hypotheses, although the authors either do not properly attempt to answer the hypotheses or do it insufficiently. Example: “Legumes affect emissions by providing organic N or by modulating the competition between roots and microbes for soil N”. The authors could have added how these processes are ‘modulated’, and use the appropriate methods to quantify species competition and microbial processes.

Response: Studying legume-rhizobia interactions is not trivial (see for example Raji et al., 2019). Species competition and microbial processes were not the primary focus of our study. Instead, we were interested in the overall effect of forage legume intercropping and its management on N₂O and CH₄ fluxes. We rephrased the sentence to “...or by modulating the competition between plants and microbes for soil N, for example by

C2

acting as an additional N sink prior to nodulation”.

2. The methods are poorly described to assess the value of the experimental data. I indicated shortcomings in Specific comments below.

Response: We address these shortcomings in response to the specific comments below.

3. The discussion is mostly a compilation of literature conducted elsewhere reporting GHG fluxes from intercropping including legumes. I would expect a reflection of the results against the relevant literature.

Response: Comparing our flux estimates with those found in other GHG studies in Sub-Saharan Africa is an important first step to scrutinize and contextualize our measurements. The remainder of the discussion tries to interpret treatment effects by linking fluxes to measured variables (weather, legume biomass, etc.), necessarily drawing on the general literature. We are not entirely sure what the reviewer means by “relevant literature”. Even though intercropping with forage legumes is a common practice in the Ethiopian Rift Valley, there are no published studies on how these practices affect N₂O and CH₄ fluxes. We therefore compared our N₂O fluxes and emission factors to those reported for humid tropical maize production systems with intercropping, which – at least geographically – come closest to the system studied by us. We would be grateful to learn about relevant literature we have missed out.

A modest aim for this paper could have been simply documenting the GHG flux measurements and explaining the patterns observed, using all the data collected and conducting a sensitivity analysis for the fluxes that have been roughly estimated, such as the contribution of the legumes to N inputs, the emission factors and the emissions intensity.

Response: Estimating emission factors necessitates estimating N inputs, which is particularly challenging in experiments involving N input from BNF, green manuring or

C3

crop residue retention. Our estimates of N input by BNF are based on assumptions of legume shoot-root ratios and residue decomposition rates, which we anchored in the literature, as outlined in chapter 2.4. We believe that this approach does not lend itself to “sensitivity analysis for the fluxes” as we do not use statistical models to explain variations in flux. We decided to abstain from such models because of the inherent insecurity of underlying variables such as legume N input. Instead, we resorted to simple linear regression using measured aboveground legume N yield and N₂O emission intensity (Fig. 3). We want to emphasize that all variables and their derivations (cumulative flux, emission intensity and factors) were estimated or calculated on a per plot level before averaging them, giving at least some measure of dispersion (e.g. Figures 2, 5 and Table 2).

Because there are very few experiments measuring GHG fluxes in Africa, I would suggest a thorough revision addressing the shortcomings, to re-consider this manuscript for publication.

Response: A thoroughly revised manuscript will be provided addressing all points raised by the reviewers.

These are the most important issues to be addressed:

Specific comments Introduction

L39: The use of inorganic fertilisers does not necessarily reduce the soil methane sink. Please explain.

Response: No, it does not. Our introduction tries to detail the conditions potentially leading to reduced CH₄ uptake by citing a meta-study that found an overall higher propensity for reduction in CH₄ uptake at mineral N fertilization rates > 100kg N ha⁻¹ y⁻¹ (Aronson and Helliker, 2010). We further outline possible mechanisms regulating CH₄ uptake in fields with intercropping in Lines 81-91 of the original text. At no point, we claim that inorganic fertilizers invariably reduce the soil’s sink strength for methane.

C4

The sentence in line 39 now reads: “Abundant ammonium (NH₄⁺) may also reduce the soil CH₄ sink by competing with CH₄ for the active binding site of methane monooxygenase, the key enzyme of CH₄ oxidation (Bédard and Knowles, 1989)”

L40 remove ‘by contrast’. It doesn’t follow naturally from the previous sentence

Removed

L41 the concept of CSA – coined by FAO – doesn’t talk about profits. Please revisit original source

Response: The reviewer is right. We remove ‘profits’.

L43: I don’t think the understanding of GHG fluxes in SSA is limited. There is a scarcity of quantified GHG fluxes in SSA, and limited experimentation on which CSA practices would be suitable for the SSA context. Please rephrase.

Response: We agree with the reviewer that sources and sinks of GHGs in SSA are well understood, in principal, and rephrase the sentence to “However, greenhouse gas emission measurements in SSA crop production systems are scarce and proof-of-concept for the mitigation potential of specific CSA practices is missing (Kim et al., 2016, Hickman et al., 2014b).”

L49: Crop production can be a major source of N₂O emissions when fertilisers are used. This is not often the cause in East African agriculture. There are empirical studies that show that

Response: Food production in SSA has to double by 2050 to feed a growing population. This requires intensification of crop production, be it by increasing nitrogen fertilization or by other approaches, such as legume intercropping. We therefore believe that studying and documenting intensification effects on N₂O emissions are important in the wider context of GHG mitigation in the global agrifood system. We agree that N₂O emissions in rainfed SSA crop production appear small per date, but given the enormous productivity increase needed, also crop production in SSA may become a

C5

major source of N₂O. In the revised version, we state explicitly “Emission rates of N₂O reported for SSA crop production so far are low (Kim et al., 2016) owing to low fertilization rates, but may increase with increasing intensification.”

L53: strange reference to ‘upland soils’ here. Please explain why the focus is suddenly shifted towards upland soils

Response: The term “upland” was removed as the statement refers to factors that control N₂O production in soils in general.

L58: soil management practices are not the only controls of the factors affecting soil N₂O fluxes. Soil type and climate are major determinants, which don’t depend on management

Response: The reviewer is right! We rephrase the statement and add soil type and climate as important factors for N₂O emissions, including two new references. The sentence added reads: “Other important factors are soil type (Davidson et al., 2000) and temperature (Schaufler et al., 2010).”

L59: The position of the two first references in this sentence is not logical. Please revise.

Revised

L68: diversification, rotation and intercropping do not always enhance productivity. Please rephrase

Response: The reviewer is right. We rephrased the sentence to: “Crop diversification by combining legumes with cereals, both in rotation and intercropping, enhances overall productivity and resource use efficiency, if managed properly (Ehrmann and Ritz, 2014)”

L71 please add reference that shows that legume improve N uptake of the cereal crop in the Rift Valley (this is a large area across countries!). There is evidence in favor and

C6

against this.

Response: A reference was added (Sime and Aune, 2018), describing the general benefits of legume inclusion in farming systems in the region.

L86 rates of 100 kg N per hectare are very uncommon in Africa. Please consult the literature on fertilizer use for the continent.

Response: This statement refers to the general relationship between N rates and methane uptake as elucidated by the meta-study of Aronson and Helliker (2010) and not to common N fertilization rates in SSA.

L89: increasing. Remove or replace 'accordingly', doesn't seem to fit the meaning of the sentence.

Done

L93: add 'the' to 'the' release. Please explain how root exudates release 'extra N'

Done

L95-96: are these the hypotheses this study wanted to test?

Response: No. This sentence refers to the background of how legumes may cause extra N₂O emissions. The hypotheses of the study are given in L109 -115 in the original text.

L110-112: these hypotheses don't have any mechanistic underpinning, and are therefore weak. Time measured in weeks is unlikely to be a fixed effect, since the effects of management such as sowing date, choice of species and cultivar on yields and GHG fluxes will depend on soil and weather.

Response: The competition for nutrients after under-sowing the intercrop, as well as the benefit of the main crop from N transfer depend in deed on a variety of factors, particularly those which control the initial growth of maize and hence its shading effect

C7

on the legume. Our study is a good illustration for this: equal sowing dates produced vastly different legume aboveground biomasses in a dry and a wet year (Table 1). Yet, among all factors, the sowing date of the legume (relative to the main crop) is the one, which potentially could be controlled by the farmer, preferably in response to prevailing weather conditions. In response to the reviewer's righteous remark, we modify the respective conclusion in the discussion section to "Our study therefore points to optimizing the sowing date in response to expected emergence and growth of maize as a promising option to control growth of the intercrop and hence to deal with the risk of increased N₂O emissions associated with high legume biomass".

L115: because there are so few experiments measuring GHG fluxes in Africa, and more modest aim for this paper could have been simply documenting these measurements and explaining the patterns observed.

Response: Our study served two aims, documenting fluxes and evaluating intercropping strategies with respect to GHG mitigation. We agree that a merely descriptive study of fluxes in different treatments would have been the least risky approach, but mitigation needs causation if it is to be widely adopted. Therefore, we chose to link seasonal emissions to stipulated legume N input and climatic variability, which we believe are the key drivers for N₂O emissions in sub-Saharan intercropping systems. Methods

L121-126 please report soil type using a known classification, e.g. WSD. And please add measure of dispersion to the reported soil properties, and weather variables.

Response: We now include the soil type and SD for the bulk density. For analysis of soil texture and chemical composition, we used composite soil samples and hence cannot give measures of dispersion.

L128: Please explain the 6 treatments clearly here. No clear which are the treatment is Table1, and how they were imposed. Treatments seem to be listed in Table2, although there is no consistency in labels used in Tables and Figures.

C8

Response: A treatment list is now included in the Materials and Method section; label inconsistencies in tables and figures are corrected.

L130: only one cultivar? Wouldn't the researchers have expected cultivar effects on the treatments?

Response: Farmers' preference was considered in choosing the maize cultivar for the trial. This cultivar is widely used and we focused on legume species and intercropping times rather than maize cultivar as a factor.

L131: only one sowing date each year? I understood from the objective and hypothesis that the authors wanted to test the effect of sowing date (L110) on GHG fluxes.

Response: Our objective was to test the effect of legume species and sowing date of the legumes relative to maize in combination with interannual weather variation, and not the effect of the sowing date of maize itself.

L133: fertiliser rates per hectare? I am surprised to read that N fertiliser was applied to the intercropping treatment. Was there a scientific basis to half the rate? If yes, please add reference to previous experimental work.

Response: Mineral N fertilization followed national recommendations, which are low. Annual legume intercrops are used, among others, to bring additional nitrogen into the soil, both during growth and after harvest. As outlined in line 134 ff., the rate of annual mineral N fertilization was halved in the second year there where legume mulch was applied, to test whether biologically fixed N could replace mineral N, which in itself would be a climate-smart approach. Cutting down on mineral fertilization is a common goal and practice when using catch or cover crops as green manure.

L136 I would have expected an effect of plant density. These were fixed.

Response: The numbers given for legume density are the planting densities, which did not result in "fixed" densities during the growing season. Much to the contrary, in terms of legume aboveground biomass, there was a huge variability across the two years.

C9

Aboveground dry matter varied from 186 to 2221 kg ha⁻¹ for lablab and 65 to 1516 kg ha⁻¹ for crotalaria across the two years. We used this variation to explore the effect of legume biomass on N₂O emissions, which indeed showed a significant effect in the dry year 2015 (Figure 3).

L141: why half removed? did you measure this variable amount of mulching applied to the plots? This is not really a welcome variation to the treatments, and could have affected the data analysis and assumptions on treatment effects.

Response: The idea of removing 50% of the biomass and mulching the rest was motivated by livestock feed shortage in the mixed farming systems of the region. Providing feed through intercropping provides an added "climate smart" value by alleviating the pressure on crop residues otherwise used as feed, thus increasing residue retention and building/stabilizing soil carbon. It is true that different mulching rates introduced additional variation. However, applying equal mulching rates to all plots would have negated plot-specific differences in soil fertility and hence belowground input. We therefore decided to scale the rate of mulch applied according to the plot-wise legume yield, as would be done by practitioners in real fields. In this way, we created a wide range of legume biomasses and likely also of N inputs, which allowed us to explore the effect of legume growth on N₂O emissions (Fig. 3).

L151: why didn't the measurements of GHG fluxes start before planting to capture background GHG fluxes?

Response: The flux study was restricted to two growing seasons due to logistic reasons. Two control treatments with maize monocrops were included, one with recommended mineral fertilization and one without. Thus, background fluxes are captured. We agree that flux measurements outside the cropping season would be desirable.

L152: what was the frequency of sampling? Weekly? There is evidence that less than weekly sampling doesn't capture the variation of GHG fluxes in a crop's cycle. See Barton et al. 2015 Scientific Reports volume 5, Article number: 15912 (2015)

C10

Response: Flux sampling was conducted weekly as indicated in line 151 of the original manuscript

L159: Helium filled? Yes. Corrected.

L185: these treatments were not introduced before.

Response: The treatments M+Cr3w and M+Lb3w are now introduced at the beginning of the Materials and Method section.

L187 Was bulk density measured? If yes, how?

Response: A description of how it was measured is now added.

L195-L198: Not having assessed belowground biomass and the amount of N fixed by the legumes is an important shortcoming of this study. Especially because the authors pose the hypothesis in the introduction (L95-96) that “Legumes affect emissions by providing organic N or by modulating the competition between roots and microbes for soil N”. Without having quantified belowground N and N₂ fixation, the results are less useful as a contribution to test this hypothesis.

Response: The question of whether and how biologically fixed N affects N₂O emissions is a long-standing issue. Our study is a modest attempt to address this issue for sub-Saharan conditions. It was however not designed to capture the exact mechanisms of competition between crops and microbes, nor did we hypothesize that it would. Instead, our working hypothesis was that legumes inter-cropped early in the season would increase N₂O emissions if fertilized at the same time (L. 113). The reviewer is right that determining the amount and N content of belowground biomass would strengthen our approach, but given the number of field plots and the clayey soil (which makes it difficult to extract roots), the effort to do so would have been exorbitant. We therefore used aboveground biomass and its N content as a proxy for “potential” legume N input by scaling up literature based shoot-root ratios for lablab and crotalaria and estimating N release factors from literature.

C11

L199: until here, it wasn't indicated that there were different sowing times for maize and legumes. Treatments must be clearly explained at the beginning.

Response: Additional explanations about the treatments have been added to the Materials and Methods section

L202-204: this is another shortcoming, having assumed the ‘release’ of 50% and 30% of the N during the growing season doesn't help with hypothesis testing. The authors could have followed at least inorganic N in the soil.

Response: We agree that mineral N contents would have supplemented our dataset in a meaningful way, but frozen storage of extracts prior to shipment out of the country was not possible due to frequent power cuts. As to the estimated release factors for legume N in the two years, we give detailed rationale for the underlying assumption (L.201 – 212, original version).

L213: this emission factor is not meaningful given all the assumptions used to estimate N input.

Response: We agree in principal, but seasonal emission factors for N₂O have been used in the literature previously and may be considered useful for comparing different crop management strategies in regions with scarce flux data (see f. ex. Kim et al., 2016)

L221 Was grain moisture content measured?

Response: Yes, we used a digital grain moisture meter. Results

L236-237: to be able to measure peaks, N₂O fluxes must be measured continuously after fertiliser application. There is typically a peak 6-48 hours after application. The dataset unfortunately doesn't show baseline emissions that happened before the treatments were imposed.

Response: The term “peaks” has been removed

C12

L 280-295: I find this section on EFs speculative because there are large uncertainties in the estimation of N input as described in the methods section.

Response: See answer to L. 213 above.

L318: this should be explained in the methods section with all assumptions and reported as absolute emissions not GWP. This section is not clear, and need to consistently explain Fig 2 and 5. Fig 5 doesn't include letters showing the contrasts.

Response: We thank the reviewer for drawing our attention to missing indicators in Figure 5. GWP is replaced with "total non-CO2 GHG emissions" as suggested. Discussion

L330-340: this belongs more to results than to discussion.

Response: See our response to your general comment #3

L349-354: because the researchers didn't measure N2 fixation, this sentence is speculative. Also attributing the lack of relationship between N input and legume N yield and N2O fluxes to the variability of fluxes is speculative, since the estimation of the N input and yield are very uncertain and based on strong assumptions.

Response: Therefore, we talk about "potential leguminous N input" and not actual N input. We agree that our estimates of N input are insecure, but our analysis does not do more than examining the relationship of cumulative N2O emissions and "potential leguminous N input" on a plot for plot basis, before problematizing this approach in the discussion following L. 349.

L375-378: the data shown in Fig 2 doesn't show that intercropping legumes increases emissions 'risk' further than cultivating fertilized maize. If that were the case, there would be a consistent effect across years, and all legumes would increase emissions

Response: We believe that the sentence "Our data suggest that excessive accumulation of leguminous biomass in SSA maize cropping enhances the risk for elevated

C13

N2O emissions" summarizes our findings in an appropriate way based on the analysis shown in Figure 3. In the discussion following L. 275, we are explicit about other factors such as rainfall early in the season potentially overriding this relationship.

L381: unfortunately the experimental data of the one experiment in Ethiopia presented here is insufficient to claim that N2O fluxes in the sub-sequent year are negligible under SSA conditions. It is unfortunate that the researchers didn't follow the dynamics of inorganic N in the soil or plant N uptake when they sampled GHG fluxes.

Response: We agree. Therefore, we added a disclaimer to this paragraph (L. 383-395). Future studies will examine N carry over between cropping seasons following mulching of the legumes in more detail, involving mineral N measurements and nutrient modelling.

L385: it is also unfortunate that the researchers don't present data of N2O fluxes and soil N dynamics off-season. So this observation remains speculative.

Response: Unfortunate, yes, but at least we draw N leaching into consideration.

L385: not clear what is meant with 'emissions were at par', neither why this is striking.

Response: Emission "at par" means emissions were at the same level. To avoid further confusion, we replace this expression with "comparable".

L395: the lack of explanation to the effect on mulching actually calls to explain this by measuring consistently the factors driving N2O fluxes such as moisture content and availability of substrate (inorganic N) over time.

Response: Absolutely! The sentence now reads: "... calls for studies tracing cumulative mulching effects over multiple years and exploring their driving factors in more detail."

L397: the relative effect of soil moisture vs inorganic N could have been tested if the researchers would have collected such data. Now this conclusion leads to speculation.

C14

Response: Yes; see answer above.

L398-410: this study doesn't present solid evidence to sustain this claim, because sowing date doesn't control per se GHG fluxes, but determines the state of soil and weather that the soil+crop system will experience. So giving prescriptions of sowing dates that are not tied to indications of environmental conditions wouldn't be useful at all. In addition, this research didn't find any consistent evidence that legumes increase the emissions beyond the fertilized crop according to Fig 2, which shows that one treatment had higher N₂O fluxes than the control.

Response: Sowing date of legumes in our study had a clear effect on legume development (aboveground biomass yield). We agree that our data provide no basis for prescribing sowing dates, and it was never our intention to do so. The sentence now reads: "... emerge as viable management factors for controlling the accumulation of legume biomass between the maize rows and hence the risk for increased N₂O emissions". See also our response to L. 110.

L412-420, this section needs re-writing to make a comparison instead of a list of studies and their findings.

Response: We do compare emission factors; see line 423 in the original version

L420-424 for this comparison to be useful, please report the biomass measured that was added in year 2 across treatments.

Response: See Table 1

L428, in my opinion the EFs should be re-worked with uncertain parameter ranges to be able to assess how far there are from IPCC. This statement is too crude given the procedures used to estimate the EF.

Response: We agree but want to point out that we use our emission factors solely to compare our numbers with emission factors compiled by Kim et al. (2016) for other SSA agricultural systems, which also were scaled up from a limited number of mea-

C15

surements. It is by no means our intention to challenge IPCC default values. One may be critical to the concept of IPCC Tier 1 emission factors for regions with few flux data, but they are the only tool, for the time being, to compare systems with respect to their propensity to emit N₂O from added reactive N and hence an important criterion when studying intensification effects.

L433 the levels of N inputs could have been underestimated because there were no measurements of the real contributions of the legumes. Which soil has been used over decades? Not clear. Intense use of soils usually leads to loss of fertility not enrichment.

Response: Smallholder farmer in the Rift Valley use little if any fertilization and remove all crop residues for feeding animals, thus they have a negative nutrient balance and lose soil fertility. In comparison, the experimental fields of the university farm are relatively "fertile" as they have experienced N and P fertilization, residue retention and N input from legumes BNF for many years. We believe it is important to point this out, when generalizing our findings for the region.

L441: dynamics of inorganic N not measured.

Response: Yes; that is why we have to speculate here.

L454-474: this piece of text is not needed because it cannot be compared with the experimental results reported here. I would suggest contrasting the experimental results with the literature and avoiding listing all that is known for legumes in completely different climates.

Response: This part discusses benefits and risks of legumes intercropping from the perspective of smallholder farming in the region. We believe that this discussion is important and integral to CSA and the question how to sustainably intensify crop production. Since there are no published studies on legume intercropping covering GHG emissions from this region so far, we resort to similar studies in other regions, trying to relate legume quality and management to nutrient release and N₂O emissions.

C16

L482-482: I understood that the researchers didn't measure the N 'carry over effects'. So this point is speculative.

Response: Yes, we speculate here.

L485-487: this statement could be verified at least against the soil moisture data.

Response: Daily rainfall is shown in figures 1d, 1g and 4d, 4g.

L494: please consider environmental conditions instead of referring to sowing date alone. You could also discuss what would be the incentives for farmers to reduce N₂O emissions.

Response: We agree. We added a sentence reading: "This is complicated by the annual variability in growth conditions and requires active planning of sowing and mulching time by the farmers."

L500: indeed more studies would be needed to confirm and to explain the results obtained. I would suggest reflecting on the need to quantify N₂ fixation, and to follow N mineralisation, especially key for legumes.

Response: We fully agree and have changed the sentence accordingly: "Future studies should attempt to combine flux measurements with inorganic N dynamics and BNF measurements".

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2019-303>, 2019.