

# Supplementary Materials

Effect of legume intercropping on N<sub>2</sub>O emission and CH<sub>4</sub> uptake during maize production in the Ethiopian Rift Valley

Shimelis G Raji<sup>1,2</sup> and Peter Dörsch<sup>1</sup>

<sup>1</sup> Faculty for Environmental Sciences and Resource Management, Norwegian University of Life Sciences (NMBU), 1432 Ås, Norway

<sup>2</sup> College of Agriculture, Hawassa University, P.O, Box 05, Hawassa, Ethiopia

Correspondence to: Peter Dörsch (peter.doersch@nmbu.no)

Table S1: Experimental design of the intercropping trial

No.	Crop type	Treatment	2015		C:N forage legume	2016	
			Mineral N kg ha <sup>-1</sup>	P kg ha <sup>-1</sup>		Mineral N kg ha <sup>-1</sup>	P kg ha <sup>-1</sup>
1	Maize monocrop	Unfertilized	0	-	-	0	-
		Fertilized	64	20	-	64	10
2	Maize+Cr <sup>a</sup>	3 weeks	64	20	12.6	32	10
		6 weeks	64	20		32	10
3	Maize+Lb <sup>b</sup>	3 weeks	64	20	10.07	32	10
		6 weeks	64	20		32	10

Cr<sup>a</sup> Crotalaria, Lb<sup>b</sup> Lablab




Figure S1: Cumulative seasonal CH<sub>4</sub>-C in 2015 (a) and 2016 (b). Error bars denote SEM (n=4). Different letters indicate insignificant differences between treatments. For treatment names, see Fig. 2.