

## Authors response

We thank the Editor for the opportunity to improve the quality of our manuscript and his valuable feedback. We considered all comments and remarks by the editor and the referees and improved our manuscript accordingly. In addition, we clarified some phrasing, corrected typos, and revised punctuation.

## Response to Editor's comments:

*When revising your manuscript I think you should at least discuss how information on microbial biomass would add to an interpretation of your findings. It is not enough to simply state that they were not in the scope of the research objective.*

We added a paragraph explaining how analysis of microbial biomass could give further insights in lines 402-409:

Under N limiting conditions, a higher portion of N is recovered in soil microbial biomass in relation to litter N input (Bending and Turner, 1999, Troung and Marschner, 2018). When N is abundant relative to C availability, excess N is released by soil microorganisms and can be lost as  $N_2O$ . In -Rt, where N availability was low, N was immobilized by soil microorganisms and  $N_2O$  emission were low. When more easily degradable N was added with maize shoots, N released from decomposition of maize shoots presumably fostered decomposition of maize roots (Robertson and Groffman, 2015) and denitrification of excess N leading to strongly increased  $CO_2$  and  $N_2O$  emissions in -RS. To estimate the contribution of plant litter N to mineralization, immobilization, and denitrification,  $^{15}N$  labeled litter together with analysis of microbial biomass N and  $^{15}N_2O$  emissions could be used (e.g. Frimpong and Baggs, 2010; Ladd et al., 1981).

*In addition to the comments by the reviewers I would like to strongly encourage you to use SI-Units throughout your manuscript (including the Figures), eg. a flux should be expressed per g s instead of per kg h.*

We changed figures 2 and 3 a+b to show fluxes per g s and updated the description accordingly.

## Response to Referee Comment #1, uploaded November 14<sup>th</sup>, 2019:

Updates (e.g. line numbers in **bold**).

We are grateful for the positive evaluation of our work and the constructive suggestions to improve the quality of our manuscript. In the following, we reply as comprehensively as possible to each of the comments:

*The manuscript submitted by Rummel and coworkers for publication in Biogeosciences describes the role of litter quality for  $N_2O$  as well as  $CO_2$  emissions as well as bacterial community structure. The*

authors used litter material from maize roots and shoots which were grown under different fertilization levels, applied the materials in a pot experiment to soil which was obtained from an agricultural field and measured for a period of 22 days gas fluxes as well as chemical parameters. At the end of the incubation period also bacterial community structure was analysed. As expected depending on the C:N ratio of the litter material and the availability of easily degradable materials gas emissions and N pools in soil changed, which was also reflected by shifts in bacterial community structure. The study is nicely performed and the data presented of interest, although not totally new. The paper is nicely written and the figures are clear. Like always in such experiments, there is the issue of water content, which was fixed to 50% max WHK, however other water contents would for sure change the results (mainly fluctuation water levels like observed in the field) and also the use of other soil types may induce different response patter. I think here the discussion must be adapted accordingly to make sure that this is showcase but not a general response.

We agree that soil moisture is an important control of N<sub>2</sub>O emissions and changes in water content would affect results. Certainly, fluctuations of water content would induce different response patterns. We **addressed** this topic in a respective paragraph in the discussion (**L. 426-430**):

In addition to soil mineral N concentration and plant litter, soil type and soil moisture may have influenced our results (e.g. Aulakh et al., 1991). Increasing soil moisture leads to increasing N<sub>2</sub>O emissions, but relative contribution of nitrification and denitrification to N<sub>2</sub>O formation may change with increasing soil moisture (Bateman and Baggs, 2005; Baral et al., 2016; Li et al., 2016). Therefore, future experiments with different soil moisture contents should include methods to differentiate between N<sub>2</sub>O formation pathways.

Furthermore there are several issues that need to be considered during revision 1. The description of the sequencing data is very poor. Neither basic data on reads quality rarefraction subsampling etc is given

In addition to the information you find in the manuscript, we included raw reads, reads after filtering, subsample size, observed ASVs, and diversity indices in Table S4 in the Supplementary that was published alongside the manuscript. Rarefaction curves of the observed amplicon sequence variants (ASVs) of the soil inhabiting bacterial communities are displayed in Supplementary Fig. S1.

nor analysis of core microbiomes (together with responders) were made. I guess this is somehow a missed change and the paper would much benefit from a better integration of the molecular data.

As suggested, we analyzed the core microbiomes and respective responders and **included** the following paragraphs in the manuscript. Core microbiome tables (**Table S6-S8**) and Venn diagrams (Fig. S5) **are** included in the Supplementary.

#### 2.5.2 Sequence processing

In addition, core microbiomes and respective responders have been analyzed on genus level, grouped by either the applied litter treatment or N fertilizer levels using *ampvis2* v2.4.7 (**L. 211-212**).

#### 3.5 Bacterial community structure

At the genus level, *Pseudomonas*, *Altererythrobacter*, *Gaiella*, *Nocardioides*, *Agromyces*, *Bacillus*, and *Lysobacter* were most abundant accounting for up to 5.7 % of all ASVs. Accordingly, these were also most abundant genera attributed to the core microbiome (Tables S6 and S8). Overall, 80 genera represented the core microbiome, when grouped by N levels, while 21 genera and 6 genera were identified as responders to N1 and N2, respectively (Fig. S5). In detail, the classified responders to the applied N treatments were the genera *Chthonibacter*, *Luteimonas*, *Sphingobium*, *Novosphingobium*, *Adhaeribacter*, *Nitrospira*, *Gemmata*, and *Devosia* for N1 and *Conexibacter* for N2 samples (Table S. 8). The genera *Bacillus*, *Gaiella*, *Altererythrobacter*, *Blastococcus*, and *Pseudomonas* showed highest abundance in N2 samples, while *Lysobacter*, and *Sphingomonas* were more abundant in N1 samples (Fig. S3). When grouped by litter treatment, the core microbiome comprised 77 genera accounting for 73 % of the relative abundance, while 9, 3 and 10 genera were identified as responders to the applied litter treatments Control, Root and Root+Shoot, respectively (Fig. S5). *Nonomuraea*, *Fluviicola*, and *Nitrospira* responded to the Root+Shoot treatment, while the genera *Lapillicoccus* and *Adhaeribacter* responded to the Root treatment (Table S7). The genera *Litorilinea*, *Gemmata*, *Novosphingobium*, and *Opitutus* were identified as responders to the Control treatment. For N levels and litter treatments respectively, 833 and 838 genera were identified as non-core microbiomes, accounting for 20 % and 19.5 % of relative abundance (Fig. S5) (L. 316-330).

#### 4.3 Bacterial community structures as affected by maize litter and soil N level

The most abundant phyla in our soil samples were the *Actinobacteria*, *Proteobacteria*, and *Chloroflexi*. Among these phyla, the genera *Pseudomonas* (*Proteobacteria*) and *Gaiella* (*Actinobacteria*) were also affiliated to the core microbiomes (L. 444-445). [...]

In our treatments, *Actinobacteria*, *Chloroflexi*, and *Firmicutes* were more abundant in N2 samples, whereas *Bacteroidetes*, and *Nitrospirae* were more abundant in N1 samples which may indicate that the latter are more competitive under conditions of very low mineral nitrogen availability in soil. This was further validated as *Nitrospira* (*Nitrospirae*), known to oxidize nitrite (Koch et al., 2015), was identified as a responder for N1 and -RS (L. 466-469). [...]

Species belonging to the genus *Agromyces* (*Actinobacteria*), which was affiliated to the core microbiomes, are also known to reduce nitrate (Zgurskaya et al., 2008). In addition, species capable of denitrification under anaerobic, O<sub>2</sub>-limited and aerobic conditions can be found in the genera *Bacillus* and *Micromonospora*, as well as *Pseudomonas* and *Rhodococcus* (Verbaendert et al., 2011) that were affiliated to the core microbiome but were more abundant in N2 samples. The genus *Opitutus* was identified as responder to -Cn and comprises the bacterium *Opitutus terrae* that was only found in anoxic habitats in soils (Chin et al., 2001) (L. 471-476). [...]

*Further the sequencing data needs to be submitted to a public database.*

The information on sequence data availability can be found under "Data availability" in line 490: "The 16S rRNA gene sequences were deposited in the National Centre for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under bioproject number PRJNA557843."

*Finally it is general accepted that all DNA extraction kits contain contaminating DNA. Thus a water extraction control would be essential to remove contaminating OTUs from the data.*

As described in line 169ff, we did not use a commercial DNA extraction kit. Instead the DNA extraction protocol by Griffiths et al. (2000) was used. All solutions were sterilized by either autoclaving or sterile filtration. In addition, controls for contamination were carried out during the extractions and in subsequent PCRs.

*2. I miss data on bacterial abundance microbial biomass C and N etc. This information is required and the one hand as soil microbes are an important storage device for N. On the other hand all molecular data is relative, thus to translate the data to absolute numbers biomass values are needed.*

We agree that it is highly likely that microbial biomass varies depending on litter input. Strong differences in CO<sub>2</sub> emissions between treatments indicate differences in microbial activity which could be reasoned by variations in microbial biomass and taxonomy. Nonetheless, the here presented study shows how microbial community composition responds to different litter inputs and whether the relative differences in microbial community structure can be related to CO<sub>2</sub> and N<sub>2</sub>O emissions. However, in the here presented work, changes of microbial biomass were not in the scope of the research objective and therefore this data was not collected. **Nevertheless, we included a paragraph describing how microbial biomass could contribute to interpretation in lines 402-409.**

Under N limiting conditions, a higher portion of N is recovered in soil microbial biomass in relation to litter N input (Bending and Turner, 1999, Troung and Marschner, 2018). When N is abundant relative to C availability, excess N is released by soil microorganisms and can be lost as N<sub>2</sub>O. In -Rt, where N availability was low, N was immobilized by soil microorganisms and N<sub>2</sub>O emission were low. When more easily degradable N was added with maize shoots, N released from decomposition of maize shoots presumably fostered decomposition of maize roots (Robertson and Groffman, 2015) and denitrification of excess N in -RS led to strongly increased CO<sub>2</sub> and N<sub>2</sub>O emissions. To estimate the contribution of plant litter N to mineralization, immobilization, and denitrification, <sup>15</sup>N labeled litter together with analysis of microbial biomass N and <sup>15</sup>N<sub>2</sub>O emissions could be used (e.g. Frimpong and Baggs, 2010; Ladd et al., 1981).

*3. I am quite confused that only three replicates were used for molecular analysis, despite 4 replicates were used for each treatment.*

We used all four replicates of each treatment for molecular analysis. However, for one replicate of N2-Rt, DNA concentration was too low and the 16S rRNA gene PCR was not successful, thus only the remaining three replicates of this treatment were evaluated. In addition to figure and table captions, we included this information in Material & Methods section 2.5.2: **For one replicate of N2-Rt, DNA concentration was very low and the 16S rRNA gene could not be amplified. Thus, we only evaluated the remaining three replicates of this treatment (L. 213-214).**

*Further I wonder why only shoots from N2 were used and not shoots from N1 treatment.*

To be able to compare the litter treatments over soil conditions, we had to use the same litter types for both soil N levels. We **specified** our choices in Materials & Methods section 2.2 (L. 118-122):

The incubation experiment consisted of a two-factorial setup comprising two N levels (N1 and N2) and three litter levels (Control = Cn, Root = Rt, Root+Shoot = RS) (see Table 1 and Figure 1 for details). To allow comparison of litter treatments over soil conditions, the same litter types for both soil N levels were used. As N2 plants had produced greater and healthier biomass during pre-experimental growth phase, only N2 shoots were used for both soils. Roots from N1 and N2 plants were mixed to ensure sufficient amounts for all replicates. [...]

*4. The provided hypothesis is very generic and I guess it must be specified as it is quite obvious that the degree of label materials influences process rates in soil.*

We **specified** our hypotheses as following: We hypothesize that differences in N<sub>2</sub>O emissions between treatments can be related to degradability of maize litter with easier degradable shoot litter leading to higher N<sub>2</sub>O formation. We further expect that differences in litter chemical quality are reflected in the structural composition of the soil microbial community with higher availability of N and C leading to a more specialized community (L. 78-82).

## Response to referee comment #2, uploaded November 14<sup>th</sup>, 2019

Updates (e.g. line numbers in **bold**).

We thank you for thoroughly reading our manuscript and the detailed and constructive comments to improve the quality of our paper. In the following, we will reply to each remark in detail:

*It is a thoroughly conducted study and a well written manuscript, that warrants publication. I have several only minor questions and requests for some clarifications: 1) My biggest confusion when reading the manuscript was the regression analyses between the emissions and the amounts of added litter. Were not the same amounts of litter with the same properties added to each treatment? If that is correct, with only two treatments, how is it possible to do a regression? If that is not correct, better explanations are needed in the Methods.*

We used data of all 24 pots for the regression analyses. The amounts of litter added differed between the three litter treatments as described in Table 1. We agree that having only three litter levels does not allow to draw general conclusions. However, for the soil and litter used in our study, the regressions summarize the relationship between litter quality, mineralization, and N<sub>2</sub>O and CO<sub>2</sub> emissions.

We **included** the missing information in L. **228-230**:

For cumulative CO<sub>2</sub> emissions, regression models included the factors total C input, water-extractable C input, hemicellulose fraction, cellulose fraction, and lignin fraction from all litter treatments (-Cn, -Rt, -RS, n=24).

*2) There is a need to describe the reasoning for some of the experimental choices and decisions that the authors made. a. What was the purpose of growing plants at two different N rates? I presumed that since you had plants grown at two different N levels you would use their litter separately. If the point was that the plants grown at two different rates will generate different N levels in the soil, would it not be just easier to add N to the soil prior to the incubation?*

The main purpose of growing plants at two N rates was to obtain soils with different background mineral N levels for the incubation experiment. We did not add any fresh mineral N immediately before onset of the incubation because we wanted to simulate conditions comparable to agricultural practice in Europe where in most countries farmers are not allowed to add mineral N with crop residues/catch crops. In addition, soil microorganisms adapt to different N availability during plant growth phase.

We **specified** this information in the introduction (L. **84ff**):

Maize plants were grown in a greenhouse to produce root and shoot litter. As **in many European countries law prohibits addition of** mineral N with incorporation of crop residues or catch crops, we applied two N fertilizer regimes (low vs. high) to realize differences in soil N<sub>min</sub> concentration at harvest. We then set up a laboratory incubation experiment with fresh maize root or root and shoot litter under fully controlled conditions and determined hourly CO<sub>2</sub> and N<sub>2</sub>O fluxes for 22 days.

We decided to use a two-factorial design for the incubation experiment. Thus, we used the same litter types for both soil N levels to be able to compare the litter treatments over soil conditions. We will clarify this in Material and Methods section 2.2 (see improved section 2.2 below).

*b. Why the samples were not just incubated in the dark as, commonly done?*

We agree that the information given in L. 137ff **was** misleading and **corrected** it accordingly:

**Pots** were covered with PVC lids, to minimize evaporation from the soil and to incubate samples in the dark.

*3) Some improvement in organization might be warranted. Section 2.2 - I would start the section with a general description of the experiment (what is currently located on ll. 119-120); then add the specific details about shoot and root plant preparations later. As is, it is confusing.*

We improved this section according to your suggestions starting with a general description of the experimental design and explanation of the experimental choices. Then, we describe preparations of treatments and setting up of the experiment:

**L. 117-140:**

## **2.2 Incubation experiment**

The incubation experiment consisted of a two-factorial setup comprising two N levels (N1 and N2) and three litter levels (Control = Cn, Root = Rt, Root+Shoot = RS) (see Table 1 and Figure 1 for details). To allow comparison of litter treatments over soil conditions, the same litter types for both soil N levels were used. As N2 plants had produced greater and healthier biomass during pre-experimental growth phase, only N2 shoots were used for both soils. Roots from N1 and N2 plants were mixed to ensure sufficient amounts for all replicates. Control soils (N1-Cn and N2-Cn) did not receive plant biomass, yet they contained C input from rhizodeposition of the previous maize growth. C remaining from rhizodeposition, root hairs and small root fragments was calculated as the difference in soil C concentration before and after maize growth. For the root treatment, 100 g fresh root biomass was added per kg dry soil (N1-Rt and N2-Rt), and in the root and shoot treatment, 100 g fresh root and 100 g fresh shoot biomass was added per kg dry soil (N1-RS, N2-RS). Each treatment was replicated four times.

Within each N level, soil was homogenized to ensure similar starting conditions. Subsamples of both soils were taken for analysis of mineral N, water extractable C<sub>org</sub> concentration, and total soil C. Soil mineral N concentrations were 0.93 and 1.97 mg N kg<sup>-1</sup> for N1 and N2, respectively. Plant litter was cut to a size of 2 cm and homogeneously mixed with the soil, simulating residue incorporation and tillage. PVC pots with a diameter of 20 cm and a total volume of 6.8 L were filled with fresh soil equivalent to 3.5 kg dry weight previously mixed with plant litter. Soil was compacted in a stepwise mode by filling a 2 cm-layer of soil in pots and compacting it with a plunger. To ensure continuity between soil layers, the surface of the compacted layer was gently scratched before adding the next soil layer. Due to high litter input, target bulk density was 1.1 g cm<sup>-3</sup>. Actual bulk density was determined by measuring headspace height, and these values were used for calculations.

To adjust soil moisture of all pots to 70 % WHC, equivalent to 49 % WFPS, water was dripped on the soil surface through hollow needles (outer diameter 0.9 mm). Pots were covered with PVC lids to minimize evaporation from the soil surface and to incubate samples in the dark. The incubation experiment was carried out under controlled temperature (16 h day at 25 °C, 8 h night at 19 °C) for 22 days. Volumetric water content (VWC) sensors (EC-5, Decagon Devices, Pullman, USA) were used to monitor soil water content.

4) Minor suggestions: a. L.273-274 – this information will be more visible when reported in a table, instead of being buried in the text.

Data on soil  $\text{NO}_3^-$  and  $\text{NH}_4^+$  concentrations are shown in Figures 4 a and b. We **added** a table showing mineralization during the incubation period (**Table 4**).

b. In some places you talk about statistical significance and provide p-values, in others you say how things are different but without mentioning the statistical significance. I suggest being consistent and either only talk about statistically significant differences or specify what is being regarded as numeric and what as statistically significant difference.

We **added** p-values for differences between cumulative  $\text{CO}_2$  and  $\text{N}_2\text{O}$  emission in the text (L. 261ff.). Currently, these values are depicted in Table 3. In all other cases, p-values are given in the text and in the respective tables. We did not conduct statistics on hourly  $\text{N}_2\text{O}$  and  $\text{CO}_2$  fluxes or soil  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , and WEOC concentrations. Thus, we do not provide p-values for these.

#### L. 261-266

To account for different C inputs in treatments, cumulative  $\text{CO}_2$  and  $\text{N}_2\text{O}$  emissions were standardized against the C input per treatment (Table 1). Still, cumulative  $\text{CO}_2$  emissions were almost twice as high in -Rt and about four times higher in -RS compared to -Cn ( $p < 0.05$ ), indicating that differences between litter treatments cannot simply be explained by differences in C input. Addition of maize root and shoot litter increased cumulative  $\text{N}_2\text{O}$  emissions by roughly 100-times compared to control treatments ( $p < 0.05$ ). In contrast, root litter increased cumulative  $\text{N}_2\text{O}$  emissions only by a factor of 5.4 (N1-Rt) and 7 (N2-Rt) compared to the respective controls ( $p < 0.05$ ).

c. L. 351-354 and l. 368-370 – I don't believe that just the correlation results can warrant the conclusions that are stated in these two cases.

We improved these paragraphs as following:

#### L. 351-354 now L. 366-376

Denitrification in soil is largely controlled by the supply of readily decomposable organic matter (Azam et al., 2002; Burford and Bremner, 1975; Loecke and Robertson, 2009), leading to significant correlations between both hourly and cumulative  $\text{N}_2\text{O}$  and  $\text{CO}_2$  emissions (Azam et al., 2002; Fiedler et al., 2017; Frimpong and Baggs, 2010; Huang et al., 2004; Millar and Baggs, 2004, 2005). Hourly  $\text{CO}_2$  fluxes increased directly with onset of incubation and started to decline after day 10, thus mostly C compounds with a short turnover time, i.e. sugars, proteins, starch, and hemicellulose were



decomposed and contributed to CO<sub>2</sub> fluxes. Availability of easily degradable C compounds stimulates microbial respiration, limiting O<sub>2</sub> at the microsite level and thus increasing N<sub>2</sub>O emissions from denitrification (Azam et al., 2002; Chen et al., 2013; Miller et al., 2008). Accordingly, hourly N<sub>2</sub>O fluxes increased after a lag phase of two days. The strong positive correlation ( $R^2=0.9362$ ,  $p \leq 7.632 \times 10^{-15}$ ) between cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions (Table 6) further supports our hypothesis that litter quality, in particular degradability of C compounds, affects N<sub>2</sub>O fluxes from denitrification by creating plant litter associated microsites with low O<sub>2</sub> concentrations.

**L. 368-371 now L. 390-393**

High correlation of cumulative N<sub>2</sub>O emissions and mineralized N during the incubation period ( $R^2=0.5791$ ,  $p < 9.551 \times 10^{-6}$ ) indicates that, in addition to denitrification, heterotrophic nitrification may have contributed to N<sub>2</sub>O production in our study. However, to further differentiate between processes contributing to N<sub>2</sub>O production, stable isotope methods need to be used (Baggs, 2008; Butterbach-Bahl et al., 2013; Van Groenigen et al., 2015; Wrage-Mönnig et al., 2018).

# Maize root and shoot litter quality controls short-term CO<sub>2</sub> and N<sub>2</sub>O emissions and bacterial community structure of arable soil

Pauline Sophie Rummel<sup>1</sup>, Birgit Pfeiffer<sup>1,2</sup>, Johanna Pausch<sup>3</sup>, Reinhard Well<sup>4</sup>, Dominik Schneider<sup>2</sup>, Klaus Dittert<sup>1</sup>

<sup>1</sup>Division of Plant Nutrition and Crop Physiology, Department of Crop Science, University of Göttingen, Germany

<sup>2</sup>Institute of Microbiology and Genetics, Dept. of Genomic and Applied Microbiology, University of Göttingen, Germany

<sup>3</sup>Agroecology, Faculty for Biology, Chemistry, and Earth Sciences, University of Bayreuth, Germany

<sup>4</sup>Thünen Institute, Climate-Smart Agriculture, Braunschweig, Germany

Correspondence to: Pauline Sophie Rummel (pauline.rummel@uni-goettingen.de)

**Abstract.** Chemical composition of root and shoot litter controls decomposition and, subsequently, C availability for biological nitrogen transformation processes in soils. While aboveground plant residues have been proven to increase N<sub>2</sub>O emissions, studies on root litter effects are scarce. This study aimed 1) to evaluate how fresh maize root litter affects N<sub>2</sub>O emissions compared to fresh maize shoot litter, 2) to assess whether N<sub>2</sub>O emissions are related to the interaction of C and N mineralization from soil and litter, and 3) to analyze changes in soil microbial community structures related to litter input and N<sub>2</sub>O emissions.

To obtain root and shoot litter, ~~m~~Maize plants (*Zea mays* L.) were cultivated with two N fertilizer levels in a greenhouse and harvested. A two-factorial 22-day laboratory incubation experiment was set up with soil from both N levels (N1, N2) and three litter addition treatments (Control, Root, Root+Shoot). We measured ~~hourly~~ CO<sub>2</sub> and N<sub>2</sub>O fluxes, analyzed soil ~~nitrate~~mineral N and water extractable organic C (WEOC) concentrations, and determined quality parameters of maize litter.

Bacterial community structures were analyzed using 16S rRNA gene sequencing.

Maize litter quality controlled NO<sub>3</sub><sup>-</sup> and WEOC availability and decomposition related CO<sub>2</sub> emissions. ~~Emissions induced by maize root litter remained low, while H~~high bioavailability of maize shoot litter strongly increased CO<sub>2</sub> and N<sub>2</sub>O emissions ~~when both root and shoot litter were added. -while emissions induced by maize root litter remained low.~~ We identified a strong positive correlation between cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions, supporting our hypothesis that litter quality affects denitrification by creating plant litter associated anaerobic microsites. The interdependency of C and N availability was validated by analyses of regression. Moreover, there was a strong positive interaction between soil NO<sub>3</sub><sup>-</sup> and WEOC concentration resulting in much higher N<sub>2</sub>O emissions, when both NO<sub>3</sub><sup>-</sup> and WEOC were available. A significant correlation was observed between total CO<sub>2</sub> and N<sub>2</sub>O emissions, the soil bacterial community composition and the litter level, showing a clear separation of Root+Shoot samples of all remaining samples. Bacterial diversity decreased with higher N level and higher input of easily available C. Altogether, changes in bacterial community structure reflected degradability of maize litter with easily degradable C from maize shoot litter favoring fast growing C cycling and N reducing bacteria of the phyla *Actinobacteria*, *Chloroflexi*, *Firmicutes* and *Proteobacteria*.

## 1 Introduction

Chemical composition controls decomposition of both roots (Birouste et al., 2012; Redin et al., 2014; Silver and Miya, 2001) and plant litter (Jensen et al., 2005; Kögel-Knabner, 2002; Zhang et al., 2008) and, subsequently, C availability for biological nitrogen transformation processes in soils. When O<sub>2</sub> concentrations are low, denitrifying soil microorganisms may use nitrate (NO<sub>3</sub><sup>-</sup>) as electron acceptor in the respiratory chain to break down organic compounds (Zumft, 1997). This leads to loss of plant available N (Müller and Clough, 2014) and makes soils an important source of the greenhouse gas N<sub>2</sub>O (Ciais et al., 2013).

Plant residues have been proven to increase N<sub>2</sub>O emissions upon incorporation into soil. When different types of litter were compared, quality parameters of plant residues, such as C:N ratio, lignin:N ratio and chemical composition of structural components explained a large share of variances in N<sub>2</sub>O emissions (Baggs et al., 2000; Chen et al., 2013; Millar and Baggs, 2004). Especially in drier soils, denitrification is largely controlled by the supply of readily decomposable organic matter (Azam et al., 2002; Burford and Bremner, 1975; Loecke and Robertson, 2009). Availability of easily degradable C compounds stimulates microbial respiration, limiting O<sub>2</sub> at the microsite level and increasing N<sub>2</sub>O emissions (Azam et al., 2002; Chen et al., 2013; Miller et al., 2008). Furthermore, plant litter enhances local anaerobicity by absorbing water from surrounding pores and retaining high moisture concentrations (Kravchenko et al., 2017, 2018).

While effects of aboveground plant residues on N<sub>2</sub>O emissions have been studied extensively, studies of root residues on N<sub>2</sub>O emissions are scarce. In a temperate forest soil, fine root litter of maize and native tree species did not cause any N<sub>2</sub>O emissions, but a very close interrelation between C mineralization of fine root litter and N<sub>2</sub>O emission was found in other biomes (Hu et al., 2016). In other studies, lower cumulative N<sub>2</sub>O emissions were reported after addition of sugar beet roots compared to leaves (Velthof et al., 2002) and rice roots compared to rice straw (Lou et al., 2007). Furthermore, decomposition dynamics of roots have been studied in great detail, revealing that chemical composition explains most of its variation (Birouste et al., 2012; Johnson et al., 2007; Machinet et al., 2011; Redin et al., 2014; Silver and Miya, 2001; Zhang and Wang, 2015). In general, decomposition rates of hemicelluloses and pectin are higher than that of cellulose, while among cell wall components lignin is most resistant against microbial decomposition (Kögel-Knabner, 2002).

Soil microorganisms are often specialized in specific substrates with fungi being regarded as the main decomposers of plant materials rich in cellulose and lignin, while hemicelluloses and pectin are decomposed by many aerobic and anaerobic bacteria and fungi (Kögel-Knabner, 2002). While the phyla *Firmicutes*, *Proteobacteria*, and *Bacteroidetes* are described as fast growing copiotrophic bacteria that are stimulated by input of easily degradable C compounds (Fierer et al., 2016; Pascault et al., 2013), abundance of *Acidobacteria* decreased following the addition of dissolved organic matter into the soil (Fierer et al., 2016). Similarly, denitrifying microorganisms are found in bacteria, fungi and archaea depending on substrate availability and environmental conditions (Zumft, 1997). Fungi are seen as major contributors to denitrification under aerobic and weakly anaerobic conditions, while bacterial denitrification predominates under strongly anaerobic conditions (Hayatsu et al., 2008). Denitrifying bacteria can be found in most phyla (Zumft, 1997), with dominant populations in

*Pseudomonas* and *Alcaligenes* (Gamble et al., 1977; Megonigal et al., 2013). The most abundant denitrifying bacteria in soil are heterotrophic, and, as such, require a source of electrons or reducing equivalents contained in C compounds of organic matter or plant residues. Availability of organic C may thus affect both decomposing and denitrifying soil microorganisms.

In most reported studies on decomposition and N<sub>2</sub>O emissions, dried and often ground plant material was used. This facilitates a homogenous distribution in soil and minimizes differences between replicates. Nevertheless, drying of fine roots prior to incubation increased their decomposition rate and led to overestimation of decomposition and nutrient cycling rates (Ludovici and Kress, 2006). Additionally, formation of plant litter associated anaerobic hotspots was reduced when ground plant material was homogeneously mixed with the soil, while litter aggregation significantly increased soil N<sub>2</sub>O emissions (Loecke and Robertson, 2009). Differences in N<sub>2</sub>O emissions between two clover species were observed only with intact (but dried) leaves, but not when ground material was used (Kravchenko et al., 2018).

The aim of this study was 1) to evaluate how fresh maize root litter affects N<sub>2</sub>O emissions compared to fresh maize shoot litter, 2) to assess to what extent N<sub>2</sub>O emissions are related to the interaction of C and N mineralization from soil and litter, and 3) to analyze the changes in soil microbial community structures related to litter input and N<sub>2</sub>O emissions. ~~We hypothesize that easily degradable C compounds stimulate microbial respiration in plant litter associated hotspots leading to high N<sub>2</sub>O formation when both C and N availability is high. We hypothesize that differences in N<sub>2</sub>O emissions between treatments can be related to degradability of maize litter with easier degradable shoot litter leading to higher N<sub>2</sub>O formation.~~ We further expect that differences in litter chemical quality are reflected in the structural composition of the soil microbial community with higher availability of N and C leading to a more specialized community.

Maize plants were grown in a greenhouse to produce root and shoot litter. ~~To realize differences in soil N<sub>min</sub> concentration at harvest, two N fertilizer regimes (low vs. high) were applied. As in many European countries law prohibits addition of mineral N with incorporation of crop residues or catch crops, we applied two N fertilizer regimes (low vs. high) to realize differences in soil N<sub>min</sub> concentration at harvest.~~ We then set up a laboratory incubation experiment with fresh maize root or root and shoot litter under fully controlled conditions and determined hourly CO<sub>2</sub> and N<sub>2</sub>O fluxes for 22 days. Soil samples were taken in regular intervals and analyzed for soil mineral N and water-extractable organic C (WEOC) concentrations. At the end of the incubation experiment, soil microbial community structures were analyzed to identify adaptations to litter input.

## 2 Material and Methods

### 2.1 Preparation of plants and soils prior to incubation experiment

The soil for the experiment was collected 10 km south of Göttingen, Germany at the experimental farm Reinshof of the University of Göttingen (51.484°N, 9.923°E). Soil was classified as Gleyic Fluvisol (21 % clay, 68 % silt, 11 % sand) containing 1.5 % C and 2.81 % humus, with a pH (CaCl<sub>2</sub>) = 7.44.

Prior to the incubation experiment, maize plants were cultivated to obtain shoot and root biomass. For maize cultivation, Mitscherlich pots were filled with 5 kg air dried and sieved (2 mm) soil previously mixed with fertilizers (0.2 g N kg<sup>-1</sup> as

NH<sub>4</sub>NO<sub>3</sub>, 0.14 g P kg<sup>-1</sup> as Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub>, 0.2 g K kg<sup>-1</sup> as K<sub>2</sub>SO<sub>4</sub> and 0.04 g Mg kg<sup>-1</sup> as MgSO<sub>4</sub> \* 7 H<sub>2</sub>O including 0.135 g S kg<sup>-1</sup>). Soil moisture was adjusted to 25 Vol. % and volumetric water content (VWC) sensors (EC-5, Decagon  
100 Devices, Pullman, USA) were used to monitor soil water content. Six maize plants (*Zea mays* L. var. Ronaldinio) were sown per pot and cultivated in a greenhouse with 16 h light and 8 h dark cycles. Pots were randomized in regular intervals to avoid microclimatic effects in the greenhouse.

To get different soil mineral N concentrations in soil, a second N fertilizer dose (0.2 g N kg<sup>-1</sup> as Ca(NO<sub>3</sub>)<sub>2</sub> \* 4 H<sub>2</sub>O) was applied to half of the pots six weeks after sowing. Soil with one N dose is referred to as N1 (0.2 g N kg<sup>-1</sup>) and soil with two  
105 N doses is referred to as N2 (2 x 0.2 g N kg<sup>-1</sup>). Plants were harvested 8 weeks after sowing: Maize plants were cut above the soil surface and roots were removed from soil by sieving and handpicking. Fresh roots were shaken and slightly brushed to remove adhering soil.

A subsample of aboveground maize biomass and maize roots was dried at 60 °C to determine dry matter contents and milled to a particle size <1mm. To determine water-extractable C and N concentrations, subsamples were extracted with H<sub>2</sub>O<sub>bidest</sub>  
110 (maize root 1:1000 w/v, maize shoot 1:10000 w/v) for 16 h and analyzed using a multi N/C® Analyzer (Model 3100, Analytik Jena, Jena, Germany). Another subsample was analyzed for the sum of structural components following established feedstuff analysis protocols based on the method proposed by Goering and Van Soest (1970), namely ash free neutral detergent fiber aNDFom, (VDLUF, 2012a) acid detergent fiber ADFom, (VDLUF, 2011), and acid detergent lignin ADL, (VDLUF, 2012b). According to the definitions, hemicellulose, cellulose, and lignin contents were calculated as  
115 following: Hemicellulose = aNDFom - ADFom; Cellulose = ADFom - ADL; Lignin = ADL. Another subsample was milled using a ball mill and total carbon and nitrogen concentrations were analyzed using a C/N analyzer (Model 1110, Carlo Erba, Milano, Italy).

## 2.2 Incubation experiment

The incubation experiment consisted of a two-factorial setup comprising two N levels (N1 and N2) and three litter levels (Control = Cn, Root = Rt, Root+Shoot = RS) (see Table 1 and Figure 1 for details). To allow comparison of litter treatments over soil conditions, the same litter types for both soil N levels were used. As N2 plants had produced greater and healthier biomass during pre-experimental growth phase, only N2 shoots were used for both soils. Roots from N1 and N2 plants were mixed to ensure sufficient amounts for all replicates. Control soils (N1-Cn and N2-Cn) did not receive plant biomass, yet they contained C input from rhizodeposition of the previous maize growth. C remaining from rhizodeposition, root hairs and  
125 small root fragments was calculated as the difference in soil C concentration before and after maize growth. For the root treatment, 100 g fresh root biomass was added per kg dry soil (N1-Rt and N2-Rt), and in the root and shoot treatment, 100 g fresh root and 100 g fresh shoot biomass was added per kg dry soil (N1-RS, N2-RS). Each treatment was replicated four times.

Within each N level, soil was homogenized to ensure similar starting conditions. Subsamples of both soils were taken for analysis of mineral N, water extractable C<sub>org</sub> concentration, and total soil C. Soil mineral N concentrations were 0.93 and  
130

1.97 mg N kg<sup>-1</sup> for N1 and N2, respectively. Plant litter was cut to a size of 2 cm and homogeneously mixed with the soil, simulating residue incorporation and tillage. PVC pots with a diameter of 20 cm and a total volume of 6.8 L were filled with fresh soil equivalent to 3.5 kg dry weight previously mixed with plant litter. Soil was compacted in a stepwise mode by filling a 2 cm-layer of soil in pots and compacting it with a plunger. To ensure continuity between soil layers, the surface of the compacted layer was gently scratched before adding the next soil layer. Due to high litter input, target bulk density was 1.1 g cm<sup>-3</sup>. Actual bulk density was determined by measuring headspace height, and these values were used for calculations. For the incubation experiment, shoot litter was taken from N2 plants only, and cut to a size of 2 cm. Roots of all plants (N1+N2) were mixed and also cut to 2 cm. Within each N level, all soil was mixed to ensure homogeneous initial conditions. Subsamples of both soils were taken for analysis of mineral N, water extractable C<sub>org</sub> concentration, and total soil C. C remaining from rhizodeposition, root hairs and small root fragments was calculated as the difference in soil C concentration before and after maize growth. The incubation experiment consisted of a two-factorial setup comprising two N levels (N1 and N2) and three litter levels (Control = Cn, Root = Rt, Root+Shoot = RS) (see Table 1 and Figure 1 for details). Each treatment was replicated four times. Soil mineral N concentrations were 0.93 and 1.97 mg N kg<sup>-1</sup> for N1 and N2, respectively. Control soils (N1-Cn and N2-Cn) did not receive plant biomass, yet they contained C input from rhizodeposition of the previous maize growth. For the root treatment, 100 g fresh root biomass was added per kg dry soil (N1-Rt and N2-Rt), and in the root and shoot treatment, 100 g fresh root and 100 g fresh shoot biomass was added per kg dry soil (N1-RS, N2-RS). Plant litter was homogeneously mixed with the soil, simulating residue incorporation and tillage. PVC pots with a diameter of 20 cm and a total volume of 6.8 L were filled with fresh soil equivalent to 3.5 kg dry weight previously mixed with plant litter. Soil was compacted in a stepwise mode by filling a 2cm layer of soil in pots and compacting it with a plunger. To ensure continuity between soil layers, the surface of the compacted layer was gently scratched before adding the next soil layer. Due to high litter input, target bulk density was 1.1 g cm<sup>-3</sup>. Actual bulk density was determined by measuring headspace height, and these values were used for calculations. To adjust soil moisture of all pots to 70 % WHC, equivalent to 49 % WFPS, water was dripped on the soil surface through hollow needles (outer diameter 0.9 mm). Pots were covered with PVC lids to minimize evaporation from the soil surface and to incubate samples in the dark. The incubation experiment was carried out under controlled temperature and light conditions (16 h day at 25 °C, 8 h night at 19 °C) for 22 days. Volumetric water content (VWC) sensors (EC-5, Decagon Devices, Pullman, USA) were used to monitor soil water content.

### 2.3 Gas sampling and analysis

Gas fluxes were measured using the closed chamber method (Hutchinson and Mosier, 1981). Gas samples were taken every 12 hours (morning and evening) for the first 15 days and every 24 hours (midday) for the remaining 7 days. Due to technical issues, gas samples taken in the morning of day 10 to day 15 had to be discarded. Before gas sampling, all pots were opened for ventilation to ensure homogenous ambient air background conditions. Pots were closed with gastight PVC lids and 30 ml gas samples were taken from each pot 0, 20, and 40 minutes after closure and filled into pre-evacuated 12 ml-Exetainer

glass bottles (Labco, High Wycombe, UK). Samples were analyzed on a Bruker gas chromatograph (456-GC, Bruker, Billerica, USA) deploying an electron capture detector (ECD) for N<sub>2</sub>O and a thermal conductivity detector (TCD) for CO<sub>2</sub>. Samples were introduced using a Gilson Autosampler (Gilson Inc., Middleton, WI, USA). Data processing was performed using CompassCDS software. The analytical precision was determined by repeated measurements of standard gases (2500 and 550 ppm CO<sub>2</sub>, 307, 760, and 6110 ppm N<sub>2</sub>O) and was consistently < 2 %.

## 2.4 Soil analyses

Soil samples were taken from the pots using a soil auger of 16 mm diameter on 5, 9, 14 and 22 DAO (days after onset of experiment). Holes were closed with glass tubes to avoid variation in the soil surface. Fresh subsamples were analyzed for water extractable C<sub>org</sub> concentration (WEOC), and a subsample was frozen at -20 °C for soil mineral N analysis. Total soil carbon and nitrogen concentrations were analyzed using a C/N analyzer (Model 1110, Carlo Erba, Milano, Italy). For determination of soil mineral N content, frozen samples were extracted with a 0.0125 M CaCl<sub>2</sub> solution (1:5 w/v) for 60 min on an overhead shaker (85 rpm). The extracts were filtered with 615 ¼ filter paper (Macherey - Nagel GmbH & Co. KG, Düren, Germany) and stored at -20 °C. The extracts were analyzed colorimetrically for the concentrations of NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> using the San++Continuous-Flow Analyzer (Skalar Analytical B.V., Breda, The Netherlands). Soil water content was determined with a parallel set of samples. Net N mineralization was calculated as the difference between the NH<sub>4</sub><sup>+</sup>-N + NO<sub>3</sub><sup>-</sup> N concentrations at the start and end of the incubation period plus N lost as N<sub>2</sub>O-N (Eq. 1).

$$\text{Net mineralization} = (\text{NO}_3^- + \text{NH}_4^+)_{\text{end}} - (\text{NO}_3^- + \text{NH}_4^+)_{\text{start}} + \text{N}_2\text{O} \quad (1)$$

WEOC was determined according to Chantigny et al. (2007). Briefly, fresh soil was homogenized with deionized water (1:2 w/v), samples were centrifuged and filtered with 0.45 µm polyether sulfone syringe filters (Labsolute, Renningen, Germany) and stored at -20 °C. The extracts were analyzed using a multi N/C® Analyzer (Analytik Jena, Jena, Germany).

## 2.5 Analysis of bacterial community structures

### 2.5.1 DNA isolation and 16S rRNA gene amplification

To analyze the soil inhabiting bacterial communities, DNA was extracted from 0.5 g (fresh weight) soil sample taken at the end of the incubation experiment (22 DAO) using the DNA extraction protocol described by Griffiths et al. (2000). Plant litter was removed from samples prior to extraction. In brief, cells were mechanically disrupted using bead beating and nucleic acids were extracted using phenol:chloroform:isoamyl alcohol (25:24:1; Carl Roth, Karlsruhe, Germany). Nucleic acids were then precipitated using polyethylene glycol (Carl Roth, Karlsruhe, Germany) and washed with 70 % ice-cold ethanol (VWR, Radnor, Pennsylvania, USA). Subsequently, RNA was removed by RNase A digestion (Thermo Fischer Scientific, Waltham, Massachusetts, USA) as described by the manufacturer. The RNA-free DNA was used for amplification of the V3 to V4 region of the 16S rRNA gene. We used the bacterial primer pair S-D-Bact-0341-b-S-17 and S-D-Bact-0785-a-A-21 targeting the V3-V4 region of the 16S rRNA gene described by Klindworth et al. (2013) with adapters

for Illumina MiSeq sequencing. The PCR reaction mixture contained 5-fold Phusion GC buffer, 200 µM of each of the four deoxynucleoside triphosphates, 5% DMSO, 0.4 µM of each primer, 1 U of Phusion HF DNA polymerase (Fisher Scientific GmbH, Schwerte, Germany), and 25 ng of RNA-free DNA as template. The following cycling scheme was used for DNA amplification: initial denaturation at 98 °C for 5 min and 25 cycles of denaturation at 98 °C for 45 s, annealing at 60 °C for 30 s, and extension at 72 °C for 30 s, followed by a final extension at 72 °C for 10 min. For each sample, PCR reactions were performed in triplicate. Resulting PCR products were pooled in equimolar amounts and purified using the QIAquick Gel Extraction kit (Qiagen, Hilden, Germany) as recommended by the manufacturer. Quantification of the PCR products was performed using the Quant-iT dsDNA HS assay kit and a Qubit fluorometer as described by the manufacturer (Invitrogen GmbH, Karlsruhe, Germany). Indexing of the PCR products was performed by the Göttingen Genomics Lab (G2L, Göttingen, Germany) using the Nextera XT Index kit as recommended by the supplier (Illumina, San Diego, CA, USA) and sequencing of 16S rRNA amplicons was performed using the dual index paired-end approach (2 × 300 bp) with v3 chemistry for the Illumina MiSeq platform.

### 2.5.2 Sequence processing

All bioinformatic processing of sequence data was done using Linux based software packages. Adapter removal and quality filtering of raw paired-end sequences was done using fastp v0.19.6 (Chen et al., 2018), with base correction in overlapped regions, a qualified quality phred of 20, size exclusion of sequences shorter than 50 bp and per read trimming by quality (phred 20). Merging of quality filtered paired-end reads was done by PEAR v0.9.11 (64 bit) with default parameters (Zhang et al., 2014). Primer removal was conducted using cutadapt v1.18 (Martin, 2013). Subsequently, dereplication, denoising, as well as chimera detection and removal (denovo followed by reference based against the SILVA 132 SSU database), was performed with VSEARCH v2.13.0 (64 bit) (Rognes et al., 2016). Taxonomic classification of the amplicons sequence variants (ASVs, 100% sequence identity) was performed with BLAST+ v2.7.1 against the SILVA 132 SSU reference database (Quast et al., 2013). Subsequently, extrinsic domain ASVs and chloroplasts were removed from the dataset. Sample comparisons were performed at the same surveying effort of 5601-0200 sequences. Statistical analyses were done using ASVs in R version 3.5.3 (R Core Team, 2019). ~~All graphs were prepared using the R package ampvis2 v2.4.7 (Andersen et al., 2018) was used to determine species richness, alpha diversity estimates, and rarefaction curves, and to prepare all graphs. were determined using ampvis2 v2.4.7 as well.~~ To visualize the multivariate constrained dispersion, Canonical Correspondence Analysis (CCA) was conducted with Hellinger transformed data (Legendre and Gallagher, 2001), ASV's with a relative abundance lower than 0.1% in any sample were removed. Correlations of environmental parameters to the bacterial communities were analyzed using the *envfit* function of the *vegan* package v2.5-4 (Oksanen et al., 2015) and projected into the ordination with arrows with a p-value cutoff of 0.005. For further statistical analysis of the microbial community composition (on phyla, order and genus level) and diversity (Shannon, Simpson and PD index) multivariate generalized linear models (MGLM; with N level and litter addition as factors) as implemented in the *mvabund* R package v4.0.1 were employed with adjusted p-values (Wang et al., 2019). For the generalized linear model analysis of variance



(MGLM-ANOVA) tests, p-values < 0.05 were considered to be significant. In addition, core microbiomes and respective responders were analyzed on genus level, grouped by either the applied litter treatment or N fertilizer levels using *ampvis2* v2.4.7.

For one replicate of N2-Rt, DNA concentration was very low and the 16S rRNA gene could not be amplified. Thus, we only evaluated the remaining three replicates of this treatment.

In addition, we attempted to analyze the soil inhabiting fungal community using the fungal specific primer set ITS3\_KYO2 and ITS4 (Toju et al., 2012), but were not able to amplify them.

## 2.6 Calculations and statistical analyses

All statistical analyses were performed using the statistical software R version 3.5.2 (R Core Team, 2018). Arithmetic means and standard error of the four replicates were calculated for ~~hourly~~ CO<sub>2</sub> and N<sub>2</sub>O fluxes. Cumulative gas emissions were calculated by linear interpolation between measured fluxes. To account for different C input in treatments, cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions were standardized against the C input per treatment (see Table 1 for details on C input). Tukey's HSD test was used after analysis of variance to test for treatment effects (i.e., N level and litter addition) on cumulative CO<sub>2</sub> emissions. An interaction was identified between N level and litter addition on cumulative N<sub>2</sub>O emissions using interaction plots from the package *HH* v3.1-35 (Heiberger, 2018). A linear model using generalized least squares (glS) was fitted between cumulative N<sub>2</sub>O as response variable and N level, litter addition, and their interaction as fixed effects. Additionally, the model was fitted to account for inhomogeneous within-class variances. Estimated marginal means were then computed to analyze treatment effects using the R package *emmeans* v1.3.4 (Lenth, 2018). Several regression models were tested to analyze the effect of maize litter on cumulative N<sub>2</sub>O emissions including the factors cumulative CO<sub>2</sub> emissions, initial soil NO<sub>3</sub><sup>-</sup> concentration, and net N mineralization during incubation period. For cumulative CO<sub>2</sub> emissions, regression models included the factors total C input, water-extractable C input, hemicellulose fraction, cellulose fraction, and lignin fraction from all litter treatments (-Cn, -Rt, -RS, n=24), total C input from litter, water-extractable C input from litter, hemicellulose fraction from litter input, cellulose fraction from litter input, and lignin fraction from litter input.

To evaluate effects of soil environmental variables on ~~hourly~~ N<sub>2</sub>O and CO<sub>2</sub> fluxes, a linear mixed effect model (lme) was fitted between ~~hourly~~ N<sub>2</sub>O fluxes (ln transformed), soil NO<sub>3</sub><sup>-</sup>-N and WEOC concentrations using the *lme* function from the package *nlme* v3.1-131 (Pinheiro et al., 2017). Pseudo-R<sup>2</sup> for lme was calculated using *r.squaredGLMM* from the package *MuMIn* v1.42.1 (Barton, 2018). Soil NO<sub>3</sub><sup>-</sup>-N and WEOC concentrations between sampling dates were estimated by linear interpolation. Only evening/midday gas measurements were included in model calculations. To account for repeated measurements, incubation vessel and sampling day were set as random effects. Models were compared using maximum likelihood (ML), selected using AIC (Akaike's information criterion), and fitted using restricted maximum likelihood (REML).

All plots were made using the statistical software R version 3.5.2 (R Core Team, 2018) including the packages *plotrix* v3.7.4 (Lemon, 2006), *plot3D* v1.1.1 (Soetaert, 2017), and *viridisLite* v0.3.0 (Garnier, 2018).

### 3 Results

#### 3.1 Chemical analyses of maize litter

Maize root and shoot litter differed in their chemical compositions (Table 2). Dry matter content of maize roots was much higher compared to shoot as roots had not been washed prior to analyses, so some soil adhering to roots was included in dry matter determinations. Thus, we calculated water-extractable concentrations in relation to total C instead of dry matter. Maize shoot litter was characterized by higher concentrations of water-soluble C and N, and a higher share of easily degradable compounds like hemicellulose and cellulose compared to maize roots.

#### 3.2 CO<sub>2</sub> and N<sub>2</sub>O fluxes Hourly and cumulative CO<sub>2</sub> and N<sub>2</sub>O fluxes emissions

Addition of maize litter increased hourly-CO<sub>2</sub> fluxes compared to Control (Fig. 2), where addition of root and shoot litter (N1-RS, N2-RS) resulted in much higher fluxes compared to roots only (N1-Rt, N2-Rt). While absolute emission rates were strongly affected by litter input, time courses were similar in all litter treatments without visible differences between N1 and N2. CO<sub>2</sub> fluxes stayed on a similar level for the first ten days after onset of incubation showing fluctuations between morning and evening sampling times, and then constantly decreased until the end of the experiment.

After a short lag phase right after onset of experiment, hourly-N<sub>2</sub>O emissions increased in all litter treatments compared to control treatments (Fig. 3 a+b). Highest hourly-fluxes were measured in N2-RS, reaching 7.828 µg N<sub>2</sub>O-N kg<sup>-1</sup> h<sup>-1</sup> on day 5. Fluxes stayed on a similar level from day 7 to day 15, and then declined until the end of the experiment. Hourly-N<sub>2</sub>O fluxes from root (N1-Rt, N2-Rt) and control treatments (N1-Cn, N2-Cn) remained on a low level during the whole incubation period (≤ 0.591 µg and ≤ 0.041 µg N<sub>2</sub>O-N kg<sup>-1</sup> h<sup>-1</sup>, for -Rt and -Cn respectively). N<sub>2</sub>O fluxes from N1 were slightly lower than from N2 in both litter treatments. Over all treatments and sampling dates, hourly-CO<sub>2</sub> and N<sub>2</sub>O fluxes were positively correlated (R<sup>2</sup>=0.5993, p<0.001, data not shown).

To account for different C inputs in treatments, cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions were standardized against the C input per treatment (Table 1). Still, cumulative CO<sub>2</sub> emissions were almost twice as high in -Rt and about four times higher in -RS compared to -Cn (p<0.05), indicating that differences between litter treatments cannot simply be explained by differences in C input. Addition of maize root and shoot litter increased cumulative N<sub>2</sub>O emissions by roughly 100-times compared to control treatments (p<0.05). In contrast, root litter increased cumulative N<sub>2</sub>O emissions only by a factor of 5.4 (N1-Rt) and 7 (N2-Rt) compared to the respective controls (p<0.05).

#### 3.3 Soil NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> and water-extractable Corg concentrations

Addition of maize litter affected the time course of soil NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> and WEOC concentrations (Fig. 4 a-c). In control treatments, initial soil NO<sub>3</sub><sup>-</sup> concentrations of 0.93 (N1-Cn) and 1.97 mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup> dry soil (N2-Cn) continuously increased until the end of the experiment reaching concentrations of 8.24 mg N kg<sup>-1</sup> (N1-Cn) and 11.74 mg N kg<sup>-1</sup> (N2-Cn) respectively. Soil NH<sub>4</sub><sup>+</sup> concentrations showed variations on a low level only. Soil NO<sub>3</sub><sup>-</sup> concentrations were continuously

higher in N2 than in N1 and differences in soil  $\text{NH}_4^+$  concentration were small. Higher fertilization in N2 during previous plant growth led to higher residual organic N and higher net N mineralization (7.61 and 10.08 mg N  $\text{kg}^{-1}$  for N1-Cn and N2-Cn, respectively, Table 4) during the incubation experiment. In treatments with litter, soil  $\text{NO}_3^-$  concentrations decreased after an initial increase. In root treatments, soil  $\text{NO}_3^-$  concentrations continuously decreased until the end of the incubation experiment to 1.9 (N1-Rt) and 2.5 mg N  $\text{kg}^{-1}$  (N2-Rt), while in root plus shoot treatments soil  $\text{NO}_3^-$  concentrations increased again until the end of the experiment, reaching concentrations of 9.46 (N1-RS) and 9.52 mg N  $\text{kg}^{-1}$  (N2-RS). During the whole incubation period, soil  $\text{NO}_3^-$  concentrations in -RS were higher than in -Rt. Soil  $\text{NH}_4^+$  concentrations only marginally increased for -Rt. Contrary to -Rt and -Cn, soil  $\text{NH}_4^+$  concentrations increased until the end of the incubation experiment to 1.68 (N1-RS) and 1.52 mg N  $\text{kg}^{-1}$  (N2-RS) in root and shoot treatments. Net N mineralization was 1.44 (N1-Rt) and 1.10 mg N  $\text{kg}^{-1}$  (N2-Rt) in root treatments, and 14.32 (N1-RS) and 14.14 mg N  $\text{kg}^{-1}$  (N2-RS) in root and shoot treatments (Table 4). Maize root litter did not affect WEOC, as concentrations were similar to ~~-Control~~ throughout the incubation period. However, in -RS treatments, WEOC increased after onset of incubation, reaching highest values (45.32 mg C  $\text{kg}^{-1}$ ) for N1-RS at day 9, after which they decreased until the end of the experiment.

### 3.4 Relations between $\text{N}_2\text{O}$ emissions and C and N parameters of plant litter and soil

To identify the effect of N and C availability on hourly- $\text{N}_2\text{O}$  fluxes, a linear mixed effect model was applied. The best model included a significant interaction between soil  $\text{NO}_3^-$  and WEOC ( $p < 0.0394$ , Pseudo- $R^2 = 0.8888$ , Table 54), and incubation vessel and sampling time as random parameters. Predictions of  $\text{N}_2\text{O}$  fluxes based on this model are shown in (Fig. 5). Linear regression analyses were used to identify relations between cumulative  $\text{CO}_2$  and  $\text{N}_2\text{O}$  emissions, litter quality, and N parameters. Either hemicellulose + cellulose fraction or water-extractable C fraction of plant litter explained more than 96 % of variance of total cumulative  $\text{CO}_2$  emissions ( $p < 2.2 \times 10^{-16}$ ) (Table 65). Regression analyses of the relationships between total cumulative  $\text{N}_2\text{O}$  emissions and influencing factors identified a strong positive relationship between total cumulative  $\text{N}_2\text{O}$  emissions and total cumulative  $\text{CO}_2$  emissions ( $R^2 = 0.9362$ ,  $p < 7.632 \times 10^{-15}$ ) (Table 76), and between cumulative  $\text{N}_2\text{O}$  emissions and mineralized N ( $R^2 = 0.5791$ ,  $p < 9.551 \times 10^{-6}$ ), while initial soil  $\text{NO}_3^-$  concentration did not explain any variance.

### 3.5 Bacterial community structure

The comparison over all maize litter treatments revealed that the bacterial diversity was slightly higher in N1 than in N2 soil as shown by a higher number of amplicon sequence variants (ASVs,  $R^2 = 0.1195$ ,  $p = 0.059$ , Fig. S1). In addition, the alpha diversity indices Shannon ( $R^2 = 0.1844$ ,  $p = 0.023$ ) and Simpson ( $R^2 = 0.1131$ ,  $p = 0.065$ ), as well as Faith's phylogenetic diversity (PD;  $R^2 = 0.1844$ ,  $p = 0.059$ ) were higher for N1 than for N2 samples (Table S4). The canonical correspondence analysis revealed a significant correlation ( $p < 0.001$ ) of the bacterial community composition with total  $\text{CO}_2$  ( $R^2 = 0.6758$ ) and  $\text{N}_2\text{O}$  ( $R^2 = 0.6179$ ) emissions, and the litter level, expressed by a clear separation of the N1-RS and N2-RS samples of all other samples (Fig. 6). With increasing C input, N2 samples cluster more closely than N1 samples. No significant correlation of litter level and microbial diversity was observed, PD index increased in N1 samples

with increasing C input, while the opposite was found for N2 samples. Comparison of N1-C<sub>n</sub> and N1-RS revealed no difference in diversity indices (Shannon and Simpson), while N1-R<sub>t</sub> showed lower Shannon and Simpson diversity indices (Table S4). The Shannon diversity index was lowest in N2-R<sub>t</sub> comparing all N2 treatments, while the Simpson index was lowest for N2-RS.

Overall, the soil bacterial communities were dominated by *Actinobacteria*, *Proteobacteria* and *Chloroflexi* accounting for 15 to 31 % (Fig. S2). The highest relative abundance of *Actinobacteria* and *Chloroflexi* was found in N2-R<sub>t</sub> and of *Proteobacteria* in N1-R. Among these phyla, the order *Gaiellales* (*Actinobacteria*), *Sphingomonadales* (*Proteobacteria*) and *Thermomicrobiales* (*Chloroflexi*) showed the highest relative abundance, especially in N2-R<sub>t</sub> (9.3 %), N1-R<sub>t</sub> (7.5 %) and N2-RS (9 %), respectively. Nevertheless, the phyla *Acidobacteria*, *Planctomycetes*, *Verrucomicrobia*, *Gemmatimonadetes*, *Firmicutes*, *Patescibacteria* and *Bacteroidetes* were also detected (>1 %) (Fig. 7). In detail, *Bacteroidetes* and *Gemmatimonadetes* decreased (with a negative slope, but not significant) with increasing N level, while the abundance of *Firmicutes* increased significantly ( $p=0.038$ ). In addition, although present only in low relative abundance, the *Cyanobacteria* decreased significantly ( $p=0.003$ ) with increasing N levels. At the genus level, *Pseudomonas*, *Altererythrobacter*, *Gaiella*, *Nocardioides*, *Agromyces*, *Bacillus*, and *Lysobacter* were most abundant accounting for up to 5.7 % of all ASVs. Accordingly, these were also most abundant genera attributed to the core microbiome (Tables S6 and S8). The genera *Bacillus*, *Gaiella*, *Altererythrobacter*, *Blastococcus*, and *Pseudomonas* showed highest abundance in N2 samples, while *Lysobacter* and *Sphingomonas* were more abundant in N1 samples (Fig. S3). The most abundant classified species found were *Agromyces* sp., *Bacillus* sp. and *Sphingomonas* sp. Nevertheless, species such as *Pseudomonas* sp., *Nitrospira* sp., *Nitrospira briensis*, *Alcaligenes* sp. and *Mesorhizobium* sp. were also identified. Overall, the bacterial community composition was significantly influenced by N level ( $p=0.005$ ) and maize litter treatment ( $p=0.033$ ). Overall, 80 genera represented the core microbiome, when grouped by N levels, while 21 genera and 6 genera were identified as responders to N1 and N2, respectively (Fig. S5). In detail, the classified responders to the applied N treatments were the genera *Chthonibacter*, *Luteimonas*, *Sphingobium*, *Novosphingobium*, *Adhaeribacter*, *Nitrospira*, *Gemmata*, and *Devosia* for N1 and *Conexibacter* for N2 samples (Table S. 8). The genera *Bacillus*, *Gaiella*, *Altererythrobacter*, *Blastococcus*, and *Pseudomonas* showed highest abundance in N2 samples, while *Lysobacter* and *Sphingomonas* were more abundant in N1 samples (Fig. S3). When grouped by litter treatment, the core microbiome comprised 77 genera accounting for 73 % of the relative abundance, while 9, 3 and 10 genera were identified as responders to the applied litter treatments Control, Root and Root+Shoot, respectively (Fig. S5). *Nonomuraea*, *Fluviicola*, and *Nitrospira* responded to the Root+Shoot treatment, while the genera *Lapillicoccus* and *Adhaeribacter* responded to the Root treatment (Table S7). The genera *Litorilinea*, *Gemmata*, *Novosphingobium*, and *Opitutus* were identified as responders to the Control treatment. For N levels and litter treatments respectively, 833 and 838 genera were identified as non-core microbiomes, accounting for 20 % and 19.5 % of relative abundance (Fig. S5).

The most abundant classified species found were *Agromyces* sp., *Bacillus* sp. and *Sphingomonas* sp. Nevertheless, species such as *Pseudomonas* sp., *Nitrospira* sp., *Nitrospira briensis*, *Alcaligenes* sp. and *Mesorhizobium* sp. were also

identified. Overall, the bacterial community composition was significantly influenced by N-level ( $p=0.005$ ) and maize litter treatment ( $p=0.033$ ).

## 4 Discussion

### 4.1 Decomposability of maize litter

Maize root and shoot litter quality controlled  $\text{NO}_3^-$  and WEOC availability and decomposition related  $\text{CO}_2$  emissions during the initial phase of maize litter decomposition. Harvest of plants, removal of roots and mixing of soil fostered mineralization and nitrification, as reflected by gradually increasing soil  $\text{NO}_3^-$  concentrations. The absence of changes in soil  $\text{NH}_4^+$  concentrations in control treatments without litter addition (N1-Cn, N2-Cn) indicate that all  $\text{NH}_4^+$  was directly nitrified. Also in controls, available C was low as indicated by low  $\text{CO}_2$  emissions and decreasing WEOC concentrations.

The potential for mineralization in soil is known to be high after tillage (Höper, 2002) and positive net mineralization has been reported in control soil without litter addition (Machinet et al., 2009; Velthof et al., 2002), and in the fallow period after rice harvest (Aulakh et al., 2001).

Maize shoot litter was characterized by a high share of easily degradable compounds. High percentages of water-soluble N and water-soluble  $\text{C}_{\text{org}}$  from maize shoot litter strongly increased soil WEOC and  $\text{NO}_3^-$  concentrations. Availability of easily degradable compounds was also reflected by strongly increased  ~~$\text{CO}_2$  fluxes hourly~~ and cumulative ~~emission  $\text{CO}_2$  fluxes~~ from N1-RS and N2-RS. While net mineralization in -RS was similar to -Cn, it was very small in -Rt indicating that N from mineralization was immobilized by soil microorganisms to decompose root C compounds (Robertson and Groffman, 2015). Cumulative  $\text{CO}_2$  emissions in litter treatments were clearly higher than in control, but ~~hourly~~- $\text{CO}_2$  fluxes continuously decreased after onset of incubation, as easily degradable C was consumed. This is in accordance with results of Hu et al. (2016), who reported that maize fine root input initially increased  $\text{CO}_2$  fluxes, which then decreased during the first 20 days of incubation.

Mineralization of plant litter may increase soil  $\text{NO}_3^-$  concentrations especially when C:N ratios are low (Li et al., 2013; Millar and Baggs, 2004). However, net N immobilization has been reported after addition of roots of maize (Machinet et al., 2009; Mary et al., 1993; Velthof et al., 2002), wheat (Jin et al., 2008; Velthof et al., 2002), barley and sugar beet (Velthof et al., 2002), reaching a maximum around day 21 (Mary et al., 1993). Chemical composition has been proven to be the primary controller of decomposition rates of both roots (Birouste et al., 2012; Redin et al., 2014; Silver and Miya, 2001) and aboveground plant litter (Jensen et al., 2005; Zhang et al., 2008) of many different species. Slower decomposition of roots compared to leaves and stems was related to differences in chemical composition of plant organs (Jenkinson, 1965; Johnson et al., 2007). Accordingly, decomposition of roots from 16 maize genotypes was controlled by soluble residue components in the short term whereas lignin and the interconnections between cell wall polymers were important in the long-term

(Machinet et al., 2011). In our study, regression analyses identified a strong positive relationship between cumulative CO<sub>2</sub> emissions and water-extractable C fraction of plant litter ( $R^2=0.966$ ,  $p<2.2\cdot10^{-16}$ ) (Table 56).

#### 4.2 N<sub>2</sub>O emissions as affected by biodegradability of maize litter and soil N level

Denitrification in soil is largely controlled by the supply of readily decomposable organic matter (Azam et al., 2002; Burford and Bremner, 1975; Loecke and Robertson, 2009), leading to significant correlations between both N<sub>2</sub>O and CO<sub>2</sub> fluxes and cumulative emissions (Azam et al., 2002; Fiedler et al., 2017; Frimpong and Baggs, 2010; Huang et al., 2004; Millar and Baggs, 2004, 2005). CO<sub>2</sub> fluxes increased directly with onset of incubation and started to decline after day 10, thus mostly C compounds with a short turnover time, i.e. sugars, proteins, starch, and hemicellulose were decomposed and contributed to CO<sub>2</sub> fluxes. Availability of easily degradable C compounds stimulates microbial respiration, limiting O<sub>2</sub> at the microsite level and thus increasing N<sub>2</sub>O emissions from denitrification (Azam et al., 2002; Chen et al., 2013; Miller et al., 2008). Accordingly, N<sub>2</sub>O fluxes increased after a lag phase of two days. The strong positive correlation ( $R^2=0.9362$ ,  $p<7.632\cdot10^{-15}$ ) between cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions (Table 7) further supports our hypothesis that litter quality, in particular degradability of C compounds, affects N<sub>2</sub>O fluxes from denitrification by creating plant litter associated microsites with low O<sub>2</sub> concentrations.

Denitrification in soil is largely controlled by the supply of readily decomposable organic matter (Azam et al., 2002; Burford and Bremner, 1975; Loecke and Robertson, 2009). Availability of easily degradable C compounds stimulates microbial respiration, limiting O<sub>2</sub> at the microsite level and increasing N<sub>2</sub>O emissions (Azam et al., 2002; Chen et al., 2013; Miller et al., 2008) leading to significant correlations between hourly and cumulative N<sub>2</sub>O and CO<sub>2</sub> emissions (Azam et al., 2002; Fiedler et al., 2017; Frimpong and Baggs, 2010; Huang et al., 2004; Millar and Baggs, 2004, 2005). We identified a strong positive correlation ( $R^2=0.9362$ ,  $p<7.632\cdot10^{-15}$ ) between cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions (Table 6), supporting our hypothesis that litter quality, especially quality of C compounds, affects N<sub>2</sub>O fluxes from denitrification by creating plant litter associated microsites with low O<sub>2</sub> concentrations.

High mineralization in -RS treatments may have especially favored coupled nitrification-denitrification where NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup> are produced by nitrifiers in aerobic habitats and subsequently denitrified by denitrifiers in close-by anaerobic habitats (Butterbach-Bahl et al., 2013; Wrage et al., 2001). Here, N<sub>2</sub>O is mainly produced in the interface of aerobic and anaerobic zones, which are typically found in plant litter associated hotspots (Kravchenko et al., 2017). In addition, N<sub>2</sub>O can also be produced aerobically during heterotrophic and autotrophic nitrification (Anderson et al., 1993; Van Groenigen et al., 2015; Wrage et al., 2001; Zhang et al., 2015). In both processes, N<sub>2</sub>O can be formed as byproduct from chemical hydroxylamine oxidation (Butterbach-Bahl et al., 2013; Van Groenigen et al., 2015). Nitrifier denitrification as a pathway of autotrophic nitrification has been reported mostly under soil conditions differing from our study, namely high NO<sub>2</sub><sup>-</sup>, NH<sub>3</sub> or urea concentrations, and low organic C availability (Wrage-Mönnig et al., 2018; Wrage et al., 2001). In contrast, with high availability of organic C and N compounds, high N<sub>2</sub>O emissions from heterotrophic nitrification have been reported (Anderson et al., 1993; Hu et al., 2016; Papen et al., 1989; Wrage et al., 2001). Zhang et al. (2015) reported 72-77% of N<sub>2</sub>O

being produced by heterotrophic nitrification ~~with-from~~ an arable soil under incubation conditions similar to our study. However, Li et al. (2016) estimated that denitrification was the dominant source of N<sub>2</sub>O in residue-amended soil at 40-60 % WFPS. High correlation of cumulative N<sub>2</sub>O emissions and mineralized N during the incubation period ( $R^2=0.5791$ ,  $p<9.551 \times 10^{-6}$ ) indicates that, in addition to denitrification, heterotrophic nitrification may have contributed to N<sub>2</sub>O production in our study. ~~However, To~~ further differentiate ~~further~~ between processes contributing to N<sub>2</sub>O production, stable isotope methods ~~can-need to~~ be used (Baggs, 2008; Butterbach-Bahl et al., 2013; Van Groenigen et al., 2015; Wrage-Mönnig et al., 2018).

Another aim of this study was to investigate the effect of residual mineral N on plant litter induced N<sub>2</sub>O emissions. To this end, we included two N levels that were obtained by different N fertilization during the pre-experimental plant growth phase (N1: 0.2 g N kg<sup>-1</sup>, N2: 2 x 0.2 g N kg<sup>-1</sup>). At the onset of the incubation experiment, soil mineral N concentration was twice as high in N2 compared to N1, but generally very low (0.93 and 1.97 mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup> dry soil for N1 and N2, respectively). Higher N fertilizer input in N2 during plant growth led to lower C input from rhizodeposition (Tab. 1), which is consistent with literature findings (Kuzyakov and Domanski, 2000; Paterson and Sim, 1999). Cumulative N<sub>2</sub>O emissions were in tendency higher in N2 than in N1, suggesting that NO<sub>3</sub><sup>-</sup> was limited, especially in -RS treatments where C availability was highest. In addition, litter chemical quality strongly affected N availability.

Under N limiting conditions, a higher portion of N is recovered in soil microbial biomass in relation to litter N input (Bending and Turner, 1999, Troung and Marschner, 2018). When N is abundant relative to C availability, excess N is released by soil microorganisms and can be lost as N<sub>2</sub>O. In -Rt, where N availability was low, N was immobilized by soil microorganisms and N<sub>2</sub>O emission were low. When more easily degradable N was added with maize shoots, N released from decomposition of maize shoots presumably fostered decomposition of maize roots (Robertson and Groffman, 2015) and denitrification of excess N leading to strongly increased CO<sub>2</sub> and N<sub>2</sub>O emissions in -RS. To estimate the contribution of plant litter N to mineralization, immobilization, and denitrification, <sup>15</sup>N labeled litter together with analysis of microbial biomass N and <sup>15</sup>N<sub>2</sub>O emissions could be used (e.g. Frimpong and Baggs, 2010; Ladd et al., 1981).

The interdependency of C and N availability was further validated by analyses of regression highlighting a strong positive interaction between soil NO<sub>3</sub><sup>-</sup> and WEOC concentrations resulting in much higher N<sub>2</sub>O emissions only when both NO<sub>3</sub><sup>-</sup> and WEOC were available. This further supports our findings that high bioavailability of maize shoot litter increased microbial respiration by heterotrophic microorganisms resulting in plant litter associated hotspots with high N<sub>2</sub>O formation.

Variation in N<sub>2</sub>O emissions is often related to quality parameters of plant residues, mostly the C:N ratio (Baggs et al., 2000; Chen et al., 2013; Millar and Baggs, 2004; Novoa and Tejeda, 2006). Especially easily degradable fractions, such as water-soluble C (Burford and Bremner, 1975) or the holocellulose fraction (hemicelluloses + cellulose) (Jensen et al., 2005), explained a large share of variability of C mineralization and N<sub>2</sub>O emissions, while lignin content was not relevant (Redin et al., 2014; Silver and Miya, 2001). Comparing 28 laboratory and field studies, Chen et al. (2013) reported that microbial growth-induced microsite anaerobicity could be the major driver for the dynamic change in soil N<sub>2</sub>O emissions following residue amendment and Kravchenko et al. (2017) showed that water absorption by plant residues further enhances formation of plant-litter associated anaerobic hotspots. In the initial phase of decomposition, water-soluble compounds (sugars, amino

acids) are leached from litter providing easily degradable compounds for microbial metabolism. After litter addition, CO<sub>2</sub> fluxes increased immediately due to increased respiration, rapidly reducing pO<sub>2</sub>, and creating anaerobic microsites. We anticipate that formation of such hotspots was further enhanced by the amount of litter addition, as litter input was higher in -RS than in -Rt, and higher compared to other studies (Chen et al., 2013).

In addition to soil mineral N concentration and plant litter, soil type and soil moisture may have influenced our results (e. g. Aulakh et al., 1991). Increasing soil moisture leads to increasing N<sub>2</sub>O emissions, but relative contribution of nitrification and denitrification to N<sub>2</sub>O formation may change with increasing soil moisture (Bateman and Baggs, 2005; Baral et al., 2016; Li et al., 2016). Therefore, future experiments with different soil moisture contents should include methods to differentiate between N<sub>2</sub>O formation pathways.

#### 4.3 Bacterial community response to maize litter input and soil N level

After litter addition, the bacterial community adapts within a few days to substrate availability (Pascault et al., 2013). The canonical correspondence analysis (CCA) showed a clear correlation of the soil inhabiting bacterial community, litter input and total CO<sub>2</sub> and N<sub>2</sub>O emissions. As shown by the CCA, the bacterial community structure in N1-RS and N2-RS was distinct from those in the control samples and soil with addition of root residues. Combined addition of root and shoot litter affected the soil bacterial community leading to a less diverse and more specialized community structure, which was also shown by the alpha diversity indices (see supplemental Table S1). A significant reduction of soil bacterial diversity was induced by different N levels, as previously shown by Zeng et al. (2016). In addition, Rousk and Bååth (2007) observed a negative correlation between mineral N addition and bacterial growth, while the addition of barley straw and alfalfa correlated positively. The phylogenetic diversity (PD) supports these findings by showing a more complex picture. While PD in N1 samples increased with increasing C input, it decreased in N2 samples with increasing C input, indicating a shift of the influencing factors from the C input to the N level. Accordingly, the increase of N<sub>2</sub>O emissions from N2 compared to N1 was smaller in -RS where C availability was highest indicating that N was limited here.

The most abundant phyla in our soil samples were the Actinobacteria, Proteobacteria, and Chloroflexi. Among these phyla, the genera Pseudomonas (Proteobacteria) and Gaiella (Actinobacteria) were also affiliated to the core microbiomes. Thermomicrobiales (Chloroflexi) showed the highest abundance in N2 samples, indicating their involvement in N-cycling. ~~The most abundant phyla in our soil samples were the Actinobacteria, Proteobacteria and Chloroflexi. Among these phyla, the genera Pseudomonas (Proteobacteria), Gaiella (Actinobacteria) and Thermomicrobiales (Chloroflexi) showed the highest abundance in the N2 samples, indicating their involvement in N-cycling.~~ ~~Pseudomonas~~ species such as *Pseudomonas aeruginosa*, *P. stutzeri*, and *P. denitrificans* are known to reduce NO<sub>3</sub><sup>-</sup> and to contribute to N<sub>2</sub>O and N<sub>2</sub> emissions (Carlson and Ingraham, 1983). *Gaiella occulta*, belonging to the Actinobacteria, is also known for the reduction of NO<sub>3</sub><sup>-</sup> to NO<sub>2</sub><sup>-</sup> (Albuquerque et al., 2011). The genus *Thermomicrobiales* comprises species which can grow on nitrate, ammonia and



alanine as sole nitrogen sources and are able to hydrolyze cellulose or starch (Houghton et al., 2015). Relative abundance of *Thermomicrobiales* increased with N and C input, indicating favorable growth conditions for this genus (Fig. 7).

We further identified several genera involved in C cycling including members of *Agromyces*, *Bacillus*, and *Micromonospora*, which were also affiliated to the core microbiome. *Agromyces ulmi* was present in low abundance in our samples and it is known to contribute to C cycling in soils through xylanolytic activity (Rivas et al., 2004). Members of the genus *Bacillus* (*Firmicutes*) have been reported to play a crucial role in carbon cycling in a wide range of environments by functions such as plant growth promotion or production of amylases and cellulases (Lyngwi and Joshi, 2014). Among the genus *Bacillus*, we found one species, *Bacillus sp.* KSM-N252, in relatively high abundance (1-2 %) in N2 samples. This species encodes an alkaline endoglucanase, which can hydrolyze cellulose (Endo et al., 2001). Similarly, *Micromonospora* (*Actinobacteria*) are known to produce hydrolytic enzymes showing cellulolytic and xylanolytic activity (Carro et al., 2018; de Menezes et al., 2012). Abundance of *Bacillus sp.* KSM-N252 (N2-Cn 2 %, N2-Rt 1.1 % and N2-RS 0.8 %) and *Micromonospora* (N2-R 1.9 %, N2-RS 1 %) decreased with increasing input of water-extractable C indicating that cellulose was only decomposed when no easily degradable C was available.

Culture-independent sequence techniques have revealed that members of the phyla *Actinobacteria*, *Chloroflexi*, *Firmicutes*, *Bacteroidetes*, and *Nitrospirae* possess *nirK* or *nirS* genes, and can reduce nitrite to nitric oxide (Cantera and Stein, 2007; Nolan et al., 2009). In our treatments, *Actinobacteria*, *Chloroflexi*, and *Firmicutes* were more abundant in N2 samples, whereas *Bacteroidetes* and *Nitrospirae* were more abundant in N1 samples which may indicate that the latter are more competitive under conditions of very low mineral nitrogen availability in soil. This was further validated as *Nitrospira* (*Nitrospirae*), known to oxidize nitrite (Koch et al., 2015), was identified as a responder for N1 and -RS. The reduction of nitrate has been shown for *Mesorhizobium sp.* (Okada et al., 2005) and *Rhizobium sp.* (Daniel et al., 1982). Although only in low abundance, we found these species predominantly in N2 samples. Species belonging to the genus *Agromyces* (*Actinobacteria*), which was affiliated to the core microbiomes, are also known to reduce nitrate (Zgurskaya et al., 2008). In addition, species capable of denitrification under anaerobic, O<sub>2</sub>-limited and aerobic conditions can be found in the genera *Bacillus* and *Micromonospora*, as well as *Pseudomonas* and *Rhodococcus* (Verbaendert et al., 2011) that were affiliated to the core microbiome but were more abundant in N2 samples. The genus *Opitutus* was identified as responder to -Cn and comprises the bacterium *Opitutus terrae* that was only found in anoxic habitats in soils (Chin et al., 2001).

Altogether, the higher relative abundances of C cycling and N reducing bacteria in N2 samples and their affiliation with the core microbiomes reflects the tendency of increased N<sub>2</sub>O emissions with increasing N level and further supports our hypothesis that C and N availability from plant litter were the main drivers of N<sub>2</sub>O emissions in our study.

hat formatiert: Schriftart: Kursiv

hat formatiert: Schriftart: Kursiv

## 5 Conclusions

We examined CO<sub>2</sub> and N<sub>2</sub>O emissions after simulated post-harvest incorporation of maize root or root plus shoot litter in a laboratory incubation study. High bioavailability of maize shoot litter strongly increased microbial respiration in plant litter associated hotspots leading to increased N<sub>2</sub>O emissions when both C and NO<sub>3</sub><sup>-</sup> were available. Coupled nitrification-denitrification and heterotrophic nitrification presumably contributed to N<sub>2</sub>O formation. Maize root litter was characterized by a higher share of slowly degradable C compounds and lower concentrations of water-soluble N, hence formation of anaerobic hotspots was limited and microbial N immobilization restricted N<sub>2</sub>O emissions. Bacterial community structures reflected degradability of maize litter types. Its diversity decreased with increasing C and N availability, favoring fast growing C cycling and N reducing bacteria, namely *Actinobacteria*, *Chloroflexi*, *Firmicutes* and *Proteobacteria*.

**Data availability:** The 16S rRNA gene sequences were deposited in the National Centre for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under bioproject number PRJNA557843. Data from measurements are available upon request from the corresponding author.

**Supplement:** The supporting information related to this study will be published online.

**Author contributions:** PSR, RW and KD designed the experiments and PSR carried them out. BP and DS carried out microbial analyses, sequence processing and provided figures. JP, RW and KD contributed to interpretation of results. PSR prepared the manuscript with contributions from all co-authors.

**Competing interests:** The authors declare no competing interests.

**Acknowledgments:** The authors thank Jakob Streuber, Simone Urstadt, and Finn Malinowski for gas sampling and laboratory analyses, as well as Alexander Silbersdorff (ZfS Statistical Consulting) and Oliver Caré for advice on data handling and statistical analysis.

This study was funded by the Deutsche Forschungsgemeinschaft through the research unit DFG-FOR 2337: Denitrification in Agricultural Soils: Integrated Control and Modelling at Various Scales (DASIM).

## References

Albuquerque, L., França, L., Rainey, F. A., Schumann, P., Nobre, M. F., and Da Costa, M. S.: *Gaiella occulta* gen. nov., sp. nov., a novel representative of a deep branching phylogenetic lineage within the class *Actinobacteria* and proposal of

550 *Gaiellaceae* fam. nov. and *Gaiellales* ord. nov., Syst. Appl. Microbiol., 34(8), 595–599, doi:10.1016/j.syapm.2011.07.001, 2011.

Andersen, K. S., Kirkegaard, R. H., Karst, S. M., and Albertsen, M.: ampvis2: an R package to analyse and visualise 16S rRNA amplicon data, bioRxiv, 299537, doi:10.1101/299537, 2018.

Anderson, I. C., Poth, M., Homstead, J., and Burdige, D.: A comparison of NO and N<sub>2</sub>O production by the autotrophic nitrifier *Nitrosomonas europaea* and the heterotrophic nitrifier *Alcaligenes faecalis*, Appl. Environ. Microbiol., 59(11), 3525–3533, 1993.

555 [Aulakh, M. S., Doran, J. W., Walters, D.T., and Power, J.F.: Legume residue and soil water effects on denitrification in soils of different textures, Soil. Biol. Biochem., 23, 1161–1167, doi:10.1016/0038-0717\(91\)90029-J, 1991.](#)

Aulakh, M. S., Khera, T. S., Doran, J. W., and Bronson, K. F.: Denitrification, N<sub>2</sub>O and CO<sub>2</sub> fluxes in rice-wheat cropping system as affected by crop residues, fertilizer N and legume green manure, Biol. Fertil. Soils, 34, 375–389, doi:10.1007/s003740100420, 2001.

560 Azam, F., Müller, C., Weiske, A., Benckiser, G., and Ottow, J. C. G.: Nitrification and denitrification as sources of atmospheric nitrous oxide - Role of oxidizable carbon and applied nitrogen, Biol. Fertil. Soils, 35(1), 54–61, doi:10.1007/s00374-001-0441-5, 2002.

Baggs, E. M.: A review of stable isotope techniques for N<sub>2</sub>O source partitioning in soils: Recent progress, remaining challenges and future considerations, Rapid Commun. Mass Spectrom., 22(11), 1664–1672, doi:10.1002/rcm.3456, 2008.

565 Baggs, E. M., Rees, R. M., Smith, K. A., and Vinten, A. J. A.: Nitrous oxide emission from soils after incorporating crop residues, Soil Use Manag., 16, 82–87, doi:10.1111/j.1475-2743.2000.tb00179.x, 2000.

[Baral, K. R., Arthur, E., Olesen, J. E., and Petersen, S. P.: Predicting nitrous oxide emissions from manure properties and soil moisture: An incubation experiment, Soil. Biol. Biochem., 97, 112–120, doi:10.1016/j.soilbio.2016.03.005, 2016.](#)

570 Barton, K.: MuMIn: Multi-Model Inference, R package version 1.43.6, <https://cran.r-project.org/package=MuMIn>, 2018.

[Bateman, E. J. and Baggs, E. M.: Contributions of nitrification and denitrification to N<sub>2</sub>O emissions from soils at different water-filled pore space, Biol. Fertil. Soils, 41, 379–388, doi:10.1007/s00374-005-0858-3, 2005.](#)

[Bending, G. D. and Turner, M. K.: Interaction of biochemical quality and particle size of crop residues and its effect on the microbial biomass and nitrogen dynamics following incorporation into soil, Biol. Fertil. Soils, 29, 319–327, doi:10.1007/s003740050559, 1999.](#)

575 Birouste, M., Kazakou, E., Blanchard, A., and Roumet, C.: Plant traits and decomposition: Are the relationships for roots comparable to those for leaves?, Ann. Bot., 109(2), 463–472, doi:10.1093/aob/mcr297, 2012.

Burford, J. R. and Bremner, J. M.: Relationships between the denitrification capacities of soils and total, water-soluble and readily decomposable soil organic matter, Soil Biol. Biochem., 7(6), 389–394, doi:10.1016/0038-0717(75)90055-3, 1975.

580

hat formatiert: Englisch (Vereinigtes Königreich)

- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philos. Trans. R. Soc. B*, 368, 1–13, doi:10.1098/rstb.2013.0122, 2013.
- Cantera, J. J. L. and Stein, L. Y.: Molecular diversity of nitrite reductase genes (*nirK*) in nitrifying bacteria, *Environ. Microbiol.*, 9(3), 765–776, doi:10.1111/j.1462-2920.2006.01198.x, 2007.
- Carlson, C. A. and Ingraham, J. L.: Comparison of denitrification by *Pseudomonas stutzeri*, *Pseudomonas aeruginosa*, and *Paracoccus denitrificans*, *Appl. Environ. Microbiol.*, 45(4), 1247–1253, 1983.
- Carro, L., Nouioui, I., Sangal, V., Meier-Kolthoff, J. P., Trujillo, M. E., Montero-Calasanz, M. D. C., Sahin, N., Smith, D. L., Kim, K. E., Peluso, P., Deshpande, S., Woyke, T., Shapiro, N., Kyrpides, N. C., Klenk, H. P., Göker, M., and Goodfellow, M.: Genome-based classification of micromonosporae with a focus on their biotechnological and ecological potential, *Sci. Rep.*, 8(1), 1–23, doi:10.1038/s41598-017-17392-0, 2018.
- Chantigny, M. H., Angers, D. A., Kaiser, K., and Kalbitz, K.: Extraction and Characterization of Dissolved Organic Matter, in *Soil Sampling and Methods of Analysis*, 2007.
- Chen, H., Li, X., Hu, F., and Shi, W.: Soil nitrous oxide emissions following crop residue addition: a meta-analysis, *Glob. Chang. Biol.*, 19(10), 2956–2964, doi:10.1111/gcb.12274, 2013.
- Chen, S., Zhou, Y., Chen, Y., and Gu, J.: Fastp: An ultra-fast all-in-one FASTQ preprocessor, *Bioinformatics*, 34(17), i884–i890, doi:10.1093/bioinformatics/bty560, 2018.
- Chin, K.-J., Liesack, W., and Janssen, P. H.: *Opitutus terrae* gen. nov., sp. nov., to accommodate novel strains of the division ‘*Verrucomicrobia*’ isolated from rice paddy soil, *Int. J. Syst. Evol. Microbiol.*, 51, 1965–1968, doi:10.1099/00207713-51-6-1965, 2001.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 465–570, 2013.
- Daniel, R. M., Limmer, A. W., Stelle, K. W., and Smith, I. M.: Anaerobic Growth, Nitrate Reduction and Denitrification in 46 Rhizobium Strains, *J. Gen. Microbiol.* 128, 1811–1815, doi:10.1099/00221287-128-8-1811, 1982.
- Endo, K., Hakamada, Y., Takizawa, S., Kubota, H., Sumitomo, N., Kobayashi, T., and Ito, S.: A novel alkaline endoglucanase from an alkaliphilic *Bacillus* isolate: Enzymatic properties, and nucleotide and deduced amino acid sequences, *Appl. Microbiol. Biotechnol.*, 57(1–2), 109–116, doi:10.1007/s002530100744, 2001.
- Fiedler, S. R., Augustin, J., Wrage-Mönnig, N., Jurasinski, G., Gusovius, B., and Glatzel, S.: Potential short-term losses of N<sub>2</sub>O and N<sub>2</sub> from high concentrations of biogas digestate in arable soils, *SOIL*, 3, 161–176, doi:10.5194/soil-3-161-2017, 2017.
- Fierer, N., Bradford, M. A., and Jackson, R. B.: Toward an Ecological Classification of Soil Bacteria, *Ecology*, 88(6), 1354–1364, doi:10.1890/05-1839, 2016.

615 Frimpong, K. A. and Baggs, E. M.: Do combined applications of crop residues and inorganic fertilizer lower emission of  
N<sub>2</sub>O from soil?, *Soil Use Manag.*, 26(4), 412–424, doi:10.1111/j.1475-2743.2010.00293.x, 2010.

Gamble, T. N., Betlach, M. R., and Tiedje, J. M.: Numerically dominant denitrifying bacteria from world soils, *Appl.*  
*Environ. Microbiol.*, 33(4), 926–939, 1977.

Garnier, S.: viridisLite: Default Color Maps from “matplotlib” (Lite Version), R package version 0.3.0, [https://cran.r-](https://cran.r-project.org/package=viridisLite)  
620 [project.org/package=viridisLite](https://cran.r-project.org/package=viridisLite), 2018.

Goering, H. K. and Van Soest, P. J.: Forage fiber analyses, *Agric. Handb. No. 379*, (379), 12–20, 1970.

Griffiths, R. I., Whiteley, A. S., Anthony, G., Donnell, O., Bailey, M. J., Griffiths, R. I., Whiteley, A. S., and Donnell, A. G.  
O.: Rapid Method for Coextraction of DNA and RNA from Natural Environments for Analysis of Ribosomal DNA- and  
rRNA-Based Microbial Community Composition, *Appl. Environ. Microbiol.* 66(12), 1–5, doi:10.1128/AEM.66.12.5488-  
625 5491.2000.Updated, 2000.

Van Groenigen, J. W., Huygens, D., Boeckx, P., Kuypers, T. W., Lubbers, I. M., Rütting, T., and Groffman, P. M.: The soil N  
cycle: New insights and key challenges, *Soil*, 1(1), 235–256, doi:10.5194/soil-1-235-2015, 2015.

Hayatsu, M., Tago, K., and Saito, M.: Various players in the nitrogen cycle: Diversity and functions of the microorganisms  
involved in nitrification and denitrification, *Soil Sci. Plant Nutr.*, 54(1), 33–45, doi:10.1111/j.1747-0765.2007.00195.x,  
630 2008.

Heiberger, R. M.: HH: Statistical Analysis and Data Display: Heiberger and Holland, R package version 3.1-35,  
<https://cran.r-project.org/package=HH>, 2018.

Höper, H.: Carbon and nitrogen mineralisation rates of fens in Germany used for agriculture, in *Wetlands in Central Europe*,  
edited by Broll G., M. W., and P. EM., pp. 149–164, Springer, Berlin, Heidelberg., 2002.

635 Houghton, K. M., Morgan, X. C., Lagutin, K., Mackenzie, A. D., Vyssotskii, M., Mitchell, K. A., McDonald, I. R., Morgan,  
H. W., Power, J. F., Moreau, J. W., Hanssen, E., and Stott, M. B.: *Thermorudis pharmacophila* sp. Nov., a novel member  
of the class *Thermomicrobia* isolated from geothermal soil, and emended descriptions of *Thermomicrobium roseum*,  
*Thermomicrobium carboxidum*, *Thermorudis peleae* and *Sphaerobacter thermophilus*, *Int. J. Syst. Evol. Microbiol.*,  
65(12), 4479–4487, doi:10.1099/ijsem.0.000598, 2015.

640 Hu, X., Liu, L., Zhu, B., Du, E., Hu, X., Li, P., Zhou, Z., Ji, C., Zhu, J., Shen, H., and Fang, J.: Asynchronous responses of  
soil carbon dioxide, nitrous oxide emissions and net nitrogen mineralization to enhanced fine root input, *Soil Biol.*  
*Biochem.*, 92, 67–78, doi:10.1016/j.soilbio.2015.09.019, 2016.

Huang, Y., Zou, J., Zheng, X., Wang, Y., and Xu, X.: Nitrous oxide emissions as influenced by amendment of plant residues  
with different C:N ratios, *Soil Biol. Biochem.*, 36(6), 973–981, doi:10.1016/j.soilbio.2004.02.009, 2004.

645 Hutchinson, G. and Mosier, A.: Improved soil cover method for field measurement of nitrous oxide fluxes, *Soil Sci. Soc.*  
*Am. J.*, 45, 311–316, 1981.

Jenkinson, D. S.: Studies on the Decomposition Of Plant Material in Soil. I. Losses of Carbon from <sup>14</sup>C labelled Ryegrass  
incubated with soil in the field, *J. Soil Sci.*, 16(1), 104–115, doi:10.1111/j.1365-2389.1965.tb01424.x, 1965.

- Jensen, L. S., Salo, T., Palmason, F., Breland, T. A., Henriksen, T. M., Stenberg, B., Pedersen, A., Lundström, C., and Esala, M.: Influence of biochemical quality on C and N mineralisation from a broad variety of plant materials in soil, *Plant Soil*, 273(1–2), 307–326, doi:10.1007/s11104-004-8128-y, 2005.
- Jin, K., Sleutel, S., De Neve, S., Gabriels, D., Cai, D., Jin, J., and Hofman, G.: Nitrogen and carbon mineralization of surface-applied and incorporated winter wheat and peanut residues, *Biol. Fertil. Soils*, 44(4), 661–665, doi:10.1007/s00374-008-0267-5, 2008.
- Johnson, J. M.-F., Barbour, N. W., and Weyers, S. L.: Chemical Composition of Crop Biomass Impacts Its Decomposition, *Soil Sci. Soc. Am. J.*, 71(1), 155, doi:10.2136/sssaj2005.0419, 2007.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glöckner, F. O.: Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies., *Nucleic Acids Res.*, 41(1), 1–11, doi:10.1093/nar/gks808, 2013.
- Koch, H., Lückner, S., Albertsen, M., Kitzinger, K., Herbold, C., Spieck, E., Nielsen, P. H., Wagner, M., and Daims, H.: Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*, *PNAS*, 112, 11371–11376, doi:10.1073/pnas.1506533112, 2015.
- Kögel-Knabner, I.: The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter, *Soil Biol. Biochem.*, 34, 139–162, doi:10.1016/S0038-0717(01)00158-4, 2002.
- Kravchenko, A. N., Toosi, E. R., Guber, A. K., Ostrom, N. E., Yu, J., Azeem, K., Rivers, M. L., and Robertson, G. P.: Hotspots of soil N<sub>2</sub>O emission enhanced through water absorption by plant residue, *Nat. Geosci.*, 10, 496–500, doi:10.1038/NGEO2963, 2017.
- Kravchenko, A. N., Fry, J. E., and Guber, A. K.: Water absorption capacity of soil-incorporated plant leaves can affect N<sub>2</sub>O emissions and soil inorganic N concentrations, *Soil Biol. Biochem.*, 121, 113–119, doi:10.1016/j.soilbio.2018.03.013, 2018.
- Kuzyakov, Y. and Domanski, G.: Carbon input by plants into the soil. Review, *J. Plant Nutr. Soil Sci.*, 163(4), 421–431, doi:10.1002/1522-2624(200008)163:4<421::AID-JPLN421>3.0.CO;2-R, 2000.
- Ladd, J. N., Oades, J. M., and Amato, M.: Microbial biomass formed from <sup>14</sup>C, <sup>15</sup>N-labelled plant material decomposing in soils in the field, *Soil. Biol. Biochem.*, 13, 119–162, doi:10.1016/0038-0717(81)90007-9, 1981.
- Legendre, P. and Gallagher, E. D.: Ecologically meaningful transformations for ordination of species data, *Oecologia*, 129(2), 271–280, doi:10.1007/s004420100716, 2001.
- Lemon, J.: Plotrix: a package in the red light district of R, *R-News*, 6(4), 8–12, 2006.
- Lenth, R.: Emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.4, <https://cran.r-project.org/package=emmeans>, 2018.
- Li, X., Sørensen, P., Olesen, J. E., and Petersen, S. O.: Evidence for denitrification as main source of N<sub>2</sub>O emission from residue-amended soil, *Soil Biol. Biochem.*, 92, 153–160, doi:10.1016/j.soilbio.2015.10.008, 2016.

Li, X., Hu, F., and Shi, W.: Plant material addition affects soil nitrous oxide production differently between aerobic and oxygen-limited conditions, *Appl. Soil Ecol.*, 64, 91–98, doi:10.1016/j.apsoil.2012.10.003, 2013.

Loecke, T. D. and Robertson, G. P.: Soil resource heterogeneity in terms of litter aggregation promotes nitrous oxide fluxes and slows decomposition, *Soil Biol. Biochem.*, 41(2), 228–235, doi:10.1016/j.soilbio.2008.10.017, 2009.

Lou, Y., Ren, L., Li, Z., Zhang, T., and Inubushi, K.: Effect of Rice Residues on Carbon Dioxide and Nitrous Oxide Emissions from a Paddy Soil of Subtropical China, *Water. Air. Soil Pollut.*, 178, 157–168, doi:10.1007/s11270-006-9187-x, 2007.

Ludovici, K. H. and Kress, L. W.: Decomposition and nutrient release from fresh and dried pine roots under two fertilizer regimes, *Can. J. For. Res.*, 36(1), 105–111, doi:10.1139/X05-227, 2006.

Lyngwi, N. A. and Joshi, S.: Economically important *Bacillus* and related genera: a mini review, in *Biology of Useful Plants and Microbes*, 33–43, 2014.

Machinet, G. E., Bertrand, I., Chabbert, B., and Recous, S.: Decomposition in soil and chemical changes of maize roots with genetic variations affecting cell wall quality, *Eur. J. Soil Sci.*, 60(2), 176–185, doi:10.1111/j.1365-2389.2008.01109.x, 2009.

Machinet, G. E., Bertrand, I., Barrière, Y., Chabbert, B., and Recous, S.: Impact of plant cell wall network on biodegradation in soil: Role of lignin composition and phenolic acids in roots from 16 maize genotypes, *Soil Biol. Biochem.*, 43(7), 1544–1552, doi:10.1016/j.soilbio.2011.04.002, 2011.

Martin, M.: Cutadapt removes adapter sequences from high-throughput sequencing reads, *EMBnet.journal*, 17(1), 10–12, doi:10.14806/ej.17.1.200, 2013.

Mary, B., Fresneau, C., Morel, J. L., and Mariotti, A.: C and N cycling during decomposition of root mucilage, roots and glucose in soil, *Soil Biol. Biochem.*, 25(8), 1005–1014, doi:10.1016/0038-0717(93)90147-4, 1993.

Megonigal, J. P., Hines, M. E., and Visscher, P. T.: Anaerobic Metabolism: Linkages to Trace Gases and Aerobic Processes, in *Treatise on Geochemistry: Second Edition*, vol. 10, 273–359., 2013.

de Menezes, A. B., McDonald, J. E., Allison, H. E., and McCarthy, A. J.: Importance of *Micromonospora* spp. as colonizers of cellulose in freshwater lakes as demonstrated by quantitative reverse transcriptase PCR of 16s rRNA, *Appl. Environ. Microbiol.*, 78(9), 3495–3499, doi:10.1128/AEM.07314-11, 2012.

Millar, N. and Baggs, E. M.: Chemical composition, or quality, of agroforestry residues influences N<sub>2</sub>O emissions after their addition to soil, *Soil Biol. Biochem.*, 36, 935–943, doi:10.1016/j.soilbio.2004.02.008, 2004.

Millar, N. and Baggs, E. M.: Relationships between N<sub>2</sub>O emissions and water-soluble C and N contents of agroforestry residues after their addition to soil, *Soil Biol. Biochem.*, 37, 605–608, doi:10.1016/j.soilbio.2004.08.016, 2005.

Miller, M. N., Zebarth, B. J., Dandie, C. E., Burton, D. L., Goyer, C., and Trevors, J. T.: Crop residue influence on denitrification, N<sub>2</sub>O emissions and denitrifier community abundance in soil, *Soil Biol. Biochem.*, 40(10), 2553–2562, doi:10.1016/j.soilbio.2008.06.024, 2008.

715 Müller, C. and Clough, T. J.: Advances in understanding nitrogen flows and transformations: Gaps and research pathways, J. Agric. Sci., 152, S34–S44, doi:10.1017/S0021859613000610, 2014.

Nolan, M., Tindall, B. J., Pomrenke, H., Lapidus, A., Copeland, A., Del Rio, T. G., Lucas, S., Chen, F., Tice, H., Chen, J.-F., Saunders, E., Han, C., Bruce, D., Goodwin, L., Chain, P., Pitluck, S., Ovchinnikova, G., Pati, A., Ivanova, N., Mavromatis, K., Chen, A., Palaniappan, K., Land, M., Hauser, L., Chang, Y.-J., Jeffries, C. D., Brettin, T., Göker, M., Bristow, J., Eisen, J. A., Markowitz, V., Hugenholtz, P., Kyrpides, N. C., Klenk, H.-P., and Detter, J. C.: Complete genome sequence of *Rhodothermus marinus*, Stand. Genomic Sci., 1, 283–291, doi:10.4056.sigs46736, 2009.

720 Novoa, R. S. A. and Tejeda, H. R.: Evaluation of the N<sub>2</sub>O emissions from N in plant residues as affected by environmental and management factors, Nutr. Cycl. Agroecosystems, 75(1–3), 29–46, doi:10.1007/s10705-006-9009-y, 2006.

Okada, N., Nomura, N., Nakajima-Kambe, T., and Uchiyama, H.: Characterization of the Aerobic Denitrification in *Mesorhizobium* sp. Strain NH-14 in Comparison with that in Related Rhizobia, Microbes Environ., 20(4), 208–215, doi:10.1264/jsme2.20.208, 2005.

725 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O’Hara, R. B., Simpson, G. L., Solymos, P., Henry, M., Stevens, H., Szoecs, E., and Wagner, H.: vegan: Community Ecology Package, R Package version 2.3-2, <https://cran.r-project.org/web/packages/vegan/index.html>, 2015.

730 Papen, H., von Berg, R., Hinkel, I., Thoene, B., and Rennenberg, H.: Heterotrophic Nitrification by *Alcaligenes faecalis*: NO<sub>2</sub><sup>-</sup>, NO<sub>3</sub><sup>-</sup>, N<sub>2</sub>O, and NO Production in Exponentially Growing Cultures, Appl. Environ. Microbiol., 55(8), 2068–2072, 1989.

Pascual, N., Ranjard, L., Kaisermann, A., Bachar, D., Christen, R., Terrat, S., Mathieu, O., Lévêque, J., Mougél, C., Henault, C., Lemanceau, P., Péan, M., Boiry, S., Fontaine, S., and Maron, P. A.: Stimulation of Different Functional Groups of Bacteria by Various Plant Residues as a Driver of Soil Priming Effect, Ecosystems, 16(5), 810–822, doi:10.1007/s10021-013-9650-7, 2013.

735 Paterson, E. and Sim, A.: Rhizodeposition and C-partitioning of *Lolium perenne* in axenic culture affected by nitrogen supply and defoliation, Plant Soil, 216(1–2), 155–164, doi:10.1023/A:1004789407065, 1999.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team: nlme: Linear and Nonlinear Mixed Effects Models, R package version 3.1-140, <https://cran.r-project.org/package=nlme>, 2017.

740 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, Nucleic Acids Res., 41(D1), 590–596, doi:10.1093/nar/gks1219, 2013.

R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>, 2018.

745 R Core Team: R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>, 2019.



- Redin, M., Guénon, R., Recous, S., Schmatz, R., de Freitas, L. L., Aita, C., and Giacomini, S. J.: Carbon mineralization in soil of roots from twenty crop species, as affected by their chemical composition and botanical family, *Plant Soil*, 378, 205–214, doi:10.1007/s11104-013-2021-5, 2014.
- 750 Rivas, R., Trujillo, M. E., Mateos, P. F., Martínez-Molina, E., and Velázquez, E.: *Agromyces ulmi* sp. nov., xylanolytic bacterium isolated from *Ulmus nigra* in Spain, *Int. J. Syst. Evol. Microbiol.*, 54(6), 1987–1990, doi:10.1099/ijs.0.63058-0, 2004.
- Robertson, G. P. and Groffman, P. M.: Nitrogen Transformations, in *Soil microbiology, ecology and biochemistry*, edited by E. A. Paul, pp. 421–446, Academic Press, Burlington, Massachusetts, USA., 2015.
- 755 Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F.: VSEARCH: a versatile open source tool for metagenomics, *PeerJ*, 4, e2584, doi:10.7717/peerj.2584, 2016.
- Rousk, J. and Bååth, E.: Fungal and bacterial growth in soil with plant materials of different C/N ratios, *FEMS Microbiol. Ecol.*, 62(3), 258–267, doi:10.1111/j.1574-6941.2007.00398.x, 2007.
- 760 Silver, W. L. and Miya, R. K.: Global patterns in root decomposition: Comparisons of climate and litter quality effects, *Oecologia*, 129(3), 407–419, doi:10.1007/s004420100740, 2001.
- Soetaert, K.: plot3D: Plotting Multi-Dimensional Data, R package version 1.1.1, <https://cran.r-project.org/package=plot3D>, 2017.
- Toju, H., Tanabe, A. S., Yamamoto, S., and Sato, H.: High-coverage ITS primers for the DNA-based identification of Ascomycetes and Basidiomycetes in environmental samples, *PLoS One*, 7(7), doi:10.1371/journal.pone.0040863, 2012.
- 765 Truong, T. H. H. and Marschner, P.: Respiration, available N and microbial biomass N in soil amended with mixes of organic materials differing in C/N ratio and decomposition stage, *Geoderma*, 319, 167-174, doi:10.1016/j.geoderma.2018.01.012, 2018.
- VDLUFA, Ed.: 6.5.2 Bestimmung der Säure-Detergentien-Faser (ADF), in *Methodenbuch Band III Die chemische Untersuchung von Futtermitteln*, pp. 1–4, VDLUFA-Verlag, Darmstadt., 2011.
- 770 VDLUFA, Ed.: 6.5.1 Bestimmung der neutral-Detergentien-Faser (NDF), in *Methodenbuch Band III Die chemische Untersuchung von Futtermitteln*, pp. 1–4, VDLUFA-Verlag, Darmstadt., 2012a.
- VDLUFA, Ed.: 6.5.3 Bestimmung des Säure-Detergentien-Lignings (“Rohlignin”), in *Methodenbuch Band III Die chemische Untersuchung von Futtermitteln*, pp. 1–3, VDLUFA-Verlag, Darmstadt., 2012b.
- 775 Velthof, G. L., Kuikman, P. J., and Oenema, O.: Nitrous oxide emission from soils amended with crop residues, *Nutr. Cycl. Agroecosystems*, 62, 249–261, doi:10.1023/A:1021259107244, 2002.
- Verbaendert, I., De Vos, P., Boon, N., and Heylen, K.: Denitrification in Gram-positive bacteria: an underexplored trait, *Biochem. Soc. Trans.*, 39(1), 254–258, doi:10.1042/bst0390254, 2011.
- Wang, Y., Naumann, U., Eddebuettel, E., Wilshire, J., Warton, D., Byrnes, J., dos Santos Silva, R., Niku, J., Renner, I., and Wright, S.: mvabund: Statistical Methods for Analysing Multivariate Abundance Data, R package version 4.0.1, <https://cran.r-project.org/web/packages/mvabund/index.html>, 2019.
- 780

Wrage-Mönnig, N., Horn, M. A., Well, R., Müller, C., Velthof, G., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited, *Soil Biol. Biochem.*, 123, A3–A16, doi:10.1016/j.soilbio.2018.03.020, 2018.

785 Wrage, N., Velthof, G. L., van Beusichem, M. L., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide, *Soil Biol. Biochem.*, 33, 1723–1732, doi:10.1016/S0038-0717(01)00096-7, 2001.

Zeng, J., Liu, X., Song, L., Lin, X., Zhang, H., Shen, C., and Chu, H.: Nitrogen fertilization directly affects soil bacterial diversity and indirectly affects bacterial community composition, *Soil Biol. Biochem.*, 92, 41–49, doi:10.1016/j.soilbio.2015.09.018, 2016.

790 Zgurskaya, H. I., Evtushenko, L. I., Akimov, V. N., Voyevoda, H. V., Dobrovolskaya, T. G., Lysak, L. V., and Kalakoutskii, L. V.: Emended Description of the Genus *Agromyces* and Description of *Agromyces cerinus* subsp. *cerinus* sp. nov., subsp. nov., *Agromyces cerinus* subsp. *nitratus* sp. nov., subsp. nov., *Agromyces fucosus* subsp. *fucosus* sp. nov., subsp. nov., and *Agromyces fucosus* subsp. *hippuratus* sp. nov., subsp. nov., *Int. J. Syst. Bacteriol.*, 42(4), 635–641, doi:10.1099/00207713-42-4-635, 1992.

Zhang, D., Hui, D., Luo, Y., and Zhou, G.: Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors, *J. Plant Ecol.*, 1(2), 85–93, doi:10.1093/jpe/rtn002, 2008.

795 Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A.: PEAR: A fast and accurate Illumina Paired-End reAd mergeR, *Bioinformatics*, 30(5), 614–620, doi:10.1093/bioinformatics/btt593, 2014.

Zhang, J., Müller, C., and Cai, Z.: Heterotrophic nitrification of organic N and its contribution to nitrous oxide emissions in soils, *Soil Biol. Biochem.*, 84, 199–209, doi:10.1016/j.soilbio.2015.02.028, 2015.

800 Zhang, X. and Wang, W.: The decomposition of fine and coarse roots: Their global patterns and controlling factors, *Sci. Rep.*, 5, 1–10, doi:10.1038/srep09940, 2015.

Zumft, W.: Cell biology and molecular basis of denitrification, *Microbiol. Molecular Biol. Rev.*, 61(4), 533–616, 1997.

Table 1: Two-factorial setup of the incubation experiment. Soil  $N_{\min}$  concentrations were measured directly before onset of the incubation experiment. C input in Control is from rhizodeposition (RD) only, C input in Root is from rhizodeposition and roots, C input in Root + Shoot is from rhizodeposition, roots and shoot biomass. N input is from root and shoot biomass, respectively (**FM = fresh matter**).

N level	$N_{\min}$ [mg $\text{NO}_3^-$ -N $\text{kg}^{-1}$ dry soil]	Treatment	<del>Biomass-Litter</del> input [g FM $\text{kg}^{-1}$ dry soil]	C input [g C $\text{kg}^{-1}$ dry soil]	N input [g N $\text{kg}^{-1}$ dry soil]
N1	0.93	Control	RD	3.47	n.d.
		Root	RD + 100	$3.47 + 4.18 = 7.65$	0.25
		Root + Shoot	RD + 100 + 100	$3.47 + 4.18 + 6.16 = 13.80$	$0.25 + 0.27 = 0.52$
N2	1.97	Control	RD	2.74	n.d.
		Root	RD + 100	$2.74 + 4.18 = 6.92$	0.25
		Root + Shoot	RD + 100 + 100	$2.74 + 4.18 + 6.16 = 13.07$	$0.25 + 0.27 = 0.52$

Table 2: Chemical characteristics of maize root and shoot litter used in the incubation experiment. Hemicellulose and cellulose are expressed relative to lignin content.

	Root	Shoot
Dry matter [%]	62.9	14.7
C:N ratio	17.0	23.2
Lignin:N ratio	2.82	1.44
Water soluble C <sub>org</sub> [% of total C]	11.6	23.4
Water soluble N [% of total N]	8.8	25.8
Hemicellulose (relative content)	3.36	9.08
Cellulose (relative content)	3.18	11.5
Lignin (relative content)	1	1

810

**Table 3: Absolute cumulative N<sub>2</sub>O and CO<sub>2</sub> emissions and relative to C input and N<sub>2</sub>O / CO<sub>2</sub> ratio of 22-day incubation experiment with two pre-incubation-N-levels (N1, N2) and three litter addition treatments (Control = no litter input, Root = 100 g root FM kg<sup>-1</sup> dry soil, Root+Shoot = 100 g root FM + 100 g shoot FM kg<sup>-1</sup> dry soil).**

N level	Treatment	N <sub>2</sub> O [μg N <sub>2</sub> O-N kg <sup>-1</sup> ]		N <sub>2</sub> O [μg N <sub>2</sub> O-N kg <sup>-1</sup> g <sup>-1</sup> C input]		CO <sub>2</sub> [mg CO <sub>2</sub> -C kg <sup>-1</sup> ]		CO <sub>2</sub> [mg CO <sub>2</sub> -C kg <sup>-1</sup> g <sup>-1</sup> C input]		N <sub>2</sub> O/CO <sub>2</sub> ratio [μg N mg <sup>-1</sup> C]
N1	Control	10.21 ± 4.23	a	2.95 ± 1.22	a	141.89 ± 29.74	a	40.94 ± 8.58	a	0.07
	Root	120.91 ± 24.09	b	15.81 ± 3.15	b	533.51 ± 83.19	b	69.78 ± 10.88	b	0.23
	Root+Shoot	4337.31 ± 424.98	c	314.25 ± 30.95	c	2287.23 ± 289.48	c	165.72 ± 20.97	c	1.91
N2	Control	11.35 ± 6.75	a	4.15 ± 2.47	a	129.44 ± 47.47	a	47.30 ± 17.35	a	0.08
	Root	201.14 ± 105.62	ab	29.08 ± 15.27	ab	647.48 ± 196.13	ab	93.61 ± 28.36	b	0.31
	Root+Shoot	5357.87 ± 1193.50	c	409.82 ± 91.30	c	2361.19 ± 287.20	c	180.63 ± 21.97	c	2.25

815 Values represent means (n=4) ± standard deviation. Different letters in the same column indicate a significant difference according to the Tukey's HSD post-hoc tests at  $p \leq 0.05$ .

**Table 4: N mineralization during the incubation period.**

<u>N level</u>	<u>Treatment</u>	<u>N mineralized during incubation</u> <u>[mg N kg<sup>-1</sup>]</u>	
<u>N1</u>	<u>Control</u>	<u>7.61 ± 0.98</u>	<u>b</u>
	<u>Root</u>	<u>1.44 ± 0.72</u>	<u>a</u>
	<u>Root+Shoot</u>	<u>14.32 ± 2.66</u>	<u>c</u>
<u>N2</u>	<u>Control</u>	<u>10.08 ± 1.76</u>	<u>b</u>
	<u>Root</u>	<u>1.10 ± 0.68</u>	<u>a</u>
	<u>Root+Shoot</u>	<u>14.14 ± 4.83</u>	<u>c</u>

Values represent means (n=4) ± standard deviation. Different letters in the same column indicate a significant difference according to the Tukey's HSD post-hoc tests at p ≤ 0.05.

**Table 54:** Significance of fixed effects of soil NO<sub>3</sub><sup>-</sup>-N (mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup>), water extractable organic C (WEOC, mg C kg<sup>-1</sup>) and first-order interaction on N<sub>2</sub>O fluxes (μg N<sub>2</sub>O-N kg<sup>-1</sup> h<sup>-1</sup>; ln transformed) using linear mixed effect model.

	Estimate	Standard error	p value
Intercept	-0.4237	0.2074	0.0416
NO <sub>3</sub> <sup>-</sup> -N	0.0152	0.0248	0.5401
WEOC	0.0210	0.0077	0.0065
NO <sub>3</sub> <sup>-</sup> -N : WEOC	0.0023	0.0011	0.0394

825

**Table 65:** Results of regression analyses of the relationship between total cumulative CO<sub>2</sub> emissions and C quality parameters of plant litter (AICc = Akaike's information criterion).


Regression model	residual standard error	degrees of freedom	adjusted R <sup>2</sup>	p value	AICc
CO <sub>2</sub> ~ total litter C input	274.5	22	0.9213	7.65*e <sup>-14</sup>	342.73
CO <sub>2</sub> ~ water-soluble C input	181.9	22	0.9655	< 2.2*e <sup>-16</sup>	322.98
CO <sub>2</sub> ~ Hemicellulose	272.4	22	0.9225	6.497*e <sup>-14</sup>	342.38
CO <sub>2</sub> ~ Cellulose	221.1	22	0.9489	6.478*e <sup>-16</sup>	332.35
CO <sub>2</sub> ~ Lignin	496.6	22	0.7425	3.873*e <sup>-08</sup>	371.19
CO <sub>2</sub> ~ Hemicellulose + Cellulose	180.2	21	0.9661	< 2.2*e <sup>-16</sup>	324.32




**Table 7.6:** Results of regression analyses of the relationship between total cumulative N<sub>2</sub>O emissions, total cumulative CO<sub>2</sub> emissions and N parameters of plant litter and soil (AIC = Akaike's information criterion).

Regression model	residual standard error	degrees of freedom	adjusted R <sup>2</sup>	p value	AICc
N <sub>2</sub> O ~ CO <sub>2</sub>	593.9	22	0.9366	7.073*e <sup>-15</sup>	379.78
N <sub>2</sub> O ~ initial soil NO <sub>3</sub> <sup>-</sup>	2404	22	-0.03885	0.7119	446.89
N <sub>2</sub> O ~ Mineralized N	2191	22	0.5791	9.551*e <sup>-06</sup>	425.21

Plant cultivation 8 weeks



Incubation experiment 22 days



830 **Figure 1: Preparation and experimental setup of the incubation experiment. N1 (0.2 g N kg<sup>-1</sup>) and N2 (2 x 0.2 g N kg<sup>-1</sup>)**  
 | **referring to the N levels during plant growth. Control soil (N1-C and N2-C) without addition of plant biomasslitter.**  
**Root treatment with addition of 100g fresh root biomass per kg dry soil (N1-R and N2-R) and Root + Shoot treatment**  
**with addition of 100 g root and 100 g shoot biomass per kg dry soil (N1-RS, N2-RS).**

835





Figure 2: Hourly CO<sub>2</sub> emissions fluxes from soils with two N levels (N1, N2) after incorporation of maize root litter (-Rt), maize root + shoot litter (-RS) and control (-Cn) without litter. Error bars show standard error of mean values (n = 4). When not visible, error bars are smaller than the symbols.






Figure 3 a+b: Hourly  $\text{N}_2\text{O}$  emissions fluxes from soils with two N levels (N1, N2) after incorporation of maize root litter (-Rt), maize root + shoot litter (-RS) and control (-Cn) without litter. Error bars show standard error of mean values ( $n = 4$ ). When not visible, error bars are smaller than the symbols. Note: data of figure 3 b are excerpt from 3 a, and are shown with different scaling.




Figure 4 a-c: NO<sub>3</sub><sup>-</sup>, WEOC, and NH<sub>4</sub><sup>+</sup> concentration from soils with two N levels (N1, N2) after incorporation of maize root litter (-Rt), maize root + shoot litter (-RS) and control (-Cn) without litter. Error bars show standard error of mean values (n = 4) (day 0: n=3). When not visible, error bars are smaller than the symbols.





Figure 5: Prediction of  $\text{N}_2\text{O}$  fluxes ( $\mu\text{g N}_2\text{O-N kg}^{-1} \text{ h}^{-1}$ ) ( $\ln$  transformed) based on soil  $\text{NO}_3^-$  and water extractable  $\text{C}_{\text{org}}$  concentrations based on linear mixed-effect model (Pseudo- $R^2=0.89$ ).



**Figure 6:** Canonical Correspondence Analysis (CCA) displaying the compositional distribution of the soil inhabiting bacterial communities between the control (N1-C and N2-C; n=4), root (N1-R and N2-R; n=4 and n=3) and root + shoot treatment (N1-RS and N2-RS; n=4). Significant correlations of total CO<sub>2</sub> and N<sub>2</sub>O emissions are shown by black arrows ( $p \leq 0.005$ ). The relative contribution (eigenvalue) of each axis to the total inertia in the data as well as to the constrained space only, respectively, are indicated in percent at the axis titles.






Figure 7: Heatmap of the 16 most abundant bacterial orders of the soil inhabiting bacterial community in the analyzed soil samples grouped by N levels and litter input treatments (n=4, except for N2 Root: n=3).