

Quantifying the contributions of riverine vs. oceanic nitrogen to hypoxia in the East China Sea

Fabian Große^{1,2}, Katja Fennel¹, Haiyan Zhang^{1,3}, and Arnaud Laurent¹

¹Department of Oceanography, Dalhousie University, Halifax, NS, Canada

²Department of Mathematics and Statistics, University of Strathclyde, Glasgow, United Kingdom

³School of Marine Science and Technology, Tianjin University, Tianjin, China

Correspondence: Fabian Große (fabian.grosse@dal.ca)

Abstract. In the East China Sea, hypoxia (oxygen $\leq 62.5 \text{ mmol m}^{-3}$) is frequently observed off the Changjiang (or Yangtze) River estuary covering up to about 15,000 km². The Changjiang River is a major contributor to hypoxia formation because it discharges large amounts of freshwater and nutrients into the region. However, modelling and observational studies have suggested that intrusions of nutrient-rich oceanic water from the Kuroshio Current also contribute to hypoxia formation. The relative contributions of riverine versus oceanic nutrient sources to hypoxia have not been estimated before. Here, we combine a three-dimensional, physical-biogeochemical model with an element tracing method to quantify the relative contributions of nitrogen from different riverine and oceanic sources to hypoxia formation during 2008–2013. Our results suggest that the hypoxic region north of 30 °N is dominated by Changjiang River inputs, with its nitrogen loads supporting 74% of oxygen consumption. South of 30 °N, oceanic nitrogen sources become more important supporting 39% of oxygen consumption during the hypoxic season, but the Changjiang River remains the main control of hypoxia formation also in this region. Model scenarios with reduced Changjiang River nitrogen loads and reduced open-ocean oxygen levels suggest that nitrogen load reductions can significantly reduce hypoxia in the East China Sea and counteract a potential future decline in oxygen supply from the open ocean into the region.

Copyright statement. TEXT

1 Introduction

In the East China Sea (ECS), hypoxic conditions (i.e. dissolved oxygen (O₂) concentrations $\leq 62.5 \text{ mmol m}^{-3}$) are frequently observed off the Changjiang River (or Yangtze) Estuary covering up to about 15,000 km² (Li et al., 2002; Zhu et al., 2017). Hypoxia was first reported in 1959 (Zhu et al., 2011), and a significant increase in its spatial extent has been observed since the 1980s (Li et al., 2011; Wang, 2009; Wang et al., 2015; Zhu et al., 2011). The Changjiang River is the fifth largest river in the world in terms of freshwater (FW) discharge ($9 \times 10^{11} \text{ m}^3 \text{ y}^{-1}$) and its nutrient concentrations are comparable to other strongly anthropogenically affected rivers (Liu et al., 2003). The observed increase in hypoxic area since the 1980s has been

attributed to elevated nutrient loads due to fertilizer use in the Changjiang watershed (Siswanto et al., 2008; Wu et al., 2019; Yan et al., 2003).

Various studies have suggested that oceanic nutrients also play a role in hypoxia formation in the ECS (Chi et al., 2017; Li et al., 2002; Wang et al., 2018; Zhou et al., 2017b, 2018; Zhu et al., 2011). The importance of oceanic nutrient supply distinguishes hypoxia in the ECS from the otherwise comparable situation in the northern Gulf of Mexico (NGoM), where a similar spatial extent of hypoxic conditions is fueled by freshwater and anthropogenic nutrient inputs from a major river, the Mississippi (Fennel and Testa, 2019). Observations in the ECS indicate that south of 30 °N, intrusions of nutrient-rich water from the Kuroshio Current influence the shelf dynamics (Wang et al., 2018; Zhou et al., 2017b, 2018). These intrusions vary seasonally, with stronger intrusions in winter (Bian et al., 2013; Guo et al., 2006). However, northward water mass transport on the ECS shelf is stronger in summer than in winter (Guo et al., 2006), supported by the weak northeastward winds during the East Asian summer monsoon.

The complexity of the circulation and importance of different nutrient sources for hypoxia development make this system particularly amenable to model analyses with high spatio-temporal resolution (Fan and Song, 2014; Zhao and Guo, 2011; Zhang et al., same issue; Zheng et al., 2016; Zhou et al., 2017a; Zhang et al., 2018). Fan and Song (2014) used simulated salinity and nutrient distributions to show that Changjiang River inputs are transported southward in winter and northward in summer. Zhao and Guo (2011) and Zhou et al. (2017a) used model sensitivity studies to highlight the role of oceanic nutrient sources on productivity and hypoxia in the ECS, respectively. Zhang et al. (2019) combined a physical-biogeochemical model with an element tracing method (e.g. Ménesguen et al., 2006) to quantify the relative contributions of different nutrient sources to primary production (PP). They found that riverine nitrogen (N) supports 56% of water column integrated PP in the ECS regions shallower than 50 m. With organic matter degradation being the main sink of O₂ in the subsurface waters of the ECS (Li et al., 2002), this suggests that riverine N also dominates O₂ consumption. However, a quantification of the relative contributions of riverine versus oceanic nutrient sources to hypoxia has not been available until now.

By combining a high-resolution biogeochemical model with this active element tracing method expanded for the quantification of the contributions to O₂ processes (Große et al., 2017, 2019), we provide such an analysis here. We apply the element tracing method to an implementation of the Regional Ocean Modeling System (ROMS; Fennel et al., 2006; Haidvogel et al., 2008) configured for the ECS (Zhang et al., same issue). This allows us to quantify the contributions of N from different riverine and oceanic sources to hypoxia formation in the ECS and analyze year-to-year and seasonal variability in the individual contributions resulting from the East Asian monsoon cycle. In addition to supplying nutrients, the intrusions of open-ocean subsurface waters are relevant to hypoxia by preconditioning O₂ concentrations in the region. Subsurface O₂ has declined over the past decades in the northwest Pacific (Schmidtko et al., 2017) and is projected to continue decreasing in the future (Bopp et al., 2017), which may further exacerbate hypoxia in the region (Qian et al., 2017). We analyze the effect of reduced open-ocean O₂ concentrations on hypoxia under current and reduced N loads from the Changjiang River and compare our results to previous findings for the NGoM's hypoxic zone.

2.1 The physical-biogeochemical model

We used an implementation of ROMS (Haidvogel et al., 2008) for the ECS (Bian et al., 2013). The model covers the region from 116 °E to 134 °E and from 20 °N to 42 °N with a resolution of 1/12 ° (Fig. 1) with 30 terrain-following σ -layers.




Figure 1. Model domain and bathymetry, with sub-domain used for nitrogen tracing, rivers inside the tracing region, and northern and southern regions used for time series analysis. © The GMT Team, 2018.

The biogeochemical component is based on the N-cycle model of Fennel et al. (2006, 2011) but was expanded to include phosphate Laurent et al. (2012), oxygen (Fennel et al., 2013) and riverine dissolved organic matter (Yu et al., 2015). The model state variables are: nitrate (NO_3^-), ammonium (NH_4^+), phosphate (PO_4^{3-}), one functional group each for phyto- and zooplankton, small and large detritus, riverine dissolved organic matter, and dissolved O_2 . Riverine dissolved organic matter is explicitly represented to account for the more refractory nature of river-borne organic matter compared to organic matter produced in the marine environment. This is also reflected in the one order of magnitude lower remineralization rate compared to small detritus (Yu et al., 2015). Instantaneous benthic remineralization was applied at the sediment-water interface (Fennel et al., 2006, 2011). This implies that all organic matter that sinks to the seafloor is remineralized immediately, with 75% of the deposited N being lost to dinitrogen via benthic denitrification (Fennel et al., 2006). Sediment O_2 consumption is calculated

from the benthic remineralization flux multiplied by a molar ratio of $\sim 115:4$ between O_2 uptake and release of NH_4^+ (Fennel et al., 2013). Light attenuation was expanded by including a term dependent on bottom depth (Zhang et al., same issue). For a complete set of model equations, we refer the reader to the Appendix of Laurent et al. (2017).

Initial and open boundary conditions for temperature and salinity were derived from World Ocean Atlas 2013 version 2 (WOA13-v2) climatologies (Locarnini et al., 2013; Zweng et al., 2013). Temperature and salinity were nudged weekly toward the climatology using a nudging scale of 120 days. Horizontal velocities and sea surface elevation at the open boundaries are based on the SODA reanalysis (Carton and Giese, 2008). Eight tidal constituents (M_2 , S_2 , N_2 , K_2 , K_1 , O_1 , P_1 and Q_1) are imposed using tidal elevations, and tidal currents are derived from the global tide model of Egbert and Erofeeva (2002). Initial and open boundary conditions for NO_3^- , PO_4^{3-} and O_2 are also based on WOA13-v2 (Garcia et al., 2013a, b), while small positive values are prescribed for all other biogeochemical variables. In regions deeper than 100 m and in the Yellow Sea north of $34^\circ N$, NO_3^- concentrations are nudged toward the climatology using time constants of 7 and 10 days, respectively. The results are not sensitive to the exact choice of the nudging time scale and, in fact, are almost indistinguishable if the times scales are varied between one week and two weeks.

The model was run for the period 2006–2013, with the first two years used as spin-up, and forced by 6-hourly wind stress, surface heat and FW fluxes from the ECMWF ERA-Interim dataset (Dee et al., 2011). Daily FW discharge and nutrient loads for 11 rivers were imposed (see supplement Table S1 for river locations), with the Changjiang River being by far the largest. Discharge for the Changjiang is obtained from Datong Hydrological Station (<http://www.cjh.com.cn/en/>). Concentrations of NO_3^- , NH_4^+ and PO_4^{3-} for the Changjiang River were obtained from the monthly Global NEWS data set (Seitzinger et al., 2005). For the other rivers, FW discharge and nutrient loads were prescribed using climatologies (Liu et al., 2009; Tong et al., 2015; Zhang, 1996). Due to the lack of data on organic matter loads, river load concentrations of small and large detritus and dissolved organic N were assumed conservatively at $0.5 \text{ mmol N m}^{-3}$, $0.2 \text{ mmol N m}^{-3}$ and 15 mmol N m^{-3} , respectively. Riverine concentrations for phyto- and zooplankton were assumed equal to their pelagic concentrations. The model demonstrates good skill in representing the hydrography and biogeochemistry of the ECS (Zhang et al., same issue; Figs. S2–S4) and reproduces the main features of the ECS circulation (Fig. S1). The simulation using this setup is hereafter referred to as the reference simulation.

In addition to the reference simulation, three scenario simulations were performed: First, to assess the influence of a reduction in riverine N load on hypoxia, a nutrient reduction scenario was run with 50% smaller N concentrations in the Changjiang River input. All other forcing remained the same. The reduction of riverine N only reflects the assumption that nutrient reductions are achieved primarily by reduced application of industrial fertilizer in agriculture—the main source of excess N loads to the ECS (Yan et al., 2003). Second, to investigate the potential effect of reduced open-ocean O_2 supply, a scenario similar to the reference simulation but with reduced O_2 in the open ocean was performed where the initial and open-boundary O_2 concentrations were reduced by 20% throughout the water column in regions deeper than 200 m. This implies that simulated O_2 transport into the model domain is reduced by 20% relative to the reference simulation, which corresponds to the changes projected by Earth System Models under an RCP8.5 scenario for the Northwest Pacific Ocean at the end of the 21st century (Bopp et al., 2017). Third, the reduced open-ocean O_2 scenario was repeated for reduced river N as in the N reduction scenario.

2.2 Passive freshwater and active nitrogen tracing

Passive dye tracers were used to track FW inputs from the Changjiang River similar to previous ROMS applications (Große et al., 2019; Hetland and Zhang, 2017; Rutherford and Fennel, 2018; Zhang et al., 2010, 2012). The rate of change of the concentration of a passive dye tracer from the i -th source (C_p^i) is described as:

$$\frac{\partial C_p^i}{\partial t} = \nabla \cdot (\overline{\overline{D}} \nabla C_p^i) - \nabla \cdot (C_p^i \mathbf{v}) + S_{C_p^i}. \quad (1)$$

Here, $\overline{\overline{D}}$ is the second-order diffusion tensor (or diffusivity), \mathbf{v} is the velocity vector, and $S_{C_p^i}$ represents the external dye tracer sources (i.e. riverine FW discharge). The dye tracer was initialized with zero in the entire model domain. Changjiang FW discharge had a dye tracer concentration of 1 and the dye tracer was used to analyze the influence of FW from the Changjiang River on stratification (i.e. potential energy anomaly; Simpson, 1981).

In addition, we applied an active element tracing method (Große et al., 2017; Ménesguen et al., 2006; Radtke et al., 2012) to quantify the contributions of different N sources to O_2 consumption. For this purpose, each model variable containing N is subdivided into fractions from the different source regions or rivers. The rate of change of a specific source fraction is described as:

$$\frac{\partial C_X^i}{\partial t} = \nabla \cdot (\overline{\overline{D}} \nabla C_X^i) - \nabla \cdot (C_X^i \mathbf{v}) + R_{C_X} \cdot \frac{C_{X_{con}}^i}{C_{X_{con}}}. \quad (2)$$

C_X and C_X^i represent the concentrations of state variable X (e.g. NO_3^-) and its labeled fraction from the i -th source (e.g. NO_3^- from the Changjiang River), respectively. R_{C_X} describes internal and external sources and sinks of X . The index ‘con’ in the source/sink term implies that the relative concentration of the variable consumed by a process is used; e.g. source-specific nitrification is calculated based on the relative concentration of NH_4^+ . In our study, Eq. (2) is solved diagnostically for N tracers from each traced source using the daily model output for the complete N cycle and the post-processing software of Große et al. (2017), which has been adapted for ROMS (Große et al., 2019). Essentially, a source-specific flux over a calculation time step is the product of the ‘bulk’ flux (e.g. total NO_3^- uptake during PP) with the relative fraction of the source-specific state variable (e.g. NO_3^- containing N from the Changjiang River) divided by its corresponding ‘bulk’ state variable (e.g. NO_3^-).

N tracing is only applied in the region without NO_3^- nudging (see ‘tracing region’ in Fig. 1). Inside this region, we simultaneously traced N from five different sources: the Changjiang River, three other smaller rivers (Minjiang, Oujiang and Qiantangjiang Rivers; grouped into one source; see Table S1), the Taiwan Strait (at the southern boundary of the tracing region), the Kuroshio Current (at the eastern boundary), and the Yellow Sea (at the northern boundary; see Fig. 1). As all N tracers that enter the tracing region across the Taiwan Strait, Kuroshio or Yellow Sea boundaries are labeled as such, this implies that N tracers leaving the tracing region cannot reenter. This tracer setup is similar to that of Zhang et al. (2019), with the difference that we separate the Changjiang River from smaller rivers and that we do not account for atmospheric nutrient deposition.

We applied the N tracing to the reference simulation. Since the initial distributions of N tracers from the different sources are not known, we apply a spin-up procedure to the tracing, which provides the initial distributions of the labeled N tracers at the beginning of the analysis period 2008–2013. We first re-ran year 2006 three times. For the first iteration, all N mass already in

the system was arbitrarily attributed to the small rivers, while subsequent iterations were initialized from the final distribution of the previous iteration. Comparison of the final states of two subsequent iterations confirmed that a dynamic steady state of the labeled N tracer distributions was reached after three iterations. We then ran year 2007 as an additional spin-up year to ensure that the spatial distributions at the beginning of 2008 are not affected by re-running year 2006.

3 Results

3.1 Changjiang freshwater discharge and nitrogen concentrations

The Changjiang River is the main source of FW and nutrients in the ECS. Daily time series of its FW discharge and total nitrogen (TN) concentrations (i.e. the sum of the riverine concentrations of all N variables: NO_3^- , NH_4^+ , small and large detritus, dissolved organic matter, phyto- and zooplankton) are shown in Fig. 2. The FW discharge has a distinct seasonal cycle (highest in summer, lowest in winter) due to the monsoon season with high precipitation in summer (Wang et al., 2008), and significant year-to-year variability. The TN concentrations also show high year-to-year variability. TN concentrations are dominated by NO_3^- concentrations, with detritus and dissolved organic N contributing only 8–13%.




Figure 2. Time series of daily freshwater discharge (black, left y-axis) and monthly total nitrogen (TN) concentration (i.e. the sum of the riverine concentrations of all N variables: NO_3^- , NH_4^+ , small and large detritus, dissolved organic matter, phyto- and zooplankton) in the Changjiang River (red, right y-axis).

3.2 Spatial patterns in oxygen consumption

We focus our analysis of O_2 dynamics on gross O_2 consumption (GOC), i.e. the sum of sediment O_2 consumption (SOC) and water column respiration (WR; incl. nitrification), and integrate over the 12 deepest pelagic layers (analogous to Zhang et al., same issue). This is reasonable as Zhang et al. show that both SOC and WR are relevant O_2 sinks in the hypoxic bottom boundary layer of the ECS. Hypoxia in the ECS is most pronounced between July and November; we hence focus most of our




Figure 3. (a) Gross O_2 consumption (GOC) and (b) relative contribution supported by nitrogen from rivers (Changjiang and other rivers) averaged from July to November 2008–2013. Green line: area affected by hypoxia. Boxes: analysis regions used in sections 3.3 and 3.4.

analyses on this time period. First, we analyze the general spatial patterns in GOC and the relative contributions from riverine and oceanic N sources.

Figures 3a and b show maps of GOC and its relative riverine contribution averaged from July to November over the years 2008 to 2013, respectively. GOC is highest (up to $70 \text{ mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$) in the northern part (30°N to 31°N) of the region typically affected by hypoxia (indicated by the green line). Farther north, GOC is still high (50 to $60 \text{ mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$), but decreases significantly in the offshore and southward directions with the strongest gradient roughly between 29° , 122.5°E and 31°N , 123°E . The riverine contribution to GOC is highest ($>95\%$) in the coastal regions between 30°N and 32°N and steadily decreases offshore and southward. South of 32°N , the strongest gradient in the riverine contribution corresponds to the maximum gradient in GOC.

Based on the hypoxic area locations simulated in our model and the spatial patterns in GOC and its riverine contribution, we defined two distinct analysis regions (black boxes in Fig. 3): a northern region where GOC is supported mainly by riverine N ($>70\%$) and a southern region where the riverine contribution declines strongly from 75% to 20% in southeastward direction.

These regions are used to quantify how the relative contributions of riverine (Changjiang River and smaller rivers) and oceanic
 165 N sources (Kuroshio, Taiwan Strait and Yellow Sea) to GOC differ regionally.

3.3 Year-to-year variability in source-specific oxygen consumption

In order to provide insight in the relative importance of the different N sources for hypoxia formation in the northern and south-
 ern hypoxic regions, Fig. 4 shows time series of average source-specific GOC and total hypoxic area from July to November
 for 2008 to 2013 in both regions. Here, we define total hypoxic area as the area experiencing hypoxia at any time during July
 170 to November. The corresponding values of total GOC and the relative contributions of the different sources are provided in
 Table S2.




Figure 4. Average source-specific gross O₂ consumption (GOC) and total hypoxic area during July–November of 2008 to 2013 in the (a) northern and (b) southern regions (see Fig. 1). Same legend for both panels.

In the northern region (Fig. 4a), riverine N sources (Changjiang and other rivers) account for $78.0 \pm 5.9\%$ of GOC averaged
 over 2008–2013, while oceanic sources support only $22.0 \pm 5.9\%$. The Changjiang River constitutes the largest contribution
 ranging between 63.3% in 2013 and 78.2% in 2010, which are also the years of the smallest and largest hypoxic areas in this
 175 time series, respectively. This indicates that the Changjiang River is the main control on O₂ consumption in the northern region.

In the southern region (Fig. 4b), total GOC is about 24% lower than in the northern region, and the riverine contribution is
 also lower ($61.6 \pm 5.7\%$). The average Changjiang contribution is $56.9 \pm 5.1\%$, while the contributions from the Kuroshio and
 Taiwan Strait account for $19.5 \pm 2.2\%$ and $18.9 \pm 3.2\%$, respectively. Clearly, N sources other than from the rivers play a role
 in the southern region.

180 Interestingly, larger hypoxic areas tend to coincide with high contributions from the Changjiang River and small oceanic
 contributions (2008–2010), while the tends to be Changjiang is less important in years of small hypoxic areas (2011 and 2013).
 This suggests that the water mass distribution is an important factor for controlling the extent of hypoxia in the southern region,
 with a higher Changjiang contribution supporting larger hypoxic areas.

The water mass distribution in the southern region is strongly influenced by large-scale wind patterns, i.e. the East Asian monsoon, and variations in the wind field may affect both seasonal and year-to-year variability in GOC and thus hypoxia. Next, we analyze the seasonality of source-specific GOC and hypoxic area in relation to the large-scale winds in the southern region in years of small and large hypoxic areas.

3.4 Seasonal cycle of oxygen consumption and hypoxia in the southern region

In the southern region, the largest and smallest hypoxic areas are simulated in 2008 and 2013, respectively. Figure 5 presents monthly time series of source-specific GOC and total hypoxic area for both years. In addition, it shows the 6-year monthly average of the meridional wind speed 10 m above sea level (v_{10}) and the corresponding anomalies for both years.




Figure 5. Monthly time series of source-specific contributions to GOC and total hypoxic area (A_H) in the southern region (see Fig. 1) in (a) 2008 (year of largest A_H) and (b) 2013 (smallest A_H), and anomaly of northward wind speed 10 m above sea level (v_{10}) relative to 2008–2013 (‘climatology’) averaged over the ECS (25–33 °N, 119–125 °E). Same legend and axes for both panels.

In general, the Changjiang’s GOC fraction tends to be high in spring (March and April) and in late summer (September). This seasonal pattern is explained by shifts in the dominant wind direction (Yang et al., 2012), where the transition from southward to northward winds from March to July supports a northward transport of both the Changjiang River plume and Kuroshio and Taiwan Strait waters increasing the oceanic contribution to GOC. In contrast, the reversal from northward to southward winds in September results in a southward movement of river-influenced water masses. Year-to-year differences in hypoxia, e.g. in June or September/October, result from differences in the Changjiang contribution. The large Changjiang contribution to GOC in September/October 2008 (Fig. 5a) and in June 2013 (Fig. 5b) are followed by large hypoxic areas, while hypoxia almost vanishes in August 2013 when the Changjiang contribution diminishes. These increases (decreases) in the Changjiang contribution also coincide with significant increases (decreases) in FW thickness and thus stratification (see Fig. S5).

The significant year-to-year differences in both the Changjiang contribution to GOC and thus the hypoxic area can be related to year-to-year variability in the wind field and also in FW discharge. The anomalously weak southward winds in September/October of 2008 resulted in a weaker southward coastal current (see Fig. S6). This allowed for a longer presence

of Changjiang FW and nutrients in the region stimulating organic matter production and stabilizing vertical density stratification (see Fig. S5). The higher FW discharge in 2008 compared to 2013 (see Fig. 2) also supported stronger stratification. Consequently, the Changjiang contribution to GOC and thus hypoxic area remained large through October and only dropped in November. In June 2013, the anomalously weak northward winds allowed Changjiang water to be transported into the region (see Fig. S6). This caused an increase in GOC and – supported by less wind-induced mixing – stratification, and thus hypoxia. The opposite occurred in July/August of 2013, when anomalously strong northward winds pushed the Changjiang River water northward resulting in a decrease in GOC, stratification and thus hypoxia.

The results from sections 3.3 and 3.4 highlight the importance of Changjiang River nutrients for hypoxia formation in both the northern and the southern regions and suggest that N load reductions in the Changjiang River would mitigate hypoxia also in the southern region. This raises the question how a potential future decline in open-ocean O₂ concentrations would affect hypoxia in the ECS, which is analyzed next.

215 3.5 Potential future changes in hypoxia

The open-ocean water masses travelling to the hypoxic region are not only relevant as N sources but may also act as sources of low-O₂ water, thus preconditioning the region for hypoxia. Consequently, a future decline in subsurface open-ocean O₂ concentrations may exacerbate hypoxia in the ECS. To investigate how reduced open-ocean O₂ concentrations and reduced Changjiang River N loads would affect hypoxia, we performed the scenario simulations described in section 2.1. Figure 6 shows the cumulative hypoxic exposure in the bottom layer from July to November averaged over 2008–2013 for the reference simulation (Fig. 6a), the nutrient reduction scenario with 50% lower Changjiang River N loads (Fig. 6b), the scenario with 20% lower open-ocean O₂ concentrations (Fig. 6c) and the combined scenario with reduced N load and lower open-ocean O₂ (Fig. 6d). Table 1 presents the corresponding changes in different hypoxia metrics between the reference case and the scenarios.

In the reference case (Fig. 6a), hypoxia occurs between 28–32 °N and from the coast to about 123 °E, on average affecting 225 $19.3 \pm 8.1 \times 10^3 \text{ km}^2$. Maximum hypoxic exposure is 28 days in the central parts of the hypoxic region. Hypoxia is still present in the simulation with 50% lower Changjiang River N loads (Fig. 6b), but its areal extent and hypoxic exposure are significantly reduced. Areas affected by continuous hypoxia of more than one and two weeks are reduced by >60%. Figure S7 shows that the 50% N load reduction in the Changjiang River significantly reduces the Changjiang contribution to GOC and the hypoxic area in the southern analysis region throughout the seasonal cycles of 2008 and 2013 (analogous to Fig. 5). Hypoxia vanishes entirely in most months with small hypoxic areas under current N loads (May to August 2008, September/October 2013), 230 suggesting a relevant role of riverine N load reductions for hypoxia mitigation.

Under reduced open-ocean O₂ concentrations (Fig. 6c), hypoxic area increases by about 50% and regions affected by continuous hypoxia for more than one and two weeks expand by 86% and 118%, respectively (Table 1). Under reduced open-ocean O₂ and reduced Changjiang River N load (Fig. 6d), simulated changes relative to the reference are still negative, although 235 comparably small (see Table 1). This suggests that N load reductions constitute a potent means to mitigate hypoxia under present conditions and to counteract deterioration of O₂ conditions in ECS due to open-ocean deoxygenation.




Figure 6. Average cumulative hypoxia during 2008–2013 derived from simulated bottom O_2 concentrations for (a) the reference simulation, (b) the nitrogen reduction scenario, (c) the reference simulation with reduced oceanic O_2 concentrations, and (d) the nitrogen reduction scenario with reduced oceanic O_2 concentrations.

4 Discussion

4.1 Model validation and study limitations

The here applied ROMS model demonstrates good skill in reproducing the hydrography and general circulation of the ECS (Zhang et al., same issue; Fig. S1) and agrees well with both observations (Zhou et al., 2017b, 2018) and other modeling studies (Bian et al., 2013; Guo et al., 2006; Yang et al., 2011, 2012).

The simulated spatial distributions of dissolved inorganic nitrogen (DIN) are in agreement with observations of Gao et al. (2015) (see Figs. S2–S4). This, together with the good representation of the circulation, is essential for reliable results of the N tracing. Monthly averaged simulated SOC rates in the analysis regions result in $0.3\text{--}69.8 \text{ mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$ and are usually within the observed range of $1.7\text{--}62.5 \text{ mmol } O_2 \text{ m}^{-2} \text{ d}^{-1}$ (Song et al., 2016; Zhang et al., 2017). Simulated rates higher than the observed ones only occur in July 2008 and August 2010, which is also the time of year of highest observed rates (Zhang et al., 2017). Zhang et al. (same issue) further show that simulated (daily) hypoxic areas are in agreement with observations, which range between $2,500 \text{ km}^2$ and $15,000 \text{ km}^2$ (Zhu et al., 2017). It should be noted that our values reported for the reference simulation in Table 1 exceed these values as they state the total area affected by hypoxia during each year, while observations are limited to individual months.

Table 1. Hypoxia metrics for the four different cases presented in Fig. 6. A_H : hypoxic area; $A_{H,1w}/A_{H,2w}$: areas with continuous hypoxia for one and two weeks, respectively; A_A : anoxic area (i.e. sum of areas of all grid cells with at least one day of $O_2 = 0 \text{ mmol m}^{-3}$; all in 10^3 km^2). Values for the reference simulation are total areas. Other values are changes in area (Δ) relative to the reference simulation.

Metric	2008	2009	2010	2011	2012	2013	mean \pm stdev
Reference							
A_H	23.3	20.8	31.0	16.3	18.2	6.4	19.3 ± 8.1
$A_{H,1w}$	11.2	11.5	18.7	2.8	5.2	0.1	8.2 ± 6.8
$A_{H,2w}$	4.0	5.7	11.8	0.6	0.9	0.0	3.8 ± 4.5
A_A	1.8	2.2	3.5	0.1	0.6	0.0	1.4 ± 1.4
N reduction							
ΔA_H	-11.2	-8.3	-12.4	-14.9	-13.8	-4.9	-10.9 ± 3.8
$\Delta A_{H,1w}$	-6.9	-7.4	-11.7	-2.8	-4.7	-0.1	-5.6 ± 4.0
$\Delta A_{H,2w}$	-3.0	-4.4	-8.8	-0.6	-0.9	0.0	-2.9 ± 3.3
ΔA_A	-1.3	-1.9	-3.5	-0.1	-0.6	0.0	-1.4 ± 1.3
O_2 reduction							
ΔA_H	13.6	10.7	9.3	10.6	11.4	9.3	10.8 ± 1.6
$\Delta A_{H,1w}$	10.4	6.6	7.5	6.5	9.3	2.3	7.1 ± 5.1
$\Delta A_{H,2w}$	6.8	6.8	7.0	3.0	3.1	0.0	4.5 ± 2.9
ΔA_A	0.5	0.5	2.8	0.1	0.3	0.0	0.7 ± 1.1
N and O_2 reduction							
ΔA_H	-2.3	-2.5	-3.9	-11.8	-5.8	-3.8	-5.0 ± 3.6
$\Delta A_{H,1w}$	-2.1	-2.5	-4.4	-2.8	-3.2	-0.1	-2.5 ± 1.4
$\Delta A_{H,2w}$	-1.1	-1.6	-4.1	-0.6	0.0	0.0	-1.4 ± 1.4
ΔA_A	-0.7	-1.5	-3.2	-0.1	0.0	0.0	-1.0 ± 1.2

The instantaneous benthic remineralization (Fennel et al., 2006) used in our model does not take into account sediment burial. Song et al. (2016) estimated that averaged over the ECS shelf about 45% of deposited organic matter (ca. 14% of total primary production) is buried permanently in sediments, although with high spatial variability. This partly explains the slight overestimation of SOC rates. However, we consider it having only a small effect on the relative contributions of individual
255 sources to GOC, as this limitation equally applies to all labeled N sources.

Shi et al. (2011) showed that the amount of suspended matter is significant in the coastal regions of the ECS. Sediment resuspension is not taken into account in this study. Its inclusion may result in a lower riverine contribution near the river mouths

and higher contributions in more distant areas (vice versa for oceanic contributions). Therefore, we recommend considering sediment resuspension in future studies.

260 It should further be noted that the NO_3^- nudging to a climatology in the off-shelf areas and in the Yellow Sea does not resolve interannual variability in open-ocean NO_3^- concentrations; however, we believe that these are small. Nutrient supply from the Kuroshio occurs primarily in the subsurface. With open-ocean subsurface concentrations showing significantly lower absolute values (e.g. Liu et al., 2016) and lower variability than coastal waters that are directly influenced by river inputs, neglecting interannual variations in open-ocean NO_3^- concentrations seems justifiable. Variations in volume transport of Kuroshio intrusions are likely the main cause for interannual variations in N supply from the Kuroshio and are resolved by the model. 265 Similarly, N concentrations in Taiwan Strait (e.g. Chen et al., 2004) are significantly lower than in the river inputs (by factor 10 to 100).

The diagnostic calculation of the source-specific N fluxes as the product of the fluxes for the non-labeled N tracers and the ratios of the labeled, source-specific tracers over the non-labeled ones implies that the diffusive fluxes of all labeled N tracers follow the gradient of the non-labeled (or bulk) N tracers. In reality, diffusive fluxes of matter from N tracers from different sources may be opposed. However, numerical diffusion is known to be higher than background diffusion. Therefore, 270 we consider the impact on the results to be small.

In summary, the model and N tracing applied here provide a useful basis for a meaningful analysis of our research questions.

4.2 Hypoxia under current environmental conditions

275 Previous observational studies of the hypoxic region off the Changjiang River Estuary have concluded that the regions north and south of 30°N differ with respect to the nutrient sources contributing to hypoxia formation (Chi et al., 2017; Li et al., 2002; Zhu et al., 2011). The northern region is considered to be dominated by Changjiang River inputs, while oceanic nutrient sources are thought to be important in the southern region. Here, we present the first quantification of the relative contributions of the different nutrient sources to hypoxia formation. Our results show that the riverine contribution to GOC during the hypoxia 280 season (July to November) during 2008–2013 steadily decreases from the northwest to the southeast in the region typically affected by hypoxia (see Fig. 3). The riverine contribution to GOC dominates the northern region, supporting $78.0 \pm 5.9\%$ of GOC ($73.9 \pm 5.4\%$ attributed to Changjiang River), while oceanic N supports only $22.0 \pm 5.9\%$ ($15.2 \pm 3.7\%$ from Kuroshio; see Fig. 4a and Table S2) confirming that Changjiang River nutrient loads control O_2 consumption and thus hypoxia formation in the northern region.

285 When considering July–November averages in the southern region, the relative contribution of riverine N to GOC is $61.6 \pm 5.7\%$ ($56.9 \pm 5.1\%$ from Changjiang River), while the oceanic contribution is $38.4 \pm 5.7\%$, nearly twice the oceanic contribution in the northern region, with roughly equal contributions from Kuroshio and Taiwan Strait (see Fig. 4b and Table S2). However, analysis of the seasonal cycles of the source-specific contributions to GOC reveals that hypoxia expands whenever the Changjiang contribution to GOC increases (see Fig. 5 and Table S3). This demonstrates that the Changjiang 290 River also is the major factor for hypoxia development in the southern region.

High Changjiang contributions to GOC correspond to high FW thicknesses and thus strong stratification (see Fig. S5), the other essential factor for hypoxia formation and maintenance than biological O₂ consumption. In contrast, the hypoxic area is smaller during periods with high oceanic contributions as a result of reduced GOC and weaker stratification (see Figs. 5 and S5). Nevertheless, the relatively low subsurface O₂ concentrations in the oceanic water masses precondition the southern region for hypoxia formation.

Our analyses of the seasonal cycles of GOC and stratification in relation to the large-scale meridional winds further show that the East Asian monsoon and its year-to-year variability result in seasonal and year-to-year variability in the contributions to GOC, in stratification, and thus in hypoxia in the southern region (see Fig. 5). During winter monsoon (September–April), the southward winds transport the Changjiang River plume towards the southern region, where it arrives in August/September supporting the formation of hypoxia. The opposite occurs during summer monsoon (May–August), when the northward winds transport oceanic water masses into the southern region and push the river plume northward out of the southern region.

With respect to year-to-year variability, our analysis further shows that anomalies in the meridional winds during summer and during the transition from summer to winter monsoon (August–October) significantly affect the water mass distribution, and thus GOC and hypoxia on subseasonal scales, especially south of 30°N (see Fig. 5 and Table S3). Particularly weak southward winds in September/October of 2008 resulted in a longer presence of the Changjiang River plume in the southern region, increasing stratification and GOC, which in turn caused the formation of the largest hypoxic area in that region during 2008–2013. The opposite occurred in July/August of 2013, when particularly strong northward winds pushed the Changjiang River plume north. This resulted in a drop in GOC and stratification, and in the vanishing of hypoxia, which had started forming in June through early July as anomalously weak northward winds allowed a southward transport of the Changjiang River plume. This is in agreement with Zhang et al. (2018) who simulated similar almost immediate responses of hypoxia to changes in the wind field as a result of the redistribution of Changjiang FW causing changes in stratification. Our study expands on their work by additionally demonstrating the importance of the redistribution of Changjiang nutrients for GOC.

These short-term changes in GOC, stratification and hypoxic area in response to variations in the large-scale winds illustrate the complexity of the region with respect to atmosphere–ocean interaction and their effect on hypoxia. They further highlight the necessity of a high spatio-temporal resolution of both model approaches and observations to understand the causes of hypoxia formation and maintenance in the ECS under current environmental conditions.

In general, our results are in agreement with Zhang et al. (2019) who found that riverine N supports 56% of water column integrated PP (8-year average) in the ECS regions shallower than 50 m, which are most susceptible to hypoxia (see Fig. 3). This confirms the high contribution of Changjiang N to GOC found for our analysis regions. Similarly, the comparably low oceanic contributions, especially in the northern region, are in line with their results. Zhang et al. (2019) also found a relatively high contribution (22%) of atmospheric N to PP suggesting that atmospheric N deposition should be considered in future biogeochemical modeling studies of the ECS.

4.3 Mitigation of hypoxia in the present and future

Our analysis of the changes in hypoxic area and hypoxic exposure under reduced Changjiang River N loads (see Fig. 6 and Table 1) suggests a high potential of riverine N load reductions to mitigate hypoxia. The simulated average reduction in hypoxic area by 56.5% is almost proportional to the N load reduction of 50% but varies significantly from year to year (39.9% in 2009 to 91.4% in 2011). Hypoxia vanishes almost entirely during months of small hypoxic areas (May to August 2008, September/October 2013; see Fig. S7) and maximum hypoxic areas during the seasonal cycle are significantly reduced in response to decreased GOC supported by Changjiang River N inputs. However, comprehensive further studies are needed to assess the effect of Changjiang River nutrient reductions on hypoxia. In this context, it is important to consider all processes relevant for hypoxia formation. Zhou et al. (2017b) conducted a scenario with 50% lower Changjiang River nutrient loads and found a reduction in hypoxic area by only about 20%. However, they did not consider SOC, a major sink of O_2 in the ECS (Zhang et al., 2017), which may explain this discrepancy.

Analysis of the scenario where open-ocean O_2 concentrations were 20% lower than at present shows that the reduced lateral supply due to declining open-ocean O_2 concentrations (Bopp et al., 2017) may significantly exacerbate hypoxia in the ECS, indicated by the expansion of both hypoxic and anoxic areas by about 50% (see Fig. 6 and Table 1). However, we further found that a 50% reduction of the Changjiang River N loads more than compensates for this increase in hypoxia.

The 20% reduction of open-ocean O_2 concentrations that was applied here corresponds to changes in subsurface O_2 of about 30–40 $mmol\ O_2\ m^{-3}$ projected by earth system models (Bopp et al., 2017), but does not consider other climate change effects such as local changes in water temperature or stratification. For instance, an increase in temperature would reduce O_2 solubility, which would worsen O_2 conditions further as shown, e.g. for the NGoM (Laurent et al., 2018). Therefore, more comprehensive studies on climate change impacts on hypoxia in the ECS are required.

4.4 Comparison with the northern Gulf of Mexico

In the following, we compare the hypoxic zones of the ECS and the NGoM with respect to their sizes and timing, based on observations, and the main sources of O_2 consumption, based on the results of this study and an analogous nutrient tracing study for the NGoM (Große et al., 2019). The key messages of this comparison are summarized in Table 2.

The spatial extent of observed hypoxia in the ECS is similar to that in the NGoM, affecting areas on the order of 15,000 km^2 . In the NGoM, hypoxia is frequently observed from May to September when seasonal stratification is strongest due to high riverine freshwater inputs. In the ECS, the hypoxic season starts about one month later in June, with hypoxia forming directly off the Changjiang River Estuary. Hypoxia can be observed until October/November, though the hypoxic zone moves southward over summer.

In the NGoM, riverine nutrients from the Mississippi and Atchafalaya Rivers dominate O_2 consumption. In the ECS, only in the region north of 30°N is O_2 consumption dominated by riverine nutrient inputs that almost exclusively originate from the Changjiang River. South of 30°N, the oceanic contribution to O_2 consumption is comparable to that of riverine nutrients. However, driven by year-to-year variability in the East Asian monsoon, the Changjiang River appears to be the main control

Table 2. Comparison of hypoxia in the northern Gulf of Mexico (NGoM) and the East China Sea (ECS).

Characteristic	NGoM	ECS
Hypoxic area (km ²)	15,000 ¹	15,000 ²
Hypoxic season	May–September ¹	June–November ³
GOC controls (seasonal average)	riverine ≫ oceanic	>30 °N: riverine ≫ oceanic <30 °N: riverine ≈ oceanic
Main riverine source(s)	Mississippi & Atchafalaya	Changjiang
Controls of year-to-year variability	Mississippi & wind field	Changjiang & wind field

¹ Rabalais et al. (2002); ² Li et al. (2002), Zhu et al. (2011, 2017); ³ Wang et al. (2012)

of hypoxia formation also in the southern region. Similarly, the Mississippi River in concert with the local wind field seems to be the main control of year-to-year variability in hypoxic area in the NGoM. There, episodic upwelling-favorable winds can result in an offshore transport of Mississippi FW and nutrients leading to a reduction in stratification and O₂ consumption and thus hypoxia (Feng et al., 2014).

360 **5 Conclusions**

To our knowledge, this is the first study quantifying the relative contributions of individual riverine and oceanic nutrient sources on hypoxia formation in ECS using active element tracing. Therefore, it constitutes an important milestone towards the quantification of the contributions of riverine vs. oceanic nutrient sources to hypoxia formation in the ECS.

Our results suggest that N from the Changjiang River is the dominant driver of O₂ consumption ($73.9 \pm 5.4\%$) north of 30 °N
365 under present-day environmental conditions (2008–2013). Contrary to observational insights and despite high contributions of N from Kuroshio and Taiwan Strait to O₂ consumption on seasonal time scales ($19.5 \pm 2.2\%$ and $18.9 \pm 3.2\%$, respectively), the Changjiang River waters are also the main factor for hypoxia formation south of 30 °N.

Our analysis highlights the importance of considering subseasonal time scales for understanding the controls of hypoxia formation and its year-to-year variability in this region. The East Asian monsoon and its associated change in large-scale wind
370 patterns control water mass distribution and thus O₂ consumption and stratification. Year-to-year variability in the intensity of the winds can lead to significant differences in the amount of Changjiang River water in the southern region explaining year-to-year variability in hypoxia.

Reductions in the Changjiang River nutrient loads appear to have a high potential for mitigating hypoxia and to counteract the likely future decline of open-ocean O₂ supplied to the region. However, more comprehensive studies of both the effect of
375 riverine nutrient load reductions and climate change effects are required.

Code and data availability. A modified version of the ROMS source code that facilitates writing of model diagnostics needed by the element tracing software (ETRAC) is available at https://github.com/FabianGrosse/ROMS_3.7_for_ETRAC.

ETRAC is available via <https://github.com/FabianGrosse/ETRAC>. Due to the large data size, ROMS and ETRAC output is freely available upon request from the corresponding author (fabian.grosse@dal.ca).

380 *Author contributions.* FG and KF conceived the study. FG set up the element tracing and conducted the analyses. HZ set up the model. AL helped with validation. FG wrote the manuscript with input from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. All figures in this article were created with MATLAB, benefiting significantly from the toolboxes for TEOS-10 (Mc-Dougall and Barker, 2011), the *cmocean* color schemes (Thyng et al., 2016) and *m_map* (<https://www.eoas.ubc.ca/~rich/map.html>). The
385 authors of these toolboxes are thankfully acknowledged. The geographical information used for creation of Fig. 1 were obtained from the Global Self-consistent, Hierarchical, High-resolution Geography Database (GSHHG; <https://www.soest.hawaii.edu/wessel/gshhg/>). We thank Compute Canada for providing computing resources under the resource allocation project qqh-593-ac. KF acknowledges funding from the Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery program. Financial support to HZ from the China Scholarship Council (CSC) is gratefully acknowledged. We thank Hagen Radtke and one anonymous reviewer for their constructive criticism,
390 which helped to improve this manuscript.

References

- Bian, C., Jiang, W., and Greatbatch, R. J.: An exploratory model study of sediment transport sources and deposits in the Bohai Sea, Yellow Sea, and East China Sea, *J. Geophys. Res.-Oceans*, 118, 5908–5923, <https://doi.org/10.1002/2013JC009116>, 2013.
- Bopp, L., Resplandy, L., Untersee, A., Le Mezo, P., and Kageyama, M.: Ocean (de)oxygenation from the Last Glacial Maximum to the
395 twenty-first century: insights from Earth System models, *Philos. T. Roy. Soc. A*, 375, <https://doi.org/10.1098/rsta.2016.0323>, 2017.
- Carton, J. A. and Giese, B. S.: A Reanalysis of Ocean Climate Using Simple Ocean Data Assimilation (SODA), *Mon. Weather Rev.*, 136, 2999–3017, <https://doi.org/10.1175/2007MWR1978.1>, 2008.
- Chen, C.-T. A., Hsing, L.-Y., Liu, C.-L., and Wang, S.-L.: Degree of nutrient consumption of upwelled water in the Taiwan Strait based on dissolved organic phosphorus or nitrogen, *Mar. Chem.*, 87, 73–86, <https://doi.org/10.1016/j.marchem.2004.01.006>, 2004.
- 400 Chi, L., Song, X., Yuan, Y., Wang, W., Zhou, P., Fan, X., Cao, X., and Yu, Z.: Distribution and key influential factors of dissolved oxygen off the Changjiang River Estuary (CRE) and its adjacent waters in China, *Mar. Pollut. Bull.*, 125, 440–450, <https://doi.org/10.1016/j.marpolbul.2017.09.063>, 2017.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L.,
405 Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Egbert, G. D. and Erofeeva, S. Y.: Efficient Inverse Modeling of Barotropic Ocean Tides, *J. Atmos. Ocean. Tech.*, 19, 183–204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002.
- 410 Fan, W. and Song, J.: A numerical study of the seasonal variations of nutrients in the Changjiang River estuary and its adjacent sea area, *Ecol. Model.*, 291, 69–81, <https://doi.org/10.1016/j.ecolmodel.2014.07.026>, 2014.
- Feng, Y., Fennel, K., Jackson, G. A., DiMarco, S. F., and Hetland, R. D.: A model study of the response of hypoxia to upwelling-favorable wind on the northern Gulf of Mexico shelf, *J. Marine Syst.*, 131, 63–73, <https://doi.org/10.1016/j.jmarsys.2013.11.009>, 2014.
- Fennel, K. and Testa, J. M.: Biogeochemical Controls on Coastal Hypoxia, *Annu. Rev. Mar. Sci.*, 11, 105–130,
415 <https://doi.org/10.1146/annurev-marine-010318-095138>, 2019.
- Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., and Haidvogel, D.: Nitrogen cycling in the Middle Atlantic Bight: Results from a three-dimensional model and implications for the North Atlantic nitrogen budget, *Global Biogeochem. Cy.*, 20, 1–14, <https://doi.org/10.1029/2005GB002456>, 2006.
- Fennel, K., Hetland, R., Feng, Y., and DiMarco, S.: A coupled physical-biological model of the Northern Gulf of Mexico shelf: model
420 description, validation and analysis of phytoplankton variability, *Biogeosciences*, 8, 1881–1899, <https://doi.org/10.5194/bg-8-1881-2011>, 2011.
- Fennel, K., Hu, J., Laurent, A., Marta-Almeida, M., and Hetland, R.: Sensitivity of hypoxia predictions for the northern Gulf of Mexico to sediment oxygen consumption and model nesting, *J. Geophys. Res.-Oceans*, 118, 990–1002, <https://doi.org/10.1002/jgrc.20077>, 2013.
- Gao, L., Li, D., Ishizaka, J., Zhang, Y., Zong, H., and Guo, L.: Nutrient dynamics across the river-sea interface in the Changjiang (Yangtze
425 River) estuary—East China Sea region, *Limnol. Oceanogr.*, 60, 2207–2221, <https://doi.org/10.1002/lno.10196>, 2015.

- Garcia, H. E., Boyer, T. P., Locarnini, R. A., Antonov, J. I., Mishonov, A. V., Baranova, O. K., Zweng, M. M., Reagan, J. R., Johnson, D. R., and Levitus, S.: World Ocean Atlas 2013. Volume 3: Dissolved oxygen, apparent oxygen utilization, and oxygen saturation, Tech. rep., National Oceanic and Atmospheric Administration (NOAA), Silver Spring, MD, <https://doi.org/10.7289/V5XG9P2W>, 2013a.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., Johnson, D. R., Mishonov, A. V., and Levitus, S.: World Ocean Atlas 2013. Volume 4: Dissolved inorganic nutrients (phosphate, nitrate, silicate), Tech. rep., National Oceanic and Atmospheric Administration (NOAA), Silver Spring, MD, <https://doi.org/10.7289/V5J67DWD>, 2013b.
- Große, F., Kreus, M., Lenhart, H.-J., Pätsch, J., and Pohlmann, T.: A Novel Modeling Approach to Quantify the Influence of Nitrogen Inputs on the Oxygen Dynamics of the North Sea, *Front. Mar. Sci.*, 4, 383, <https://doi.org/10.3389/fmars.2017.00383>, 2017.
- Große, F., Fennel, K., and Laurent, A.: Quantifying the Relative Importance of Riverine and Open-Ocean Nitrogen Sources for Hypoxia Formation in the Northern Gulf of Mexico, *J. Geophys. Res.-Oceans*, <https://doi.org/10.1029/2019JC015230>, 2019.
- Guo, X., Miyazawa, Y., and Yamagata, T.: The Kuroshio onshore intrusion along the shelf break of the East China Sea: The origin of the Tsushima Warm Current, *J. Phys. Oceanogr.*, 36, 2205–2231, <https://doi.org/10.1175/JPO2976.1>, 2006.
- Haidvogel, D. B., Arango, H., Budgell, W. P., Cornuelle, B. D., Curchitser, E., Di Lorenzo, E., Fennel, K., Geyer, W. R., Hermann, A. J., Lanerolle, L., Levin, J., McWilliams, J. C., Miller, A. J., Moore, A. M., Powell, T. M., Shchepetkin, A. F., Sherwood, C. R., Signell, R. P., Warner, J. C., and Wilkin, J.: Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System, *J. Comput. Phys.*, 227, 3595–3624, <https://doi.org/10.1016/j.jcp.2007.06.016>, 2008.
- Hetland, R. D. and Zhang, X.: Interannual Variation in Stratification over the Texas–Louisiana Continental Shelf and Effects on Seasonal Hypoxia, in: *Modeling Coastal Hypoxia*, edited by Dubravko, J., Rose, K. A., Hetland, R. D., and Fennel, K., pp. 49–60, Springer, https://doi.org/10.1007/978-3-319-54571-4_3, 2017.
- Laurent, A., Fennel, K., Hu, J., and Hetland, R.: Simulating the effects of phosphorus limitation in the Mississippi and Atchafalaya River plumes, *Biogeosciences*, 9, 4707–4723, <https://doi.org/10.5194/bg-9-4707-2012>, 2012.
- Laurent, A., Fennel, K., Cai, W.-J., Huang, W.-J., Barbero, L., and Wanninkhof, R.: Eutrophication-induced acidification of coastal waters in the northern Gulf of Mexico: Insights into origin and processes from a coupled physical-biogeochemical model, *Geophys. Res. Lett.*, 44, 946–956, <https://doi.org/10.1002/2016GL071881>, 2017.
- Laurent, A., Fennel, K., Ko, D. S., and Lehrter, J.: Climate Change Projected to Exacerbate Impacts of Coastal Eutrophication in the Northern Gulf of Mexico, *J. Geophys. Res.-Oceans*, 123, 3408–3426, <https://doi.org/10.1002/2017JC013583>, 2018.
- Li, D., Zhang, J., Huang, D., Wu, Y., and Liang, J.: Oxygen depletion off the Changjiang (Yangtze River) estuary, *Sci. China Ser. D*, 45, 1137–1146, <https://doi.org/10.1360/02yd9110>, 2002.
- Li, X., Bianchi, T., Yang, Z., Osterman, L. E., Allison, M. E., DiMarco, S. F., and Yang, H.: Historical trends of hypoxia in Changjiang River estuary: Applications of chemical biomarkers and microfossils, *J. Marine Syst.*, 86, 57–68, <https://doi.org/10.1016/j.jmarsys.2011.02.003>, 2011.
- Liu, S. M., Zhang, J., Chen, H. T., Wu, Y., Xiong, H., and Zhang, Z. F.: Nutrients in the Changjiang and its tributaries, *Biogeochemistry*, 62, 1–18, <https://doi.org/10.1023/A:1021162214304>, 2003.
- Liu, S. M., Hong, G.-H., Zhang, J., Ye, X. W., and Jiang, X. L.: Nutrient budgets for large Chinese estuaries, *Biogeosciences*, 6, 2245–2263, <https://doi.org/10.5194/bg-6-2245-2009>, 2009.
- Liu, S. M., Qi, X. H., Li, X., Ye, H. R., Wu, Y., Ren, J. L., Zhang, J., and Xu, W. Y.: Nutrient dynamics from the Changjiang (Yangtze River) estuary to the East China Sea, *J. Marine Syst.*, 154, 15–27, <https://doi.org/10.1016/j.jmarsys.2015.05.010>, 2016.

Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., Seidov, D., and Levitus, S.: World Ocean Atlas 2013. Volume 1: Temperature, Tech. rep., National Oceanic and Atmospheric Administration (NOAA), Silver Spring, MD, <https://doi.org/10.7289/V55X26VD>, 2013.

McDougall, T. J. and Barker, P. M.: Getting started with TEOS-10 and the Gibbs Seawater (GSW) oceanographic toolbox, Tech. rep., SCOR/IAPSO, http://www.teos-10.org/pubs/gsw/v3_04/pdf/Getting_Started.pdf, 2011.

Ménesguen, A., Cugier, P., and Leblond, I.: A new numerical technique for tracking chemical species in a multi-source, coastal ecosystem, applied to nitrogen causing *Ulva* blooms in the Bay of Brest (France), *Limnol. Oceanogr.*, 51, 591–601, https://doi.org/10.4319/lo.2006.51.1_part_2.0591, 2006.

Qian, W., Dai, M., Xu, M., Kao, S.-J., Du, C., Liu, J., Wang, H., Guo, L., and Wang, L.: Non-local drivers of the summer hypoxia in the East China Sea off the Changjiang Estuary, *Estuar. Coast. Shelf S.*, 198, 393–399, <https://doi.org/10.1016/j.ecss.2016.08.032>, 2017.

Rabalais, N. N., Turner, R. E., and Wiseman Jr., W. J.: Gulf of Mexico Hypoxia, A.K.A. “The Dead Zone”, *Annu. Rev. Ecol. Syst.*, 33, 235–263, <https://doi.org/10.1146/annurev.ecolsys.33.010802.150513>, 2002.

Radtke, H., Neumann, T., Voss, M., and Fennel, W.: Modeling pathways of riverine nitrogen and phosphorus in the Baltic Sea, *Journal of Geophysical Research: Oceans*, 117, 1–15, <https://doi.org/10.1029/2012JC008119>, 2012.

Rutherford, K. and Fennel, K.: Diagnosing transit times on the northwestern North Atlantic continental shelf, *Ocean Sci.*, 14, 1207–1221, <https://doi.org/10.5194/os-14-1207-2018>, 2018.

Schmidtko, S., Stramma, L., and Visbeck, M.: Decline in global oceanic oxygen content during the past five decades, *Nature*, 542, 335, <https://doi.org/10.1038/nature21399>, 2017.

Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., and Bouwman, A. F.: Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application, *Global Biogeochem. Cy.*, 19, <https://doi.org/10.1029/2005GB002606>, 2005.

Shi, W., Wang, M., and Jiang, L.: Spring-neap tidal effects on satellite ocean color observations in the Bohai Sea, Yellow Sea, and East China Sea, *J. Geophys. Res.-Oceans*, 116, 1–13, <https://doi.org/10.1029/2011JC007234>, 2011.

Simpson, J. H.: The shelf-sea fronts: implications of their existence and behaviour, *Philos. T. Roy. Soc. A*, 302, 531–546, <https://doi.org/10.1098/rsta.1981.0181>, 1981.

Siswanto, E., Nakata, H., Matsuoka, Y., Tanaka, K., Kiyomoto, Y., Okamura, K., Zhu, J., and Ishizaka, J.: The long-term freshening and nutrient increases in summer surface water in the northern East China Sea in relation to Changjiang discharge variation, *J. Geophys. Res.-Oceans*, 113, <https://doi.org/10.1029/2008JC004812>, 2008.

Song, G., Liu, S., Zhu, Z., Zhai, W., Zhu, C., and Zhang, J.: Sediment oxygen consumption and benthic organic carbon mineralization on the continental shelves of the East China Sea and the Yellow Sea, *Deep Sea Res. Pt. II*, 124, 53–63, <https://doi.org/10.1016/j.dsr2.2015.04.012>, 2016.

Thyng, K. M., Greene, C. A., Hetland, R. D., Zimmerle, H. M., and DiMarco, S. F.: True Colors of Oceanography: Guidelines for Effective and Accurate Colormap Selection, *Oceanography*, 29, <https://doi.org/10.5670/oceanog.2016.66>, 2016.

Tong, Y., Zhao, Y., Zhen, G., Chi, J., Liu, X., Lu, Y., Wang, X., Yao, R., Chen, J., and Zhang, W.: Nutrient loads flowing into coastal waters from the main rivers of China (2006–2012), *Sci. Rep.*, 5, 16 678, <https://doi.org/10.1038/srep16678>, 2015.

Wang, B.: Hydromorphological mechanisms leading to hypoxia off the Changjiang estuary, *Mar. Environ. Res.*, 67, 53–58, <https://doi.org/10.1016/j.marenvres.2008.11.001>, 2009.

- 500 Wang, B., Wei, Q., Chen, J., and Xie, L.: Annual cycle of hypoxia off the Changjiang (Yangtze River) Estuary, *Mar. Environ. Res.*, 77, 1–5, <https://doi.org/10.1016/j.marenvres.2011.12.007>, 2012.
- Wang, H., Yang, Z., Wang, Y., Saito, Y., and Liu, J. P.: Reconstruction of sediment flux from the Changjiang (Yangtze River) to the sea since the 1860s, *J. Hydrol.*, 349, 318–332, <https://doi.org/10.1016/j.jhydrol.2007.11.005>, 2008.
- Wang, J., Yan, W., Chen, N., Li, X., and Liu, L.: Modeled long-term changes of DIN:DIP ratio in the Changjiang River in relation to Chl- α and DO concentrations in adjacent estuary, *Estuar. Coast. Shelf S.*, 166, 153–160, <https://doi.org/10.1016/j.ecss.2014.11.028>, 2015.
- 505 Wang, W., Yu, Z., Song, X., Yuan, Y., Wu, Z., Zhou, P., and Cao, X.: Intrusion Pattern of the Offshore Kuroshio Branch Current and Its Effects on Nutrient Contributions in the East China Sea, *J. Geophys. Res.-Oceans*, 123, 2116–2128, <https://doi.org/10.1002/2017JC013538>, 2018.
- Wu, G., Cao, W., Wang, F., Su, X., Yan, Y., and Guan, Q.: Riverine nutrient fluxes and environmental effects on China's estuaries, *Sci. Total Environ.*, 661, 130–137, <https://doi.org/10.1016/j.scitotenv.2019.01.120>, 2019.
- 510 Yan, W., Zhang, S., Sun, P., and Seitzinger, S. P.: How do nitrogen inputs to the Changjiang basin impact the Changjiang River nitrate: A temporal analysis for 1968–1997, *Global Biogeochem. Cy.*, 17, 1091, <https://doi.org/10.1029/2002GB002029>, 2003.
- Yang, D., Yin, B., Liu, Z., and Feng, X.: Numerical study of the ocean circulation on the East China Sea shelf and a Kuroshio bottom branch northeast of Taiwan in summer, *J. Geophys. Res.-Oceans*, 116, 1–20, <https://doi.org/10.1029/2010JC006777>, 2011.
- Yang, D., Yin, B., Liu, Z., Bai, T., Qi, J., and Chen, H.: Numerical study on the pattern and origins of Kuroshio branches in the bottom water of southern East China Sea in summer, *J. Geophys. Res.-Oceans*, 117, 1–16, <https://doi.org/10.1029/2011JC007528>, 2012.
- 515 Yu, L., Fennel, K., Laurent, A., Murrell, M. C., and Lehrter, J. C.: Numerical analysis of the primary processes controlling oxygen dynamics on the Louisiana shelf, *Biogeosciences*, 12, 2063–2076, <https://doi.org/10.5194/bg-12-2063-2015>, 2015.
- Zhang, H., Zhao, L., Sun, Y., Wang, J., and Wei, H.: Contribution of sediment oxygen demand to hypoxia development off the Changjiang Estuary, *Estuar. Coast. Shelf S.*, 192, 149–157, <https://doi.org/10.1016/j.ecss.2017.05.006>, 2017.
- 520 Zhang, H., Fennel, K., Laurent, A., and Bian, C.: A numerical model study of the main factors contributing to hypoxia and its sub-seasonal to interannual variability off the Changjiang Estuary, *Biogeosciences Discussions*, same issue.
- Zhang, J.: Nutrient elements in large Chinese estuaries, *Cont. Shelf Res.*, 16, 1023–1045, [https://doi.org/10.1016/0278-4343\(95\)00055-0](https://doi.org/10.1016/0278-4343(95)00055-0), 1996.
- Zhang, J., Guo, X., and Zhao, L.: Tracing external sources of nutrients in the East China Sea and evaluating their contributions to primary production, *Prog. Oceanogr.*, 176, 102 122, <https://doi.org/10.1016/j.pocean.2019.102122>, <http://www.sciencedirect.com/science/article/pii/S0079661118301046>, 2019.
- 525 Zhang, W., Wu, H., and Zhu, Z.: Transient Hypoxia Extent Off Changjiang River Estuary due to Mobile Changjiang River Plume, *J. Geophys. Res.-Oceans*, 123, 9196–9211, <https://doi.org/10.1029/2018JC014596>, 2018.
- Zhang, W. G., Wilkin, J. L., and Schofield, O. M. E.: Simulation of Water Age and Residence Time in New York Bight, *J. Phys. Oceanogr.*, 40, 965–982, <https://doi.org/10.1175/2009JPO4249.1>, 2010.
- 530 Zhang, X., Hetland, R. D., Marta-Almeida, M., and DiMarco, S. F.: A numerical investigation of the Mississippi and Atchafalaya freshwater transport, filling and flushing times on the Texas-Louisiana Shelf, *J. Geophys. Res.-Oceans*, 117, C11 009, <https://doi.org/10.1029/2012JC008108>, 2012.
- Zhao, L. and Guo, X.: Influence of cross-shelf water transport on nutrients and phytoplankton in the East China Sea: a model study, *Ocean Sci.*, 7, 27–43, <https://doi.org/10.5194/os-7-27-2011>, 2011.
- 535 Zheng, J., Gao, S., Liu, G., Wang, H., and Zhu, X.: Modeling the impact of river discharge and wind on the hypoxia off Yangtze Estuary, *Nat. Hazard. Earth Sys.*, 16, 2559–2576, <https://doi.org/10.5194/nhess-16-2559-2016>, 2016.

- Zhou, F., Chai, F., Huang, D., Xue, H., Chen, J., Xiu, P., Xuan, J., Li, J., Zeng, D., Ni, X., and Wang, K.: Investigation of hypoxia off the Changjiang Estuary using a coupled model of ROMS-CoSiNE, *Prog. Oceanogr.*, 159, 237–254, <https://doi.org/10.1016/j.pocean.2017.10.008>, 2017a.
- 540 Zhou, P., Song, X., Yuan, Y., Wang, W., Cao, X., and Yu, Z.: Intrusion pattern of the Kuroshio Subsurface Water onto the East China Sea continental shelf traced by dissolved inorganic iodine species during the spring and autumn of 2014, *Mar. Chem.*, 196, 24–34, <https://doi.org/10.1016/j.marchem.2017.07.006>, 2017b.
- Zhou, P., Song, X., Yuan, Y., Wang, W., Chi, L., Cao, X., and Yu, Z.: Intrusion of the Kuroshio Subsurface Water in the southern East China Sea and its variation in 2014 and 2015 traced by dissolved inorganic iodine species, *Prog. Oceanogr.*, 165, 287–298, <https://doi.org/10.1016/j.pocean.2018.06.011>, 2018.
- 545 Zhu, Z.-Y., Zhang, J., Wu, Y., Zhang, Y.-Y., Lin, J., and Liu, S.-M.: Hypoxia off the Changjiang (Yangtze River) Estuary: oxygen depletion and organic matter decomposition, *Mar. Chem.*, 125, 108–116, <https://doi.org/10.1016/j.marchem.2011.03.005>, 2011.
- Zhu, Z.-Y., Wu, H., Liu, S.-M., Wu, Y., Huang, D.-J., Zhang, J., and Zhang, G.-S.: Hypoxia off the Changjiang (Yangtze River) estuary and in the adjacent East China Sea: Quantitative approaches to estimating the tidal impact and nutrient regeneration, *Mar. Pollut. Bull.*, 125, 103–114, <https://doi.org/10.1016/j.marpolbul.2017.07.029>, 2017.
- 550 Zweng, M. M., Reagan, J. R., Antonov, J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P., Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., Biddle, M. M., and Levitus, S.: *World Ocean Atlas 2013. Volume 2: Salinity*, Tech. rep., National Oceanic and Atmospheric Administration (NOAA), Silver Spring, MD, <https://doi.org/10.7289/V5251G4D>, 2013.