

In this response letter the *referee comments are listed in blue italic font*, followed by the authors response.

Dear colleagues,

The efforts made to introduce more nuance to their statements regarding uncertainties of flux calculations from different methods used here is appreciable. However, I think some more work should be done.

For instance, suggestions made by Michael Staubwasser to show better pore water profiles by zooming around the interface and try to fit more than two points are crucial here. The arguments advanced to stick on 2 points modelling are, in my opinion, not really convincing.

argument 1. I am not sure that saying "many others did, we should do" is very scientific.

In addition, looking back to papers from Sundby and collaborators (Sundby et al, 1986 and Westerlund et al 1986), it seems that they do not calculate fluxes out of porewaters. In the second one, they compare overlying water and the first layer of porewater to predict flux direction, they do not go further.

argument 2

the authors say that "advanced curve fitting methods would fail to capture sharp gradients..." in order to justify why they do a simple 2 points fitting to calculate diffusive flux. I agree that there is no problem using linear regressions with pour-resolved vertical profiles. The problem here is that between a 2-points and a 3-points regression, there is a big gap in terms of modelling validity.

argument 3

Indeed, if bottom water concentration is low so it does not affect the gradient, however the first porewater concentration does. The latter concentration is the result of a certain volume of porewater mixed altogether during sampling and extraction. We can imagine that some of it had oxygen, some of it sulphide and the final concentration is the result of thermodynamic equilibrium, not only due to mixing porewaters of different concentrations.

This is better said by Warnken et al., 2001 as follows:

"The diffusive fluxes of Fe agreed with measured fluxes at station 1, but diffusive flux estimates of Fe at stations 2 and 3 were greater than the measured chamber fluxes, possibly indicating that pore water Fe concentration gradients did not accurately represent actual interfacial concentration gradients at these stations. This could be due to less steep gradients in the upper mm of sediment suggesting that the 1 cm sampling resolution used here was not adequate. Accounting for the reduced slope very near

the sediment–water interface would result in lower calculated diffusive fluxes that would possibly agree more closely with measured fluxes. Additionally, rapid oxidative processes at or near the sediment–water interface could also reduce the amount of Fe released from the sediments, as concentration increases during the incubation experiments conducted at stations 2 and 3 were reduced when compared to station 1."

Having a third point, helps to show that the trend is maintained despite interface processes.

I would suggest the authors to simply explain that linear regressions were made because of scarcity of data due to sharpness of gradients at a centimetre resolution profiling. In addition, I would put a warning about the precision of numbers obtained.

We would like to thank the reviewer for providing us with additional referee comments and we appreciate the reviewer's remaining concerns regarding methodology.

He rightly pointed out that a three-point regression would be preferable for the calculation of benthic diffusive fluxes because of a lower statistical uncertainty. He also acknowledged problems related to the coarse vertical resolution of our pore water profiles and the sharpness of the concentration gradients within the surface sediment.

Due to the soft texture of the sediment, it was not possible to sample the pore water at a higher resolution within the first centimeter of the sediment. Furthermore, as stated in the first response to the reviewer, Fe concentrations in pore water drop sharply shortly below the sediment surface because of high rates of bacterial sulfate reduction and Fe sulfide precipitation within the surface sediment. Consequently, choosing an additional concentration point, that would be located below this drop in concentration, would simply lead to artificially low fluxes. Although, the uncertainty of a three-point approach would be lower from a statistical point of view, it would clearly lead to erroneous results. To illustrate this and following the reviewers' recommendation, we added close-up profiles of pore water concentrations within the upper 5 cm of the surface sediment to the supplement (Fig. S1 and S2). Furthermore, we added a more detailed explanation, including the reviewers' suggestions, to state why we applied the two-point gradient in lines 323 – 330 in the revised manuscript "Due to the coarse resolution of our pore water profiles and the steep gradients between the uppermost pore water and bottom water sample (see close-up profiles, Fig. S1 and S2, in the supplement), we chose to follow previous studies (e.g., Noffke et al., 2012; Lenstra et al., 2019; Scholz et al., 2019) and calculate diffusive benthic fluxes based on a two point concentration gradient. Including deeper samples into a linear regression or applying more advanced curve fitting methods would reduce the statistical uncertainty but fail to capture the sharp concentration gradients at the sediment surface, thus, leading to erroneous flux estimates (cf. Shibamoto and Harada, 2010). " We hope that this will clarify, why we have to choose the applied method. A more detailed discussion of the early diagenetic processes leading to steep and variable gradients within the surface sediment (bacterial sulfate reduction, precipitation of Fe monosulfides, etc.) would be inappropriate in the method section.

Another point that was not fully addressed is the uncertainties on incubation calculations. You must have errors over 100 % with a R² of 0.5. You should look in detail what are uncertainties and which regression has statistical significance.

Following the reviewer's recommendation, we calculated the uncertainty of our in-situ benthic fluxes by propagating the errors of the linear regressions of the concentration change over time (see Tables 2 and 3 in the revised version). Furthermore, following previous studies (e.g., Friedrich et al. (2002); Lenstra et al. (2019)), only fluxes with a linear regressions with a $R^2 > 0.3$ are reported in Tables 2 and 3 (See lines 338 - 339 in the revised manuscript).

All this should lead to a discussion about how we can do better. As it is, it gives me the idea that since 1986 no analytical progress was made to improve spatial resolution and to limit incubation time and increase incubation replicates. Maybe that is the case, but it has to be said somewhere

We agree that the advantages and disadvantages of the various methods and approaches for the determination of benthic fluxes is an exciting topic. However, the focus of our study is the benthic biogeochemical cycling of Fe and Cd in oxygen minimum zones and not the technical aspects of the methods applied. Therefore, we feel that providing an extended discussion about technical limitations and potential improvements to the methods is beyond the scope of our work.

All the methods and approaches applied in our study are commonly applied in benthic biogeochemical studies by various research groups around the world. We agree that improvements to these methods are possible. On the other hand, for a comparison of benthic fluxes from different locations or seasons (note that we compare our Fe flux data to those determined in the same area a couple of years earlier), it is quite useful to stick to methodologies that have been applied before. For this reason, we refer to previous studies in the description of our methodology.

1 **The control of hydrogen sulfide on benthic iron and**
2 **cadmium fluxes in the oxygen minimum zone off Peru**

3

4 Anna Plass^{1*}, Christian Schlosser¹, Stefan Sommer¹, Andrew W. Dale¹, Eric P.
5 Achterberg¹, Florian Scholz^{1*}

6 ¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Wischhofstraße 1-3, 24148
7 Kiel, Germany

8 *Correspondence to: Anna Plass (aplass@geomar.de), Florian Scholz
9 (fscholz@geomar.de)

10

11 **Abstract**

12 Sediments in oxygen-depleted marine environments can be an important sink or
13 source of bio-essential trace metals in the ocean. However, the key mechanisms
14 controlling the release from or burial of trace metals in sediments are not exactly
15 understood. Here, we investigate the benthic biogeochemical cycling of Fe and Cd in
16 the oxygen minimum zone off Peru. We combine bottom water and pore water
17 concentrations, as well as benthic fluxes determined from pore water profiles and in-
18 situ from benthic chamber incubations, along a depth transect at 12° S. In agreement
19 with previous studies, both concentration-depth profiles and in-situ benthic fluxes
20 indicate a release of Fe from sediments to the bottom water. Diffusive Fe fluxes and
21 Fe fluxes from benthic chamber incubations ($-0.3 - -17.5 \text{ mmol m}^{-2} \text{ y}^{-1}$) are broadly
22 consistent at most stations, indicating that diffusion is the main transport mechanism
23 of dissolved Fe across the sediment-water interface. The occurrence of mats of sulfur
24 oxidizing bacteria on the seafloor represents an important control on the spatial
25 distribution of Fe fluxes by regulating hydrogen sulfide (H_2S) concentrations and,
26 potentially, Fe sulfide precipitation within the surface sediment. Rapid removal of
27 dissolved Fe after its release to anoxic bottom waters hints to oxidative removal by
28 nitrite and interactions with particles in the near-bottom water column. Benthic flux
29 estimates of Cd suggest a flux into the sediment within the oxygen minimum zone.
30 Fluxes from benthic chamber incubations (up to $22.6 \text{ } \mu\text{mol m}^{-2} \text{ y}^{-1}$) exceed diffusive
31 fluxes ($< 1 \text{ } \mu\text{mol m}^{-2} \text{ y}^{-1}$) by a factor > 25 , indicating that downward diffusion of Cd
32 across the sediment-water interface is of subordinate importance for Cd removal from
33 benthic chambers. As Cd removal in benthic chambers co-varies with H_2S
34 concentrations in the pore water of surface sediments, we argue that Cd removal is
35 mediated by precipitation of CdS within the chamber water or directly at the sediment-
36 water interface. A mass balance approach, taking into account the contributions of
37 diffusive and chamber fluxes as well as Cd delivery with organic material, suggests
38 that CdS precipitation in the near-bottom water could make an important contribution
39 to the overall Cd mass accumulation in the sediment solid phase. According to our
40 results, the solubility of trace metal sulfide minerals ($\text{Cd} \ll \text{Fe}$) is a key-factor
41 controlling trace metal removal and consequently the magnitude as well as the
42 temporal and spatial heterogeneity of sedimentary fluxes. We argue that depending on
43 their sulfide solubility, sedimentary source or sink fluxes of trace metals will change
44 differentially as a result of declining oxygen concentrations and an associated

45 expansion of sulfidic surface sediments. Such a trend could cause a change in the
46 trace metal stoichiometry of upwelling water masses with potential consequences for
47 marine ecosystems in the surface ocean.

48

49

50 **1. Introduction**

51

52 **1.1 Scientific rationale**

53 The world's oceans are losing oxygen (e.g. Keeling et al. 2010; Stramma et al.
54 2010; Helm et al. 2011). In total around 2 % of oxygen has been lost over the past five
55 decades (Schmidtko et al., 2017) and an expansion of oxygen minimum zones (OMZs)
56 in the tropical oceans has been documented over the same timespan (Stramma et al.,
57 2008). The biogeochemical cycling of several nutrient-type trace metals (TMs) is likely
58 to be particularly susceptible to changing oxygen concentrations as they occur in
59 different oxidation states (e.g. Fe, Mn, Co) and/or are precipitated as sulfide mineral in
60 anoxic-sulfidic environments (e.g. Fe, Zn, Cd; listed in the order of decreasing sulfide
61 solubility). However, with the exception of Fe (Dale et al., 2015a; Lohan and Bruland,
62 2008; Rapp et al., 2018; Schlosser et al., 2018; Scholz et al., 2014a), little information
63 is available on how other TM fluxes will respond to ocean deoxygenation. As certain
64 TMs are essential for the growth of marine organisms (e.g. Fe, Mn, Co, Ni, Zn, Cd),
65 TM availability can (co-)limit primary productivity and therefore affect oceanic carbon
66 sequestration through the biological pump (Saito et al., 2008; Moore et al., 2013; Morel
67 et al., 2014). As a consequence, a better understanding of how TMs respond to low
68 oxygen conditions is essential for predicting how marine ecosystems and the carbon
69 cycle will evolve in the future ocean, with modelling scenarios predicting a continuation
70 of ocean deoxygenation (Bopp et al., 2002; Oschlies et al., 2008; Keeling et al., 2010)

71 Marine sediments are an important source or sink of TMs to the ocean under
72 low oxygen conditions (Böning et al., 2004; Brumsack, 2006; Scor Working Group,
73 2007; Severmann et al., 2010; Noble et al., 2012; Biller and Bruland, 2013; Conway
74 and John, 2015b; Klar et al., 2018). In the OMZ off the coast of Peru, substantial fluxes
75 of reduced Fe and other TMs across the sediment-bottom water interface have been

76 documented (Noffke et al., 2012; Scholz et al., 2016) or inferred (Hawco et al., 2016).
77 While a number of studies have addressed biogeochemical processes that lead to
78 benthic Fe release, the key biogeochemical processes and conditions that control the
79 sedimentary release or burial of other TMs in open marine systems are still poorly
80 constrained. Moreover, a detailed picture of removal or stabilization processes and
81 rates that take place in the highly dynamic water layer overlying the seafloor is lacking.

82 In this article, we compare the benthic biogeochemical cycling of Fe and Cd. It
83 has been established that the Peruvian OMZ represents a source of dissolved Fe to
84 the ocean (Noffke et al., 2012; Fitzsimmons et al., 2016; John et al., 2018). In contrast,
85 earlier studies have demonstrated that OMZs represent a sink for Cd (Janssen et al.,
86 2014; Böning et al., 2004). Because of their contrasting tendency to form sulfide
87 minerals and different supply pathways to the sediment, Fe and Cd can serve as
88 prototypes to provide information about how sedimentary fluxes of different TMs may
89 respond to declining oxygen concentrations. Under more reducing conditions the
90 mobility of TMs can either be enhanced or diminished, e.g., through precipitation of
91 sulfide minerals that are buried in the sediments (e.g. Westerlund et al., 1986; Rigaud
92 et al., 2013; Olson et al., 2017). Increased burial or release of TMs at the seafloor can
93 have an impact on the amplitude of primary productivity, especially at the eastern
94 ocean boundaries where the near-bottom water column is connected to the surface
95 ocean via upwelling. Moreover, since the inventories of TMs in the ocean are generally
96 dependent on the respective input and output fluxes, changes in the balance between
97 trace metal recycling and burial can have an impact on oceanic TM reservoirs on longer
98 timescales. By comparing the benthic biogeochemical cycling of Fe and Cd across
99 spatial and temporal redox gradients, we aim to provide general constraints on how
100 the stoichiometry of bio-essential TMs in seawater may be affected by ocean
101 deoxygenation.

102

103 **1.2. Marine biogeochemistry of iron**

104 Iron is the most abundant TM in phytoplankton and part of a range of
105 metalloenzymes that are involved in important biological functions, such as
106 photosynthesis or nitrogen fixation (Twining and Baines, 2013). Despite Fe being
107 highly abundant in the continental crust, its low availability limits primary productivity in
108 up to 30 % of the surface ocean area (Moore et al., 2013). This limitation arises from

109 the low solubility of its thermodynamically stable form in oxic waters, Fe(III).
110 Concentrations can reach up to ~ 1 nM when Fe(III) is kept in solution through
111 complexation with organic ligands (Rue and Bruland, 1997; Liu and Millero, 2002; Boyd
112 and Ellwood, 2010; Raiswell and Canfield, 2012). The thermodynamically stable form
113 of Fe under anoxic conditions, Fe(II), is more soluble and therefore anoxic waters are
114 typically characterized by higher dissolved Fe concentrations (up to tens of nM)
115 (Conway and John, 2014; Vedamati et al., 2014; Fitzsimmons et al., 2016; Schlosser
116 et al., 2018).

117 Sediments within OMZs are considered an important source of dissolved Fe and
118 some of the highest sedimentary Fe fluxes have been observed in these regions
119 (Severmann et al., 2010; Noffke et al., 2012). Under anoxic conditions, Fe(II) can be
120 liberated from the sediments into pore waters from Fe-(oxyhydr)oxides through
121 reductive dissolution by microbes or abiotic reduction with H₂S (Canfield, 1989). In the
122 absence of oxygen, dissolved Fe(II) escapes the rapid re-oxidation and subsequent
123 (oxyhydr)oxide precipitation and can, therefore, diffuse from pore waters into bottom
124 waters. However, in anoxic OMZs, where denitrification takes place, Fe(II) can also be
125 re-oxidized with nitrate as a terminal electron acceptor, either mediated by nitrate-
126 reducing microbes or abiotically through reaction with nitrite (Straub et al., 1996;
127 Carlson et al., 2013; Scholz et al., 2016; Heller et al., 2017). The solubility of Fe in
128 sulfidic (i.e. NO₃⁻ and NO₂⁻ depleted) water is relatively high (Rickard et al., 2006) and
129 during sulfidic events dissolved Fe can accumulate in the water column (up to
130 hundreds of nM) because of decreased Fe oxidation (Scholz et al., 2016) and
131 stabilization as aqueous Fe sulfide complexes and clusters (Schlosser et al., 2018).
132 However, Fe fluxes across the benthic boundary have also been hypothesized to
133 decrease under strongly sulfidic conditions in the surface sediments, when pore waters
134 become oversaturated with respect to Fe monosulfide (Scholz et al., 2014), which is
135 the precursor for pyrite (FeS₂) (Raiswell and Canfield, 2012).

136

137 **1.3. Marine biogeochemistry of cadmium**

138 Cd is abundant in phytoplankton despite concentrations that are one order of
139 magnitude lower than Fe (Moore et al., 2013; Twining and Baines, 2013). A function
140 for Cd as a catalytic metal atom in the carbonic anhydrase protein has been found in
141 diatoms (Lane and Morel, 2000) and it can also substitute Zn and enhance

142 phytoplankton growth under Zn limitation in different phytoplankton species (Price and
143 Morel, 1990; Lee and Morel, 1995; Sunda and Huntsman, 2000; Xu et al., 2008). In
144 marine sediments Cd can be released from the solid phase to the pore waters through
145 the remineralization of organic matter (Klinkhammer et al., 1982; Collier and Edmond,
146 1984; Gendron et al., 1986; Gerringa, 1990; Audry et al., 2006; Scholz and Neumann,
147 2007). After its release to the pore water, Cd can diffuse across the sediment-water
148 interface. Under anoxic and sulfidic conditions, Cd is thought to be precipitated as CdS
149 (Greenockite) and retained in the sediment (Westerlund et al., 1986; Gobeil et al.,
150 1987; Rosenthal et al., 1995; Audry et al., 2006). Due to its low sulfide solubility, CdS
151 can precipitate at much lower H₂S concentrations than FeS (mackinawite) (Morse and
152 Luther, 1999).

153 Most previous studies have focused on the benthic cycling of Cd in near- and
154 in-shore environments such as estuaries and lagoons (e.g. Westerlund et al., 1986;
155 Colbert et al., 2001; Audry et al., 2006b; Metzger et al., 2007; Point et al., 2007; Scholz
156 and Neumann, 2007). By contrast, little is known about Cd cycling in open-marine
157 sedimentary environments, where the redox- and sediment-dynamics are different.
158 Previous studies on sedimentary Cd cycling generally concluded that the flux of organic
159 material and the presence of H₂S are the most important factors controlling the balance
160 between Cd recycling versus precipitation and burial (e.g. Westerlund et al., 1986;
161 Colbert et al., 2001; Audry et al., 2006; Metzger et al., 2007; Scholz and Neumann,
162 2007). Low oxygen regions in the ocean are considered an important sink for Cd
163 (Janssen et al., 2014; Conway and John, 2015a; Xie et al., 2019) and sediments below
164 OMZs are highly enriched in Cd (Ragueneau et al., 2000; Böning et al., 2004; Borchers
165 et al., 2005; Muñoz et al., 2012; Little et al., 2015). However, the respective
166 contributions of different Cd removal mechanisms to Cd accumulation in the sediment
167 have not been quantified.

168

169 **1.4. Study area**

170 Seasonal upwelling of nutrient-rich waters off the Peruvian coast in austral
171 winter leads to high rates of primary productivity in the photic zone ($\sim 300 \text{ mmol C m}^{-3}$
172 d^{-1}) (Pennington et al., 2006). The combination of oxygen consumption through the
173 respiration of this organic matter and low oxygen concentrations in water masses that
174 supply upwelling regions, leads to the formation of one of the world's most intense

175 OMZs, with complete oxygen consumption in the OMZ core between ~ 100 m – 300 m
176 water depth (Karstensen et al., 2008; Thamdrup et al., 2012). Upon oxygen depletion,
177 NO_3^- can serve as an electron acceptor for respiration. Therefore, denitrification,
178 dissimilatory reduction of NO_3^- to ammonium (DNRA) and anaerobic ammonium
179 oxidation (anammox) with NO_2^- are important biogeochemical processes within the
180 anoxic and nitrogenous water column (Lam et al., 2009; Lam and Kuypers, 2011;
181 Dalsgaard et al., 2012). The OMZ overlying the Peruvian shelf is a temporally and
182 spatially dynamic system where biogeochemical conditions can range from fully oxic
183 to anoxic and sulfidic. Occasional shelf oxygenation events occur mostly during El Niño
184 events and are linked to the propagation of coastal trapped waves (Gutiérrez et al.,
185 2008). During such events, oxygenated water can be found on the upper slope to 200
186 m – 300 m water depth (Levin et al., 2002). By contrast, sulfidic events can occur
187 during periods of stagnation, when oxygen, NO_3^- and NO_2^- become depleted in the
188 water column due to sluggish ventilation. Once NO_3^- and NO_2^- are depleted,
189 chemolithoautotrophic H_2S oxidation is impeded. Hydrogen sulfide produced by
190 bacterial sulfate reduction in sediments can then be released to the water column
191 (Schunck et al., 2013) at rates reaching several $\text{mmol m}^{-2} \text{d}^{-1}$ (Sommer et al., 2016).

192 Our sampling campaign (cruises M136 and M137) took place in April and May
193 2017, during the decline of a coastal El Niño event. A coastal El Niño is a local
194 phenomenon that refers to reduced upwelling and increased sea surface temperatures
195 off the coasts of Peru and Ecuador, with typically heavy rainfall on land. During this
196 event in austral summer, coastal waters off Peru showed a strong positive sea surface
197 temperature anomaly of up to 2 to 4 °C (Echevin et al., 2018; Garreaud, 2018). The
198 warming is proposed to be a result of strong local alongshore wind anomalies and
199 equatorial Kelvin waves propagating towards the Peruvian coast (Echevin et al., 2018;
200 Peng et al., 2019).

201

202

203 **2. Methods**

204

205 **2.1 Sampling and sample handling**

206 In this study, data from three different types of samples were combined: (1) pore
207 waters for the determination of benthic diffusive fluxes and to study TM cycling in
208 sediments; (2) Benthic chamber incubations, to determine in-situ fluxes across the
209 sediment-water interface; (3) Bottom water concentration-depth profiles to determine
210 the fate of TMs in the particle-rich and dynamic near-bottom water column.

211 The sampling took place during RV Meteor cruises M136 and M137 in austral
212 autumn between April and May 2017. We also compared our data to benthic diffusive
213 Fe(II) fluxes from RV Meteor cruise M92 that took place in austral summer during
214 January 2013. Our sampling stations covered the entire Peruvian shelf and slope
215 across a transect at 12°S (Fig. 1) with water depths from 75 – to 950 m, thus including
216 stations above, inside and below the permanent OMZ. Our sampling of pore waters
217 and sample collection from benthic chamber incubations generally followed the
218 methodology described in Noffke et al. (2012).

219 Short sediment cores of 30 – 40 cm length were retrieved with a multiple corer
220 (MUC). Upon recovery, the cores were directly transferred into the ship's cool room
221 (4°C). The supernatant bottom water was instantly sampled and filtered through 0.2
222 µm cellulose acetate filters (Sartorius) and acidified to pH < 1 with subboiled distilled
223 HNO₃. The sediment cores were subsequently sampled in vertical sections in a glove
224 bag under Ar atmosphere to prevent any contact with oxygen. The sediment samples
225 were centrifuged to separate the pore waters from the sediment solid phase. Pore
226 waters were then filtered in another Ar-filled glove bag through 0.2 µm cellulose acetate
227 filters (Sartorius). An 8 ml aliquot was acidified to pH < 1 with subboiled distilled HNO₃
228 and stored in acid cleaned low-density polyethylene (LDPE) bottles for TM analysis.
229 Another aliquot was taken for analysis of H₂S concentrations. Additional sediment
230 subsamples were collected in pre-weighed cups for water content and porosity
231 determination as well as for Cd and organic C concentrations measurements in the
232 solid phase.

233 Benthic landers, constructed from titanium frames, containing two circular
234 benthic chambers for in-situ incubations, were deployed on the seafloor (see Sommer
235 et al. (2009) for details). After placement of the lander on the seafloor, the benthic
236 chambers (internal diameter of 28.8 cm) were partially driven into the sediment,
237 covering a sediment area of 651.4 cm². A volume between 12 – 18 l, overlying the first
238 20 – 30 cm of the seafloor, was enclosed in the chamber, depending on the insertion

239 depth of the chamber into the sediment. Prior to the incubation, the seawater contained
240 in the chamber was repeatedly replaced with ambient seawater to replace solutes and
241 flush out particles that might have been mobilized during the insertion of the chamber
242 into the sediment. Over the incubation time of around 32 hours, 8 consecutive samples
243 of 12 ml were filtered in-situ through 0.2 μm cellulose acetate filters (Sartorius) via
244 peristaltic pumps and collected in quartz glass tubes. All sampling tubes were acid
245 cleaned prior to use to guarantee a TM clean sampling. After recovery of the lander,
246 the quartz glass tubes were transferred to the laboratory and samples were stored in
247 acid cleaned LDPE bottles and acidified to $\text{pH} < 2$ with subboiled distilled HNO_3 . Other
248 samples were collected simultaneously for analysis of nitrogen species. The incubated
249 sediments within the benthic chamber were sampled after recovery of the lander and
250 pore waters were extracted to analyze H_2S concentrations for comparison with pore
251 water profiles from parallel MUCs.

252 To determine TM concentrations across the near-bottom water column, water
253 samples were collected at 0.5, 1.0, 2.0, 3.0 and 4.0 m above the seafloor using
254 sampling apparatus attached to the landers. Filter holders with 0.2 μm polyether
255 sulfone filters (Supor) were attached at the various depths and connected to sampling
256 tubes that went through peristaltic pumps into gas sampling bags (Tedlar). Sampling
257 at 3.0 m and 4.0 m above the seafloor was realized by attaching the filter holders and
258 tubing to an arm that was automatically unfolded upon placement of the lander at the
259 seafloor. The peristaltic pumps transferred the seawater from the sampling depths into
260 the sampling bags over the same time period as the lander incubations of around 32
261 hours. This resulted in an average sample volume of 1.5 l per depth. All filters, tubing
262 and sampling bags were acid cleaned prior to deployment to guarantee a TM clean
263 sampling. Directly after sample retrieval, a 60 ml aliquot was stored in acid cleaned
264 LDPE bottles and acidified to $\text{pH} < 2$ for TM analysis. Another aliquot was taken for
265 analysis of silicic acid ($\text{Si}(\text{OH})_4$).

266

267 **2.2 Analytical methods**

268 Concentrations of $\text{Fe}(\text{II})$ in pore waters were measured on board directly after
269 sample retrieval by photometry using the ferrozine method (Stookey, 1970). Other
270 geochemical parameters in our different samples were also determined photometrical
271 (U-2001 Hitachi spectrometer) using standard techniques (Grasshoff et al., 1999).

272 Hydrogen sulfide concentrations were determined using the methylene blue method
273 and Si(OH)_4 concentrations were determined using a heptamolybdate solution as
274 reagent. Concentrations of nitrogen species were determined by an auto-analyzer
275 (QuAatro, SEAL Analytical) using sulfanilamide as reagent (Hydes et al., 2010).

276 For TM analysis of bottom water samples we followed the procedure described
277 by Rapp et al. (2017), whereby the TMs were pre-concentrated by a fully automated
278 device (SeaFAST). After raising the sample pH to 6.4 with an ammonium acetate buffer
279 (1.5 M), 15 ml of sample was loaded onto a chelating resin column, where the seawater
280 matrix was rinsed off, before the TMs were collected into 1ml elution acid (1 M
281 subboiled HNO_3). Due to the smaller size of pore water samples and samples from
282 benthic lander incubations, a half-automated device (Preblab) with a smaller sample
283 loop and thus dead volume was used. On this device, sample loading and collection
284 as well as the addition of buffer was done manually. For samples from benthic lander
285 incubations and pore waters, an amount of 3 ml and 1 ml, respectively, was needed
286 for pre-concentration. The samples were diluted with de-ionised water (MilliQ,
287 Millipore) to increase the sample volume to 5 ml for samples from benthic chamber
288 incubations and to 3 ml for pore waters. The pre-concentrated samples were measured
289 by ICP-MS (HR-ICP-MS; Thermo Fisher Element XR) and TM concentrations were
290 quantified by isotope dilution. The detection limits were 28.8 pM for Fe and 0.8 pM for
291 Cd (Rapp et al., 2017). Accuracies for replicate measurements of reference seawater
292 certified for TMs are listed in Table 1.

293 For the calculation of sedimentary Cd enrichments (Cd_{xs}), Cd and Al contents
294 in sediments were determined following total digestions of freeze dried and ground
295 sediment samples. The sediment was digested in 40 % HF (suprapure), 65 % HNO_3
296 (suprapure) and 60 % HClO_4 (suprapure). Concentrations were measured by ICP-OES
297 (VARIAN 720-ES). The reference standard MESS was used to check the digestion
298 procedure. The accuracy was ± 0.3 % for Cd and ± 1.3 % for Al (MESS-3 Cd: $0.24 \pm$
299 $0.01 \mu\text{g g}^{-1}$, recommended value $0.24 \pm 0.01 \mu\text{g g}^{-1}$, MESS-3 Al: $8.59 \pm 0.11 \mu\text{g g}^{-1}$,
300 recommended value $8.59 \pm 0.23 \mu\text{g g}^{-1}$).

301 Organic carbon content in the sediment was determined using an Elemental
302 Analyzer (Euro EA) after removal of inorganic carbon with 0.25 mM HCl. Precision of
303 the measurement was ± 1 %.

304

305 **2.3 Diffusive flux calculations**

306 Benthic diffusive fluxes (F_D) were determined using Fick's first law of diffusion
307 using concentration gradients between the uppermost pore water sample (0 – 1 cm)
308 and the overlying bottom water (dC/dx) (Boudreau, 1997):

$$309 \quad F_D = -\Phi D_{sed}(dC/dx) \quad (1)$$

310 The effective molecular diffusion coefficients of Fe and Cd for sediments (D_{sed}) were
311 calculated from the molecular diffusion coefficient in seawater (D_{sw}) under standard
312 conditions (Li and Gregory, 1974) by adjusting it to in-situ temperature, pressure and
313 salinity applying the Stokes-Einstein Equation. We determined the diffusion
314 coefficients for sediments as follows:

$$315 \quad D_{sed} = D_{sw}/\theta^2 \quad (2)$$

316 Tortuosity (θ) was calculated from porosity (Φ) as follows (Boudreau, 1997):

$$317 \quad \theta^2 = 1 - \ln(\phi^2) \quad (3)$$

318 Positive values represent a flux from the bottom water into the sediment pore
319 water, negative values a flux from the sediment pore water into the bottom water. All
320 input values for the diffusive flux calculations are listed in Tables S1 and S2 in the
321 supplement.

322 ~~We chose to use the commonly applied approach of a two point concentration~~
323 ~~gradient for the determination of diffusive fluxes, as more advanced curve fitting~~
324 ~~methods would fail to capture sharp concentration gradients at the sediment surface~~
325 ~~and, thus, lead to erroneous flux estimates (Shibamoto and Harada, 2010).~~

326 ~~—————The fluxes from benthic lander incubations were calculated from the slopes of~~
327 ~~linear regressions from the concentration-time data. The relevant equations are listed~~
328 ~~together with the coefficients of determination (R^2) in Table S4 in the supplement.~~
329 ~~Fluxes were determined using the water volume enclosed in the benthic chamber,~~
330 ~~estimated for each deployment from the insertion depth of the benthic chamber into~~
331 ~~the sediment.~~

332 Due to the coarse resolution of our pore water profiles and the steep gradients
333 between the uppermost pore water and bottom water sample (see close-up profiles,
334 Fig. S1 and S2 in the supplement), we chose to follow previous studies (e.g. Noffke et

335 [al., 2012; Lenstra et al., 2019; Scholz et al., 2019](#)) and calculate diffusive benthic fluxes
336 [based on a two point concentration gradient. Including deeper samples into a linear](#)
337 [regression or applying more advanced curve fitting methods would reduce the](#)
338 [statistical uncertainty but fail to capture the sharp concentration gradients at the](#)
339 [sediment surface, thus, leading to erroneous flux estimates \(cf. Shibamoto and](#)
340 [Harada, 2010\).](#)

341 [The fluxes from benthic lander incubations were calculated by fitting a linear](#)
342 [regression to the concentration change over time. The relevant equations are listed](#)
343 [together with the coefficients of determination \(\$R^2\$ \) in Table S4 in the supplement.](#)
344 [Concentration changes over time were converted to fluxes by taking into account the](#)
345 [water volume enclosed in the benthic chamber, estimated for each deployment from](#)
346 [the insertion depth of the benthic chamber into the sediment. The uncertainties of](#)
347 [fluxes were estimated by propagating the uncertainties of the linear regressions.](#)
348 [Following previous studies \(e.g. Friedrich et al. \(2002\); Lenstra et al. \(2019\)\), only](#)
349 [fluxes where the linear regression has an \$R^2 > 0.3\$ are reported in Tables 2 and 3.](#)

352 **3. Results**

354 **3.1 Biogeochemical conditions in the water column**

355 Due to the particular atmospheric and oceanographic conditions, the decline of
356 a coastal El Niño during our sampling campaign (cruises M136 and M137), the water
357 column overlying the Peruvian shelf was oxygenated. Oxygen concentrations were >
358 20 μM in the water column down to around 100 m water depth. However, bottom water
359 oxygen concentrations directly above the seafloor, measured using optodes attached
360 to lander, were below the detection limit (> 1 μM) at the shallowest station (Station 1).
361 The OMZ, with O_2 concentrations < 5 μM , extended from around 120 to 400 m water
362 depth. The water column within the OMZ was nitrogenous (i.e. NO_3^- reducing) as
363 indicated by the presence of NO_2^- ($\geq 4 \mu\text{M}$), an intermediate product of denitrification
364 (Zumft, 1997). Oxygen gradually increased to > 50 μM below 400 m towards 950 m
365 water depth (Fig. 2). As we will compare some of our data to those of an earlier cruise

366 (M92), the corresponding oxygen distribution across the Peruvian continental margin
367 is shown for comparison (Fig. 2).

368

369 **3.2 Bottom water, pore water and benthic flux data**

370

371 **3.2.1 Iron**

372 Iron concentrations in near-bottom waters decreased from near-shore to off-
373 shore stations, from > 100 nM at the shallowest shelf station at 75 m water depth
374 (Station 1) to 6 nM at 750 m water depth (Station 9) (Fig. 3). At a number of stations
375 within the OMZ (Stations 3 and 4), vertical concentration gradients were observed.
376 Here Fe concentrations decreased by 15 – 20 nM from 0.5 to 4 m above the seafloor.
377 Multiple sampling at the shallowest shelf station (Station 1) revealed that Fe
378 concentrations were temporally variable and ranged from ~ 100 nM at the end of April
379 to < 60 nM at the end of May 2017.

380 Concentrations of Fe(II) in pore waters were highest (up to a few μM) in the
381 upper 5 – 10 cm of the sediment cores. Deeper in the sediment cores, concentrations
382 decreased to > 0.2 μM (Fig. 4). At all stations, sharp concentration gradients between
383 the uppermost pore water and bottom water sample were observed, with higher
384 concentrations in pore waters at the sediment surface (μM) than in the overlying bottom
385 water (nM). This observation implies a diffusive flux from pore waters into bottom
386 waters. The steepest concentration gradients across the sediment-water interface
387 were observed within the OMZ. The highest Fe(II) concentrations at the sediment
388 surface (> 6 μM) were observed at Station 4 (145 m water depth). At this station, the
389 benthic diffusive flux into the bottom waters was also highest at -17.5 $\text{mmol m}^{-2} \text{y}^{-1}$.
390 The lowest diffusive fluxes of 0.0 (due to concentrations below the detection limit) and
391 -0.3 $\text{mmol m}^{-2} \text{y}^{-1}$ were observed on the upper slope below the OMZ at Stations 9 and
392 10 respectively (Table 2). An accumulation of H_2S in pore waters coincided with a
393 depletion of Fe(II) concentrations (Fig. 4). At Station 1, we observed the highest H_2S
394 concentrations throughout the core and in particular at the sediment surface, with
395 maximum concentrations reaching > 4 mM. At Stations below the OMZ (Stations 9 and
396 10), no H_2S was detected within pore waters (Fig. 4).

397 Iron concentrations inside the benthic chambers reached maximum
398 values > 300 nM. At Stations 4 and 6, located inside the OMZ, concentrations in the
399 chambers increased in a linear way during the incubation. At stations above and below
400 the OMZ, we did not observe a similar trend over time. For comparison with diffusive
401 fluxes, we estimated benthic Fe fluxes from linear regressions of Fe concentrations
402 versus time (Table 2). We also calculated the theoretical concentration gradients over
403 time in the benthic chambers based on our diffusive flux estimates (Fig. 5). The
404 incubation data were largely consistent in direction and slope with the diffusive fluxes.
405 In particular, the projected and observed concentration gradients were in good
406 agreement at stations inside the OMZ (Station 4 and 6), where the highest diffusive
407 fluxes of -17.5 and -8.0 mmol m⁻² y⁻¹ were observed. At these stations also the highest
408 R² for the linear regressions of the concentration change over the incubation time were
409 calculated (Station 4: R² = 0.7, Station 6: R² = 0.5) (Table S4). At stations below the
410 OMZ, diffusive fluxes of < 1 mmol m⁻² y⁻¹ were too low to be detected over the
411 incubation time of 32 hours.

412

413 **3.2.2 Cadmium**

414 In near-bottom waters Cd concentrations increased with distance from the
415 coast, from 0.4 nM at the shallowest station at 75 m water depth (Station 1) to 1.1 nM
416 below the OMZ at 750 m water depth (Station 9). Cadmium concentrations were
417 constant at each station between 0.5 and 4 m above the seafloor (Fig. 3).

418 Cadmium concentrations in pore waters ranged between 0.1 – 2 nM (Fig. 6).
419 Within the OMZ, bottom water concentrations were higher than concentrations in pore
420 water at the sediment surface (0 - 1 cm), indicating a downward diffusive flux into the
421 sediments. The benthic diffusive fluxes inside the OMZ were on the order of 0.6 – 0.8
422 μmol m⁻² y⁻¹ (Table 3). In contrast, at Stations 1 and 9 an upward-directed
423 concentration gradient was observed, indicating a diffusive flux from the sediments into
424 bottom waters. The upward diffusive flux was -1.9 μmol m⁻² y⁻¹ above the permanent
425 OMZ and -0.2 μmol m⁻² y⁻¹ below the OMZ (Table 3). Pore water Cd concentrations at
426 greater sediment depths were mostly higher than bottom water concentrations. In
427 some cases (Stations 3 and 4), elevated pore water Cd concentrations (up to 2 nM)
428 coincided with elevated H₂S concentrations (few hundred μM).

429 In the benthic chambers three different Cd trends were observed (Fig. 7). Above
430 the permanent OMZ (Station 1), Cd concentrations in the chambers were low (< 0.2
431 nM) throughout the incubation period, indicating no Cd flux. At sites within the OMZ
432 (Stations 4, 5 and 6), concentrations decreased from ~ 0.6 – 0.3 nM over the course
433 of the incubation. Below the OMZ (Stations 9 and 10), Cd concentrations in the
434 chamber were high (~ 1 nM) and remained constant or increased slightly during the
435 incubation. At sites within the OMZ, Cd removal within the chamber was near-linear
436 (Stations 4, 5 and 6: $R^2 = \geq 0.9$) (Table S4), which translates to a removal flux of 13 –
437 23 $\mu\text{mol m}^{-2} \text{y}^{-1}$. The Cd removal fluxes in benthic chambers were more than one order
438 of magnitude higher than diffusive benthic fluxes (0.6 – 0.8 $\mu\text{mol m}^{-2} \text{y}^{-1}$) (Table 3).

439

440

441 **4. Discussion**

442

443 **4.1 Benthic iron cycling**

444 **4.1.1 Comparison of diffusive and in-situ benthic chamber iron fluxes**

445 Concentrations of Fe in bottom waters from benthic chamber incubations are
446 mostly higher than in ambient bottom waters because of Fe release from the sediment
447 and an accumulation in the enclosed water volume inside the benthic chamber. In the
448 absence of oxygen and, thus, bottom-dwelling macrofauna at stations within the OMZ,
449 bioturbation and bioirrigation are unlikely to exert an important control on sedimentary
450 Fe release. Consistent with this notion, the slope calculated from benthic diffusive
451 fluxes is largely consistent with the concentration gradients observed within the benthic
452 chambers (Fig. 5). Moreover, our fluxes from benthic chamber incubations and
453 diffusive fluxes are generally of the same order of magnitude (Table 2). Therefore,
454 diffusive transport of dissolved Fe from the sediment into the bottom water seems to
455 be the main control on the concentration evolution observed within the benthic
456 chamber.

457 Some of the concentration gradients in benthic chambers are non-linear,
458 indicating that the Fe flux was not constant during the incubations. This is a common
459 observation in Fe flux data from benthic chamber incubations and higher Fe fluxes

460 generally have higher R^2 values for the linear regressions (Friedrich et al., 2002;
461 Turetta et al., 2005; Severmann et al., 2010; Lenstra et al., 2019). However, the non-
462 linearity can be used to identify additional processes affecting Fe concentrations and
463 fluxes within the benthic chamber, which may also affect fluxes under natural
464 conditions. One possible process that can remove dissolved Fe(II) under anoxic
465 conditions is Fe oxidation with NO_3^- as the terminal electron acceptor or oxidation with
466 NO_2^- (Straub et al., 1996; Carlson et al., 2013; Klueglein and Kappler, 2013). The
467 oxidation of reduced Fe in the absence of oxygen, either microbially mediated with
468 NO_3^- or abiotically with NO_2^- , has been hypothesized to be important in the water
469 column of the Peruvian OMZ (Scholz et al., 2016; Heller et al., 2017). During our
470 incubation at Station 4 (Fig. 8), we observed a decline in Fe concentrations during the
471 first ten hours of the incubation time. Concurrently, NO_3^- concentrations were
472 decreasing, while NO_2^- accumulated, presumably due to progressive denitrification and
473 release from the sediments. Once NO_3^- and NO_2^- were depleted, Fe concentrations
474 started to rise again, resulting in the highest in-situ Fe flux observed throughout our
475 sampling campaign (Table 2). The coincidence in timing of Fe accumulation and NO_2^-
476 decrease suggest that depletion of Fe at the beginning of the incubation was most
477 likely caused by Fe oxidation with NO_2^- . The incubation at Station 4 was the only one
478 where NO_3^- and NO_2^- were substantially removed during the incubation. However, the
479 high Fe flux cannot be interpreted as a natural flux estimate at steady state. In general,
480 we argue that bottom water NO_2^- concentrations exert a first order control on the
481 intensity of Fe efflux at the absence of oxygen and, therefore, need to be considered
482 in the evaluation of sedimentary Fe mobility in anoxic-nitrogenous OMZs.

483 During the incubations at Stations 1, 9 and 10, Fe concentrations did not
484 continuously increase but fluctuated between high and low values. This observation
485 could be explained by a combination of bioirrigation and bioturbation at stations where
486 oxygen was present (stations 9 and 10), as well as rapid Fe oxidation and precipitation
487 processes. Under oxic conditions, bottom-dwelling macrofauna is likely to increase the
488 transfer of dissolved Fe from the sediments into the bottom water (Elrod et al., 2004;
489 Lenstra et al., 2019). During episodes of oxygenation a population of macrofauna that
490 can enhance bioturbation and bioirrigation was observed on the Peruvian shelf
491 (Gutiérrez et al., 2008). However, under oxic conditions, any Fe delivered to the
492 chamber is prone to rapid oxidative removal. Moreover, ex-situ experiments have
493 demonstrated a fast and efficient removal of up to 90% of dissolved Fe in incubated

494 bottom waters due to particle resuspension (Homoky et al., 2012). Bioturbation and
495 bioirrigation could also contribute to particle resuspension at oxic stations, thus leading
496 to removal of dissolved Fe.

497 Furthermore, colloidal Fe could modify Fe concentrations within our samples
498 and explain some of the fluctuations observed during the incubations. Colloids are
499 quite reactive, they are much more soluble than larger particles and can be rapidly
500 reduced and dissolved in anoxic environments, but they can also aggregate into larger
501 particles (Raiswell and Canfield, 2012). The transfer of Fe between dissolved, colloidal
502 and particulate pools is likely to affect the balance between Fe transport and re-
503 precipitation and -deposition to some extent. However, since did not differentiate
504 between colloidal and truly dissolved fractions during our sampling, we cannot discuss
505 this aspect further based on our data.

506 Oxidation processes and interactions with particles can efficiently remove Fe
507 shortly after its transfer to bottom waters and this process is likely to be most intense
508 close to the seafloor where the highest particle concentrations prevail. We argue that
509 the same processes are reflected by declining Fe concentrations away from the
510 seafloor in some of the bottom water profiles (at Stations 3 and 4) (Fig. 3).

511

512 **4.1.2 Removal rates of dissolved iron in the near-bottom water column**

513 We observed declining Fe concentrations in the first 4 m away from the seafloor
514 at Stations 3 and 4, which hints at removal of dissolved Fe in the near bottom waters
515 after its release from the sediments. To differentiate between dilution with ambient
516 bottom water (by currents) from Fe removal from the dissolved phase, Fe
517 concentrations were normalized by Si(OH)_4 measured in the same samples (Fig. 3).
518 Due to opal dissolution within Peru margin sediments, Si(OH)_4 is released into bottom
519 waters (Ehlert et al., 2016). In contrast to Fe, we assume that Si(OH)_4 behaves
520 conservatively and precipitation reactions within the bottom waters are of subordinate
521 importance. The decreasing Fe to Si(OH)_4 ratios at Station 3 and 4 with distance from
522 the seafloor indicate that there is Fe removal within the near-bottom water column that
523 must be related to precipitation processes or scavenging.

524 We further constrained rates of dissolved Fe removal at stations with a
525 discernable Fe to Si(OH)_4 gradient within the first 4 m distance from the seafloor. To

526 this end, we first determined an eddy diffusion coefficient (K_y) using $\text{Si}(\text{OH})_4$ fluxes
527 from benthic chamber incubations (F_{Si}) (see chapter 2.3 for methodology) and the
528 known concentration gradient of dissolved $\text{Si}(\text{OH})_4$ within the bottom water (d_{Si}/d_x),
529 where x is the height above the seafloor. At the seafloor, the flux of $\text{Si}(\text{OH})_4$ from the
530 sediment is equal to the flux in the water column.

$$531 \quad F_{\text{Si}} = -K_y(d_{\text{Si}}/d_x) \quad (4)$$

532 This equation can be solved for the eddy diffusion coefficient.

533 Dissolved Fe in the bottom water (DFe) can be described by the solving the
534 diffusion-reaction equation for DFe (ignoring advection and assuming a steady-state
535 first-order consumption of dissolved Fe):

$$536 \quad DFe = C_{\text{BW}} * \exp.(-\sqrt{k_{\text{Feox}}/K_y}) \quad (5)$$

537 The equation can be fitted to the measured DFe concentrations in the bottom water by
538 adjusting the Fe concentration directly above the seafloor (C_{BW}) and the Fe oxidation
539 constant (k_{Feox}). From the fitted first-order rate constant k_{Feox} , the half-life for dissolved
540 Fe in bottom waters can be calculated.

541 The half-lives of dissolved Fe in the first 4 m away from the seafloor are 2.5 min
542 and 0.3 min at Stations 3 and 4, respectively (Table 4). Another study reported a
543 dissolved Fe half-life of 17 hours under nitrogenous conditions in the first 10 – 20 m
544 above the seafloor in the Peruvian OMZ (Scholz et al., 2016). Our calculations suggest
545 that Fe removal in near-bottom waters is much faster. The approach assumes that
546 $\text{Si}(\text{OH})_4$ is transported vertically by eddy diffusion and eddy diffusion and oxidation
547 control the half-life of Fe in the first 4 m above the seafloor. It is possible that our
548 assumption of solute transport by eddy diffusion is not correct. Alternatively,
549 decreasing Fe and $\text{Si}(\text{OH})_4$ concentration above the seafloor could be due to super-
550 imposed water layers with different Fe and $\text{Si}(\text{OH})_4$ concentrations but little vertical
551 exchange. In this case our calculated half-life would be an underestimation.

552 As mentioned above (chapter 4.1.1), in the absence of oxygen, removal
553 processes of dissolved Fe could be related to oxidation of dissolved Fe with NO_2^- or to
554 interactions with suspended particles, which are likely to be most abundant directly
555 above the seafloor. Further research on dissolved-particulate interactions, including

556 the role of colloidal Fe, in bottom waters is needed to better constrain how sedimentary
557 Fe fluxes are modified in the near-bottom water column.

558

559 **4.1.3 Controls on the temporal variability of benthic iron fluxes**

560 The Peruvian OMZ is known to experience high-amplitude fluctuations in
561 upwelling intensity as well as variability in bottom water oxygen, NO_3^- , NO_2^- and H_2S
562 concentrations (Pennington et al., 2006; Gutiérrez et al., 2008; Graco et al., 2017;
563 Ohde, 2018). To get an insight into how different biogeochemical conditions control
564 benthic diffusive Fe(II) fluxes, we compared the fluxes from our recent cruise with
565 fluxes from our earlier cruise M92 (Fig. 9). Cruise M92 took place in austral autumn
566 2013 following the main upwelling season and during a period of intense primary
567 productivity. Due to reduced upwelling and stable density stratification, the water
568 column on the shallow shelf was not only depleted in oxygen but also in NO_3^- and NO_2^-
569 during cruise M92 (Sommer et al., 2016). Under such conditions,
570 chemolithoautotrophic H_2S oxidation with NO_3^- or NO_2^- was impeded so that pore water
571 H_2S could be released from the sediment into the water column. As a result, the water
572 column during M92 was sulfidic between around 50 – 150 m water depth with the
573 highest H_2S concentration of 13 μM observed at 50 m depth (Fig. 2). While the
574 biogeochemical conditions on the shallow shelf were fundamentally different to those
575 during M136 and M137, below 150 m water depth the conditions were largely
576 comparable (oxygen-depleted, NO_3^- : 20 – 30 μM , NO_2^- up to 9 μM between 150 – 300
577 m). At the stations with similar biogeochemical water column conditions, the Fe(II)
578 fluxes during both sampling campaigns were remarkably similar (Fig. 9). However,
579 similar to the temporal variability of Fe concentrations in bottom waters at Station 1
580 (Fig. 3), we observed a pronounced difference in the diffusive flux magnitude on the
581 shallow shelf where the biogeochemical conditions differed between both cruises. The
582 highest diffusive flux during M92 in 2013 of $-22.7 \text{ mmol m}^{-2} \text{ y}^{-1}$ was measured at Station
583 1. By contrast, during M136/137 in 2017 we determined a much lower flux of -2.6 mmol
584 $\text{m}^{-2} \text{ y}^{-1}$ at this station. During M136 and M137 the highest flux of $-17.5 \text{ mmol m}^{-2} \text{ y}^{-1}$ was
585 measured at Station 4 at 145 m water depth.

586 Diffusive fluxes are a function of the concentration gradient between pore water
587 and bottom water (Eq. (1)). As dissolved Fe concentrations in bottom waters are
588 generally much lower (nM) compared to those observed in pore waters (μM), the flux

589 magnitude is chiefly determined by differences in pore water Fe concentrations. During
590 M92, pore waters at the sediment surface were characterized by high dissolved Fe
591 concentrations (4.8 μM in the upper pore water sample), which resulted in a steep
592 gradient and a comparably high Fe flux. Under the slightly sulfidic conditions that
593 prevailed in the water column during M92, oxidative removal of dissolved Fe(II) with
594 NO_3^- or NO_2^- was impeded (Scholz et al., 2016) and dissolved Fe(II) could be stabilized
595 as aqueous iron sulfide (Schlosser et al., 2018). Therefore, the bottom water was
596 characterized by high dissolved Fe concentrations (up to 0.7 μM in the supernatant
597 bottom water of MUCs).

598 Despite oxic conditions in the water column during M136 and M137, we
599 observed much higher H_2S concentrations in surface sediments at Station 1 compared
600 to M92 (4100 μM during M136 and M137 versus 1800 μM during M92 within the first 8
601 cm of the core) (Fig. 4). Because of higher H_2S concentrations, Fe concentrations were
602 controlled by the solubility of Fe monosulfide minerals (FeS). It may seem
603 counterintuitive that the surface sediment was highly sulfidic, while the overlying water
604 column was oxygenated. In order to explain this observation, we need to consider the
605 role of mats of filamentous sulfur oxidizing bacteria in controlling H_2S concentrations
606 in surface sediments. (Gutiérrez et al., 2008; Noffke et al., 2012; Yücel et al., 2017).
607 During M92 these mats were generally abundant on the shelf and upper slope
608 (Sommer et al., 2016), thus limiting the extent of H_2S accumulation within surface
609 sediments. Previous studies demonstrated that mats of sulfur oxidizing bacteria can
610 disappear during periods of oxygenation (Gutiérrez et al., 2008). Consistent with this
611 previous finding, visual inspection of the seafloor using the video-guided MUC revealed
612 that the abundance of bacterial mats on the seafloor seemed greatly reduced, which
613 is most probably related to oxic bottom water conditions on the shallow shelf during
614 the coastal El Niño event. As these microaerophilic organisms tend to avoid high
615 oxygen concentrations they probably started to die off or withdraw into the sediment
616 once oxygen levels raised. We suggest that the disappearance of sulfide-oxidizing
617 bacteria under oxic conditions created a situation where H_2S accumulation in the
618 surface sediment and FeS precipitation limited the extent of Fe release into the bottom
619 water.

620

621 **4.2 Benthic cadmium cycling**

622 4.2.1 Comparison of diffusive and in-situ benthic chamber cadmium fluxes

623 At stations above and below the permanent OMZ (Stations 1, 9 and 10),
624 ~~diffusive the slopes of Cd fluxes and fluxes in concentrations versus time during~~ benthic
625 ~~chambers chamber incubations~~ were largely consistent (~~Table 3 with theoretical Cd~~
626 ~~concentration gradients over time based on our diffusive flux estimates (Fig. 7)~~). In
627 contrast, the fluxes determined with benthic chambers at stations within the OMZ
628 (Station 4, 5 and 6) were 25 to 40 times higher than the diffusive flux (Table 3). This
629 discrepancy demonstrates that diffusion cannot be the dominant process leading to
630 the continuous decrease of dissolved Cd during benthic chamber incubations.
631 Alternatively, Cd could be precipitated within the benthic chamber and removed
632 through downward sinking of Cd-rich particles. Cadmium sulfide (greenockite) has a
633 relatively low solubility compared to sulfide minerals of other TMs ($\text{CdS} \ll \text{FeS}$). It is
634 generally agreed that CdS precipitation can take place at trace amounts of H_2S (H_2S
635 $< 1 \mu\text{M}$, i.e., below the detection limit of the method applied in this study) (Davies-
636 Colley et al., 1985; Rosenthal et al., 1995). Previous studies using in-situ benthic flux
637 chambers have concluded that production of H_2S in the sediment or the accumulation
638 of H_2S in benthic chambers during incubations can switch the direction of the Cd flux
639 or intensify Cd removal through CdS precipitation (Westerlund et al., 1986; Colbert et
640 al., 2001). Precipitation of CdS during the incubation is, therefore, a viable explanation
641 for the discrepancy between diffusive Cd flux and Cd fluxes in benthic chambers
642 observed in our study. Furthermore, the three different trends of Cd concentrations
643 observed in benthic chamber incubations can be related to H_2S concentrations in the
644 surface sediment below the benthic chambers (Table 3). At stations within the OMZ
645 (Stations 4, 5 and 6), pore water H_2S concentrations in surface sediments were
646 moderate (few μM). It is likely that there was a continuous leakage of trace amounts of
647 H_2S from the pore water into the bottom waters during the incubation, thus leading to
648 CdS precipitation and declining Cd concentrations. On the shallowest shelf station
649 (Station 1), where pore water H_2S concentrations in the surface sediment were high
650 (hundreds of μM), a potentially large amount could have been released at the
651 beginning of the incubation, thus explaining pronounced Cd depletion in the chamber
652 compared to the surrounding bottom water (0.1 nM within the chamber compared to
653 0.4 nM outside the chamber). Below the OMZ (Stations 9 and 10), where there was no
654 H_2S present in surface sediments, there was no Cd depletion in the chamber during
655 the incubation and, consistent with previous studies in oxic settings (Westerlund et al.,

1986; Ciceri et al., 1992; Zago et al., 2000; Turetta et al., 2005), both diffusive and benthic chamber flux data were indicative of an upward-directed flux out of the sediment. Due to the absence of H₂S, dissolved Cd released from biogenic particles in the surface sediment could accumulate in the pore water thus driving a diffusive flux out of the sediment.

661

4.2.2 Quantification of the sedimentary cadmium sink

Consistent with our Cd flux data there is general consent that OMZs are a sink for Cd. Several water column studies have observed Cd depletion in water masses within the Peruvian and other OMZs, which was mostly attributed to Cd removal via CdS precipitation in sulfidic micro-niches within particles in the water column (Janssen et al., 2014; Conway and John, 2015b). Sedimentary studies showed that Cd is highly enriched in OMZ sediments, which has mostly been attributed to the delivery of Cd with organic material and subsequent fixation as CdS within sulfidic sediments (Ragueneau et al., 2000; Böning et al., 2004; Borchers et al., 2005; Muñoz et al., 2012; Little et al., 2015). Based on our data, we can quantify the delivery of Cd to the sediments via three different pathways: (1) diffusion across the sediment-water interface and CdS precipitation within the sediment; (2) Cd incorporation by phytoplankton and delivery to the sediment with organic matter; (3) CdS precipitation in the water column and particulate delivery to the sediment (Table 3).

The enrichment of Cd in the sediment relative to the lithogenic background (expressed as excess Cd concentration; Cd_{xs}) was calculated using the following equation (Brumsack, 2006):

$$Cd_{xs} = Cd_{sample} - Al_{sample} * (Cd/Al)_{crust} \quad (6)$$

The Cd/Al ratio of the upper continental crust (1.22*10⁻⁶) was used as lithogenic background reference (Taylor and McLennan, 2009). To calculate the flux of Cd to the sediment, Cd_{xs} was multiplied with the mass accumulation rate (MAR) from published data for each individual site (Dale et al., 2015b). To approximate the amount of Cd delivered to the sediment with organic material, the average concentration ratio of Cd to C in phytoplankton (Moore et al., 2013) was multiplied by published particulate organic carbon rain rates (maximum estimate) or burial rates (minimum estimate) for each individual site (Dale et al., 2015b). The Cd delivery via precipitation in the water

688 column was determined as the remainder of $Cd_{xs} * MAR$ after subtraction of the two
689 other sources (i.e., diffusive flux and minimum/maximum delivery by organic material).

690 Sediments at all stations on the Peruvian shelf and slope are enriched in Cd
691 relative to the lithogenic background. The accumulation rate of Cd decreases with
692 distance from the coast from $250 \mu\text{mol m}^{-2} \text{y}^{-1}$ at Station 1 to $4 \mu\text{mol m}^{-2} \text{y}^{-1}$ at Station 9
693 (Table 3). These fluxes generally exceed the amount of Cd delivered to the sediments
694 via diffusion and associated with organic material. Together these mechanisms of Cd
695 delivery can only account for ~ 20 % of the Cd enrichment at stations above and inside
696 the permanent OMZ, with the delivery with organic material being of greater
697 importance. The remaining Cd enrichment in the sediment (~ 80 %), after subtraction
698 of diffusive and minimum/maximum organic Cd sources, must be related to CdS
699 precipitation in the water column and delivery of Cd-rich particles to the sediment. This
700 removal process can be a combination of CdS precipitation in sulfidic micro-niches
701 around sinking particles (Janssen et al., 2014; Bianchi et al., 2018), CdS precipitation
702 in sulfide plumes (Xie et al., 2019) when sedimentary H_2S can spread throughout the
703 water column (Schunck et al., 2013; Ohde, 2018), and precipitation of CdS in the near-
704 bottom water (this study). Our estimated CdS precipitation in the water column within
705 the OMZ agrees with the Cd fluxes determined from benthic chamber incubations,
706 where dissolved Cd removal takes place in the 20 – 30 cm of overlying water above
707 the seafloor. These Cd removal fluxes from benthic chambers alone are sufficient to
708 account for 41 % – 68 % of the estimated particulate Cd removal from the water column
709 and 38 % – 60 % of total Cd enrichment in the sediment within the OMZ (Table 3).
710 Considering that Cd precipitation in near-bottom water is unlikely to be restricted to the
711 20 – 30 cm above the seafloor, covered by our benthic chambers, the removal flux
712 associated with this process is likely to be even higher. At Station 1, where the surface
713 sediment below the benthic chamber was highly sulfidic, the particulate Cd removal
714 calculated from the concentration difference between the bottom water (0.5 m) and the
715 first sample from the benthic chamber incubation (taken after 0.25 h) was high enough
716 to explain the total Cd enrichment in the sediment.

717 Below the OMZ, at Station 9, where the smallest Cd enrichment was observed, the
718 relative contribution of Cd delivery with organic material increases. About half of the
719 Cd enrichment can be attributed to organic material at this station.

720 Once Cd is delivered to the sediment, it can either stay fixed in the solid phase
721 or be released to the pore waters. Cadmium concentrations in pore waters of
722 subsurface sediments (> 10 cm sediment depth) were mostly higher than bottom water
723 concentrations (Fig. 6), indicating a transfer of Cd from the solid phase into pore waters
724 during early diagenesis. Cadmium sulfides are considered highly insoluble and stable
725 within sediments (Elderfield et al., 1981), even upon re-oxygenation (Rosenthal et al.,
726 1995). Therefore, Cd release through re-dissolution of CdS is ruled out as a potential
727 source of dissolved Cd. Alternatively, Cd liberation upon remineralization of organic
728 material could explain elevated Cd concentrations in the pore water. Elevated Cd
729 concentrations in sulfidic pore waters have been observed in previous studies and
730 attributed to Cd stabilization through formation of organic and inorganic complexes
731 (Gobeil et al., 1987; Sundby et al., 2004). Experimental data gave evidence for the
732 presence of dissolved Cd bisulfide and polysulfide complexes in pore waters. An
733 increase of electrochemically active Cd after UV irradiation, was explained by the
734 destruction of electrochemically inactive bisulfide and polysulfide complexes (Gobeil et
735 al., 1987). At very high H₂S concentrations (> 10⁻³ M) the solubility of Cd may increase
736 due to an increase in these bisulfide and polysulfide complexes. Under such highly
737 sulfidic conditions, Cd solubility may even exceed the solubility in oxygenated waters
738 and highly sulfidic sediment can eventually lead to a diffusive source of Cd to the
739 bottom water (Davies-Colley et al., 1985). Such a scenario may explain the negative
740 (i.e., upward-directed) diffusive Cd flux at Station 1, where the pore waters of surface
741 sediments are highly sulfidic.

742

743

744 **5. Conclusions and implications for trace metal sources and sinks in the future** 745 **ocean**

746 Consistent with earlier work, our results demonstrate that that OMZ sediments
747 are a source for Fe and a sink for Cd. Moreover, based on our findings, biogeochemical
748 conditions and processes that control the benthic fluxes of these TM across the
749 Peruvian OMZ can be further constrained.

750 Within the OMZ, where bottom dwelling macrofauna is absent, diffusion is the
751 main process that transports Fe from the sediment pore water into the bottom water.

752 The accumulation of high levels of H₂S in pore waters, modulated by the abundance
753 of sulfur oxidizing bacteria, can reduce diffusive Fe release through sulfide precipitation
754 within pore waters. In anoxic bottom waters Fe can be rapidly removed, likely via
755 oxidation with NO₂⁻ and/or interaction with particles. Benthic Cd fluxes are directed
756 from the bottom water into the sediment within the OMZ. Diffusive fluxes and delivery
757 of Cd via organic material cannot account for the sedimentary Cd enrichment. Instead
758 CdS precipitation in near-bottom waters could be the most important pathway that
759 delivers Cd to the sediments.

760 According to our results, H₂S concentrations in surface sediments exert a first
761 order control on the magnitude and direction of Fe and Cd fluxes across the sediment-
762 water interface. With generally decreasing oxygen concentrations in the ocean and an
763 expansion of OMZs (Stramma et al., 2008; Schmidtko et al., 2017), sulfidic surface
764 sediments will likely also expand. With regard to the solubility of their sulfide minerals,
765 Fe and Cd represent two opposite end members. The solubility of sulfide minerals of
766 other important nutrient-type TMs, such as Ni and Zn, is intermediate between those
767 of Fe and Cd (Fe > Ni > Zn > Cd). An expansion of sulfidic surface sediments is thus
768 likely to affect sedimentary TM fluxes in a differing manner. This notion is illustrated in
769 Fig. 10, showing saturation indices calculated based on the range of TM
770 concentrations observed in the ocean and typical H₂S concentrations observed in
771 anoxic marine environments (nM – μM concentrations represent sulfidic events in the
772 water column; μM – mM concentrations are typical for pore waters). Cadmium sulfide
773 minerals become oversaturated at nM to μM H₂S concentrations, which explains why
774 Cd removal can take place in the bottom water in OMZs. By contrast, FeS is highly
775 undersaturated under the typical biogeochemical conditions in the water column.
776 Therefore, FeS precipitation is unlikely to take place in the water column, even under
777 somewhat more reducing conditions. Other sulfide-forming TMs have an intermediate
778 sulfide solubility (e.g. Zn, Ni), which could imply that the direction and magnitude of
779 their sedimentary fluxes is susceptible to expanding ocean anoxia. The differing
780 response of TMs to an expansion of sulfidic conditions may cause a change in the TM
781 stoichiometry of upwelling water masses with potential consequences for TM-
782 dependent marine ecosystems in surface waters.

783

784

785 **Data availability**

786 The data will be made available at Pangaea upon publication of the article.

787

788

789 **Author contribution**

790 AP and FS conceived the study. AP, FS, AD, SS conducted the sampling at sea. AP
791 analyzed the trace metal concentrations. AP and FS prepared the manuscript with
792 contributions from all co-authors.

793

794

795 **Competing Interests**

796 The authors declare that they have no conflict of interest.

797

798

799 **Acknowledgements**

800 We are grateful for the support of the crew of RV Meteor during the fieldwork. For their
801 technical and analytical assistance we thank A. Beck, A. Bleyer, B. Domeyer, D.
802 Jasinski, A. Petersen, T. Steffens, R. Surberg and M. Türk. This study was supported
803 by the German Research Foundation through the Emmy Noether
804 Nachwuchsforschergruppe ICONOX (Iron Cycling in Continental Margin Sediments
805 and the Nutrient and Oxygen Balance of the Ocean) and Sonderforschungsbereich
806 754 (Climate-Biogeochemistry Interactions in the Tropical Ocean). We also would like
807 to thank Edouard Metzger and Michael Staubwasser for their constructive reviews, as
808 well as S. Wajih A. Naqvi for the editorial handling.

809 **References**

810

811 Audry, S., Blanc, G., Schäfer, J., Chaillou, G. and Robert, S.: Early diagenesis of
812 trace metals (Cd, Cu, Co, Ni, U, Mo, and V) in the freshwater reaches of a macrotidal
813 estuary, *Geochim. Cosmochim. Acta*, 70(9), 2264–2282,
814 doi:10.1016/j.gca.2006.02.001, 2006.

815 Ball, J. W. and Nordstrom, D. K.: WATEQ4F -- User's manual with revised
816 thermodynamic data base and test cases for calculating speciation of major, trace
817 and redox elements in natural waters, US Geol. Surv., (Open-File Rep.), 91–183,
818 doi:10.3133/ofr90129, 1991.

819 Bianchi, D., Weber, T. S., Kiko, R. and Deutsch, C.: Global niche of marine anaerobic
820 metabolisms expanded by particle microenvironments, *Nat. Geosci.*, 11(April), 1–6,
821 doi:10.1038/s41561-018-0081-0, 2018.

822 Biller, D. V. and Bruland, K. W.: Sources and distributions of Mn, Fe, Co, Ni, Cu, Zn,
823 and Cd relative to macronutrients along the central California coast during the spring
824 and summer upwelling season, *Mar. Chem.*, 155, 50–70,
825 doi:10.1016/j.marchem.2013.06.003, 2013.

826 Böning, P., Brumsack, H. J., Böttcher, M. E., Schnetger, B., Kriete, C., Kallmeyer, J.
827 and Borchers, S. L.: Geochemistry of Peruvian near-surface sediments, *Geochim.
828 Cosmochim. Acta*, 68(21), 4429–4451, doi:10.1016/j.gca.2004.04.027, 2004.

829 Bopp, L., Le Quéré, C., Heimann, M., Manning, A. C. and Monfray, P.: Climate-
830 induced oceanic oxygen fluxes: Implications for the contemporary carbon budget,
831 *Global Biogeochem. Cycles*, 16(2), 6-1-6–13, doi:10.1029/2001GB001445, 2002.

832 Borchers, S. L., Schnetger, B., Böning, P. and Brumsack, H.-J.: Geochemical
833 signatures of the Namibian diatom belt: Perennial upwelling and intermittent anoxia,
834 *Geochemistry, Geophys. Geosystems*, 6(6), doi:10.1029/2004GC000886, 2005.

835 Boudreau, B. P.: *Diagenetic Models and Their Implementation*, Springer., 1997.

836 Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nat.
837 Geosci.*, 3(10), 675–682, doi:10.1038/ngeo964, 2010.

838 Bruland, K. W. and Lohan, M. C.: Controls of Trace Metals in Seawater, in *Treatise*

839 on Geochemistry, pp. 23–47, Elsevier., 2003.

840 Brumsack, H. J.: The trace metal content of recent organic carbon-rich sediments:
841 Implications for Cretaceous black shale formation, *Palaeogeogr. Palaeoclimatol.*
842 *Palaeoecol.*, 232(2–4), 344–361, doi:10.1016/j.palaeo.2005.05.011, 2006.

843 Canfield, D. E.: Reactive iron in marine sediments, *Geochim. Cosmochim. Acta*,
844 53(3), 619–632, doi:10.1016/0016-7037(89)90005-7, 1989.

845 Carlson, H. K., Clark, I. C., Blazewicz, S. J., Iavarone, A. T. and Coates, J. D.: Fe(II)
846 Oxidation Is an Innate Capability of Nitrate-Reducing Bacteria That Involves Abiotic
847 and Biotic Reactions, *J. Bacteriol.*, 195(14), 3260–3268, doi:10.1128/JB.00058-13,
848 2013.

849 Ciceri, G., Maran, C., Martinotti, W. and Queirazza, G.: Geochemical cycling of heavy
850 metals in a marine coastal area: benthic flux determination from pore water profiles
851 and in situ measurements using benthic chambers, *Hydrobiologia*, 235–236(1), 501–
852 517, doi:10.1007/BF00026238, 1992.

853 Colbert, D., Coale, K. ., Berelson, W. . and Johnson, K. .: Cadmium Flux in Los
854 Angeles/Long Beach Harbours and at Sites along the California Continental Margin,
855 *Estuar. Coast. Shelf Sci.*, 53(2), 169–180, doi:10.1006/ecss.2001.0802, 2001.

856 Collier, R. and Edmond, J.: The trace element geochemistry of marine biogenic
857 particulate matter, *Prog. Oceanogr.*, 13(2), 113–199, doi:10.1016/0079-
858 6611(84)90008-9, 1984.

859 Conway, T. M. and John, S. G.: Quantification of dissolved iron sources to the North
860 Atlantic Ocean, *Nature*, 511(7508), 212–215, doi:10.1038/nature13482, 2014.

861 Conway, T. M. and John, S. G.: Biogeochemical cycling of cadmium isotopes along a
862 high-resolution section through the North Atlantic Ocean, *Geochim. Cosmochim.*
863 *Acta*, 148, 269–283, doi:10.1016/j.gca.2014.09.032, 2015a.

864 Conway, T. M. and John, S. G.: The cycling of iron, zinc and cadmium in the North
865 East Pacific Ocean - Insights from stable isotopes, *Geochim. Cosmochim. Acta*, 164,
866 262–283, doi:10.1016/j.gca.2015.05.023, 2015b.

867 Dale, A. W., Nickelsen, L., Scholz, F., Hensen, C., Oschlies, A. and Wallmann, K.: A
868 revised global estimate of dissolved iron fluxes from marine sediments, *Global*

869 Biogeochem. Cycles, 29(5), 691–707, doi:10.1002/2014GB005017, 2015a.

870 Dale, A. W., Sommer, S., Lomnitz, U., Montes, I., Treude, T., Liebetrau, V., Gier, J.,
871 Hensen, C., Dengler, M., Stolpovsky, K., Bryant, L. D. and Wallmann, K.: Organic
872 carbon production, mineralisation and preservation on the Peruvian margin,
873 Biogeosciences, 12(5), 1537–1559, doi:10.5194/bg-12-1537-2015, 2015b.

874 Dalsgaard, T., Thamdrup, B., Farías, L. and Revsbech, N. P.: Anammox and
875 denitrification in the oxygen minimum zone of the eastern South Pacific, Limnol.
876 Oceanogr., 57(5), 1331–1346, doi:10.4319/lo.2012.57.5.1331, 2012.

877 Davies-Colley, R. J., Nelson, P. O. and Williamson, K. J.: Sulfide control of cadmium
878 and copper concentrations in anaerobic estuarine sediments, Mar. Chem., 16(2),
879 173–186, doi:10.1016/0304-4203(85)90021-0, 1985.

880 Echevin, V., Colas, F., Espinoza-Morriberon, D., Vasquez, L., Anculle, T. and
881 Gutierrez, D.: Forcings and Evolution of the 2017 Coastal El Niño Off Northern Peru
882 and Ecuador, Front. Mar. Sci., 5(October), 1–16, doi:10.3389/fmars.2018.00367,
883 2018.

884 Ehlert, C., Doering, K., Wallmann, K., Scholz, F., Sommer, S., Grasse, P., Geilert, S.
885 and Frank, M.: Stable silicon isotope signatures of marine pore waters – Biogenic
886 opal dissolution versus authigenic clay mineral formation, Geochim. Cosmochim.
887 Acta, 191, 102–117, doi:10.1016/j.gca.2016.07.022, 2016.

888 Elderfield, H., McCaffrey, R. J., Luedtke, N., Bender, M. and Truesdale, V. W.:
889 Chemical diagenesis in Narragansett Bay sediments, Am. J. Sci., 281(8), 1021–1055,
890 doi:10.2475/ajs.281.8.1021, 1981.

891 Elrod, V. A., Berelson, W. M., Coale, K. H. and Johnson, K. S.: The flux of iron from
892 continental shelf sediments: A missing source for global budgets, Geophys. Res.
893 Lett., 31(12), n/a-n/a, doi:10.1029/2004GL020216, 2004.

894 Fitzsimmons, J. N., Conway, T. M., Lee, J.-M., Kayser, R., Thyng, K. M., John, S. G.
895 and Boyle, E. A.: Dissolved iron and iron isotopes in the southeastern Pacific Ocean,
896 Global Biogeochem. Cycles, 30(10), 1372–1395, doi:10.1002/2015GB005357, 2016.

897 Friedrich, J., Dinkel, C., Friedl, G., Pimenov, N., Wijsman, J., Gomoiu, M.-T.,
898 Cociasu, A., Popa, L. and Wehrli, B.: Benthic Nutrient Cycling and Diagenetic
899 Pathways in the North-western Black Sea, Estuar. Coast. Shelf Sci., 54(3), 369–383,

900 doi:10.1006/ecss.2000.0653, 2002.

901 Garreaud, R. D.: A plausible atmospheric trigger for the 2017 coastal El Niño, *Int. J.*
902 *Climatol.*, 38(January 2017), e1296–e1302, doi:10.1002/joc.5426, 2018.

903 Gendron, A., Silverberg, N., Sundby, B. and Lebel, J.: Early diagenesis of cadmium
904 and cobalt in sediments of the Laurentian Trough, *Geochim. Cosmochim. Acta*,
905 50(5), 741–747, doi:10.1016/j.ijmachtools.2007.10.013, 1986.

906 Gerringa, L. J. A.: Aerobic degradation of organic matter and the mobility of Cu, Cd,
907 Ni, Pb, Zn, Fe and Mn in marine sediment slurries, *Mar. Chem.*, 29(C), 355–374,
908 doi:10.1016/0304-4203(90)90023-6, 1990.

909 Gobeil, C., Silverberg, N., Sundby, B. and Cossa, D.: Cadmium diagenesis in
910 Laurentian Trough sediments, *Geochim. Cosmochim. Acta*, 51(3), 589–596,
911 doi:10.1016/0016-7037(87)90071-8, 1987.

912 Graco, M. I., Purca, S., Dewitte, B., Castro, C. G., Morón, O., Ledesma, J., Flores, G.
913 and Gutiérrez, D.: The OMZ and nutrient features as a signature of interannual and
914 low-frequency variability in the Peruvian upwelling system, *Biogeosciences*, 14(20),
915 4601–4617, doi:10.5194/bg-14-4601-2017, 2017.

916 Grasshoff, M., Erhardt, M. and Kremling, K.: *Methods of seawater analysis.*, Wiley-
917 VCH, Weinheim, doi:10.1002/ange.19770890738, 1999.

918 Gutiérrez, D., Enríquez, E., Purca, S., Quipúzcoa, L., Marquina, R., Flores, G. and
919 Graco, M.: Oxygenation episodes on the continental shelf of central Peru: Remote
920 forcing and benthic ecosystem response, *Prog. Oceanogr.*, 79(2–4), 177–189,
921 doi:10.1016/j.pocean.2008.10.025, 2008.

922 Hawco, N. J., Ohnemus, D. C., Resing, J. A., Twining, B. S. and Saito, M. A.: A
923 dissolved cobalt plume in the oxygen minimum zone of the eastern tropical South
924 Pacific, *Biogeosciences*, 13(20), 5697–5717, doi:10.5194/bg-13-5697-2016, 2016.

925 Heller, M. I., Lam, P. J., Moffett, J. W., Till, C. P., Lee, J. M., Toner, B. M. and
926 Marcus, M. A.: Accumulation of Fe oxyhydroxides in the Peruvian oxygen deficient
927 zone implies non-oxygen dependent Fe oxidation, *Geochim. Cosmochim. Acta*, 211,
928 174–193, doi:10.1016/j.gca.2017.05.019, 2017.

929 Helm, K. P., Bindoff, N. L. and Church, J. A.: Observed decreases in oxygen content

930 of the global ocean, *Geophys. Res. Lett.*, 38(23), 1–6, doi:10.1029/2011GL049513,
931 2011.

932 Homoky, W. B., Severmann, S., McManus, J., Berelson, W. M., Riedel, T. E.,
933 Statham, P. J. and Mills, R. A.: Dissolved oxygen and suspended particles regulate
934 the benthic flux of iron from continental margins, *Mar. Chem.*, 134–135, 59–70,
935 doi:10.1016/j.marchem.2012.03.003, 2012.

936 Hydes, D., Aoyama, M., Aminot, A., Bakker, K., Becker, S., Coverly, S., Daniel, A.,
937 Dickson, A. G., Grosso, O., Kerouel, R., van Ooijen, J., Sato, K., Tanhua, T.,
938 Woodward, E. M. S. and Zhang, J. Z.: Determination of dissolved nutrients (N, P, Si)
939 in seawater with high precision and inter-comparability using gas-segmented
940 continuous flow analysers, *Go-sh. Repeat Hydrogr. Man. IOCCP Rep. A Collect.*
941 *Expert Reports Guidel.*, 134(14), 1–87 [online] Available from:
942 <http://archimer.ifremer.fr/doc/00020/13141/>, 2010.

943 Janssen, D. J., Conway, T. M., John, S. G., Christian, J. R., Kramer, D. I., Pedersen,
944 T. F. and Cullen, J. T.: Undocumented water column sink for cadmium in open ocean
945 oxygen-deficient zones, *Proc. Natl. Acad. Sci.*, 111(19), 6888–6893,
946 doi:10.1073/pnas.1402388111, 2014.

947 John, S. G., Helgoe, J., Townsend, E., Weber, T., DeVries, T., Tagliabue, A., Moore,
948 K., Lam, P., Marsay, C. M. and Till, C.: Biogeochemical cycling of Fe and Fe stable
949 isotopes in the Eastern Tropical South Pacific, *Mar. Chem.*, 201(March), 66–76,
950 doi:10.1016/j.marchem.2017.06.003, 2018.

951 Karstensen, J., Stramma, L. and Visbeck, M.: Oxygen minimum zones in the eastern
952 tropical Atlantic and Pacific oceans, *Prog. Oceanogr.*, 77(4), 331–350,
953 doi:10.1016/j.pocean.2007.05.009, 2008.

954 Keeling, R. F., Körtzinger, A. and Gruber, N.: Ocean Deoxygenation in a Warming
955 World, *Ann. Rev. Mar. Sci.*, 2(1), 199–229,
956 doi:10.1146/annurev.marine.010908.163855, 2010.

957 Klar, J. K., Schlosser, C., Milton, J. A., Woodward, E. M. S., Lacan, F., Parkinson, I.
958 J., Achterberg, E. P. and James, R. H.: Sources of dissolved iron to oxygen minimum
959 zone waters on the Senegalese continental margin in the tropical North Atlantic
960 Ocean: Insights from iron isotopes, *Geochim. Cosmochim. Acta*, 236, 60–78,

961 doi:10.1016/j.gca.2018.02.031, 2018.

962 Klinkhammer, G., Heggge, D. T. and Graham, D. W.: Metal diagenesis in oxic marine
963 sediments, *Earth Planet. Sci. Lett.*, 61(2), 211–219, doi:10.1016/0012-
964 821X(82)90054-1, 1982.

965 Klueglein, N. and Kappler, A.: Abiotic oxidation of Fe(II) by reactive nitrogen species
966 in cultures of the nitrate-reducing Fe(II) oxidizer *Acidovorax* sp. BoFeN1 - questioning
967 the existence of enzymatic Fe(II) oxidation, *Geobiology*, 11(2), 180–190,
968 doi:10.1111/gbi.12019, 2013.

969 Lam, P. and Kuypers, M. M. M.: Microbial Nitrogen Cycling Processes in Oxygen
970 Minimum Zones, *Ann. Rev. Mar. Sci.*, 3(1), 317–345, doi:10.1146/annurev-marine-
971 120709-142814, 2011.

972 Lam, P., Lavik, G., Jensen, M. M., van de Vossenberg, J., Schmid, M., Woebken, D.,
973 Gutierrez, D., Amann, R., Jetten, M. S. M. and Kuypers, M. M. M.: Revising the
974 nitrogen cycle in the Peruvian oxygen minimum zone, *Proc. Natl. Acad. Sci.*, 106(12),
975 4752–4757, doi:10.1073/pnas.0812444106, 2009.

976 Lane, T. W. and Morel, F. M. M.: A biological function for cadmium in marine diatoms,
977 *Proc. Natl. Acad. Sci.*, 97(9), 4627–4631, doi:10.1073/pnas.090091397, 2000.

978 Lee, J. and Morel, F.: Replacement of zinc by cadmium in marine phytoplankton,
979 *Mar. Ecol. Prog. Ser.*, 127(1–3), 305–309, doi:10.3354/meps127305, 1995.

980 Lenstra, W. K., Hermans, M., Séguret, M. J. M., Witbaard, R., Behrends, T., Dijkstra,
981 N., van Helmond, N. A. G. M., Kraal, P., Laan, P., Rijkenberg, M. J. A., Severmann,
982 S., Teacă, A. and Slomp, C. P.: The shelf-to-basin iron shuttle in the Black Sea
983 revisited, *Chem. Geol.*, 511(April), 314–341, doi:10.1016/j.chemgeo.2018.10.024,
984 2019.

985 Levin, L., Gutiérrez, D., Rathburn, A., Neira, C., Sellanes, J., Muñoz, P., Gallardo, V.
986 and Salamanca, M.: Benthic processes on the Peru margin: a transect across the
987 oxygen minimum zone during the 1997–98 El Niño, *Prog. Oceanogr.*, 53(1), 1–27,
988 doi:10.1016/S0079-6611(02)00022-8, 2002.

989 Li, Y.-H. and Gregory, S.: Diffusion of ions in sea water and in deep-sea sediments,
990 *Geochim. Cosmochim. Acta*, 38(5), 703–714, doi:10.1016/0016-7037(74)90145-8,
991 1974.

992 Little, S. H., Vance, D., Lyons, T. W. and McManus, J.: Controls on trace metal
993 authigenic enrichment in reducing sediments: Insights from modern oxygen-deficient
994 settings, *Am. J. Sci.*, 315(2), 77–119, doi:10.2475/02.2015.01, 2015.

995 Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77(1), 43–54,
996 doi:10.1016/S0304-4203(01)00074-3, 2002.

997 Lohan, M. C. and Bruland, K. W.: Elevated Fe(II) and dissolved Fe in hypoxic shelf
998 waters off Oregon and Washington: An enhanced source of iron to coastal upwelling
999 regimes, *Environ. Sci. Technol.*, 42(17), 6462–6468, doi:10.1021/es800144j, 2008.

1000 Metzger, E., Simonucci, C., Viollier, E., Sarazin, G., Prévot, F., Elbaz-Poulichet, F.,
1001 Seidel, J.-L. and Jézéquel, D.: Influence of diagenetic processes in Thau lagoon on
1002 cadmium behavior and benthic fluxes, *Estuar. Coast. Shelf Sci.*, 72(3), 497–510,
1003 doi:10.1016/j.ecss.2006.11.016, 2007.

1004 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W.,
1005 Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J.,
1006 Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka,
1007 T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A. and Ulloa, O.: Processes
1008 and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6(9), 701–710,
1009 doi:10.1038/ngeo1765, 2013.

1010 Morel, F. M. M., Milligan, A. J. and Saito, M. A.: *Marine Bioinorganic Chemistry: The*
1011 *Role of Trace Metals in the Oceanic Cycles of Major Nutrients*, in *Treatise on*
1012 *Geochemistry*, vol. 197, pp. 123–150, Elsevier., 2014.

1013 Morse, J. W. and Luther, G. W.: Chemical influences on trace metal-sulfide
1014 interactions in anoxic sediments, *Geochim. Cosmochim. Acta*, 63(19–20), 3373–
1015 3378, doi:10.1016/S0016-7037(99)00258-6, 1999.

1016 Muñoz, P., Dezileau, L., Cardenas, L., Sellanes, J., Lange, C. B., Inostroza, J.,
1017 Muratli, J. and Salamanca, M. A.: Geochemistry of trace metals in shelf sediments
1018 affected by seasonal and permanent low oxygen conditions off central Chile, SE
1019 Pacific (~36°S), *Cont. Shelf Res.*, 33, 51–68, doi:10.1016/j.csr.2011.11.006, 2012.

1020 Noble, A. E., Lamborg, C. H., Ohnemus, D. C., Lam, P. J., Goepfert, T. J., Measures,
1021 C. I., Frame, C. H., Casciotti, K. L., DiTullio, G. R., Jennings, J. and Saito, M. A.:
1022 Basin-scale inputs of cobalt, iron, and manganese from the Benguela-Angola front to

1023 the South Atlantic Ocean, *Limnol. Oceanogr.*, 57(4), 989–1010,
1024 doi:10.4319/lo.2012.57.4.0989, 2012.

1025 Noffke, A., Hensen, C., Sommer, S., Scholz, F., Bohlen, L., Mosch, T., Graco, M. and
1026 Wallmann, K.: Benthic iron and phosphorus fluxes across the Peruvian oxygen
1027 minimum zone, *Limnol. Oceanogr.*, 57(3), 851–867, doi:10.4319/lo.2012.57.3.0851,
1028 2012.

1029 Ohde, T.: Coastal Sulfur Plumes off Peru During El Niño, La Niña, and Neutral
1030 Phases, *Geophys. Res. Lett.*, 45(14), 7075–7083, doi:10.1029/2018GL077618, 2018.

1031 Olson, L., Quinn, K. A., Siebecker, M. G., Luther, G. W., Hastings, D. and Morford, J.
1032 L.: Trace metal diagenesis in sulfidic sediments: Insights from Chesapeake Bay,
1033 *Chem. Geol.*, 452, 47–59, doi:10.1016/j.chemgeo.2017.01.018, 2017.

1034 Oschlies, A., Schulz, K. G., Riebesell, U. and Schmittner, A.: Simulated 21st
1035 century's increase in oceanic suboxia by CO₂-enhanced biotic carbon export, *Global*
1036 *Biogeochem. Cycles*, 22(4), 1–10, doi:10.1029/2007GB003147, 2008.

1037 Peng, Q., Xie, S.-P., Wang, D., Zheng, X.-T. and Zhang, H.: Coupled ocean-
1038 atmosphere dynamics of the 2017 extreme coastal El Niño, *Nat. Commun.*, 10(1),
1039 298, doi:10.1038/s41467-018-08258-8, 2019.

1040 Pennington, J. T., Mahoney, K. L., Kuwahara, V. S., Kolber, D. D., Calienes, R. and
1041 Chavez, F. P.: Primary production in the eastern tropical Pacific: A review, *Prog.*
1042 *Oceanogr.*, 69(2–4), 285–317, doi:10.1016/j.pocean.2006.03.012, 2006.

1043 Point, D., Monperrus, M., Tessier, E., Amouroux, D., Chauvaud, L., Thouzeau, G.,
1044 Jean, F., Amice, E., Grall, J., Leynaert, A., Clavier, J. and Donard, O. F. X.: Biological
1045 control of trace metal and organometal benthic fluxes in a eutrophic lagoon (Thau
1046 Lagoon, Mediterranean Sea, France), *Estuar. Coast. Shelf Sci.*, 72(3), 457–471,
1047 doi:10.1016/j.ecss.2006.11.013, 2007.

1048 Price, N. M. and Morel, F. M. M.: Cadmium and cobalt substitution for zinc in a
1049 marine diatom, *Nature*, 344(6267), 658–660, doi:10.1038/344658a0, 1990.

1050 Ragueneau, O., Tréguer, P., Leynaert, A., Anderson, R. F., Brzezinski, M. A.,
1051 DeMaster, D. J., Dugdale, R. C., Dymond, J., Fischer, G., François, R., Heinze, C.,
1052 Maier-Reimer, E., Martin-Jézéquel, V., Nelson, D. M. and Quéguiner, B.: A review of
1053 the Si cycle in the modern ocean: Recent progress and missing gaps in the

1054 application of biogenic opal as a paleoproductivity proxy, *Glob. Planet. Change*,
1055 26(4), 317–365, doi:10.1016/S0921-8181(00)00052-7, 2000.

1056 Raiswell, R. and Canfield, D. E.: The Iron Biogeochemical Cycle Past and Present,
1057 *Geochemical Perspect.*, 1(1), 1–220, doi:10.7185/geochempersp.1.1, 2012.

1058 Rapp, I., Schlosser, C., Rusiecka, D., Gledhill, M. and Achterberg, E. P.: Automated
1059 preconcentration of Fe, Zn, Cu, Ni, Cd, Pb, Co, and Mn in seawater with analysis
1060 using high-resolution sector field inductively-coupled plasma mass spectrometry,
1061 *Anal. Chim. Acta*, 976, 1–13, doi:10.1016/j.aca.2017.05.008, 2017.

1062 Rapp, I., Schlosser, C., Menzel Barraqueta, J.-L., Wenzel, B., Lüdke, J., Scholten, J.,
1063 Gasser, B., Reichert, P., Gledhill, M., Dengler, M. and Achterberg, E. P.: Controls on
1064 redox-sensitive trace metals in the Mauritanian oxygen minimum zone,
1065 *Biogeosciences Discuss.*, (November), 1–49, doi:10.5194/bg-2018-472, 2018.

1066 Rickard, D., Griffith, A., Oldroyd, A., Butler, I. B., Lopez-Capel, E., Manning, D. A. C.
1067 and Apperley, D. C.: The composition of nanoparticulate mackinawite, tetragonal
1068 iron(II) monosulfide, *Chem. Geol.*, 235(3–4), 286–298,
1069 doi:10.1016/j.chemgeo.2006.07.004, 2006.

1070 Rigaud, S., Radakovitch, O., Couture, R. M., Deflandre, B., Cossa, D., Garnier, C.
1071 and Garnier, J. M.: Mobility and fluxes of trace elements and nutrients at the
1072 sediment-water interface of a lagoon under contrasting water column oxygenation
1073 conditions, *Appl. Geochemistry*, 31(April 2013), 35–51,
1074 doi:10.1016/j.apgeochem.2012.12.003, 2013.

1075 Rosenthal, Y., Lam, P., Boyle, E. A. and Thomson, J.: Precipitation and
1076 postdepositional mobility, *Earth Planet. Sci. Lett.*, 132, 99–111, doi:10.1016/0012-
1077 821X(95)00056-I, 1995.

1078 Rue, E. L. and Bruland, K. W.: The role of organic complexation on ambient iron
1079 chemistry in the equatorial Pacific Ocean and the response of a mesoscale iron
1080 addition experiment, *Limnol. Oceanogr.*, 42(5), 901–910,
1081 doi:10.4319/lo.1997.42.5.0901, 1997.

1082 Saito, M. A., Goepfert, T. J. and Ritt, J. T.: Some thoughts on the concept of
1083 colimitation: Three definitions and the importance of bioavailability, *Limnol.*
1084 *Oceanogr.*, 53(1), 276–290, doi:10.4319/lo.2008.53.1.0276, 2008.

1085 Schlosser, C., Streu, P., Frank, M., Lavik, G., Croot, P. L., Dengler, M. and
1086 Achterberg, E. P.: H₂S events in the Peruvian oxygen minimum zone facilitate
1087 enhanced dissolved Fe concentrations, *Sci. Rep.*, 8(1), 1–8, doi:10.1038/s41598-
1088 018-30580-w, 2018.

1089 Schmidtko, S., Stramma, L. and Visbeck, M.: Decline in global oceanic oxygen
1090 content during the past five decades, *Nature*, 542(7641), 335–339,
1091 doi:10.1038/nature21399, 2017.

1092 Scholz, F. and Neumann, T.: Trace element diagenesis in pyrite-rich sediments of the
1093 Achterwasser lagoon, SW Baltic Sea, *Mar. Chem.*, 107(4), 516–532,
1094 doi:10.1016/j.marchem.2007.08.005, 2007.

1095 Scholz, F., McManus, J., Mix, A. C., Hensen, C. and Schneider, R. R.: The impact of
1096 ocean deoxygenation on iron release from continental margin sediments, *Nat.*
1097 *Geosci.*, 7(6), 433–437, doi:10.1038/ngeo2162, 2014.

1098 Scholz, F., Löscher, C. R., Fiskal, A., Sommer, S., Hensen, C., Lomnitz, U., Wuttig,
1099 K., Göttlicher, J., Kossel, E., Steininger, R. and Canfield, D. E.: Nitrate-dependent
1100 iron oxidation limits iron transport in anoxic ocean regions, *Earth Planet. Sci. Lett.*,
1101 454, 272–281, doi:10.1016/j.epsl.2016.09.025, 2016.

1102 [Scholz, F., Schmidt, M., Hensen, C., Eroglu, S., Geilert, S., Gutjahr, M. and](#)
1103 [Liebetrau, V.: Shelf-to-basin iron shuttle in the Guaymas Basin, Gulf of California,](#)
1104 [Geochim. Cosmochim. Acta, 261, 76–92, doi:10.1016/j.gca.2019.07.006, 2019.](#)

1105 Schunck, H., Lavik, G., Desai, D. K., Großkopf, T., Kalvelage, T., Löscher, C. R.,
1106 Paulmier, A., Contreras, S., Siegel, H., Holtappels, M., Rosenstiel, P., Schilhabel, M.
1107 B., Graco, M., Schmitz, R. A., Kuypers, M. M. M. and LaRoche, J.: Giant Hydrogen
1108 Sulfide Plume in the Oxygen Minimum Zone off Peru Supports
1109 Chemolithoautotrophy, *PLoS One*, 8(8), doi:10.1371/journal.pone.0068661, 2013.

1110 Scor Working Group: GEOTRACES – An international study of the global marine
1111 biogeochemical cycles of trace elements and their isotopes, *Geochemistry*, 67(2),
1112 85–131, doi:10.1016/j.chemer.2007.02.001, 2007.

1113 Severmann, S., McManus, J., Berelson, W. M. and Hammond, D. E.: The continental
1114 shelf benthic iron flux and its isotope composition, *Geochim. Cosmochim. Acta*,
1115 74(14), 3984–4004, doi:10.1016/j.gca.2010.04.022, 2010.

- 1116 ~~Shibamoto, Y. and Harada, K.: Silicon flux and distribution of biogenic silica in deep-~~
1117 ~~sea sediments in the western North Pacific Ocean, Deep Sea Res. Part I Oceanogr.~~
1118 ~~Res. Pap., 57(2), 163–174, doi:10.1016/j.dsr.2009.10.009, 2010.~~
- 1119 Sommer, S., Linke, P., Pfannkuche, O., Schleicher, T., Schneider v. D, D., Reitz, A.,
1120 Haeckel, M., Flögel, S. and Hensen, C.: Seabed methane emissions and the habitat
1121 of frenulate tubeworms on the Captain Arutyunov mud volcano (Gulf of Cadiz), Mar.
1122 Ecol. Prog. Ser., 382, 69–86, doi:10.3354/meps07956, 2009.
- 1123 Sommer, S., Gier, J., Treude, T., Lomnitz, U., Dengler, M., Cardich, J. and Dale, A.
1124 W.: Depletion of oxygen, nitrate and nitrite in the Peruvian oxygen minimum zone
1125 cause an imbalance of benthic nitrogen fluxes, Deep. Res. Part I Oceanogr. Res.
1126 Pap., 112(3), 113–122, doi:10.1016/j.dsr.2016.03.001, 2016.
- 1127 Stookey, L. L.: Ferrozine---a new spectrophotometric reagent for iron, Anal. Chem.,
1128 42(7), 779–781, doi:10.1021/ac60289a016, 1970.
- 1129 Stramma, L., Johnson, G. C., Sprintall, J. and Mohrholz, V.: Expanding Oxygen-
1130 Minimum Zones in the Tropical Oceans, Science (80-.), 320(5876), 655–658,
1131 doi:10.1126/science.1153847, 2008.
- 1132 Stramma, L., Schmidtko, S., Levin, L. A. and Johnson, G. C.: Ocean oxygen minima
1133 expansions and their biological impacts, Deep. Res. Part I Oceanogr. Res. Pap.,
1134 57(4), 587–595, doi:10.1016/j.dsr.2010.01.005, 2010.
- 1135 Straub, K. L., Benz, M., Schink, B. and Widdel, F.: Anaerobic, nitrate-dependent
1136 microbial oxidation of ferrous iron. Appl Environ Microbiol, Appl. Environ. Microbiol.,
1137 62(4), 1458–60, 1996.
- 1138 Sunda, W. G. and Huntsman, S. A.: Effect of Zn, Mn, and Fe on Cd accumulation in
1139 phytoplankton: Implications for oceanic Cd cycling, Limnol. Oceanogr., 45(7), 1501–
1140 1516, doi:10.4319/lo.2000.45.7.1501, 2000.
- 1141 Sundby, B., Martinez, P. and Gobeil, C.: Comparative geochemistry of cadmium,
1142 rhenium, uranium, and molybdenum in continental margin sediments, Geochim.
1143 Cosmochim. Acta, 68(11), 2485–2493, doi:10.1016/j.gca.2003.08.011, 2004.
- 1144 Taylor, S. R. and McLennan, S. M.: Planetary crusts: Their composition, origin and
1145 evolution , by Stuart Ross Taylor and Scott M. McLennan, Meteorit. Planet. Sci.,
1146 44(3), 465–466, doi:10.1111/j.1945-5100.2009.tb00744.x, 2009.

1147 Thamdrup, B., Dalsgaard, T. and Revsbech, N. P.: Widespread functional anoxia in
1148 the oxygen minimum zone of the Eastern South Pacific, *Deep Sea Res. Part I*
1149 *Oceanogr. Res. Pap.*, 65, 36–45, doi:10.1016/j.dsr.2012.03.001, 2012.

1150 Turetta, C., Capodaglio, G., Cairns, W., Rabar, S. and Cescon, P.: Benthic fluxes of
1151 trace metals in the lagoon of Venice, *Microchem. J.*, 79(1–2), 149–158,
1152 doi:10.1016/j.microc.2004.06.003, 2005.

1153 Twining, B. S. and Baines, S. B.: The Trace Metal Composition of Marine
1154 Phytoplankton, *Ann. Rev. Mar. Sci.*, 5(1), 191–215, doi:10.1146/annurev-marine-
1155 121211-172322, 2013.

1156 Vedamati, J., Goepfert, T. and Moffett, J. W.: Iron speciation in the eastern tropical
1157 south pacific oxygen minimum zone off peru, *Limnol. Oceanogr.*, 59(6), 1945–1957,
1158 doi:10.4319/lo.2014.59.6.1945, 2014.

1159 Westerlund, S. F. G., Anderson, L. G., Hall, P. O. J., Iverfeldt, Å., Van Der Loeff, M.
1160 M. R. and Sundby, B.: Benthic fluxes of cadmium, copper, nickel, zinc and lead in the
1161 coastal environment, *Geochim. Cosmochim. Acta*, 50(6), 1289–1296,
1162 doi:10.1016/0016-7037(86)90412-6, 1986.

1163 Xie, R. C., Rehkämper, M., Grasse, P., van de Flierdt, T., Frank, M. and Xue, Z.:
1164 Isotopic evidence for complex biogeochemical cycling of Cd in the eastern tropical
1165 South Pacific, *Earth Planet. Sci. Lett.*, 512, 134–146, doi:10.1016/j.epsl.2019.02.001,
1166 2019.

1167 Xu, Y., Feng, L., Jeffrey, P. D., Shi, Y. and Morel, F. M. M.: Structure and metal
1168 exchange in the cadmium carbonic anhydrase of marine diatoms, *Nature*, 452(7183),
1169 56–61, doi:10.1038/nature06636, 2008.

1170 Yücel, M., Sommer, S., Dale, A. W. and Pfannkuche, O.: Microbial sulfide filter along
1171 a benthic redox gradient in the Eastern Gotland Basin, Baltic Sea, *Front. Microbiol.*,
1172 8(FEB), 1–16, doi:10.3389/fmicb.2017.00169, 2017.

1173 Zago, C., Capodaglio, G., Ceradini, S., Ciceri, G., Abelmoschi, L., Soggia, F.,
1174 Cescon, P. and Scarponi, G.: Benthic fluxes of cadmium, lead, copper and nitrogen
1175 species in the northern Adriatic Sea in front of the River Po outflow, Italy, *Sci. Total*
1176 *Environ.*, 246(2–3), 121–137, doi:10.1016/S0048-9697(99)00421-0, 2000.

1177 Zumft, W. G.: Cell biology and molecular basis of denitrification., *Microbiol. Mol. Biol.*

1178 Rev., 61(4), 533–616 [online] Available from:
1179 <http://www.ncbi.nlm.nih.gov/pubmed/9409151>[http://www.pubmedcentral.nih.gov/](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC232623)
1180 [articlerender.fcgi?artid=PMC232623](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC232623), 1997.

1181

1213 **Figure captions**

1214

1215 Fig. 1: Sampling stations on the Peruvian continental margin during cruises M136 &
1216 M137 along a latitudinal depth transect at 12° S. The sampling stations for pore
1217 waters are depicted by white stars, for bottom waters by yellow dots and for benthic
1218 chamber incubations by red dots.

1219 Fig. 2: Oxygen, nitrate, nitrite and hydrogen sulfide concentrations on the Peruvian
1220 slope (Station 10, 1000 m depth), crossing the oxygen minimum zone (upper panel),
1221 and the upper shelf (Station 1, 75 m depth) (lower panel) during cruises M136 &
1222 M137 and M92 along the 12° S transect.

1223 Fig. 3: Near-bottom water concentrations of dissolved Fe and Cd and dissolved Fe to
1224 silicic acid ratios 0.5 m to 4 m above the seafloor across the 12° S transect. The red
1225 diamonds show results from a second sampling at Station 1 one month later.
1226 Concentrations of silicic acid are listed in Table S3 in the supplement.

1227 Fig. 4: Pore water dissolved Fe(II) and hydrogen sulfide concentrations. Data from
1228 an earlier cruise, M92, at Station 1 (75 m water depth) are displayed for comparison.
1229 The uppermost sample represents the bottom water concentration. The analytical
1230 error is smaller than the symbol size.

1231 Fig. 5: Dissolved Fe concentrations in incubated bottom waters from benthic chamber
1232 incubations. ~~The grey~~The black dashed line represents the linear regressions of the
1233 concentration change over the incubation time. The equations for these linear
1234 regressions are listed together with the coefficients of determination (R^2) in Table S4
1235 in the supplement. ~~The grey~~ dashed line represents the linear regressions of the
1236 ~~concentration change over the incubation time. The equations for these linear~~
1237 ~~regressions are listed together with the coefficients of determination (R^2) in Table S4~~
1238 ~~in the supplement.~~ The black dashed line represents theoretical concentration
1239 gradients over the incubation time based on our benthic diffusive fluxes (Table 2).
1240 The analytical error is smaller than the symbol size.

1241 Fig. 6: Pore water dissolved Cd and hydrogen sulfide concentrations. The uppermost
1242 sample represents the bottom water concentrations. The analytical error is smaller
1243 than the symbol size.

Formati

Formati

1270 Fig. 7: Dissolved Cd concentrations in incubated bottom waters from benthic
1271 chamber incubations. ~~The grey~~The black dashed line represents the linear
1272 regressions of the concentration change over the incubation time. The equations for
1273 these linear regressions are listed together with the coefficients of determination (R^2)
1274 in Table S4 in the supplement. ~~The grey~~ dashed line represents the linear
1275 ~~regressions of the concentration change over the incubation time. The equations for~~
1276 ~~these linear regressions are listed together with the coefficients of determination (R^2)~~
1277 ~~in Table S4 in the supplement.~~ The black dashed line represents theoretical
1278 concentration gradients over the incubation time based on our benthic diffusive fluxes
1279 (Table 3). The analytical error is smaller than the symbol size.

1280 Fig. 8: Dissolved Fe, nitrate and nitrite concentrations in incubated bottom waters
1281 from the benthic chamber incubation at Station 4 (145 m water depth).

1282 Fig. 9: Comparison of benthic diffusive Fe(II) fluxes between cruises M136 & M137
1283 and M92 on the Peruvian shelf. Negative values represent fluxes from the sediment
1284 pore water into the bottom waters. Shaded bars on the upper panel display the
1285 geochemical conditions in the water column during the time of sampling.

1286 Fig. 10: Schematic overview of the possible mobility of different trace metal to an
1287 expansion of sulfidic conditions. Saturation indices (SI) were calculated for different
1288 H_2S concentrations and reported minimum and maximum concentrations of trace
1289 metals in the water column (data from Bruland and Lohan 2003). Equilibrium
1290 constants (log K under standard conditions) for Fe (FeS ppt: -3.92), Ni (millerite: -
1291 8.04), Zn (sphalerite: -11.62) and Cd (greenokite: -15.93) were taken from the
1292 PHREEQC WATEQ4F database (Ball and Nordstrom, 1991). The results are
1293 approximate since concentrations instead of activities were used for calculations. A
1294 positive SI is indicative of oversaturation whereas a negative SI is indicative of
1295 undersaturation.

Formati

Formati

1307 Table 1: Accuracy of replicate concentration measurements (n = 7) of certified
 1308 reference seawater for trace metals NASS-7 and CASS-6 by ICP-MS.

	NASS-7 certified value	NASS-7 measured value	CASS-6 certified value	CASS-6 measured value
Fe (µg/L)	0.351 ± 0.026	0.352 ± 0.017	1.56 ± 0.12	1.56 ± 0.03
Cd (µg/L)	0.0161 ± 0.0016	0.0162 ± 0.0024	0.0217 ± 0.0018	0.0216 ± 0.0016

1309

1310

1311 Table 2: Comparison of benthic diffusive Fe(II) fluxes out of the sediment and
 1312 geochemical bottom water conditions between M136 & M137 and M92 on the
 1313 Peruvian shelf. Fluxes during M92 correspond to similar depth (see Fig. 9).

station	M136 & M137	M136 & M137	M136 & M137	M136 & M137	M136 & M137	M136 & M137	M92	M92
	water depth	latitude	longitude	water column condition	Fe(II) flux diffusive	Fe flux benthic chamber	water column condition	Fe(II) flux diffusive
	(m)	(S)	(W)		(mmol m ⁻² y ⁻¹)	(mmol m ⁻² y ⁻¹)		(mmol m ⁻² y ⁻¹)
1	75	12°13.52	77°10.93	O ₂ < 5 µM	-2.56	-1.74 ₋	slightly sulfidic	-22.69
3	130	12°16.68	77°14.95	nitrogenous	-0.81	-	slightly sulfidic	-3.16
4	145	12°18.71	77°17.80	nitrogenous	-17.45	-8.57 ± 2.18	nitrogenous	-5.77
5	195	12°21.50	77°21.70	nitrogenous	-2.49	2.01 ₋	nitrogenous	-1.51
6	245	12°23.30	77°24.82	nitrogenous	-7.96	-5.43 ± 2.36	nitrogenous	-10.20
9	750	12°31.35	77°35.01	O ₂ > 5 µM	0.00	-6.11 ± 3.12	O ₂ > 5 µM	0.00
10	970950	12°34.90	77°40.32	O ₂ > 5 µM	-0.26	-1.68 ₋	O ₂ > 5 µM	-0.12

1314

1315

1316

1317

1337 Table 3: Comparison of sedimentary Cd excess compared to the lithogenic
 1338 background and the contribution of Cd delivery to the sediment via different
 1339 pathways: (1) diffusion across the sediment-water interface and Cd sulfide
 1340 precipitation within the sediment; (2) Cd incorporation by phytoplankton and delivery
 1341 to the sediment with organic matter; (3) Cd sulfide precipitation in the water column
 1342 and particulate delivery to the sediment.

station	water depth (m)	Cd excess sediment ¹ ($\mu\text{mol m}^{-2} \text{y}^{-1}$)	(1) Cd flux diffusive ($\mu\text{mol m}^{-2} \text{y}^{-1}$)	Cd flux benthic chamber ($\mu\text{mol m}^{-2} \text{y}^{-1}$)	H ₂ S in surface sediment below benthic chamber (μM)	(2) Cd from organic matter ² ($\mu\text{mol m}^{-2} \text{y}^{-1}$)	(3) CdS precipitation in water column ³ ($\mu\text{mol m}^{-2} \text{y}^{-1}$)
1	75	248.87	-1.85	-1.6 (3109.5) ⁴	641.02	8.34 – 49.04	199.83 – 240.53
3	130	153.41	0.83	=	=	4.87 – 17.40	135.19 – 147.72
4	145	35.07	0.54	13.4 ± 1.05	1.30	1.55 – 6.48	28.07 – 32.99
5	195	44.76	0.63	22.6 ± 3.24	9.52	5.71 – 7.71	36.36 – 38.36
6	245	35.15	0.55	21.2 ± 3.31	0.40	3.60 – 6.54	28.06 – 31.00
9	750	4.44	-0.30	0.00 ± 0.02	0.00	1.48 – 3.21	1.23 – 2.96
10	950	=	=	=	0.00	=	=

1343
 1344 ¹ Calculated after Brumsack (2006) and multiplied by the mass accumulation rate for
 1345 each site (Dale et al., 2015b).

1346 ² Determined by multiplication of Cd/C ratio in average phytoplankton (Moore et al.,
 1347 2013) with particulate organic carbon rain rates (maximum values) and organic
 1348 carbon accumulation rates (minimum values) for each individual site (data from Dale
 1349 et al., 2015b).

1350 ³ Remainder of Cd excess in sediment after subtraction of diffusive and minimum and
 1351 maximum organic Cd sources.

1352 ⁴ Flux calculated from the concentration difference between the bottom water (0.5 m)
 1353 and the first sample from the benthic chamber incubation (taken after 0.25 h).

1354

1355

1356 Table 4: Modelled half-lives ($t_{1/2}$) of dissolved Fe within the first 4 m distance from the
 1357 seafloor at Stations 3 and 4 and data used for determination of $t_{1/2}$ using Eq. (4) and
 1358 Eq. (5).

station	water depth (m)	Si(OH) ₄ flux benthic chamber (F_{Si}) ($\mu\text{mol cm}^{-2} \text{d}^{-1}$)	Si(OH) ₄ concentration gradient (d_{Si}) ($\mu\text{mol cm}^{-3} \text{cm}^{-1}$)	eddy diffusion coefficient (K_y) ($\text{m}^2 \text{s}^{-1}$)	modelled Fe at sediment surface (C_{BW}) (nM)	Fe oxidation constant (k_{Feox}) (d^{-1})	half-life in near- bottom water column ($t_{1/2}$) (min)
3	130	0.73	$-4.05 \cdot 10^{-6}$	$1.55 \cdot 10^6$	70	400	2.5
4	145	0.33	$-1.44 \cdot 10^{-6}$	$1.96 \cdot 10^6$	81	3500	0.3

1359