

References

- Blackadar, A. K.: Boundary layer wind maxima and their significance for the growth of nocturnal inversions, *Bull. Amer. Meteorol. Soc.*, 38, 283-290, 1957.
- Hicks, B. B., N. S. Eash, D. L. O'Dell and N. S. Oetting, 2020. Augmented Bowen ratio analysis – I: Site adequacy, fetch and heat storage. *Agric. and Forest Meteorol.*, in review.
- Mahrt, L., Pfister, L. Thomas, C. K.: Small-scale variability in the nocturnal boundary layer. *Boundary-Layer Meteorol.*, 174, 81-98, 2020.
- O'Dell, D. L., Sauer, T. J., Hicks, B. B., Thierfelder, C., Lambert, D. M., Logan, J. and Eash, N. S.: A short-term assessment of carbon dioxide fluxes under contrasting agricultural and soil management practices in Zimbabwe, *J. Agric. Sci.*, 7, 32-48, 2015.
- O'Dell, D. L., Eash, N. S., Hicks, B. B., Oetting, J. N., Sauer, T. J., Lambert, D. M., Logan, J., Wright, W. C. and Zahn, J. A.: Reducing CO₂ flux by decreasing tillage in Ohio: overcoming conjecture with data, *J. Agric. Sci.*, 7, 1-15, 2018.