

Reply to the referees (see also the respective on-line Discussion for more detail)

Referee #1

1. The referee briefly remarks that the findings have little novelty. This may be the case with the conclusions, which emphasize the potential importance of acetate oxidation coupled either syntrophically to hydrogenotrophic methanogenesis or to the reduction of organic substances. Such conclusions have indeed been made before. However, to our knowledge they have so far not been supported by showing that the methyl group of acetate is indeed oxidized to CO₂ rather than reduced to CH₄. Here, we addressed this point by determining the fate of radioactively labeled acetate-methyl, and indeed found that in some of the lake sediments a large percentage of the acetate methyl was oxidized to CO₂. In our opinion, this is a crucial and novel experimental result.

2. The referee criticizes the presentation of the methods, in particular the determination of parameters. We have now presented the relevant methods in more detail including the equations used for calculation. We also better explain the determination of f_{H_2} as done in our previous publication (Ji et al. 2016) versus the present one, which used ¹³C versus ¹⁴C methods, respectively. We have now also added more explicit statements which incubation was used for which determination. This concerns in particular the determination of f_{H_2} and rates of CH₄ production on the one hand, and of RI and acetate turnover on the other hand. Finally, we have added to the Discussion by arguing about the use of different incubations and different incubation times for the individual parameters.

Minor issues:

General: We made several changes in the references as suggested by the referee.

Methods: The CH₄ production rates without addition of radiotracers (Fig. 1A) are basically those from our previous study (Ji et al., 2016), analogously to the values of f_{H_2} shown in Fig. 1B. This has now been explained in the Methods. It is also stated in the figure legend.

The addition of ¹⁴C was almost carrier free, i.e., the specific radioactivities of [2-¹⁴C]acetate and ¹⁴C-bicarbonate were on a range of about 50 Ci/mol. Thus, the addition of 1 μCi acetate-methyl was equivalent to an amount of about 20 nmol acetate. Hence, the addition of radioactive acetate was small (<5%) compared to the in-situ concentration of acetate. The same is true for bicarbonate. This information has been added to the Methods.

Fig. 2: The specific radioactivity is the relevant number for determining f_{H_2} . Therefore, we preferred showing these values instead of showing only Bq of CO₂ and CH₄ without reference to total CH₄ and CO₂.

L.32: ok

L.86: ok

L.106: There was a significant amount of radioactivity in the sediment carbonates, which has to be considered for the correct determination of the RI. In some of the sediments the released radioactive CO₂ more than doubled upon acidification. Actually, we also calculated RI values using the final radioactivity in CO₂ before acidification. These RI values were (of course) generally smaller than those after acidification, ranging between RI = 0.05-0.30 compared to RI = 0.06-0.48. The use of the non-acidified values of RI for calculation of acetate dependent CH₄ production would result in even higher rates than those given in our paper. Hence, the rates of acetate-dependent CH₄ production given in our paper were conservative. A brief statement has been added to the Discussion.

L.124: Averages were calculated since the values of f_{H_2} were not constant (Fig. 2C) and we wanted to compare the values with those determined earlier using ^{13}C (comparison in Fig. 1B). No changes were made.

Referee #2

1. The referee criticizes that the comparison between different methods is confusing. The comment is similar to one of referee #1. See there for action taken.

2. The referee questions whether sulfate, which was initially present in the sediments, could have contributed to acetate oxidation. To avoid sulfate reduction, we preincubated the sediments. This information is given in the Methods. We have now also briefly mentioned it in the Results and have emphasized that CH_4 production started without lag phase indicating that suppressive concentrations of sulfate (or Fe(III)) were not available.

3. Acetate oxidation and hydrogenotrophic methanogenesis are indeed two independent processes, that are coupled syntrophically via H_2 (or perhaps other electron carriers). The turnover rate of radioactive acetate comprises also such syntrophic acetate oxidation. However, syntrophic acetate oxidation results in stoichiometric amounts of hydrogenotrophically formed CH_4 . If turnover of radioactive acetate is larger than CH_4 production (as was the case for several sediments), the surplus cannot be due to syntrophically produced CH_4 . Hence this amount of acetate oxidation must be caused by other oxidants, i.e., organic compounds if inorganic ones are not available. The conclusion is summarized in Fig. 5, to which we will make better reference in the revision. The oxidation of acetate can either be directly coupled to reduction orgC (reaction 4 in Fig.5) or acetate is oxidized to CO_2 plus H_2 followed by the oxidation of H_2 with orgC (reactions 2 plus 5 in Fig. 5). However, acetate oxidation coupled to hydrogenotrophic methanogenesis (reactions 2 plus 3 in Fig. 5) cannot be larger than CH_4 production. We have added a few sentences to the Discussion for clarity. We have also amended the references to Fig. 5 with the respective reactions shown in the scheme of pathways.

Specific comments:

Line 4: we have used only the term acetoclastic.

Line 107-108: We have added the equations describing how the parameters were calculated.

Line 135: Thank you, we have corrected the figure numbers.

1 **Acetate turnover and methanogenic pathways in Amazonian lake sediments**

2

3 Ralf Conrad^a, Melanie Klose^a, Alex Enrich-Prast^{b,c}

4

5 ^aMax Planck Institute for Terrestrial Microbiology, Karl-von-Frisch-Str. 10, 35043 Marburg,
6 Germany

7 ^bDepartment of Thematic Studies - Environmental Change, Linköping University, Linköping,
8 Sweden

9 ^cDepartamento de Botânica, Instituto de Biologia, University Federal do Rio de Janeiro
10 (UFRJ), Rio de Janeiro, Brazil

11

12

13 Corresponding author:

14 Ralf Conrad, Max Planck Institute for Terrestrial Microbiology, Karl-von-Frisch-Str. 10,
15 35043 Marburg, Germany

16 Tel. +49-6421-178801; Fax: +49-6421-178809;

17 email: Conrad@mpi-marburg.mpg.de

18

19 **Abstract**

20 Lake sediments in Amazonia are a significant source of CH₄, a potential greenhouse gas.
21 Previous studies of sediments using ¹³C analysis found that the contribution of
22 hydrogenotrophic versus acetoclastic methanogenesis to CH₄ production was relatively high.
23 Here, we determined the methanogenic pathway in the same sediments (n = 6) by applying
24 [¹⁴C]bicarbonate or [2-¹⁴C]acetate, and confirmed the high relative contribution (50-80%) of
25 hydrogenotrophic methanogenesis. The respiratory index (RI) of [2-¹⁴C]acetate, which is
26 ¹⁴CO₂ relative to ¹⁴CH₄ + ¹⁴CO₂, divided the sediments into two categories, i.e., those with an
27 RI < 0.2 being consistent with the operation of acetoclastic methanogenesis, and those with an
28 RI > 0.4 showing that a large percentage of the acetate-methyl was oxidized to CO₂ rather
29 than reduced to CH₄. Hence, part of the acetate was probably converted to CO₂ plus H₂ via
30 syntrophic oxidation, thus enhancing hydrogenotrophic methanogenesis. This happened
31 despite the presence of potentially acetoclastic *Methanosaetaceae* in all the sediments.
32 Alternatively, acetate may have been oxidized with a constituent of the sediment organic
33 matter (humic acid) serving as oxidant. Indeed, apparent acetate turnover rates were larger
34 than CH₄ production rates except in those sediments with a R < 0.2. Our study demonstrates
35 that CH₄ production in Amazonian lake sediments was not simply caused by a combination of
36 hydrogenotrophic and acetoclastic methanogenesis, but probably involved additional acetate
37 turnover.
38

39 **1. Introduction**

40 Acetate is an important intermediate in the anoxic degradation of organic matter and is
41 produced by fermentation processes and chemolithotrophic homoacetogenesis. The
42 contribution of these two processes to acetate production is difficult to determine, but seems
43 to be quite different for different environments (Fu et al., 2018; Hädrich et al., 2012; Heuer et
44 al., 2010; Lokshina et al., 2019; Ye et al., 2014). The degradation of acetate requires a suitable
45 oxidant such as oxygen, nitrate, ferric iron or sulfate. If such oxidants are not or no longer
46 available, such as in many freshwater environments (e.g., paddy fields, lake sediments, peat),
47 acetate sometimes accumulates until suitable electron acceptors become again available.
48 Temporal accumulation and subsequent oxidative consumption has for example been
49 observed in peatlands during increase and decrease, respectively, of the water table
50 (Duddleston et al., 2002). ~~Normally, b~~However, it is generally assumed that acetate
51 degradation in the absence of inorganic electron acceptors is accomplished by aceticlastic
52 methanogenesis (Zinder, 1993). If aceticlastic methanogenesis is operative, the methyl group
53 of the acetate is converted to CH₄.

Formatted: Font color: Red

Formatted: Font color: Red

54 If methanogenesis is the exclusive final step in the anaerobic degradation of organic
55 matter, polysaccharides (one of the most important compounds from primary production) will
56 be dismutated to equal amounts of CH₄ and CO₂. Furthermore, acetate usually accounts for
57 more than two third of total methane production, especially if polysaccharides are the
58 predominant degradable organic matter (Conrad, 1999). However, CO₂ has often been found
59 to be the main product in many anoxic environments despite the absence of inorganic electron
60 acceptors (O₂, nitrate, ferric iron, sulfate) (Keller et al., 2009; Yavitt and Seidmann-Zager,
61 2006). Such results have been explained by the assumption that organic substances (e.g.
62 humic acids) may also serve as electron acceptors (Gao et al., 2019; Keller et al., 2009;
63 Klüpfel et al., 2014). Organic electron acceptors also allow the oxidation of acetate (Coates et
64 al., 1998; Lovley et al., 1996). The role of organic electron acceptors during anaerobic
65 degradation of organic matter is potentially important, but still not well known (Corbett et al.,
66 2013)

Formatted: Font color: Red

67 There are also many reports that methane production in lake sediments is dominated by
68 hydrogenotrophic rather than aceticlastic methanogenesis (Conrad, 1999; Conrad et al., 2011;
69 Ji et al., 2016). Such observations were explained (1) by incomplete degradation of organic
70 matter producing predominantly H₂ and CO₂ without concomitant acetate production (Conrad
71 et al., 2010; Hodgkins et al., 2014; Liu et al., 2017), (2) by acetate oxidation coupled to the
72 reduction of organic substances (see above), or (3) by syntrophic acetate oxidation coupled
73 with hydrogenotrophic methanogenesis (Lee and Zinder, 1988; Vavilin et al., 2017). If
74 ~~syntrophic~~ acetate oxidation is operative, the methyl group of the acetate is converted to CO₂;
75 ~~similarly as found during acetate oxidation with external inorganic or organic electron~~
76 ~~acceptors~~. However, if ~~syntrophic~~ acetate oxidation is syntrophic, it does not require a
77 chemical compound (other than H⁺) as electron acceptors, since it is the hydrogenotrophic
78 methanogenesis that eventually accepts the electrons released during acetate oxidation.

79 Syntrophic acetate oxidation can replace aceticlastic methanogenesis and thus, has been
80 found when aceticlastic methanogenic archaea were not present in the microbial community
81 of lake sediment (Nüsslein et al., 2001). This may also happen in other anoxic environments
82 when conditions are not suitable for aceticlastic methanogens, e.g., at elevated temperatures
83 (Conrad et al., 2009; Liu and Conrad, 2010; Liu et al., 2018), in the presence of high
84 concentrations of ammonium (Müller et al., 2016; Schnürer et al., 1999; Zhang et al., 2014),
85 or phosphate (Conrad et al., 2000). However, syntrophic acetate oxidation has also been found
86 in lake sediments that contained populations of putatively aceticlastic methanogens (Vavilin
87 et al., 2017). It is presently unknown under which conditions syntrophic acetate oxidizers can
88 successfully compete with aceticlastic methanogens and co-occur with acetate oxidation that
89 is coupled to the reduction of organic substances.

90 As a further step in understanding the ecology of ~~syntrophic~~ acetate oxidizers (syntrophic
91 or non-syntrophic ones) versus aceticlastic methanogens, we attempted to document their
92 coexistence by studying lake sediments, which had been reported containing 16S rRNA genes
93 of putatively aceticlastic *Methanosaetaceae* (*Methanotrichaceae* (Oren, 2014)) (Ji et al.,
94 2016). We used these sediments and measured the fractions of hydrogenotrophic

Formatted: Font color: Red

95 methanogenesis and of the methyl group of acetate being oxidized to CO₂ rather than reduced
96 to CH₄, and compared the turnover of acetate to the production rate of CH₄.

97

98 2. Materials and Methods

99 The sediment samples were obtained from floodplain lakes in the Amazon region and have
100 already been used for a study on structure and function of methanogenic microbial
101 communities (Ji et al., 2016). In particular, these sediment have been assayed for the
102 percentage of hydrogenotrophic methanogenesis ~~using values of δ¹³C of CH₄, CO₂ and~~
103 ~~acetate-methyl,~~ and for the percentage contribution of putatively aceticlastic methanogens to
104 the total archaeal community (Ji et al., 2016). Here, we used six of these sediments for
105 incubation experiments with radioactive tracers. These ~~are the same sediments samples as are~~
106 ~~identical to~~ those listed in our previous publication (Ji et al., 2016). The identity of the lake
107 sediments and the percentage content of putatively aceticlastic methanogens is summarized in
108 Table 1.

109 The experiments were carried out at the same time as those in our previous publication (Ji
110 et al., 2016) and were basically using the same incubation techniques. ~~However, the~~
111 ~~experimental approach to determine the fractions of hydrogenotrophic methanogenesis (f_{H2})~~
112 ~~were different. In our previous experiment, values of f_{H2} were determined from the δ¹³C of~~
113 ~~CH₄ in the presence (δ¹³C_{CH4-mc}) and absence (δ¹³C_{CH4}) of methyl fluoride, an inhibitor of~~
114 ~~aceticlastic methanogenesis, and from the δ¹³C of the methyl group of acetate (δ¹³C_{ac-methyl}):~~

$$115 f_{H2} = (\delta^{13}C_{CH4} - \delta^{13}C_{ac-methyl}) / (\delta^{13}C_{CH4-mc} - \delta^{13}C_{ac-methyl}) \quad (1)$$

116 The CH₄ production rates and f_{H2} values from this experiment are shown in Fig. 1 for
117 comparison.

118 In the present experiment, however, values of f_{H2} were determined by addition of
119 NaH¹⁴CO₃ and measurement of the specific radioactivities in CH₄ and CO₂. Briefly, ~~for~~
120 ~~determination of CH₄ production rates and the fractions of hydrogenotrophic methanogenesis~~
121 ~~(f_{H2})~~ about 10-15 ml of each replicate (n =3) were filled into 27-ml sterile tubes, flushed with
122 N₂, closed with butyl rubber stoppers, and incubated at 25°C. After preincubation for 12 days
123 (in order to deplete eventually present inorganic oxidants), 0.5 ml of a solution of carrier-free

Formatted: Subscript

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Subscript

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Superscript

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Subscript

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Subscript

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted

Formatted: Indent: First line: 0 cm

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

124 NaH¹⁴CO₃ (about 1 μCi; 50 Ci mol⁻¹) was added, the tubes flushed again with N₂, and
 125 incubation was continued at 25°C for about 100 days. Partial pressures of CH₄ and CO₂ as
 126 well as their contents of ¹⁴C were measured at different time points after mixing the slurries
 127 by heavy manual shaking. The gas partial pressures were measured by gas chromatography
 128 with a flame ionization detector (Ji et al., 2016), the radioactivities were analyzed with a
 129 radiodetector (RAGA) (Conrad et al., 1989). The data were used to calculate the fractions of
 130 hydrogenotrophic methanogenesis (f_{H2}) from the specific radioactivities of gaseous CH₄
 131 (SR_{CH4}) and CO₂ (SR_{CO2}):

$$132 \quad f_{H2} = SR_{CH4} / SR_{CO2} \quad (2).$$

133 For determination of acetate turnover, the same conditions were used, except that
 134 preincubation was for 25 days, 0.5 ml of a solution of carrier-free Na[2-¹⁴C]acetate (about 2
 135 μCi; 50 Ci mol⁻¹), equivalent to about 20 nmol acetate, was added, and incubation was
 136 continued for about 8 h. During this time gas samples were repeatedly taken and the
 137 radioactivities in CH₄ and CO₂ were analyzed in a gas chromatograph with a radiodetector
 138 (RAGA) (Conrad et al., 1989). In the end, the sediment samples were acidified with 1 ml of
 139 1M H₂SO₄ to liberate CO₂ from carbonates, and the radioactivities in CH₄ and CO₂ were
 140 analyzed again. The data were used to calculate the fractions of hydrogenotrophic
 141 methanogenesis (f_{H2}), the acetate turnover rate constants (k_{ac}) and the respiratory index (RI)
 142 values from the radioactivities of gaseous CH₄ and CO₂ as described by Schütz et al. (1989).

143 The RI is defined as:

$$144 \quad RI = {}^{14}CO_2 / ({}^{14}CO_2 + {}^{14}CH_4) \quad (3).$$

145 Both ¹⁴CH₄ and ¹⁴CO₂ were measured at the end of the incubation after acidification. These
 146 data were asleThe acetate turnover rate constants was were determined from the change of
 147 ¹⁴CH₄ and ¹⁴CO₂ with incubation time (t) and the maximal values of ¹⁴CH₄ and ¹⁴CO₂ at the
 148 end of the incubation before acidification:

$$149 \quad k_{ac} = [\ln(1 - ({}^{14}CH_4 + {}^{14}CO_2) / ({}^{14}CH_{4-max} + {}^{14}CO_{2-max}))] / t \quad (4).$$

150 The acetate turnover rates (v_{ac}) were calculated by as the product of k_{ac} time the acetate
 151 concentration (ac), which was analyzed in the sediments at the end of the incubation using
 152 high pressure liquid chromatography:

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Indent: First line: 0,5 cm

Formatted: Subscript

Formatted: Indent: First line: 0 cm

Formatted: Subscript

153 $v_{ac} = k_{ac} \cdot ac$ (5).

154 The acetate concentration (ac) was analyzed in the sediments at the end of the incubation
155 using high pressure liquid chromatography. The acetate concentrations are summarized in
156 Table 1. The rates of acetate-dependent CH₄ production (P_{ac}) were calculated from the acetate
157 turnover rates and the RI:

158 $P_{ac} = v_{ac} \cdot (1 - RI)$ (6).

160 3. Results

161 Six different lake sediments from Amazonia were incubated in the ~~absence and the~~
162 presence of H¹⁴CO₃. Methane production started without lag phase indicating that the
163 inorganic electron acceptors, which were present in the original sediment (Ji et al., 2016) had
164 been depleted during the anaerobic preincubation and did not suppress methanogenesis. The
165 CH₄ production rates were compared to those obtained in our previous experiments without
166 addition of H¹⁴CO₃ (Ji et al., 2016). Although the rates of CH₄ production were different in
167 the two different incubations, the orders of magnitude were similar for the different lake
168 sediments (Fig. 1A). The incubations in the presence of H¹⁴CO₃ were used to follow the
169 specific radioactivities of CH₄ (Fig. 2A) and CO₂ (Fig. 2B) over the incubation time. The
170 specific radioactivities of CH₄ changed only little but were slightly different for the different
171 lake sediments. The specific radioactivities of CO₂ decreased with time as expected due to the
172 production of non-radioactive CO₂. Both specific radioactivities were used to calculate the
173 fractions of hydrogenotrophic methanogenesis (f_{H2}), which increased with incubation time
174 and eventually reached a plateau. The values of f_{H2} averaged between 30 and 60 d of
175 incubation are summarized in Fig. 1B. Only the incubations of sediment “Grande” did not
176 reach a plateau but still increased after 260 d of incubation due to the continuously decreasing
177 specific radioactivities of CO₂ (data not shown). Averaging these values over the 4 data points
178 between 160 and 260 d resulted in f_{H2} of about 60% (Fig. 1B). The thus determined values of
179 f_{H2} were comparable to those determined in the absence of H¹⁴CO₃ using values of δ¹³C,
180 which have already been published (Ji et al. 2016) (Fig.1B).

Formatted: Not Superscript/ Subscript

Formatted: Indent: First line: 0 cm

Formatted: Subscript

Formatted: Subscript

Formatted: Not Superscript/ Subscript

Formatted: Subscript

181 The same sediments were used to determine the turnover of [2-¹⁴C]acetate by measuring
182 the increase of radioactive CH₄ (Fig. 3A) and CO₂ (Fig. 3B). These data were used to
183 determine the rate constants of acetate turnover (Fig. 3C), which ranged between 0.02 and 1.7
184 h⁻¹. The respiratory indices (RI) were generally larger than 0.2 except those of the sediments
185 Tapari and Verde, which were smaller than 0.2 (Fig. 4B). The RI values and the acetate
186 turnover rate constants were used to calculate the rates of CH₄ production from acetate in
187 comparison to the rates of total CH₄ production (Fig. 4A). Interestingly, acetate-dependent
188 CH₄ production was always larger than total CH₄ production, except in those sediments
189 exhibiting a RI <0.2.

190

191 4. Discussion

192 The RI value quantifies the fraction of the methyl group of acetate that is oxidized to CO₂
193 rather than reduced to CH₄. Since some oxidation of acetate methyl is also happening in pure
194 cultures of aceticlastic methanogens (Weimer and Zeikus, 1978), and since a RI of around 0.2
195 has often been found in environments where acetate turnover was dominated by aceticlastic
196 methanogenesis (Phelps and Zeikus, 1984; Rothfuss and Conrad, 1993; Winfrey and Zeikus,
197 1979), an RI value of 0.2 may in practice be used as the threshold for the change of
198 methanogenic to oxidative acetate turnover. Based on this criterion, i.e. RI < 0.2, the lake
199 sediments of Tapari and Verde behaved as when acetate turnover was exclusively caused by
200 aceticlastic methanogenesis. The percentage of acetate-dependent CH₄ production was fairly
201 consistent with the fraction of hydrogenotrophic methanogenesis, which made up the
202 remainder of total CH₄ production. In conclusion, the acetate turnover and CH₄ production in
203 these lake sediments behaved as expected as when aceticlastic methanogenesis was the sole
204 process of acetate consumption (reaction 1 in Fig. 5).

205 However, the sediments of Jua and in particular those of Jupinda, Cataldo, and Grande
206 exhibited RI values >0.2, showing that a substantial fraction of the acetate-methyl was
207 oxidized to CO₂. Hence, acetate was not exclusively consumed by aceticlastic
208 methanogenesis, but it was oxidized, for example by syntrophic acetate oxidation producing
209 H₂ and CO₂. The H₂ and CO₂ may subsequently have been used as methanogenic substrates,

210 thus supporting CH₄ production (reactions 2 and 3 in Fig. 5). Such support would be
211 consistent with the relatively high fractions (f_{H_2}) of hydrogenotrophic methanogenesis
212 observed in these sediments. However, it would not explain why acetate turnover rates were
213 higher than necessary for supporting the observed rates of total CH₄ production. A possible
214 conclusion is that acetate was converted to CO₂ without concomitant production of H₂.
215 Possibly, electrons from acetate were transferred to organic electron acceptors (reaction 4 in
216 Fig. 5), such as suggested in the literature (Coates et al., 1998; Lovley et al., 1996).

217 Alternatively, acetate may have first been converted to H₂ plus CO₂ followed by the oxidation
218 of H₂ with organic electron acceptors (reactions 2 and 5 in Fig. 5) rather than syntrophic
219 formation of CH₄ from H₂ plus CO₂ (reactions 2 and 3 in Fig. 5). In conclusion, these lake
220 sediments behaved as when acetate consumption was accomplished not only by ~~aceticlastic~~
221 acetate-dependent methanogenesis, but also by oxidative consumption.

222 Our conclusions are mainly based on radiotracer measurements, which may be biased. For
223 example, acetate turnover rate constants are calculated from acetate concentrations and
224 turnover rate constants. Acetate concentrations were only measured at the end of incubation
225 and thus, may not have been representative for ~~much of the~~ entire incubation time.

226 Furthermore, acetate in the sediment may occur in several pools with different turnover
227 (Christensen and Blackburn, 1982). Therefore, acetate turnover rates and acetate-dependent
228 CH₄ production rates may be overestimated, if the actual acetate turnover depends on a pool
229 size that is smaller than that analyzed. Overestimation may also result from too low RI values,
230 such as when carbonate-bound radioactivity is neglected. However, such bias was avoided by
231 acidification prior to determination of the RI. ~~Such overestimation in sediments of Jua,~~
232 ~~Jupinda, Cataldo and Grande cannot be completely excluded, although rates in sediments of~~
233 ~~Tapari and Verde were in a realistic range.~~ Finally, a potential bias may arise from the fact
234 that the rates of CH₄ production and the acetate turnover rates were measured in two different
235 sets of incubation, with different incubation times. While CH₄ production (and f_{H_2}) was
236 measured over tens of days (Fig. 2), acetate turnover was determined within 8 h (Fig. 3).
237 Nevertheless, the data in the lake sediments of Tapari and Verde resulted in CH₄ production
238 and acetate turnover consistent with the operation of aceticlastic methanogenesis, which is the

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

239 canonical acetate consumption pathway for methanogenic sediments. Therefore, we are
240 confident that our results obtained from ~~Such overestimation in the~~ sediments of Jua, Jupinda,
241 Cataldo and Grande cannot be completely excluded, although rates in sediments of Tapari and
242 Verde were also in a realistic range.

243 The determination of fractions of hydrogenotrophic methanogenesis (f_{H_2}) depends on the
244 specific radioactivity of the dissolved CO_2 pool that is involved in CH_4 production. However,
245 it is the pool of gaseous CO_2 that is analyzed in the assay, assuming that its specific
246 radioactivity is identical to that of the active dissolved pool. Since non-radioactive CO_2 is
247 permanently produced from oxidation of organic matter, there may be disequilibrium.
248 Nevertheless, determinations of f_{H_2} using radioactive bicarbonate exhibited the same
249 tendencies as those based on $\delta^{13}C$ values, and thus are probably quite reliable. Furthermore,
250 the f_{H_2} values were fairly consistent with the fractions of acetate-dependent methanogenesis
251 determined from the turnover of radioactive acetate.

252 Despite these reservations, our results collectively demonstrated that acetate turnover in
253 tropical lake sediments did not necessarily follow a canonical pattern with aceticlastic
254 methanogenesis as sole or predominant process of acetate turnover, despite the fact that all
255 these sediments contained populations of putative aceticlastic methanogenic archaea. Acetate
256 consumption in *Methanosaeta* species is known to have a relatively high affinity and a low
257 threshold for acetate (Jetten et al., 1992). Therefore, the question arises why oxidative
258 processes, including syntrophic acetate oxidation, could successfully compete with
259 aceticlastic methanogenesis.

261 5. Author contribution

262 RC designed the experiments, evaluated the data and wrote the manuscript; MK did the
263 experiments; AEP provided the samples and contributed to the discussion of the data.

265 6. Competing interests

266 The authors declare that they have no conflict of interest.

Formatted: Subscript

268 **7. Acknowledgements**
269 AEP acknowledges funding from the Swedish Research Council Vinnova and Linköping
270 University and for funding from the Brazilian Research Council FAPERJ.
271

272 **8. References**

273 Christensen, D. and Blackburn, T. H.: Turnover of ¹⁴C-labelled acetate in marine
274 sediments, *Mar. Biol.*, 71, 113-119, 1982.

275 Coates, J. D., Ellis, D. J., Blunt-Harris, E. L., Gaw, C. V., Roden, E. E., Lovley,
276 D. R.: Recovery of humic-reducing bacteria from a diversity of environments,
277 *Appl. Environ. Microbiol.*, 64, 1504-1509, 1998.

278 Conrad, R.: Contribution of hydrogen to methane production and control of
279 hydrogen concentrations in methanogenic soils and sediments [review], *FEMS*
280 *Microbiol. Ecol.*, 28, 193-202, 1999.

281 Conrad, R., Claus, P., Casper, P.: Stable isotope fractionation during the
282 methanogenic degradation of organic matter in the sediment of an acidic bog
283 lake, Lake Grosse Fuchskuhle, *Limnol. Oceanogr.*, 55, 1932-1942, 2010.

284 Conrad, R., Klose, M., Claus, P.: Phosphate inhibits acetotrophic methanogenesis
285 on rice roots, *Appl. Environ. Microbiol.*, 66, 828-831, 2000.

286 Conrad, R., Klose, M., Noll, M.: Functional and structural response of the
287 methanogenic microbial community in rice field soil to temperature change,
288 *Environ. Microbiol.*, 11, 1844-1853, 2009.

289 Conrad, R., Mayer, H. P., Wüst, M.: Temporal change of gas metabolism by
290 hydrogen-syntrophic methanogenic bacterial associations in anoxic paddy soil,
291 *FEMS Microbiol. Ecol.*, 62, 265-274, 1989.

292 Conrad, R., Noll, M., Claus, P., Klose, M., Bastos, W. R., Enrich-Prast, A.: Stable
293 carbon isotope discrimination and microbiology of methane formation in
294 tropical anoxic lake sediments, *Biogeosciences*, 8, 795-814, 2011.

295 Corbett, J., Tfaily, M. M., Burdige, D. J., Cooper, W. T., Glaser, P. H., Chanton,
296 J. P.: Partitioning pathways of CO₂ production in peatlands with stable carbon
297 isotopes, *Biogeochem.*, 114, 327-340, 2013.

298 Duddleston, K. N., Kinney, M. A., Kiene, R. P., Hines, M. E.: Anaerobic
299 microbial biogeochemistry in a northern bog: Acetate as a dominant metabolic
300 end product, *Global Biogeochem. Cycles*, 16, 1063-
301 doi:10.1029/2001GB001402, 2002.

302 Fu, B., Conrad, R., Blaser, M.: Potential contribution of acetogenesis to anaerobic
303 degradation in methanogenic rice field soils, *Soil Biol. Biochem.*, 119, 1-10,
304 2018.

Formatted: Indent: First line: 0 cm, Keep with next

305 Gao, C., Sander, M., Agethen, S., Knorr, K. H.: Electron accepting capacity of
 306 dissolved and particulate organic matter control CO₂ and CH₄ formation in
 307 peat soils, *Geochim. Cosmochim. Acta*, 245, 266-277, 2019.

308 Hädrich, A., Heuer, V. B., Herrmann, M., Hinrichs, K. U., Küsel, K.: Origin and
 309 fate of acetate in an acidic fen, *FEMS Microbiol. Ecol.*, 81, 339-354, 2012.

310 Heuer, V. B., Krüger, M., Elvert, M., Hinrichs, K. U.: Experimental studies on the
 311 stable carbon isotope biogeochemistry of acetate in lake sediments, *Org.*
 312 *Geochem.*, 41, 22-30, 2010.

313 Hodgkins, S. B., Tfaily, M. M., McCalley, C. K., Logan, T. A., Crill, P. M.,
 314 Saleska, S. R., Rich, V. I., Chanton, J. P.: Changes in peat chemistry associated
 315 with permafrost thaw increase greenhouse gas production, *Proc. Natl. Acad.*
 316 *Sci. USA*, 111, 5819-5824, 2014.

317 Jetten, M. S. M., Stams, A. J. M., Zehnder, A. J. B.: Methanogenesis from acetate
 318 - A comparison of the acetate metabolism in *Methanotherix soehngenii* and
 319 *Methanosarcina* spp., *FEMS Microbiol. Rev.*, 88, 181-197, 1992.

320 Ji, Y., Angel, R., Klose, M., Claus, P., Marotta, H., Pinho, L., Enrich-Prast, A.,
 321 Conrad, R.: Structure and function of methanogenic microbial communities in
 322 sediments of Amazonian lakes with different water types, *Environ. Microbiol.*,
 323 18, 5082-5100, 2016.

324 Keller, J. K., Weisenhorn, P. B., Megonigal, J. P.: Humic acids as electron
 325 acceptors in wetland decomposition, *Soil Biol. Biochem.*, 41, 1518-1522, 2009.

326 Klüpfel, L., Piepenbrock, A., Kappler, A., Sander, M.: Humic substances as fully
 327 regenerable electron acceptors in recurrently anoxic environments, *Nature*
 328 *Geoscience*, 7, 195-200, 2014.

329 Lee, M. J. and Zinder, S. H.: Isolation and characterization of a thermophilic
 330 bacterium which oxidizes acetate in syntrophic association with a methanogen
 331 and which grows acetogenically on H₂-CO₂, *Appl. Environ. Microbiol.*, 54,
 332 124-129, 1988.

333 Liu, F. H. and Conrad, R.: *Thermoanaerobacteriaceae* oxidize acetate in
 334 methanogenic rice field soil at 50°C, *Environ. Microbiol.*, 12, 2341-2354,
 335 2010.

336 Liu, P. F., Klose, M., Conrad, R.: Temperature effects on structure and function of
 337 the methanogenic microbial communities in two paddy soils and one desert
 338 soil, *Soil Biol. Biochem.*, 124, 236-244, 2018.

339 Liu, Y., Conrad, R., Yao, T., Gleixner, G., Claus, P.: Change of methane
 340 production pathway with sediment depth in a lake on the Tibetan plateau,
 341 *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 474, 279-286, 2017.

342 Lokshina, L., Vavilin, V., Litt, Y., Glagolev, M., Sabrekov, A., Kotsyurbenko,
 343 O., Kozlova, M.: Methane production in a West Siberian eutrophic fen is much
 344 higher than carbon dioxide production: incubation of peat samples,

345 stoichiometry, stable isotope dynamics, modeling, *Water Resources*, 46, S110-
346 S125, 2019.

347 Lovley, D. R., Coates, J. D., Blunt-Harris, E. L., Phillips, E. J. P., Woodward, J.
348 C.: Humic substances as electron acceptors for microbial respiration, *Nature*,
349 382, 445-448, 1996.

350 Müller, B., Sun, L., Westerholm, M., Schnürer, A.: Bacterial community
351 composition and fhs profiles of low- and high-ammonia biogas digesters reveal
352 novel syntrophic acetate-oxidising bacteria, *Biotechnol. Biofuels*, 9, 48-
353 doi:10.1186/s13068-016-0454-9, 2016.

354 Nüsslein, B., Chin, K. J., Eckert, W., Conrad, R.: Evidence for anaerobic
355 syntrophic acetate oxidation during methane production in the profundal
356 sediment of subtropical Lake Kinneret (Israel), *Environ. Microbiol.*, 3, 460-
357 470, 2001.

358 Oren, A.: The family *Methanotrichaceae*, in: *The Prokaryotes*, edited by:
359 Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E., Thompson, F.,
360 Springer, Berlin, 298-306, 2014.

361 Phelps, T. J. and Zeikus, J. G.: Influence of pH on terminal carbon metabolism in
362 anoxic sediments from a mildly acidic lake, *Appl. Environ. Microbiol.*, 48,
363 1088-1095, 1984.

364 Rothfuss, F. and Conrad, R.: Vertical profiles of CH₄ concentrations, dissolved
365 substrates and processes involved in CH₄ production in a flooded Italian rice
366 field, *Biogeochem.*, 18, 137-152, 1993.

367 Schnürer, A., Zellner, G., Svensson, B. H.: Mesophilic syntrophic acetate
368 oxidation during methane formation in biogas reactors, *FEMS Microbiol.*
369 *Ecol.*, 29, 249-261, 1999.

370 Schütz, H., Seiler, W., Conrad, R.: Processes involved in formation and emission
371 of methane in rice paddies, *Biogeochem.*, 7, 33-53, 1989.

372 Vavilin, V., Rytov, S., Conrad, R.: Modeling methane formation in sediments of
373 tropical lakes, focusing on syntrophic acetate oxidation: dynamics and static
374 isotope equations, *Ecol. Modeling*, 363, 81-95, 2017.

375 Weimer, P. J. and Zeikus, J. G.: Acetate metabolism in *Methanosarcina barkeri*,
376 *Arch. Microbiol.*, 119, 175-182, 1978.

377 Winfrey, M. R. and Zeikus, J. G.: Anaerobic metabolism of immediate methane
378 precursors in Lake Mendota, *Appl. Environ. Microbiol.*, 37, 244-253, 1979.

379 Yavitt, J. B. and Seidmann-Zager, M.: Methanogenic conditions in northern peat
380 soils, *Geomicrobiol. J.*, 23, 119-127, 2006.

381 Ye, R., Jin, Q., Bohannon, B., Keller, J. K., Bridgham, S. D.: Homoacetogenesis:
382 A potentially underappreciated carbon pathway in peatlands, *Soil Biol.*
383 *Biochem.*, 68, 385-391, 2014.

- 384 Zhang, C., Yuan, Q., Lu, Y.: Inhibitory effects of ammonia on methanogen *mcrA*
385 transcripts in anaerobic digester sludge, *FEMS Microbiol. Ecol.*, 87, 368-377,
386 2014.
- 387 Zinder, S. H.: Physiological ecology of methanogens, in: *Methanogenesis.*
388 *Ecology, Physiology, Biochemistry and Genetics*, edited by: Ferry, J. G.,
389 Chapman & Hall, New York, 128-206, 1993.
390
391

392 Table 1: Identity of sediment samples (same as those in Ji et al. (2016)),
393 percentage content of putatively acetoclastic methanogens (*Methanosaetaceae*)
394 relative to total archaea, and concentrations of acetate; mean \pm SE.

395

Lake #	Name	Type	<i>Methanosaetaceae</i> (%)	Acetate (nmol g ⁻¹ dry weight)
P1	Jua	clear water	21 \pm 1	93 \pm 5
P8	Tapari	clear water	19 \pm 3	261 \pm 39
P9	Verde	clear water	19 \pm 11	126 \pm 12
P10	Jupinda	clear water	27 \pm 4	110 \pm 6
A1	Cataldo	white water	42 \pm 1	50 \pm 3
A2	Grande	white water	36 \pm 3	35 \pm 1

396

397

398 **Figure captions**

399 Fig. 1: Methane production in sediments of different Amazonian lakes: (A) rates
400 of CH₄ production, and (B) fractions of hydrogenotrophic methanogenesis,
401 both determined in the absence and the presence of radioactive bicarbonate.
402 The data in the absence of radioactive bicarbonate are the same as published in
403 Ji et al. (2016), when f_{H₂} was determined from values of δ¹³C; mean ±SE.

404 Fig. 2: Conversion of radioactive bicarbonate in sediments of different Amazonian
405 lakes: (A) specific radioactivities in CH₄; (B) specific radioactivities in gaseous
406 CO₂; and (C) fractions (f_{H₂}) of hydrogenotrophic methanogenesis; mean ±SE.

407 Fig. 3: Conversion of [2-¹⁴C]acetate in sediments of different Amazonian lakes:
408 (A) accumulation of radioactive CH₄; (B) accumulation of radioactive gaseous
409 CO₂; and (C) acetate turnover rate constants; mean ±SE.

410 Fig. 4: (A) Rates of total and acetate-derived CH₄ production in sediments of
411 different Amazonian lakes and (B) respiratory indices (RI) of the turned over
412 [2-¹⁴C]acetate; mean ±SE.

413 Fig. 5: Scheme of the pathways involved in acetate turnover in sediments of
414 Amazonian lakes; (1) aceticlastic methanogenesis; (2) syntrophic acetate
415 oxidation; (3) hydrogenotrophic methanogenesis; (4) acetate oxidation with
416 organic electron acceptors.