

# 5 **Wind-driven stratification patterns and dissolved oxygen depletion off the Changjiang (Yangtze) Estuary**

Taavi Liblik<sup>1,2</sup>, Yijing Wu<sup>1</sup>, Daidu Fan<sup>1,3,\*</sup>, Dinghui Shang<sup>1</sup>

10 <sup>1</sup> State Key Laboratory of Marine Geology, Tongji University, Shanghai, 200092, China

<sup>2</sup> Department of Marine Systems, Tallinn University of Technology, Tallinn, 12618, Estonia

<sup>3</sup> Laboratory of Marine Geology, Qingdao National Laboratory for Marine Science and Technology, 387P+WW Shinan, Qingdao, Shandong, China

*Correspondence to:* Daidu Fan (ddf@tongji.edu.cn)

15

**Abstract.** The area off the Changjiang Estuary is under strong impact of fresh water and anthropogenic nutrient load from the Changjiang River. The seasonal hypoxia in the area has variable location and range, but the decadal trend reveals expansion and intensification of the dissolved oxygen (DO) depletion.

Two oceanographic cruises, conducted in summers 2015 and 2017, were complemented by river discharge, circulation simulation, and remotely sensed wind, salinity and sea level anomaly data. Two distinct situations of stratification and DO distributions in summers 2015 and 2017 can be explained by wind forcing and concurrent features in surface and deep layer circulation, upwelling and downwelling events.

Enhanced primary production in the upper layer in the Changjiang Diluted Water (CDW) or the upwelled water determined the location and extent of oxygen depletion. However, the pycnocline created by Kuroshio subsurface water intrusion was an essential precondition for hypoxia formation. Spreading of the CDW, the occurrence of upwelling and intrusion of deep layer water are strongly influenced by wind forcing.

Intensification of Chinese Coastal Current and CDW spreading to the south together with coastal downwelling caused by northerly wind were observed in 2015. This physical forcing leads to a well-ventilated area in the north and a hypoxic area of  $1.3 \times 10^4$  km<sup>2</sup> in the southern part. Alteration of the Chinese Coastal Current due to Ekman surface flow and reversal of the geostrophic current related to upwelling induced by summer monsoon (southerly winds) were observed in 2017. Wind-driven offshore transport in the surface layer caused a subsurface intrusion of Kuroshio derived water to the shallow coastal areas (<10 m depth). Intense hypoxia (DO down to 0.6 mg l<sup>-1</sup>) starting from 4-8 m depth connected to CDW and deep water intrusion in the north, and coastal hypoxia linked to upwelling were observed in 2017.

Variability in the wind forcing and river run-off and related changes in a physical environment considerably shape the extent and location of hypoxia in synoptic, seasonal and inter-annual time-scale off the Changjiang Estuary.

40

Dead zones in the coastal ocean have spread since the 1960s (Diaz and Rosenberg, 2008). Besides eutrophication (Diaz and Rosenberg, 2008), climate change (Altieri and Gedan, 2015) intensifies dissolved oxygen (DO) depletion. Estuaries and other regions of fresh water influence are typically very productive areas and often experience hypoxia (Conley et al., 2009; Cui et al., 2019; Obenour et al., 2013; Testa and Kemp, 2014). Natural and anthropogenic nutrient load in these areas leads to intensive sedimentation of organic matter. DO is used by concurrent decomposition of detritus by bacteria in the sub-surface layer. Deep layers below the euphotic zone can receive DO by physical processes only – lateral advection and vertical mixing. If the decline of DO exceeds DO import or production, the DO concentrations decrease.

Hypoxia is a natural phenomenon off the Changjiang Estuary, as evidence of its existence extends back to 2600 years at least (Ren et al., 2019). However, the hypoxic area has been expanded in recent decades (Chen et al., 2017; Ning et al., 2011; Zhu et al., 2011), which can be mostly related to eutrophication (Wang et al., 2016). A low DO zone off the estuary can be found from late spring (Zuo et al., 2019) until autumn as the stratification decays (Wang et al., 2012). The formation and maintenance of seasonal hypoxia have been related to various physical, biogeochemical and biological processes (Ning et al., 2011). Hypoxia off the Changjiang Estuary is in the literature often divided into the southern and northern areas around 30° N (Wei et al., 2017a; Zhu et al., 2011). The northern part features a shallow (20–40 m) and flat sea bottom, while the southern part is characterized by a deep trough with water depth > 60 m and a steep slope. Numerous small peninsulas and islands occupy the southern part with irregular coastline.

The region is strongly impacted by huge discharge from the Changjiang River (Beardsley et al., 1985; Xu et al., 2018). Correlation between the CDW plume area and river discharge has been suggested by Kang et al. (2013). Nutrient-rich freshwater mixes with ocean water to form the Changjiang Diluted Water (CDW). Enhanced primary production in the CDW causes intense detritus accumulation and DO consumption in the near-bottom layers (Chen et al., 2017; Li et al., 2018b; Wang et al., 2017, 2016; Zhou et al., 2019). The CDW is separated from the deeper water by a shallow halocline, which often coincides with the seasonal thermocline (Zhu et al., 2016). The shallow pycnocline impedes vertical mixing and consequently DO transport downwards. Thus, the presence of the CDW provides favorable conditions for hypoxia formation in the layer below the pycnocline. Moreover, the shallow pycnocline supports accelerated warming in summer (Moon et al., 2019), which in turn strengthens the pycnocline even more. Therefore, the spreading of the CDW strongly determines oxygen depletion in the near-bottom layer.

Shallower areas near the river mouth are strongly impacted by tidal forcing (Li et al., 2018). However, in the open sea, the contribution of tides in the vertical mixing budget is low comparing to that of wind stirring (Ni et al., 2016). Abrupt wind mixing events can weaken stratification and considerably increase DO concentrations in the deep layer (Ni et al., 2016). However, such vertical mixing also causes the nutrient flux to the upper layer (Hung et al., 2013), consequently enhancing primary production in the upper layer and then DO consumption in the deep layer (Ni et al., 2016).

Besides the CDW and wind mixing, circulation and hydrography near the Changjiang Estuary are further influenced by the Chinese Coastal Current and the Taiwan Warm Current. The latter originates from the Taiwan Strait warm water (>25° C, e.g. Zhang et al. (2007a) at the surface and the shelf-intrusion water of Kuroshio subsurface water (KSSW, <25° C, e.g. Zhang et al. (2007a) at the bottom (Lie and Cho, 2002, 2016; Liu et al.,

85 2017; Zhang et al., 2007b; Zuo et al., 2019). Geostrophic currents in the area are modified by wind forcing, which  
exhibits seasonal alternation of winter and summer monsoons (Lie and Cho, 2002). Winds from north and northeast  
prevail during the winter monsoon, and southerly winds dominate during the summer monsoon (Chu et al., 2005).  
Summer monsoon favors upwelling in the East China Sea (ECS) (Hu and Wang, 2016), including the offshore  
area near the Changjiang Estuary (Xu et al., 2017; Yang et al., 2019). Upwelling events bring large amounts of  
90 nutrients to the surface layer (Wang and Wang, 2007), which cause increase of primary production and subsequent  
decrease of DO concentrations in the near-bottom layer (Chen et al., 2004; Zhou et al., 2019). Moreover, coupling  
of the CDW plume and upwelled KSSW has a cumulative effect on primary production and hypoxia formation in  
the near bottom layer (Wei et al., 2017a). CDW spreading and upwelling of subsurface water are both sensitive to  
wind forcing, which gives rise to the hypothesis that hypoxic conditions are strongly dependent on wind conditions.  
95 Zhang et al. (2018) demonstrated that high mobility of the CDW strongly controls the variation of location and  
areal extent of bottom hypoxia. Thus, distributions of oceanographic fields mapped during an occasional research  
cruise could depend on simultaneous synoptic-scale forcing conditions and do not necessarily represent the  
hydrographical and chemical situation in a particular summer. In order to put oceanographic cruise results in a  
more general context, observed ocean variables must be linked to the forcing. The role of nutrient consumption  
100 and primary production in the CDW and the upwelled water, and their interaction have been previously  
investigated. Enhanced primary production due to upwelling and the CDW spreading produces higher oxygen  
consumption rates in the near-bottom layers (Chen et al., 2017, 2004; Große et al., 2019; Li et al., 2018b; Wang et  
al., 2017, 2016; Wei et al., 2017a; Zhou et al., 2019). The focus of this work will be on the physical forcing  
mechanisms that create favorable conditions for those mentioned above biogeochemical and biological processes.  
105 We hypothesize that prevalent physical factors, wind forcing and river run-off mainly determines the stratification  
patterns and consequently, the location and areal extent of the DO depletion off the Changjiang Estuary. That is,  
changes in forcing over various time scales (synoptic, inter-annual, and decadal) cause changes in the patterns of  
stratification and bottom hypoxia. To testify this hypothesis, we (1) describe spatial patterns of temperature,  
salinity, chlorophyll *a* (Chl *a*) and DO off the Changjiang Estuary by two cruises in the summers of 2015 and  
110 2017, (2) explore the potential link between physical variables and DO distributions, and (3) investigate the forcing  
mechanism behind the observed patterns.

115

## 2 Data and Methods

We performed two surveys in the Changjiang Estuary and the adjacent sea onboard the RV Zhehaike-1 from 27  
August to 5 September in 2015 and from 24 to 29 July in 2017, respectively. CTD (Conductivity, Temperature,  
120 Depth) and DO profiles were obtained from 110 stations in 2015 and 83 stations in 2017. However, we have  
excluded stations inside the estuary; therefore, 65 and 49 stations were included in the present study, respectively,  
in 2015 and 2017 (Fig. 1). At all stations, vertical profiles of temperature, salinity and DO were recorded with the

SBE 25plus Sealogger CTD with DO sensor SBE43. The DO sensor was calibrated against water sample analyses conducted by the Winkler titration method. Linear regression between sensor and sample DO concentration data for 2015 and 2017 were analyzed:  $DO = DO_{SBE} \times 0.98 - 0.09$  ( $r^2 = 0.93$ ,  $n = 191$ ) and  $DO = DO_{SBE} \times 0.98 - 0.06$  ( $r^2 = 0.99$ ,  $n = 98$ ), where  $DO_{SBE}$  is the DO concentration recorded by SBE43 sensor (Wu et al., 2019).

For the 2015 cruise, Chl *a* concentrations were measured from water samples as follows. About 500 ml surface water was filtered through 0.7  $\mu\text{m}$  (pore size) Whatman glass fiber filters at every station with pressure lower than 15 Kpa to get the particles for further analysis. To avoid the influence of light during the storage, all filters after filtration were folded up and wrapped by the aluminium foil before preserved in the  $-20^\circ\text{C}$  freezer. The pigments in each filter were extracted by 90% acetone and measured by fluorometric analysis using the Hitachi Designs fluorescence spectrophotometer (F-2700).

Since in-situ Chl *a* was not sampled in 2017, the daily satellite (Aqua MODIS) derived Chl *a* data were downloaded from <https://coastwatch.pfeg.noaa.gov/>. According to data availability, mean Chl *a* field on 21–22 July, i.e. 3–4 days before our survey, was presented to illustrate primary production in 2017.

Salinity data in the present work was given in the Practical Salinity Scale (Fofonoff and Millard Jr, 1983) and density as potential density anomaly ( $\sigma_\theta$ ) to reference pressure of 0 dbar (Association for the Physical Sciences of the Sea, 2010).

8-day running average Sea Surface Salinity (SSS) data with a resolution of  $0.25^\circ$  calculated from Soil Moisture Active Passive (SMAP) mission (Meissner et al., 2018) were used to estimate the spatial extent of CDW from 2015 to 2018. Remotely sensed salinity agreed with the in-situ salinity measurements (Fig. 2) at the mooring station M1 (Fig. 1).

Reprocessed 6-hourly wind observations with a horizontal resolution of  $0.25^\circ$  from 1993 to 2018 downloaded from Copernicus Marine Service (product ID WIND\_GLO\_WIND\_L4\_REP\_OBSERVATIONS\_012\_006, <http://marine.copernicus.eu/>) were used to describe wind conditions. Wind stress was calculated following Large and Pond (1981):

$$\tau = \rho_{air} C_D |U_{10}|U_{10}, \quad (1)$$

where  $\rho_{air}$  is the air density,  $C_D$  is the drag coefficient, and  $U_{10}$  is the wind speed at the reference height of 10 m above sea level.

Current speed and direction from 1993 to 2018 were downloaded from the Copernicus Marine Service reanalysis product GLORYS12V1. The global eddy-resolving model has a regular horizontal grid of approximately 8 km ( $1/12^\circ$ ) and standard vertical levels with resolutions increasing from 1 m near the surface to 8 m around 50 m depth, and then to 17 m around 100 m depth. Remotely sensed and observed in-situ temperature, salinity, and sea level were assimilated to the model. The model has a too coarse resolution to estimate details of meso- and finer-scaled features in the area. However, it can be used to estimate the general current patterns.

Daily mean sea level anomaly and meridional and zonal components of absolute geostrophic velocity based on satellite altimetry were downloaded from the Copernicus Marine Service (product DATASET-DUACS-REP-GLOBAL-MERGED-ALLSAT-PHY-L4). The sea level anomaly is defined as the sea surface height above the mean sea surface during the reference period 1993–2012. The horizontal and temporal resolution of the dataset is  $0.25^\circ$  and one day, respectively.

Depending on marine organisms, hypoxia can have various definitions (Vaquer-Sunyer and Duarte, 2008). In the present study, hypoxia is defined as DO concentration of  $<3.0 \text{ mg l}^{-1}$ . In order to estimate oxygen consumption, the apparent oxygen utilization (AOU) between DO concentration at the saturation level and measured DO concentration in water with the same temperature and salinity was estimated as follows:

$$AOU = DO_s - DO_M; \quad (2)$$

where  $DO_s$  is DO concentration at the saturation level (Weiss, 1970) and  $DO_M$  is the measured DO concentration.

Total AOU ( $\text{g m}^{-2}$ ) in the water column was calculated as:

$$AOU_{TOT} = \sum_{z=0}^{z=h} \begin{cases} 0, & \text{if } AOU \leq 0 \\ AOU(z)dz, & \text{if } AOU > 0 \end{cases} \quad (3)$$

where  $AOU(z)$  is the AOU profile and  $h$  is the water column depth.

The upper boundary of the DO depletion was defined as the AOU isoline  $2 \text{ mg l}^{-1}$ . The  $2 \text{ mg l}^{-1}$  criterion was visually chosen from the AOU profiles in order to describe the center of oxycline. Different authors have used various criteria (Lie and Cho, 2002; Liu et al., 2017; Zhang et al., 2007a, 2007b; Zuo et al., 2019) to define water masses in the region. The cold bottom water in the area mostly originates from the KSSW (Liu et al., 2017).

Therefore we define deep layer water as the upper boundary of the Kuroshio subsurface water mass by the isotherm  $24.5 \text{ }^\circ\text{C}$  (Zhang et al., 2007a).

Daily (2015 and 2017 summer) and monthly (2001–2018) river discharge data from Datong hydrological station (see location in Xu et al. (2018)) were used in this study.

The width of the buoyant coastal current was estimated according to Lentz and Helfrich (2002):

$$W_p = (c_w/f) (1 + c_w/c_a), \quad (4)$$

$$\text{where } c_w = (g' h_p)^{1/2}; c_a = \alpha g'/f; h_p = (2Qf/g')^{1/2}; g' = g d\rho/\rho \quad (5)$$

where  $f$  is Coriolis force,  $Q$  is the volume transport (taken as river discharge),  $g$  is the gravitational acceleration,  $d\rho$  is the density difference between the plume and ambient fluids ( $\rho$ ), and  $\alpha$  is the coastal slope.

### 3 Results

#### 3.1. Spatial patterns of hydrography, Chlorophyll *a* and oxygen in 2015 and 2017

In order to link the thermohaline structure and DO observations, we analyzed temperature, salinity, stratification and DO distributions observed in the summers 2015 and 2017. In order to explore the relationship between primary production and subsequent organic matter degradation and oxygen depletion, distributions of Chl *a* and AOU were compared.

The water of SSS  $<30$  was observed along the coastal zone of the entire study area in 2015 (Fig. 3a). The 25-isohaline reached  $29.3^\circ \text{ N}$  in the southern part of the study area, while its northern boundary was at  $31.7^\circ \text{ N}$  near the mouth of the Changjiang Estuary in 2015. In 2017, the 25-isohaline reached only  $30.6^\circ \text{ N}$  in the southern part of the study area (Fig. 3b), and the water fresher than 25 was also observed in the east and northeast of the Changjiang Estuary. Thus, the southward spreading CDW was indicated to prevail before the survey in 2015, but the eastern and northeastern diversion of CDW was prevalent prior to the survey in 2017.

Colder surface water was observed along the coast between 28.5° N and 31° N, and in the northern part of the study area (>31.5° N) in both years. The SST was much lower in these regions, but colder water covered smaller areas in 2017 (Fig. 3d) than in 2015.

The near-bottom temperature difference between the two surveys was particularly large in the northern part of the study area (Figs. 3g–h, 4a, 5a). The near-bottom temperature was 24–25 °C in 20–30 m sea depth in 2015 (Fig. 4a) while it was 21–22 °C in 2017 (Fig. 5a). Likewise, near-bottom water was much saltier in 2017 than in 2015 (Figs. 3e–f, 4d, 5d). The minimum near-bottom temperature zone was located at 50–60 m sea depth in the southern and central part of the study area in 2015 (Fig. 3g and Fig. 4b–c), but it shifted onshore to much shallower depths in 2017 (Fig. 3h and Fig. 5a–c).

Spatial patterns of stratification in the study area were described by the density difference between the surface and near-bottom layer (Fig. 6a–b). Weaker stratification (< 3 kg m<sup>-3</sup>) was observed in the northern part of the study area in 2015 (Fig. 6a) when KSSW was present only at its offshore part (Figs. 4a). Stratification was stronger (>3 kg m<sup>-3</sup>) in the area south of the Changjiang Estuary, being particularly strongest (>6 kg m<sup>-3</sup>) in areas of 20–30 m water depth (Fig. 6a), where the CDW covered the surface (Figs. 4e–f), and a cold and salty water mass occupied the bottom (Fig. 4b–c, e–f). From there strong stratification decline rapidly onshore and offshore (Fig. 6a), because the dense, deep layer water could not intrude into the shallower areas (Fig. 4a–c) and the CDW could not spread further offshore and Fig. 3a, c, e, g). The upper boundary of KSSW had considerable inclination from 24 m depth near the coast to 48 m depth at 80–85 km offshore distance in the southern section in 2015 (Figs. 4c, f and 7c). Opposite inclination (i.e., downwelling) was observed at the central station in 2015, with the boundary at 38 m depth near the coastal slope, tilting offshore towards 17 m depth (Figs. 4b, e). Downwelling had pushed the salinity 30 isoline down to 40 m depth along the central section in 2015 (Fig. 4e). Probably, onshore transport of more saline offshore water over fresher coastal water caused vertical mixing.

Strongest stratification (>6 kg m<sup>-3</sup>) was observed in 2017 in the northern part of the study area (Fig. 6b), where the cold upper boundary of KSSW was usually located at depths of 4–8 m (Figs. 5a, 7d). Above the strong pycnocline was warm and fresh CDW (Fig. 5a, d), whose extent coincided well with the very strongly stratified (>6 kg m<sup>-3</sup>) region (Fig. 3b). Stratification was weaker (<3 kg m<sup>-3</sup>) in the very shallow areas (< 10 m, Fig. 6b) where cold and saltier water mass did not exist, and in the southeastern part of the study area where was influenced by the CDW, but bottom water was warmer than the core of the cold water mass (Figs. 3f, h and 5a–f). Stratification pattern in the central and southern section was characterized by the strong inclination of the isopycnals. The upper boundary of KSSW was located respectively at 50 m and 7–8 m depth at the eastern and western end of the 110-km long southern section (Fig. 5c, f). Surfaced upper boundary of KSSW (i.e., upwelling) was observed at 70–80 km distance in the central section (Fig. 5b). Warmer and fresher surface water was found both onshore and offshore from the upwelling core (Fig. 5b, e). Thus, the coupling of the riverine fresher water and upwelling water could be expected there.

Low DO concentrations (2–3 mg l<sup>-1</sup>) in the near-bottom layer occurred from 25 to 60 m water depth in the southern and central part of the study area in 2015, affecting an area of  $1.3 \times 10^4$  km<sup>2</sup> approximately (Figs. 4h–i, 6e). Higher Chl *a* concentration near the coast in the central and southern part illustrated the role of primary production in the CDW and subsequent processes resulting in oxygen consumption in the deeper layers (Fig. 6c). Although due to downwelling, the low DO water mass was pressed deeper and further offshore while the CDW was constrained closer to the coast (Figs. 4e h). Higher Chl *a* concentration in one station in the northern part (at 31.6° N and 122.6°

E) coincided with the lower (<30) surface salinity intrusion (Fig. 6a). Elevated Chl *a* concentration further offshore (around 31.5° N and 123.2–123.5° E) did not coincide with the CDW (Figs. 3b, 6c). The upper boundary of the DO depletion and upper boundary of KSSW were strongly linked in the southern and central section in 2015 (Figs. 4j–k, 7a, c). However, the boundaries coincided only on the offshore side of the northern section (Fig. 4j). That was the only section shown in Figs. 4–5, where the upper boundary of KSSW and the upper boundary of the DO depletion did not coincide (Fig. 4j). Likewise, this region differed from the rest (except shallower areas with water depth <20 m) of the observations by a high near bottom DO values (Figs. 4g, 6e). Near-bottom DO concentrations were 3–5 mg l<sup>-1</sup> offshore the northern section but increased onshore rapidly in 2015 (Figs. 6e, 4g–i). Low DO area (2–3 mg l<sup>-1</sup>) only overlapped partly with the CDW distribution, but fitted well with the area occupied by cold (< 20 °C) water mass in the near-bottom layer (Figs. 4a–c, 4g–l). The vertically integrated AOU maximum (>150 g m<sup>2</sup>) pattern was observed between 30–31° N and 123–124° E in 2015 (Fig. 6g), which did not coincide with the Chl *a* distribution. In fact, very low Chl *a* concentration was observed at 30.5–31° N and 123–124° E (Fig. 6c) while oxygen depletion (total AOU) was rather high (Fig. 6e). It was therefore hypothesized that oxygen depletion in the southern and central part could have developed over longer periods.

Two low DO concentration zones were observed in the near-bottom layer in 2017 (Figs. 5g–i and 6f). The one in the north well overlapped with the low salinity CDW and strong stratification area (Fig. 6b, 6f). Hypoxia there started at 4–8 m depth already and was closely linked to the location of the pycnocline (Figs. 5a, 5d, 5g). Lowest DO and highest AOU of all our measurements were observed in the northern part in 2017 (Figs. 5g, j). Some profiles showed DO nearly 0.6 mg l<sup>-1</sup> and AOU over 6 mg l<sup>-1</sup>. High Chl *a* concentration was observed in the northern part, displaying the intense production in the upper layer (Fig. 6d). The second low DO concentration zone was found in shallow depths (10–30 m) at the coastal slope in the southern and central part of the study area and overlapped with lower SST and SSS upwelling region (Figs. 5b–c, e–f, h–i). This indicates that coupling between coastal upwelling, which bring subsurface water to shallower depths (euphotic zone), and CDW spreading created favorable conditions for the primary production in the upper layer and subsequent DO consumption and depletion in the deep layer (Figs. 5h, i, k, l). Higher Chl *a* concentration in the upwelling areas were revealed from the satellite image, although not as high as those in the north (Fig. 6d). The vertically integrated AOU had the highest values (>150 g m<sup>2</sup>) in the northern part of the study area in 2017 (Fig. 6h).

The high AOU zones in the near-bottom layer could be found at different water depths, e.g., as shallow as 4 m in 2017 (Figs. 4j–l, 5j–l). Strong and shallow enough stratification makes DO depletion possible in such a shallow depth. The question is whether the DO depletion is linked to a certain physical property of water, e.g. to a temperature-, salinity-, or density isocline. We found high correlation ( $r^2 = 0.81$ ,  $p < 10^{-10}$ ,  $n = 44$  in 2015, and  $r^2 = 0.98$ ,  $p < 10^{-10}$ ,  $n = 34$  in 2017) between upper boundary of KSSW and upper boundary of the DO depletion depth. Thus, the upper boundary of KSSW determines the vertical location of the upper boundary of the DO depletion largely. The cold KSSW is denser than the water above, and the resultant thermocline/pycnocline provides a barrier of vertical mixing to develop DO depletion below. There was an inclination of both isoclines in both years (Fig. 7). The inclination was particularly steep in 2017. Both isoclines were located >50 m depth in the southeastern corner of the study area, while in the shallow areas in the west and north the same isoclines were at 4–8 m depth (Figs. 7b, d). Such an inclination of isotherms and cold SST near the coast indicated upwelling. Deeper colder water intruded along the coastal slope shallower to replace the former upper layer water. The inclination of isoclines was not that pronounced and it was rather north-south directed in 2015 (Figs 7a, c). Lower correlation in

280 2015 was caused by the discrepancy of the two isolines in the northern part in 2015. This indicates some other processes than Kuroshio water intrusion, e.g. vertical mixing, vertical movement of pycnoclines or advection altered subsurface conditions there.

It is noteworthy that the high vertically integrated AOU (Fig. 6g) area overlapped with a shallow position of isolines in 2015 (Figs. 4k, 7a, 7c.). It means that high total AOU there (comparing to neighbouring areas) was related both to the thick subsurface DO depleted layer and to the higher (comparing to the surrounding area) AOU. 285 We have captured two completely distinct situations of stratification and DO distributions in summers 2015 and 2017. The differences between the two years are summarized in Table 1. Prevailing of either pattern in the physical fields should lead to varied biogeochemical fields and ecosystems in the area. The main questions addressed next are: what are the main drivers causing the distinct patterns? and what is the forcing behind the formation of the patterns?

290

### 3.2. Forcing behind distinct patterns in 2015 and 2017

The observed discrepancies of general features in the two cruise surveys can be driven by several forcing mechanisms. Two main factors can be considered to affect the distribution of CDW in the region: Changjiang River discharge and wind forcing. The riverine water is a source of buoyancy flux, causing the barotropic pressure gradient. Latter drives a current to flow south along the coast. Mean discharge at Datong hydrological station in the two summers (1 June to 31 August) was rather similar, 44 000–45 000 m<sup>3</sup>s<sup>-1</sup>. We consider the time lag of 1 week for water propagation from Datong to the river mouth (Li and Chen, 2019) in further discussion.

300 In comparison, river discharge in June was higher in 2015 and in July was higher in 2017. Daily discharge peaked at 58 000 m<sup>3</sup>s<sup>-1</sup> on 1 July in 2015 and at 70 000 m<sup>3</sup>s<sup>-1</sup> on 13 July in 2017. According to typical residual current velocities off the estuary (Peng et al., 2017) and along the coast in the south (Wu et al., 2013), it should take a few days to weeks for the water from the river mouth to reach offshore as registered in 2017 or to the southern part in 2015. We assume that seven days of forcing (accumulated run-off or wind forcing) is enough to alter the field 305 distributions off the estuary. Mean river flow in the week ahead of the surveys was 30 000 m<sup>3</sup>s<sup>-1</sup> in 2015 and 65 000 m<sup>3</sup>s<sup>-1</sup> in 2017, respectively. Variations in discharge influence the cross-shore extent of plume and the width of the buoyant current. According to the Eq. (4), the width  $W_p$  of buoyant current would be 77 km in the case of ambient and CDW water density difference of 10 kg m<sup>-3</sup>, coastal slope of 10<sup>-4</sup> and discharge of 65 000 m<sup>3</sup> s<sup>-1</sup> before our survey in 2017. The width would be 55 km for the inflow volume of 30 000 m<sup>3</sup> s<sup>-1</sup> in 2015. Thus, the 310 variability in the river discharge only partly explains the discrepancy between the two surveys, particularly more extensive spreading of the CDW eastward in 2017. However, it does not explain the northeast advection of the CDW that far in 2017. Likewise, it remains unclear why southward transport of the CDW was much more limited in 2017 than 2015. Several indicators hint that wind-driven transport could have caused the differences between the 2015 and 2017 observations, including stronger crossshore inclination of the pycnocline, offshore advection 315 of the CDW, and onshore advection of the colder sub-pycnocline water in 2017. Such features were not found during the survey in 2015, but southward advection of the CDW revealed instead. Since our surveys are conducted annually one month apart, the differences between the two surveys might be associated with the annual cycle in



wind climate. Our hypothesis is that the dominant factor behind the discrepancy of the two surveys is wind forcing and possibly its seasonality.

320 As described previously, we assume the thermohaline fields in the study area were established by the forcing one week before the sampling. Mean wind, sea level anomaly, surface currents and bottom currents during seven days before the surveys are presented in Fig. 8. The mean wind direction was from the northeast before the 2015 survey (Fig. 8a) and from the south before the 2017 survey (Fig. 8b). Northeasterly wind should cause downwelling and accelerate alongshore buoyant current toward the south. This current was clearly visible in the mean surface current  
325 plot in 2015 (Fig. 8c), along the coast from 32.5° N to 28° N. Such coastal current did not prevail before the 2017 survey (Fig. 8d). Weak mean flow towards the east or northeast could be noted near the mouth of the Changjiang River in 2017 (Fig. 8d). Probably, the buoyant coastal current was altered, and the CDW was transported across-shore by southerly wind forcing. Thus, simulated current patterns and wind forcing before the two surveys were qualitatively in accordance with observed salinity fields in the surface layer (Figs. 3a–b). The main difference  
330 between the two years in the bottom layer was the stronger northward flow in 2017 (Figs. 8e–f). Penetration of the deep layer water to the shallower areas, towards the river mouth, occurred in 2017 (Fig. 8f). This coincided with our observations in terms of the stronger inclination of clines (Figs. 7b, d) and presence of the colder KSSW water mass in the northern part of the study area in 2017 (Fig. 3h). Because southerly wind forced offshore transport of the surface water, deep layer water intruded to the shallower depths for compensation in 2017. This suggested the  
335 pattern of processes based on field surveys and modelling data can be confirmed by mean sea level anomaly during the surveys (Figs. 8g, h). There is a strong sea level gradient between nearshore and offshore in both years. Higher sea level can be found along the coast from the northern boundary of the study area to 30° N in the south in 2015, and as a result, the geostrophic current is southward. This is the combined effect of the lighter riverine water and convergence of the shoreward transported upper layer water. This is in accordance with our observation of the  
340 downwelling in the central section (Figs. 4b, e, h, k) while such a phenomenon did not occur in the southern section (Figs. 4c, f, i, l) in 2015. Downwelling co-occurred with the alongshore southward current in the northern part according to the sea level anomaly (Fig. 8g) and modelled current field (Fig. a, c). This could explain the well ventilated and warmer near-bottom waters in the north in 2015 (Fig. 4a, g, j). An opposite gradient, i.e. lower sea level anomaly near the coastline, was observed in 2017. This is a typical sign of divergence, i.e. offshore transport  
345 of the surface water and consequent upwelling along the coastline. The zonal pattern of sea level from the river mouth to offshore (high-low-high sea level, Fig. 8h) agrees with our observation of the upwelling in the central section (Figs. 5b, e). As a result of sea level anomalies, geostrophic current toward the north at 28.5–31° N and to offshore near the river mouth were observed in 2017.

### 3.3. Variability and drivers of the Changjiang Diluted Water spreading

350 In order to verify the sensitivity of the buoyant coastal current to wind forcing, simulated current and wind data from June to September in 1993–2018 were used. First, we calculated daily mean meridional (alongshore) upper layer (0–5 m) current velocity component  $v_m$  at the zonal section of 31° N (section S1) from 122.1° to 122.6° E (see red line in Figs. 8c–d). Secondly, the correlation was calculated between the current velocity component  $v_m$   
355 and wind (see the location in Fig. 1) from different directions (full circle by 10° steps) and by different time lags shifted by one day step. The best correlation ( $r^2 = 0.76$ ,  $p < 10^{-10}$ ,  $n = 3025$ ) was found with the one-day (prior to

the modeled current value) mean SE–NW wind velocity component  $w_c$  (Fig. 9). Thus, the meridional current along the coast correlates well with wind.

360 Simulated near-bottom currents (Fig. 8f) suggest that cold and saltier water (comparing to 2015 observation) was advected to the northern part of the study area as compensation flow to the near-surface offshore flow in 2017. Moderate winds before the 2017 survey (Fig. 10a) could not mix the cold and DO depleted deep layer water into the upper layer. However, stronger wind impulse with speeds up to 11–12 m s<sup>-1</sup> occurred a few days before the start of the survey in 2015 (Fig. 10a). Upper mixed layer depth created by wind stirring was estimated by the formula describing the turbulent Ekman boundary layer  $h = 0.1 u_* / f$  (Csanady, 1981), where  $u_* = (|\tau| / \rho_w)^{1/2}$  is  
365 friction velocity,  $\rho_w$  is water density,  $\tau$  is wind stress, and  $f$  is Coriolis parameter. Vertical mixing reached down to 15–16 m depth according to this empirical method before the 2015 survey. Note that our estimation is based on 6-h average wind speed. Stronger winds, which occur on shorter time scales, probably causing deeper mixing. Thus, wind stirring could weaken the stratification before the 2015 survey, as evident by the presence of warmer, fresher water with higher DO concentration in the deep layer. However, advection from the north (Fig. 8c) and  
370 downwelling (Fig. 8g) likely also contributed to the weakening of stratification and ventilation of the northern part in 2015. Water from the north is colder and causes convective mixing and weakening of the thermocline. The effect of downwelling can be seen in the central section (Fig. 4b, e, h, k). Sea bottom is flatter in the northern part and therefore the sub-thermocline distributions in the case of downwelling showed up there as lateral gradients (Figs. 4a, d, g, j).

375 We further analyzed time series of wind speed (Fig. 10a), SE–NW wind stress component ( $\tau_c$ , Fig. 10b), meridional current velocity component ( $v_m$ , Fig. 10c) at the section S1, and the Changjiang river run-off (Fig. 10d) in June–September 2015 and 2017. Wind stress  $\tau_c > 0$  and a northward current component prevailed from June to early August in 2017 (Fig. 10b). Larger wind stress events alternated with calmer periods from June to late August in 2015. During the periods of  $\tau_c < 0$  or weak northward winds, the current was often directed southward. The  $v_m < 0$   
380 occurred more frequently in June–July 2015 compared to the same period in 2017 (Fig. 10c). Latter is reflected in salinity distributions (Fig. 3a–b). We defined the CDW as the salinity of  $< 30$  and calculated its monthly spatial occurrence from June to September in 2015–2018 using satellite surface salinity distributions (Fig. 11). The lowest areal extent of the CDW was observed in September in both 2015 and 2017. The CDW was mainly transported to the northeast and east in all summer months in 2017. Northeastward transport of the CDW could also be noted in  
385 2015, but the CDW occupied areas in the southern part of the study area as well. Thus, the satellite-derived salinity confirms our in-situ observations of the southward flow of the CDW. We can conclude that our survey in 2017 well described the prevailing situation of the CDW spreading in summer 2017. In 2015, we captured the situation, where the CDW spread to the south, but during summer, northeastward spreading occurred as well (Fig. 11). The CDW spreading in 2016 and 2018 was rather northeastward (similar to 2017). It is noteworthy that coverage of  
390 the CDW was clearly larger in 2016 than we observed in 2017 while it was more limited in 2018. The CDW spreading presented in Fig. 11 is further analyzed in relation to inter-annual wind and river forcing, and in the context to previous studies in the discussion.

Next, we attempted to link the CDW spreading, i.e. coastal current velocities to the wind forcing in the period  
395 June–September in 1993–2018. We grouped the simulated meridional currents  $v_m$  at section S1 (see Fig. 8c-d) to the wind velocity  $w_c$  classes with the step of 0.25 ms<sup>-1</sup> and took an average of each group. By doing this, we found

relation between mean  $w_c$  and  $v_m$ . If the daily mean wind velocity component  $w_c=0 \text{ m s}^{-1}$ , current velocity component  $v_m$  would be on average  $-13 \text{ cm s}^{-1}$  (i.e., southward) at S1. We can assume that this is the mean geostrophic component of the current velocity. Daily mean wind velocity component  $w_c$  of  $5.6 \text{ m s}^{-1}$  is required to reverse the current and have the same magnitude at S1 ( $v_m = +13 \text{ cm s}^{-1}$ ). This corresponds roughly to the wind stress  $\tau_c$  of  $0.04 \text{ N m}^{-2}$ . Wind stress  $\tau_c$  of  $0.02 \text{ N m}^{-2}$  corresponds to the  $v_m=0 \text{ cm s}^{-1}$ , i.e., coastal current is altered. We define  $v_m=0 \text{ cm s}^{-1}$  as the term “alteration of the coastal current” in further analysis/discussion in this paper. There are two main wind-driven processes that contribute to the alteration of the coastal current. First is the direct effect of wind stress and resulting Ekman surface current. Secondly, southerly winds cause divergence and upwelling at the coast and alter sea level distributions. When upwelling is evoked, alongshore geostrophic current, resulting from the cross-shore gradient between colder-denser and warmer-lighter offshore water, is rather to the north. The mean sea level anomalies during our surveys in 2015 and 2017 illustrate well this phenomenon (Fig. 8g–h). The joint effect of the two processes explains why relatively low wind stress is required for the alteration of the circulation regime. We analyzed the simulated current data in June–September from 1993 to 2018 and averaged the cases when daily mean wind stress component  $\tau_c$  was  $< 0$  and  $\geq 0.02 \text{ N m}^{-2}$  (Fig. 12). The same circulation features, but more pronounced, stand out as before the surveys in 2015 and 2017 (Fig. 8). In the case of  $\tau_c \geq 0.02 \text{ N m}^{-2}$  surface flow was northward along the coast, to the northeast near river mouth (Fig. 12a) while shifted north or towards the river mouth in the deep layer (Fig. 12c) as before our survey in 2017 (Fig. 8d, f). In the case of  $\tau_c < 0 \text{ N m}^{-2}$  southerly flow prevailed in the surface layer along the coast (Fig. 12b), and the mean currents in the near-bottom layer were rather weak (Fig. 12d) as before our survey in 2015 (Figs. 8c, e). This supports our hypothesis that the wind field is a key forcing driver for the direction of the Chinese Coastal Current controlling the CDW distribution off the Changjiang Estuary, and thus hypoxia formation (Fig. 8a–f).

## Discussion

The balance between oxygen consumption caused by the decomposition of organic matter by bacteria in the sub-surface layer and oxygen supply controls the formation of hypoxia. There is a strong link between primary production in the upper layer and oxygen consumption in the near-bottom layer (Chen et al., 2017, 2004; Wang et al., 2017, 2016; Wei et al., 2017a; Zhou et al., 2019). The processes accompanied by enhanced primary production, the spreading of the riverine nutrient-rich CDW and KSSW upwelling, are dependent on wind and river forcing. Likewise, physical processes, which can supply deep layers with oxygen– lateral advection and vertical mixing, are partially driven by wind forcing.

Two distinct stratification, Chl *a* and DO distribution patterns were observed in the area off the Changjiang Estuary in the summers of 2015 and 2017. There was a stronger DO depletion in the southern part of the study area in August–September 2015, while intense hypoxia was observed in the northern part of the study area in July 2017. Likewise, remarkable DO depletion was found in the southern part in the upwelling region in 2017. Similar hypoxia patterns in August and October were observed in 2006 (Zhu et al., 2011). The main reasons for the different spatial hypoxia patterns are variations in the wind forcing and river discharge. Southeasterly winds and higher discharge during summer monsoon both favor cross-shore transport of CDW and development of hypoxia in the northern part of the study area. Both the maximum frequency of southerly wind and river discharge in the

435 annual cycle occurred in July–August (Figs. 13a, c). In accordance with the annual cycle of wind (Fig. 13a), the highest frequency of upwelling events occur in July–August as well (Xu et al., 2017). Thus, our observations conducted in late August/early September 2015 and late July 2017 illustrate the seasonal cycle of forcing and are reflected in oxygen and stratification patterns. Summer monsoon and river discharge were close to average in the whole summer of 2017, while the summer monsoon was clearly weaker in 2015 (Fig. 13a–b). Wang et al. (2012) explained the annual cycle of the location of near-bottom DO minimum by the seasonal change of stratification due to river discharge and warming/cooling of the upper layer. We suggest that the southern location of the DO minimum in June and September, and the northern location, as observed in July–August 2006 (Fig. 3 in Wang et al. (2012), are also related to the seasonality of wind forcing, surface transport of the CDW and compensating KSSW intrusion in the deep layer.

440 The monthly occurrence of CDW clearly shows that CDW transport offshore (to northeast) prevailed during summer periods (Fig. 11). However, one can see that 2015 differed with low values in the inter-annual time series of wind stress of  $\tau_c \geq 0.02 \text{ N m}^{-2}$  (Fig. 13a–b) and with considerable southward advection of the CDW (Fig. 11). Monthly maps of bottom oxygen in 2015 (Li et al., 2018b) do not show significant oxygen depletion in the north in any month while deteriorated hypoxia (compared to our observations) occurred in the south in October (Li et al., 2018b). This well demonstrates that the wind regime plays an important role in the inter-annual variations of the hypoxic area location (Fig. 6e–f). The year 2016 stands out in terms of wind forcing close to the inter-annual mean (Fig. 13b), but with very high river discharge (Fig. 13d). One can see that offshore advection prevailed in 2016 (Fig. 11). The latter indicates that if wind forcing is close to the long-term average, hypoxia will more likely occur east of the river mouth (northern part of the study area) and hypoxia occurrence in the south is more limited.

450 That was confirmed by our observations in 2017. River discharge and wind stress  $\tau_c \geq 0.02 \text{ N m}^{-2}$  were both similar to the long term mean in 2017 (Fig. 13), and we observed hypoxia and CDW water in the northern part. It means, hypoxia in the southern part (as we found in 2015) occurs if the summer monsoon is considerably weaker than the long-term mean. Indeed, hypoxia in the south has been quite rare in 1998–2013 (Chen et al., 2017). As a consequence of large river runoff, the largest CDW extent occurred in the year 2016 out of the four years (Fig. 11). The occurrence of wind stress  $\tau_c \geq 0.02 \text{ N m}^{-2}$  was higher than average (Fig. 13b) and river discharge was lower than average in 2018 (Fig. 13d), and this reflected well in the CDW distribution (Fig. 11). One can see that CDW covered the smallest area out of four years, and the prevailing transport direction was towards the north or northeast (Fig. 11). From our stratification and DO distribution observations (Figs. 3–7) and their dependence on wind forcing (Figs. 8–10, 12), and CDW spreading direction and extent (Fig. 11), DO depleted area could have occurred in the northern part both in 2016 and 2018, with larger extent in the former year. Hypoxia in the northern part in 2016 has reported (Zhang et al., 2019). Lower than average river discharge and close to average wind stress of  $\tau_c \geq 0.02 \text{ N m}^{-2}$  (as in 2016) occurred in 2006 (Figs. 13b, d) when hypoxia registered in north (Zhu et al., 2011). We estimated that wind stress of  $\tau_c \geq 0.02 \text{ N m}^{-2}$  is necessary to alter the geostrophic coastal current and create favorable conditions for hypoxia in the northern part. In accordance with higher inter-annual occurrences of wind stress  $\tau_c \geq 0.02 \text{ N m}^{-2}$  in July–August (Fig. 13a), the larger hypoxic area in the north was documented for instance in August 2012 (Li, 2015) and August 2013 (Ye et al., 2016; Zhu et al., 2017). Contrary, in summers when the occurrence of  $\tau_c \geq 0.02 \text{ N m}^{-2}$  was lower considerable oxygen depletion was also expected to have occurred in the southern part, as observed, e.g. in 1998 (Wang and Wang, 2007) and 1999 (Li et al., 2002). However, on the top of the inter-annual variability and seasonality, synoptic-scale changes of wind-driven currents and river forcing

475 likely to influence the distributions on shorter time scales (Fig. 10). Thus, when planning measurement campaigns, it is worthwhile to take into account wind-driven transport, river discharge, remotely sensed salinity and altimetry to forecast spreading of the CDW; as well upwelling occurrence and deep water intrusion. That allow to estimate potential hypoxic area and therefore more efficient use of ship time and more detailed sampling of the hypoxic area.

480 The fate of the river plume can be separated by two regions and processes: circulating bulge near the river mouth and downstream coastal current (Fong and Geyer, 2002; Horner-Devine, 2009). It has been estimated that about 80–90% of the river discharge accounts to freshwater transport of the coastal current (Li and Rong, 2012; Wu et al., 2013). That means that most of the river discharge does not remain in the river plume bulge, but impacts the surrounding areas off the estuary. We estimated the southward geostrophic component of the current velocity to

485 be on average  $13 \text{ cm s}^{-1}$  at section S1. The southward coastal current was measured in winter by Wu et al. (2013). They estimated the maximum current speed up to  $50 \text{ cm s}^{-1}$ , including also the wind-driven component, which in winter supports flow to the south (Wu et al., 2013).

Offshore, east- or northeastward advected CDW caused by southerly wind, as we observed in 2017, might form detached eddies due to interaction of the Ekman flow and density-driven frontal currents (Xuan et al., 2012). Those

490 eddies bring CDW further offshore and alter physical, chemical characteristics (including oxygen conditions) and primary production in the water column (Wei et al., 2017).

Offshore transported CDW co-occurred with coastal upwelling caused by southerly wind in 2017. Shoreward, upslope penetration of the sub-thermocline KSSW and hypoxia in the upwelling – CDW coupling zone (Wei et al., 2017a) were observed. Upwelling, induced by southerly winds and its relaxation supported by northerly winds,

495 was also found by Yang et al. (2019). Wei et al. (2017) showed that the coupling of the CDW plume front and KSSW upwelling caused DO minimum at the sloping bottom. Likewise, nitrogen-to-phosphorus ratio and primary production in the CDW is considerably modified by upwelling (Tseng et al., 2014). Phosphate transport by upwelling reduces phosphorus deficiency in the CDW water and therefore promote phytoplankton growth and nitrate uptake (Chen et al., 2004; Zhou et al., 2019).

500 Time series of wind displayed large variations in wind forcing on shorter time scales (days to weeks) (Fig. 10) which may alter the stratification pattern and DO distributions considerably. Numerical simulation by Zhang et al. (2018) showed that wind-induced redistribution of the Changjiang River plume changes near-bottom DO conditions rapidly. Also, it has been shown that vertical mixing caused considerable variations in DO concentrations in the near-bottom layer (Ni et al., 2016). In July 2015, 1.5 months before our survey, hypoxia was

505 terminated by a typhoon, but two days later, hypoxic conditions were re-established (Guo et al., 2019). Our field measurements showed that the upper boundary of the DO depletion is strongly linked to the upper boundary of KSSW (Fig. 7). Besides enhanced (or impeded) vertical diapycnal mixing, DO conditions can be altered by vertical movement of this water mass. Ni et al. (2016) published a valuable time-series of DO in the near-bottom layer. They linked the increase of DO concentrations in the near-bottom layer with the vertical mixing and DO decline

510 in the near-bottom layer with the primary production and consequent decomposition of detritus. One can see several cases from their time series data (Fig. 2 in Ni et al., 2016) when the near-bottom temperature drops. Those events must be related to the advection of colder water and uplift of the KSSW as DO declined during those events. Penetration of the cold, low DO water upwards along the coastal slope appears when temperature and DO decline in the point measurement time series. There are some registered events when the near-bottom layer temperature

515 rises, but sea surface temperature does not change much (e.g. Fig. 2 in Ni et al. (2016)). In these cases, DO concentrations increased in the near-bottom layer. Such events can be associated to the downward movement of the thermocline or advection of warmer water as we observed in the central section in 2015 (Figs. 4b, e, h, k) than vertical mixing. Thus, vertical location and movement of the thermocline have an important role in the near-bottom DO distributions at the coastal slope. The importance of KSSW thickness on the oxygen depletion is also revealed  
520 when near-bottom oxygen maps are compared with the maps of vertically integrated AOU (Figs. 6g–h). Bottom hypoxia in the north in 2017 was much more intense compared to hypoxia in the south in 2015. However, the total AOU was similar in hypoxic zones in both years due to thicker oxygen-depleted layer, i.e. thicker KSSW in the south in 2015.

Pycnocline dynamics, including downwelling studies, as suggested by Hu and Wang (2016), are essential in future  
525 investigations. Those studies are difficult to arrange with conventional research vessel surveys only. Autonomous measurement platforms, such as profiling moorings (Lips et al., 2016; Sun et al., 2016), moored sensor chains (Bailey et al., 2019; Venkatesan et al., 2016), which allow capturing the variability in necessary spatio-temporal resolutions can be used. Underwater gliders (Liblik et al., 2016; Rudnick, 2016) might be complicated to use due to strong tidal velocities and heavy ship traffic, but are worth considering as well.

530 Our observations indicated that two conditions in the water column must be present for the development of hypoxia. First, high AOU occurred only below the thermocline, in the cold KSSW. Secondly, primary production in the fresher CDW or upwelling is needed to cause high DO consumption that leads to hypoxia. Interestingly, these two conditions for hypoxia were valid in very different situations. Shallow (4–6 m depth) and sharp thermocline coincided with the halocline (related to the CDW) in the northern part of the study area in 2017.  
535 Contrary, thermocline and halocline were separated in the southern part of the study area in 2015. Thus, the vertical coincidence of halocline and thermocline is not necessary for the hypoxia formation. The thermocline acts as a physical barrier, which impedes vertical mixing and DO exchange with upper layers. Primary production in the CDW or the upwelled water and related DO consumption in the near-bottom layers (Wang et al., 2017, 2016) cause DO decline. Zhu et al. (2016) found that the area of DO concentration  $< 3.0 \text{ mg l}^{-1}$  fits relatively well with  
540 the region of the pycnocline strength  $> 2.0 \text{ kg m}^{-3}$ . This relationship between pycnocline strength and low DO only partly holds according to our analysis (Figs. 6a–b, e–f). For instance, there was strong stratification (Fig. 6a) near the river mouth (around  $122.5^\circ \text{ E}$  and  $31^\circ \text{ N}$ ) in 2015 due to presence of the CDW, but hypoxia was not observed (Fig. 6e) since the colder KSSW was not present. Most of the study area, except very shallow areas, had strong enough stratification (pycnocline strength  $> 2.0 \text{ kg m}^{-3}$ ) in 2017 (Fig. 6b). However, hypoxia in the southern  
545 part in 2017 was only observed near the coast and in connection to the coastal upwelling (Fig. 6f). Rest of the study area in the south was free of hypoxia. In the latter area, KSSW was present, but there was no CDW in the surface layer. That means that strong stratification as such does not necessarily lead to hypoxia. Two features must be present for hypoxia formation: 1) KSSW, 2) CDW and/or subsurface water upwelling. We can conclude that colder KSSW determines where (including at what depths) hypoxia could develop, i.e. provides the necessary  
550 precondition for hypoxia. The CDW spreading and/or sub-surface water upwelling (and related biogeochemical, biological processes) determine the magnitude, exact location and timing of oxygen depletion. Both features are strongly impacted by wind.

Besides the barrier effect of the thermocline, the intrusion of KSSW has another impact on oxygen dynamics. First, the subsurface water is oxygen-depleted already before the local impact of oxygen consumption (Qian et al., 2017).

555 The furthest stations in the southeast (Fig. 1) had AOU of 2–2.5 mg l<sup>-1</sup> in the deep layer in 2017, i.e. in the same order that has been estimated in the KSSW before (Qian et al., 2017). The total AOU in the water column there was 50–60 g m<sup>-2</sup> (Fig. 6h). This water is still rather well ventilated compared to the deep layer waters that had been impacted by upwelling or CDW induced production in the surface layer (Fig. 6g–h). Despite its initial oxygen depletion, Kuroshio intrusion is an important source of oxygen import to the study area (Zuo et al., 2019). Without  
560 this lateral DO supply, hypoxia could form much faster in a larger area (Zuo et al., 2019). Kuroshio intrusion is nutrient-rich (Zhang et al., 2007b; Zhou et al., 2019) and its upwelling or vertical mixing could intensify primary production in the surface layer; consequently organic matter sinking causes DO consumption in the near-bottom layers.

We have already outlined the main difference in the wind forcing, causing the formation of the hypoxia in the  
565 southern and northern parts of our study area. The one in the north develops under the conditions of the summer monsoon. Intensive hypoxia can start from very shallow depths (at 4–8 m) in the northern area (Fig. 5g), and it can develop very fast under favorable conditions (Guo et al., 2019). At the same time, the bottom layer there can be easily ventilated by wind stirring or downwelling as we observed at N15 section in 2015 (Fig. 4g). Thus, hypoxia in the northern part can be very pronounced but disappears fast if forcing changes (Ni et al., 2016).

570 Hypoxia in the southern part of the study area was not as pronounced but quite stable as noted already by Zhu et al. (2011). The decline of the DO from March to October in the southern part was demonstrated by monthly measurements (Li et al., 2018b). We suggest three main reasons for this. First, favorable wind conditions for the southward transport are not that frequent in summer. Second, the mixture of CDW with ambient ocean water promotes primary production and subsequent settling of organic matter on its way south. So less detritus sinks to  
575 the bottom layer and DO consumption is lower in the southern part compared to the northern region. Third, the KSSW and related thermocline are located deeper in the south, and therefore wind mixing does not diminish the thermocline. The first two reasons account for weaker hypoxia and the third for long-lasting hypoxia in the southern part.

580

## Conclusions

585 Two main conditions in the water column must be present for the occurrence of hypoxia in the near-bottom layer off the Changjiang Estuary: enhanced primary production in the upper layer and KSSW intrusion in the near-bottom layer. Pycnocline created by KSSW intrusion or CDW spreading is the precondition and determines where hypoxia could develop. Primary production in the CDW and/or in the upwelled water, and consequent oxygen consumption by sinking organic matter below the pycnocline determine the magnitude, exact location and timing  
590 of oxygen depletion. Both the distribution of CDW and KSSW and the occurrence of KSSW upwelling are controlled by the wind field.

Summer monsoon (wind from south) alters Chinese Coastal Current by creating an Ekman surface flow and by changing the geostrophic current flowing to north or northeast. As a result, the CDW spreads offshore, and KSSW

intrudes northwards and upwards on the coastal slope, resulting in upwelling. The joint effect of these processes  
595 can lead to intense and shallow (4–8 m) oxygen depletion in the north and hypoxia at the coastal slope in the south.  
Northerly wind intensifies Chinese Coastal Current and CDW spreading to south, and causes downwelling.  
Consequently, the northern part is well ventilated and hypoxia rather occurs further offshore in the southern part.  
Wind forcing and river runoff are important contributors of inter-annual and seasonal variations, determining the  
size and location of low DO areas. The DO minimum is located more likely in the northern part in July–August  
600 and in the southern part during rest of the stratified period.

There is a strong connection between the upper boundaries of KSSW intrusion and oxygen depletion. The  
sensitivity of the boundaries to wind forcing shapes oxygen conditions considerably in the area. Autonomous  
measurement campaigns by mooring arrays and underwater gliders could considerably improve the knowledge  
about related processes. Our results further suggest that a combination of wind field data, river discharge, remotely  
605 sensed sea-surface salinity and sea level could be used to forecast the hydrographic conditions and potential  
hypoxic area before field work.

610

615

*Code availability.* Scripts to analyze the results are available upon request. Please contact TL.

*Author contributions.* TL led the analyzes of the data and writing of the manuscript with contributions of DF and  
620 YW. DF was responsible for arranging oceanographic cruises. DS measured chlorophyll *a* during the 2015 cruise.

*Competing interests.* We declare that no competing interests are present.

*Acknowledgments.* This research was funded by the National Natural Science Foundation of China (41776052),  
625 Qingdao National Laboratory for Marine Science and Technology (MGQNLN-TD201802), and the Research  
Fund of State Key Laboratory of Marine Geology at Tongji University (MG20190104).

River data was downloaded from Bulletin of River Sediment in China provided by the Ministry of Water Resources  
of the People's Republic of China (MWR). Website: <http://www.mwr.gov.cn/sj/tjgb/zghlnsgb/> (visited 7<sup>th</sup>  
September 2019). We thank Yue Zhang for gathering CTD data in 2015, Junbiao Tu for providing mooring data,  
630 Huiping Xu for agreeing to use their chlorophyll *a* data. We would like to thank our colleagues, who helped us in  
performing measurements. We thank Jaan Laanemets for his comments on the manuscript. We thank reviewer



Fabian Große and an anonymous reviewer for the valuable comments and suggestions that helped to improve the manuscript.

635


## References

- Altieri, A. H. and Gedan, K. B.: Climate change and dead zones, *Glob. Chang. Biol.*, 21(4), 1395–1406, doi:10.1111/gcb.12754, 2015.
- 640 Association for the Physical Sciences of the Sea, I.: The international thermodynamic equation of seawater-2010: Calculation and use of thermodynamic properties Intergovernmental Oceanographic Commission., 2010.
- Bailey, K., Steinberg, C., Davies, C., Galibert, G., Hidas, M., McManus, M. A., Murphy, T., Newton, J., Roughan, M. and Schaeffer, A.: Coastal Mooring Observing Networks and Their Data Products: Recommendations for the Next Decade, *Front. Mar. Sci.*, 6, 180, doi:10.3389/fmars.2019.00180, 2019.
- 645 Beardsley, R. C., Limeburner, R., Yu, H. and Cannon, G. A.: Discharge of the Changjiang (Yangtze River) into the East China Sea, *Cont. Shelf Res.*, 4(1–2), 57–76, doi:10.1016/0278-4343(85)90022-6, 1985.
- Chen, J., Pan, D., Liu, M., Mao, Z., Zhu, Q., Chen, N., Zhang, X. and Tao, B.: Relationships Between Long-Term Trend of Satellite-Derived Chlorophyll-a and Hypoxia Off the Changjiang Estuary, *Estuaries and Coasts*, 40(4), 1055–1065, doi:10.1007/s12237-016-0203-0, 2017.
- 650 Chen, Y.-L. L., Chen, H.-Y., Gong, G.-C., Lin, Y.-H., Jan, S. and Takahashi, M.: Phytoplankton production during a summer coastal upwelling in the East China Sea, *Cont. Shelf Res.*, 24, 1321–1338, doi:10.1016/j.csr.2004.04.002, 2004.
- Chu, P., Yuchun, C. and Kuninaka, A.: Seasonal variability of the Yellow Sea/East China Sea surface fluxes and thermohaline structure, *Adv. Atmos. Sci.*, 22(1), 1–20, doi:10.1007/BF02930865, 2005.
- 655 Conley, D. J., Björck, S., Bonsdorff, E., Carstensen, J., Destouni, G., Gustafsson, B. G., Hietanen, S., Kortekaas, M., Kuosa, H., Markus Meier, H. E., Müller-Karulis, B., Nordberg, K., Norkko, A., Nürnberg, G., Pitkänen, H., Rabalais, N. N., Rosenberg, R., Savchuk, O. P., Slomp, C. P., Voss, M., Wulff, F. and Zillén, L.: Hypoxia-Related Processes in the Baltic Sea, *Environ. Sci. Technol.*, 43(10), 3412–3420, doi:10.1021/es802762a, 2009.
- Csanady, G. T.: Circulation in the Coastal Ocean, *Adv. Geophys.*, 23, 101–183, doi:10.1016/S0065-2687(08)60331-3, 1981.
- 660 Cui, Y., Wu, J., Ren, J. and Xu, J.: Physical dynamics structures and oxygen budget of summer hypoxia in the Pearl River Estuary, *Limnol. Oceanogr.*, 64(1), 131–148, doi:10.1002/lno.11025, 2019.
- Czeschel, R., Schütte, F., Weller, R. A. and Stramma, L.: Transport, properties, and life cycles of mesoscale eddies in the eastern tropical South Pacific, *Ocean Sci.*, 14(4), 731–750, doi:10.5194/os-14-731-2018, 2018.
- 665 Diaz, R. J. and Rosenberg, R.: Spreading dead zones and consequences for marine ecosystems., *Science*, 321(5891), 926–9, doi:10.1126/science.1156401, 2008.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E. and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system, *Remote Sens. Environ.*, 116, 140–158, doi:10.1016/J.RSE.2010.10.017, 2012.
- 670 Fofonoff, N. P. and Millard Jr, R. C.: Algorithms for the computation of fundamental properties of seawater., UNESCO. [online] Available from: <https://www.oceanbestpractices.net/handle/11329/109> (Accessed 30 August 2019), 1983.
- Fong, D. A. and Geyer, W. R.: The Alongshore Transport of Freshwater in a Surface-Trapped River Plume\*, *J. Phys. Oceanogr.*, 32(3), 957–972, doi:10.1175/1520-0485(2002)032<0957:TATOFI>2.0.CO;2, 2002.
- 675 Große, F., Fennel, K., Zhang, H. and Laurent, A.: Quantifying the contributions of riverine vs. oceanic nitrogen to hypoxia in the East China Sea, *Biogeosciences Discuss.*, doi:10.5194/bg-2019-342, 2019.


- Guo, Y., Rong, Z., Li, B., Xu, Z., Li, P. and Li, X.: Physical processes causing the formation of hypoxia off the Changjiang estuary after Typhoon Chan-hom, 2015, *J. Oceanol. Limnol.*, 37(1), 1–17, doi:10.1007/s00343-019-7336-5, 2019.
- 680 Horner-Devine, A. R.: The bulge circulation in the Columbia River plume, *Cont. Shelf Res.*, 29(1), 234–251, doi:10.1016/j.csr.2007.12.012, 2009.
- Hu, J. and Wang, X. H.: Progress on upwelling studies in the China seas, *Rev. Geophys.*, 54(3), 653–673, doi:10.1002/2015RG000505, 2016.
- 685 Hung, C.-C., Chung, C.-C., Gong, G.-C., Jan, S., Tsai, Y., Chen, K.-S., Chou, W. C., Lee, M.-A., Chang, Y., Chen, M.-H., Yang, W.-R., Tseng, C.-J. and Gawarkiewicz, G.: Nutrient supply in the Southern East China Sea after Typhoon Morakot, *J. Mar. Res.*, 71(1), 133–149, doi:10.1357/002224013807343425, 2013.
- Kang, Y., Pan, D., Bai, Y., He, X., Chen, X., Chen, C.-T. A. and Wang, D.: Areas of the global major river plumes, *Acta Oceanol. Sin.*, 32(1), 79–88, doi:10.1007/s13131-013-0269-5, 2013.
- 690 Large, W. G. and Pond, S.: Open Ocean Momentum Flux Measurements in Moderate to Strong Winds, *J. Phys. Oceanogr.*, 11(3), 324–336, doi:10.1175/1520-0485(1981)011<0324:OOMFMI>2.0.CO;2, 1981.
- Lentz, S. J. and Helfrich, K. R.: Buoyant gravity currents along a sloping bottom in a rotating fluid, *J. Fluid Mech.*, 464, 251–278, doi:10.1017/S0022112002008868, 2002.
- Li, D., Jing, Z., Daji, H., Ying, W. and Jun, L.: Oxygen depletion off the Changjiang (Yangtze River) Estuary, *Sci. CHINA (Series D)*, 45(12), 2002.
- 695 Li, L., He, Z., Xia, Y. and Dou, X.: Dynamics of sediment transport and stratification in Changjiang River Estuary, China, *Estuar. Coast. Shelf Sci.*, 213, 1–17, doi:10.1016/J.ECSS.2018.08.002, 2018a.
- Li, M. and Chen, Z.: An Assessment of Saltwater Intrusion in the Changjiang (Yangtze) River Estuary, China, *Coasts and Estuaries*, 31–43, doi:10.1016/B978-0-12-814003-1.00002-2, 2019.
- 700 Li, M. and Rong, Z.: Effects of tides on freshwater and volume transports in the Changjiang River plume, *J. Geophys. Res.*, 117, 6027, doi:10.1029/2011JC007716, 2012.
- Li, Y. L.: Seasonal Hypoxia and its Affecting Factors in the Yangtze River Estuary, Ocean University of China, Qingdao (in Chinese), 2015.
- 705 Li, Z., Song, S., Li, C. and Yu, Z.: The sinking of the phytoplankton community and its contribution to seasonal hypoxia in the Changjiang (Yangtze River) estuary and its adjacent waters, *Estuar. Coast. Shelf Sci.*, 208, 170–179, doi:10.1016/J.ECSS.2018.05.007, 2018b.
- Liblik, T., Karstensen, J., Testor, P., Alenius, P., Hayes, D., Ruiz, S., Heywood, K. J., Pouliquen, S., Mortier, L. and Mauri, E.: Potential for an underwater glider component as part of the Global Ocean Observing System, *Methods Oceanogr.*, 17, 50–82, doi:10.1016/j.mio.2016.05.001, 2016.
- 710 Lie, H.-J. and Cho, C.-H.: Recent advances in understanding the circulation and hydrography of the East China Sea, *Fish. Oceanogr.*, 11(6), 318–328, doi:10.1046/j.1365-2419.2002.00215.x, 2002.
- Lie, H.-J. and Cho, C.-H.: Seasonal circulation patterns of the Yellow and East China Seas derived from satellite-tracked drifter trajectories and hydrographic observations, *Prog. Oceanogr.*, 146, 121–141, doi:10.1016/J.POCEAN.2016.06.004, 2016.
- 715 Lips, U., Kikas, V., Liblik, T. and Lips, I.: Multi-sensor in situ observations to resolve the sub-mesoscale features in the stratified Gulf of Finland, Baltic Sea, *Ocean Sci.*, 12(3), 715–732, doi:10.5194/os-12-715-2016, 2016.
- Liu, W., Song, J., Huamao, Y., Li, N., Xuegang, L. I. and Liqin, D.: Dissolved barium as a tracer of Kuroshio incursion in the Kuroshio region east of Taiwan Island and the adjacent East China Sea, *Sci. China Earth Sci.*, 60(7), 1356–1367, doi:10.1007/s11430-016-9039-7, 2017.
- 720 Liu, Z. and Gan, J.: Modeling Study of Variable Upwelling Circulation in the East China Sea: Response to a Coastal Promontory, *J. Phys. Oceanogr.*, 44(4), 1078–1094, doi:10.1175/JPO-D-13-0170.1, 2014.
- Meissner, T., Wentz, F. J. and Manaster, A.: Remote Sensing Systems SMAP Ocean Surface Salinities [Level 2C, Level 3 Running 8-day, Level 3 Monthly], Version 3.0 validated release. Remote Sensing Systems, Santa

- Rosa, CA, USA., , doi:10.5067/SSSSS-TTTTT, 2018.
- 725 Moon, J.-H., Kim, T., Son, Y. B., Hong, J.-S., Lee, J.-H., Chang, P.-H. and Kim, S.-K.: Contribution of low-salinity water to sea surface warming of the East China Sea in the summer of 2016, *Prog. Oceanogr.*, 175, 68–80, doi:10.1016/j.pocean.2019.03.012, 2019.
- Ni, X., Huang, D., Zeng, D., Zhang, T., Li, H. and Chen, J.: The impact of wind mixing on the variation of bottom dissolved oxygen off the Changjiang Estuary during summer, *J. Mar. Syst.*, 154, 122–130, doi:10.1016/j.jmarsys.2014.11.010, 2016.
- 730 Ning, X., Lin, C., Su, J., Liu, C., Hao, Q. and Le, F.: Long-term changes of dissolved oxygen, hypoxia, and the responses of the ecosystems in the East China Sea from 1975 to 1995, *J. Oceanogr.*, 67(1), 59–75, doi:10.1007/s10872-011-0006-7, 2011.
- Obenour, D. R., Scavia, D., Rabalais, N. N., Turner, R. E. and Michalak, A. M.: Retrospective Analysis of Midsummer Hypoxic Area and Volume in the Northern Gulf of Mexico, 1985–2011, *Environ. Sci. Technol.*, 47(17), 9808–9815, doi:10.1021/es400983g, 2013.
- Peng, L., Benwei, S., Yaping, W., Weihua, Q., Yangang, L. and Jian, C.: Analysis of the characteristics of offshore currents in the Changjiang (Yangtze River) estuarine waters based on buoy observations, *Acta Ocean. Sin.*, 36(4), 13–20, doi:10.1007/s13131, 2017.
- 740 Qian, W., Dai, M., Xu, M., Kao, S. J., Du, C., Liu, J., Wang, H., Guo, L. and Wang, L.: Non-local drivers of the summer hypoxia in the East China Sea off the Changjiang Estuary, *Estuar. Coast. Shelf Sci.*, 198, 393–399, doi:10.1016/j.ecss.2016.08.032, 2017.
- Ren, F., Fan, D., Wu, Y. and Zhao, Q.: The evolution of hypoxia off the Changjiang Estuary in the last 3000 years: Evidence from benthic foraminifera and elemental geochemistry, *Mar. Geol.*, 417, 106039, doi:10.1016/J.MARGEO.2019.106039, 2019.
- 745 Rudnick, D. L.: Ocean Research Enabled by Underwater Gliders, *Ann. Rev. Mar. Sci.*, 8(1), 519–541, doi:10.1146/annurev-marine-122414-033913, 2016.
- Sun, H., Yang, Q., Zhao, W., Liang, X. and Tian, J.: Temporal variability of diapycnal mixing in the northern South China Sea, *J. Geophys. Res. Ocean.*, 121(12), 8840–8848, doi:10.1002/2016JC012044, 2016.
- 750 Testa, J. M. and Kemp, W. M.: Spatial and Temporal Patterns of Winter–Spring Oxygen Depletion in Chesapeake Bay Bottom Water, *Estuaries and Coasts*, 37(6), 1432–1448, doi:10.1007/s12237-014-9775-8, 2014.
- Tseng, Y.-F., Lin, J., Dai, M. and Kao, S.-J.: Joint effect of freshwater plume and coastal upwelling on phytoplankton growth off the Changjiang River, *Biogeosciences*, 11(2), 409–423, doi:10.5194/bg-11-409-2014, 2014.
- 755 Vaquer-Sunyer, R. and Duarte, C. M.: Thresholds of hypoxia for marine biodiversity., *Proc. Natl. Acad. Sci. U. S. A.*, 105(40), 15452–7, doi:10.1073/pnas.0803833105, 2008.
- Venkatesan, R., Lix, J. K., Phanindra Reddy, A., Arul Muthiah, M. and Atmanand, M. A.: Two decades of operating the Indian moored buoy network: significance and impact, *J. Oper. Oceanogr.*, 9(1), 45–54, doi:10.1080/1755876X.2016.1182792, 2016.
- 760 Wang, B. and Wang, X.: Chemical hydrography of coastal upwelling in the East China Sea, *Chinese J. Oceanol. Limnol.*, 25(1), 16–26, doi:10.1007/s00343-007-0016-x, 2007.
- Wang, B., Wei, Q., Chen, J. and Xie, L.: Annual cycle of hypoxia off the Changjiang (Yangtze River) Estuary, *Mar. Environ. Res.*, 77, 1–5, doi:10.1016/j.marenvres.2011.12.007, 2012.
- 765 Wang, B., Chen, J., Jin, H., Li, H., Huang, D. and Cai, W.-J.: Diatom bloom-derived bottom water hypoxia off the Changjiang estuary, with and without typhoon influence, *Limnol. Oceanogr.*, 62(4), 1552–1569, doi:10.1002/lno.10517, 2017.
- Wang, H., Dai, M., Liu, J., Kao, S.-J., Zhang, C., Cai, W.-J., Wang, G., Qian, W., Zhao, M. and Sun, Z.: Eutrophication-Driven Hypoxia in the East China Sea off the Changjiang Estuary, *Environ. Sci. Technol.*, 50(5), 2255–2263, doi:10.1021/acs.est.5b06211, 2016.
- 770 Wei, Q., Wang, B., Yu, Z., Chen, J. and Xue, L.: Mechanisms leading to the frequent occurrences of hypoxia and a preliminary analysis of the associated acidification off the Changjiang estuary in summer, *Sci. China Earth*


- Sci., 60(2), 360–381, doi:10.1007/s11430-015-5542-8, 2017a.
- 775 Wei, Q., Yu, Z., Wang, B., Wu, H., Sun, J., Zhang, X., Fu, M., Xia, C. and Wang, H.: Offshore detachment of the Changjiang River plume and its ecological impacts in summer, *J. Oceanogr.*, 73(3), 277–294, doi:10.1007/s10872-016-0402-0, 2017b.
- Weiss, R. F.: The solubility of nitrogen, oxygen and argon in water and seawater, *Deep Sea Res. Oceanogr. Abstr.*, 17(4), 721–735, doi:10.1016/0011-7471(70)90037-9, 1970.
- 780 Wu, H., Deng, B., Yuan, R., Hu, J., Gu, J., Shen, F., Zhu, J., Zhang, J., Wu, H., Deng, B., Yuan, R., Hu, J., Gu, J., Shen, F., Zhu, J. and Zhang, J.: Detiding Measurement on Transport of the Changjiang-Derived Buoyant Coastal Current, *J. Phys. Oceanogr.*, 43(11), 2388–2399, doi:10.1175/JPO-D-12-0158.1, 2013.
- Wu, Y., Fan, D., Wang, D. and Yin, P.: Increasing hypoxia in the Changjiang Estuary during the last three decades deciphered from sedimentary redox-sensitive elements, *Mar. Geol.*, 106044, doi:10.1016/J.MARGEO.2019.106044, 2019.
- 785 Xu, Q., Zhang, S., Cheng, Y. and Zuo, J.: Interannual Feature of Summer Upwelling around the Zhoushan Islands in the East China Sea, *J. Coast. Res.*, 331, 125–134, doi:10.2112/JCOASTRES-D-15-00197.1, 2017.
- Xu, Z., Ma, J., Wang, H., Hu, Y., Yang, G., Deng, W., Xu, Z., Ma, J., Wang, H., Hu, Y., Yang, G. and Deng, W.: River Discharge and Saltwater Intrusion Level Study of Yangtze River Estuary, China, *Water*, 10(6), 683, doi:10.3390/w10060683, 2018.
- 790 Xuan, J.-L., Huang, D., Zhou, F., Zhu, X.-H. and Fan, X.: The role of wind on the detachment of low salinity water in the Changjiang Estuary in summer, *J. Geophys. Res. Ocean.*, 117(C10), n/a-n/a, doi:10.1029/2012JC008121, 2012.
- Yang, L., Chen, Z., Sun, Z. and Hu, J.: Sectional characteristics of temperature, salinity and density off the central Zhejiang coast in the spring of 2016, *Acta Oceanol. Sin.*, 38(4), 175–182, doi:10.1007/s13131-019-1421-7, 2019.
- 795 Ye, W., Zhang, G., Zhu, Z., Han, Y., Wang, L. and Sun, M.: Methane distribution and sea-to-air flux in the East China Sea during the summer of 2013: Impact of hypoxia, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 124, 74–83, doi:10.1016/J.DSR2.2015.01.008, 2016.
- 800 Zhang, J., Liu, S. M., Ren, J. L., Wu, Y. and Zhang, G. L.: Nutrient gradients from the eutrophic Changjiang (Yangtze River) Estuary to the oligotrophic Kuroshio waters and re-evaluation of budgets for the East China Sea Shelf, *Prog. Oceanogr.*, 74(4), 449–478, doi:10.1016/j.pocean.2007.04.019, 2007a.
- Zhang, L., Liu, Z., Zhang, J., Hong, G. H., Park, Y. and Zhang, H. F.: Reevaluation of mixing among multiple water masses in the shelf: An example from the East China Sea, *Cont. Shelf Res.*, 27(15), 1969–1979, doi:10.1016/J.CSR.2007.04.002, 2007b.
- 805 Zhang, W., Wu, H. and Zhu, Z.: Transient Hypoxia Extent Off Changjiang River Estuary due to Mobile Changjiang River Plume, *J. Geophys. Res. Ocean.*, 123(12), 9196–9211, doi:10.1029/2018JC014596, 2018.
- Zhang, W., Wu, H., Hetland, R. D. and Zhu, Z.: On Mechanisms Controlling the Seasonal Hypoxia Hot Spots off the Changjiang River Estuary, *J. Geophys. Res. Ocean.*, 2019JC015322, doi:10.1029/2019JC015322, 2019.
- 810 Zhou, Z., Yu, R., Sun, C., Feng, M. and Zhou, M.: Impacts of Changjiang River Discharge and Kuroshio Intrusion on the Diatom and Dinoflagellate Blooms in the East China Sea, *J. Geophys. Res. Ocean.*, 124(7), 2019JC015158, doi:10.1029/2019JC015158, 2019.
- Zhu, J., Zhu, Z., Lin, J., Wu, H. and Zhang, J.: Distribution of hypoxia and pycnocline off the Changjiang Estuary, China, *J. Mar. Syst.*, 154, 28–40, doi:10.1016/j.jmarsys.2015.05.002, 2016.
- 815 Zhu, Z.-Y., Zhang, J., Wu, Y., Zhang, Y.-Y., Lin, J. and Liu, S.-M.: Hypoxia off the Changjiang (Yangtze River) Estuary: Oxygen depletion and organic matter decomposition, *Mar. Chem.*, 125(1–4), 108–116, doi:10.1016/J.MARCHEM.2011.03.005, 2011.
- Zhu, Z. Y., Wu, H., Liu, S. M., Wu, Y., Huang, D. J., Zhang, J. and Zhang, G. Sen: Hypoxia off the Changjiang (Yangtze River) estuary and in the adjacent East China Sea: Quantitative approaches to estimating the tidal impact and nutrient regeneration, *Mar. Pollut. Bull.*, 125(1–2), 103–114, doi:10.1016/j.marpolbul.2017.07.029, 2017.




825 **Figure 1.** Map of the study area off the Changjiang Estuary. Crosses represent CTD cast stations conducted on 27 August–5 September in 2015 and circles show CTD cast stations on 24–29 July in 2017. Lines represent northern sections (N15 in 2015 and N17 in 2017), central sections (C15 in 2015 and C17 in 2017), and southern sections (S15 in 2015 and S17 in 2017). Larger circle represents the mooring M1 location. The larger cross shows the location, where wind data were gathered. The color scale shows depth (m) of the study area. The inlay shows the study area in the East China Sea.



830 **Figure 2.** Time series of in-situ and remotely sensed 8-day running mean salinity at the M1 location in 2017.




**Figure 3.** Maps of surface salinity (a, b), surface temperature (c, d), near-bottom salinity (e, f) and near-bottom temperature (g, h) from the surveys in 2015 (left panel) and 2017 (right panel). 30 and 24.5 °C isolines are shown as white lines.




**Fig. 4.** Vertical distributions of temperature, salinity, density, DO concentration, and Apparent Oxygen Utilization (AOU) along sections N15, C15, and S15 in 2015 (Fig. 1). Temperature isoline 24.5 °C is shown with the thicker dashed line in the temperature, oxygen and AOU plots. AOU isoline 2 mg l<sup>-1</sup> is shown with the thicker solid line in AOU plots. Hypoxia border (3.0 mg l<sup>-1</sup>) is shown with a thicker solid line in oxygen plots. Thin lines represent isolines of temperature with 2 °C step, salinity with 2 step, oxygen and AOU with 2 mg l<sup>-1</sup> step both. Locations of CTD stations are

shown as crosses in the top panels. Values on the x-axis indicate the distance from the westernmost point of a section (Fig. 1).




**Fig. 5.** Vertical distributions of temperature, salinity, density, DO concentration and AOU along sections N17, C17, S17 in 2017 (Fig. 1). Temperature isoline 24.5 °C is shown with the dashed line in the temperature, oxygen and AOU plots. AOU isoline 2 mg l<sup>-1</sup> is shown with a solid line in AOU plots. Hypoxia border (3.0 mg l<sup>-1</sup>) is shown with a solid line in oxygen plots. Locations of CTD stations are shown as crosses in the top panels. Thin lines represent isolines of temperature with 2 °C step, salinity with 4 step, oxygen and AOU with 2 mg l<sup>-1</sup> step both. Values on the x-axis indicate the distance from the westernmost point of a section (Fig. 1).





860 **Fig 6.** Maps of density difference between the near-bottom layer and surface layer (a,b), in-situ Chl *a* (c), satellite-derived Chl *a* (d), near-bottom DO (e,f) and total AOU in the water column (g,h). Maps of 2015 are shown on left panels and 2017 on the right. Remotely sensed Chl *a* map (f) is a mean field of two daily images of 21–22 July. AOU was calculated according to Eq. (3).



865 **Fig. 7.** Maps of AOU isoline 2 mg l<sup>-1</sup> (upper boundary of the DO depletion, UBD) depth (upper panels) and temperature isoline 24.5 °C (upper boundary of KSSW, UBK) depth (lower panels) in 2015 (left panels) and in 2017 (right panels).







Figure 8. Maps of mean wind (panels a–b,  $\text{m s}^{-1}$ ), simulated surface current (panels c–d,  $\text{cm s}^{-1}$ ) and bottom current (e–f,  $\text{cm s}^{-1}$ ) s currents in the sea bottom depths down to 100 m during seven days period before the CTD surveys. Bottom bathymetry is shown as a background in panels a–f. Red lines in panels c and d show the section S1, where current time series (presented in Fig. 10c) were calculated. Every fourth current vector is shown only in the panels c–f. Mean sea level anomaly (m) as contours and geostrophic velocities ( $\text{cm s}^{-1}$ ) as vectors derived from satellite altimetry during the surveys are shown in the panels g–h.



875 **Figure 9.** Relation of daily mean SE–NW wind velocity component  $w_c$  (positive towards NW) and meridional current velocity component  $v_m$  (positive northward) in the section at section S1 (see Fig. 8) in 1993–2018. The color scale shows the co-occurrence (number of cases) of respective wind velocity and current velocity component combinations.




880 **Figure 10.** Time series of wind, current, and river discharge. Mean wind speed (6-hours) (a), 1-day running mean SE-NW wind stress  $\tau$  (positive towards NW;  $\text{N m}^{-2}$ ) (b), daily mean meridional current velocity component  $v_m$  (positive northward) at section S1 (c), and daily Changjiang river discharge at Datong station shifted by seven days to represent flow in the river mouth (d). Blue and red dots represent the start and end of the CTD surveys, respectively in 2015 and 2017.



885

Figure 11. Maps of monthly occurrence (%) of (remotely sensed) salinity < 30 from June to September in 2015–2018.



890 **Figure 12. Maps of surface (upper panels) and near-bottom (lower panels) currents in the sea bottom depths down to 100 m in the cases of wind stress component  $\tau_c \geq 0.02 \text{ N m}^{-2}$  (left panels) and  $\tau_c < 0 \text{ N m}^{-2}$  (right panels) in 1993–2018. Every fourth current vector is shown.**




Figure 13. Annual cycle (a, c) and inter-annual variability (b, d) of wind stress component (1992–2018)  $\tau_c \geq 0.02 \text{ N m}^{-2}$  (a, b) and river discharge (c, d) (2001–2018). Blue line shows the overall mean in (a) and (c), and June–September long-term mean in (c) and (d). Red line shows the July–August long-term mean in (b) and (d).

895

Table 1. Summary of general features observed in the study area in summers 2015 and 2017



<b>Survey on 27 August–5 September 2015</b>	<b>Survey on 24–29 July 2017</b>
Northerly winds prevailing before the survey	Southerly winds prevailing before the survey
Southward transport of the CDW	East/northeastward transport of the CDW
Stronger stratification in the south	Stronger stratification in the north
Downwelling along the coast	Upwelling along the coast
Higher Chl <i>a</i> in south	Higher Chl <i>a</i> in north
No low DO concentration zone in the north	Strong and shallow hypoxic zone in the north
Bottom water warmer and fresher in the north	Bottom water colder and saltier in the north
Lower DO concentrations further offshore at sea depths $\geq 30$ m in the south	Lower DO concentrations in the coastal upwelling zone in the south