



## Distribution of chlorine and fluorine in benthic foraminifera

Anne Roepert<sup>1</sup>, Lubos Polerecky<sup>1</sup>, Esmee Geerken<sup>2</sup>, Gert-Jan Reichart<sup>1,2</sup>, and Jack J. Middelburg<sup>1</sup>

<sup>1</sup>Department of Earth Sciences, Utrecht University, PO Box 80021, 3508 TA Utrecht, The Netherlands

<sup>2</sup>Department of Ocean Systems, NIOZ Royal Netherlands Institute for Sea Research, and Utrecht University, 1790 AB Den Burg, The Netherlands

**Correspondence:** Anne Roepert (a.roepert@uu.nl)

**Abstract.** Over the last decades a suite of inorganic proxies based on foraminiferal calcite have been developed, of which some are now widely used for paleoenvironmental reconstructions. Studies of foraminiferal shell chemistry have largely focused on cations and oxyanions, while much less is known about the incorporation of anions. The halogens fluoride and chloride are conservative in the ocean, which makes them candidates for reconstructing paleoceanographic parameters. However, their potential as a paleoproxy has hardly been explored, and fundamental insight in their incorporation is required. Here we used nano-scale secondary ion mass spectrometry (NanoSIMS) to investigate, for the first time, the distribution of Cl and F within shell walls of four benthic species of foraminifera. In the rotaliid species *Ammonia tepida* and *Amphistegina lessonii* Cl and F were highly heterogeneous and correlated within the shell walls, forming bands that were co-located with the banded distribution of phosphorus. In the miliolid species *Sorites marginalis* and *Archaias angulatus* the distribution of Cl and F was much more homogeneous without discernible bands. In these species Cl and P were correlated, whereas no correlation was observed between Cl and F or between F and P. Additionally, their F content was about an order of magnitude higher than in the rotaliid species. The high variance in the Cl and F content in the studied foraminifera could not be attributed to environmental parameters. Based on these findings we suggest that in the rotaliid species Cl and F are predominately associated with organic linings. We further propose that in the miliolid species Cl may be incorporated as a solid solution of chlorapatite or associated with organic molecules in the calcite. The high F content together with the lack of correlation between Cl and F or P in the miliolid foraminifera suggests a fundamentally different incorporation mechanism. Overall, our data clearly show that the calcification pathway employed by the studied foraminifera governs the incorporation and distribution of Cl, F, P and other elements in their calcite shells.

### 1 Introduction

Foraminifera are widely used to reconstruct paleoenvironments and climates based on relative species abundances and the chemical or isotopic composition of their shells. Apart from the carbon and oxygen isotopic composition, most inorganic proxies based on foraminiferal calcite involve cations or their isotopes (Boyle, 1981; Elderfield et al., 1996; Lea et al., 1999; Allen et al., 2016), which can substitute for calcium in the calcite lattice. While there are also proxies based on the incorporation of oxyanions such as iodate, sulphate and borate, and their isotopes (Lu et al., 2010; Paris et al., 2014; Yu et al., 2007; Rae



25 et al., 2011), much less is known about the incorporation of anions such as the halogens Cl, F and Br that are less likely to substitute for the carbonate ion (Kitano et al., 1975; Okumura et al., 1983, 1986).

The halogen anions are conservative in the ocean, which makes them potential candidates for reconstructing paleoceanographic parameters. Chlorine as a major constituent of sea salt is the most abundant halogen in seawater, followed by bromine, while fluorine concentrations are in the ppm range (e.g., Kendrick, 2018, and references therein). However, even though Cl<sup>-</sup> is  
30 the most abundant (an)ion in seawater, the Cl content of inorganic and biogenic calcites is low as the Cl<sup>-</sup> anion does not fit well into the calcite lattice (Tokuyama et al., 1972; Kitano et al., 1975). In contrast, the F<sup>-</sup> anion has a much higher compatibility in many minerals compared to the heavier halogens, which can be attributed to its much smaller ionic radius (Kendrick, 2018). While seawater is enriched with the halogens chlorine and bromine with respect to the primary mantle (Kendrick, 2018), fluorine is strongly depleted in seawater with calcium carbonates being the major sink of dissolved fluoride in the oceans  
35 (Carpenter, 1969).

Marine carbonates have F/Ca ratios ranging between 0.1–3.5 mmol mol<sup>-1</sup> and 4.0–6.5 mmol mol<sup>-1</sup> in calcite and aragonite, respectively (Carpenter, 1969), while Cl/Ca ratios range between 9–18 μmol mol<sup>-1</sup> and 53–71 μmol mol<sup>-1</sup> in inorganically precipitated calcite and aragonite, respectively (Kitano et al., 1975). The fluorine content of planktic foraminifera co-varies not only with δ<sup>18</sup>O (Rosenthal and Boyle, 1993; Opdyke et al., 1993; Rosenthal et al., 1997), but also with Mg and Sr content  
40 (Opdyke et al., 1993), and varies mostly in response to species-specific biological factors (Rosenthal and Boyle, 1993; Opdyke et al., 1993; Rosenthal et al., 1997). Furthermore, foraminiferal F content appears to be highly susceptible to diagenesis. Indeed, the release of F to porewater has been shown to be a proxy for carbonate dissolution during early diagenesis (Rude and Aller, 1991). The chlorine content of biogenic calcite has been proposed as a potential direct salinity proxy (Wit et al., 2013), but there are very few data on Cl in foraminifera (Szafrank and Erez, 1993; Erez, 2003).

45 Not much is known about how these anions are incorporated into foraminiferal shells. Possible incorporation modes include lattice-bound, interstitial, solid solution (e.g. fluorite, apatite), or bound to organic templates within the foraminiferal shell walls. Possible environmental factors playing a role during the incorporation processes of Cl and F may be related to pH homeostasis in the calcification fluid as Cl<sup>-</sup> and F<sup>-</sup> are the dissociated compounds of a strong and weak acid, respectively. Tanaka et al. (2013) showed that fluorine incorporation in non-symbiotic corals was governed by the carbonate ion concentration in  
50 solution because of ion-exchange between fluoride and carbonate. A similar link has been suggested between foraminiferal fluorine content and carbonate ion concentration (Rosenthal and Boyle, 1993; Opdyke et al., 1993) as well as the presence of symbionts (Opdyke et al., 1993).

Furthermore, the spatial distribution of incorporated elements within foraminiferal shells can shed light on the incorporation mechanism. To the best of our knowledge, the spatial distribution of F has not been investigated yet, while Cl was previously  
55 found at the location of organic linings in shell walls of *A. lobifera* (Szafrank and Erez, 1993; Erez, 2003). Recently, also the iodine micro-distribution in shell walls of *U. striata* was found to spatially correlate with organics (Glock et al., 2019). This suggests that halogen incorporation may be regulated by foraminiferal biomineralization pathways, although biogenic carbonate formation is ultimately expected to be governed by the same controlling factors as inorganic carbonate formation. The biomineralization pathway has consequences for trace element incorporation, as reflected in the occurrence of bands of



60 higher and lower concentrations of cations (e.g. Mg, Sr, Ba, Na) in perforate hyaline foraminiferal shell walls (Eggins et al.,  
2004; Sadekov et al., 2005; Kunioka et al., 2006; Branson et al., 2015; Jonkers et al., 2016; Geerken et al., 2018, 2019).  
This banding has been attributed to chamber addition in perforate hyaline foraminifera: an additional high-concentration band  
appears each time a new calcite lamella is added to the shell, which also covers previous chambers (Nehrke et al., 2013).  
Similarly, spine and the spine attachment zone of planktic foraminifera are enriched in Na while depleted in Mg, and vice  
65 versa, respectively (Branson et al., 2016; Mezger et al., 2019; Bonnin et al., 2019).

Here we investigated, for the first time, the Cl and F distribution within foraminiferal shell walls of four benthic species  
on a sub-micron scale with NanoSIMS. Two rotaliid benthic species (*Ammonia tepida* and *Amphistegina lessonii*) and two  
miliolid benthic species (*Sorites marginalis* and *Archaias angulatus*) were investigated to cover benthic foraminifera with  
fundamentally different biomineralization pathways. Furthermore, one species is symbiont-barren (*Ammonia tepida*), whereas  
70 the other three species are bearing photosynthetic symbionts.

## 2 Materials and Methods

### 2.1 Specimen selection and basic characterization

Six specimens of four species (two rotaliid and two miliolid species, respectively) from culturing experiments were investigated  
in this study (Table 1). The use of laboratory-grown specimens has the advantage that their growth conditions were controlled  
75 and that potential effects of diagenesis on their shell halogen chemistry can be excluded.

Parameters of the carbonate system in the culture media, including alkalinity, carbonate ion concentration and saturation  
state with respect to calcite, were calculated from the measured DIC concentration, pH, salinity and temperature (Table 1).  
This was done in R (R Core Team, 2018) using the package *seacarb* (Gattuso et al., 2018). The equilibrium constants for the  
carbonate system  $K_1$  and  $K_2$  were taken from Lueker et al. (2000). Because DIC was not measured in the culture of *A. lessonii*,  
80 we first applied the salinity-alkalinity relationship for the North Atlantic (Lee et al., 2006) to estimate the culture medium  
alkalinity, based on which we then calculated the remaining carbonate system parameters.

Prior to sample preparation for NanoSIMS analyses, each specimen was characterized with respect to its Na, Mg, Sr and Ba  
content. This was done by LA-ICP-MS as previously described (Geerken et al., 2018; van Dijk et al., 2019). For each specimen  
El/Ca ratios were determined on two or more chambers and averaged (Table A1).

### 85 2.2 Sample preparation and SEM imaging

The selected specimens of *A. tepida*, *A. lessonii*, *S. marginalis* and *A. angulatus* were embedded in epoxy resin (Araldite 2020)  
under vacuum and subsequently polished to expose cross-sections perpendicular to the shell walls (see Geerken et al. (2019)  
for details). Although epoxy resins contain high amounts of chlorine, we are confident that the halogens measured in the  
foraminiferal calcite are not due to contamination with resin nor an artifact of redeposition during the sputtering process  
90 (Roepert, 2019). The first polishing steps used wet grinding papers of decreasing coarseness (HERMES, WS Flex 18C,



230 mm, P 800 and 219 ATM, SiC wet grinding paper, grain 4000) followed by agglomerated alpha alumina powder (Struers AP-A powder, grain size 0.3  $\mu\text{m}$ ) and SiO<sub>2</sub> powder (Logitech SF1 Polishing Suspension, grain size 0.035  $\mu\text{m}$ ). The polished samples were sputter-coated with 20 nm of Au using a sputter coater (JEOL JFC-2300HR high resolution fine coater with JEOL FC-TM20 thickness controller), after which they were imaged using a JEOL Neoscope II JCM-6000 table-top SEM to  
95 identify suitable areas for NanoSIMS analysis (Figure A1).

### 2.3 NanoSIMS imaging

NanoSIMS analysis was performed with the Cameca NanoSIMS 50L instrument available at Utrecht University. Using an element standard (SPI Supplies, 02757-AB 59 Metals & Minerals Standard), magnetic field and exact positions of the electron multiplier detectors were adjusted to enable detection of negative secondary ions <sup>12</sup>C<sup>-</sup>, <sup>16</sup>O<sup>-</sup>, <sup>19</sup>F<sup>-</sup>, <sup>31</sup>P<sup>-</sup>, <sup>35</sup>Cl<sup>-</sup>, <sup>37</sup>Cl<sup>-</sup>, and  
100 <sup>40</sup>Ca<sup>16</sup>O<sup>-</sup>. The secondary ions were sputtered from the sample surface using an 8 kV primary Cs<sup>+</sup> ion source.

Because the primary ion beam was positive, calcium had to be detected as <sup>40</sup>Ca<sup>16</sup>O<sup>-</sup> and not as <sup>40</sup>Ca<sup>+</sup>. However, because the Ca:O stoichiometry in calcite with trace amounts of organics is fixed, we assumed that the measured distribution of <sup>40</sup>Ca<sup>16</sup>O well approximated the distribution of Ca. This assumption was supported by the good correlation between the secondary ions <sup>40</sup>Ca<sup>16</sup>O<sup>-</sup> and <sup>16</sup>O<sup>-</sup> detected from the calcite (Figure A2). <sup>31</sup>P was measured as a tracer for organics in the calcite (Geerken  
105 et al., 2019), whereas <sup>12</sup>C was measured to help distinguish between resin and calcite. To ensure that the detection of <sup>35</sup>Cl was not influenced by possible isobaric interferences such as <sup>16</sup>O<sup>18</sup>O<sup>1</sup>H and <sup>34</sup>S<sup>1</sup>H, we used sufficiently high mass resolution power (MRP > 5113) and additionally measured the isotope <sup>37</sup>Cl as well. The obtained <sup>37</sup>Cl/<sup>35</sup>Cl ratio differed from the natural abundance ratio of 0.320 by no more than 0.015, confirming that the influence of isobaric interferences for the detection of Cl was negligible. Similarly, separation of <sup>19</sup>F from the possible interference by <sup>18</sup>O<sup>1</sup>H was achieved by using MRP > 2214,  
110 whereas interferences from molecules such as <sup>12</sup>C<sup>7</sup>Li, <sup>13</sup>C<sup>6</sup>Li or <sup>16</sup>O<sup>1</sup>H<sub>3</sub> are highly unlikely.

Before each analysis the area of interest was pre-sputtered for 10–15 min until secondary ion counts stabilized. Subsequently, ion count images were acquired by rastering the primary beam (dwell time of 1 ms pixel<sup>-1</sup>) over the sample surface using the diaphragm and slit settings listed in Table A2. The primary beam current on the sample surface ranged between 0.5–2 pA depending on the size of the imaged area. The spatial resolution ranged between 50–100 nm pixel<sup>-1</sup>. All analyses employed  
115 an e-gun to avoid charging of the sample surface. Because some of the secondary ion counts were very low, the imaged areas were measured multiple times (250–1000, depending on the sample) and the signals from the individual planes were aligned and accumulated.

### 2.4 Data processing

NanoSIMS data were processed with the freeware program Look@NanoSIMS (Polerecky et al., 2012). After alignment and accumulation of the image data, regions of interest (ROIs) corresponding to foraminiferal calcite were drawn by hand based  
120 on the <sup>40</sup>Ca<sup>16</sup>O<sup>-</sup> image. With the additional use of the <sup>12</sup>C<sup>-</sup> and <sup>35</sup>Cl<sup>-</sup> images, areas of exposed resin or pores within the foraminiferal walls were identified and excluded from the final analysis.



Due to the lack of reliable calibration standards, and because Ca was measured as the molecular ion  $^{40}\text{Ca}^{16}\text{O}^-$ , the EI/Ca ratios are reported in this study by the ratios between the raw data, i.e., the secondary ion counts  $\text{EI}^-$  and  $^{40}\text{Ca}^{16}\text{O}^-$ . Although  
125 not fully quantitative, the ratios calculated in this way are comparable between the different foraminifera specimens and species because the secondary ion counts detected by NanoSIMS are linearly proportional to the concentration of the corresponding element, and because all the measurements were done in a similar biogenic calcite matrix using the same pre-sputtering and measuring protocol.

To ensure that the EI/Ca ratios were not affected by insufficient pre-sputtering, depth profiles of the  $\text{EI}^- / ^{40}\text{Ca}^{16}\text{O}^-$  ion count ratios were inspected for each ROI, and the planes where the ratios showed a significant trend with depth were excluded from  
130 the final analysis. Lateral profiles of the  $\text{EI}^- / ^{40}\text{Ca}^{16}\text{O}^-$  ion count ratios perpendicular to the shell surface were extracted from the accumulated NanoSIMS images. The width of the profile line, which corresponds to the amount of averaged lateral profiles, was set to 20 pixel to increase the signal-to-noise-ratio.

To investigate the spatial correlation of  $\text{EI}^- / ^{40}\text{Ca}^{16}\text{O}^-$  ion count ratios, ROIs were drawn on the NanoSIMS images in  
135 Look@NanoSIMS in such a way, that regions of higher and lower ion count ratios on the foraminifera were separated into different ROIs to separate the spatial variability. That way, 40 to 47 separate data points (ROIs) per species, grouped per specimen, were extracted from the NanoSIMS images. Subsequently, correlation matrices were calculated for the accumulated ion count ratios in those ROIs using the corrpilot package (Wei and Simko, 2017) in R (R Core Team, 2018).

### 3 Results

#### 140 3.1 Spatial distribution of chlorine and fluorine

The halogens Cl and F show distinct banding in the rotaliids, in particular in *A. tepida* (Figure 1). Moreover, in the rotaliids, maxima of Cl/Ca and F/Ca are co-located with those of P/Ca, and correlate well in *A. tepida*, while the correspondence between P/Ca and F/Ca is moderate in *A. lessonii*, and spatially rather complex for Cl/Ca (see lateral profiles in Figure 1). Furthermore, the contrast between the high-intensity and low-intensity bands in F/Ca, Cl/Ca and P/Ca is higher in *A. tepida* than in *A. lessonii*.  
145 In the miliolid foraminifers no banding of neither halogens nor P is visible, with the exception of a slight elevation in Cl and P in areas of an image that were identified as a suture in SEM images. Lateral profiles in *A. angulatus* show a correlation of Cl with P, and no correlation of F with Cl or P. The lateral profile through the shell wall of *S. marginalis* shows similar patterns as the one of *A. angulatus*. The F/Ca ion count ratios in the miliolids *A. angulatus* and *S. marginalis* are in the same range and one order of magnitude higher than those in rotaliid *A. tepida* and *A. lessonii* (Figures 1 and 2). The Cl/Ca and P/Ca ion count  
150 ratios are in the same order of magnitude in all four species (Figures 1 and 2).

Ion count ratios of F/Ca and Cl/Ca correlate with each other in *A. tepida*, while there is no correlation in *A. lessonii* and the miliolids (Figure 2A and A3). Both F/Ca and Cl/Ca are correlated with P/Ca in the rotaliids, while only Cl/Ca is correlated with P/Ca in the miliolids (Figure 2B,C). All correlations described here are significant to a level of  $p < 0.001$ ;  $R^2$  and p-values of the correlations are reported in Figure A3. These correlations are also seen when the elements F, Cl and P are normalized to O  
155 instead being normalized to Ca (Figure A5).



### 3.2 Relation with cation incorporation and culture media properties

In all four species, the Cl content does not correlate to any of the elemental ratios measured by LA-ICP-MS (upper panels in Figure 3). However, the elevated F/Ca ratios in the miliolid foraminifera coincide with elevated Mg/Ca and Ba/Ca, which also are an order of magnitude higher in these species than in the rotaliid foraminifera (lower panels in Figure 3). Our data show no correlation of Cl/Ca or F/Ca with neither Na/Ca nor Sr/Ca (Figure 3).

Our data show no correlation of Cl/Ca or F/Ca with salinity or temperature (Figure A4). Furthermore, crossplots of the NanoSIMS Cl/Ca and F/Ca ion count ratios show no correlation with carbonate system parameters for Cl/Ca (Figure 4). However, NanoSIMS F/Ca ion count ratios are higher in the miliolid foraminifera, which were cultured at higher DIC, corresponding to higher alkalinity and carbonate ion concentration as well (Figure 4).

## 4 Discussion

### 4.1 Limited environmental control on Cl and F incorporation into foraminiferal shells

The NanoSIMS data presented clearly show that biomineralization pathways govern the incorporation and distribution of Cl and F within foraminiferal shells: the rotaliid species show distinct banding in Cl, F and P, while the F-rich miliolid species do not. Biologic control is known to affect incorporation of most elements into foraminiferal shells, while at the same time relationships with environmental conditions have proven robust tools for paleo reconstructions (Eggins et al., 2004; Kunioka et al., 2006; Paris et al., 2014; Spero et al., 2015; Fehrenbacher et al., 2017; Geerken et al., 2019). In our data set comprised of a very limited amount of specimens, we see no overall trend in Cl/Ca ratios in foraminiferal calcite with chemical properties of the culture media. The high intra-shell variability in rotaliid foraminifera and the spatial correlation with P on the location of organic linings suggest that Cl is associated with organic linings in rotaliid foraminiferal shells. Furthermore, Cl/Ca is highly variable within a single section through a foraminiferal wall in all the species measured and the range of Cl/Ca ratios is similar in all investigated specimens.

The overall absence of trends with environmental conditions as well as the high intra-specimen variability lowers the confidence in potentially using Cl/Ca for paleo reconstructions. However, as this study does not cover a range of physicochemical parameters for a single species, but rather presents a collection of different species that were also grown in different conditions, we cannot exclude that any of the presented physicochemical parameters may exert an influence on the Cl content of foraminifera on a species-specific level. A definite conclusion regarding proxy applicability would require culturing studies including 20–30 specimens per species per environmental condition.

Moreover, there is no discernable trend of Cl/Ca ratios in foraminiferal calcite with any of the measured trace elements (Figure 3). Cl/Ca ratios are in the same range for species with low-Mg calcite (rotaliid) as they are for those with high-Mg calcite (miliolid), suggesting that chlorine incorporation is systematically different from the incorporation of these cations. In inorganically precipitated calcite, chlorine contents are an order of magnitude lower than sodium contents (Kitano et al.,





1975), suggesting that Cl is incorporated neither as fluid inclusions (Wit et al., 2013) nor as solid solutions of NaCl into calcite. As chlorine content seems not to reflect any environmental parameter, and Cl/Ca correlates well with P/Ca in all the species investigated here, we suspect that chlorine incorporation into foraminiferal calcite is closely related to organic molecules involved in calcification or to a solid-solution between calcite and chlorapatite ( $\text{Ca}_5(\text{PO}_4)_3(\text{OH},\text{Cl})$ ).

The F/Ca ratios in the miliolid species are about an order of magnitude higher than those in the rotaliid species. The elevated F/Ca ratios in miliolids coincide with overall higher  $\text{CO}_3^{2-}$ -ion concentration in the culture media of the miliolid species. This might indicate a relation between foraminiferal F/Ca ratios and carbonate ions, but the relationship is inverse to what one would expect for inorganic ion exchange (Ichikuni, 1979) and what has been observed in corals (Tanaka et al., 2013). However, the high intra-shell variability in F/Ca ratios of single specimens and the co-variation of environmental conditions with mineralization pathway complicates attributing F/Ca ratios to environmental parameters. Species-specific culturing studies could provide more insight into whether F/Ca ratios of benthic foraminifera on a species-specific level correlate with environmental conditions, as then the effect of different biomineralization pathways would not hamper interpretation as is the case in our data set.

Opdyke et al. (1993) suggested that the presence of photosynthetic symbionts in foraminifera impacts their F/Ca ratio. Symbionts influence the intracellular carbonate chemistry by photosynthesis, which could link to fluoride incorporation via the intracellular  $\text{CO}_3^{2-}$  ion activity. *A. tepida* is the only symbiont-barren species we investigated, and indeed its F/Ca ratios are lower than in the miliolid symbiont-bearing species. However, the symbiont-bearing rotaliid *A. lessonii* exhibits the lowest mean F/Ca ratios, which are in the same order of magnitude as in *A. tepida*, but less variable throughout the shell wall. We therefore conclude that F/Ca ratio is unlikely directly related to the presence of symbionts in foraminifera.

Notably, F/Ca ratios are higher in specimens with a higher Mg and Ba content (Figure 3). A correlation of trace element content with Mg content within and between species has been found for several elements, including Sr, Zn, and Na (Evans et al., 2015; van Dijk et al., 2017; Geerken et al., 2018), and also F (Opdyke et al., 1993). The miliolid species have generally a much higher Mg content than the rotaliid species and their biomineralization mechanisms are thought to be substantially different (Figure 5). The fact that higher F content corresponds with higher Mg content may point towards a strong biological control on F incorporation. Fluorine may be incorporated in solid solutions. Fluorite ( $\text{CaF}_2$ ) solid solution has been suggested as the incorporation mechanism for fluorine in calcite (Carpenter, 1969), but also fluorapatite ( $\text{Ca}_5(\text{PO}_4)_3(\text{OH},\text{F})$ ) solid solution would be a possible option.

#### 215 4.2 Cl and F incorporation in foraminifera is primarily controlled by biomineralisation pathway

In the two species of rotaliid foraminifera that were investigated here, Cl and F show strong banding. The Cl and F bands are co-located with P in the foraminiferal shell walls. Since phosphorus is present in organic molecules like phospholipids in membranes, P/Ca can be used to image organic linings in between the lamella in hyaline foraminiferal shell walls, as demonstrated in Geerken et al. (2019). In *A. tepida*, the correlation of Cl and F with P is tight, and we conclude that both elements are primarily associated with the organic linings. In *A. lessonii*, the peaks in Cl/Ca and F/Ca lateral profiles are also co-located with peaks in P/Ca. However, in the specimens we analyzed, there seems to be substantial additional Cl and F also



in some calcite lamella, as can be seen in the lateral profiles. Moreover, the contrast between high-intensity bands and low-intensity bands is less prominent in *A. lessonii* than in *A. tepida*. We suggest that also in *A. lessonii*, association with organic linings is the primary mode of incorporation of both elements in the foraminiferal shells. Using NanoSIMS and the very same  
225 species and specimens, Geerken et al. (2019) reported co-occurrence of organic linings and banding of Mg, Na, Sr, K, S, P and N. Moreover, they showed that elemental incorporation in *A. lessonii* was overall higher than in *A. tepida*, consistent with our observations for the halogens (Figures 1, 2).

In the miliolids, the distributions of Cl and F are distinctly different from those of the rotaliids: since miliolids do not calcify by adding subsequent lamella of calcite (Figure 5), no patterns of alternating high and low concentration banding are  
230 visible. Moreover, Cl and F are spatially not correlated throughout the shell walls of these miliolids, indicating that Cl and F may have different modes of incorporation. The correlation of Cl and P within the shell wall supports the hypothesis that Cl is incorporated in the calcite as a solid solution of chlorapatite, a mineral containing both Cl and P, or associated with organic molecules as for the rotaliids, but then distributed in a less organized way (no banding) within the calcite. The type of organics being present in foraminiferal calcite is determined by the precipitation pathway: rotaliids initiate calcification around  
235 a primary organic sheet (POS) and cover their shell with organic linings, while miliolid shells comprise of randomly oriented calcite needles held together in an organic matrix. As these organics may differ in their P content, it is possible to measure comparable P contents in both rotaliid and miliolid calcite, even though their absolute organic matter content within the calcite is expected to be different. Similar Cl content in both rotaliids and miliolids may thus be due to differences in the composition of the organics or may hint to an incorporation via a different pathway. Alternatively, apatite may form via the adsorption  
240 of phosphate to calcite and amorphous calcium phosphates at low Mg concentrations in solution (Martens and Harriss, 1970; Millero et al., 2001). The incorporation of Cl via chlorapatite could also explain the spatial correlation of Cl and P in the species we analysed. If F would be incorporated similarly as Cl, we would expect the F and Cl distribution to be comparable. As this is not the case, we conclude that F is incorporated primarily in a different way than Cl, e.g. as a solid solution of fluorite, as suggested before (Rosenthal and Boyle, 1993).

245 The observed difference between F-rich miliolids lacking organic linings and clear banding of trace elements on the one hand, and the rotaliids with F, Cl and P rich bands on the other hand, is consistent with known differences in calcification mechanisms (Figure 5) that have developed during the evolution of foraminifera (Debenay et al., 1996; Bentov and Erez, 2006). Hyaline (including the rotaliid) foraminifera precipitate calcite onto organic templates within an extracellular but confined space, and add a new lamella to the entire shell each time they produce a new chamber (Hemleben et al., 1986; de Nooijer  
250 et al., 2014). For some intermediate Mg-calcite producing foraminifera (like *A. lobifera* or also *A. lessonii*), transport of vesicles to the site of calcification has been observed suggesting controlled biomineralization (at least partly) from internal Ca and carbonate pools (e.g., de Nooijer et al., 2009). Furthermore, in hyaline foraminifera, selective ion transport to the site of calcification via trans-membrane pumping of elements is discussed as the biological control on biomineralization (Nehrke et al., 2013; de Nooijer et al., 2014; Toyofuku et al., 2017). In contrast, porcelaneous miliolid foraminifera produce a wall  
255 of high-Mg calcite with thin inner and outer layers connected by a thick middle layer of crystal needles (Hemleben et al., 1986; Debenay et al., 1996), which are glued together by an organic matrix. Miliolid foraminifera are generally thought to





precipitate calcium crystals in intracellular vesicles prior to arranging them in the shape of the new chamber wall (Angell, 1980; Hemleben et al., 1986; Debenay et al., 1998, 2000; Bentov and Erez, 2006). Furthermore, there are species within the suborder of the miliolids that show features of both biomineralization pathways, such as *Archaias angulatus*, which appears to precipitate calcite at the site of the new chamber wall, opposed to other miliolid species (Wetmore, 1999).

During calcification, miliolids enclose seawater vesicles (Figure 5) and then produce calcite rich in various cations (van Dijk et al. (2017), Table A1) and F (Figure 1). The rotaliids such as *A. tepida* may use highly selective ion channels and organic layers to deposit new calcite. As a consequence, these calcites are low in Mg, Ba and the halogens (F, Cl) and show distinct banding. The rotaliid *A. lessonii* produces calcite with intermediate Mg contents and less distinct banding for cations (Geerken et al., 2019), indicative of less biological control on ion transport and calcite deposition than in *A. tepida*. Our results for P, Cl and F support this.

## 5 Conclusions

Here we investigated for the first time the spatial distribution of the halogens Cl and F in foraminiferal shell walls. In the rotaliid benthic species *Ammonia tepida* and *Amphistegina lessonii*, Cl and F are distributed in bands within the chamber walls, which co-locate with P banding. In the miliolid benthic species *Sorites marginalis* and *Archaias angulatus* Cl and F were not found to occur in bands. However, the rather homogeneously-distributed Cl was found to correlate with P content, while F did not correlate with either P or Cl. Based on these findings we suggest that Cl and F are predominately associated with organic linings in the rotaliid species. We further propose that Cl may be incorporated in miliolid species as a solid solution of chlorapatite or be associated with organics. Our data in the miliolid species suggests that F is incorporated in a different way than Cl, as F does not correlate with P.

*Data availability.* The data presented in this study are available at doi:10.4121/uuid:9951e801-5574-498e-b375-fa6941a0f071.

*Author contributions.* JJM, GJR and AR designed the experiments. EG prepared the samples for NanoSIMS analyses. EG and AR conducted the SEM analyses. AR conducted the NanoSIMS analyses. AR and LP performed the data analysis. AR interpreted the data, prepared the figures and wrote the manuscript text with contributions from all authors.

*Competing interests.* The authors declare no competing interests.

*Acknowledgements.* We thank Inge van Dijk and Lennart de Nooijer for providing *A. lessonii* specimens, and Michiel Kienhuis for analytical support. The NanoSIMS facility at Utrecht University was financed through a large infrastructure grant by the Netherlands Organisation for

<https://doi.org/10.5194/bg-2019-424>  
Preprint. Discussion started: 20 November 2019  
© Author(s) 2019. CC BY 4.0 License.



Scientific Research (NWO) (grant no. 175.010.2009.011). This work was carried out under the programme of the Netherlands Earth System Science Centre (NESSC), financially supported by the Ministry of Education, Culture and Science (OCW) (grant no. 024.002.001).



## 285 References

- Allen, K. A., Hönisch, B., Eggins, S. M., Haynes, L. L., Rosenthal, Y., and Yu, J.: Trace element proxies for surface ocean conditions: A synthesis of culture calibrations with planktic foraminifera, *Geochimica et Cosmochimica Acta*, 193, 197–221, 2016.
- Angell, R. W.: Test morphogenesis (chamber formation) in the foraminifer *Spiroloculina hyalina* Schulze, *The Journal of Foraminiferal Research*, 10, 89–101, 1980.
- 290 Bentov, S. and Erez, J.: Impact of biomineralization processes on the Mg content of foraminiferal shells: A biological perspective, *Geochemistry, Geophysics, Geosystems*, 7, 2006.
- Bonnin, E. A., Zhu, Z., Fehrenbacher, J. S., Russell, A. D., Hönisch, B., Spero, H. J., and Gagnon, A. C.: Submicron Sodium Banding in Cultured Planktic Foraminifera Shells, *Geochimica et Cosmochimica Acta*, 2019.
- Boyle, E. A.: Cadmium, zinc, copper, and barium in foraminifera tests, *Earth and Planetary Science Letters*, 53, 11 – 35,
- 295 [https://doi.org/http://dx.doi.org/10.1016/0012-821X\(81\)90022-4](https://doi.org/http://dx.doi.org/10.1016/0012-821X(81)90022-4), <http://www.sciencedirect.com/science/article/pii/0012821X81900224>, 1981.
- Branson, O., Kaczmarek, K., Redfern, S. A., Misra, S., Langer, G., Tyliszczak, T., Bijma, J., and Elderfield, H.: The coordination and distribution of B in foraminiferal calcite, *Earth and Planetary Science Letters*, 416, 67–72, 2015.
- Branson, O., Bonnin, E. A., Perea, D. E., Spero, H. J., Zhu, Z., Winters, M., Hönisch, B., Russell, A. D., Fehrenbacher, J. S., and Gagnon,
- 300 A. C.: Nanometer-scale chemistry of a calcite biomineralization template: Implications for skeletal composition and nucleation, *Proceedings of the National Academy of Sciences*, 113, 12 934–12 939, 2016.
- Carpenter, R.: Factors controlling the marine geochemistry of fluorine, *Geochimica et Cosmochimica Acta*, 33, 1153–1167, [https://doi.org/https://doi.org/10.1016/0016-7037\(69\)90038-6](https://doi.org/https://doi.org/10.1016/0016-7037(69)90038-6), 1969.
- de Nooijer, L. J., Toyofuku, T., and Kitazato, H.: Foraminifera promote calcification by elevating their intracellular pH, *Proceedings of the*
- 305 *National Academy of Sciences*, 106, 15 374–15 378, 2009.
- de Nooijer, L. J., Spero, H. J., Erez, J., Bijma, J., and Reichart, G. J.: Biomineralization in perforate foraminifera, *Earth-Science Reviews*, 135, 48–58, 2014.
- Debenay, J.-P., Guillou, J.-J., and Lesourd, M.: Colloidal calcite in foraminiferal tests; crystallization and texture of the test, *The Journal of Foraminiferal Research*, 26, 277–288, 1996.
- 310 Debenay, J.-P., Guillou, J.-J., Geslin, E., Lesourd, M., and Redois, F.: Processus de cristallisation de plaquettes rhomboédriques à la surface d'un test porcelané de foraminifère actuel, *Geobios*, 31, 295–302, 1998.
- Debenay, J.-P., Guillou, J.-J., Geslin, E., and Lesourd, M.: Crystallization of calcite in foraminiferal tests, *Micropaleontology*, 46, 87–94, 2000.
- Eggins, S. M., Sadekov, A., and De Deckker, P.: Modulation and daily banding of Mg/Ca in *Orbulina universa* tests by symbiont photosynthesis and respiration: a complication for seawater thermometry?, *Earth and Planetary Science Letters*, 225, 411–419, 2004.
- Elderfield, H., Bertram, C., and Erez, J.: A biomineralization model for the incorporation of trace elements into foraminiferal calcium carbonate, *Earth and Planetary Science Letters*, 142, 409–423, 1996.
- Erez, J.: The source of ions for biomineralization in foraminifera and their implications for paleoceanographic proxies, *Reviews in mineralogy and geochemistry*, 54, 115–149, 2003.
- 320 Evans, D., Erez, J., Oron, S., and Müller, W.: Mg/Ca-temperature and seawater-test chemistry relationships in the shallow-dwelling large benthic foraminifera *Operculina ammonoides*, *Geochimica et Cosmochimica Acta*, 148, 325–342, 2015.



- Fehrenbacher, J. S., Russell, A. D., Davis, C. V., Gagnon, A. C., Spero, H. J., Cliff, J. B., Zhu, Z., and Martin, P.: Link between light-triggered Mg-banding and chamber formation in the planktic foraminifera *Neogloboquadrina dutertrei*, *Nature communications*, 8, 15 441, 2017.
- 325 Gattuso, J.-P., Epitalon, J.-M., Lavigne, H., and Orr, J.: seacarb: Seawater Carbonate Chemistry, <https://CRAN.R-project.org/package=seacarb>, r package version 3.2.10, 2018.
- Geerken, E., de Nooijer, L. J., van Dijk, I., and Reichart, G.-J.: Impact of salinity on element incorporation in two benthic foraminiferal species with contrasting magnesium contents, *Biogeosciences*, 15, 2205–2218, 2018.
- Geerken, E., de Nooijer, L. J., Roepert, A., Polerecky, L., King, H. E., and Reichart, G.-J.: Element banding and organic linings within chamber walls of two benthic foraminifera, *Scientific Reports*, 9, 3598, <https://doi.org/10.1038/s41598-019-40298-y>, 2019.
- 330 Glock, N., Liebetrau, V., Vogts, A., and Eisenhauer, A.: Organic Heterogeneities in Foraminiferal Calcite Traced Through the Distribution of N, S, and I Measured With NanoSIMS: A New Challenge for Element-Ratio-Based Paleoproxies?, *Frontiers in Earth Science*, 7, 2019.
- Hemleben, C., Erson, O., Berthold, W., and Spindler, M.: Calcification and chamber formation in Foraminifera—a brief overview, 1986.
- Ichikuni, M.: Uptake of fluoride by aragonite, *Chemical Geology*, 27, 207–214, [https://doi.org/10.1016/0009-2541\(79\)90039-1](https://doi.org/10.1016/0009-2541(79)90039-1), 1979.
- 335 Jonkers, L., Buse, B., Brummer, G.-J. A., and Hall, I. R.: Chamber formation leads to Mg/Ca banding in the planktonic foraminifer *Neogloboquadrina pachyderma*, *Earth and Planetary Science Letters*, 451, 177–184, 2016.
- Kendrick, M. A.: Halogens in seawater, marine sediments and the altered oceanic lithosphere, in: *The role of halogens in terrestrial and extraterrestrial geochemical processes*, pp. 591–648, Springer, 2018.
- Kitano, Y., Okumura, M., and Idogaki, M.: Incorporation of sodium, chloride and sulfate with calcium carbonate, *Geochem. J.*, 9, 75–84, 1975.
- 340 Kunioka, D., Shirai, K., Takahata, N., Sano, Y., Toyofuku, T., and Ujiie, Y.: Microdistribution of Mg/Ca, Sr/Ca, and Ba/Ca ratios in *Pulleniatina obliquiloculata* test by using a NanoSIMS: Implication for the vital effect mechanism, *Geochemistry, Geophysics, Geosystems*, 7, 2006.
- Lea, D. W., Mashiotta, T. A., and Spero, H. J.: Controls on magnesium and strontium uptake in planktonic foraminifera determined by live culturing, *Geochimica et Cosmochimica Acta*, 63, 2369–2379, 1999.
- 345 Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G.-H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, *Geophysical research letters*, 33, <https://doi.org/10.1029/2006GL027207>, 2006.
- Lu, Z., Jenkyns, H. C., and Rickaby, R. E.: Iodine to calcium ratios in marine carbonate as a paleo-redox proxy during oceanic anoxic events, *Geology*, 38, 1107–1110, 2010.
- 350 Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean  $p\text{CO}_2$  calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of  $p\text{CO}_2$  in gas and seawater at equilibrium, *Marine Chemistry*, 70, 105–119, 2000.
- Martens, C. S. and Harriss, R. C.: Inhibition of apatite precipitation in the marine environment by magnesium ions, *Geochimica et Cosmochimica Acta*, 34, 621–625, 1970.
- 355 Mezger, E. M., de Nooijer, L. J., Bertlich, J., Bijma, J., Nürnberg, D., and Reichart, G.-J.: Planktonic foraminiferal spine versus shell carbonate Na incorporation in relation to salinity, *Biogeosciences*, 16, 1147–1165, 2019.
- Millero, F., Huang, F., Zhu, X., Liu, X., and Zhang, J.-Z.: Adsorption and desorption of phosphate on calcite and aragonite in seawater, *Aquatic Geochemistry*, 7, 33–56, 2001.



- Nehrke, G., Keul, N., Langer, G., De Nooijer, L., Bijma, J., and Meibom, A.: A new model for biomineralization and trace-element signatures of Foraminifera tests, *Biogeosciences*, 10, 6759–6767, 2013.
- Okumura, M., Kitano, Y., and Idogaki, M.: Incorporation of fluoride ions into calcite—Effect of organic materials and magnesium ions in a parent solution, *Geochemical Journal*, 17, 257–263, <https://doi.org/https://doi.org/10.2343/geochemj.17.257>, 1983.
- Okumura, M., Kitano, Y., and Idogaki, M.: Behavior of bromide ions during the formation of calcium carbonate, *Marine Chemistry*, 19, 109–120, 1986.
- Opdyke, B. N., Walter, L. M., and Huston, T. J.: Fluoride content of foraminiferal calcite: Relations to life habitat, oxygen isotope composition, and minor element chemistry, *Geology*, 21, 169–172, 1993.
- Paris, G., Fehrenbacher, J. S., Sessions, A. L., Spero, H. J., and Adkins, J. F.: Experimental determination of carbonate-associated sulfate  $\delta^{34}\text{S}$  in planktonic foraminifera shells, *Geochemistry, Geophysics, Geosystems*, 15, 1452–1461, 2014.
- Polerecky, L., Adam, B., Milucka, J., Musat, N., Vagner, T., and Kuypers, M. M.: Look@NanoSIMS—a tool for the analysis of nanoSIMS data in environmental microbiology, *Environmental microbiology*, 14, 1009–1023, 2012.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2018.
- Rae, J. W. B., Foster, G. L., Schmidt, D. N., and Elliott, T.: Boron isotopes and B/Ca in benthic foraminifera: Proxies for the deep ocean carbonate system, *Earth and Planetary Science Letters*, 302, 403–413, 2011.
- Roepert, A.: Imaging element distributions within small marine calcifiers: a NanoSIMS perspective, PhD Thesis, Utrecht Series of Earth Sciences 201, 2019.
- Rosenthal, Y. and Boyle, E. A.: Factors controlling the fluoride content of planktonic foraminifera: An evaluation of its paleoceanographic applicability, *Geochimica et Cosmochimica Acta*, 57, 335–346, 1993.
- Rosenthal, Y., Boyle, E. A., and Slowey, N.: Temperature control on the incorporation of magnesium, strontium, fluorine, and cadmium into benthic foraminiferal shells from Little Bahama Bank: Prospects for thermocline paleoceanography, *Geochimica et Cosmochimica Acta*, 61, 3633–3643, [https://doi.org/https://doi.org/10.1016/S0016-7037\(97\)00181-6](https://doi.org/https://doi.org/10.1016/S0016-7037(97)00181-6), 1997.
- Rude, P. D. and Aller, R. C.: Fluorine mobility during early diagenesis of carbonate sediment: An indicator of mineral transformations, *Geochimica et Cosmochimica Acta*, 55, 2491–2509, [https://doi.org/https://doi.org/10.1016/0016-7037\(91\)90368-F](https://doi.org/https://doi.org/10.1016/0016-7037(91)90368-F), 1991.
- Sadekov, A. Y., Eggins, S. M., and de Deckker, P.: Characterization of Mg/Ca distributions in planktonic foraminifera species by electron microprobe mapping, *Geochemistry, Geophysics, Geosystems*, 6, 2005.
- Spero, H. J., Eggins, S. M., Russell, A. D., Vetter, L., Kilburn, M. R., and Hönisch, B.: Timing and mechanism for intratest Mg/Ca variability in a living planktic foraminifer, *Earth and Planetary Science Letters*, 409, 32–42, 2015.
- Szafranek, D. and Erez, J.: chemistry of Mg,  $\text{SO}_4^{2-}$ , Sr, Na, and Cl in live foraminifera shells, in: *The 7th International Symp. on Biomineralization, Program and Abstracts*, 1993.
- Tanaka, K., Ono, T., Fujioka, Y., and Ohde, S.: Fluoride in non-symbiotic coral associated with seawater carbonate, *Marine Chemistry*, 149, 45–50, 2013.
- Tokuyama, A., Kitano, Y., and Kaneshima, K.: Geochemical behavior of chemical species in the processes of limestone formation. Part I. Chemical composition of corals and limestones in the Ryukyu Islands, *Geochem. J.*, 6, 83–92, 1972.
- Toyofuku, T., Matsuo, M. Y., de Nooijer, L. J., Nagai, Y., Kawada, S., Fujita, K., Reichart, G.-J., Nomaki, H., Tsuchiya, M., Sakaguchi, H., and Kitazato, H.: Proton pumping accompanies calcification in foraminifera, *Nature communications*, 8, 14 145, 2017.



- van Dijk, I., de Nooijer, L. J., and Reichart, G.-J.: Trends in element incorporation in hyaline and porcelaneous foraminifera as a function of  $p\text{CO}_2$ , *Biogeosciences*, 14, 497–510, <https://doi.org/10.5194/bg-14-497-2017>, <https://www.biogeosciences.net/14/497/2017/>, 2017.
- van Dijk, I., Barras, C., de Nooijer, L. J., Mouret, A., Geerken, E., Oron, S., and Reichart, G.-J.: Coupled Ca and inorganic carbon uptake suggested by magnesium and sulfur incorporation in foraminiferal calcite, *Biogeosciences*, 16, 2115–2130, <https://doi.org/10.5194/bg-16-2115-2019>, 2019.
- Wei, T. and Simko, V.: R package "corrplot": Visualization of a Correlation Matrix, <https://github.com/taiyun/corrplot>, (Version 0.84), 2017.
- Wetmore, K. L.: Chamber formation in *Archaias angulatus*, *The Journal of Foraminiferal Research*, 29, 69–74, 1999.
- Wit, J. C., De Nooijer, L., Wolthers, M., and Reichart, G.-J.: A novel salinity proxy based on Na incorporation into foraminiferal calcite, *Biogeosciences*, 10, 6375–6387, 2013.
- 405 Yu, J., Elderfield, H., and Hönisch, B.: B/Ca in planktonic foraminifera as a proxy for surface seawater pH, *Paleoceanography*, 22, 2007.





**Table 1.** Studied foraminifera specimens and the corresponding culture conditions.


#	species <sup>1</sup>	salinity	T [°C]	DIC [μmol/kg]	alkalinity [μmol/kg]	pH	CO <sub>3</sub> <sup>2-</sup> [μmol/kg]	Ω <sub>calcite</sub>
1 <sup>†</sup>	<i>A. tepida</i> (R)	25.2	25	1087	1350	8.32	162	4.22
2 <sup>‡</sup>	<i>A. lessonii</i> (R)	35.2	21.2	2069	2314	8.00	177	4.24
3*	<i>A. angulatus</i> (M)	40	25	3861	4477	8.10	506	11.67
4*	<i>A. angulatus</i> (M)	30	25	2644	3153	8.27	399	10.00
5*	<i>S. marginalis</i> (M)	30	25	2644	3153	8.27	399	10.00
6*	<i>S. marginalis</i> (M)	40	25	3861	4477	8.10	506	11.67

<sup>1</sup> R = rotaliid, M = miliolid.


<sup>†</sup> Selected from the culture experiment by Geerken et al. (2018).

<sup>‡</sup> Selected from the culture experiment by van Dijk et al. (2019).


\* Selected from an unpublished culture experiment (see Appendix B).




**Figure 1.** Spatial distribution of Cl/Ca, F/Ca, and P/Ca ratios in the calcite shells of the studied foraminifera species. Shown are representative images as well as lateral profiles along a line going from the inside to the outside of the shell as depicted by an arrow in the images. Note that the displayed ratio images are log-transformed, and that the color-scale for the given ratio is the same for all species. Blacked out areas correspond to resin. The scale bar in each image is 5  $\mu\text{m}$ .




**Figure 2.** Variability of the different EI/Ca ratios within shells of the studied foraminifera. Individual data-points represent multiple ROIs drawn on the images as described in Section 2.4. Symbol shapes depict different specimens.



**Figure 3.** Crossplots of NanoSIMS  $^{35}\text{Cl}/^{40}\text{Ca}^{16}\text{O}$  and  $^{19}\text{F}/^{40}\text{Ca}^{16}\text{O}$  ion count ratios with LA-ICP-MS EI/Ca ratios in the same specimen. The error bars depict one standard error of the mean NanoSIMS ion count ratios where more than one image was analysed per specimen, and one standard error of duplicate or triplicate LA-ICP-MS measurements.



**Figure 4.** Crossplots of NanoSIMS  $^{35}\text{Cl}/^{40}\text{Ca}^{16}\text{O}$  and  $^{19}\text{F}/^{40}\text{Ca}^{16}\text{O}$  ion count ratios with carbonate system parameters of the culture media. The error bars depict one standard error of the mean NanoSIMS ion count ratios where more than one image was analysed per specimen.




**Figure 5.** Scheme highlighting differences of the calcification mechanisms in miliolid and rotaliid foraminifera. Miliolid foraminifera precipitate calcium crystals in intracellular vesicles prior to arranging them in the shape of the new chamber wall (A). Rotaliid foraminifera precipitate calcite onto organic templates within an extracellular but confined space, and add a new lamella to the entire shell each time they produce a new chamber (B, C). In intermediate Mg-calcite producing rotaliid foraminifera (like *A. lessonii*), transport of vesicles to the site of calcification has been observed suggesting controlled biomineralization (at least partly) from internal Ca and carbonate pools (B). In rotaliid low Mg-calcite producing foraminifera, selective ion transport to the site of calcification via trans-membrane pumping of elements is discussed as the biological control on biomineralization (C).







## Appendix A: Supplementary tables and figures




**Figure A1.** SEM images of the analysed specimens with the areas imaged by NanoSIMS indicated by white or yellow squares: *Ammonia tepida* (#1), *Amphistegina lessonii* (#2), *Archaias angulatus* (#3, #4), *Sorites marginalis* (#5, #6). The specimen numbers correspond to those in Table 1 and A1. The yellow squares indicate the locations of the NanoSIMS images shown in Figure 1. SEM images are flipped horizontally to facilitate navigation in the NanoSIMS instrument, where the secondary ion images are horizontally mirrored.




**Figure A2.** NanoSIMS  $^{40}\text{Ca}^{16}\text{O}$ /plane/pixel and  $^{16}\text{O}$ /plane/pixel in four foraminifera species. The NanoSIMS ion count rates of  $^{40}\text{Ca}^{16}\text{O}$  and  $^{16}\text{O}$  highly correlate in calcite (Figure A2), which is why elemental ratios can be normalized to both for display. Since elemental ratios determined in bulk calcite are normalized to calcium, we present the NanoSIMS data mostly as  $\text{El}/^{40}\text{Ca}^{16}\text{O}$  ratios.



**Figure A3.** Correlation matrices of NanoSIMS  $^{19}\text{F}/^{40}\text{Ca}^{16}\text{O}$ ,  $^{31}\text{P}/^{40}\text{Ca}^{16}\text{O}$ , and  $^{35}\text{Cl}/^{40}\text{Ca}^{16}\text{O}$  ion count ratios per foraminiferal species across n ROIs. Coloured values in the matrix are  $R^2$  and the correlation is significant to a level of  $p < 0.001$  wherever no black cross present. The plot was created using the corrplot package in R (Wei and Simko, 2017; R Core Team, 2018).



**Figure A4.** NanoSIMS  $^{35}\text{Cl}/^{40}\text{Ca}^{16}\text{O}$  and  $^{19}\text{F}/^{40}\text{Ca}^{16}\text{O}$  ion count ratios in the measured specimen versus temperature and salinity of the culture media. The error bars depict the standard deviation of the mean NanoSIMS ion count ratios for those specimen where more than one image was analysed on the shell.



**Figure A5.** Illustration to show that the patterns in figure 2 do not change much when normalizing to  $^{16}\text{O}$  instead of  $^{40}\text{Ca}^{16}\text{O}$ .



**Table A1.** Additional information of specimens. LA-ICP-MS data of average single shell E/Ca ratios were determined on several chambers.

#	specimen ID*	species	symbionts	Na/Ca [mmol/mol]	Mg/Ca [mmol/mol]	Sr/Ca [mmol/mol]	Al/Ca [mmol/mol]	Ba/Ca [mmol/mol]
1	34_61	<i>Ammonia tepida</i>	no	3.48 ± 0.03	2.0 ± 0.1	1.26 ± 0.05	0.009 ± < 0.000	0.0026 ± 0.0002
2	13_110	<i>Amphistegina lessonii</i>	yes	9.27 ± 1.10	22.7 ± 3.8	1.57 ± 0.10	0.008 ± < 0.000	0.0016 ± 0.0001
3	32_89	<i>Archaias angulatus</i>	yes	3.75 ± 0.22	142.0 ± 1.0	2.05 ± 0.06	0.011 ± < 0.000	0.0057 ± 0.0001
4	46_87	<i>Archaias angulatus</i>	yes	4.73 ± 0.22	143.0 ± 4.2	0.01 ± < 0.00	0.008 ± 0.001	0.0070 ± 0.0002
5	01_87	<i>Sorites marginalis</i>	yes	3.47 ± 0.27	150.0 ± 5.2	1.92 ± 0.07	0.153 ± 0.093	0.0081 ± < 0.000
6	30_89	<i>Sorites marginalis</i>	yes	5.24 ± 0.19	146.3 ± 2.5	2.08 ± 0.02	0.044 ± 0.013	0.0075 ± 0.0005

\* Lab internal specimen ID based on stub#\_number. For *A. tepida* as in Geerken et al. (2018).





**Table A2.** NanoSIMS 50L acquisition settings.

NanoSIMS settings*	<i>Ammonia tepida</i>	<i>Amphistegina lessonii</i>	<i>Archaias angulatus</i>	<i>Sorites marginalis</i>
<b>Pre-sputtering</b>				
beam current (in $FC_o$ ) [pA]	~ 140 (preset 10 in D1–3)	~ 280 (preset 20 in D1–3)	~ 280 (preset 20 in D1–3)	~ 140–280 (preset 10–20 in D1–3)
diaphragm and slits	D1–1	D1–1	D1–1	D1–1
FOV [ $\mu\text{m}^2$ ]	8 × 8	12 × 12 to 23 × 23	55 × 55	25 × 25 to 40 × 40
time [min]	10	10	15	10–15
eGun	on	on	on	on
<b>Image Aquisition</b>				
beam current (in $FC_o$ ) [pA]	0.5	1	2	1
diaphragm and slits	D1–3, ES–3, AS–2, EnS–1	D1–3, ES–3, AS–2, EnS–1	D1–3, ES–3, AS–2, EnS–1	D1–3, ES–3, AS–2, EnS–1
Detected masses	$^{12}\text{C}$ , $^{16}\text{O}$ , $^{19}\text{F}$ , $^{31}\text{P}$ , $^{35}\text{Cl}$ , $^{37}\text{Cl}$ , $^{40}\text{Ca}$ $^{16}\text{O}$	$^{12}\text{C}$ , $^{16}\text{O}$ , $^{19}\text{F}$ , $^{31}\text{P}$ , $^{35}\text{Cl}$ , $^{37}\text{Cl}$ , $^{40}\text{Ca}$ $^{16}\text{O}$	$^{12}\text{C}$ , $^{16}\text{O}$ , $^{19}\text{F}$ , $^{31}\text{P}$ , $^{35}\text{Cl}$ , $^{37}\text{Cl}$ , $^{40}\text{Ca}$ $^{16}\text{O}$	$^{12}\text{C}$ , $^{16}\text{O}$ , $^{19}\text{F}$ , $^{31}\text{P}$ , $^{35}\text{Cl}$ , $^{37}\text{Cl}$ , $^{40}\text{Ca}$ $^{16}\text{O}$
dwell time [ $\mu\text{s pixel}^{-1}$ ]	1000	1000	1000	1000
image FOV [ $\mu\text{m}^2$ ]	6 × 6	8 × 8 to 20 × 20	50 × 50	23 × 23 to 30 × 30
image resolution [px × px]	128 × 128	128 × 128, once 256 × 256	256 × 256	256 × 256
number of planes	1000	1000, once 250	250	250
eGun	on	on	on	on

\* Abbreviations: objective current ( $FC_o$ ); field of view (FOV); electron flood gun (eGun).

## Appendix B: Culture conditions of the foraminiferal species

The *A. angulatus* and *S. marginalis* specimens were collected in Sint Eustatius (Oranjestad Bay, 17.479751°N –62.987273°W). The culture experiments with *A. angulatus* and *S. marginalis* were conducted in the same manner as described in van Dijk et al. (2017), with the exception of media preparation. Culture media of different salinities were prepared by mixing natural 0.2  $\mu\text{m}$  filtered seawater with deionized water and 'instant ocean' salt, to obtain a range in salinities between 25–45. The *A. lessonii* specimens are from Burger's Zoo (van Dijk et al., 2019); the culture conditions are reported in van Dijk et al. (2019). The specimens of *A. tepida* were collected on a tidal flat near Den Oever, the Wadden Sea, NL (Hayward et al. 2004); the culture conditions are reported in Geerken et al. (2018).