# Population dynamics of modern planktonic foraminifera in the western Barents Sea

Julie Meilland<sup>1</sup>, Hélène Howa<sup>2</sup>, Vivien Hulot<sup>3,4</sup>, Isaline Demangel<sup>5,6</sup>, Joëlle Salaün<sup>7</sup>, Thierry Garlan<sup>7</sup>

<sup>1</sup> MARUM - Center for Marine Environmental Sciences, Leobener Str. 8, D-28359, Bremen, Germany

<sup>2</sup> LPG-BIAF, UMR-CNRS 6112; University of Angers, France

<sup>3</sup> University of French Polynesia, UMR-241 EIO, Labex Corail, FAA'A, Tahiti, French Polynesia

<sup>4</sup> Ifremer, UMR-241 EIO, Labex Corail, Departement Ressources Biologiques et Environnement, Vairao, Tahiti, French Polynesia

<sup>5</sup> Institute of Earth Sciences, University of Graz, NAWI Graz Geocenter, Graz, Austria

<sup>6</sup> Department of Geology, University of Lund, Lund, Sweden
 <sup>7</sup> SHOM –Sciences et Techniques Marines/Géologie Marine, Brest, France

Correspondence to: Julie Meilland (jmeilland@marum.de)

**Abstract.** This study reports on species diversity and distribution of planktonic foraminifera (PF) at the Barents Sea Opening (BSO). Populations of PF living in late summer (collected by means of stratified plankton tows) and recently settled

- 15 individuals (sampled by interface corer) were studied and compared. High abundances reaching up to 400 ind.m<sup>-3</sup> in tow samples and 8000 ind.cm<sup>-3</sup> in surface sediments were recorded in the centre of the studied area while low abundances were observed in coastal areas, likely hampered by continental influences. The living and core-top assemblages are mainly composed of the same four species *Neogloboquadrina pachyderma*, *Neogloboquadrina incompta*, *Turborotalita quiqueloba* and *Globigerinita uvula*. The two species *G. uvula* and *T. quiqueloba* largely dominate the upper water column whereas
- 20 surface sediment assemblages display especially high concentrations of *N. pachyderma*. The unusual dominance of *G. uvula* in the water sample assemblages compared to its low occurrence in surface sediments might be the signature of 1) a seasonal signal due to summer phytoplankton composition changes at the BSO, linked to the increase of summer temperature at the study site, and/or 2) a signal of a larger time-scale and wide geographical reach phenomenon inducing poleward temperate/subpolar species migration and consecutive foraminiferal assemblage diversification at high latitudes under global
- 25 climate forcing. Protein concentrations were measured on single specimens and used as a proxy of individual carbon biomass. Specimens of all species show the same trend, i.e. a northward decrease of their size-normalized-protein concentration suggesting foraminiferal biomass to be potentially controlled by different constituents of their organelles (e.g. lipids). The originality of coupling data from plankton tows, protein measurements and surface sediments allows us to hypothesise that PF dynamics (seasonality and distribution) are decoupled from their metabolism.

30

5

# **1** Introduction

Polar areas are sensitive to global temperature changes, particularly in the Arctic where warming occurs faster than in the rest of the world and has accelerated over the past 50 years (Shepherd, 2016). This Arctic amplification appears to be mainly

- 35 caused by sea-ice loss under increasing CO<sub>2</sub> (Dai et al., 2019). Recent increased advection of Atlantic Water in the Barents Sea modifies its physico-chemical properties (Smedsrud et al., 2013), which gets directly reflected in the entire ecology of the region. Higher temperatures lead to increased rates of planktonic primary production (Vaquer-Sunyer et al., 2013) and increased CO<sub>2</sub> concentrations are expected to have a fertilization effect on marine autotrophs (Holding et al., 2015). Enhanced primary production is accompanied by lateral shifts of the spring and summer phytoplankton blooms in the 40 European Arctic Ocean (Oziel et al., 2017). As a response some taxa of different calcifying groups (i.e. foraminifera,
- 40 European Arctic Ocean (Oziel et al., 2017). As a response some taxa of different calcifying groups (i.e. foraminifera, coccolithophores, molluscs and echinoderms; Beaugrand et al., 2013) exhibit a poleward movement in agreement with expected biogeographical changes under sea temperature warming. Both satellite images (Smyth et al., 2004; Burenkov et al., 2011) and *in situ* measurements (Dylmer et al., 2013; Giraudeau et al., 2016; AMAP 2018) have recorded rapid expansion of temperate species of coccolithophores in the Arctic. For example, *Emiliania huxleyi* shows a striking poleward
- 45 shift (>5°) in the distribution of its blooms (Neukermans et al., 2018). Such phenomenon, called "atlantification" (Årthun et al., 2012), are expected to impact every trophic levels of the food web, from small phytoplanktonic species (Neukermans et al., 2018) to larger organisms (Dalpadado et al., 2012). Recent studies have investigated the ecology and biodiversity of planktonic foraminifera from the high-latitude North Atlantic (i.e. Schiebel et al., 2017). Eynaud, (2011) noticed that the species *N. pachyderma*, the most characteristic high-latitude taxon, dominates the past interglacial assemblages for the last
- 50 1.8 Ma. while the subpolar species *Turborotalia quinqueloba* records northward penetration of Atlantic warm water masses in the Arctic, especially during interglacial periods. The species *N. pachyderma* comprises more than 90% of recent assemblages (i.e. found in surface sediments) from the Polar Region, North of Iceland (MARGO data base; Kucera et al., 2005). Rather few studies on living planktonic foraminifera (PF) communities have concentrated on (sub-) Arctic regions. Pados and Spielhagen (2014) observed PF by the means of plankton tows, in the cold and fresh Polar waters of the Fram
- 55 Strait during early summer. They report a large dominance of *N. pachyderma* and a co-occurrence of *T. quinqueloba*, accounting for 90 and 5% of all tests, respectively. Volkmann, (2000) also documented this large dominance of *N. pachyderma* and the co-occurrence with *T. quinqueloba*, overall in the Arctic. Through the compilation of population density profiles from 104 stratified plankton tow hauls collected in the Arctic and the North Atlantic Oceans, Greco et al., (2019) deeply investigated the ecology of *N. pachyderma*. In particular, the variability of its habitat depth, and finally underlined the
- 60 knowledge gap on its ecological preferences. In the western subpolar North Atlantic (Irminger Sea), the maximum production of *N. pachyderma* shows two peaks, in spring and late summer, while winter shows a low production (Jonkers et al., 2010; 2013). Following the extensive review of Schiebel et al., (2017), diversity of planktonic foraminifera has increased

in polar waters over the past decades, even though it remains low in comparison to lower latitudes. Some species from lower latitudes are described as new components of formerly high-latitude assemblages (Southern Indian Ocean; Meilland et al.,

- 65 2016). The shift of planktonic foraminifera assemblages to warmer conditions, since the pre-industrial stage, has been very recently highlighted more globally in the Northern hemisphere (Jonkers et al., 2019). These major modifications in PF distribution patterns display changes more in primary production than in water temperature itself (e.g. Jonkers et al., 2010; Schiebel et al., 2017). Planktonic foraminifera, being sensitive to ambient water geochemistry, are considered good indicators of the polar changing environments (Schiebel et al., 2017). More studies on living PF communities in the Arctic
- 70 regions are needed to assess the spatial and temporal variability in their population dynamics and to better constrain the today's polar and subpolar species ecological preferences. Taking the opportunity of a cruise dedicated to the exploration of the physical oceanography of the western Barents Sea (MOCOSED 2014 cruise), we investigated the connections between the spatial variability of living planktonic foraminifera, phytoplankton communities (Giraudeau et al., 2016), and the hydrological system through a South-to-North transect,
- 75 between Northern Norway and Spitsbergen [68-76]°N. Along this transect, we compared PF living faunas (from plankton tow) to the assemblages found on the sea floor (from core-top sediments) in order to investigate eventual recent changes in their population dynamics. This latitudinal transect also gave us the opportunity to quantify protein concentrations of individual living PF in this area for the first time and along a physico-chemical gradient to see if and how it varies and explore how planktonic foraminifera from a same species may adjust to different environments.

# 80 2. Oceanographic setting

The studied area covers the western Barents Sea margin, i.e. Barents Sea Opening (BSO), where surface and intermediate ocean circulations are characterised by the confrontation of the North Atlantic and the Arctic Waters (Figure 1). The seasonal and interannual dynamics of these two water masses, interacting with the complex topography of the western margin (Spitsbergen Banken and shallow Bjørnøya; Storfjordrenna and Bjømøyrenna glacial troughs), determine position and

- 85 meandering of the Polar Front (Loeng, 1991). The Norwegian Atlantic Current (NwAC) carries Atlantic Water into the Barents Sea. Along the western Barents Sea margin, Atlantic Water is then transported to the Fram Strait by the West Spitsbergen current (Skagseth et al., 2008; Oziel et al., 2017; Figure 1). Environmental parameters (temperature, salinity and fluorescence) as well as phytoplankton composition were obtained during the MOCOSED 2014 cruise using a total of 32 vertical casts deployed ≈ 20 km apart from each other (Figure 1 and 2,
- 90 Giraudeau et al., 2016). In the southern first half of the transect a strong thermohalocline clearly underlined a surface mixed layer of about 30-35 m depth. This cline slightly deepened northwards and blurred out north of 74.5°N where no more stratification was observed in the water column. From South to North of the transect: i) high temperatures and low salinities reflected the Norwegian Coastal Water (NwCW), also enriched in Chl-a. The relatively warm NwCW (8.5 to 11°C) extended northwards up to 74.5°N overlying the colder Norwegian Atlantic Water (NwAW). Less saline (33.5) to the South,
- 95 NwCW became saltier (34.9) in the vicinity of the Spitsbergen Banken; ii) at the northern end of the transect, the NwAW

penetrated the Barents Sea through the Storfjordrenna trough with temperatures from 6 to 8°C and an open marine salinity of 35.1.

The Chl-*a* content followed the hydrological pattern above described (Figure 2 c). Relatively high concentrations (mean  $\approx$  0.8 mg.m<sup>-3</sup>) were located in the surface mixed layer composed of NwCW. The highest values around 1.25 mg.m<sup>-3</sup> were

- 100 recorded off the Norwegian coast. Chl-*a* content decreased northwards (north of 74.5°N) to reach  $\approx 0.4$  mg.m<sup>-3</sup> in the upper layer (0-60m) of the well-mixed NwAW. The composition of the phytoplankton community observed in surface water at 7 stations along the studied transect was essentially dominated by three algal groups (Giraudeau at al., 2016): Fuco-flagelattes (25 to 43%; major component *Phaeocystis pouchetii*), Prasinophytes (15 to 30%; major components *Micromonas pusilla* and *Bathycoccus pusilla*) and Prymnesiophytes (13 to 24%; major component *Emiliana huxleyi*). Three other features are
- 105 noteworthy (Figure 2 d): i) the dominance of dinoflagelattes (24%) at the southernmost station of the transect (close to the Norwegian coast) contrasted with its total absence in the well mixed NwAW, North of 74.5; ii) the presence of diatoms (10-20 %) in the surficial NwCW, but rare (<5%) to the North; iii) the constant increase in relative abundance of Cyanobacteria from < 5% to more than 15%, along the South-to-North transect.</p>

# 3. Material and Methods

110 In late summer 2014 from August 8 to September 20, the SHOM (French Hydrographic Office) operated the oceanographic cruise MOCOSED 2014, on board the "R/V *Pourquoi pas* ?". Along a 700 km South-to-North transect from the Norwegian (68°N) to the Spitzberg (76°N) coasts, investigations of hydrological processes at the BSO were carried out coupled with the exploration of the phytoplankton and foraminiferal communities (Figure 1).

# 3.1. Living planktonic foraminifera from stratified plankton samples (MultiNet)

- 115 Living PF were collected at 7 of the 32 CTD South-to-North transect stations (#3 to #9), and at 2 stations (#1 and #2) located West-to-East ≈ 20 km apart from the central point of the main South-to-North CTD transect (Figure 1; Table 1), using a stratified plankton tow (MultiNet Hydro-Bios type Midi, opening of 0.25 m2) equipped with five nets (mesh size 100 µm to avoid nets clogging in case of intense phytoplankton bloom). This collection was set in order to observe the potential effect of latitudinal changes but also of bathymetry, longitudinally, on PF distribution. At each station, one single vertical haul sampled five successive water layers from the sea surface to 100 m depth. For each of the five depth intervals (0–20 m, 20–40 m, 40–60 m, 60–80 m, 80–100 m), the filtered water volume was measured by means of a flowmeter attached to the MultiNet mouth. Each MultiNet sample was preserved in a 250 mL vial with ethanol (90%) buffered with hexamethylenetetramine until processing at the land-based laboratory. Back at the laboratory, MultiNet samples were washed over a 100 µm mesh, all foraminifera were removed from the sample and dried in an oven at 50 °C. All living PF,
- 125 distinguished by their coloured cytoplasm visible through the shell, were individually picked, stored in counting cells and identified at the species level, following the SCOR WG138 taxonomy as implemented in Siccha and Kucera (2017).

Correlations following a non-metric multidimensional scaling ordination (NMDS) were carried out with the R package Vegan (Oksanen et al., 2013). Using the Bray-Curtis distance these correlations were tested between PF species absolute abundances, the latitude of the station and parameters of the ambient waters (temperature, salinity, Chl-*a* concentration).

130 Empty tests, considered as dead individuals were separately numbered. Results are given in relative abundances (% of the total, live or dead fauna) or in absolute abundances in number of individuals per m<sup>3</sup> of filtered water (ind.m<sup>-3</sup>).

### Protein biomass and test size measurements

Immediately after sampling (on board), a few living individuals ( $\approx 60$ ) were picked out of the shallowest water samples (0 – 20 m) of the 7 stations sampled along the main CTD transect (stations 3 to 9) for protein extraction and measurement. Each 135 for a minifer picked for protein measurement was carefully selected under the strict condition of its shell to be fully filled with cytoplasm. After picking, individuals were immediately cleaned with a brush and filtered seawater to remove all particles stuck to the test including organic matter. Individual were stored in a 1.5 mL Eppendorf cup and analysed on board, using the bicinchoninic acid (BCA) method as explained in Meilland et al., (2016). Morphometric analyses on single foraminiferal tests were carried out at the University of Angers with an automated incident light microscope (Bollmann et al., 2004; Clayton et al., 2009) at a resolution of 1.4  $\mu$ m<sup>2</sup> (pixel size). Images were analysed for their two dimensional (silhouette) 140 morphometry (Beer et al., 2010), including foraminiferal test minimum diameter being the shortest distance wall to wall passing through the centre of the proloculus (the initial chamber of a foraminifer). Protein-to-size relations were determined for the minimum diameter of each test providing size-normalized protein content (SNP) for data analyses and handling. Foraminifera protein concentrations were linearly normalized to 1 µm minimum test diameter, being aware of any 145 unavoidable errors related to non-linear increments of biomass at volumetric test growth (cf. Beer et al., 2010).

#### 3.2. Fossil planktonic foraminifera assemblages from core-tops (Multitube)

At 5 sites of the main CTD transect, an interface corer (Multitube type Oktopus GmbH, INSU<sup>1</sup> division of Brest, France) was implemented to obtain simultaneously 8 short sediment cores (less than 1 m in length) (Figure 1; Table 1). At each station, the core with the more horizontal and undisturbed water-sediment interface was selected. The core-top sediment (0-0.5 cm slice) was sampled and fixed with 95 % ethanol and Rose Bengal. This organic stain reacting with cytoplasm was used here to distinguish PF still bearing cytoplasm (fresh or in degradation) and thus very recently deposited from empty tests of fossil PF. The last study on Rose Bengal-stained individuals of Foraminifera, focussing only on benthic ones did not remove the uncertainty about the exact duration of complete cytoplasm degradation in the tests (Schönfeld et al., 2013), thus we cannot be precise on the time-scale pointed out by Rose Bengal-stained specimens. Based on the hydrology, sites depth 155 (Table 1), and sediment oxidation over the studied area, we can reasonably think that Rose Bengal coloration is in our

5

situation highlighting spring and summer population that recently felt. The discussion will be based on this assumption.

<sup>&</sup>lt;sup>1</sup>INSU : Institut National des Sciences de l'Univers

For the purpose of this study, the core-top sediment has been wet sieved on a 100  $\mu$ m mesh (same as the plankton net mesh size), and analysed for the planktonic foraminiferal assemblages. Every picked planktonic foraminifer has been identified consistently with those collected by plankton tows.

# 160 4. Results

170

# 4.1. Planktonic foraminifera diversity and distribution in the water column

Data from the 7 stations of the South-to-North CDT transect with 5 values per station, were compiled to display the repartition of PF absolute abundances (for total assemblage and species-specific) in the upper 100 meters of the vertical section across the BSO. Data from station 1 (Western) and station 2 (Eastern) located 20 km on either side of the S-N transect, were compared to the data of station 6 at the middle of the transect.

Total absolute abundances of living PF fauna varied between 0 and 400 ind.m<sup>-3</sup> (Figure 3). Along the South-to-North CTD transect, the highest concentrations were all observed above 20 m water depth, in the surface mixed layer of the well stratified water area, i.e. in NwCW. The two stations located at the south and north extremities of the transect (# 3, off Norwegian coast, and #9, off the Spitsbergen coast) displayed low densities (10 to 50 ind.m<sup>-3</sup>). At station 1, located above

- the Barents Sea margin slope, the maximal abundance of 220 ind.m<sup>-3</sup> was recorded in a deeper habitat (20-40 m, Figure 3). Station 2, located relatively inside the Barents Sea, was very poor in PF (<10 ind.m<sup>-3</sup>). In total, 10 species were observed. The studied area was characterised by the large occurrence of subpolar to polar species (Figure 4), listed in descending order: *Globigerinita uvula* (45%), *Turborotalita quinqueloba* (26%), *Neogloboquadrina incompta* (15%) and *Neogloboquadrina*
- 175 pachyderma (9%). There was a noticeable presence of the temperate water species Globigerina bulloides (3%), and negligible percentages (<1%) of Globigerinita glutinata, Neogloboquadrina dutertrei, Globigerinoides ruber, Globigerinoides sacculifer and Orcadia riedeli.

In the surface waters of the South-to-North transect (0-20 m depth), except at the septentrional station 9, *G. uvula* was the most abundant species reaching 64 % of the total fauna at station 7. The second major species, *T. quinqueloba* dominated the

- PF fauna only at station 9, with 45 % of the total assemblage and 26 ind.m<sup>-3</sup>. Both *G. uvula* and *T. quinqueloba* have a patchy repartition with two patches of maximum abundances located in the first 0-20 m, at station 4 and 7 for *G. uvula* (175 and 245 ind.m<sup>-3</sup>, respectively) and at station 5 and 7 for *T. quinqueloba* (53 and 80 ind.m<sup>-3</sup>, respectively). The northernmost and common patch for both species is the more intense. In the 20-40 m deep layer at station 1 (West of the transect), these two species showed also relatively high concentrations (121 ind.m<sup>-3</sup> for *G. uvula*, and 30 ind.m<sup>-3</sup> for *T. quinqueloba*). The
- 185 two other major species *N. pachyderma* and *N. incompta* presented similar low relative abundances (4 to 22%, and 8 to 25%, respectively). For both species, maximum absolute abundance of about 40-45 ind.m<sup>-3</sup> occurred in the central part of the transect between [72-74°N].

The NMDS analyse of species abundances with regard to environmental parameters (latitude of the station, temperature, salinity, Chl-*a* concentration) indicates that none of the species-specific distribution displays a significant correlation to any

190 of the tested variables (p-values > 0.1). NMDS documents distributional affinity (Figure 5), with *N. pachyderma* and *N. incompta* plotting in the same area whereas *T. quinqueloba* and *G. uvula* plot separately from each other but also from the *N. pachyderma* / *N. incompta* area.

#### 4.2. Planktonic foraminifera protein biomass

- Individual protein content (BCA method) and associated test minimum diameter (i.e. the shortest distance wall to wall
  passing through the centre of the proloculus) were successfully measured for a total of 272 specimens of the 4 major species encountered along the South-North transect, including 32 specimens of *Neogloboquadrina pachyderma*, 58 *Neogloboquadrina incompta*, 72 *Globigerinita uvula* and 110 *Turborotalita quinqueloba*. A minimum of 5 individuals per species (until a max of 25) was selected at each sampled depth-interval of the 7 stations along the S-N transect, paying careful attention to sample the whole range of size variation observed in the population present in a net sample. For station 7,
  the protein extraction was successful for only one specimen of *N. pachyderma*. Therefore, no value is display for this species
- at this station in Figure 6.

Minimum diameters of the 272 selected tests cover a large size range, from 65 to 315  $\mu$ m with a median value of 160  $\mu$ m. *N. incompta* represent the biggest species with a median of 200  $\mu$ m, and *G. uvula* the smallest one with a median of 110  $\mu$ m (Table 2). For each studied species, the mean size is equal to the median size indicating that the size distribution of the picked tests is symmetric, thus making us confident that our test selection represents properly the natural test size range of

205 picked tests is symmetric, thus making us confident that our test selection represents properly the natural test size range of each studied species. The biomass of a single individual normalized by its test size (SNP), averages out about 0.0055 μm of protein per μm of foraminiferal shell diameter. It varies depending on species from 0.0004 (*G. uvula*) to 0.0426 μg (*T. quinqueloba*). The 4 studied species display similar trends in their SNP per station, characterized by a northward decrease from 70 to 74° (Figure 6). *T. quiqueloba* and *G. uvula* have slightly (but not significantly) higher relative protein 210 concentrations than *N. pachyderma* and *N. incompta*.

# 4.3. Planktonic foraminifera diversity and distribution in surface sediments

215

Interface undisturbed cores were retrieved from 5 stations of the South-to-North CTD transect (Table 1) to investigate the core-top sediment (0-0.5 cm slice). The dead PF assemblages were studied making the difference between Rose Bengal-stained showing recently dead individuals still bearing a non-degraded cytoplasm after post-mortem deposition, and colourless empty tests of individuals dead for longer periods of time.

Concentrations of planktonic foraminifera with colourless empty tests (Figure 7 a) varied from a maximum of 6200 ind.cm<sup>-3</sup> at station 4 (71.3°N) to a minimum of 200 ind.cm<sup>-3</sup> at the septentrional station 9. All along the S-N transect, the fossil

assemblages were dominated by *Neogloboquadrina pachyderma* (31 to 59%). Assemblages were more balanced at the two ends of the transect where *N. pachyderma* showed off at its lowest occurrence. At the southernmost point, station 3 presents

- 220 a co-occurrence with *Turborotalita quinqueloba* (33%) and *Neogloboquadrina incompta* (24%). While at the northernmost point, station 9, *T. quinqueloba* (23%) co-occurred with *Globigerinita uvula* (25%). Concentrations of planktonic foraminifera bearing a coloured cytoplasm (Figure 7 b) varied from 100 to 300 ind.cm<sup>-3</sup>. All along the transect, the relative abundance of *N. pachyderma* remained between 10 and 26 %. The species *T. quinqueloba* occurred everywhere above 20% and up to 40% South of 72°N. The central station 6 was largely dominated by *G. uvula* (38%). North of 74°, the fauna was
- balanced between *N. incompta* (33 and 9 %) and *G. uvula* (8 and 34%).

### 5. Discussion

# Distribution pattern of living planktonic foraminifera at the Barents Sea Opening

In late summer 2014 the hydrology at the BSO was characterised, from 69.8°N to 74.5°N, by a strong water stratification with a 30 m thick Chl-a enriched lens of NwCW overlapping northwards the NwAW (saltier and colder). Further North, a 230 well-mixed water column with characteristics of the NwAW occupied the Storfjordrenna trough where a coccolithophore bloom (Giraudeau et al., 2016) and the highest concentration of cyanobacteria were recorded in the upper water column. Despite these marked features the global pattern of planktonic foraminifera abundance did not correlate with any of the studied environmental parameters (Figure 5). These observations confirm the low influence of commonly imputed parameters such as temperature, salinity and primary production on PF density (Schiebel et al., 2017). In accordance with 235 Retailleau et al., (2018) conclusions, multiples indices however highlight the possible importance of water turbidity in PF abundance distribution. The highest densities of planktonic foraminifera occurred in the 0-20m upper water layer between 70.5 and 74.5°N, and very low abundances were recorded nearby the Norwegian and Spitzbergen coasts. The low abundances at the two ends of the studied transect could reflect planktonic foraminifera patchiness pattern of distribution (Meilland et al., 2019) or highlight the fact that waters under continental influences (nutrient-enriched, more turbid) likely 240 hamper the foraminiferal production. In line with this, the abrupt decrease in abundances from West to East (stations 2, to 6, to 1) may be ascribed to the decrease in depth of the Bjømøyrenna trough up to the Barents Sea shelf (from 1850 to 430 m),

as foraminifera are suspected to avoid neritic waters over continental shelves (Schmuker, 2000).

The remarkable point of our results is the dominance of *Globigerinita uvula* in the high-latitude (> 70°N) waters at the BSO. This species, described as a temperate to polar species (Schiebel and Hemleben, 2017), is known to occupy less than 2% of

245 the assemblages in marginal Arctic Seas based on material collected with a 63 μm plankton net mesh size (Volkmann, 2000). *Neogloboquadrina pachyderma* is considered the dominant species in polar regions, making up more than 90% of the total planktonic foraminifera assemblages (e.g. Schiebel et al., 2017). The high densities of *G. uvula* recorded at the BSO in 2014 seem to be inconsistent with the former statements but are consistent with a recent study reporting *G. uvula* as one of the

dominant species in southern high latitudes, South of the Polar Front (Meilland et al., 2017). A possible explanation to these 250 observations could be the warming experienced by the western Barents Sea (SST anomalies  $\approx +2^{\circ}$ C) and its increase in salinity (SSS anomalies  $\approx +0.3$ ) over the last decades (Dobrynin and Pohlmann, 2015). These hydrological changes impact the plankton dynamics and biogeography, with a northwards shift of the natural range of biological communities (Barton et al., 2016). Thus the species distribution of planktonic foraminifera could be affected by an eventual expansion of subpolar/temperate species towards high latitudes leading to phytoplankton composition changes, in response to sea temperature warming under global climate change. Our observations from the North Polar Region support the shift of 255 planktonic foraminifera assemblages to warmer conditions already asserted from North Atlantic (Jonkers et al., 2019) and from the southern Indian Ocean data (Meilland et al., 2017). However, a single observational dataset is the Barents Sea is not sufficient to robustly validate this assumption and a second hypothesis for the dominance of G. uvula in our sampling area could be a response to specific phytoplankton composition and ambient water conditions by pulsed reproduction events only 260 in summer conditions. This seasonal pattern is known to occur in polar regions for Turborotalita quinqueloba (Schiebel and Hemleben, 2017). In fact, this species is the second dominant one in our late summer 2014 samples. As observed in this study, T. quinqueloba is also known to display high concentrations in the Barents Sea and western Spitsbergen (Volkmann 2000) and to co-occur with the typically polar species Neogloboquadrina pachyderma in the high-latitude cold-water assemblages (Volkmann, 2000; Eynaud, 2011).

265 Discrepancy between the species-specific distribution patterns was observed in late summer 2014 at the BSO. The low abundances of *Neogloboquadrina pachyderma* and *Neogloboquadrina incompta* consistent over the studied area versus the patchy distribution and high densities of *Globigerinita uvula* and *Turborotalita quinqueloba*, suggest differences in the ecological strategy and behaviour between these two pairs of species. The patchy pattern of planktonic foraminifera distribution has been observed before (Boltovskoy, 1971; Siccha et al., 2012; Meilland et al., 2019) suggesting that high

270 densities are not exclusively constrained by the physical structure of the (sub-) surface layers.

Potential differences in diet preferences could explain the observed species distribution in late summer 2014 at the BSO. Both *G. uvula* and *T. quinqueloba* are supposed to follow food availability and primary production (Volkmann 2000, Schiebel and Hemleben 2017), but we did not observe any correlation between their distribution and Chl-*a* concentrations (Figure 5). In late summer 2014, *G. uvula* and *T. quinqueloba* showed high concentrations especially at station 8, located at the cross road of the Atlantic (NwAW) and Arctic waters flowing out of the Storfjordrenna (Figure 1), at the edge of the polar front (Oziel et al., 2017). For this particular location, the concentration of phytoplankton was relatively low and the phytoplankton community showed singular characteristics, in comparison to the southern part of the studied transect: fuco-flagelattes became dominant and diatom concentrations decreased. The fuco-flagelatte blooms (mainly *Phaeocystis pouchetii* in late summer 2014; Giraudeau et al., 2016) are well known to occur in the Barents Sea (Wassmann et al., 1990; Vaquer-280 Sunyer et al., 2013). Our hypothesis thus is that *G. uvula* and *T. quinqueloba* high densities reflect more food composition

(quality) than food concentrations (quantity). This also implies that satellite-derived chlorophyll concentrations, considered as indices for algal bloom, may not always be good indicators to perceive neither lateral extension nor intensity of foraminiferal production.

# Planktonic foraminifera protein concentration, potential marker of their metabolism

- 285 Proteins are the main component of zooplankton biomass (Corg) in all oceanographic regions, from the tropics to polar areas (e.g., Percy and Fife 1981; Donnelly et al., 1994; Kumar et al., 2013; Yun et al., 2015). Their role is essential to organisms' growth and their concentration and composition likely reflect the environment individuals grew in and how well they adjust to it. Based on previous studies, the protein concentration of PF can be used as a proxy of its biomass (Corg) and for a given test size (Schiebel and Movellan, 2012). However, in our study, 290 the SNP (size normalized protein content) of PF appears to decrease with higher latitude and with related decrease in Chl-a concentration and temperature, whereas the sizes of individuals picked for these analyses remain constant. This observation could suggest foraminifera metabolisms (i.e. ability to consume/degrade food and to grow) is decreasing along the South to North transect and is supported by the observation of lower metabolism for zooplankton with decreasing salinity and temperature in the Arctic (Alcaraz et al., 2010). If proteins are the main component of zooplankton biomass (Corg) they are 295 closely followed by lipids. Lipids in zooplankton organisms are very variable geographically, showing a latitudinal pattern with high percentages in polar areas and low percentages in warm tropical waters, but also seasonal features, with higher percentages in summer than in winter (Falk-Petersen et al., 1999; Mayzaud et al., 2011; Kumar et al., 2013). It is thus possible that a part of energy (biomass / Corg) of the PF collected along the South-to-North transect shifts from being stocked as protein in warmer waters to being stocked as lipids in colder waters. This strategy would allow foraminifera to resist the
- 300 cold to potentially overwinter. This hypothesis is supported by analogous observations made on different size fraction of zooplankton in the Southern Indian Ocean showing variability in protein and lipid percentages among the 80 to 200 μm populations (Harmelin-Vivien et al., 2019) and also by observations made on pteropods in the Arctic (Kattner et al., 1998; Phleger et al., 2001; Böer et al., 2005). The fact that higher SNP of foraminifera were observed where Chl-*a* is higher is compatible with the fact that polar organisms are supposed to rely on their protein catabolism when food is easily accessible
- 305 rather than on their lipid storage (Brockington & Clarke 2001). It has also been shown that a single organism in a cold environment is able to switch between predominantly protein or lipid catabolism across his life (Mayzaud, 1976). This suggests that individuals from a same species can display more or less proteins for a same biomass in different locations. With the reduction of PF protein concentration (and likely metabolism) going North, one could expect lower densities. However we observe no link between PF concentrations, which appeared to be species specific, and protein concentrations
- 310 (evolving similarly for all four species) suggesting a decoupling between individuals metabolism and densities.

# Discrepancy between upper water column and interface sediment samples

The PF species compositions recorded during the late summer 2014 in the water column and in surficial sediments are similar while species relative abundances are drastically different. Indeed, the living fauna (collected by plankton net)

- 315 displays large relative and absolute abundances of the two species *Globigerinita uvula* and *Turborotalita quinqueloba* whereas the fossil assemblages (found in core-tops) are largely dominated by *Neogloboquadrina pachyderma* or, at the southernmost station, co-dominated by *T. quinqueloba* and *N. pachyderma* (Figure 7 a). Affected by differential settling velocities (200 and 500 m.day<sup>-1</sup> in normal conditions) and test sizes of different species, the foraminiferal fluxes exported from the upper productive surface and reaching the sea bottom depend on direction and intensity of currents (Takahashi and
- 320 Bé, 1984). Lateral advection may smear shells over long distances > 25 km for *N. pachyderma* and > 50 km for *T. quinqueloba*, respectively (Von Gyldenfeldt et al., 2000). Lateral advection of shells is also strengthened by water stratification that increases resident time at the shear boundary between superposed water masses (Kuhnt at al. 2013). The BSO present a complex hydrography with the buoyant NwCW flowing northwards above the NwAW that is entering eastwards the Barents Sea when cold BSAW is creeping westwards. In such area, the PF settling velocities and extension of
- 325 lateral advection are completely unknown. Consequently, the sediment core records cannot match exactly with the place and/or the intensity of production (Von Gyldenfeldt et al., 2000; Jonkers et al., 2015). However, aware of the eventual lateral advection of shells, Pados and Spielhagen, (2014) concluded from a study through the dynamic Fram Strait that the distribution pattern obtained by plankton tows was clearly reflected on the sediment surface, and that the assemblage on the sediment surface can be used as an indicator for modern planktonic foraminiferal fauna. This suggests that the large
- discrepancy between upper water column and interface sediment samples collected at the BSO in late summer 2014 should be taken into consideration. As a sedimentation rate of  $1.3 \pm 0.6$  mm.yr<sup>-1</sup> has been recently measured in the Storfjordrenna outlet [76°N-17°E], close to our station 9 (Fossile et al., 2019), the upper core-top sediments (0-0.5 cm slice) may have recorded less than a decade. Staying aware about the eventual variability of the sedimentation rate along the studied transect and of the limitations of our dataset, we hypothesise that the sea surface-bottom differences in the foraminiferal assemblages
- 335 along the South-to-North transect at the BSO might reflect a community change within a short period of time (about a decade).

Furthermore, the analysis of sediment from the 5 core-tops collected during the MOCOSED cruise demonstrated important differences between the assemblages of fossil fauna, i.e. empty tests of individuals dead for a while, and recently settled tests
(likely coming from surface Spring/Summer production), i.e. Rose-Bengal stained tests bearing not yet decomposed cytoplasm. For example, at 71.3°N, the percentages of coloured *T. quinqueloba* and *G. uvula* are twice higher than the ones observed for the fossil faunas (Figure 7). At 72.9°N in the surficial sediment, *G. uvula* reaches up to 38% of the coloured assemblages (Figure 7 b) whereas it never exceeds 25% in the non-coloured ones (Figure 7 a). The large representation of the two species *G. uvula* and *T. quinqueloba* in the living fauna as well as in the recently settled shells but not in the fossil

faunas suggest that they likely present a seasonal character with a production period focussed in late summer as a response to environmental and trophic conditions. This is supported by previous studies in the Arctic where *T. quinqueloba* has been



found to dominate assemblages sampled in August (Carstens et al., 1997; Volkmann, 2000) but not in June/ early July (Pados and Spielhagen, 2014), and by sediment trap observations from the subpolar North Atlantic where T. quinqueloba reaches its maximum in autumn (Jonkers et al., 2010). The dominance of N. pachyderma in the fossil faunas collected at the

- 350 BSO and its low but constant presence in the coloured shells of surficial sediment and plankton tow sampled in late summer 2014 suggests that this species may demonstrate a more sustainable behaviour with a regular production throughout the year. This hypothesis is supported by a recent study showing that abundances and distribution of the species N. pachyderma are not significantly perturbed by seasonal seawater temperature, productivity or salinity variations occurring in the Arctic (Greco et al., 2019). Thus it makes sense that N. pachyderma production appears to be yearly sustained and constant in the
- 355 area whereas other species clearly respond to a local seasonal signal.

# 6. Conclusion

The sampling and analytic approaches deployed during the MOCOSED14 cruise and combining the use of plankton net, core-top, molecular biology (protein measurement), environmental parameters and phytoplankton characterisation provides us with a unique dataset to better constrain the distribution of planktonic foraminifera within the highly complex studied area of the western Barents Sea.

360

The observed abundances of PF in the studied area are high and the lower densities were recorded nearby the Norwegian and Spitzbergen coasts. These observations highlight the fact that waters under continental influences (nutrient-enriched, more turbid) are rather inhospitable for PF production. The PF species composition observed at the BSO happens to be diverse, with more than 10 different species including Globigerinita uvula (45%), Turborotalita quinqueloba (26.2%),

- Neogloboquadrina incompta (15%) and Neogloboquadrina pachyderma (8.9%). The two species G. uvula and T. 365 quinqueloba clearly dominate the living (water sample) population and display high patchy abundances suggesting they occur in late summer in response to physico-chemical conditions and related specific primary productivity. The dominance of G. uvula in water samples could also be a signal of the temperature increase experienced over the last decades by the Barents Sea and the North Atlantic Ocean under global climate change. Further sampling in the area is thus needed to
- explore this hypothesis. The species N. pachyderma and N. incompta show low densities but a continuous distribution 370 pattern in the water samples. They also dominate the core-top assemblages suggesting that both species present a more consistent production over the course of the year.

Unlike their species specific abundances pattern of distribution, size-normalized protein concentrations of all four major species decrease with the increasing latitude (and a decrease in temperature and Chl-a concentration). This observation leads

us to hypothesise that 1) PF densities and metabolism are decouple and 2) foraminifera metabolism is the North of the 375 studied region is lower than in the South. It opens the following question: Can individuals of the same species balance the ratio between their protein and lipid concentrations (known to be the major components of zooplankton C<sub>org</sub>) in order to adapt to their environmental conditions (e.g. temperature change)? Further analyses on planktonic foraminifera lipid concentration and composition are thus needed and would help us to better understand the metabolism of these organisms

380 and their fate in a context of climate change.

# Data availability

Data will be made available on request to the main author until their online publication on PANGAEA (https://pangaea.de/).

# **Author contributions**

JM and HH designed the study. JM, VH, ID and JS generated the data and carried out the analyses. TG provided access to the MOCOSED 2014 cruise. All authors contributed to writing the manuscript.

### **Competing interests**

The authors declare that they have no conflict of interest.

# Acknowledgements

The authors are thankful to the crew and captain of R/V *Pourquoi pas* ?, as well as the scientific participants from the 390 SHOM (French Hydrographic Office) for their support in sampling during the cruise "MOCOSED 2014". We fully acknowledge the efficient technical help provided by Sophie Sanchez at LPG-BIAF laboratory. Grant was kindly provided to Vivien Hulot by the SHOM to study phytoplankton and PF communities.

# References

Alcaraz, M., Almeda, R., Calbet, A., Saiz, E., Duarte, C. M., Lasternas, S., Agustí, S., Santiago, S., Movilla, J., Alonso, A.:
The role of arctic zooplankton in biogeochemical cycles: respiration and excretion of ammonia and phosphate during summer. Polar Biology, 33(12), 1719-1731, 2010.

AMAP Assessment 2018: Arctic Ocean Acidification. Arctic Monitoring and Assessment Programme (AMAP). Tromsø, Norway: Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway (<u>www.amap.no</u>).

Årthun, M., Eldevik, T., Smedsrud, L. H., Skagseth, Ø., Ingvaldsen R. B.: Quantifying the Influence of Atlantic Heat on
Barents Sea Ice Variability and Retreat. Journal of Climate, 25, 4736-4743, https://doi.org/10.1175/JCLI-D-11-00466.1, 2012.

Barton, A.D., Irwin, A.J., Finkel, Z.V., Stock, C.A.: Anthropogenic climate change drives shift and shuffle in North Atlantic phytoplankton communities. PNAS, 113 (11) 2964-2969, https://doi.org/10.1073/pnas.1519080113, 2016.

Beaugrand, G., McQuatters-Gollop, A., Edwards, M., Goberville, E.: Long-term responses of North Atlantic calcifying 405 plankton to climate change. Nature Climate Change, 3(3), 263, 2013.

Beer, C.J., Schiebel, R., Wilson, P.A.: Technical note: on methodologies for determining the size-normalised weight of planktic foraminifera. Biogeosciences, http://dx.doi.org/10.5194/bg-7-2193-2010, 2010.

Böer, M., Gannefors, C., Kattner, G., Graeve, M., Hop, H., Falk-Petersen, S.: The Arctic pteropod *Clione limacina*: seasonal lipid dynamics and life-strategy. Marine biology, 147(3), 707-717, 2005.

410 Bollmann, J., Quinn, P.S., Vela, M., Brabec, B., Brechner, S., Cortés, M.Y., Hilbrecht, H., Schmidt, D.N., Schiebel, R., Thierstein, H.R.: Automated particle analysis: calcareous microfossils. Francus, P. (Ed.), Image Analysis, Sediments and Paleoenvironments. Springer, pp. 229–252 chap. 12, 2004

Boltovskoy, E.: Patchiness in the distribution of planktonic foraminifera. In: A. Farinacci (Ed.), Proceedings of the Second Planktonic Conference, Rome 1970. Edizioni Technoscienza, Roma, pp. 107–116, 1971.

415 Brockington, S., Clarke, A.: The relative influence of temperature and food on the metabolism of a marine invertebrate. Journal of Experimental Marine Biology and Ecology, 258(1), 87-99, 2001.

Burenkov, V. I., Kopelevich, O. V., Rat'kova, T. N., Sheberstov, S. V.: Satellite observations of the coccolithophorid bloom in the Barents Sea. Oceanology, 51 (5), 766–774, https://doi.org/10.1134/s0001437011050043, 2011.

Carstens, J., Hebbeln, D., Wefer, G.: Distribution of planktic foraminifera at the ice margin in the Arctic (Fram Strait). 420 Marine Micropaleontology, 29(3-4), 257-269, 1997.

Clayton, C.R.I., Abbireddy, C.O.R., Schiebel, R.: A method of estimating the form of coarse particulates. Geotechnique, http://dx.doi.org/10.1680/geot.2007.00195, 2009.

Dai, A., Luo, D, Song, M, and Liu, J.: 'Arctic amplification is caused by sea-ice loss under increasing CO<sub>2</sub>', Nature Communications. Springer US, 10(1), pp. 1–13. https://doi.org/10.1038/s41467-018-07954-9, 2019.

425 Dalpadado, P., Ingvaldsen, R. B., Stige, L. C., Bogstad, B., Knutsen, T., Ottersen, G., Ellertsen, B.: Climate effects on Barents Sea ecosystem dynamics. ICES Journal of Marine Science, 69, 1303–1316, https://doi.org/10.1093/icesjms/fss063, 2012.

Dobrynin, M., Pohlmann, T.: Anomalous hydrographic conditions in the western Barents Sea observed in March 2014. Cont. Shelf Res. 111, 69–82. doi:10.1016/j.csr.2015.10.020, 2015.

430 Donnelly, J., Torres, J. J., Hopkins, T. L., Lancraft, T. M.: Chemical composition of Antarctic zooplankton during austral fall and winter. Polar Biology, 14(3), 171-183, 1994.

Dylmer, C.V., Giraudeau, J., Eynaud, F., Husum, K., De Vernal, A.: Northward advection of Atlantic water in the eastern Nordic Seas over the last 3000 yr. Clim. Past, 9, 1505–1518, <u>https://doi.org/10.5194/cp-9-1505-2013</u>, 2013.

Eynaud F.: Planktonic foraminifera in the Arctic: potentials and issues regarding modern and quaternary populations. IOP 435 Conf. Series: Earth and Environmental Science, 14, O12005. doi:10.1088/1755-1315/14/1/012005, 2011.

Falk-Petersen, S., Sargent, J. R., Lønne, O. J., Timofeev, S.: Functional biodiversity of lipids in Antarctic zooplankton: Calanoides acutus, Calanus propinquus, Thysanoessa macrura and Euphausia crystallorophias. Polar Biology, 21(1), 37-47, 1999.

Fossheim, M., Primicerio, R., Johannesen, E., Ingvaldsen, R. B., Aschan, M.M., Dolgov, A. V.: Recent warming leads to a 440 rapid borealization of fish communities in Climate Change, 5, the Arctic. Nature 673-677. https://doi.org/10.1038/nclimate2647, 2015.

Fossile, E., Nardelli, M.P., Jouini, A., Lansard, B., Pusceddu, A., Moccia, D., Michel, E., Péron, O., Howa, H., Mojtahid,
M.: Benthic foraminifera as tracers of brine production in Storfjorden sea ice factory, Biogeosciences Discuss.,
https://doi.org/10.5194/bg-2019-405, in review, 2019.

445 Giraudeau, J., Hulot, V., Hanquiez, V., Devaux, L., Howa, H., Garlan, T.: A survey of the summer coccolithophore community in the western Barents Sea. Journal of Marine Systems, 158, 93-105, https://doi.org/10.1016/j.jmarsys.2016.02.012, 2016.

Greco, M., Jonkers, L., Kretschmer, K., Bijma, J., Kucera, M.: Depth habitat of the planktonic foraminifera *Neogloboquadrina pachyderma* in the northern high latitudes explained by sea-ice and chlorophyll
 450 concentrations. Biogeosciences, 16(17), 3425-3437, 2019.

Harmelin-Vivien, M., Bănaru, D., Dromard, C. R., Ourgaud, M., Carlotti, F.: Biochemical composition and energy content of size-fractionated zooplankton east of the Kerguelen Islands. Polar Biology, 42(3), 603-617, 2019.

Holding, J.M., Duarte, C.M., Sanz-Martín, M., Mesa, E., Arrieta, J.M., Chierici, M., Hendriks, I.E., García-Corral, L.S., Regaudie-de-Gioux, A., Delgado, A., Reigstad, M., Wassmann, P., Agustí, S.: Temperature dependence of CO<sub>2</sub>-enhanced primary production in the European Arctic Ocean. Nat. Clim. Change 5, 1079–1082. doi:10.1038/nclimate2768, 2015.

455

Jonkers, L., Brummer, G.-J. A., Peeters, F. J. C., van Aken, H. M., de Jong, M. F.: Seasonal stratification, shell flux, and oxygen isotope dynamics of left-coiling *N. pachyderma* and *T. quinqueloba* in the western subpolar North Atlantic. Paleoceanography, 25, PA2204, https://doi.org/10.1029/2009PA001849, 2010.

Jonkers, L., van Heuven, S., Zahn, R., Peeters, F.J.C.: Seasonal patterns of shell flux, d18O and d13C of small and large *N*. 460 *pachyderma* (s) and *G. bulloides* in the subpolar North Atlantic. Paleoceanography, 28(1), 164-174, https://doi.org/10.1002/palo.20018, 2013.

Jonkers, L., Reynolds, C. E., Richey, J., Hall, I. R.: Lunar periodicity in the shell flux of planktonic foraminifera in the Gulf of Mexico, Biogeosciences, 12, 3061–3070, https://doi.org/10.5194/bg-12-3061-2015, 2015.

Jonkers, L., Hillebrand, H., Kucera, M.: Global change drives modern plankton communities away from the pre-industrial state. Nature, 570(7761), 372–375. https://doi.org/10.1038/s41586-019-1230-3, 2019.

Kattner, G., Hagen, W., Graeve, M., Albers, C.: Exceptional lipids and fatty acids in the pteropod *Clione limacina* (Gastropoda) from both polar oceans. Marine Chemistry, 61(3-4), 219-228, 1998.

Kucera M., Weinelt M., Kiefer T., Pflaumann U., Hayes A., Weinelt M., Chen M.-T., Mix A.C., Barrows T.T., Cortijo E., Duprat J., Juggins S., Waelbroeck C.: Reconstruction of sea-surface temperatures from assemblages of planktonic

470 foraminifera: Multi-technique approach based on geographically constrained calibration data sets and its application to glacial Atlantic and Pacific Oceans Quat. Sci. Rev. 24 95198, 2005.

Kuhnt T., Howa H., Schmidt S., Marié L., Schiebel R.: Flux dynamics of planktic foraminifer tests at a hemipelagic site of the inner Bay of Biscay (Northeast Atlantic margin). Journal of Marine Systems, 109-110, S169-S181, doi:10.1016/j.jmarsys.2011.11.026, 2013.

475 Kumar, M. A., Padmavati, G., Anandavelu, I.: Biochemical composition and calorific value of zooplankton from the coastal waters of South Andaman. Proceedings of the International Academy of Ecology and Environmental Sciences, 3(3), 278-287, 2013.

Loeng, H.: Features of the physical oceanographic conditions of the Barents Sea. Polar Res. 10 (1), 5-18, 1991.

Mackey, M.D., Mackey, D.J., Higgins, H.W., Wright, S.W.: CHEMTAX — a program for estimating class abundances from 480 chemical markers: application to HPLC measurement of phytoplankton. Mar. Ecol. Prog. Ser. 144, 265–283, 1996.

Mayzaud, P., Lacombre, S., Boutoute, M.: Seasonal and growth stage changes in lipid and fatty acid composition in the multigeneration copepod *Drepanopus pectinatus* from Iles Kerguelen. Antarctic Science, 23(1), 3-17, 2011.

Mayzaud, P.: Respiration and nitrogen excretion of zooplankton. IV. The influence of starvation on the metabolism and the biochemical composition of some species. Marine Biology, 37(1), 47-58, 1976.

485 Meilland, J., Howa, H., LoMonaco, C., Schiebel, R.: Individual planktic foraminifer protein-biomass affected by trophic conditions in the Southwest Indian Ocean, 30°S–60°S. Mar. Micropaleontol. 124, 63–74. http://dx.doi.org/10.1016/j.marmicro.2016.02.004, 2016.

Meilland, J., Schiebel, R., Monaco, C. L., Sanchez, S., Howa, H.: Abundances and test weights of living planktic foraminifers across the Southwest Indian Ocean: Implications for carbon fluxes. Deep Sea Research Part I: Oceanographic Research Papers, 131, 27-40, 2018.

490

Meilland, J., Siccha, M., Weinkauf, M. F. G., Jonkers, L., Morard, R., Baranowski, U., Baumeister, A., Bertlich, J., Brummer, G. J., Debray, P., Fritz-Endres, T., Groeneveld, J., Magerl, L., Munz, P., Rillo, M. C., Schmidt, C., Takagi, H., Theara, G., and Kucera, M.: Highly replicated sampling reveals no diurnal vertical migration but stable species-specific vertical habitats in planktonic foraminifera. J. Plank. Res., 41, 127–141, 2019.

495 Neukermans, G., Oziel, L., Babin, M.: Increased intrusion of warming Atlantic water leads to rapid expansion of temperate phytoplankton in the Arctic. Global Change Biology, Wiley, 24 (6), pp.2545-2553. doi: 10.1111/gcb.14075, 2018.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Wagner, H.: Vegan: Community Ecology Package [available on internet at http://www. CRAN.R-project.org/package=vegan], 2013.

500 Oziel, L., Neukermans, G., Ardyna, M., Lancelot, C., Tison, J-L., Wassmann, P., Sirven, J., Ruiz-Pino, D., Gascard, J-C.: Role for Atlantic inflows and sea ice loss on shifting phytoplankton blooms in the Barents Sea, J. Geophys. Res. Oceans, 122, 5121–5139, doi:10.1002/2016JC012582, 2017.

Pados, T., Spielhagen, R.F.: Species distribution and depth habitat of recent planktic foraminifera in Fram Strait, Arctic Ocean. Polar Res. 33. doi:10.3402/polar.v33.22483, 2014.

505 Percy, J. A., Fife, F. J.: The biochemical composition and energy content of Arctic marine macrozooplankton. Arctic, 307-313, 1981.

Phleger, C. F., Nelson, M. M., Mooney, B. D., Nichols, P. D.: Interannual variations in the lipids of the Antarctic pteropods Clione limacina and Cliopyramidata. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 128(3), 553-564, 2001.

510 Retailleau S., Eynaud F., Mary Y., Schiebel R., Howa H.: Canyon heads and river plumes: how might they influence neritic planktonic foraminifera communities in the SE Bay of Biscay? Journal of Foraminiferal Research, 42 (3), 257-269, 2012.

Schiebel, R., Hemleben, C.: Planktic Foraminifers in the Modern Ocean. 357 pp, Ed. Springer-Verlag GmbH, Berlin. doi : 10.1007/978-3-662-50297-6, 2017.

Schiebel, R., Spielhagen, R.F., Garnier, J., Hagemann, J., Howa, H., Jentzen, A., Martínez-Garcia, A., Meilland, J., Michel,
E., Repschläger, J., Salter, I., Yamasakig, M., Haug, G.: Modern planktic foraminifers in the high-latitude ocean. Marine Micropaleontology, 136, 1–13, 2017.

Schiebel, R., Movellan, A.: First-order estimate of the planktic foraminifer biomass in the modern ocean. Earth System Science Data, 4(1), 75-89, 2012.

Schlitzer, R.: Ocean data view, odv. awi. de, edited, 2014.

520 Schmuker, B.: The influence of shelf vicinity on the distribution of planktic foraminifera south of Puerto Rico. Marine Geology, 166(1-4), 125-143, 2000.

Schönfeld, J., Elena Golikova, E., Korsun, S., Spezzaferri, S.: The Helgoland Experiment - assessing the influence of methodologies on Recent benthic foraminiferal assemblage composition. Journal of Micropalaeontology 32(2):161-182. DOI: 10.1144/jmpaleo2012-022, 2013.

525 Shepherd T.G.: Effects of a warming Arctic. Science 02 Sep 2016, vol. 353, Issue 6303, pp. 989-990. DOI: 10.1126/science.aag2349, 2016.

Siccha, M., Kucera, M.: ForCenS, a curated database of planktonic foraminifera census counts in marine surface sediment samples. Sci. Data, 4, 170109, 2017.

Siccha M., Schiebel R., Schmidt S., Howa H.: Short-term and small scale variability in planktic foraminifera test flux in the 530 Bay of Biscay. Deep-Sea Research I, 64, 146-156, doi:10.1016/j.dsr.2012.02.004, 2012.

Skagseth, Ø., Furevik, T., Ingvaldsen, R., Loeng, H., Mork, K. A., Orvik, K. A., Ozhigin, V.: Volume and heat transports to the Arctic Ocean via the Norwegian and Barents Seas. In Arctic–Subarctic Ocean Fluxes (pp. 45-64). Springer, Dordrecht, 2008.

Smedsrud, L. H., Esau, I., Ingvaldsen, R. B., Eldevik, T., Haugan, P. M., Li, C., Lien, V.S., Olsen, A., Omar, A.M., Otterå,
O.H., Risebrobakken, B., Sandø, A.B., Semenov, V.A., Sorokina, S.A.: The role of the Barents Sea in the Arctic climate system. Reviews of Geophysics, 51(3), 415-449, 2013.

Smyth, T.J., Tyrell, T., Tarrant, B.: Time series of coccolithophore activity in the Barents Sea, from twenty years of satellite imagery. Geophysical Research Letters, 31, L11302, 2004.

Takahashi, K., Bé, A. W. H.: Planktonic foraminifera: factors controlling sinking speeds, Deep-Sea Res., 31, 1477–1500, doi:10.1016/0198-0149(84)90083-9, 1984.

Troupin, C., Barth, A., Sirjacobs, D., Ouberdous, M., Brankart, J.- M., Brasseur, P., Rixen, M.: Generation of analysis and consistent error fields using the Data Interpolating Variational Analysis (DIVA). Ocean Modelling, 52, 90-101, 2012.

Van Heukelem, L., Thomas, C.S.: Computer-assisted high performance liquid chromatography method development with applications to the isolation and analysis of phytoplankton pigments. J. Chromatogr. 910, 31–49, 2001.

545 Vaquer-Sunyer, R., Duarte, C. M., Holding, J., Regaudie-de-Gioux, A., García-Corral, L. S., Reigstad, M., Wassmann, P.: Seasonal patterns in Arctic planktonic metabolism (Fram Strait – Svalbard region), Biogeosciences, 10, 1451–1469, https://doi.org/10.5194/bg-10-1451-2013, 2013.

Volkmann, R.: Planktic foraminifers in the outer Laptev Sea and the Fram Strait modern distribution and ecology. J. Foraminifer. Res. 30, 157–176, 2000.

550 Von Gyldenfeldt, A.B., Carstens, J., Meincke, J.: Estimation of the catchment area of a sediment trap by means of current meters and foraminiferal tests. Deep-Sea Res II 47:1701–1717, 2000.

Wassmann P, Vernet M, Mitchell BG, Reyh F. Mass sedimentation of Phaeocystis pouchettii in the Barents Sea. Mar Ecol, Prog Ser; 66:183–95, 1990.

Yun, M. S., Lee, D. B., Kim, B. K., Kang, J. J., Lee, J. H., Yang, E. J., Park, W. G., Chung, K. H., Lee, S. H.: Comparison of
phytoplankton macromolecular compositions and zooplankton proximate compositions in the northern Chukchi Sea. Deep
Sea Research Part II: Topical Studies in Oceanography, 120, 82-90, 2015.

**Table 1:** Location (Latitude and Longitude), sampling date and water depth, of the 9 MultiNet and 5 Multitube stations, incremented from South to North (stations 1 and 2 being positioned aside the main transect mid-point #6). Stations where phytoplankton analyses were performed are also indicated.

					Sample collection		
Station	Latitude (°N)	Longitude (°E)	Date	Depth of sampling site (m)	Multinet	Phytoplankton	Multitube
3	69.845	13.879	22.08.14	2675	Х	Х	Х
4	71.308	13.942	23.08.14	1940	х	Х	х
5	72.138	14.098	23.08.14	1253	х	Х	
1	72.893	11.762	16.08.14	1839	х		
6	72.897	14.207	24.08.14	990	х	Х	х
2	72.912	19.487	17.08.14	430	х		
7	73.736	14.376	24.08.14	1320	Х	Х	
8	74.537	14.508	25.08.14	1978	х	Х	х
9	75.602	14.705	26.08.14	445	х	х	Х

**Table 2:** Descriptive statistics [minimum, maximum, average, and median values in  $\mu$ m] of the minimum (i.e. the shortest) diameter of all the 272 BCA measured specimens, per species.

Species	Minimum	Average	Median	Maximum	
All	65.12	158.45	160.2	315.18	
G. uvula	65.5	108.8	108.8	160.8	
T. quinqueloba	65.12	166.13	166.56	249.18	
N. incompta	75.08	191.29	196.66	280.46	
N. pachyderma	100.7	184.3	180.7	315.18	

Test size distribution parameters in µm



**Figure 1:** Sampling map of the MOCOSED cruise in the western Barents Sea with schematic surface circulation (red arrows = Atlantic Water; blue arrows = Arctic Water; green arrow = Norwegian Coastal Current; adapted from Oziel et al., 2017). Little green dots display the 32 CTD/Niskin stations along the South-to-North transect; large blue circles show the location of the 9 MultiNet stations; medium red circles underline the 5 multicore stations.



**Figure 2:** South-North, 0-100 m deep sections across the Barents Sea Opening, compiled from data of the 32 CTD casts: (a) Temperature (°C); (b) Salinity; and (c) Chl-*a* concentrations (mg.m<sup>-3</sup>). Vertical dashed lines and associated numbers correspond to the location of the MultiNet hauls. The lower right panel (d) displays pigment concentrations translated into the relative abundances of the major phytoplankton species at 7 sampling stations along the transect (Giraudeau et al., 2016).



**Figure 3:** Distribution of planktonic foraminifera total abundances (ind.m<sup>-3</sup>) in the 0-100 m depth section across the Barents Sea Opening (upper panel) and in the West-East transect (stations 1, 6 and 2; lower panel). Station names are indicated above vertical alignments of 5 dots representing the middle points of the 5 net sampling intervals.



```
590
```

**Figure 4:** Distribution of the four major species abundances (ind.m<sup>-3</sup>) in the 0-100 m depth section across the Barents Sea Opening. (a) For *Globigerinita uvula* with species abundances (z-axes) going from 0 to 250 ind.m<sup>-3</sup>; (b) for *Turborotalita quinqueloba* (c) *Neogloboquadrina incompta* (d) and *Neogloboquadrina pachyderma* values goes from 0 to 80 ind.m<sup>-3</sup>. Station names are indicated above vertical alignments of 5 dots representing the middle points of the 5 net sampling intervals.



**Figure 5:** Standard nonmetric multidimensional scaling ordination analysis (NMDS) of planktonic foraminifera species distribution (red dots) with temperature, salinity, Chl-*a* and station location as factor (blue arrows).





**Figure 6:** Boxplot of the size normalized protein biomass (SNP,  $\mu g.\mu m^{-1}$ ), along the South-to-North transect (from station 3 = 69.8°N to station 9 = 75.6°N) (a) for *Globigerinita uvula* (b), *Turborotalita quinqueloba* (c), *Neogloboquadrina incompta* (d) and *Neogloboquadrina pachyderma*. Blue dots highlight the data dispersion. Potential outliers were removed such as data for *N. pachyderma* at station 7 as analyses were only run on one individual. Thick lines indicate median, boxes extend to interquartile range (IQR) and whiskers indicate 1.5\*IQR.



Figure 7: Relative species occurrence (% of the total fauna) of planktonic foraminifera found in the upper 0.5 cm core-top sediment (histograms) and total concentration of individuals per cm<sup>3</sup> of dry sediment (logarithmic Y axis on the right): (a)
610 colourless empty tests and (b) individuals bearing a Rose Bengal coloured cytoplasm.