

# Rare Earth Elements in oyster shells: provenance discrimination and potential vital effects

Vincent Mouchi<sup>1</sup>, Camille Godbillot<sup>1</sup>, Vianney Forest<sup>2</sup>, Alexey Ulianov<sup>3</sup>, Franck Lartaud<sup>4</sup>, Marc de Rafélis<sup>5</sup>, Laurent Emmanuel<sup>1</sup>, Eric P. Verrecchia<sup>6</sup>

5 <sup>1</sup> Sorbonne Université, CNRS-INSU, Institut des Sciences de la Terre Paris, ISTeP, F-75005 Paris, France

<sup>2</sup> INRAP-Occitanie, UMR 5068, TRACES, Toulouse, France

<sup>3</sup> University of Lausanne, Institut des Sciences de la Terre, CH-1015, Lausanne, Switzerland

<sup>4</sup> Sorbonne Université, CNRS, Laboratoire d'Ecogéochimie des Environnements Benthiques, LECOB, F-66650, Banyuls, France

10 <sup>5</sup> Géosciences Environnement Toulouse, CNRS, IRD, Université Paul Sabatier Toulouse 3, 14 Avenue Edouard Belin, 31400 Toulouse, France

<sup>6</sup> University of Lausanne, Institut des Dynamiques de la Surface Terrestre, CH-1015, Lausanne, Switzerland

*Correspondence to:* Vincent Mouchi (vmouchi@gmail.com)

**Abstract.** Rare Earth Elements (REE) and yttrium in seawater originate from atmospheric fallout, continental weathering, and transport from rivers, as well as hydrothermal activity. Previous studies reported the use of REE and Y measurements in biogenic carbonates as a means to reconstruct these surface processes in ancient times. As coastal seawater REE and Y concentrations partially reflect those of nearby rivers, it may be possible to obtain a regional fingerprint of these concentrations from bivalve shells for seafood traceability and environmental monitoring studies. Here, we present a dataset of 297 measurements of REE and Y abundances by LA-ICP-MS from two species (*Crassostrea gigas* and *Ostrea edulis*). We measured a total of 49 oyster specimens from six locations in France (Atlantic Ocean and Mediterranean Sea). Our study reports that there is no significant difference in concentrations from shell parts corresponding to winter and summer periods for both species. Moreover, interspecific vital effects are reported from specimens from both species and from the same locality. REE and Y profiles and t-distributed Stochastic Neighbour Embedding processing (t-SNE; a discriminant statistical method) indicate that REE and Y measurements from *C. gigas* shells can be discriminated from one locality to another, but this is not the case for *O. edulis*, which presents very similar concentrations in all studied localities. Therefore, provenance studies using bivalve shells based on REE and Y have to be first tested for the species. Other methods have to be investigated to be able to find the provenance of some species such as *O. edulis*.

## 1 Introduction

Rare Earth Elements (REE) are a group of 15 elements (La to Lu) with similar electronic configuration of the atoms, similar properties and chemical behavior (Elderfield, 1988), with the exceptions of Ce and Eu, which present multiple oxidation states. The main sources of REE in seawater are atmospheric fallout (Elderfield & Greaves, 1982; De Baar et al., 1983) and riverine input through continental weathering (Goldstein et al., 1984; Frost et al., 1986), as well as hydrothermal activity (Olivarez &

Owen, 1991). In addition to these various sources, the concentrations of REE in seawater are impacted by adsorption processes of REE to mineral surfaces and complexation with organic ligands, especially near the ocean surface (Sholkovitz et al., 1994; Schijf et al., 2015).

Reconstruction of REE compositions of seawater is generally used to provide information on past continental weathering, tectonic activity, and water mass circulation (Greaves et al., 1991; Censi et al., 2004; Haley et al., 2005; Piper & Bau, 2013). Moreover, specific elemental ratios, such as Y/Ho, have been investigated as potential provenance proxies. Indeed, although the average Y/Ho value is equivalent to that of chondritic meteorites and mid-ocean-ridge basalts (MORB; Jochum et al., 1986; Taylor & McLennan, 1988), it has been shown that Y/Ho fractionates in sediment particles, not only in seawater with depth, but also probably in waterbodies from watersheds (rivers and estuaries), whose composition has been modified depending on the weathered continental rocks (Bau et al., 1995; Nozaki et al., 1997; Prajith et al., 2015). The Y/Ho ratios in estuaries could therefore exhibit different values according to the regional inputs related to the mineralogical variability in continental covers. It has been demonstrated that the seawater composition in REE and yttrium is at least partially recorded in carbonate materials, such as ooids (Li et al., 2019), brachiopod shells (Zaky et al., 2015, 2016), foraminifera tests (Osborne et al., 2017), and coral skeletons (Sholkovitz & Shen, 1995). In addition, the anthropogenic REE contamination of the Rhine River (Germany) has been demonstrated by the shell composition of freshwater mussels (Merschel & Bau, 2015). The reconstruction of the REE and Y fingerprints in a coastal environment from mollusk shells could therefore be useful, not only to monitor potential contaminations from anthropic activities (Le Goff et al., 2019), but also as a provenance proxy for quality control of cultured organisms prior to human consumption (Bennion et al., 2019; Morrison et al., 2019). Indeed, public interest is rising regarding the origin of food due to various reasons including decreased confidence in the quality and safety of remote food supply (Kelly et al., 2005; Gopi et al., 2019). The geographic origin of seashells can also be of interest in archaeology, since mollusk shells can be unearthed sometimes very far from the nearest shoreline (Bardot-Cambot, 2014), in order to rebuild historic trade routes. Seafood traceability is therefore important for both modern and archaeological contexts. However, ‘vital effects’ (compositional shifts between inorganic and biogenic carbonate due to metabolic activity; Urey et al., 1951) have been reported to alter this regional fingerprint of REE between species (Akagi et al., 2004); therefore, a feasibility study needs to be conducted on mollusks before any extended seafood traceability perspective is considered.

To this end, REE and Y measurements have been performed by laser-ablation inductively-coupled mass spectrometry (LA-ICP-MS) on modern and archaeological oyster shells from various French localities. The aim of this study is to assess the interspecific effects using specimens from two edible species: the flat oyster *Ostrea edulis* (Linnaeus, 1758), commonly found in antique sites but still found on modern shores, and the cupped oyster *Crassostrea gigas* (Thunberg, 1793), which first appeared on the French coastline during the 20<sup>th</sup> century. Multiple measurements were performed on each of the specimens to evaluate the intraspecific variations, in particular regarding the potential seasonal fluctuations of rainfall intensity, weathering, and rate of river flows. A recent statistical method, the t-SNE, standing for “t-distributed Stochastic Neighbour Embedding” (van der Maaten & Hinton, 2008), is introduced as an attempt to discriminate the nine oyster groups of the study, in order to highlight significant interspecific and inter-regional differences between REE incorporations.

## 2. Material and methods

### 2.1 Modern-day settings and specimens

#### 2.1.1 Baie des Veys (Normandy)

70 The G fosse area, in Baie des Veys (Normandy, France), is currently used as a commercial oyster farm location. This open sea area is characterized by a semidiurnal tidal range of 8 m from the English Channel and an overall siltation due to weaker ebb than flow, inducing a poor resuspension of sediment particles (Le Gall, 1970). The Baie des Veys is recharged in freshwater by the Isigny Channel, formed by the Vire and Aure rivers, and the Carentan Channel, constituted of the Douve and the Taute rivers, which drain a large part of the Bessin plain and the Cotentin. The watershed comprises limestone as well as basalt, 75 acidic and alkaline metavolcanic rock and diorite (Baize et al., 1997). Respective flows of the Isigny and Carentan Channels are  $19 \text{ m}^3 \text{ s}^{-1}$  and  $33 \text{ m}^3 \text{ s}^{-1}$ , these rates having no significant impact on the salinity of the shoreline (Sylvand, 1995). Oyster specimens from this locality were gathered during a previous rearing experiment conducted between 2005 and 2006 (Lartaud et al., 2010a, 2010b; Mouchi et al., 2013). Although these specimens have been transplanted in several localities during their lives (for details, see Lartaud et al., 2010a), the part of the shells analyzed in the present study is restricted to that 80 corresponding to their Baie des Veys stay. This period is recognized in the shells owing to *in vivo* chemical labeling performed during the rearing experiment (Huyghe et al., 2019). Both *Crassostrea gigas* (n=5) and *Ostrea edulis* (n=5) specimens from this locality and having shared the same breeding location, have been considered (Figure 1, Table 1, Appendix A). *Crassostrea gigas* was recently renamed *Magallana gigas* by Salvi and Mariottini (2016); however, as this genus change is still debated (Bayne et al., 2017) and *C. gigas* is the most commonly found occurrence in the literature, this paper refers to this species by 85 using its original genus name. As modern specimens, these are referred to as Mod\_BDV\_Cgig and Mod\_BDV\_Oedu groups in this paper for *C. gigas* and *O. edulis*, respectively.

#### 2.1.2 Marennes-Ol ron bay (Charente-Maritime)

The Marennes-Ol ron bay is located on the Atlantic Ocean coast of France. It is bordered by the Ol ron island on the western side, and opens to the Atlantic on its northern and southern borders. Its surface area is approximately 180 km<sup>2</sup>. Atlantic seawater 90 runs through the bay from North to South (Dechambenoy et al., 1977). In the area, the tidal range is 5 m with a semi-diurnal rhythm. The large watershed covers 10,000 km<sup>2</sup> of land and comprises Cenozoic river deposits, limestones, and clays (Bourgueil et al., 1968, 1972; Platel et al., 1977, 1978; Bambier et al., 1982; Hantzpergue et al., 1984; Mourier et al., 1989). The bay is recharged in freshwater by two rivers, the Charente (North side), with a  $36 \text{ m}^3 \cdot \text{s}^{-1}$  flow, and the Seudre (South side) whose flow is 30 times less important (Soletchnik et al., 1998). 95 The bay hosts 30 km<sup>2</sup> of aquaculture domains, amongst these oysters. This site was also used during a rearing experiment (Lartaud et al., 2010a, 2010b): both *C. gigas* (n=3) and *O. edulis* (n=3) shells parts analyzed in the present study refer to the shell portions corresponding to their stay at this site, highlighted by *in vivo* labels. These two groups are referred to as Mod\_MO\_Cgig and Mod\_MO\_Oedu for *C. gigas* and *O. edulis* specimens, respectively.

### 100 **2.1.3 Tès (Arcachon basin)**

The Arcachon basin is a lagoon of 156 km<sup>2</sup> on the French Atlantic coastline. The area is subdivided into a subtidal zone and an intertidal zone with a semidiurnal tidal range of 3 m, where the studied oysters grew. Freshwater is provided from a watershed of 4,138 km<sup>2</sup> by three main channels, the Eyre, the Porge and the Landes, as well as twenty-six other streams and local groundwater, for a total supply of 1,340 million m<sup>3</sup> freshwater per year (Lamour & Balades, 1979; Auby et al., 1994).

105 The largest part of this watershed consists of Cenozoic river deposits, with, to a lesser extent, limestone, clay (Dubreuilh & Bouchet, 1992), and some iron oxide deposits historically used as building material (Gourdon-Platel & Maurin, 2004).

*Crassostrea gigas* specimens (n=8; Figure 1, Table 1) from this locality originate from the same rearing experiment as those that were placed at Baie des Veys and measurements were restricted to the parts of the shells corresponding to the period spent at this locality (Lartaud et al., 2010a). These specimens are referred to as Mod\_TES\_Cgig.

### 110 **2.1.4 Leucate (Aude)**

The Salses-Leucate lagoon is located on the southwestern French Mediterranean coast. It corresponds to a shallow coastal basin 14 km long and 5 km wide, separated from the Mediterranean Sea by a sandy barrier interrupted by three narrow marine inlets. The average water depth is 1.7 m and the hydrology is a balance between entrance of marine waters from the Mediterranean Sea, supply of groundwater discharges from two main karstic springs with flows of 3 x 10<sup>5</sup> m<sup>3</sup> d<sup>-1</sup> and 2 x 10<sup>5</sup> m<sup>3</sup> d<sup>-1</sup>, respectively (Fleury et al., 2007), and rainfall of approximately 500 mm y<sup>-1</sup> restricted to the fall and spring periods. The superficial watershed covers 162 km<sup>2</sup> but the total area including the karstic waters is not yet known accurately, likely extended to 60 km inland far from the shore (Salvayre, 1989; Ladagnous & Le Bec, 1997), with karstic waters penetrating Jurassic and Cretaceous limestone and dolomite. While tidal range is restricted, seawater level changes in the lagoon are controlled by strong northwesterly winds, regularly exceeding 10 m s<sup>-1</sup> (Rodellas et al., 2018).

120 *Crassostrea gigas* oysters (n=5; Figure 1, Table 1) originate from a wild brood stock in the vicinity of the local oyster farming area. We were not able to collect reliable *O. edulis* specimens from the Mediterranean Sea shoreline for comparison, as only aquaculture specimens of *C. gigas* are now available here. Specimens from this group are referred to as Mod\_LEU\_Cgig.

## **2.2 Archaeological sites and specimens**

### **2.2.1 Lyon, Auvergne-Rhône-Alpes**

125 In the area of the Fourvière hill of Lyon city, where remains of a building were tentatively identified as a sanctuary to the goddess Cybele, a pit was filled by food wastes which included around 200 valves of the flat oyster *O. edulis* (Bardot-Cambot, 2013). Absolute dating of this pit is currently being re-evaluated, and is approximated to the beginning of the current era or during the 1<sup>st</sup> century CE. The provenance of these oysters is debated (Bardot-Cambot, 2013). Two groups of animals, one

originating from the Mediterranean Sea coastline and the other from the Atlantic Ocean coastline, were identified based on morphometric measurements and associated mollusc shells (Bardot-Cambot, 2013). Six *O. edulis* specimens were selected from the first group (later referred to as Anc\_CYB1\_Oedu group) and seven more from the second group (Anc\_CYB2\_Oedu group) for the preservation quality of their umbo (Figure 1, Table 1).

### 2.2.2 La Malène, Occitany

This medieval site (circa the 6<sup>th</sup> century CE) is located on top of a cliff and is comprised of the remains of a fortified construction (Schneider & Clément, 2012). This *castrum* corresponds to one of the last antique sites where oysters were consumed by the elite, with supposedly spatially-restricted commercial travel (Bardot-Cambot & Forest, 2014). Still, the shells found in a dump, along with other proofs of the high social status of the occupants (such as golden currency and silver nails; Schneider & Clément, 2012), had been transported for over 120 km from the Mediterranean Sea. This origin is certified because some valves are fixed on valves of *Flexopecten glaber*, which is endemic to the Mediterranean Sea. Six *O. edulis* specimens were selected for the preservation quality of their umbo (Figure 1, Table 1). As ancient specimens, those are referred to as Anc\_MAL\_Oedu.

(Figure 1)

145 (Table 1)

### 2.3 Sample preparation

All specimens were mechanically cleaned from any epibiont and were selected according to the preservation state of their umbo region. The umbo was cut from the rest of the shell and embedded in Huntsman Araldite 2020 epoxy resin. Longitudinal thick sections (approx. 750  $\mu\text{m}$  thick) were manufactured to expose the preserved internal structures (Figure 2), in order to perform geochemical analyses on this protected region, away from shell surface organic or chemical contaminants. An extensive chemical cleaning of the section surfaces, as advised by Zaky et al. (2015), was not performed as the analytical surface was preserved from any external contaminants over the history of the shell. However, the influence of organic matter occluded in the crystal lattice cannot be discarded, as in any LA-ICP-MS work on biominerals.

155

(Figure 2)

All sections were observed under cathodoluminescence using a Cathodyne-OPEA cold cathode at IStEP, Sorbonne Université (Paris, France). Observation settings were 15-20 kV and 200-400  $\mu\text{A mm}^{-2}$ , at a pressure of 0.05 Torr. Areas potentially affected by diagenesis or damaged were identified in order to avoid any analysis of these regions by LA-ICP-MS. In addition,

160

cathodoluminescence (CL) observations were used to define seasonal calibration of the umbo, according to Langlet et al. (2006) and Lartaud et al. (2010a). Only Leucate specimens did not have seasonal calibration as the CL signal was uniform and nearly absent for these specimens. A second seasonal calibration method from Kirby et al. (1998), based on a sclerochronological record on the ligamental area in the form of external convex and concave bands, was attempted. 165 Unfortunately, umbos from Leucate specimens did not exhibit the necessary curved surface to conduct such a study. Consequently, measurement data from Leucate specimens were removed from the dataset used for the study of seasonal contrasts of the REE and Y fingerprints.

## 2.4 Geochemical analyses

Chemical analyses were carried out by LA-ICP-MS at ISTE (University of Lausanne, Switzerland). Measurements were 170 performed using an Element XR (ThermoScientific) ICP-MS coupled with a RESOLUTION 193 nm ArF excimer ablation system equipped with an S155 two-volume ablation cell (Australian Scientific Instruments). A pulse repetition rate of 20 Hz and an on-sample energy density of 4 J cm<sup>2</sup> were used. Pre-ablation of spots was first conducted in order to clean the surface of potential contaminants that could possibly be introduced during the sanding and polishing of the samples. The analytical spots were 200 µm in diameter. Ablation was performed on the areas of each sample section corresponding to winter and summer 175 periods (according to the cathodoluminescence seasonal calibration). This protocol allows the REE and Y incorporation to be compared at different seasons throughout the life of the oysters (Figure 2c), with the exception of Leucate specimens, which did not exhibit seasonal cathodoluminescence signals. Sections from Leucate specimens were analysed at random positions over the umbo region instead. Multiple measurements were performed on each section to avoid bias from potential internal variability. Repeated measurements of NIST SRM 612 prior to and following each 15-samples analytical series were used for 180 external standardisation. Accuracy was checked against measurements of the BCR-2 basalt reference material from USGS according to the GeoReM preferred values (Jochum et al., 2005). Relative standard deviation from 22 measurements of BCR-2 was always better than 2.8% for all REE and Y. Measured and expected values are indicated in Appendix B. Measured elements were La, Ce, Pr, Nd, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, Lu, Hf, Y, and Ca as the internal standard. Data reduction was performed using the LAMTRACE software (Jackson, 2008). A total of 297 measurements were made.

## 185 2.5 Data processing

Data processing was conducted using the Matlab software (MathWorks, www.mathworks.com, v. R2017a). None of the measured elements exhibit a normal distribution (Kolmogorov-Smirnov test): the distribution is right-skewed, at larger element abundances. To facilitate further statistical treatment of such data, they need to be transformed to normality, for which several mathematical transforms can be used. Here, we use the cubic root transform (Chen & Deo, 2004). Seasonal differences (from 190 the seasonal age models from cathodoluminescence) were estimated by hierarchical cluster analyses of 30 measurements from *C. gigas* specimens (n=5) and 41 measurements from *O. edulis* specimens (n=5) from Baie des Veys, and 24 measurements from *C. gigas* specimens (n=3) and 13 measurements from *O. edulis* specimens (n=3) from Marennes-Oléron (Appendix C).

For both species, two methods for calculating cluster distances were tested, (i) unweighted average distance and (ii) “Ward” inner squared distance. A cophenetic correlation coefficient has been calculated, as it is the linear correlation coefficient between the distances obtained from the cluster tree and the original distances (in the multivariate space). This coefficient is an indicator of the accuracy of the distances (estimated on the tree) to faithfully represent the dissimilarities among the observations. Multivariate analysis of variance (MANOVA) was used to compare the REE and Y fingerprints obtained from measurements of *C. gigas* and *O. edulis* from Baie des Veys, against the null hypothesis that both datasets belong to the same population. Kruskal-Wallis tests were performed to compare the Y/Ho ratios of multiple groups, against the null hypothesis that all groups belong to the same population. A recent statistical method, the t-SNE (t-distributed Stochastic Neighbour Embedding; van der Maaten & Hinton, 2008), was used to compare and classify the multivariate dataset (exact Euclidean method). The idea of the t-SNE method is to embed high-dimensional data points in low dimensions in a way that respects similarities between points. Nearby data points in the high-dimensional space correspond to nearby embedded low-dimensional points, and distant points in high-dimensional space correspond to distant embedded low-dimensional points (MathWorks, www.mathworks.com, v. R2017a).

### 3 Results

#### 3.1 Comparisons of the inter- and intra-specific seasonal record

Firstly, heavy REE (Tm, Yb, Lu) and Hf were usually not detected (below  $0.1 \text{ ng g}^{-1}$ ) in all the specimen groups, and were therefore removed from the dataset. Secondly, measurements from the Baie des Veys and Marennes-Oléron specimens allow for inter- and intra-specific comparisons, as they were performed on specimens from both *C. gigas* and *O. edulis* species. Data collected at the different seasons for each species did not show any significant difference in the incorporation of REE and Y between winter and summer. For Mod\_BDV\_Oedu, records from both winter (n=25) and summer (n=16) in *O. edulis* shell samples are mixed together, without clustering samples with respect to seasons (Appendix C); in addition, the cophenetic correlation coefficients are 0.92 and 0.76 for average and Ward methods, respectively, which emphasizes the quality of the classification. Results are similar for Mod\_BDV\_Cgig shells (n=19 and n=11 for winter and summer, respectively; Appendix C), with cophenetic correlation coefficients of 0.76 and 0.69 for average and Ward methods, respectively. Equivalent results are found for Mod\_MO\_Cgig (cophenetic correlation coefficients of 0.75 and 0.74 for average and Ward methods, respectively) and Mod\_MO\_Oedu (cophenetic correlation coefficients of 0.94 and 0.87 for average and Ward methods, respectively; Appendix C). However, a comparison between *C. gigas* and *O. edulis* (inter-specific comparison) clearly exhibits significant differences for both winter (MANOVA, p-value =  $1.10^{-8}$ ) and summer (MANOVA, p-value =  $2.10^{-5}$ ) periods between the two species. Therefore, *C. gigas* and *O. edulis* seem to record differently their respective seasonal signals.

### 3.2 The Y/Ho ratio as a provenance proxy

The Y/Ho ratio is commonly used as a provenance proxy (Bau et al., 1995; Prajith et al., 2015). The Y/Ho ratios in the present study (Figure 3) do not display significant differences between the four localities for modern *C. gigas* specimens (Kruskal-Wallis, p-values = 0.70 between groups Mod\_TES\_Cgig and Mod\_LEU\_Cgig, 0.95 between groups Mod\_TES\_Cgig and Mod\_BDV\_Cgig, 0.98 between groups Mod\_TES\_Cgig and Mod\_MO\_Cgig, 1.00 between groups Mod\_LEU\_Cgig and Mod\_BDV\_Cgig, 0.31 between groups Mod\_LEU\_Cgig and Mod\_MO\_Cgig, and 0.57 between groups Mod\_BDV\_Cgig and Mod\_MO\_Cgig). Also, all *O. edulis* modern and archaeological specimens from the other localities, except Anc\_CYB2\_Oedu, are similar to each other (Kruskal-Wallis, p-values = 0.98 between groups Anc\_CYB1\_Oedu and Anc\_MAL\_Oedu, 0.99 between groups Anc\_CYB1\_Oedu and Mod\_BDV\_Oedu, 1.00 between groups Anc\_CYB1\_Oedu and Mod\_MO\_Oedu, 1.00 between groups Anc\_MAL\_Oedu and Mod\_BDV\_Oedu, 0.97 between groups Anc\_MAL\_Oedu and Mod\_MO\_Oedu, and 0.97 between groups Mod\_BDV\_Oedu and Mod\_MO\_Oedu) but are different from *C. gigas* shells (Appendix D). Moreover, a significant difference between modern *C. gigas* and *O. edulis* shells from the same localities (Baie des Veys and Marennes-Oléron) is also reported (Appendix D). However, the Anc\_CYB2\_Oedu group does not share the homogeneity of other modern and ancient *O. edulis* populations. On the contrary, the ratios measured in *O. edulis* specimens from this group are not significantly different from the ones obtained in modern *C. gigas* specimens (Kruskal-Wallis, p-values = 0.97, 0.26, 0.48 and 1.00, when compared with specimens from Mod\_TES\_Cgig, Mod\_LEU\_Cgig, Mod\_BDV\_Cgig and Mod\_MO\_Cgig, respectively).

240 (Figure 3)

### 3.3 REE incorporation in shells

For all the specimens, a gradual decrease in REE dispersion (*i.e.*, amplitude of values relative to Y) is generally observed with their increasing atomic number (Figure 4). Indeed, several groups can be identified with light REE (*e.g.*, Pr and Sm), such as the *C. gigas* groups. On the contrary, middle REE (*e.g.*, Dy and Er) distributions appear similar for all groups, with the exception of the Mod\_LEU\_Cgig group whose measurements are isolated from those of the other groups. The REE median profiles (normalized to the Post-Archean Australian Shales; McLennan, 1989) also present similar trends of middle REE (from Gd to Er) for most groups (Figure 5), except for the Mod\_LEU\_Cgig group, which exhibits lower abundances than the other groups for all REE. However, light REE are generally substantially depleted in *C. gigas* specimens compared to modern and ancient *O. edulis* groups (approx. one order of magnitude difference). The only exceptions are La and Ce in *C. gigas* shells from Baie des Veys (Mod\_BDV\_Cgig), which present values in the range of those from modern and ancient *O. edulis* specimens. This pattern of enriched Ce (and to a lesser extent, La) is not shared by *O. edulis* specimens from this same locality



(Mod\_BDV\_Oedu). Although these two elements have similar abundances for both species in this locality, all the other REE abundances are different.

255

(Figure 4)

(Figure 5)

260 Results from the entire dataset (*i.e.*, 297 measurements and 12 elements per measurement) are compared in Figure 6 using t-SNE. Some groups are well identified by this method, such as the Mod\_LEU\_Cgig and Mod\_BDV\_Cgig groups (Mediterranean Sea and British Channel coastlines). Both Mod\_MO\_Cgig and Mod\_TES\_Cgig groups, originating from the Atlantic Ocean coastline, appear as one unique group by t-SNE, with a restricted dispersion. The modern and ancient *O. edulis* groups are however not discriminated by t-SNE and present a substantially larger dispersion than the Atlantic Ocean's *C. gigas* groups. In this sample set, Anc\_CYB1\_Oedu, Mod\_MO\_Oedu and Mod\_BDV\_Oedu groups are relatively similar in terms of range of distribution, and Anc\_CYB2\_Oedu and Anc\_MAL\_Oedu specimens share similarities; but the five groups remain poorly differentiated.

265

(Figure 6)

270

#### 4 Discussion

In both studied species, the decrease in the range of variation of REE abundances with the increasing atomic number (except for Tm, Yb, Lu, and Hf, which were not quantified in the shells) can be explained by the increased affinity of heavy REE for complexation in seawater, as it has been demonstrated in previous studies (Cantrell & Byrne, 1987; Byrne & Kim, 1990; De Baar et al., 1991). As these elements are trapped into complexed forms or ligands, their bioavailability in seawater is strongly reduced, limiting their insertion in the oyster ionic pumps leading to the mineralisation locus. These bioavailability restrictions of REE have already been demonstrated in the freshwater mussel *Corbicula fluminea* for Gd (Merschel & Bau, 2015) and heavy REE in *Mytilus edulis* (Ponnuram et al., 2016). Another explanation can be advanced regarding the technique used. LA-ICP-MS device analyses both the mineral and organic phases ablated from the biomineral without the possibility to assess their relative proportions. Although the mean proportion of organic compounds in oyster shells is limited (<0.5% for *C. gigas*; Mouchi et al., 2016), it is known that organic REE abundances are depleted in heavy REE (Freslon et al., 2014). The decreasing abundance with the increasing atomic number may then be caused by protein and polysaccharide contents. Only extensive cleaning for solution-based ICP-MS analyses would be able to remove entirely the organic molecules before measurements, but this would not fit the fast REE assessment from a large number of specimens we aimed to conduct in this study.

280

285 In this study, the Y/Ho ratios, which are usually proposed as a provenance proxy (Bau et al., 1995; Prajith et al., 2015), are  
affected by strong vital effects for both oyster species, potentially due to the decrease in REE abundance with increasing atomic  
number. In addition, significant differences between species from the same locality are also reported. Hence, the Y/Ho ratio  
generally does not depend on the original location (Figure 3). Consequently, Y/Ho should not be used directly as a provenance  
proxy (at least from LA-ICP-MS data collected on biogenic carbonates), or with extreme caution after having discarded any  
290 potential vital effect. As the decreased REE abundance with increasing atomic number discussed above prevents a locality-  
specific variation of Ho, other Y/REE ratios have been tested as alternative provenance proxies using lighter REE (Figure 4).  
Y/La, Y/Ce, Y/Pr and Y/Nd were all unsuccessful in providing identification of locality groups and also present similar values  
for all modern and ancient *O. edulis*. For Y/La and Y/Ce ratios, *C. gigas* specimens from Baie des Veys were identical to all  
modern and ancient *O. edulis* specimens, while for Y/Pr and Y/Nd ratios, *C. gigas* specimens from Baie des Veys were identical  
295 to all *C. gigas* specimens from the other localities. Overall, Y/REE ratios appear unsuccessful for provenance discrimination.  
Measurements performed on Baie des Veys and Marennes-Oléron specimens (Mod\_BDV\_Cgig and Mod\_BDV\_Oedu, as well  
as Mod\_MO\_Cgig and Mod\_MO\_Oedu) have been used to study the incorporation of REE inside a single species, in order to  
evaluate its intraspecific variation. Contrary to the models performed for different temperatures on the mussel *Mytilus edulis*  
(Ponnurangam et al., 2016), the seasonal conditions do not have any impact on the REE incorporation, neither for *O. edulis*  
300 nor *C. gigas* shells (in the range of 5-20 °C; Table 1). Other parameters than temperature and pH, used by Ponnurangam et al.  
(2016), are probably in effect, which lowers the impact of temperature on REE incorporation. This observation implies that  
any part of a shell can be sampled without necessarily having to define a temporal calibration of the umbo. However, REE  
abundances fluctuate widely within a single specimen, and the “local fingerprint” of these elements needs to be clarified and  
based on multiple measurements performed on each of several specimens. These intra-individual fluctuations cannot be due  
305 to seasonally-controlled environmental factors, such as temperature, precipitations or plankton blooms. However, a potential  
source of REE for oysters can be the porewater or resuspended sediment (Haley et al., 2004; Crocket et al., 2018), and therefore,  
the relative abundances may fluctuate without precise temporal cyclicality.

The reasons for these intra- and inter-specific vital effects remain unknown. Indeed, as far as we know, no study has ever  
shown evidence or suspicion of the use of REE in metabolic processes that could induce an effective filter of these elements  
310 between seawater and the extrapallial cavity where shell mineralization occurs. Nevertheless, it has been reported that REE,  
or other unsuspectedly useful elements, are indeed used by organisms in specific environmental settings; an example is  
provided by diatoms in Zn-depleted conditions, where Zn is used as a co-factor of carbonic anhydrase (Lee et al., 1995).  
Another example is given by methanotrophic archaea, which use Cd, a toxic element, as a co-factor of methanol dehydrogenase  
(Pol et al., 2014).

315 Incorporation of REE differ between both studied species. Not only *C. gigas* shells present different REE profiles between  
groups (unlike those of *O. edulis*), but also the Gd positive anomaly, a characteristic of modern coasts under pressure of  
anthropic activities (Bau & Dulski, 1996; Nozaki et al., 2000; Le Goff et al., 2019), is observed solely for “modern” *C. gigas*  
specimens from Mod\_BDV\_Cgig, Mod\_MO\_Cgig and Mod\_TES\_Cgig groups (Figure 5), which correspond to the only

watersheds with major cities. The Gd anomaly is neither visible in modern *C. gigas* from Mod\_LEU\_Cgig, nor in modern *O. edulis*. The systematically low abundances of REE in the Leucate shells can also be explained by the regional geology (Ladagnous & Le Bec, 1997), as watersheds of the other localities of *C. gigas* specimens present substratum types with higher REE contents than karsts (e.g., such as basalts; Baize et al., 1997). The dispersion of measurements using t-SNE seems to be ineffective to discriminate between Mod\_MO\_Cgig and Mod\_TES\_Cgig groups. Both these groups are located on the Atlantic Ocean coastline of France, and both respective watersheds comprise the same rock types (i.e., mainly limestones and some clays and river deposits). One could therefore expect a similar riverine water REE content. Overall, these species-specific characteristics indicate that *C. gigas* can be used as a sentinel species regarding REE pollution of coastal waters. On the contrary, we cannot confirm that *O. edulis* is a proper candidate for such studies.

Several reasons for these interspecific differences can be advanced. It is known that oysters can be selective in their diet, composed mainly of diatoms (Yonge, 1928; Paulmier, 1971) of a specific size range, and preferentially digest specific species of diatoms over others (Shumway et al., 1985; Cognie et al., 2001). If food is a source of REE, it may be possible that each oyster species does not feed on the same prey, which can present different abundances of these elements. Aquarium experiments reported different ingestion rates in the 5-15  $\mu\text{m}$  algal size range between these oyster species (Nielsen et al., 2017), but there is no indication on the REE content of the food. Alternatively, *O. edulis*, which exhibits generally higher abundances of REE (nearly one order of magnitude higher for light REE, except for La and Ce for Mod\_BDV\_Cgig specimens; Figure 5), could present a higher bioaccumulation of these elements in its soft tissues (and eventually shell) compared to *C. gigas*. Ong et al. (2013) presented trace element measurements from soft tissues of both species from the Baie de Quiberon (Brittany, France) indicating that soft tissues of *O. edulis* contain generally less Cu and Zn but more Cd and Pb than those of *C. gigas*. Such species-specific bioaccumulation and incorporation differences could also be in effect for REE. Finally, it is possible to explain the higher abundance of light REE in *O. edulis* shells compared to *C. gigas* by suggesting that *O. edulis* ingests more clay particles. As heavy REE are trapped in complexed form in seawater, mainly light REE must be available for adsorption on clay particles, and eventually integrate the forming carbonate shell.

In any case, this study shows that t-SNE can be used on REE and Y measurements from *C. gigas* shells to identify regions of origin of specimens from this species. However, it appears that intraspecific vital effects prevent its efficiency in *O. edulis*, whose specimens exhibit the same fingerprint for several localities of origin. For this reason, we cannot confirm or refute a different origin of the two populations of the Cybèle archaeological specimens (Anc\_CYB1\_Oedu and Anc\_CYB2\_Oedu) with these elements.

## 5 Conclusions

Multiple types of vital effects and/or other factors on REE incorporation in *C. gigas* and *O. edulis* oyster shells have been highlighted in this study. Intraspecific variations in REE abundances are significant but not related to seasonal fluctuations. A gradual decrease in REE incorporations with increasing atomic numbers has been observed, and it appears that heavy REE are

less discriminant than light REE in identifying the various studied groups. The Y/Ho ratio, previously reported as a proxy for provenance studies, is ineffective in oyster shells. Finally, interspecific variations underline the ability of t-SNE procedure to correctly separate *C. gigas* specimens of various regions of origin but not *O. edulis* specimens, which also implies that only *C. gigas* can be used as a monitor species of light REE pollution. Reconstruction of the provenance of oyster specimens will therefore have to be performed separately for each studied species, as regional geochemical fingerprints of the shells appear to be species-dependant. In order to be able to identify the regions of origin of species affected by strong vital effects (such as *O. edulis*), it is necessary to investigate other chemical elements as potential provenance proxies. Moreover, an increased efficiency in identifying the locality (and not only the region) of origin of *C. gigas* specimens could be achieved by also measuring other elements than REE.

## 360 **6 Data availability**

The data used in this manuscript have been deposited on the Zenodo data repository (Mouchi et al., 2019).

## **7 Author contribution**

VM conceived the study. VM, CG and AU performed the data analysis. VF and FL provided specimens. VM and EV performed statistics and data processing. VM, CG, MdR, LE, EV and FL interpreted the results. VM wrote the manuscript with contributions from all authors.

## **8 Competing interests**

The authors declare that they have no conflict of interest.

## **9 Acknowledgements**

The authors would like to thank Frédéric Delbès for the work he performed on the preparation of the thin sections.

## **References**

Akagi, T., Hashimoto, Y., Fu, F.-F., Tsuno, H., Tao, H., and Nakano, Y.: Variation of the distribution coefficients of rare earth elements in modern coral-lattices: Species and site dependencies, *Geochim. Cosmochim. Acta*, 68, 2265-2273, doi: 10.1016/j.gca.2003.12.014, 2004.

- Andrisoa, A.: Ecological impacts of groundwater discharges to Mediterranean coastal lagoons, PhD thesis, Univ Aix-Marseille. 187 p, 2019.
- Auby, I., Manaud, F., Maurer, D., and Trut, G.: Etude de la prolifération de algues vertes dans le bassin d'Arcachon. *IFREMER – CEMAGREF – SSA – SABARC*, 270 p, 1994.
- 380 Baize, S., Camuzard, J.-P., Freslon, M., Langevin, C., and Laignel, B.: Notice explicative, Carte géol. France (1/50 000), feuille Carentan (117). Orléans: BRGM, 83 p, Carte géologique par S. Baize et al., 1997.
- Bambier, A., Capdeville, J.-P., Cariou, E., Floc'h, J.-P., Gabilly, J., and Hantzpergue, P.: Notice explicative, Carte géol. France (1/50 000), feuille La Rochefoucauld (686). Orléans: BRGM, 30 p, Carte géologique par A. Bambier et al., 1982.
- Bardot-Cambot, A.: Les coquillages marins en Gaule romaine. Approche socio-économique et socio-culturelle, BAR IS, 2481,  
385 Archaeopress, Oxford, 270 p, 2013.
- Bardot-Cambot, A.: Consommer dans les campagnes de la Gaule romaine. *Revue du Nord*, 21, Hors-série, Collection Art et Archéologie, 109-123, 2014.
- Bardot-Cambot, A. and Forest, V.: Une histoire languedocienne des coquillages marins consommés, du Mésolithique à nos jours. In: Costamagno S. (Ed.) *Histoire de l'alimentation humaine : entre choix et contraintes*. Actes du 138<sup>e</sup> congrès national  
390 des sociétés historiques et scientifiques (Rennes, 2013). Actes des congrès nationaux des sociétés historiques et scientifiques. Édition électronique, p. 88-104, 2014.
- Bau, M., Dulski, P., and Möller, P.: Yttrium and holmium in South Pacific seawater: Vertical distribution and possible fractionation mechanisms. *Chem. Erde*, 55, 1-15, 1995.
- Bau, M. and Dulski, P.: Anthropogenic origin of positive gadolinium anomalies in river waters, *Earth Planet. Sci. Lett.*, 143,  
395 245-255, doi 10.1016/0012-821X(96)00127-6, 1996.
- Bayne, B.L., Ahrens, M., Allen, S.K., Anglès D'auriac, M., Backeljau, T., Beninger, P., Bohn, R., Boudry, P., Davis, J., Green, T., Guo, X., Hedgecock, D., Ibarra, A., Kingsley-Smith, P., Krause, M., Langdon, C., Lapègue, S., Li, C., Manahan, D., Mann, R., Perez-Paralle, L., Powell, E.N., Rawson, P.D., Speiser, D., Sanchez, J.-L., Shumway, S., and Wang, H.: The proposed dropping of the genus *Crassostrea* for all Pacific cupped oysters and its replacement by a new genus *Magallana*: A dissenting  
400 view, *J. Shellfish Res.*, 36, 545-547, doi 10.2983/035.036.0301, 2017.
- Bennion, M., Morrison, L., Brophy, D., Carlsson, J., Cortiñas Abrahantes, J., and Graham, C.T.: Trace element fingerprinting of blue mussel (*Mytilus edulis*) shells and soft tissues successfully reveals harvesting locations, *Sci. Total Environ.*, 685, 50-58, doi 10.1016/j.scitotenv.2019.05.233, 2019.
- Bourgueil, B., Moreau, P., Gabet, C., L'Homer, A., and Vouve, J.: Notice explicative, Carte géol. France (1/50 000), feuille  
405 Rochefort (658). Orléans: BRGM, 30 p, Carte géologique par B. Bourgueil et al., 1972.
- Bourgueil, B., Moreau, P., and Vouve, J.: Notice explicative, Carte géol. France (1/50 000), feuille Saintes (683). Orléans: BRGM, 19 p, Carte géologique par B. Bourgueil et al., 1968.
- Byrne, R.H. and Kim, K.H.: Rare-earth element scavenging in seawater, *Geochim. Cosmochim. Acta*, 54, 2645-2656, doi 10.1016/0016-7037(90)90002-3, 1990.

- 410 Cantrell, K.J. and Byrne, R.H.: Rare-earth element complexation by carbonate and oxalate ions, *Geochim. Cosmochim. Acta*, 51, 597-605, doi 10.1016/0016-7037(87)90072-X, 1987.
- Censi, P., Mazzola, S., Sprovieri, M, Bonanno, A., Patti, B., Punturo, R., Spoto, S.E., Saiano, F., and Alonzo, G.: Rare earth elements distribution in seawater and suspended particulate of the Central Mediterranean Sea, *Chem. Ecol.*, 20, 323-343, doi 10.1080/02757540410001727954, 2004.
- 415 Chen, W.W. and Deo, R.S.: Power transformations to induce normality and their applications, *J. R. Stat. Soc. B*, 66, 117-130, 2004.
- Cognie, B., Barillé, L., and Rincé, Y.: Selective feeding of the oyster *Crassostrea gigas* fed on a natural microphytobenthos assemblage, *Estuaries*, 24, 126-131, 2001.
- Crocket, K.C., Hill, E., Abell, R.E., Johnson, C., Gary, S.F., Brand, T., and Hathorne, E.D.: Rare Earth Element Distribution in the NE Atlantic: Evidence for Benthic Sources, Longevity of the Seawater Signal, and Biogeochemical Cycling, *Front. Mar. Sci.*, 5, 147, doi 10.3389/fmars.2018.00147, 2018.
- 420 De Baar, H.J.W., Bacon, M., and Brewer, P.G.: Rare-earth distributions with positive Ce anomaly in the western North Atlantic Ocean, *Nature*, 301, 324-327, doi 10.1038/301324a0, 1983.
- De Baar, H.J.W., Schijf, J., and Byrne, R.H.: Solution chemistry of the rare-earth elements in seawater, *Eur. J. Inorg. Chem.*, 425 28, 357-373, 1991.
- Dechambenoy, C., Pontier, L., Sirou, F., and Vouvé, J.: Apport de la thermographie infrarouge aéroportée à la connaissance de la dynamique superficielle des estuaires (système Charente – Seudre – Anse de l’Aiguillon), *C. R. Acad. Sci.*, 284, 1269-1272, 1977.
- Dubreuilh, J., Karnay, G., Bouchet, J.-M., and Le Nindre, Y.-M.: Notice explicative, Carte géol. France (1/50 000), feuille 430 Arcachon (825), Orléans, BRGM, 53 p., Carte géologique par J. Dubreuilh, J.-M. Bouchet, 1992.
- Elderfield, H.: The oceanic chemistry of the rare-earth elements, *Phil. Trans. R. Soc. A*, 325, 105-126, 1988.
- Elderfield, H. and Greaves, M.J.: The rare earth elements in seawater, *Nature*, 296, 214-219, doi 10.1038/296214a0, 1982.
- Fleury, P., Bakalowicz, M., and de Marsily, G.: Submarine springs and coastal karst aquifers: A review, *J. Hydrol.*, 339, 79–92, doi 10.1016/j.jhydrol.2007.03.009, 2007.
- 435 Freslon, N., Bayon, G., Toucanne, S., Bermell, S., Bollinger, C., Chéron, S., Etoubleau, J., Germain, Y., Khripounoff, A., Ponzevera, E., and Rouget, M.-L.: Rare earth elements and neodymium isotopes in sedimentary organic matter, *Geochim. Cosmochim. Acta*, 140, 177-198, doi 10.1016/j.gca.2014.05.016, 2014.
- Frost, C.D., O’Nions, R.K., and Goldstein, S.L.: Mass balance for Nd in the Mediterranean Sea, *Chem. Geol.*, 55, 45-50, doi 10.1016/0009-2541(86)90126-9, 1986.
- 440 Goldstein, S.L., O’Nions, R.K., and Hamilton, P.J.: An Sm-Nd isotopic study of atmospheric dusts and particulates from major river systems, *Earth Planet. Sci. Lett.*, 70, 221-236, doi 10.1016/0012-821X(84)90007-4, 1984.
- Gopi, K., Mazumder, D., Sammut, J., and Saintilan, N.: Determining the provenance and authenticity of seafood: A review of current methodologies, *Trends Food Sci. Tech.*, 91, 294-304, doi 10.1016/j.tifs.2019.07.010, 2019.

- Gourdon-Platel, N. and Maurin, B.: Le fer des marais, encroûtement superficiel holocène utilisé sur les sites archéologiques de Sanguinet (Landes, France), *Géologie de la France*, 1, 13-24, 2004.
- 445 Greaves, M.J., Rudnicki, M., and Elderfield, H.: Rare earth elements in the Mediterranean Sea and mixing in the Mediterranean Outflow, *Earth Planet. Sci. Lett.*, 103, 169-181, doi 10.1016/0012-821X(91)90158-E, 1991.
- Haley, B.A., Klinkhammer, G.P., and McManus, J.: Rare earth elements in pore waters of marine sediments, *Geochim. Cosmochim. Acta*, 68, 1265-1279, doi 10.1016/j.gca.2003.09.012, 2004.
- 450 Haley, B.A., Klinkhammer, G.P., and Mix, A.C.: Revisiting the rare earth elements in foraminiferal tests, *Earth Planet. Sci. Lett.*, 239, 79-97, doi 10.1016/j.epsl.2005.08.014, 2005.
- Hantzpergue, P., Bonnin, J., Cariou, E., Gomez de Soto, J., and Moreau, P.: Notice explicative, Carte géol. France (1/50 000), feuille Mansle (685). Orléans: BRGM, 24 p, Carte géologique par P. Hantzpergue et al., 1984.
- Huyghe, D., de Rafélis, M., Ropert, M., Mouchi, V., Emmanuel, L., Renard, M., Lartaud, F.: New insight into oyster high-resolution shell growth patterns, *Mar. Biol.*, 166, 48, doi 10.1007/s00227-019-3496-2, 2019.
- 455 Jackson, S.E.: LAMTRACE data reduction software for LA-ICP-MS. *In: Laser ablation ICP-MS in the Earth sciences: Current practices and outstanding issues*, *Can. Mineral.*, 40, 305-307, 2008.
- Jochum, K.P., Seufert, H.M., Spettel, B., and Palme, H.: The solar-system abundances of Nb, Ta, and Y, and the relative abundances of refractory lithophile elements in differentiated planetary bodies, *Geochim. Cosmochim. Acta*, 50, 1173-1183, doi 10.1016/0016-7037(86)90400-X, 1986.
- 460 Jochum, K.P., Nohl, U., Herwig, K., Lammel, E., Stoll, B., and Hofmann, A.W.: GeoReM: A new Geochemical database for reference materials and isotopic standards, *Geostand. Geoanalytical Res.*, 29, 333-338, doi 10.1111/j.1751-908X.2005.tb00904.x, 2005.
- Kelly, S., Heaton, K., and Hoogewerff, J.: Tracing the geographic origin of food: The application of multi-element and multi-isotope analysis, *Trends Food Sci. Tech.*, 16, 555-567, doi 10.1016/j.tifs.2005.08.008, 2005.
- 465 Kirby, M.X., Soniat, T.M., and Spero, H.J.: Stable isotope sclerochronology of Pleistocene and Recent oyster shells (*Crassostrea virginica*), *Palaios*, 13, 560-569, doi 10.2307/3515347, 1998.
- Ladagnous, H. and Le Bec, C.: Lagune de Sables-Leucate. I. – Analyse bibliographique, Rapport interne de l'IFREMER, 94 p., R.INT.DEL/97-02/SETE, 1997.
- 470 Lamour, J. and Balades, J.D.: Suivi de la qualité des eaux du Bassin d'Arcachon, C.E.T.E. Laboratoire régional de Bordeaux, 26 p, 1979.
- Langlet, D., Alunno-Bruscia, M., Rafélis, M., Renard, M., Roux, M., Schein, E., and Buestel, D.: Experimental and natural cathodoluminescence in the shell of *Crassostrea gigas* from Thau lagoon (France): ecological and environmental implications, *Mar. Ecol. Prog. Ser.*, 317, 143-156, doi 10.3354/meps317143, 2006.
- 475 Lartaud, F., de Rafélis, M., Ropert, M., Emmanuel, L., Geairon, P., and Renard, M.: Mn labelling of living oysters: artificial and natural cathodoluminescence analyses as a tool for age and growth rate determination of *C. gigas* (Thunberg, 1793) shells, *Aquaculture*, 300(1), 206-217, doi 10.1016/j.aquaculture.2009.12.018, 2010a.

- Lartaud, F., Emmanuel, L., de Rafélis, M., Ropert, M., Labourdette, N., Richardson, C.A., and Renard, M.: A latitudinal gradient of seasonal temperature variation recorded in oyster shells from the coastal waters of France and The Netherlands. *Facies*, 56, 13, doi 10.1007/s10347-009-0196-2, 2010b.
- 480 Le Gall, J.: La Baie des Veys : caractères principaux de la sédimentation et faciès de dépôt, Ph.D. thesis, Université de Caen, 151 p, 1970.
- Le Goff, S., Barrat, J.-A., Chauvaud, L., Paulet, Y.-M., Gueguen, B., and Ben Salem, D.: Compound-specific recording of gadolinium pollution in coastal waters by great scallops, *Sci. Rep.*, 9, 8015, doi 10.1038/s41598-019-44539-y, 2019.
- 485 Lee, J.G., Roberts, S.B., and Morel, F.M.M.: Cadmium: A nutrient for the marine diatom *Thalassiosira weissflogii*, *Limnol. Oceanogr.*, 40, 1056-1063, doi 10.4319/lo.1995.40.6.1056, 1995.
- Li, F., Webb, G.E., Algeo, T.J., Kershaw, S., Lu, C., Oehlert, A.M., Gong, Q., Pourmand, A., and Tan, X.: Modern carbonate ooids preserve ambient aqueous REE signature, *Chem. Geol.*, 509, 163-177, doi 10.1016/j.chemgeo.2019.01.015, 2019.
- McLennan, S.M.: Rare earth elements in sedimentary rocks; influence of provenance and sedimentary processes, *Rev. Mineral. Geochem.*, 21, 169-200, 1989.
- 490 Merschel, G. and Bau, M.: Rare earth elements in the aragonitic shell of freshwater mussel *Corbicula fluminea* and the bioavailability of anthropogenic lanthanum, samarium and gadolinium in river water, *Sci. Total Environ.*, 533, 91-101, doi 10.1016/j.scitotenv.2015.06.042, 2015.
- Morrison, L., Bennion, M., Gill, S., and Graham, C.T.: Spatio-temporal trace element fingerprinting of king scallops (*Pecten maximus*) reveals harvesting period and location, *Sci. Total Environ.*, 697, 134121, 2019.
- 495 Mouchi, V., de Rafélis, M., Lartaud, F., Fialin, M., and Verrecchia E.: Chemical labelling of oyster shells used for time-calibrated high resolution Mg/Ca ratios: a tool for past estimation of seasonal temperature variations, *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 373, pp. 66-74, doi 10.1016/j.palaeo.2012.05.023, 2013.
- Mouchi, V., Lartaud, F., Guichard, N., Immel, I., de Rafélis, M., Broussard, C., Crowley, Q.G., and Marin, F.: Chalky versus foliated: a discriminant immunogold labelling of shell microstructures in the edible oyster *Crassostrea gigas*, *Mar. Biol.*, 163, 1-15, doi: 10.1007/s00227-016-3040-6, 2016.
- 500 Mouchi, V., Godbillot, C., Ulyanov, A., Forest, V., de Rafélis, M., Emmanuel, L., and Verrecchia, E.P.: Data for the publication "Rare Earth Elements in oyster shells: provenance discrimination and potential vital effects", Zenodo, <http://doi.org/10.5281/zenodo.3529419>, 2019.
- 505 Mourier, J.-P., Floc'h, J.-P., and Coubès, L.: Notice explicative, Carte géol. France (1/50 000), feuille L'Isle-Jourdain (638). Orléans: BRGM, 73 p, Carte géologique par J.-P. Mourier et al., 1989.
- Nielsen, M., Hansen, B.W., and Vismann, B.: Feeding traits of the European flat oyster, *Ostrea edulis*, and the invasive Pacific oyster, *Crassostrea gigas*, *Mar. Biol.*, 164, 6, doi 10.1007/s00227-016-3041-5, 2017.
- Nozaki, Y., Zhang, J., and Amakawa, H.: The fractionation between Y and Ho in the marine environment, *Earth Planet. Sci. Lett.*, 148, 329-340, doi 10.1016/S0012-821X(97)00034-4, 1997.
- 510



- Nozaki, Y., Lerche, D., Sotto Alibo, D., and Snidvongs, A.: The estuarine geochemistry of rare earth elements and indium in the Chao Phraya River, Thailand, *Geochim. Cosmochim. Acta*, 64, 3983-3994, doi 10.1016/S0016-7037(00)00473-7, 2000.
- Olivarez, A.M. and Owen, R.M.: The europium anomaly of seawater: implications for fluvial versus hydrothermal REE inputs to the oceans, *Chem. Geol.*, 92, 317-328, doi 10.1016/0009-2541(91)90076-4, 1991.
- 515 Ong, M.C., Noor Azhar, M.S., Menier, D., and Effendy, A.W.M.: Levels of trace elements in tissue of *Ostrea edulis* and *Crassostrea gigas* from Quiberon Bay, Brittany, France, *J. Fish. Aquat. Sci.*, 8, 378-387, doi 10.3923/jfas.2013.378.387, 2013.
- Osborne, A.H., Hathorne, E.C., Schijf, J., Plancherel, Y., Böning, P., and Frank, M.: The potential of sedimentary foraminiferal rare earth element patterns to trace water masses in the past, *Geochem. Geophys. Geosyst.*, 18, 1550-1568, doi 10.1002/2016GC006782, 2017.
- 520 Paulmier, G.: Cycle des matières organiques dissoutes, du plankton et du micro-phytobenthos dans l'estuaire du Belon. Leur importance dans l'alimentation des huîtres, *Rev. Trav. Inst. Pêches Marit.*, 35, 157-200, 1971.
- Piper, D.Z. and Bau, M.: Normalized rare earth elements in water, sediments, and wine: Identifying sources and environmental redox conditions, *Am. J. Analyt. Chem.*, 4, 69-83, doi 10.4236/ajac.2013.410A1009, 2013.
- Platel, J.-P., Moreau, P., Vouvé, J., Debenath, A., Colmont, G.R., and Gabet, C.: Notice explicative, Carte géol. France (1/50 000), feuille St-Agnant (682). Orléans: BRGM, 52 p, Carte géologique par J.-P. Platel et al., 1978.
- 525 Platel, J.-P., Moreau, P., Vouvé, J., and Colmont, G.R.: Notice explicative, Carte géol. France (1/50 000), feuille Pons (707). Orléans: BRGM, 43 p, Carte géologique par J.-P. Platel et al., 1977.
- Pol, A., Barends, T.R.M., Dietl, A., Khadem, A.F., Eygensteyn, J., Jetten, M.S.M., and Op den Camp, H.J.M.: Rare earth metals are essential for methanotrophic life in volcanic mudpots, *Environ. Microbiol.*, 16, 255-264, 2014.
- 530 Ponnurangam, A., Bau, M., Brenner, M., and Koschinsky, A.: Mussel shells of *Mytilus edulis* as bioarchives of the distribution of rare earth elements and yttrium in seawater and the potential impact of pH and temperature on their partitioning behaviour, *Biogeosciences*, 13, 751-760, doi 10.5194/bg-13-751-2016, 2016.
- Prajith, A., Rao, V.P., and Kessarkar, P.M.: Controls on the distribution and fractionation of yttrium and rare earth elements in core sediments from the Mandovi estuary, western India, *Cont. Shelf Res.*, 92, 59-71, doi 10.1016/j.csr.2014.11.003, 2015.
- 535 Rodellas, V., Stieglitz, T. C., Andrisoa, A., Cook, P. G., Raimbault, P., Tamborski, J.J., van Beek, P., and Radakovitch, O.: Groundwater-driven nutrient inputs to coastal lagoons: The relevance of lagoon water recirculation as a conveyor of dissolved nutrients, *Sci. Total Environ.*, 642, 764–780, doi 10.1016/j.scitotenv.2018.06.095, 2018.
- Salvayre, H.: Les karsts des Pyrénées-Orientales (caractères hydrogéologiques et spéléologiques généraux), *Karstologia*, 13, 1-10, doi 10.3406/karst.1989.2199, 1989.
- 540 Salvi, D. and Mariottini, P.: Molecular taxonomy in 2D: a novel ITS2 rRNA sequence structure approach guides the description of the oysters' subfamily *Saccostreinae* and the genus *Magallana* (Bivalvia: Ostreidae), *Zool. J. Linn. Soc.*, doi 10.1111/zoj.12455, 2016.
- Schijf, J., Christenson, E.A., and Byrne, R.H.: YREE scavenging in seawater: A new look at an old model, *Mar. Chem.*, 177, 460-471, doi 10.1016/j.marchem.2015.06.010, 2015.


- 545 Schneider, L. and Clément, N.: Le castellum de La Malène en Gévaudan. Un "rocher monument" du premier Moyen Age (VI<sup>e</sup>-VII<sup>e</sup> s.), Académie des Inscriptions et Belles Lettres, La Lozère (48) Carte archéologique de la Gaule, Comptoir des presses d'universités, pp. 317-328, EAN 978-2-87754-277-7, 2012.
- Sholkovitz, E.R. and Shen, G.T.: The incorporation of rare earth elements in modern coral, *Geochim. Cosmochim. Acta*, 59, 2749-2756, doi 10.1016/0016-7037(95)00170-5, 1995.
- 550 Sholkovitz, E.R., Landing, W.M., and Lewis, B.L.: Ocean particle chemistry: The fractionation of rare earth elements between suspended particles and seawater, *Geochim. Cosmochim. Acta*, 58, 1567-1579, doi 10.1016/0016-7037(94)90559-2, 1994.
- Shumway, S.E., Cucci, T.L., Newell, R.C., and Yentsch, C.M.: Particle selection, ingestion, and absorption in filter-feeding bivalves, *J. Exp. Mar. Biol. Ecol.*, 91, 77-92, doi 10.1016/0022-0981(85)90222-9, 1985.
- Soletchnik, P., Faury, N., Razet, D., and Gouletquer, P.: Hydrobiology of the Marennes-Oléron bay. Seasonal indices and analysis of trends from 1978 to 1995, *Hydrobiologia*, 386, 131-146, 1998.
- 555 Sylvand, B.: La Baie des Veys, 1972-1992 : structure et évolution à long terme d'un écosystème benthique intertidal de substrat meuble sous influence estuarienne, Ph.D. thesis, Université de Caen, 409 p, 1995.
- Taylor, S.R. and McLennan, S.M.: The significance of the rare earths in geochemistry and cosmochemistry. *In: Gschneidner Jr., K.A., Eyring, L. (Eds.), Handbook on the Physics and Chemistry of Rare Earths*, 11. Elsevier, Amsterdam, 485-578, 1988.
- 560 Urey, H.C., Lowenstam, H.A., Epstein, S., and McKinney, C.R.: Measurements of paleotemperatures and temperatures of the Upper Cretaceous of England, Denmark, and the southeastern United States, *Geol. Soc. Am. Bull.*, 62, 399-416, 1951.
- van der Fliedert, T., Pahnke, K., and GEOTRACES intercalibration participants: GEOTRACES intercalibration of neodymium isotopes and rare earth element concentrations in seawater and suspended particles. Part 1: reproducibility of results for the international intercomparison, *Limnol. Oceanogr. Methods*, 10, 234-251, doi 10.4319/lom.2012.10.234, 2012.
- 565 van der Maaten, L. and Hinton, G.: Visualizing data using t-SNE, *J. Mach. Learn. Res.*, 9, 2579-2605, 2008.
- Yonge, C.M.: The absorption of glucose by *Ostrea edulis*, *J. Mar. Biol. Assoc. U. K.*, 15, 643-653, 1928.
- Zaky, A.H., Brand, U., and Azmy, K.: A new sample processing protocol for procuring seawater REE signatures in biogenic and abiogenic carbonates, *Chem. Geol.*, 416, 36-50, doi 10.1016/j.chemgeo.2015.10.015, 2015.
- Zaky, A.H., Azmy, K., Brand, U., and Svavarsson, J.: Rare earth elements in deep-water articulated brachiopods: An evaluation of seawater mass, *Chem. Geol.*, 435, 22-34, doi 10.1016/j.chemgeo.2016.04.016, 2016.
- 570

Group	Species	Locality of origin	Coordinates	Age	Annual temperature range	Annual salinity range (PSU)	Number of specimens	Number of measurements
Mod_BD V_Cgig	<i>C. gigas</i>	Géfosse, Baie des Veys (Normandy)	49°23.11 N, 01°06.05 W	Modern	5-20°C	30-34	5	30
Mod_BD V_Oedu	<i>O. edulis</i>	Géfosse, Baie des Veys (Normandy)	49°23.11 N, 01°06.05 W	Modern	5-20°C	30-34	5	41
Mod_M O_Cgig	<i>C. gigas</i>	Marennes-Oléron, Charente Maritime (Atlantic Ocean)	45°52.23 N, 01°10.60 W	Modern	5-26°C	24-38	3	24
Mod_M O_Oedu	<i>O. edulis</i>	Marennes-Oléron, Charente Maritime (Atlantic Ocean)	45°52.23 N, 01°10.60 W	Modern	5-26°C	24-38	3	13
Mod_TE S_Cgig	<i>C. gigas</i>	Tès, Arcachon (Atlantic Ocean)	44°40.01 N, 01°08.18 W	Modern	5-26°C	25-35	8	58
Mod_LE U_Cgig	<i>C. gigas</i>	Leucate, Aude (Mediterranean pond)	42°52.48 N, 03°01.50 W	Modern	2-32°C	26-42	5	14
Anc_CY B1_Oedu	<i>O. edulis</i>	Unknown	Unknown	20-30 CE	Unknown	Unknown	6	44
Anc_CY B2_Oedu	<i>O. edulis</i>	Unknown	Unknown	20-30 CE	Unknown	Unknown	7	30
Anc_MA L_Oedu	<i>O. edulis</i>	Unknown	Unknown	6 <sup>th</sup> c. CE	Unknown	Unknown	6	43

575 Table 1: Specimen groups and information on their respective localities. Temperature and salinity ranges at Marennes-Oléron, Tès and Baie des Veys are from Lartaud et al. (2010b), and at Leucate from Andrisoa (2019).




**Figure 1: Map of the localities of modern (squares) and archaeological (stars) specimens. Coordinates of the modern sites are indicated in Table 1.**



580


Figure 2: Typical archaeological *Ostrea edulis* (a) and modern *Crassostrea gigas* (b) specimens. The umbo region (c) is cut following the dashed white line. Laser ablation craters (200 μm in diameter) are indicated by circles (on c). Multiple measurements have been performed on each shell, in both winter and summer parts. Red and blue circles represent measurements corresponding to summer and winter periods, respectively, based on cathodoluminescence.



585

**Figure 3: Boxplots of Y/Ho ratios for all groups. The letters on top of the boxes (a and b) identify the significant differences between groups from Kruskal-Wallis tests. Note that for the same locality at Baie des Veys (BDV) and Marennes-Oléron (MO), the Y/Ho ratios are significantly different depending on the species considered (*C. gigas* and *O. edulis* groups). Grey bars represent median values, the lower and higher large black bars represent the 25<sup>th</sup> and 75<sup>th</sup> percentiles, respectively, and the lower and higher small black bars represent the minimum and maximum values not considered as outliers, respectively. Outliers are represented by grey crosses.**

590



**Figure 4:** Gradual decrease of REE abundances in oyster shells according to the REE atomic number, presented against Y. Values are expressed in cubic root of abundances (in  $\mu\text{g}\cdot\text{g}^{-1}$ ) to approach normality. Measurements from *C. gigas* and *O. edulis* are indicated with crosses and filled circles, respectively.




Figure 5: REE median profiles for all groups of oyster specimens. *Crassostrea gigas* groups are symbolized with dashed lines and continuous lines for *O. edulis* groups. Seawater profiles of the Atlantic Ocean (van der Fliedrt et al., 2012) and the Mediterranean Sea (Censi et al., 2004) are indicated for comparison. Values are normalized to Post-Archean Australian Shales (PAAS) according to McLennan (1989).

600






Figure 6: Visualization of shell group partitioning using t-SNE applied to all REE and Y measurements as variables. Crosses and large dots refer to *C. gigas* and *O. edulis*, respectively.