

1 **Distinguishing between early and late covering crops in the land surface model**
2 **Noah-MP: Impact on simulated surface energy fluxes and temperature**

3

4 Kristina Bohm^{ab}, Joachim Ingwersen^b, Josipa Milovac^c, Thilo Streck^b

5 ^a previously published under the name Kristina Imukova

6 ^b Institute of Soil Science and Land Evaluation, Department of Biogeophysics, University of
7 Hohenheim, 70593 Stuttgart, Germany

8 ^c Institute of Physics and Meteorology, University of Hohenheim, 70593 Stuttgart, Germany

9

10

11 Corresponding author: Kristina Bohm

12 E-Mail: imukovaks@gmail.com

13

14 Abstract

15 Land surface models are essential parts of climate and weather models. The widely used Noah-
16 MP land surface model requires information on the leaf area index (LAI) and green vegetation
17 fraction (GVF) as key inputs of its evapotranspiration scheme. The model aggregates all
18 agricultural areas into a land use class termed “Cropland and Pasture”. In a previous study we
19 showed that, on a regional scale, GVF has a bimodal distribution formed by two crop groups
20 differing in phenology and growth dynamics: early covering crops (ECC, ex.: winter wheat,
21 winter rapeseed, winter barley) and late covering crops (LCC, ex.: corn, silage maize, sugar
22 beet). That result can be generalized for Central Europe. The present study quantifies the effect
23 of splitting the land use class “Cropland and Pasture” of Noah-MP into ECC and LCC on surface
24 energy fluxes and temperature. We further studied the influence of increasing the LCC share,
25 which in the study area (the Kraichgau region, southwest Germany) is mainly the result of
26 heavily subsidized biomass production, on energy partitioning at the land surface. We used the
27 GVF dynamics derived from high-resolution (5 m x 5 m) RapidEye satellite data and measured
28 LAI data for the simulations. Our results confirm that GVF and LAI strongly influence the
29 partitioning of surface energy fluxes, resulting in pronounced differences between ECC and LCC
30 simulations. Splitting up the generic crop into ECC and LCC had the strongest effect on land
31 surface exchange processes in July-August. During this period, ECC are at the senescence
32 growth stage or already harvested, while LCC have a well-developed, ground-covering canopy.
33 The generic crop resulted in humid bias, i.e. an increase of evapotranspiration by $+0.5 \text{ mm d}^{-1}$
34 (LE: $1.3 \text{ MJ m}^{-2}\text{d}^{-1}$), decrease of **sensible heat flux** (H) by $1.2 \text{ MJ m}^{-2} \text{ d}^{-1}$ and decrease of surface
35 temperature by -1°C . The bias increased as the shares of ECC and LCC became similar. The
36 observed differences will impact the simulations of processes in the planetary boundary layer.
37 Increasing the LCC share from 28 to 38% in the Kraichgau region led to a decrease of **latent heat**
38 **flux** (LE) and a heating up of the land surface in the early growing season. Over the second part
39 of the season, LE increased and the land surface cooled down by up to 1°C .

40

41 **1 Introduction**

42 Within weather and climate models, land surface exchange processes are simulated by so-called
43 land surface models (LSMs). The main role of an LSM is to partition net radiation at the land
44 surface into sensible heat (H), latent heat (LE) and ground heat (G) fluxes and to determine the
45 land surface temperature. Surface energy partitioning has a significant influence on the evolution
46 of the Atmospheric Boundary Layer (ABL). ABL evolution strongly influences the initiation of
47 convection, cloud formation, and ultimately the location and strength of precipitation (Crawford
48 et al. 2001, Koster et al. 2006, Santanello Jr. et al. 2013, van Heerwaarden et al. 2009, Milovac et
49 al. 2016).

50
51 The surface energy partitioning depends on the physical and physiological properties of the land
52 surface (Raddatz 2007). In LSMs, the earth's surface is subdivided into different land use classes,
53 among them cropland. Physiological state variables of crops such as green vegetation fraction
54 (GVF) and leaf area index (LAI) vary significantly throughout the growing season. This alters the
55 biophysical parameters surface albedo, bulk canopy conductance, and roughness length, leading
56 to significant changes in surface energy fluxes (Crawford et al. 2001, Ghilain et al. 2012,
57 Tsvetsinskaya et al. 2001a, Wizemann et al. 2014). In many parts of the world, cropland covers a
58 considerable part of the simulation domain-area. Therefore, accurately simulating the seasonal
59 variability of surface energy fluxes highly depends on an adequate representation of plant growth
60 dynamics.

61
62 One of the widely used LSMs is Noah-MP. It is usually coupled with the Weather Research and
63 Forecasting (WRF) model, which is intended for use from the large eddy simulation (LES) scale
64 up to the global scale. Within each grid cell, Noah-MP computes net longwave radiation as well
65 as LE, H and G separately for the bare soil and the vegetated tile, whereas short-wave radiation is
66 computed over the entire grid cell (semi-tile approach; Lhomme and Chehbouni 1999, Niu et al.
67 2011).

68
69 Noah-MP collects agricultural areas into only general land use classes such as “Dryland Cropland
70 and Pasture”, “Irrigated Cropland and Pasture” or “Mixed Dryland/Irrigated Cropland and
71 Pasture” etc.. Vegetation dynamics and its seasonal development are described in the Noah-MP

72 model by the plant variables GVF and LAI. The surface energy fluxes critically depend on
73 accurately representing GVF and LAI dynamics (Chen and Xie 2011, Crawford et al. 2001,
74 Refslund et al. 2014). In Noah-MP, GVF and LAI are fixed quantities: they do not depend on the
75 weather conditions during a simulation. GVF is defined as the grid-cell fraction covered by a green
76 canopy (Gutman and Ignatov 1998). It is a function of the upper canopy (Rundquist 2002) and
77 represents the horizontal density of vegetation in each grid cell (Gutman and Ignatov 1998). LAI
78 represents the vertical density of the canopy. Certain biophysical parameters in Noah-MP such as
79 surface albedo, roughness and emissivity are considered linear functions of LAI.

80

81 By default, Noah-MP derives GVF values from the normalized difference vegetation index
82 (NDVI) obtained from the NESDIS/NOAA satellite. These data have a resolution of 15 km ×
83 15 km. Due to the mixing of croplands, forest and urban areas, the overall GVF is often positively
84 biased. Moreover, as shown by Imukova et al. (2015), seasonal GVF data are strongly smoothed
85 compared to the actual GVF dynamics. Milovac et al. (2016) and Nielsen et al. (2013) found that
86 the GVF grid data used in Noah-MP LSM are outdated and stated that these should be updated
87 given their importance for ABL evolution.

88

89 In a previous study, we derived GVF data with a resolution of 5 m x 5 m (Imukova et al. 2015) for
90 a region in southwest Germany (Kraichgau) using RapidEye satellite data. On the regional scale,
91 GVF shows a bimodal distribution mirroring the different phenology of crops. Crops could be
92 grouped into two classes. Early covering crops (ECC), such as winter wheat, winter rape, winter
93 barley and spring barley, develop early in spring, achieve maximum GVF usually between late
94 May and mid-June, and become senescent in July. Late covering crops (LCC), such as corn, silage
95 maize, and sugar beet, are drilled in spring and develop maximum ground-covering canopy from
96 July to August. They are still green in September, when the ECC are already harvested. The
97 dynamics of ECC and LCC vary to some degree from season to season and from region to region.

98

99 The shares of ECC and LCC may change over time, often reflecting economic decisions that may
100 depend on policy interventions. In Germany, a substantial change in these shares was introduced
101 by subsidizing biogas production. In 2005, 1.7 million ha of maize were cultivated in Germany.

102 Only 70,000 ha of this area were cropped with silage maize for biogas production (SRU Special
103 Report 2007). In 2009, the area cropped with maize for biogas production had increased to about
104 500,000 ha, while the total maize area remained almost constant (Huyghe et al. 2014). In 2012,
105 the total acreage of maize had increased to 2.57 million ha with 0.9 million ha intended for biogas
106 plants. The increase occurred mainly at the expense of grassland. Since then, the total maize crop
107 area has remained almost constant: 2.6 million ha in 2018 (Fachagentur Nachwachsende Rohstoffe
108 e. V. 2019). From 2005 to 2018, the maize area in Germany increased by about 53%.

109

110 The objectives of the present study were 1) to elucidate the extent to which surface energy fluxes
111 simulated with Noah-MP are affected by aggregating early and late covering crops into one generic
112 cropland class, and 2) to quantify the effect of a land use change, driven by the expansion of maize
113 cropping as a response to the increasing demand for biogas plants, on energy partitioning and
114 surface temperature in the Kraichgau region (southwest Germany). [Additionally, we tested the
115 performance of the Noah-MP on LE data measured with the Eddy Covariance technique.](#)

116

117 **2 Materials and methods**

118 **2.1 Study site and weather data measurements**

119 [The site under study is the agricultural field belonging to the farm “Katharinentalerhof”. The field
120 is located north of the city of Pforzheim \(48.92°N, 8.70°E\). The central research site is a part of
121 the Kraichgau region.](#) Kraichgau region covers about 1500 km². Mean annual temperature ranges
122 between 9-10° C and annual precipitation between 730 and 830 mm. The Neckar and Enz rivers
123 form the borders to the east. To the north and south, the region is bounded by the low mountain
124 ranges Odenwald and Black Forest. In the west, it adjoins the Upper Rhine Plain (Oberrheinisches
125 Tiefland). Kraichgau has a gently sloping landscape with elevations between 100 and 400 m above
126 sea level (a.s.l.). Soils predominantly formed from loess material. The region is intensively used
127 for agriculture: around 46 % of the total area is used for crop production. Winter wheat, winter
128 rapeseed, spring barley, corn, silage maize and sugar beet are the predominant crops.

129

130 Weather data used to force the Noah-MP model were acquired at an agricultural field (EC1, 14 ha)
131 belonging to the farm “Katharinentalerhof”. The terrain is flat (elevation a.s.l.: 319 m). The
132 predominant wind direction is south-west. The study site has been described in detail in several
133 studies (Imukova et al. 2015, Ingwersen et al. 2011, Wizemann et al. 2014).

134

135 An Eddy Covariance (EC) station was operated in the center of the EC1 field. Wind speed and
136 wind direction were measured with a 3D sonic anemometer (CSAT3, Campbell Scientific, UK)
137 installed at a height of 3.10 m. Downwelling longwave and downwelling shortwave radiation were
138 measured with a NR01 4-component sensor (NR01, Hukseflux Thermal Sensors, The
139 Netherlands). Air temperature and humidity were measured in 2 m height (HMP45C, Vaisala Inc.,
140 USA). All sensors recorded data in 30-min intervals. Rainfall was measured using a tipping bucket
141 (resolution: 0.2 mm per tip) rain gauge (ARG100, Campbell Scientific Ltd., UK). For further
142 details about instrumentation and data processing see Wizemann et al. (2014).

143

144 *2.2 Eddy Covariance measurements*

145 In order to test the Noah-MP performance, we used the EC measurements of latent heat flux over
146 maize (EC2) and winter wheat field (EC3) of 2012 growing season. EC2 and EC3 agricultural
147 fields are also belonging to the farm “Katharinentalerhof” introduced above. They are 23 ha and
148 15 ha large. The winter wheat was planted in autumn 2011 and harvested on 29th of July. The
149 maize was drilled on 2nd of May and harvested on 20th of September. The EC station was
150 operated in the center of each field. The latent heat flux was measured in a 30-min resolution.
151 For the maize, the LE data was only available till 20th of September, whereas for the winter
152 wheat field there were no missing data. A detailed information on the EC measurements is given
153 in Imukova et al. (2016). The EC flux data were processed with the TK3.1 software (Mauder M.,
154 2011). Surface energy fluxes were computed from 30-min covariances. For data quality analysis
155 we used the flag system after Foken (Mauder M., 2011). LE half-hourly values with flags from 1
156 to 6 (high and moderate quality data) were used to test the performance of the Noah-MP LSM.
157 LE data was gap filled using the mean diurnal variation method with an averaging window of 14
158 days (Falge et al., 2001). The random error of the LE flux, which consist of the instrumental
159 noise error of the EC station and the sampling error was computed by the TK3.1 software

160 (Mauder et al., 2013). For more details on EC data processing, please refer to Imukova et al.
161 (2016).

162
163 The model performance is usually tested against field measurements of sensible and latent heat
164 flux performed with Eddy Covariance (EC) technique (Ingwersen et al., 2011; El Maayar et al.,
165 2008; Falge et al., 2005). EC method is widely used method for this purpose although it has one
166 well-known problem. The energy balance of EC flux data is typically not closed, which means LE
167 and/or H fluxes measured with EC technique are most probably underestimated. Previous study
168 showed the EC technique provides reliable LE measurements at our study site and these data can
169 be used for model testing (Imukova et al. 2016).

170

171 **2.3 The Noah-MP v1.1 land surface model**

172 *2.3.1 Model parameterization*

173 Multi-physics options of Noah-MP were set as shown in the Table 1. For the simulation we used
174 the USGS land use dataset. The vegetation type index was set to 2 (Dryland cropland and
175 Pasture) and soil type index to 4 (Silt loam). The model was forced with half-hourly weather data
176 (wind speed, wind direction, temperature, humidity, pressure, precipitation, downwelling
177 longwave and shortwave radiation) measured at EC1 from 2011 to 2012. Simulations were
178 initialized with a spin up period of one year (2011) and run with a time step of 1800 seconds.

179 **Table 1.** Setting of the multi-physics options used in the Noah-MP simulation.

Multi-physics option	Setting
Vegetation model	opt_dveg = 1: prescribed [table LAI, shdfac=FVEG]
Canopy stomatal resistance	opt_crs = 2: Jarvis
Soil moisture factor for stomatal resistance	opt_btr = 1: Noah
Runoff and groundwater model	opt_run = 1: SIMGM
Surface layer drag coefficient (CH & CM)	opt_sfc = 1: based on Monin-Obukhov similarity theory
Supercooled liquid water	opt_frz = 1: NY06
Frozen soil permeability	opt_inf = 1: NY06
Radiation transfer	opt_rad = 3: gap=1—Fveg
snow surface albedo	opt_alb = 2: CLASS
rainfall & snowfall	opt_snf = 1: Jordan91
lower boundary of soil temperature	opt_tbot = 2: Noah
snow/soil temperature time scheme	opt_stc = 1: Semi-implicit

180

181

182 *2.3.2 GVF dynamics*

183 The GVF data required by the Noah-MP model were derived from high-resolution (5 m x 5 m)
 184 RapidEye satellite data. [A detailed information on the deriving of the GVF data used in the current](#)
 185 [research can be found in Imukova et al. \(2015\)](#). The GVF data were calculated from the
 186 Normalized Difference Vegetation Index (NDVI) computed from the red and near-infrared bands
 187 of the satellite images. The relationship between GVF and NDVI was established by linear
 188 regression using ground truth measurements. GVF maps [for the Kraichgau region](#) were derived in
 189 a monthly resolution.

190 **Table 2.** GVF dynamics of early covering crops (ECC) and late covering crops (LCC) in 2012 and 2013
 191 in the Kraichgau region, southwest Germany as well as the GVF dynamics of the generic crop.

GVF		15 Apr	15 May	15 Jun	15 Jul	15 Aug	15 Sep
GVF 2012	ECC	- ^b	0.74	0.83	0.37	0.01 ^c	0.01
	LCC	- ^b	0.01	0.35	0.74	0.69 ^c	0.56
GVF 2013	ECC	0.54	0.80	0.57 ^c	0.29	0.01	0.01
	LCC	0.01	0.06	0.37 ^c	0.69	0.74	0.75
Mean GVF	ECC	0.54	0.77	0.70	0.33	0.01	0.01
	LCC	0.01	0.04	0.36	0.72	0.72	0.66
Generic crop GVF^a		0.39	0.57	0.60	0.44	0.21	0.19

^a Weighted mean GVF calculated based on fractions of ECC (72%) and LCC (28%) in Kraichgau


^b No RapidEye scenes were available for April

^c No RapidEye scenes were available for these months, GVF values were derived by linear interpolation between adjacent months

192

193 Table 2 shows the observed and mean GVF dynamics of ECC and LCC over the growing seasons
 194 2012 and 2013 as well as the GVF dynamics of the generic crop [in the Kraichgau region](#). The GVF
 195 values on the 15th day of each month, as required by Noah-MP model, were calculated by linearly
 196 interpolating the monthly values derived from the GVF maps. A generic GVF dynamics was
 197 calculated as the weighted mean of ECC and LCC from 2012 and 2013. The areal distribution of
 198 ECC and LCC was determined from the GVF maps of May. All pixels with a GVF value below
 199 0.5 were counted as LCC, whereas pixels with values above that threshold were assigned to ECC.
 200 [Figure 1 shows the spatial distribution of early and late covering crops in Kraichgau](#). The estimated
 201 areal distribution of ECC and LCC was 72% and 28%, respectively. These results correspond well
 202 with data of the Statistisches Landesamt Baden-Württemberg ([http://www.statistik.baden-
 204 wuerttemberg.de/](http://www.statistik.baden-

 203 wuerttemberg.de/)).



205

206 **Figure 1.** Map of early covering (ECC) and late covering crops (LCC) in Kraichgau region, Southwestern
207 Germany.

208 **2.3.3 LAI dynamics**

209 Noah-MP requires prescribed LAI data for each month. Data were derived from field
 210 measurements. LAI was measured biweekly using a LAI-2000 Plant Canopy Analyzer (LI-COR
 211 Biosciences Inc., USA). In 2012 and 2013, LAI of the crops was measured on five permanently
 212 marked plots of 1 m² on three different fields. Detailed information about the study plots can be
 213 found in Imukova et al. (2015). In 2009-2011, LAI and the phenological development of the crops
 214 were measured on five permanently marked plots of 4 m² in the same three fields. The growth
 215 stages of crops were determined using the BBCH scale (Meier et al. 2009). More details on the
 216 measurements can be found in Ingwersen et al. (2011) and Ingwersen et al. (2015). Table 3 shows
 217 measured and mean LAI dynamics as well as generic LAI dynamics estimated considering shares
 218 of ECC (72%) and LCC (28%) in the study region. LAI dynamics of winter wheat and winter rape
 219 were assigned to ECC, those of maize to LCC. Mean LAI dynamic of ECC was estimated based
 220 on the measurements conducted in winter wheat and winter rape stands during the 2012 and 2013
 221 growing seasons. Since LAI data were not available for maize in 2013, the mean LAI dynamic of
 222 LCC were assessed using field data from the same fields collected in 2009-2012.

223 **Table 3.** LAI dynamics of early covering crops (ECC) and late covering crops (LCC) in 2012 and 2013 in
 224 the Kraichgau region, southwest Germany, as well as the LAI dynamics of the generic crop.

Green LAI		15 Apr	15 May	15 Jun	15 Jul	15 Aug	15 Sep
LAI 2012	ECC	2.4	4.4	4.6	0.0	0.0	0.0
	LCC	0.0	0.1	0.9	3.2	5.0	3.7
LAI 2013	ECC	1.7	4.2	4.3	0.0	0.0	0.0
	LCC ^b	-	-	-	-	-	-
Mean LAI	ECC	2.1	4.3	4.5	0.0	0.0	0.0
	LCC^c	0.0	0.1	0.9	3.1	4.5	3.8
Generic crop LAI^a		1.5	3.1	3.5	0.9	1.3	1.1

^a Weighted mean LAI calculated based on fractions of ECC (72%) and LCC (28%) in Kraichgau

^b LAI data for maize in 2013 were not measured

^c Since LAI data for maize in 2013 were not available, LAI dynamics were derived from the field data of 2009-2012 for maize in the Kraichgau region

225

226

227 **2.4 Simulation runs**

228 We firstly quantified the extent to which ECC and LCC differ with regard to their energy and water
229 fluxes, surface (TS) and soil temperature (TG). For this, we performed one local simulation for
230 each crop group using the mean LAI and the mean GVF dynamics observed during the two
231 growing seasons (see Table 2 and Table 3).

232

233 Secondly, to determine the effect of splitting up the vegetation dynamics of a generic crop into
234 that of ECC and LCC, we compared the following two local simulation runs:

235 Run 1: Noah-MP was forced with the GVF and LAI dynamics of the generic crop (Table 2 and
236 Table 3). Accordingly, in this simulation, we first computed the weighted mean of the vegetation
237 properties (GVF and LAI), and subsequently simulated the surface energy fluxes, TS and TG.

238 Run 2: We first simulated the energy and water fluxes [separately](#) for ECC and LCC with their
239 crop-specific vegetation dynamics. [Afterwards, we calculated the weighted averages of the
240 simulated fluxes and temperatures based on the share of early covering \(72%\) and late covering
241 crops \(28%\) in Kraichgau.](#)

242

243 Thirdly, we studied the effect of increasing the LCC share on the surface energy fluxes, surface
244 and soil temperatures. As mentioned in the Introduction, the maize cropping area in Germany
245 increased by 53% over the last decade. In our study region, this increase corresponds to a rise of
246 the LCC share from 28% to 38%. To study the effect of this land use change on the Noah-MP
247 simulations, we performed one additional generic crop simulation, but this time the generic crop
248 dynamics was computed with a LCC share of 38%.

249

250 [2.5 Statistical analysis](#)

251 [The model performance was evaluated based on the model efficiency \(EF\), root mean square error
252 \(RMSE\) and bias. EF is defined as the proportion of the total variance explained by a model:](#)

$$EF = 1 - \frac{\sum_{i=1}^N (P_i - O_i)^2}{\sum_{i=1}^N (O_i - \bar{O})^2}, \quad \text{Eq. 1}$$

253 where P_i denotes predicted values, O_i and \bar{O} – observed values and their mean, respectively, while
254 N is the number of observations. RMSE and bias were calculated as

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2} \quad \text{Eq. 2}$$

255 and

$$bias = \frac{1}{N} \sum_{i=1}^N (P_i - O_i). \quad \text{Eq. 3}$$

256

257

258 **3 Results**

259 **3.1 ECC vs. LCC**

260 Over the growing season, ECC and LCC show distinct differences with regard to energy
261 partitioning at the land surface (Figure 2). The observed shifts were strongest for LE and H. Early
262 covering crops already reached their maximum LE flux in May, after which LE declined during
263 the growing season. In contrast, LCC showed a continued increase in LE over the season, peaking
264 three months later in August. The smallest difference in evapotranspiration between both crops
265 types was on average 0.4 mm day^{-1} (LE $0.9 \text{ MJ m}^{-2}\text{day}^{-1}$) in June, while the largest mean deviation
266 of -2.3 mm day^{-1} (LE $-5.7 \text{ MJ m}^{-2}\text{day}^{-1}$) occurred in August (Table 4). With regard to the H flux,
267 the situation was opposite (Figure 2). In the case of ECC, H flux increased continuously over the
268 course of the growing season, peaking in August. In contrast, LCC already reached the H
269 maximum in May. Afterwards, H decreased continuously until late August. As for LE, the smallest
270 ($-1.2 \text{ MJ m}^{-2}\text{day}^{-1}$) and largest ($5.3 \text{ MJ m}^{-2}\text{day}^{-1}$) mean differences in H between ECC and LCC
271 were observed in June and August, respectively (Table 4). Compared with LCC, the higher latent
272 heat fluxes of ECC in May and June resulted in a cooler land surface, on average by -2.6°C and -
273 1.0°C , respectively (Table 4). From July to August the situation was reversed: because latent heat
274 fluxes of ECC are distinctly lower than that of LCC, the surface temperature at ECC sites was up
275 to 4°C warmer than at LCC sites (Figure 3).

276


277 The mean difference in daily ground heat flux between ECC and LCC during the growing season
 278 ranged between -0.2 MJ m^{-2} and 0.2 MJ m^{-2} (Table 4). Also for the ground heat flux, the smallest
 279 difference between both crops types was observed in June (0.05 MJ m^{-2}).

280 **Table 4.** Mean differences (ECC minus LCC) in latent (LE), sensible (H) and ground heat (G) fluxes,
 281 mean surface temperature (TS) and mean ground temperature (TG) between ECC and LCC simulations.


Month	DOY	LE mm d^{-1}	$\text{MJ m}^{-2} \text{ d}^{-1}$	H $\text{MJ m}^{-2} \text{ d}^{-1}$	G $\text{MJ m}^{-2} \text{ d}^{-1}$	TS $^{\circ}\text{C}$	TG $^{\circ}\text{C}$
May	121 – 151	1.3	3.3	-3.1	-0.2	-2.6	-2.2
June	152 – 181	0.4	0.9	-1.2	0.05	-1.0	-0.9
July	182 – 212	-1.5	-3.8	3.3	0.2	2.1	1.8
August	213 – 243	-2.3	-5.7	5.3	0.1	3.2	2.4
September	244 – 273	-0.7	-1.8	2.1	-0.1	1.9	1.2

DOY - day of a year


282



283



284



285


286

287

288

Figure 2. Simulation results of Noah-MP LSM for latent (LE), sensible (H) and ground heat (G) flux. Simulations were performed for two types of crops: early covering (solid line) and late covering (dashed line).

289
290
291
292
293



294
295
296
297
298
299

Figure 3. Differences (ECC minus LCC) in latent (LE), sensible (H) and ground heat (G) fluxes, mean surface temperature (TS) and mean ground temperature (TG) between simulations for ECC and LCC.

300 **3.2 Noah-MP vs. Eddy Covariance measurements**

301 The average random error of the latent heat flux measured with EC technique for the entire growing
 302 season was about 25% over the winter wheat field and about 21% over the maize field.

303
 304 The simulated latent heat flux based on ECC and LCC parametrization agreed fairly well with the
 305 Eddy Covariance data (Table 5-6, Figure 4-5). The model efficiency over the entire simulation
 306 period was 0.87 for ECC and 0.90 for LCC. The best agreement between the observations and the
 307 Noah-MP LSM using crop-type-specific sets was achieved for winter wheat in June and for maize
 308 in August and September. The generic crop parametrization showed less satisfying modeling
 309 results, particularly for the maize field (Table 5-6). For the entire growing season, EF was 0.78 for
 310 winter wheat and only 0.57 for maize. Over the winter wheat field, LE was overestimated.
 311 Overestimation of the LE was highest in in July and August. Over the maize field, LE was
 312 overestimated in May and June and underestimated in July, August and September. Particularly in
 313 May and August, the bias increased to 68.8 Wm⁻² and -56 Wm⁻², respectively. The best model
 314 performance using generic crop set was achieved for the winter in June wheat and for the maize in
 315 July.

316 **Table 5.** Root mean square error (RMSE), bias and modeling efficiency (EF) of the latent heat flux for
 317 the simulation runs for winter wheat stand (EC3 field).


Variant	May	June	July	August	September	Overall
RMSE (Wm ⁻²)						
ECC	45.4	35.4	33.0	26.3	13.5	32.5
Generic crop	36.3	33.0	59.6	63.6	20.9	45.7
Bias (Wm ⁻²)						
ECC	27.3	17.9	14.2	17.1	0.8	15.5
Generic crop	20.5	15.2	33.9	41.7	7.7	23.8
EF (1)						
ECC	0.88	0.91	0.80	0.74	0.89	0.87
Generic crop	0.91	0.92	0.62	0.41	0.85	0.78

318

319 **Table 6.** Root mean square error (RMSE), bias and modeling efficiency (EF) of the latent heat flux for
 320 the simulation runs for maize stand (EC2 field).


Variant	May	June	July	August	September	Overall
RMSE (Wm⁻²)						
LCC	53.1	37.3	31.8	28.1	18.9	35.7
Generic crop	102.0	50.9	29.8	85.8	43.7	68.0
Bias (Wm⁻²)						
LCC	37.4	21.5	13.7	-14.9	-2.5	11.0
Generic crop	68.6	29.9	-10.6	-56.0	-22.9	1.8
EF (1)						
LCC	0.59	0.87	0.94	0.96	0.96	0.90
Generic crop	0.30	0.80	0.91	0.12	0.77	0.57

321



322

323 **Figure 4.** Monthly averaged measured and simulated diurnal latent heat fluxes (LE) for May to September. The Noah-MP LSM was run with two
 324 different vegetation parametrizations: early covering crops (ECC) and generic crop.



325

326 **Figure 5.** Monthly averaged measured and simulated diurnal latent heat fluxes (LE) for May to September. The Noah-MP LSM was run with two
 327 different vegetation parametrizations: late covering crops (LCC) and generic crop.

328 **3.3 Run 1 vs. Run 2 (Generic crop vs. weighted mean of ECC and LCC)**

329 The generic crop simulation run (Run 1) generally yielded higher LE fluxes than Run 2 (i.e.
 330 splitting up the generic crop into ECC and LCC) (Figure 6). During the growing season the mean
 331 difference in evapotranspiration between two runs was 0.1 mm day⁻¹ (LE 3.7 MJ m⁻²day⁻¹) (Table
 332 5). Smallest mean monthly differences occurred in June and September: 0.02 mm day⁻¹ (LE 0.4 MJ
 333 m⁻²day⁻¹) and 0.03 mm day⁻¹ (LE 1 MJ m⁻²day⁻¹), respectively. The most pronounced differences
 334 in LE flux were recorded in late July (DOY 197-208) (Figure 7). The average difference in half-
 335 hourly fluxes over this period, between 9 a.m. and 6 p.m, was 36 W m⁻², and the highest half-
 336 hourly deviation between both runs was 83 W m⁻² (Figure 7). The highest daily deviation was 0.8
 337 mm day⁻¹ (Figure 6). Over the whole season, the cumulative difference in evapotranspiration
 338 between two runs was 20 mm, leading to a 16 percent lower seasonal water balance (SWB) in Run
 339 1 (SWB: -133 mm) than in Run 2 (SWB: -113 mm).

340 **Table 5.** Mean differences in latent (LE), sensible (H) and ground heat (G) fluxes, surface
 341 temperature (TS) and ground temperature (TG) between Run 1 and Run 2 simulations. Numbers
 342 in brackets: the relative difference between Run 1 and Run 2 simulations in percentage.

Month	DOY	LE mm d ⁻¹	MJ m ⁻² d ⁻¹	H MJ m ⁻² d ⁻¹	G MJ m ⁻² d ⁻¹	TS °C	TG °C
May	121 – 151	0.1 (3)	0.3	-0.3 (19)	-0.003 (1)	-0.3 (2)	-0.02 (0.1)
June	152 – 181	0.02 (0.4)	0.04	-0.1 (4)	0.001 (1)	-0.1 (1)	0.01 (0.05)
July	182 – 212	0.3 (7)	0.6	-0.6 (21)	-0.016 (4)	-0.4 (2)	-0.1 (0.6)
July*	197 – 208	0.5 (14)	1.3	-1.2 (46)	-0.034 (10)	-1.0 (4)	-0.2 (1)
August	213 – 243	0.2 (7)	0.5	-0.6 (18)	0.004 (2)	-0.3 (1)	0.01 (0.03)
September	244 – 273	0.03 (1)	0.1	-0.2 (5)	0.005 (3)	-0.1 (1)	0.1 (0.4)
Mean		0.1 (3.7)	0.3	-0.4 (13.2)	-0.002 (1)	-0.2 (1.4)	-0.01 (0.1)

DOY - day of a year

343
 344
 345 In contrast, H fluxes of Run 1 were mostly lower over all months than those simulated in Run 2
 346 (Figure 6). From May to September, the mean difference in H fluxes was about -0.4 MJ m⁻² (-
 347 13 %) (Table 5). The smallest difference occurred again in June, the largest difference again in
 348 late July (Figure 7). During DOY 197-208 the mean differences in half hourly H fluxes was about
 349 -29 W m⁻², the peak deviation being -72 W m⁻² (9 a.m.-6 p.m) (Figure 7). Cumulating these

350 differences over the day reduced the production of sensible heat on average in the order of
351 1.2 MJ m^{-2} , corresponding to a 46 %reduction compared to Run 2 (Table 5). Ground heat fluxes
352 as well as soil temperature were affected only moderately by the different vegetation
353 parameterization of Run 1 and 2 (Figure 7, Figure 6). As for LE and H, the largest mean differences
354 in G fluxes were observed during DOY 197-208 ($-0.034 \text{ MJ m}^{-2} = 10\%$) (Table 5).

355

356 Due to the humid bias of Run 1, the canopy surface was cooler than in Run 2 in all months. On
357 average, TS of Run 1 was $0.2 \text{ }^{\circ}\text{C}$ ($\sim 1.4\%$) lower during the growing season than in Run 2. In late
358 July (DOY 197-208) the mean daily difference was $-1 \text{ }^{\circ}\text{C}$ (Table 5, Figure 6) and reached a daytime
359 (9a.m.-6p.m.) peak difference of up to $-2.6 \text{ }^{\circ}\text{C}$ (Figure 7).

360




Figure 6. Differences in latent (LE), sensible (H) and ground heat (G) fluxes, mean surface temperature (TS) and mean ground temperature (TG) between Run 1 and Run 2 simulations (*Run 1 - Run 2*). Given percentages are relative differences between Run 1 and Run 2 simulations.

361
362
363
364




Figure 7. Differences in latent (LE), sensible (H) and ground heat (G) fluxes, mean surface temperature (TS) and mean ground temperature (TG) between Run 1 (generic crop) and Run 2 (weighted mean of early and late covering crops) simulations (*Run 1 - Run 2*).

366 **3.4 Land use change towards LCC**

367 Increasing the LCC fraction from 28% to 38% mainly led to changes in LE and H fluxes (Table
 368 6). That LCC increase lowered the LE flux (-0.3 MJ m⁻² day⁻¹-or ET 0.1 mm day⁻¹) early in the
 369 season. This was accompanied by a higher H flux (+0.3 MJ m⁻² day⁻¹), which in turn led to a 0.3
 370 °C warmer surface temperature than for the runs with the actual ECC-LCC ratio. From July to
 371 September, increasing the LCC fraction boosted evapotranspiration by about 0.2 mm day⁻¹ (LE
 372 0.4 MJ m⁻² day⁻¹) and decreased the H flux by about 0.3 MJ m⁻² day⁻¹ (Table 6). The largest half-
 373 hourly differences occurred in August (DOY 213-243, Figure 8), amounting to +40 W m⁻²
 374 and -30 W m⁻² for LE and H, respectively. The smallest deviations for both fluxes were recorded
 375 in June. Over the July–September period, the higher LE flux of the simulation run with the
 376 increased LCC fraction cooled the land surface up to -1 °C (Figure 8). In general over the growing
 377 season, increasing the LCC share by 10% led to an increase in cumulative evapotranspiration,
 378 which in turn resulted in a 10 mm lower seasonal water balance (SWB: -143 mm).

379
 380 With regard to the ground heat flux, increasing the LCC fraction led to an up to 10 W m⁻² higher
 381 flux over the noon time during the second part of the growing season (Figure 8), whereas early in
 382 the season the differences did not exceed 0.2°C (Table 6).

383 **Table 6.** Mean differences in latent (LE), sensible (H) and ground heat (G) fluxes, surface temperature
 384 (TS) and ground temperature (TG) between simulations with the LCC fraction increased by 10 % and the
 385 baseline simulation (*increased LCC share minus baseline simulation*). Numbers in brackets: the relative
 386 difference between *increased LCC share and baseline simulation* in percentage

Month	DOY	LE mm d ⁻¹	MJ m ⁻² d ⁻¹	H MJ m ⁻² d ⁻¹	G MJ m ⁻² d ⁻¹	TS °C	TG °C
May	121 – 151	-0.1 (3.3)	-0.3	0.3 (14)	0.02 (1)	0.3 (2)	0.2 (1)
June	152 – 181	-0.04 (1.0)	-0.1	0.1 (6)	-0.005 (0.5)	0.1 (1)	0.1 (1)
July	182 – 212	0.2 (4.3)	0.4	-0.3 (12)	-0.02 (6)	-0.2 (1)	-0.2 (1)
August	213 – 243	0.2 (7.6)	0.6	-0.5 (17)	-0.01 (1)	-0.3 (2)	-0.2 (1)
September	244 – 273	0.1 (3.8)	0.2	-0.2 (4)	0.01 (4)	-0.2 (1)	-0.1 (1)

DOY - day of a year

387

388




Figure 8. Impact of increasing the LCC fraction from 28% to 38% on latent (LE), sensible (H) and ground heat (G) fluxes, surface temperature (TS) and ground temperature (TG) (*Increased LCC share minus baseline simulation*).

390 4 Discussion

391 The comparison of the ECC and LCC simulations confirmed that GVF and LAI significantly affect
392 the partitioning of surface energy fluxes. LE flux increases with crop growth and peaks when the
393 canopy is fully developed, i.e. have maximum LAI and GVF. By contrast, the highest H and G
394 fluxes were observed at sparsely covered fields or on the fields with a senescent canopy. During
395 the main growth period of crops, H and G fluxes were quite low. ECC and LCC crops vary
396 significantly in sowing and harvest date, leaf area and senescence dynamics, water use efficiency
397 and phenology. Their surface energy fluxes therefore differ distinctly. Our simulation results are
398 in agreement with experimental data of Wizeman et al. (2014) as well as with modeling studies of
399 Sulis et al. (2015), Tsvetsinskaya et al. (2001b), Xue et al. (1996) or Ingwersen et al. (2018).

400
401 Simulation results based on ECC and LCC parametrization are in complete harmony with the field
402 observations at our study site. The performance test of the Noah-MP on the EC data showed the
403 crop-type-specific sets significantly improve the simulation of latent heat flux at the field scale. In
404 contrast, generic crop parametrization showed less satisfying modeling results. In general, it
405 performed better for winter wheat stand than for maize. Based on the generic crop set, simulation
406 results tend to greatly overestimate the latent heat flux for maize in the beginning of the growing
407 season when the plants are small. In August and September, the latent heat flux was in contrast
408 distinctly underestimated, during this period the maize canopy is fully developed. For wheat,
409 model overestimates the latent heat flux, particularly during July – September period, when the
410 winter wheat stand ripened and senescence or harvested.

411
412 Besides the vegetation dynamics, the simulated energy and water fluxes depend on additional
413 model settings as well. Ingwersen et al. (2011) performed a sensitivity study with the Noah model
414 for our study site. He found that among the vegetation parameters the minimum stomatal resistance
415 (RS) and a parameter used in the radiation stress function of the Jarvis scheme (RGL) are the most
416 sensitive parameters. Using constant RS, as it is implemented in Noah, results in the
417 underestimation of sensible heat flux and overestimation of latent heat flux during the ripening
418 stage of the cereals. Considering a monthly varying RS helped to distinctly improve the simulation
419 of the energy and water fluxes at the land surface. Ingwersen et al. (2010) concluded, integrating
420 the crop growth model which delivers daily RS, LAI and GVF values into Noah would greatly

421 enhance the overall performance of the land surface model. Among the soil parameters, the most
422 sensitive parameters are the soil moisture threshold where transpiration begins to stress
423 (REFSMC), maximum soil moisture content (MAXSMC) and soil moisture threshold where direct
424 evaporation from the top layer ends (DRYSMC). Considering these parameters has also a potential
425 to further improve of simulation results.

426
427 The potential increase of the LCC fraction (driven by the high demand for biogas and forage
428 production) leads to significant changes in the partitioning of the energy fluxes at the croplands.
429 In recent years the total area under maize in Germany has more than doubled. This corresponds to
430 an approximately 10% increase of the LCC fraction for the study region. In the early vegetation
431 period, the altered ECC-LCC ratio leads to a decrease of evapotranspiration, an increase of H
432 fluxes, and a warmer cropland surface because, during that period, a higher fraction of fields is
433 bare or sparsely covered with vegetation. In mid-June, the situation reverses. The higher share of
434 LCC boosts LE fluxes, decreases H fluxes and lowers surface temperatures. The increased
435 evapotranspiration over the growing season, in turn, leads to a lower seasonal water balance.

436
437 Comparing the generic crop simulation (Run 1) with the weighted mean of two separate
438 simulations for ECC and LCC (Run 2) showed the largest difference over the second half of the
439 growing season, particularly during late July/early August. In July, ECC become senescent: GVF
440 drops sharply and green LAI equals zero. In early August, ECC are usually harvested. In contrast,
441 LCC have a developed ground-covering canopy during July-August. Leaves of these crops are still
442 green in September. This transition period is very smooth in the case of the generic crop, resulting
443 on average in about 14 % higher LE and in about 46%, 10% and 4% lower H, G and surface
444 temperature, respectively, compared with Run 2.


445
446 The results presented above apply to the ECC-LCC ratio within our study area. What can we expect
447 in agricultural landscapes with different ECC-LCC ratios? The ECC-LCC ratio has nearly no effect
448 on energy partitioning in June, whereas in May, July and August its influence on the turbulent
449 fluxes is pronounced (Figure 9). The weak effect in June is because, during this period, the LAI
450 and GVF of ECC and LCC are similar (Figure 11). In the other months, however, the ECC-LCC
451 ratio heavily affects the energy partitioning. For example, increasing the LCC share from 10% to

452 90% boosts daily evapotranspiration in August from 2.5 mm d⁻¹ to 4.3 mm d⁻¹, decreases the H
 453 flux by about 4.1 MJ m⁻²d⁻¹ and cools down the cropland surface by 2 °C. Over the growing season,
 454 the increase in the LCC share leads to a general increase in evapotranspiration, which in turn
 455 lowers the seasonal water balance (Table 7). Moreover, different ECC-LCC ratios will also affect
 456 the above-mentioned humid bias of the generic crop parameterization (Figure 10). The bias is
 457 largest if ECC and LCC shares are balanced (ECC 50% and LCC 50 %), whereas combinations
 458 with one predominant crop distinctly lower the bias. In August, for instance, the LE differences
 459 between the two runs with ECC 50%- LCC 50% equal 0.27 mm day⁻¹, while ECC 10%- LCC 90
 460 % yields differences of 0.09 mm day⁻¹.

461 **Table 7.** Weather data and simulation results of Noah-MP LSM for cumulative evapotranspiration for the
 462 Kraichgau region. Simulations were performed considering different shares of early covering crops
 463 (ECC) and late covering crops (LCC).


ECC and LCC shares	Total rainfall (R), mm	Cumulative evapotranspiration (ET), mm	Seasonal water balance (R-ET), mm
ECC 90% LCC 10%	388	496	-108
ECC 70% LCC 30%	388	522	-134
ECC 50% LCC 50%	388	544	-156
ECC 30% LCC 70%	388	557	-169
ECC 10% LCC 90%	388	563	-175

464




465

466 **Figure 9.** Simulation results of Noah-MP LSM for latent (LE) and sensible (H) heat flux. Simulations
 467 were performed considering different shares of ECC and LCC.
 468



469
 470 **Figure 10.** Differences in latent (LE) and sensible heat (H) fluxes between Run 1 and Run 2 simulations
 471 (*Run 1 - Run 2*). Simulations were performed considering different shares of ECC and LCC.
 472



473
 474 **Figure 11.** GVF and LAI dynamics of early covering crops (ECC), late covering crops (LCC) and
 475 Cropland.
 476

477 Our results show that performing simulations based on single dynamics for each type of crop (ECC
 478 and LCC) improve simulations of surface fluxes during transition periods and at the end of the
 479 growing season. Lumping ECC and LCC into one land-use class (Croplands and Pasture), as done
 480 in Noah-MP, is an oversimplification. Several authors demonstrated the necessity to distinguish
 481 biophysical plant parameters between substantially different crops to obtain representative
 482 simulation results in the lower atmosphere (Sulis et al. 2015, Tsvetsinskaya et al. 2001b, Xue et
 483 al. 1996). They showed that high-resolution spatial information on various croplands and
 484 associated physiological characterizations can significantly improve the simulations of land
 485 surface energy fluxes, leading to better weather and climate predictions.

487 Changes of LAI and GVF with plant growth lead to changes in surface albedo, bulk canopy
488 conductance and roughness length, which in turn alter the partitioning of surface energy fluxes
489 (Chen and Xie 2011, Chen and Xie 2012, Crawford et al. 2001, Tsvetsinskaya et al. 2001a, Xue et
490 al. 1996). Such altered energy partitioning at the land surface then changes the thermodynamic
491 state of the atmospheric boundary layer with regard to air temperature, surface vapor pressure,
492 relative humidity and finally rainfall (Chen and Xie 2012, McPherson and Stensrud 2005, Sulis et
493 al. 2015, Tsvetsinskaya et al. 2001b). The observed differences between Run 1 and crop-type-
494 based runs will most probably influence the simulated processes in the ABL. For instance, Sulis
495 et al. (2015) significantly improved the simulations of land surface energy fluxes by using the
496 crop-specific physiological characteristics of the plant. They observed a difference of about 40%
497 between simulated fluxes using the generic and crop-specific parameter sets. The differences in
498 the land surface energy partitioning led to different heat and moisture budgets of the atmospheric
499 boundary layer for the generic and specific (sugar beet and winter wheat) croplands. In the case of
500 specific croplands, particularly sugar beet, those authors observed a larger contribution of the
501 entrainment zone to the heat budget of the ABL as well as a shallower ABL.

502

503 McPherson and Stensrud (2005) examined the impact of directly substituting the tallgrass prairie
504 land use class with winter wheat on the formation of the ABL. These crops have different growing
505 seasons. In the U.S. Great Plains, native prairie tallgrass mainly grows in summer, while winter
506 wheat grows throughout winter and reaches maturity in late spring. Simulations showed a larger
507 LE and lower H over the area with the winter wheat stand in comparison with tallgrass. By 2100
508 UTC, LE ranged from 300 to 400 $W m^{-2}$ for the wheat run and from 200 to 275 $W m^{-2}$ for the
509 tallgrass run. H ranged from 25 to 125 $W m^{-2}$ for the former and from 100 to 200 $W m^{-2}$ for the
510 latter. Substituting tallgrass prairie with winter wheat boosted the atmospheric moisture near the
511 surface above- and downstream of the study area, and resulted in a shallower ABL above- and
512 downstream of this area. The shallower ABL reduced the entrainment of higher-momentum air
513 into the ABL and therefore led to weaker winds within the ABL.

514

515 Milovac et al. (2016) performed six simulations at 2 km resolution with two local and two nonlocal
516 ABL schemes combined with two LSMs (Noah and Noah-MP) to study the influence of energy
517 partitioning at the land surface on the ABL evolution on a diurnal scale. They observed that LE

518 simulated by Noah-MP was more than 50% lower than that simulated by Noah. As expected, a
519 lower LE resulted in a drier ABL. The ABL evolution and its features strongly influence the
520 initiation of convection and cloud formation as well as the location and strength of precipitation.
521 For instance, drier and higher ABL would yield a higher lifting condensation level, leading to
522 higher clouds and a higher probability of convective precipitation.

523

524 **5 Conclusions**

525 GVF and LAI significantly affect the simulation of energy partitioning, yielding pronounced
526 differences between [simulated surface energy and water fluxes and temperatures](#) of ECC and LCC.
527 In our study area, the use of a generic crop parameterization (Croplands and Pasture in Noah-MP)
528 resulted in a humid bias along with lower surface temperatures. This humid bias will be largest in
529 landscapes with a balanced share of ECC and LCC, whereas in landscapes in which one of the two
530 crop types predominate, the bias will be weaker. We observed the strongest effects on turbulent
531 fluxes over the second part of the season, particularly in July-August. During this period, ECC are
532 at senescence growth stage or already harvested, while LCC have a fully developed ground-
533 covering canopy. We therefore expect that the observed differences will impact the simulation of
534 processes in the ABL. Our results show that splitting up croplands into ECC and LCC can improve
535 LSMs, particularly during transition periods and late in the growing season.

536
537 Increasing the LCC fraction by 10% reduces evapotranspiration and increases surface temperatures
538 over the first part of the growing season. Later in the season, this land use change leads to the
539 opposite situation: increased evapotranspiration accompanied by a slight cooling of the land
540 surface. Over the growing season, an increase of the LCC share by 10% leads to higher cumulative
541 evapotranspiration, which in turn lowers the seasonal water balance.

542

543 **6 References**

- 544 Chen, F. and Xie, Z.: Effects of crop growth and development on regional climate: A case study
545 over East Asian monsoon area, *Clim. Dyn.*, 38, 2291-2305, 2012.
- 546 Chen, F. and Xie, Z.: Effects of crop growth and development on land surface fluxes, *Adv.*
547 *Atmos. Sci.*, 28, 927-944, 2011.
- 548 Crawford, T. M., Stensrud, D. J., Mora, F., Merchant, J. W. and Wetzel, P. J.: Value of
549 incorporating satellite-derived land cover data in MM5/PLACE for simulating surface
550 temperatures, *J. Hydrometeorol.*, 2, 453-468, 2001.
- 551 [El Maayar, M., Chen, J. M. and Price, D. T.: On the use of field measurements of energy fluxes
552 to evaluate land surface models, *Ecol. Model.*, 214, 293-304, 2008.](#)
- 553 [Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G.,
554 Ceulemans, R., Clement, R., Dolman, H., Granier, A., Gross, P., Grünwald, T., Hollinger, D.,
555 Jensen, N. -, Katul, G., Keronen, P., Kowalski, A., Lai, C. T., Law, B. E., Meyers, T.,
556 Moncrieff, J., Moors, E., Munger, J. W., Pilegaard, K., Rannik, Ü, Rebmann, C., Suyker, A.,
557 Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K. and Wofsy, S.: Gap filling strategies for
558 defensible annual sums of net ecosystem exchange, *Agric. For. Meteorol.*, 107, 43-69, 2001.](#)
- 559 Ghilain, N. a., Arboleda, A. a., Sepulcre-Cantò, G., Batelaan, O., Ardö, J. d. and Gellens-
560 Meulenberghs, F. a.: Improving evapotranspiration in a land surface model using biophysical
561 variables derived from MSG/SEVIRI satellite, *Hydrology and Earth System Sciences*, 16, 2567-
562 2583, 2012.
- 563 Gutman, G. and Ignatov, A.: The derivation of the green vegetation fraction from
564 NOAA/AVHRR data for use in numerical weather prediction models, *Int. J. Remote Sens.*, 19,
565 1533-1543, 1998.
- 566 Huyghe, C., De Vlieghe, A., van Gils, B. and Peeters, A.: Grassland and herbivore production in
567 Europe and effects of common policies, Editions Quae: Versailles, France ISBN : 978-2-7592-
568 2157-8 ISSN : 1777-4624, 2014.
- 569 [Imukova, K., Ingwersen, J., Hevart, M., and Streck, T.: Energy balance closure on a winter
570 wheat stand: comparing the eddy covariance technique with the soil water balance method,
571 *Biogeosciences*, 13, 63–75, 2016.](#)
- 572 Imukova, K., Ingwersen, J. and Streck, T.: Determining the spatial and temporal dynamics of the
573 green vegetation fraction of croplands using high-resolution RapidEye satellite images, *Agric.*
574 *For. Meteorol.*, 206, 113-123, 2015.
- 575 Ingwersen, J., Imukova, K., Högy, P. and Streck, T.: On the use of the post-closure methods
576 uncertainty band to evaluate the performance of land surface models against eddy covariance
577 flux data, *Biogeosciences*, 12, 2311-2326, 2015.

578 Ingwersen, J., Steffens, K., Högy, P., Warrach-Sagi, K., Zhunusbayeva, D., Poltoradnev, M.,
579 Gäbler, R., Wizemann, H. -, Fangmeier, A., Wulfmeyer, V. and Streck, T.: Comparison of Noah
580 simulations with eddy covariance and soil water measurements at a winter wheat stand, Agric.
581 For. Meteorol., 151, 345-355, 2011.

582 Ingwersen, J., Högy, P., Wizemann, H.D., Warrach-Sagi, K., Streck, T.: Coupling the land surface
583 model Noah-MP with the generic crop growth model Gecros: Model description, calibration and
584 validation. Agric. Forest Meteorol., in print, 2018

585 Fachagentur Nachwachsende Rohstoffe e. V. Anbau und Verwendung nachwachsender Rohstoffe
586 in Deutschland. März 2019

587 Koster, R. D., Guo, Z., Dirmeyer, P. A., Bonan, G., Chan, E., Cox, P., Davies, H., Gordon, C. T.,
588 Kanae, S., Kowalczyk, E., Lawrence, D., Liu, P., Lu, C. -, Malyshev, S., McAvaney, B.,
589 Mitchell, K., Mocko, D., Oki, T., Oleson, K. W., Pitman, A., Sud, Y. C., Taylor, C. M.,
590 Verseghy, D., Vasic, R., Xue, Y. and Yamada, T.: GLACE: The Global Land-Atmosphere
591 Coupling Experiment. Part I: Overview, J. Hydrometeorol., 7, 590-610, 2006.

592 Lhomme, J. - and Chehbouni, A.: Comments on dual-source vegetation-atmosphere transfer
593 models, Agric. For. Meteorol., 94, 269-273, 1999.

594 [Mauder, M., Cuntz, M., Drüe, C., Graf, A., Rebmann, C., Schmid, H. P., Schmidt, M. and](#)
595 [Steinbrecher, R.: A strategy for quality and uncertainty assessment of long-term eddy-covariance](#)
596 [measurements, Agric. For. Meteorol., 169, 122-135, 2013.](#)

597 [Mauder M., Foken. T.: Documentation and Instruction Manual of the Eddy-Covariance Software](#)
598 [Package TK3. Arbeitsergebnisse Nr. 46, Universität Bayreuth, Abteilung Mikrometeorologie,](#)
599 [ISSN 1614-8916, Bayreuth, 2011.](#)

600 McPherson, R. A. and Stensrud, D. J.: Influences of a winter wheat belt on the evolution of the
601 boundary layer, Mon. Weather Rev., 133, 2178-2199, 2005.

602 Meier, U., Bleiholder, H., Buhr, L., Feller, C., Hack, H., Hess, M., Lancashire, P. D., Schnock,
603 U., Stauss, R., Van Den Boom, T., Weber, E. and Zwerger, P.: The BBCH system to coding the
604 phenological growth stages of plants-history and publications, Journal für Kulturpflanzen, 61,
605 41-52, 2009.

606 Miller, J., Barlage, M., Zeng, X., Wei, H., Mitchell, K. and Tarpley, D.: Sensitivity of the
607 NCEP/Noah land surface model to the MODIS green vegetation fraction data set, Geophys. Res.
608 Lett., 33, 2006.

609 Niu, G. -, Yang, Z. -, Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning,
610 K., Niyogi, D., Rosero, E., Tewari, M. and Xia, Y.: The community Noah land surface model
611 with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-
612 scale measurements, J. Geophys. Res. D Atmos., 116, 2011.

613 Raddatz, R. L.: Evidence for the influence of agriculture on weather and climate through the
614 transformation and management of vegetation: Illustrated by examples from the Canadian
615 Prairies, *Agric. For. Meteorol.*, 142, 186-202, 2007.

616 Refslund, J., Dellwik, E., Hahmann, A. N., Barlage, M. J. and Boegh, E.: Development of
617 satellite green vegetation fraction time series for use in mesoscale modeling: Application to the
618 European heat wave 2006, *Theor. Appl. Climatol.*, 117, 377-392, 2014.

619 Rundquist, B. C.: The influence of canopy green vegetation fraction on spectral measurements
620 over native tallgrass prairie, *Remote Sens. Environ.*, 81, 129-135, 2002.

621 Santanello Jr., J. A., Peters-Lidard, C. D., Kennedy, A. and Kumar, S. V.: Diagnosing the nature
622 of land-atmosphere coupling: A case study of dry/wet extremes in the U.S. southern Great
623 Plains, *J. Hydrometeorol.*, 14, 3-24, 2013.

624 SRU Special Report: Climate Change Mitigation by Biomass, Special Report, The German
625 Advisory Council on the Environment,
626 [http://www.umweltrat.de/SharedDocs/Downloads/EN/02_Special_Reports/2007_Special_Report](http://www.umweltrat.de/SharedDocs/Downloads/EN/02_Special_Reports/2007_Special_Report_Climatol.html)
627 [_Climate_Change.html](http://www.umweltrat.de/SharedDocs/Downloads/EN/02_Special_Reports/2007_Special_Report_Climatol.html), 2007.

628 Sulis, M., Langensiepen, M., Shrestha, P., Schickling, A., Simmer, C. and Kollet, S. J.:
629 Evaluating the influence of plant-specific physiological parameterizations on the partitioning of
630 land surface energy fluxes, *J. Hydrometeorol.*, 16, 517-533, 2015.

631 Tian, Y., Zou, C. Z., Mitchell, K. E., Wong, V., Kogan, F. N., Jiang, L. and Zhan, X.:
632 Improvements of numerical weather predictions using a new AVHRR green vegetation fraction
633 dataset, in: *The International Society for Optical Engineering (SPIE)*, San Diego, CA, 12 August
634 2008 through 14 August 2008, 2008.

635 Tsvetsinskaya, E. A., Mearns, L. O. and Easterling, W. E.: Investigating the effect of seasonal
636 plant growth and development in three-dimensional atmospheric simulations. Part I: Simulation
637 of surface fluxes over the growing season, *J. Clim.*, 14, 692-709, 2001a.

638 Tsvetsinskaya, E. A., Mearns, L. O. and Easterling, W. E.: Investigating the effect of seasonal
639 plant growth and development in three-dimensional atmospheric simulations. Part II:
640 Atmospheric response to crop growth and development, *J. Clim.*, 14, 711-729, 2001b.

641 van Heerwaarden, C. C., de Arellano, J. V. -, Moene, A. F. and Holtslag, A. A. M.: Interactions
642 between dry-air entrainment, surface evaporation and convective boundary-layer development,
643 *Q. J. R. Meteorol. Soc.*, 135, 1277-1291, 2009.

644 Xue, Y., Fennessy, M. J. and Sellers, P. J.: Impact of vegetation properties on U.S. summer
645 weather prediction, *J. Geophys. Res. D Atmos.*, 101, 7419-7430, 1996.

646 LUBW, 2010. Landesanstalt für Umwelt, Messungen, Naturschutz Baden-Württemberg.

- 647 Milovac, J., Warrach-Sagi, K., Behrendt, A., Späth, F., Ingwersen, J., Wulfmeyer, V.:
648 Investigation of PBL schemes combining the WRF model simulations with scanning water vapor
649 differential absorption laser measurements, *J. Geophys. Res. Atmos.*, 121, 2016.
- 650 Nielsen, J. R., D. Ebba, A. N. Hahmann, and E. Boegh (2013), Representing vegetation
651 processes in hydrometeorological simulations using the WRF model, *DTU Wind Energy*, 128 p.
652 (Riso – PhD; No. 0016(EN)).
- 653 Margulis S. A. and Entekhabi, D.: Feedback between the Land Surface Energy Balance and
654 Atmospheric Boundary Layer Diagnosed through a Model and Its Adjoint. *J. Hydrometeor.*, 2,
655 599–620., 2001
- 656 Wizemann, H.-D., Ingwersen, J., Högy, P., Warrach-sagi, K., Streck, T. and Wulfmeyer, V.:
657 Three year observations of water vapor and energy fluxes over agricultural crops in two regional
658 climates of Southwest Germany. *Meteorol.*, 2014