

# Cryptic roles of tetrathionate in the sulfur cycle of marine sediments: Microbial drivers and indicators

Subhrangshu Mandal<sup>1</sup>, Sabyasachi Bhattacharya<sup>1</sup>, Chayan Roy<sup>1</sup>, Moidu Jameela Rameez<sup>1</sup>,  
5 Jagannath Sarkar<sup>1</sup>, Tarunendu Mapder<sup>2,‡</sup>, Svetlana Fernandes<sup>3</sup>, Aditya Peketi<sup>3</sup>, Aninda  
Mazumdar<sup>3,\*</sup> and Wriddhiman Ghosh<sup>1,\*</sup>

<sup>1</sup> Department of Microbiology, Bose Institute, P-1/12 CIT Scheme VIIM, Kolkata 700054, India.

<sup>2</sup> ARC CoE for Mathematical and Statistical Frontiers, School of Mathematical Sciences,  
10 Queensland University of Technology, Brisbane, QLD 4000, Australia.

<sup>3</sup> CSIR-National Institute of Oceanography, Dona Paula, Goa 403004, India.

**Present address:** <sup>‡</sup> Department of Medicine, Indiana University School of  
Medicine, Indianapolis, IN 46202, USA.

**\* Correspondence emails:** wriman@jcbosc.ac.in / maninda@nio.org

**Running Title:** Tetrathionate metabolism in marine sediments

**Key words:** sulfur cycle, tetrathionate, tetrathionate metabolism, marine  
sediments, Arabian Sea, metaomics in biogeochemistry

---

## Abstract

To explore the potential role of tetrathionate in the sedimentary sulfur cycle, population ecology of  
25 microorganisms capable of metabolizing this polythionate was revealed at 15-30 cm resolution along two,  
~3-m-long, cores collected from 530 and 580 meters below the sea level, off India's west coast, within the  
oxygen minimum zone (OMZ) of the Arabian Sea. Metagenome analysis along the cores revealed  
widespread occurrence of genes involved in the formation, oxidation, and reduction of tetrathionate; high  
diversity and relative-abundance was also detected for bacteria that are known to render these  
30 metabolisms *in vitro*. Results of slurry-culture of the sediment-samples in thiosulfate- or tetrathionate-  
containing microbial growth media, data obtained via pure-culture isolation, and finally metatranscriptome  
analyses, corroborated the *in situ* functionality of the tetrathionate-forming, oxidizing, and reducing  
microorganisms. Ion chromatography of pore-waters revealed the presence of up to 11.1  $\mu\text{M}$  thiosulfate in  
the two cores, whereas tetrathionate remained undetected in spectroscopic assay based on its reaction  
35 with cyanide. While thiosulfate oxidation by chemolithotrophic bacteria prevalent *in situ* is the apparent  
source of tetrathionate in this ecosystem, high biochemical and geochemical reactivity of this polythionate  
could be instrumental in its cryptic status in the sulfur cycle. Potential abiotic origin of tetrathionate in the  
sediment-horizon explored could neither be ruled out nor confirmed from the geochemical information  
available. On the other hand, tetrathionate potentially present in the system can be either oxidized to  
40 sulfate or reduced back to thiosulfate/sulfide via chemolithotrophic oxidation and respiration by native  
bacterial populations respectively. Up to 2.01 mM sulfide present in the sediment-cores may also reduce  
tetrathionate abiotically to thiosulfate and elemental sulfur. However, in the absence of measured data for  
 $\text{O}_2$  or other oxyanions having possibilities of serving as electron acceptors, the biogeochemical modalities  
of the oxidative half of the tetrathionate cycle remained unresolved.

## 1 Introduction

Phylogenetically diverse microorganisms oxidize or reduce different sulfur species to meet their bioenergetic requirements, and in doing so play profound roles in biogeochemical sulfur cycling in nature (Baumgartner et al., 2006; Ghosh and Dam, 2009; Wasmund et al., 2017). Within the marine realm, microbial processes of sulfur cycle are central to benthic biogeochemistry as they are linked to the *in situ* transformations, sequestrations, and fluxes of carbon, nitrogen and iron. There have been extensive studies of benthic/sedimentary sulfur cycle across the global ocean (Jørgensen, 1990; Jørgensen and Bak 1991; Rudnicki et al., 2001; Tostevin et al., 2014), and the focus of such investigations has typically been on geomicrobial transformations of the two redox end-members sulfate and sulfide (Holmkvist et al., 2011; Jørgensen et al., 2019). Elemental sulfur and thiosulfate have also been envisaged as constituting key junctions in the network of sulfur species transformations in marine sediments (Jørgensen, 1990; Jørgensen and Bak, 1991; Thamdrup et al., 1994). However, tetrathionate or other polythionates are rarely appreciated for their potential role(s) in marine sedimentary sulfur cycle, presumably because these sulfur species are not abundant in these environments.

Overall paucity of polythionates in natural environments is largely attributable to their high reactivity under biotic (Kanao et al., 2007; Ghosh and Dam, 2009; Boden et al., 2010; Pyne et al., 2017, 2018) as well as abiotic (Schippers et al., 1999; Schippers and Jørgensen, 2001) conditions. Cryptic nature of these sulfur species notwithstanding, several bacteria capable of producing and/or utilizing tetrathionate for bioenergetic purposes have been isolated from different terrestrial and aquatic (both fresh-water and marine) habitats (Kaprálék, 1972; Oltmann and Stouthamer, 1975; Barrett and Clark, 1987; Price-Carter et al., 2001; Sorokin, 2003; Ghosh et al., 2005, 2006; Ghosh and Roy, 2007). At the same time, several enzymes catalyzing redox reactions involving tetrathionate have been characterized in taxonomically diverse microorganisms. For instance, thiosulfate dehydrogenase (TsdA), widespread in photo- or chemolithotrophic bacteria, is attributed to the formation of tetrathionate from thiosulfate (Hensen et al., 2006; Denkmann et al., 2012; Brito et al., 2015, Pyne et al., 2018; Rameez et al., 2019). In several other archaeal and bacterial chemolithotrophs, thiosulfate to tetrathionate conversion is mediated by thiosulfate:quinone oxidoreductase (TQO or DoxDA; see Muller et al., 2004; Rzhepishevskaya et al., 2007; Quatrini et al., 2009; Kikumoto et al., 2013). Chemolithotrophic oxidation of tetrathionate to sulfate, on the other hand, is rendered either (i) by the pyrroloquinoline quinone (PQQ)-binding tetrathionate hydrolase (TetH), as described in *Acidithiobacillus* species (De Jong et al., 1997; Kanao et al., 2007; Rzhepishevskaya et al., 2007; van Zyl et al., 2008; Kanao et al., 2013), or (ii) via coupling with glutathione (to form glutathione:sulfodisulfane and sulfite) by the action of another PQQ-binding protein called thiol dehydrotransferase (ThdT), followed by the oxidation of glutathione:sulfodisulfane via iterative

actions of sulfate thiol esterase (SoxB) and c-type cytochrome containing sulfur dehydrogenase (SoxCD), as reported in *Advenella kashmirensis* (Pyne et al., 2017, 2018). On the reductive half, typical tetrathionate-reducing bacteria such as *Salmonella*, *Citrobacter* and *Proteus* transform tetrathionate to thiosulfate by tetrathionate reductase (TtrABC) followed by the conversion of thiosulfate to sulfide by thiosulfate reductase (PhsAB and PsrA) (Oltmann et al., 1975; Barrett and Clark, 1987; Hensel et al., 1999; Stoffels et al., 2011).

Here we use approaches of molecular microbiology to investigate the potential involvement of tetrathionate metabolism in the sulfur cycle of sediments underlying the perennial oxygen minimum zone (OMZ) of the Arabian Sea, off the west coast of India. Community structures and functions of tetrathionate-forming, oxidizing, and reducing, microorganisms were revealed by metagenome analyses and slurry-culture of sediment-samples along two ~3-m-long cores collected from 530 and 580 meters below the sea level (mbsl), at sites having the GPS coordinates 16°50.03' N, 71°59.50' E and 16°49.88' N, 71°58.55' E respectively; sediment-samples were also analyzed for the presence of tetrathionate and thiosulfate in their pore-waters. *In situ* activity of tetrathionate-metabolizers was corroborated by pure-culture isolation and metatranscriptome analysis for the deepest sample-site, within the sulfate-methane transition zone, of one of the two cores. Correspondence was also explored between the *de novo* sequenced genomes of the isolates and the metatranscriptome or the metagenomes sequenced. The microbial ecology delineated in this way was considered in the context of the *in situ* geochemistry to infer implications for the sedimentary sulfur cycle.

## 2 Materials and methods

### 2.1 Study site and, sample collection and storage

During a comprehensive exploration of the sediment biogeochemistry of eastern Arabian Sea OMZ (ASOMZ), onboard *RV Sindhu Sankalp* (SSK42), a number of gravity cores were collected from water-depths spanning 225 and 1275 mbsl, covering the entire thickness of the eastern ASOMZ (Fernandes et al., 2018). Of these, SSK42/5 and SSK42/6, on which the present study is based, were collected from 580 mbsl (16°49.88' N, 71°58.55' E) and 530 mbsl (16°50.03' N, 71°59.50' E) water-depths respectively (Fernandes et al., 2018), i.e. the approximate center of the vertical expanse of the ASOMZ off the west coast of India. Both the cores were ~3-m-long and 12 cm in diameter; their onboard sampling was carried out at 15 to 30 cm intervals, as described previously, under constant shower of high-purity N<sub>2</sub> to minimize exposure of the native microflora to aerial O<sub>2</sub> and avoid aerial oxidation of the H<sub>2</sub>S, Fe<sup>2+</sup> and other reduced chemical substances present in the sediments (Fernandes et al., 2018).

In order to protect the sediment-samples from aerial oxidation, only one ~30-cm-long C-shaped part of PVC core-liner was removed at a time, as shown in Fig. S1. The 30 cm length

exposed at a time for sampling was constantly and closely showered with high-purity N<sub>2</sub> emitted from multiple nozzles fitted to multiple nitrogen-generators. Immediately after the C-shaped longitudinal part of the PVC core-liner was cut open, top one cm of the exposed surface was scraped off along the core-circumference, using sterile scalpels, to eliminate potential contaminations from the core-liners' inner-surfaces and/or sea-waters through which the cores had passed. Subsequently, to sample a particular sediment-depth of the core for microbiological studies, an approximately 5-mm-thick sediment-slice (spanning equally on either side of the core-height marking) was scooped out with a sterile scalpel and put into a sterile polypropylene bottle. For every sediment-depth investigated, two such slices or sample-replicates were collected for duplicate metagenome analyses (these were designated as sample-replicates 1 and 2, see Tables S1 and S2), a third one was taken for metatranscriptome analysis, while two more replicates were taken for culture-dependent investigations. All five replicates were collected in individual screw-capped bottles and those meant for metatranscriptomics were treated immediately with RNeasy (Qiagen Inc., USA). The head-space of each sample-containing bottle was flushed with high-pure N<sub>2</sub>, subsequent to which the bottles were sealed with Parafilm (Bemis Company Inc., Neenah, USA) and stored frozen or cool. Sample-replicates meant for culture-independent and culture-dependent studies were stored at -20°C and 4°C respectively. From the laboratory onboard SSK42, *en route* to Bose Institute, and over subsequent preservation of samples, these temperatures were maintained.

For onboard extraction of pore-waters, samples from a particular sediment-depth were taken out by inserting sterile 50 ml cut-off syringes deep inside the core cross-section, multiple times along the circumference on the exposed 'C half'. The samples were immediately collected in sterile 50 ml centrifuge tubes. All these operations were carried out under focused streams of high-pure N<sub>2</sub>. The tubes were centrifuged at 4700 × g for 15 minutes at 4°C, and the supernatants collected were syringe-filtered through 0.22 µm cellulose acetate membranes. Aliquots for different chemical analyses were dispensed to individual glass vials containing sodium azide that arrests microbial activity; only the vials meant for precipitating dissolved sulfide (ΣHS<sup>-</sup>) from the aliquots [in the form of cadmium sulfide (CdS)] contained cadmium nitrate [Cd(NO<sub>3</sub>)<sub>2</sub>] instead of sodium azide. All the vials were crimp sealed immediately after N<sub>2</sub> flushing, and stored at 4°C until further analysis.

## 2.2 Analytical method

Sulfide and sulfate concentrations in the pore-water samples were determined and reported previously by Fernandes et al., (2018). Concentration of dissolved thiosulfate in the pore-water samples was determined by ion chromatography using an Eco IC (Metrohm AG, Herisau, Switzerland) equipped with a conductivity detector (Metrohm, IC detector 1.850.9010). Chemical

155 suppression was used for this purpose, while separation was carried using a Metrosep A Supp5 -  
250/4.0 anion exchange column (Metrohm AG). A mixed solution of 1.0 mM sodium bicarbonate  
and 3.2 mM sodium carbonate was used as the eluent; 100 mM sulfuric acid was used as the  
regenerant; flow rate was 0.7 mL min<sup>-1</sup>, and injection volume 100 µL. Prior to analysis, pore-  
water samples were diluted 1000-fold with de-ionized water (Siemens, < 0.06 µS) and passed  
160 through 0.22 µm hydrophilic polyvinylidene fluoride membrane filters (Merck Life Science Private  
Limited, Bengaluru, India). Analytical grade thiosulfate IC Standard (Sigma Aldrich, St. Louis,  
USA) was used to prepare the calibration curve for quantification of this anion. Three different  
concentrations of thiosulfate, 0.5 µM, 5 µM and 20 µM, were measured for the construction of  
calibration curve by plotting peak height against concentration. Based on triplicate analysis of the  
standards, deviations from their actual concentrations were found to be less than 2.5%.

165 Concentration of tetrathionate in the pore-water samples was measured by cyanolytic  
method (Kelly and Wood, 1994), where tetrathionate reacts with cyanide to form thiocyanate  
according to the reaction  $S_4O_6^{2-} + 3CN^- + H_2O \leftrightarrow SCN^- + S_2O_3^{2-} + SO_4^{2-} + 2HCN$ , and the  
thiocyanate (SCN<sup>-</sup>) formed is quantified spectrophotometrically from the absorbance of ferric  
thiocyanate ( $\epsilon = 5030 \text{ M}^{-1} \text{ cm}^{-1}$ ,  $\lambda = 460 \text{ nm}$ ), which is formed by reacting the thiocyanate ion with  
170 ferric nitrate. Differences in the reactivity of thionates with cyanide enable their discrimination and  
quantitative characterization within mixtures of such compounds. For instance, trithionate is  
stable at high pH and reacts with cyanide only at elevated temperatures; thiosulfate reacts with  
cyanide at room temperature, that too only in the presence of copper(II) catalyst; in contrast, the  
higher polythionates ( $S_nO_6^{2-}$ , where  $n = 4$  or more) react rapidly with cyanide at room temperature  
175 to form SCN<sup>-</sup>, S<sub>2</sub>O<sub>3</sub><sup>2-</sup>, SO<sub>4</sub><sup>2-</sup> and HCN. Furthermore, in aqueous systems where intermediate  
sulfur compounds are cycling and dissolved sulfide is present, cyanide may also react with zero-  
valent sulfur components of the colloidal fraction of particulate elemental sulfur, polysulfides  
(S<sub>n</sub><sup>2-</sup>), and their protonated forms, albeit when administered under hot and slightly acidic  
condition (Kamyshny, 2009), which is not the case in the cyanolytic procedure currently followed  
180 for tetrathionate estimation (Kelly and Wood, 1994). Here, 0.1 ml of the pore-water sample was  
added to 0.8 ml of cyanolytic buffer (a mixture of 50 ml of 0.2 M NaH<sub>2</sub>PO<sub>4</sub> and 39 ml of 0.2 M  
NaOH; pH 7.4) and the volume was made up to 2.0 ml. The mixture was then chilled on ice for  
20 minutes and 1 ml of 0.1 M pre-chilled potassium cyanide was added and mixed rapidly and  
incubated on ice for another 20 minutes. Finally, 0.6 ml of ferric nitrate reagent [30.3 g  
185 Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O in 21.7 ml of 72% perchloric acid, made up to 50 ml with distilled water] was  
added with continuous agitation and allowed to warm to room temperature until the white  
precipitate (if any) re-dissolved. Volume was made up to 5 ml and optical density of the colour  
due to ferric thiocyanate was measured on a spectrophotometer. The standard curve used for  
tetrathionate estimation was prepared using different concentrations (0.1, 0.5 1.0, 5.0 and 10.0

190  $\mu\text{M}$ ) of analytical grade tetrathionate (Sigma Aldrich, St. Louis, USA). Based on triplicate analysis of these standards, deviations from their actual concentrations were found to be less than 5%.

### 2.3 Extraction of total DNA/RNA from sediment-samples/pure-culture isolates

195 Total community DNA (metagenome) was extracted from the sediment-samples using PowerSoil DNA Isolation Kit (MoBio, Carlsbad, USA), as per the manufacturer's protocol. Microgram-level of DNA was obtained from each batch of preparatory reaction that started with 0.5 g sediment-sample. Genomic DNA of pure culture isolates was extracted using HiPurA Bacterial Genomic DNA Purification Kit (Himedia Laboratories, Mumbai, India), following manufacturer's instructions. Quality of metagenomic/genomic DNA samples was checked by electrophoresis and considered  
200 to be of high quality when no degradation signs were apparent. DNA quantity was determined using Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific, Waltham, USA).

Total community RNA (metatranscriptome) was extracted from an RNAlater-treated sample-replicate by using RNA PowerSoil Total RNA Isolation Kit (MoBio), as per manufacturer's protocol. Nanogram-level total RNA was obtained after pooling the products of 15 individual  
205 preparatory reactions, each carried out using 2 g sediment-sample. All the individual RNA preparations were subjected to DNase digestion by RNase free DNase I (Thermo Fisher Scientific) and purified using RNeasy MinElute Cleanup Kit (Qiagen, Hilden, Germany); their concentrations were measured using Quant-iT RiboGreen RNA Assay Kit (Thermo Fisher Scientific). Integrity of RNA (RIN) within the individual preparations was determined on a  
210 TapeStation RNA ScreenTape electrophoretic system (Agilent Technologies, Santa Clara, USA) and only high-quality preparations having RIN value  $> 7.0$  were added to the RNA pool that were subsequently used for sequencing library construction.

### 2.4 Metagenome (total community DNA) / genome sequencing

215 The duplicate set of total community DNA (metagenomes) extracted for each sediment-depth explored along SSK42/5 and SSK42/6 were shotgun sequenced individually on an Ion Proton sequencing platform (Thermo Fisher Scientific) using 200 nucleotide read-chemistry, as described previously (Ghosh et al., 2015). Complete lists of sedimentary communities investigated along SSK42/5 and SSK42/6 are given in Tables S1 and S2 respectively, together  
220 with the accession numbers of the metagenomic sequence datasets.

1  $\mu\text{g}$  DNA from each metagenome-sample was taken for deep shotgun sequencing by the Ion Proton platform using 200 bp read chemistry on a PI V2 Chip. Sequencing libraries were constructed using Ion Plus Fragment Library Kit (Thermo Fisher Scientific), following the manufacturer's Ion Plus gDNA Library Preparation User Guide. The Proton library was generated  
225 using 1  $\mu\text{g}$  of genomic DNA which was fragmented to approximately 200 base pairs by the

Covaris S2 system (Covaris, Inc., Woburn, USA) and purified with 1.8X Agencourt Ampure XP Beads (Beckman Coulter, Brea, USA). Fragmentation was followed by end-repair, blunt-end ligation of the Ion Xpress Barcode and Ion P1 adaptors, and nick translation.

230 Post-ligation, size selection was done using E-Gel Size-Select 2% Agarose gels (Thermo Fisher Scientific) with 300 bp target size. Final PCR was performed using platinum PCR SuperMix High Fidelity and Library Amplification Primer Mix (Thermo Fisher Scientific), for 5 cycles of amplification. The resulting library was purified using 1.2X AMPure XP reagent (Beckman Coulter) and the concentration determined with Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific); size distribution was done with Agilent 2100 Bioanalyzer high-sensitivity DNA  
235 kit (Agilent Technologies). Libraries were pooled in equimolar concentrations and used for template preparation.

Library templates for sequencing were prepared using OneTouch 2 protocols and reagents (Thermo Fisher Scientific). Library fragments were clonally amplified onto ion sphere particles (ISPs) through emulsion PCR and then enriched for template-positive ISPs. Proton  
240 emulsion PCR reactions utilized the Ion PI Template OT2 200 Kit v3 (Thermo Fisher Scientific). Following recovery, enrichment was completed by selectively binding the ISPs containing amplified library fragments to streptavidin coated magnetic beads, removing empty ISPs through washing steps, and denaturing the library strands to allow for collection of the template-positive ISPs. For all reactions, these steps were accomplished using the ES module of the Ion  
245 OneTouch 2. The selected ISPs were loaded on PI V2 Chip and sequenced with the Ion PI 200 Sequencing Kit (Thermo Fisher Scientific) using the 500 flow (125 cycle) run format.

Whole genomes of isolated bacterial strains were sequenced on an Ion S5 platform (Thermo Fisher Scientific) using 400 nucleotide read-chemistry on a 530 or 520 Chip. Libraries were constructed by the Ion Xpress Plus Fragment library kit (Thermo Fisher Scientific) using  
250 100 ng genomic DNA from each isolate. In this procedure genomic DNA samples were fragmented using Ion Shear Plus Reagents (Thermo Fisher Scientific). The fragmented libraries were purified using 1.8X Agencourt Ampure XP Beads (Beckman Coulter), and subjected to barcode-adapter ligation and nick repair. Adapter-ligated and nick-repaired libraries were purified again by 1X Agencourt Ampure XP Beads (Beckman Coulter).

255 Size selection of the libraries was done using E-Gel Size-Select 2% Agarose gels (Thermo Fisher Scientific) with 480 bp target size. Final PCR was performed using platinum SuperMix High Fidelity PCR system and Library Amplification Primer Mix (both from Thermo Fisher Scientific), for 8 cycles of amplification. The resulting libraries were purified using 1x Agencourt AMPure XP reagent (Beckman Coulter). Concentrations of the purified libraries were determined  
260 with Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific). Libraries were then pooled in equimolar concentrations and used for template preparation.

The library template to be used for sequencing was prepared using Ion OneTouch 2 reagents (Thermo Fisher Scientific). Library fragments were clonally amplified onto ion sphere particles (ISPs) through emulsion PCR and then enriched for template-positive ISPs. Following emulsion PCR, enrichment was completed by selectively binding the ISP-containing amplified library fragments to streptavidin-coated magnetic beads, removing empty ISPs through washing steps, and denaturing the library strands to allow for collection of the template-positive ISPs using the Ion OneTouch ES instrument (Thermo Fisher Scientific). The selected ISPs were loaded on a 530 or 520 Chip and sequencing was performed with the Ion S5 Sequencing Kit (Thermo Fisher Scientific) using the 850-flow run format.

## 2.5 Metatranscriptome (community mRNA) sequencing

The pooled total RNA preparations were selectively converted to a library of template molecules using TruSeq Stranded mRNA and Total RNA kit (Illumina Inc., San Diego, USA). Depletion of rRNAs was carried out using the Ribo-Zero Gold system (Illumina Inc.), which is an integral part of the kit used for preparing the library. The rRNA-depleted RNA pool, which was expected to contain only the total mRNA, was fragmented into small pieces using divalent cations under elevated temperature. The cleaved RNA fragments were copied into first strand cDNAs using reverse transcriptase and random primers. This was followed by second strand cDNA synthesis using DNA Polymerase I and RNase H. cDNA fragments were then subjected to end-repair, addition of single 'A' bases, adaptor ligation, purification and enrichment with PCR to create the final library, which was sequenced on a HiSeq4000 platform (Illumina Inc.) using paired end, 2 x 150 nucleotide, sequencing by synthesis read-chemistry with dual indexing workflows. Furthermore, in order to extract and eliminate any rRNA read that may have remained in the raw metatranscriptomic sequence dataset, the 26,579,343 read-pairs available in all were mapped onto SILVA large subunit as well as small subunit rRNA gene sequence database (Quast et al., 2012), using the short read aligner Bowtie2 v.2.3.4.3 (Langmead and Salzberg, 2012) in default local (sensitive) alignment mode. This identified ~0.3% reads as ascribable to rRNAs, thereby leaving 26,496,769 read-pairs in the final dataset used for downstream analyses.

## 2.6 *De novo* assembly and annotation of genomes/metagenomes/metatranscriptome

High quality reads (Phred score cut-off 20) from the duplicate metagenomic sequence datasets available for each sediment-community were co-assembled using Megahit v1.2.x (Li et al., 2015) with the kmer lengths of 21, 29, 39, 59, 79, 99, 119 and 141. Contigs of > 100 bp length were searched using MetaGeneMark (Zhu et al., 2010) for genes encoding peptides having lengths of > 30 amino acids. For the genomes of pure-culture isolates, high quality reads (Phred score cut-off 20) were assembled using SPAdes 3.13.0 (Nurk et al., 2013), with kmer lengths of 21, 33, 55,



77, 99 and 127, and minimum coverage cut-off 35X. The whole genome sequences were deposited to the GenBank and annotated using Prokaryotic Genome Annotation Pipeline (PGAP located at [https://www.ncbi.nlm.nih.gov/genome/annotation\\_prok/](https://www.ncbi.nlm.nih.gov/genome/annotation_prok/)) of National Center for Biotechnology Information (NCBI), Bethesda, MD, USA. Total metagenomic or metatranscriptomic reads available for the different sediment-communities were mapped onto the individual genome sequences or the manually-curated gene-catalogs obtained from the individual genomes, using Bowtie2 v.2.3.4.3 (Langmead and Salzberg, 2012) in default local (sensitive) alignment mode. The rRNA-sequence-free metatranscriptomic dataset was assembled using the python script rnaspad.es.py, available within SPAdes 3.13.0 (Nurk et al., 2013), with default parameters. Genes encoding continuous stretches of minimum 30 amino acids were predicted in contigs longer than 100 bp using Prodigal v2.6.3 (Hyatt et al., 2010).

Gene-catalogs obtained after *de novo* assembly of the individual metagenomes, or the solitary metatranscriptome, were functionally annotated by searching against EggNOG v5.0 database ([http://eggnog5.embl.de/download/eggnog\\_5.0/](http://eggnog5.embl.de/download/eggnog_5.0/)) with EggNOG-mapper (Huerta-Cepas et al., 2016) (<http://beta-eggnogdb.embl.de/#/app/emapper>) using HMMER algorithms. Putative protein sequence catalogs obtained for the individual genomes via PGAP annotation were re-annotated by searching against EggNOG v5.0 database with EggNOG-mapper using HMMER algorithm. Genes encoding proteins involved in tetrathionate formation, oxidation and reduction were identified within the annotated genomes by their orthology numbers designated in *Kyoto Encyclopedia of Genes and Genomes* (KEGG; Kanehisa et al., 2016).

## 2.7 Direct taxonomic annotation of raw metagenomic reads

Raw reads from the duplicate metagenomic sequence datasets of each sediment-community were directly annotated for their taxonomic affiliation by searching the datasets individually against the NCBI non-redundant (*nr*) protein sequence database, using Organism Abundance tool of MG-RAST 3.6 (Meyer et al., 2008). The two independent values obtained in this way for the relative abundances of genera within a community were averaged and used for comparisons between communities; percentage allocation of reads over various genera was taken as a direct measure of the relative abundance (prevalence) of those genera within the community (Tringe et al., 2005; Ghosh et al., 2015; Roy et al., 2020). Within MG-RAST, sequences were trimmed to contain no more than 5 successive bases with Phred quality score < 15. To classify reads using Organism Abundance tool, Best Hit Classification algorithm was followed, reporting only those BlastX search results which had minimum 45 nucleotides (15 amino acids) alignment,  $\geq 60\%$  identity, and  $e\text{-value} \leq 1e^{-5}$ ; these cut-offs are stringent enough for genus-level classification of homologs of metabolically diverse genes, irrespective of their intrinsic levels of conservation.

## 2.8 Slurry-culture of sediment-samples

335 Ability of the different sedimentary microbial communities to form tetrathionate (from thiosulfate),  
oxidize tetrathionate (to sulfate) or reduce tetrathionate (to thiosulfate and/or sulfide) was tested  
via incubation of the samples in culture media supplemented with known concentrations of  
thiosulfate or tetrathionate. Within the slurry-culture set-ups, *in vitro* rates of redox  
340 transformations of sulfur involving tetrathionate were determined and expressed as  $\mu\text{mol S per L}$   
slurry per day. For each experiment testing the formation or oxidation of tetrathionate, 10% (w/v)  
sediment-sample was suspended in artificial sea water (ASW) supplemented with thiosulfate (T)  
or tetrathionate (Tr), i.e. ASWT or ASWTr broth medium (Alam et al., 2013) respectively, to form  
a slurry; the culture flask was then incubated aerobically at 15°C on a rotary shaker (150 rpm).  
For each anaerobic experiment testing tetrathionate reduction, 10% (w/v) sediment-sample was  
345 suspended in tetrathionate-supplemented Rappaport Vassiliadis tetrathionate-supplemented  
(RVTr) medium (Vassiliadis, 1983) that was already made O<sub>2</sub>-free by addition of sodium  
thioglycolate. Addition of sediment-samples to sterile O<sub>2</sub>-free RVTr media (contained in a screw-  
capped bottles) and subsequent incubation of the culture bottles were all carried out inside a  
Whitley H35 Hypoxystation (Don Whitley Scientific, West Yorkshire, UK) preset at 75% humidity,  
350 15°C temperature and 0% partial pressure of O<sub>2</sub>, using the gas mixture N<sub>2</sub>:H<sub>2</sub>:CO<sub>2</sub> = 80:10:10  
(v/v/v).

The ASWT or ASWTr medium (both having pH 7.5) contained ASW supplemented with  
Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>·5H<sub>2</sub>O (10 mM) or K<sub>2</sub>S<sub>4</sub>O<sub>6</sub> (5 mM) respectively (the two sulfur compounds were added to  
the media separately after filter-sterilization) (Alam et al., 2013). ASW contained the following L<sup>-1</sup>  
355 distilled water: 25.1 g NaCl, 1 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.3 g CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.2 g  
NaHCO<sub>3</sub>, 2.4 g Tris, 1 mL trace element solution and 0.5 g K<sub>2</sub>HPO<sub>4</sub> (added after autoclaving  
separately). 1 L trace element solution (pH 6.0), in turn, contained 50 g EDTA, 22 g  
ZnSO<sub>4</sub>·7H<sub>2</sub>O, 5.06 g MnCl<sub>2</sub>, 4.99 g FeSO<sub>4</sub>, 1.1 g (NH<sub>4</sub>)<sub>6</sub> MoO<sub>26</sub>·4H<sub>2</sub>O, 1.57 g CuSO<sub>4</sub> and 1.61 g  
CoCl<sub>2</sub>·6H<sub>2</sub>O. RVTr medium (pH 5.4) contained the following L<sup>-1</sup> of distilled water: 4.5 g soya  
360 peptone, 8.0 g NaCl, 0.4 g K<sub>2</sub>HPO<sub>4</sub>, 0.6 g KH<sub>2</sub>PO<sub>4</sub>, 29.0 g MnCl<sub>2</sub>, 0.036 g Malachite green, 10  
mM K<sub>2</sub>S<sub>4</sub>O<sub>6</sub>, and 0.5 g sodium thioglycolate (used as an O<sub>2</sub> scavenger) and 0.1 mg resazurin  
(added to indicate the presence of any dissolved O<sub>2</sub>).

During RVTr preparation, inside a Whitley H35 Hypoxystation preset to 0% partial  
pressure of O<sub>2</sub>, pre-weighed amount of potassium tetrathionate salt was first dissolved in a  
premeasured volume of anoxic, deionized water (degassed for several hours inside the H35  
365 Hypoxystation till the resazurine indicator added in the water became colorless). This anoxic  
tetrathionate solution was then added via filter-sterilization to a separate pre-autoclaved O<sub>2</sub>-free  
solution that contained the rest of the RVTr components in an appropriate volume and had  
cooled down to room temperature within the Hypoxystation. Thioglycolate that was already there

370 in the second solution had reacted irreversibly, during autoclaving, with the dissolved O<sub>2</sub> present  
in the mixed-salts solution to form dithiodiglycolate. Post-autoclave cooling of this second  
solution within the Hypoxystation, therefore, did not breakdown the S-S bonds of dithiodiglycolate  
to regenerate the SH<sup>-</sup>-containing thioglycolate, so there was no possibility of thioglycolate  
375 attacking the incoming tetrathionate solution. Moreover, (i) at neutral pH, thiol-group-containing  
reducing agents do not attack tetrathionate under non-enzymatic (abiotic) conditions, or for that  
matter in the absence of specific inorganic catalysts (Pyne et al., 2018); (ii) zero hour reading for  
all the slurry-culture sets in RVTr medium showed the intact presence of the 10 mM tetrathionate  
originally supplied in the medium; and (iii) abiotic control incubations involving autoclaved  
380 sediment-samples showed that the 10 mM tetrathionate supplied to the RVTr medium was intact  
after prolonged incubation.

Concentrations of the various redox species of sulfur were estimated in the slurry-culture  
at every 6 hr interval of incubation. End-point (final) concentrations of the different sulfur species  
in a given medium were recorded when no further changes were detected in their concentrations  
over three consecutive estimations. Concentrations of thiosulfate, tetrathionate and sulfate in the  
385 media were measured by iodometric titration, cyanolysis and gravimetric sulfate precipitation  
method respectively at different time intervals (Alam et al., 2013). Possible presence of dissolved  
sulfides was checked by precipitating as CdS by the addition of 2 M Cd(NO<sub>3</sub>)<sub>2</sub>, followed by  
spectroscopic estimation as described previously (Cline, 1969).

## 390 **2.9 Enrichment, isolation and characterization of bacterial strains**

Isolation of sulfur-oxidizing chemolithotrophs from the 275 cmbsf sediment-sample of SSK42/6  
was carried out in mineral salt-thiosulfate-yeast extract (MSTY), ASWT and ASWTY media.  
While the ASWTY (pH 7.5) medium was an yeast extract (500 mg L<sup>-1</sup>) supplemented derivative of  
ASWT, MSTY (pH 7.0) contained modified basal and mineral salts (MS) solution supplemented  
395 with 20 mM Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>·5H<sub>2</sub>O and 500 mg L<sup>-1</sup> yeast extract (Ghosh and Roy, 2006). MS, in turn,  
contained the following L<sup>-1</sup> distilled water: 1 g NH<sub>4</sub>Cl, 4 g K<sub>2</sub>HPO<sub>4</sub>, 1.5 g KH<sub>2</sub>PO<sub>4</sub>, 0.5 g  
MgSO<sub>4</sub>·7H<sub>2</sub>O and 5.0 mL trace metals solution (Vishniac and Santer, 1957). Three portions of  
the 275-cmbsf sediment-sample of SSK42/6 were added (5% w/v) individually to MSTY, ASWT  
and ASWTY broths, and incubated aerobically at 15°C until phenol red indicator present in the  
400 media turned yellow (apparently due to production of sulfuric acid from thiosulfate). Post  
yellowing, individual enrichment slurries were kept undisturbed for 1 h to allow sediment particles  
to settle down; 10 mL cell suspension from each flask was then centrifuged at 6000 g for 10 min  
and the pellet re-suspended in 1 mL of the corresponding medium, serially diluted, and spread  
onto agar plates of the corresponding medium, and incubated at 15°C. Morphologically distinct  
405 colonies were picked up and dilution-streaked until all colonies in individual plates looked similar;

representative colonies from such pure-culture plates were taken as strains and maintained in their respective isolation-medium. Only *Methylophaga*, though isolated in ASWT, was maintained in ASW supplemented with 0.3% (v/v) methanol (ASWM; this medium had a pH of 7.5) because its growth in ASWT waned after six straight sub-cultures. Chemolithotrophic abilities of the new isolates to oxidize thiosulfate to tetrathionate and/or tetrathionate to sulfate were tested in MSTY, MSTrY, ASWT, ASWTr, ASWTY, ASWTrY, ASWTM and ASWTrM media. MSTrY (pH 7.0) contained MS solution supplemented with 10 mM  $K_2S_4O_6$  and 500 mg  $L^{-1}$  yeast extract. ASWTrY (pH 7.5) was an yeast extract (500 mg  $L^{-1}$ ) supplemented derivative of ASWTr. ASWTM (pH 7.5) and ASWTrM (pH 7.5) were thiosulfate (10 mM  $Na_2S_2O_3 \cdot 5H_2O$ ) and tetrathionate (10 mM  $K_2S_4O_6$ ) supplemented variant of ASWM respectively. Concentrations of dissolved thiosulfate, tetrathionate and sulfate in the spent culture media were measured as described above.

Tetrathionate-reducing bacterial strains were isolated from the 275 cmbfs sediment-sample of SSK42/6 in RVTr medium (Vassiliadis, 1983) under strictly anaerobic condition. 2.5 g sediment-sample was added to 45 mL RVTr broth contained in a screw-capped bottle and prepared as described above. Sediment addition to the medium, and subsequent incubation of the screw-capped bottles at 15°C for one month, were all carried out inside the Whitley H35 Hypoxystation preset to zero  $O_2$  as stated above. After one month, still inside the Hypoxystation, 1 mL of the sediment-RVTr mixture was serially diluted and spread onto RVTr agar plates and incubated at 15°C. After growth appeared in the RVTr agar plates they were taken out of the Hypoxystation; biomasses were serially dilution streaked on to fresh plates and incubated aerobically until all colonies in the individual plates looked similar. Representative colonies from such pure-culture plates were taken and maintained aerobically in Luria broth medium. Tetrathionate-reducing abilities of the new isolates were tested by growing them anaerobically for 30 days in RVTr broth, inside the H35 Hypoxystation. Concentrations of dissolved thiosulfate, tetrathionate and sulfide in the RVTr cultures were measured by the methods mentioned above.

Genomic DNA extracted from the individual isolates was used as template for PCR amplification of 16S rRNA genes with the Bacteria-specific universal primer-pair 27f and 1492r (Gerhardt, 1994). 16S rRNA gene sequences were determined from the PCR products using a 3500xL Genetic Analyzer automated DNA sequencer (Thermo Fisher Scientific). The 16S rRNA gene sequence of each strain was compared against sequences available in the GenBank/EMBL/DDBJ databases, using BLASTN; strains were finally classified down to the lowest identifiable taxonomic category on the basis of their 16S rRNA gene sequence similarities with the closest, validly-published species having standing in nomenclature (<http://www.bacterio.net/>; see also Euzéby, 1997; Parte, 2013).

### 3 Results

#### 3.1 Tetrathionate-forming/oxidizing/reducing microorganisms and genes corresponding to such processes are abundant along SSK42/5 and SSK42/6

445 The duplicate metagenomic sequence datasets obtained for each of the 25 distinct sediment-  
samples explored along SSK42/5 and SSK42/6 were co-assembled and annotated individually.  
23 out of the 25 contig-collections obtained in this way were found to contain genes for  
tetrathionate formation (Table S3), while all 25 encompassed genes for tetrathionate oxidation  
(Table S4). Furthermore, 24 out of the 25 contig-collections contained genes for tetrathionate  
450 reduction (Table S5). The tetrathionate formation-related genes identified included those  
encoding for the different subunits of the thiosulfate dehydrogenases TsdA (Denkman et al.,  
2012; Pyne et al., 2018) and DoxDA (Quatrini et al., 2009), which catalyze the oxidation of  
thiosulfate to tetrathionate in taxonomically diverse bacteria and archaea. While the genes  
identified for tetrathionate oxidation encoded the sulfate thiol esterase SoxB and the sulfur  
455 dehydrogenase SoxC (Lahiri et al., 2006; Pyne et al., 2018), those detected for tetrathionate  
reduction encoded subunits of tetrathionate (TtrABC) and thiosulfate reductases (PhsAB and  
PsrA) (Barrett and Clark, 1987; Stoffels et al., 2011).

Concurrent with the above findings, direct taxonomic annotation of the raw (unassembled)  
metagenomic sequence datasets revealed that considerable proportions of the reads obtained  
460 from the individual sediment-depths of SSK42/5 and SSK42/6 were ascribable to bacterial  
genera whose members are known to render tetrathionate formation, oxidation or reduction. In  
that way, 1.3-4.36% and 3-7.8% of metagenomic reads obtained from the individual sample-sites  
of SSK42/5 (Fig. 1A) and SSK42/6 (Fig. 1B) were ascribable to the genera *Pseudomonas* and  
*Halomonas*, majority of marine strains of which are known to form tetrathionate from the  
465 oxidation of thiosulfate under aerobic or anaerobic condition (Tuttle, 1980; Mason and Kelly  
1988; Sorokin et al., 1999; 2003). 0.1-1.5 and 0.4-6.4% of metagenomic reads obtained from the  
individual sample-sites of SSK42/5 (Fig. 1A) and SSK42/6 (Fig. 1B) were ascribable to the  
genera *Acidithiobacillus*, *Guyparkeria*, *Halothiobacillus* and *Thiomicrospira*, all members of which  
oxidize tetrathionate chemolithotrophically (Hedrich and Johnson, 2013; Watsuji et al., 2016;  
470 Boden et al., 2017). 0.1-0.3 and 0.2-0.4% of metagenomic reads obtained from the individual  
sample-sites of SSK42/5 (Fig. 1A) and SSK42/6 (Fig. 1B) were ascribable to the genera  
*Citrobacter*, *Proteus* and *Salmonella*, all members of which respire by reducing tetrathionate to  
thiosulfate and/or sulfide (Kaprálék, 1972; Barrett and Clark, 1987; Price-Carter et al., 2001;  
Stoffels et al., 2011).

### 3.2 Synchronized population-fluctuation of different tetrathionate-metabolizing types, along SSK42/5 and SSK42/6

Analyses based on the direct taxonomic annotation of the unassembled metagenomic data from discrete sediment-depths of SSK42/5 revealed that the relative abundances of reads ascribed to the genera of tetrathionate-forming, oxidizing, and reducing bacteria fluctuate synchronously along the sediment-surface to core-bottom trajectory (Fig. 1A). Corroboratively, pair-wise Pearson correlation coefficients (CC) as well as Spearman rank correlation coefficients (RCC) between the prevalence of the three metabolic-types are also significantly high in SSK42/5 (Fig. 1A; Table S6), which indicate the existence of strong syntrophic interdependence between the three tetrathionate-metabolizing types in this sediment-horizon. Relative abundances of metagenomic reads ascribed to the genera of tetrathionate-forming, oxidizing, and reducing bacteria also fluctuate more or less synchronously along SSK42/6, excepting the region between 250 and 275 cmbsf (Fig. 1B), which is the sulfate-methane transition zone (SMTZ) of this sediment-horizon (notably, SMTZ in SSK42/5 laid below the 280 cmbsf sediment-depth explored in this core; see Fernandes et al., 2018 for the methane profiles of all the SSK42 cores). While lack of synchrony in the lower end of SSK42/6 apparently resulted in the lower correlation values obtained for this core (Table S7) as compared to those obtained for SSK42/5, it seems quite plausible that the changes in geochemistry and community architecture associated with the shallowing of SMTZ in SSK42/6 impacted the population ecology of tetrathionate-metabolizing microorganisms in this region. Sedimentation rate, age-depth profile and other geochemical features of the two cores separated by a distance of only one kilometer are otherwise largely comparable (Bhattacharya et al., 2019). Consistent prevalence of reads ascribed to the thiosulfate-to-tetrathionate-converting bacterial genera *Halomonas* and *Pseudomonas* in the metagenomes extracted from the different sample-sites of SSK42/5 and SSK42/6 (Figs. 1A and 1B) indicated that tetrathionate could be bioavailable in the chemical milieu of these sediment-horizons (notably, pure-culture strains belonging to these two genera were also isolated from the 275 cmbsf sample of SSK42/6; see section 3.3 below, and also Fig. 2). Apart from these two, several such genera were also found to be well represented in the metagenomes of SSK42/5 and SSK42/6, some members of which are known to produce tetrathionate as a free intermediate during the oxidation of thiosulfate to sulfate and release the same to the extra-cellular milieu (Tables S8 and S9). These organisms, affiliated to the genera *Acidithiobacillus*, *Advenella*, *Halothiobacillus*, *Paracoccus*, *Pusillimonas* and *Thiomicrospira*, can well increase tetrathionate availability in the ASOMZ sediments, even as they themselves are potential users of the tetrathionate produced (Ghosh et al., 2005; Hedrich and Johnson, 2013; Watsuji et al., 2016; Boden et al., 2017; Rameez et al., 2019). Tables S8 and S9 show the relevant references, and

the percentages of metagenomic reads that were found to be ascribed to these genera in the different sediment-samples of SSK42/5 and SSK42/6 respectively.

Tetrathionate can be oxidized *in situ* as a potential energy and electron source by members of the obligately chemolithotrophic genera *Acidithiobacillus*, *Guyparkeria*, *Halothiobacillus* and *Thiomicrospira* that were detected via direct taxonomic annotation of the unassembled metagenomic data (Figs. 1A and 1B) and/or isolated as pure cultures from the 275 cmbfs sample of SSK42/6 (Fig. 2). In addition, several such genera were detected (via direct annotation of metagenomic reads) along SSK42/5 and SSK42/6, some members of which are known to oxidize tetrathionate to sulfate. These organisms, affiliated to the genera *Advenella*, *Bosea*, *Burkholderia*, *Campylobacter*, *Hydrogenovibrio*, *Pandoraea*, *Pusillimonas*, *Pseudaminobacter*, *Sulfurivirga*, *Thiohalorhabdus*, and *Thiobacillus* may contribute to further tetrathionate depletion from the sediments (Tables S10 and S11 show the relevant references, and the metagenomic read percentages ascribed to these genera along SSK42/5 and SSK42/6 respectively).

Tetrathionate in the ASOMZ sediments can also be utilized as a respiratory substrate by bacteria such as *Citrobacter*, *Proteus* and *Salmonella*, which were detected by direct annotation of metagenomic reads (Figs. 1A and 1B). In addition, strains of *Enterobacter* such as those isolated as pure cultures from 275 cmbfs of SSK42/6 (Fig. 2) can add to the *in situ* reduction of tetrathionate to thiosulfate or sulfide. Furthermore, several such genera were also detected along SSK42/5 and SSK42/6 (via direct annotation of metagenomic reads), some members of which are known to respire via reduction of tetrathionate in the absence of O<sub>2</sub>; these included *Alteromonas*, *Alcaligenes*, *Desulfotomaculum*, *Desulfovibrio*, *Edwardsiella*, *Morganella*, *Pasteurella*, *Providencia*, *Serratia* and *Shewanella* (Tables S12 and S13 show the relevant references, and the metagenomic read percentages ascribed to these genera along SSK42/5 and SSK42/6 respectively).

### **3.3 The tetrathionate-forming/oxidizing microorganisms of the ASOMZ sediments are alive and active *in situ***

Aerobic slurry-culture of the sediment-samples of SSK42/5 and SSK42/6 in thiosulfate-containing artificial sea water (ASWT) medium resulted in either the formation of tetrathionate with no further oxidation of this polythionate, or the formation of tetrathionate followed by oxidation of the latter to sulfate, or no transformation at all. These findings, in conjunction with the results of pure culture isolation, illustrated that the sulfur-chemolithotrophic microorganisms present in these sediment-horizons are alive *in situ* and possess distinct pathways for oxidizing thiosulfate. For SSK42/5, ASWT-incubation of the 0, 15, 90 and 160 cmbfs samples resulted in the formation of tetrathionate as the sole and final product of thiosulfate oxidation, which happened *in vitro* at a

rate of 6.45-17.72  $\mu\text{mol S per L slurry per day}$  (Fig. 3A, Table S14). In contrast, ASWT-incubation of the 45, 60 and 295 cmbfs samples of SSK42/5 resulted in the initial formation of tetrathionate from thiosulfate at a rate of 1.11-6.45  $\mu\text{mol S per L slurry per day}$  (notably, no sulfate was produced during this period of incubation); subsequently, the accumulated tetrathionate was converted to sulfate at a rate of 5.86-13.75  $\mu\text{mol S per L slurry per day}$  (Fig. 3A, Table S15). Microbial communities present in the remaining five sediment-samples of SSK42/5 did not convert any thiosulfate out of the 20 mM S supplied (in the ASWT medium) to any higher oxidation state of sulfur. For SSK42/6, ASWT-incubation of the 120, 175 and 275 cmbfs samples resulted in the formation of tetrathionate as the sole and final product of thiosulfate oxidation, which happened *in vitro* at a rate of 17.2-29.71  $\mu\text{mol S per L slurry per day}$  (Fig. 3B, Table S16). In contrast, ASWT-incubation of the 2, 30 and 45 cmbfs samples of SSK42/6 resulted in the initial formation of tetrathionate from thiosulfate at a rate of 21.05-33.68  $\mu\text{mol S per L slurry per day}$  (notably, no sulfate was produced during this period of incubation); subsequently, the accumulated tetrathionate was converted to sulfate at a rate of 24-54  $\mu\text{mol S per L slurry per day}$  (Fig. 3B, Table S17). Microbial communities of the remaining seven sediment-samples of SSK42/6 did not transform any part of the thiosulfate supplied to any higher oxidation state of sulfur. This could be due to low number of metabolically active chemolithotrophic cells present in these samples. Notably, control incubation sets involving autoclaved sediment-samples in ASWT showed no change in the thiosulfate concentration of the medium.

Slurry-culture of a number of sediment-samples from SSK42/5 and SSK42/6 in ASWTr medium resulted in the oxidation of tetrathionate to sulfate. These data, together with the results of pure-culture isolation, indicated that the tetrathionate-oxidizing sulfur-chemolithotrophs present in the different sediment-samples were alive *in situ*. Of the individual microbial communities present in the different sediment-samples of SSK42/5, those from 0, 15, 45, 90, 120, 160 and 295 cmbfs oxidized tetrathionate at a rate of 2.5-23.5  $\mu\text{mol S per L slurry per day}$  (Fig. 3C, Table S18). Slurry-culture of samples from the remaining five sediment-depths of SSK42/5 did not result in any oxidation of tetrathionate, plausibly due to low number of metabolically active chemolithotrophic cells present in these samples. For SSK42/6, ASWTr-incubation of the 2, 30, 45, 60, 75 and 90 cmbfs sediment-samples resulted in the oxidation of tetrathionate to sulfate at an identical rate (approximately 140  $\mu\text{mol S per L slurry per day}$ ). While ASWTr-incubation of the samples from 120, 135, 160 and 175 cmbfs resulted in tetrathionate oxidation at a common rate of approximately 40  $\mu\text{mol S per L slurry per day}$ , the same for the 220, 250 and 275 cmbfs samples led to tetrathionate oxidation at a rate of approximately 75  $\mu\text{mol S per L slurry per day}$  (Fig. 3D, Table S19). Control incubations involving autoclaved sediment-samples in ASWTr showed no oxidation of tetrathionate.



The *in vitro* rates of tetrathionate formation, oxidation (see above) and reduction (see section 3.4 below) obtained by incubating the sediment-samples in specific media and culture conditions are expected to have little or no correspondence with the potential *in situ* rates of such processes. Rather, objective of these experiments was to check whether tetrathionate-metabolizing bacteria identified in this sediment system via culture-independent microbiological techniques were alive *in situ* (potential active state of the tetrathionate-metabolizers was subsequently corroborated by pure-culture isolations and metatranscriptome analysis). In the slurry-culture experiments it was peculiar to observe that the individual communities present within the sediment-depths spanning 2-90 cmbsf, 120-175 cmbsf, or 220-275 cmbsf, of SSK42/6 exhibited mutually identical rates of tetrathionate oxidation *in vitro*, despite having dissimilar composition/abundance of chemolithotrophic taxa. This could be explained as follows. When a natural sample is incubated in selective culture media (such as ASWTr) certain specific microbial species present in the sample often outgrow all metabolic competitors by virtue of higher substrate affinity and culture-condition suitability. Consequently, the growth/substrate-utilization phenotype(s) manifested by such enriched consortia are actually contributed to by the selected few rather than the entire community of metabolic equivalents present in the sample (Roy et al., 2016). In the light of this issue it seems plausible that distinct sets of chemolithotrophs more adept to growth in ASWTr medium are present across these sediment-samples, and it was only their characteristic rates of tetrathionate oxidation which were manifested as the *in vitro* tetrathionate oxidation rates of the communities.

Whatever may be the actual tetrathionate formation/oxidation rates of the sedimentary communities in their natural habitat, growths and activities in the slurry-culture experiments illustrated that the tetrathionate-forming and oxidizing bacteria detected along SSK42/5 and SSK42/6 are alive *in situ*. In order to further verify whether these metabolic-types were metabolically active in their native habitat, whole metatranscriptome of the 275 cmbsf sediment-sample of SSK42/6 was sequenced, and the paired end reads assembled into contigs. The gene-catalog obtained via annotation of the assembled contigs was found to encompass homologs of thiosulfate dehydrogenase (*tsdA*), which is involved in the conversion of thiosulfate to tetrathionate, and sulfate thiol esterase (*soxB*) and sulfur dehydrogenase (*soxC*), involved in tetrathionate oxidation (taxonomically, these homologs were ascribed to species of *Gramella*, *Halothiobacillus* and *Marinobacter*; see Table S20). Furthermore, from 275 cmbsf of SSK42/6, 15 such bacterial strains were isolated (Table 1) that could form tetrathionate from thiosulfate and/or oxidize tetrathionate to sulfate. 16S rRNA gene sequence-based taxonomic identification of the isolates clustered them under six species-level entities belonging to six distinct genera. Tetrathionate-forming and/or oxidizing phenotypes of one representative strain each from the six species-level clusters are shown in Fig. 2. The isolates belonging to the genera *Halomonas*,

620 *Methylophaga*, *Pseudomonas* and *Stenotrophomonas* oxidized thiosulfate only up to tetrathionate (Figs. 2A-2D); those belonging to *Pusillimonas* not only formed tetrathionate from thiosulfate but also oxidized tetrathionate to sulfate (Figs. 2E and 2F); the *Guyparkeria* isolates did not form tetrathionate from thiosulfate but oxidized tetrathionate to sulfate (Fig. 2G).

### 3.4 Active tetrathionate-reducing microorganisms in ASOMZ sediment

625 During anaerobic slurry-culture in RVTr medium, microbial communities of all the sediment-samples explored in SSK42/5 and SSK42/6 reduced tetrathionate to thiosulfate and/or sulfide at a rate of 0.5-1.5  $\mu\text{mol S per L slurry per day}$  (Figs. 3E and 3F, Tables S21 and S22); control sets involving autoclaved sediment-samples exhibited no depletion of tetrathionate from the RVTr media. No tetrathionate reductase (*ttrABC*) or thiosulfate reductase (*phsAB* or *psrA*) was  
630 detected in the gene-catalog obtained via assembly and annotation of the metatranscriptomic data from 275 cbsf of SSK42/6; but, the same catalog did contain many genes having highest sequence identities with homologs belonging to the typical tetrathionate-reducer *Salmonella*. Furthermore, anaerobic enrichment of the 275 cbsf sediment-samples of SSK42/6, followed by isolation of pure cultures, in RVTr medium yielded four such strains that reduced 30-32 mM S  
635 tetrathionate to equivalent amount of thiosulfate over 72 h of anaerobic incubation in RVTr medium (Fig. 2H shows the tetrathionate-reduction kinetics of the representative strain RVSM5a). 16S rRNA gene sequence-based taxonomic identification of the four isolates clustered them under a single species-level entity belonging to the genus *Enterobacter* (Table 1).

### 640 3.5 The tetrathionate-metabolizing bacteria isolated from 275 cbsf of SSK42/6 are widespread across SSK42/5 and SSK42/6

Whole genome sequencing and annotation was carried out for the three tetrathionate-forming isolates *Halomonas* sp. MCC 3301, *Methylophaga* sp. MTCC 12599 and *Pseudomonas bauzanensis* MTCC 12600; the two tetrathionate-oxidizing isolates *Guyparkeria* sp. SB14A, and  
645 *Pusillimonas ginsengisoli* MTCC 12558; and the tetrathionate-reducing isolate *Enterobacter* sp. RVSM5a (see Table 1 and Table S23 for GenBank accession numbers and general features of the genomes respectively). When metagenomic sequence data from the 25 distinct sediment-samples of SSK42/5 and SSK42/6 were mapped separately onto each of these six genomes, significant percentages of the metagenomic read-sets were found to match sequences from the  
650 individual genomes (Figs. 4 and 5, Table S23). In SSK42/5 and SSK42/6, 0.01-0.3% and 0.02-19.05% metagenomic reads from the individual sediment-samples mapped onto the different genomes respectively. Expectedly, prevalence of reads matching sequences from the new isolates was relatively higher for the metagenomes of SSK42/6 (Fig. 5); and within this core it was highest for 275 cbsf, i.e. the sample-site from where all the strains were isolated (Table

655 S23). These data corroborated the significant prevalence of the tetrathionate-metabolizing bacterial strains across the sediment-horizons of SSK42/5 and SSK42/6.

### 3.6 Correspondence between genomic sequences of the tetrathionate-metabolizing isolates and metatranscriptomic sequences from their habitat

660 In order to check whether the newly isolated bacterial strains were metabolically active and growing in their natural habitat, the rRNA-sequence-free metatranscriptomic sequence dataset (i.e., all paired end mRNA reads) obtained from the 275 cmbfsf sediment-sample of SSK42/6 was mapped onto six individual catalogs of tetrathionate-metabolizing and house-keeping genes curated from the genomes of the six different isolates. In this way, the metatranscriptomic  
665 sequence dataset was found to encompass reads matching the *tsdA* genes of the thiosulfate-to-tetrathionate-converting isolates *Halomonas* sp. MCC 3301 and *Pseudomonas bauzanensis* MTCC 12600, and the *soxB* and *soxC* genes of the tetrathionate-oxidizing isolates *Guyparkeria* sp. SB14A and *Pusillimonas ginsengisoli* MTCC 12558 (Table S24). In this context it is noteworthy that the annotated draft genome of the tetrathionate-forming isolate *Methylophaga* sp.  
670 MTCC 12599 did not contain any *tsdA* gene while that of the tetrathionate-reducing isolate *Enterobacter* sp. RVSM5a contained no *ttrABC* and *phsABC* genes; these could be attributable to either the incompleteness of the two genomes or the presence of novel thiosulfate-to-tetrathionate-converting and tetrathionate-reducing genes in the two bacteria respectively. On the other hand, mapping of the metatranscriptomic sequence dataset onto the individual isolates' gene-sets concerning core metabolisms such as (a) genetic information processing  
675 (transcription, translation, DNA replication and repair); (b) environmental information processing (ABC transporters, phosphotransferase system and bacterial secretion system) and (iii) cell growth and division (cell cycle), resulted in the concordant matching of up to 1.1 % metatranscriptomic read-pairs with the different sets. Table S24 shows the numbers of metatranscriptomic read-pairs that matched concordantly with representative house-keeping  
680 genes of the six isolates.

### 3.7 Thiosulfate as the key source of tetrathionate in ASOMZ sediments

685 Findings of the present culture-independent and culture-dependent investigations showed that microbe-mediated oxidation of thiosulfate to tetrathionate (by members of the bacterial genera included under group A in Fig. 6) could be a key metabolic process in the sulfur cycle of ASOMZ sediments. While genes for the metabolic conversion of thiosulfate to tetrathionate were present across the two cores, so were live microorganisms which could accomplish this process. Furthermore, metatranscriptomic data highlighted the potentially functional (metabolically active)  
690 state of thiosulfate-to-tetrathionate-converting bacteria *in situ*. Tetrathionate formed in this way

can have a number of fates: it can either be converted to sulfate (via chemolithotrophic oxidation by members of the genera included under Group B in Fig. 6) or reduced to thiosulfate and/or sulfide (by members of the genera included in Group C of Fig. 6). Whilst culture-independent as well as culture-dependent data supported the feasibility of these metabolic processes *in situ*, 695 copious hydrogen sulfide present in the pore-waters (Fernandes et al., 2018; also see Figs. 3G and 3H, Table S25) can potentially reduce tetrathionate to thiosulfate and elemental sulfur abiotically (Rowe et al., 2015). Corroborative to these possibilities, ion chromatographic analyses revealed up to 11.1  $\mu\text{M}$  thiosulfate in the pore-waters of all the sulfide-containing sample-sites of SSK42/5 and SSK42/6 (Figs. 3G and 3H, Table S25). These thiosulfate concentration values are 700 consistent with those reported from a number of physicochemically similar sediment-horizons across the global ocean (Troelsen and Jorgensen, 1982; Zopfi et al., 2004). Tandem absence of sulfide and thiosulfate in the upper 15 cmbsf of SSK42/6 could be due to potentially high rates of chemolithotrophic conversion of sulfide/thiosulfate to sulfate *in situ*. Tetrathionate was not found to exist freely in the pore-waters of SSK42/5 and SSK42/6; this is apparently attributable to the 705 fact that its build-up to measurable quantities [ $1 \mu\text{M}$  for the methods used in this as well as previous studies (Podgorsek and Imhoff, 1999)] could be debarred in natural environments due to high reactivity under the mediation of microbes (Kanao et al., 2007; Ghosh and Dam, 2009; Boden et al., 2010; Pyne et al., 2017, 2018) as well as naturally-occurring chemical substances such as sulfide (Roy and Trudinger, 1970; Podgorsek and Imhoff, 1999; Schippers et al., 1999; 710 Schippers and Jørgensen, 2001; Zopfi et al., 2004). From that perspective, the geomicrobiological information unearthed in this study illustrates the power of metaomics in discovering such cryptic interfaces between the chemosphere and the biosphere that are almost impossible to decipher from geochemical records alone.

## 715 **4 Discussions**

### **4.1 Biogeochemical perspectives of tetrathionate metabolism in ASOMZ sediments**

Concentrations and isotopic ratios of the various chemical constituents of sedimentary solid-phases and pore-fluids had long remained central to the deciphering of *in situ* biogeochemical pathways. Significant information on the carbon-sulfur cycles of modern as well as ancient, 720 marine and lacustrine sediments have been generated in this way. Currently, however, there is an increasing consensus that several questions in biogeochemistry - such as those concerning sulfur compounds oxidation/disproportionation, relative importance of simple fatty acids catabolism and anaerobic methane oxidation in sedimentary sulfate reduction, and biogeochemical processes within sulfate-methane transition zones - cannot be answered from 725 preserved geochemical records alone. In recent times a lot of advancement has taken place in our overall understanding of carbon-sulfur cycling in marine systems by virtue of data obtained

from metagenomic, metatranscriptomic, and *in situ* as well as *in vitro* geomicrobiological experiments. Forensic-level detection power of these approaches in unearthing such cryptic biogeochemical processes that do not get manifested, or leave their imprints, as detectable geological records, have been demonstrated in a number of recent papers (e.g., Canfield et al., 2010; Garcia-Robledo et al., 2017; Bhattacharya et al., 2019), which revealed such microbial community functions using metaomics approaches that were almost impossible to detect via geochemical analyses. So far as the present study is concerned, metagenome analysis along SSK42/5 and SSK42/6 revealed tetrathionate-metabolizing potentials in bacterial communities present at different depths of the sediment-horizons where tetrathionate is not detectable easily. At the same time, metatranscriptome analysis for the deepest sediment-sample of SSK42/6 indicated that these tetrathionate-metabolizing communities are potentially active *in situ*.

Regarding the biogeochemical feasibility of the three tetrathionate-metabolizing processes in the highly O<sub>2</sub>-constrained, sulfidic sediments of SSK42/5 and SSK42/6 (Fernandes et al., 2018) it is noteworthy that tetrathionate reduction is a mode of anaerobic respiration (Barrett and Clark, 1987; Hensel et al., 1999; Price-Carter et al., 2001) so has no issue with lack of O<sub>2</sub> in the chemical milieu; thiosulfate to tetrathionate conversion can also occur under both aerobic and anaerobic conditions (Sorokin et al., 1999). However, most of the sulfur chemolithotrophic bacteria known thus far, including some of those which form tetrathionate from thiosulfate and/or oxidize tetrathionate to sulfate, use O<sub>2</sub> as the terminal electron acceptor (Ghosh and Dam, 2009; Wasmund et al., 2017; Patwardhan et al., 2018). Albeit it is peculiar that such microorganisms are alive and active in these apparently O<sub>2</sub>-depleted sediment-horizons, aerobic respiration-related genes such as *aa<sub>3</sub>/cbb<sub>3</sub>*-type cytochrome-*c* oxidases (*coxABCD / ccoNOPQ*) and cytochrome-*bd* ubiquinol oxidase (*cydABX / appX*) are abundant in the metagenomes of all the sample-sites explored in SSK42/5 and SSK42/6 (Bhattacharya et al., 2019). Furthermore, when the metatranscriptomic sequence dataset obtained for the 275 cmbsf sediment-sample of SSK42/6 was assembled and annotated, the resultant contigs were found to encompass several homologs corresponding to genes for aerobic respiration by *aa<sub>3</sub>*-type and *cbb<sub>3</sub>*-type cytochrome-*c* oxidases and cytochrome-*bd* ubiquinol oxidase (Table S26), together with those for other O<sub>2</sub>-requiring (oxidase enzyme catalyzed) biochemical reactions (Table S27). These data, together with the isolation of obligately aerobic strains from this sedimentary ecosystem (Bhattacharya et al., 2019), indicated that cryptic O<sub>2</sub> source(s) supportive of aerobic metabolic processes are likely to be present *in situ*.

The nature and origin of the oxidants necessary to comprehensively realize the thiosulfate to tetrathionate, and tetrathionate to sulfate, oxidation potentials of the ASOMZ sediment microflora remain unresolved. Whilst there aren't many biochemical options for *in situ* O<sub>2</sub> production, nor there are high chances of O<sub>2</sub> influx to the system (Breuer et al. 2009; Cavan et al.

2017; Jessen et al. 2017), a plausible scenario in the form of cryptic aerobiosis by perchlorate-respiring microorganisms have been envisaged by Bhattacharya et al. (2019). In that study it has  
765 been postulated based on metagenomic, metatranscriptomic and co-culture-based data that perchlorate-respirers could be sustaining aerobic life in these sulfidic sediment-systems via cryptic O<sub>2</sub> supply-consumption partnership similar to the one reported for picocyanobacteria and nitrite-oxidizers in the acutely hypoxic waters of eastern tropical North and South Pacific (Garcia-Robledo et al. 2017). This said, in the absence of measured data for any potential oxidant,  
770 including O<sub>2</sub> or other oxyanions having possibilities of serving as electron acceptors, only future investigations of biogeochemistry focused on possible pathways of cryptic aerobiosis can resolve the modalities of sulfur compounds oxidation in these acutely O<sub>2</sub>-scarce environments.

#### **4.2 Trends of thiosulfate/sulfide concentration and prevalence of tetrathionate-metabolizing bacteria, along the sediment-cores, implicate tetrathionate as a key intermediate of the sulfur cycle**

  
775

Along both the sediment-cores SSK42/5 and SSK42/6, relative abundance of tetrathionate-forming, oxidizing, as well as reducing bacteria increased with depth (Figs. 1A and 1B). This could be attributable to the corresponding overall increase in the concentration of thiosulfate  
780 (Figs. 3G and 3H), which plausibly is the key source of biogenic tetrathionate in the sediment. Sulfide concentration also increased with depth, consistent with the increase of thiosulfate concentration (Figs. 3G and 3H). Sulfide is not only a potential direct source of thiosulfate in marine sediments (Jørgensen, 1990) but can also be a product of microbial tetrathionate reduction (Barrett and Clark, 1987; Price-Carter et al., 2001). Furthermore, sulfide, when present  
785 in sediment-cores, can chemically (abiotically) reduce tetrathionate to thiosulfate and elemental sulfur (Roy and Trudinger, 1970; Rowe et al., 2015). In view of these dependencies, depth-trends of sulfide concentration are expected to show a certain degree of correlation with trends of thiosulfate concentration.

In both the sediment-cores, relative abundance of tetrathionate-oxidizing bacteria  
790 increased with depth even as sulfate concentration decreased along the same trajectory (Figs. 1A, 1B, 3G and 3H). This indicated that the amounts of sulfate produced from potential tetrathionate oxidation at individual sediment-depths were far less than the amounts of sulfate that were reduced to sulfide *in situ*. Furthermore, in this context, it is noteworthy that in neither of the two cores, rates of *in vitro* tetrathionate formation or tetrathionate oxidation (in slurry-culture  
795 experiments) exhibited any parity with the trends of fluctuation observed for the relative abundance of tetrathionate-forming or oxidizing bacteria (Figs. 1A, 1B, 3A-3D). This could be reflective of the fact that the substrate-utilization rates manifested in the slurry-culture experiments actually resulted from the activities of the few chemolithotrophic species/strains that

were potentially enriched in the specific media used in these experiments and not that of the whole community present *in situ*. In contrast, however, rate of *in vitro* tetrathionate reduction along both the sediment-cores exhibited overall parity with the trends of fluctuation observed for the relative abundance of tetrathionate-reducing bacteria (both parameters showing overall increase with sediment-depth; Figs. 1A, 1B, 3E and 3F). This suggests that tetrathionate reduction rates observed in the slurry-cultures of the sediment-samples were attributable to all the tetrathionate-reducing species/strains present *in situ*, which in turn indicates a general ability of tetrathionate-reducers to grow in RVTr medium.

### 4.3 Potential abiotic drivers of tetrathionate formation in ASOMZ sediments

A number of other studies, based on geochemical experiments and preserved records, have also revealed the occurrence, and complex transformations, of tetrathionate and other intermediate sulfur species in ecologically diverse (but mostly non-sulfidic) environments, including marine sediments. Bak et al. (1993) measured tetrathionate, trithionate and thiosulfate in diverse natural samples, while Zopfi et al. (2004) used different techniques of analytical geochemistry to track the abiotic as well as biotic (microbe-mediated) transformations of sulfur cycle intermediates, including tetrathionate, in the sediments of Black Sea and North Sea. Findlay and Kamyshny (2017) envisaged the potential fates and transformation rates of intermediate sulfur species in lacustrine water-columns and sediments by introducing and tracking <sup>35</sup>S-labeled sulfur compounds in the samples. Furthermore, both Zopfi et al. (2004) and Findlay and Kamyshny (2017), in concurrence with previous reports (Schippers and Jørgensen, 2001), envisaged the *in situ* oxidation of pyrite (FeS<sub>2</sub>) by MnO<sub>2</sub> as an abiotic source of tetrathionate in the sediments. Such microbes-independent processes are also not improbable in the marine sediments explored in this study as (i) Fe (9232-17234 ppm), Mn (71-172 ppm) and pyrite (0.05-1.09 wt %) were all detected in the solid phase of SSK42/5 and SSK42/6 samples; (ii) genes for Mn(II) to Mn(IV) oxidation were identified in the assembled metagenomes of all the sediment-samples explored; (iii) sequences corresponding to manganese oxidase (*cotA*) and other accessory proteins involved in Mn(II) to Mn(IV) oxidation were there in the assembled metatranscriptome analyzed; and (iv) high percentage of reads ascribed to MnO-depositing bacteria such as *Aeromonas*, *Citrobacter*, *Enterobacter*, *Gallionella*, *Hyphomicrobium*, *Leptothrix* and *Proteus* (that reduce Mn<sup>+4</sup> to Mn<sup>+2</sup> for anaerobic respiration; Ghiorse, 1984), and MnO-to-MnO<sub>2</sub>-converting bacteria such as *Arthrobacter*, *Oceanospirillum* and *Vibrio* (Tebo et al., 2005; Sujith and Bharathi, 2011), were detected in all the metagenomes sequenced. Notably, however, manganese concentrations detected in the sediment-samples of SSK42/5 and SSK42/6 are orders of magnitude lower than the threshold [ $> 0.2\%$  (w/w)] reported previously for FeS<sub>2</sub> dissolution (Schippers and Jørgensen, 2001), so to confirm MnO<sub>2</sub>-FeS<sub>2</sub> interaction as a potent source of tetrathionate in these sediment-

835 horizons, future studies of geochemistry yielding fine-resolution depth-trends for pyrite and MnO<sub>2</sub>  
840 contents of the sediments are necessary alongside comprehensive data for pore-water metal  
ions concentrations and pyrite-specific (tracer) slurry incubations.

#### 4.4 Biogeochemical underpinnings of the cryptic status of tetrathionate in the sulfur cycle

840 Previous investigations based on the top few centimeters of Baltic Sea sediments had reported  
microbial production of tetrathionate and highlighted its role in the sulfur cycle (Podgorsek and  
Imhoff, 1999). Similar to the present scenario, in that study of sediments collected from coastal  
locations off Hiddensee (Germany), thiosulfate and tetrathionate were undetectable in the 7-8  
mM sulfate- and 4-20  $\mu$ M sulfide-containing native pore-water samples tested along a 6 cmbsf  
845 sediment-profile [detection limits for thiosulfate and tetrathionate, in the methods employed by  
Podgorsek and Imhoff (1999), were 0.5 and 1.0  $\mu$ M, respectively]. But when an approximately 15  
m<sup>2</sup> area of the same Baltic Sea location was covered with a plastic sheet for 5 months to  
artificially construct a stable anoxic condition, up to 5  $\mu$ M tetrathionate was found to accumulate  
within the 6 cmbsf sediment-profile explored, concomitant with the build-up of 80-280  $\mu$ M  
850 thiosulfate and 320-1200  $\mu$ M sulfide, and depletion of sulfate to a concentration of 0.4 mM within  
the first 4 cmbsf. In the above findings of Podgorsek and Imhoff (1999) it was peculiar that  
tetrathionate was absent in the native Baltic Sea (off Hiddensee coast) sediment-samples  
containing 4-20  $\mu$ M sulfide, but accumulated to a concentration of 5  $\mu$ M under induced anoxia  
involving 320-1200  $\mu$ M sulfide. Concurrently, in the other Baltic Sea study-site (Gotland Basin)  
855 explored by Podgorsek and Imhoff (1999), tetrathionate accumulation of up to 5 nmol cm<sup>-3</sup>  
sediment was recorded, during a period of annual anoxia, concomitant with increased  
concentrations of sulfide, elemental sulfur and thiosulfate. To explain these data Podgorsek and  
Imhoff (1999) invoked the cyclic sequence of reactions that was proposed and demonstrated  
previously (Roy and Trudinger, 1970; Hansen, 1974; Sorokin et al., 1996) and which involved  
860 bacterial oxidation of thiosulfate to tetrathionate on one hand and chemical reaction between  
sulfide and tetrathionate, forming elemental sulfur and thiosulfate, on the other. While it was  
added that accumulation of tetrathionate to high concentrations is debarred in euxinic marine  
environments owing to its high chemical reactivity with sulfide, and that even low concentrations  
of thiosulfate and tetrathionate acting as a catalytic couple were sufficient to promote large-scale  
865 net oxidation of sulfide to elemental sulfur (Podgorsek and Imhoff, 1999), questions remained as  
to how tetrathionate accumulation in the Baltic Sea sediments occurred only amidst heightened  
sulfide build-up. In the context of this apparent biogeochemical paradox, findings of the current  
study illuminate the centrality of microbe-mediated oxidation/reduction mechanisms (over and  
above the abiotic role of sulfide) in keeping the tetrathionate pool cryptic (below easily-detectable  
870 quantities) within the sediment systems.



## 4.5 Conclusion

Sulfur cycling is a crucial component of sediment biogeochemistry within the marine realm. Apart from controlling *in situ* sulfide-sulfate balance, microbe-mediated processes of the sulfur cycle work in conjunction with those of the carbon and nitrogen cycles to remineralize organic matter sequestered in the sea-bed, and also influence metal deposition/mobilization. Of the various redox species of sulfur, tetrathionate is seldom considered to be an important intermediate of sulfur cycling in marine sediments. The present study of the eastern Arabian Sea sediment system used microbiological data to reveal tetrathionate as a potential key intermediate of the sulfur cycle, and identify the biochemical pathways possibly involved in its formation and transformations *in situ*. Albeit the microbiological drivers and indicators underlying the potential status of tetrathionate as a key junction of sedimentary sulfur cycle were revealed here in the context of an oxygen minimum zone, there was no apparent reason to presume that such processes do not have their equivalents in other geomicrobiologically-distinct sediment-horizons of the marine realm. The absence of measured data for key *in situ* geochemical parameters disallowed any direct inference regarding the abiotic mechanisms potentially involved in the formation and transformation of tetrathionate in this sediment-system. However, a number of circumstantial evidences were there which could be indicative of pyrite (via abiotic reaction with  $\text{MnO}_2$ ) being a source of tetrathionate in the system. Consistent with previous reports from other analogous sediment-systems, tetrathionate was not detectable in the sulfidic ASOMZ sediments explored in the current study. In the context of this cryptic nature of tetrathionate in the sulfur cycle, our data revealed the key role of microbial redox metabolisms in preempting the accumulation of this highly-reactive polythionate in sediment pore-fluids, over and above the known abiotic mechanisms of tetrathionate scavenging by *in situ* sulfide. However, in the absence of any measured data for  $\text{O}_2$  or other oxyanions having possibilities of serving as electron acceptors, the biogeochemical modalities of the oxidative half of the tetrathionate cycle remain subject to future investigations of biogeochemistry focused on possible pathways of cryptic aerobiosis. Overall, appreciation of the scope and significance of our molecular and classical microbiology-based forecast of tetrathionate being a central intermediate of marine sedimentary sulfur cycle remains subject to further biogeochemical substantiation, which among other things should explore real-time the plausible presence of tetrathionate in the system.

## Supplementary material

Supplemental materials for this article may be found with the digital version of this manuscript. These include an MS Word file named “1\_Supplementary Information” and an MS Excel file named “2\_Supplementary Dataset”.

## Data availability

All nucleotide sequence data have been deposited in NCBI Sequence Read Archive (SRA) or GenBank under the BioProject accession number PRJNA309469: (i) the whole metagenome  
910 shotgun sequence datasets have the Run accession numbers SRR3646127 through  
SRR3646132, SRR3646144, SRR3646145, SRR3646147, SRR3646148, SRR3646150 through  
SRR3646153, SRR3646155 through SRR3646158, SRR3646160 through SRR3646165, and (ii)  
the metatranscriptome sequence dataset has the Run accession number SRR7991972. (iii) the  
915 whole genome sequences have the GenBank accession numbers SWAW01000000,  
SSXS01000000, RAPG00000000.1., SSXT01000000, SWAV01000000 and VTPT00000000.

**Code availability.** All data analysis codes used in this study are in the published domain, and have been appropriately cited in the text.

## Author contributions

920 W.G. conceived the study, designed the experiments, interpreted the results and wrote the  
paper. S.M., in conjunction with S.B., anchored the whole microbiological work, performed the  
experiments, analyzed the data, and contributed to paper writing. A.M led the entire SSK42  
mission and all geochemical investigations therein. C.R., M.J.R., J.S. and T.M. performed  
925 microbiological experiments and data analyses. S.F. and A.P. performed geochemical  
experiments. All authors read and vetted the manuscript.

## Competing interests

The authors declare that they have no conflict of interest.

## Acknowledgements

930 Financial support for conducting the microbiological studies was provided by given by Bose  
Institute via internal faculty grants and Earth System Science Organization, Ministry of Earth  
Sciences (MoES), Government of India (GoI) via grant number MoES/36/00IS/Extra/19/2013. We  
935 thank the Director, CSIR-National Institute of Oceanography for facilitating the geochemical  
studies and the research cruise SSK42 for acquisition of sediment-cores. MoES (GAP2303) also  
funded the research cruise. All the support received from the CSIR-NIO Ship Cell members and  
the crew members of SSK42 is gratefully acknowledged. S.B. received fellowship from Bose  
Institute. SM got fellowship from Department of Science and Technology, GoI. MJR and C.R. got  
940 fellowship from University Grants Commission, GoI. S.F. received fellowships from Council of  
Scientific and Industrial Research, GoI.

## References

- 945 Alam, M., Pyne, P., Mazumdar, A., Peketi A., and Ghosh, W.: Kinetic enrichment of  $^{34}\text{S}$  during proteobacterial thiosulfate oxidation and the conserved role of SoxB in S-S bond breaking, *Appl. Environ. Microbiol.*, 79, 4455-4464, <https://doi.org/10.1128/AEM.00956-13>, 2013.
- 950 Bak, F., Schuhmann, A., and Jansen, K. H.: Determination of tetrathionate and thiosulfate in natural samples and microbial cultures by a new, fast and sensitive ion chromatographic technique, *FEMS Microbiol. Ecol.*, 12, 257-264, <https://doi.org/10.1111/j.1574-6941.1993.tb00038.x>, 1993.
- Barrett, E. L., and Clark, M. A.: Tetrathionate reduction and production of hydrogen sulfide from thiosulfate, *Microbiol. Rev.*, 51, 192-205, 1987.
- 955 Baumgartner, L. K., Reid R. P., Dupraz, C., Decho, A. W., Buckley, D. H., Spear, J. R., Przekop, K. M., and Visscher, P. T.: Sulfate reducing bacteria in microbial mats: changing paradigms, new discoveries, *Sediment. Geol.*, 185, 131-145, <https://doi.org/10.1016/j.sedgeo.2005.12.008>, 2006.
- 960 Bhattacharya, S., Roy, C., Mandal, S., Rameez, M. J., Sarkar, J., Fernandes, S., Mapder, T., Alam, M., Roy, R., Mondal, N., Pyne, P., Halder, P. K., Peketi, A., Chakraborty, R., Mazumdar, A., and Ghosh, W.: Metabolically-active obligate aerobes in anoxic (sulfidic) marine sediments, *bioRxiv.*, <https://doi.org/10.1101/728287>, 728287, 2019.
- Boden, R., Kelly, D. P., Murrell, J. C., and Schäfer, H.: Oxidation of dimethylsulfide to tetrathionate by *Methylophaga thiooxidans* sp. nov.: a new link in the sulfur cycle, *Environ. Microbiol.*, 12, 2688-2699, <https://doi.org/10.1111/j.1462-2920.2010.02238.x>, 2010.
- 965 Boden, R., Scott, K. M., Williams, J., Russel, S., Antonen, K, Rae, A. W., and Hutt, L. P.: An evaluation of *Thiomicrospira*, *Hydrogenovibrio* and *Thioalkalimicrobium*: reclassification of four species of *Thiomicrospira* to each *Thiomicrothabodus* gen. nov. and *Hydrogenovibrio*, and reclassification of all four species of *Thioalkalimicrobium* to *Thiomicrospira*, *Int. J. Syst. Evol. Microbiol.*, 67, 1140-1151, <https://doi.org/10.1099/ijsem.0.001855>, 2017.
- 970 Breuer, E.R., Law, G.T., Woulds, C., Cowie, G.L., Shimmield, G.B., Peppe, O., Schwartz, M. and McKinlay, S.: Sedimentary oxygen consumption and microdistribution at sites across the Arabian Sea oxygen minimum zone (Pakistan margin). *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56, 296-304, <https://doi.org/10.1016/j.dsr2.2008.06.010>, 2009.
- 975 Brito, J. A., Denkmann K., Pereira, I. A., Archer, M., and Dahl, C.: Thiosulfate Dehydrogenase (TsdA) from *Allochromatium vinosum* structural and functional insights into thiosulfate oxidation, *J. Biol. Chem.*, 9222-9238, <https://doi.org/10.1074/jbc.M114.623397>, 2015.
- Canfield, D. E., Stewart, F. J., Thamdrup, B., De Brabandere, L., Dalsgaard, T., Delong, E. F., Revsbech, N. P., and Ulloa, O.: A cryptic sulfur cycle in oxygen-minimum-zone waters off the Chilean coast, *Science*, 330, 375-1378. <https://doi.org/10.1126/science.1196889>, 2010.
- 980 Cavan, E.L., Trimmer, M., Shelley, F. and Sanders, R.: Remineralization of particulate organic carbon in an ocean oxygen minimum zone. *Nat. Commun.*, 8, 1-9, <https://doi.org/10.1038/ncomms14847>, 2017.
- Cline, J. D.: Spectrophotometric determination of hydrogen sulfide in natural waters, *Limnol. Oceanogr.*, 14, 454-458, 1969.
- 985 De Jong, G. A. H., Hazeu, W., Bos, P., and Kuenen, G.: Polythionate degradation by tetrathionate hydrolase of *Thiobacillus ferrooxidans*, *Microbiology*, 143, 499-504. <https://doi.org/10.1099/00221287-143-2-499>, 1997.
- 990 Denkmann, K., Grein, F., Zigann, R., Siemen, A., Bergmann, J., van Helmont, S., Nicolai, A., Pereira, I. A., and Dahl, C.: Thiosulfate dehydrogenase: a widespread unusual acidophilic c-type cytochrome, *Environ. Microbiol.*, 14, 2673-2688, <https://doi.org/10.1111/j.1462-2920.2012.02820.x>, 2012.

- Euzéby, J. P.: List of Bacterial Names with Standing in Nomenclature: a folder available on the Internet, *Int. J. Syst. Evol. Microbiol.*, 47, 590-592, <https://doi.org/10.1099/00207713-47-2-590>, 1997.
- 995 Fernandes, S., Mazumdar, A., Bhattacharya, S., Peketi, A., Mapder, T., Roy, R., Carvalho, M. A., Roy, C., Mahalakshmi, P., Silva, R., Rao, P. S., Banik, S., and Ghosh, W.: Enhanced carbon-sulfur cycling in the sediments of Arabian Sea oxygen minimum zone center, *Sci. Rep.*, 8, 8665, <https://doi.org/10.1038/s41598-018-27002-2>, 2018.
- 1000 Findlay, A. J., and Kamysny, A.: Turnover rates of intermediate sulfur species ( $S_x^{2-}$ ,  $S^0$ ,  $S_2O_3^{2-}$ ,  $S_4O_6^{2-}$ ,  $SO_3^{2-}$ ) in anoxic freshwater and sediments. *Front. Microbiol.*, 8, 2551, <https://doi.org/10.3389/fmicb.2017.02551>, 2017.
- Garcia-Robledo, E., Padilla, C. C., Aldunate, M., Stewart, F. J., Ulloa, O., Paulmier, A., Gregori, G., and Revsbech, N. P.: Cryptic oxygen cycling in anoxic marine zones. *Proc. Natl. Acad. Sci. U.S.A.*, 114, 8319-8324, <https://doi.org/10.1073/pnas.1619844114>, 2017.
- 1005 Ghiorse, W. C.: Biology of iron-and manganese-depositing bacteria, *Annu. Rev. Microbiol.*, 38, 515-550, <https://doi.org/10.1146/annurev.mi.38.100184.002503>, 1984.
- Ghosh, W., Bagchi, A., Mandal, S., Dam, B., and Roy, P.: *Tetrathio bacter kashmirensis* gen. nov., sp. nov., a novel mesophilic, neutrophilic, tetrathionate-oxidizing, facultatively chemolithotrophic betaproteobacterium isolated from soil from a temperate orchard in Jammu and Kashmir, India, *Int. J. Syst. Evol. Microbiol.*, 55, 1779-1787, <https://doi.org/10.1099/ijs.0.63595-0>, 2005.
- 1010 Ghosh, W., Mandal, S., and Roy, P.: *Paracoccus bengalensis* sp. nov., a novel sulfur-oxidizing chemolithoautotroph from the rhizospheric soil of an Indian tropical leguminous plant. *Syst. Appl. Microbiol.*, 29, 396-403, <https://doi.org/10.1016/j.syapm.2005.10.004>, 2006.
- Ghosh, W., and Roy, P.: *Mesorhizobium thiogangeticum* sp. nov., a novel sulfur-oxidizing chemolithoautotroph from rhizosphere soil of an Indian tropical leguminous plant, *Int. J. Syst. Evol. Microbiol.*, 56, 91-97, <https://doi.org/10.1099/ijs.0.63967-0>, 2006.
- 1015 Ghosh, W., and Roy, P.: Chemolithoautotrophic oxidation of thiosulfate, tetrathionate and thiocyanate by a novel rhizobacterium belonging to the genus *Paracoccus*. *FEMS Microbiol. Lett.*, 270, 124-131, <https://doi.org/10.1111/j.1574-6968.2007.00670.x>, 2007.
- 1020 Ghosh, W., and Dam, B.: Biochemistry and molecular biology of lithotrophic sulfur oxidation by taxonomically and ecologically diverse bacteria and archaea, *FEMS Microbiol. Rev.*, 33, 999-1043, <https://doi.org/10.1111/j.1574-6976.2009.00187.x>, 2009, 2009.
- Ghosh, W., Roy, C., Roy, R., Nilawe, P., Mukherjee, A., Haldar, P. K., Chauhan, N. K., Bhattacharya, S., Agarwal, A., George, A., and Pyne, P.: Resilience and receptivity worked in tandem to sustain a geothermal mat community amidst erratic environmental conditions, *Sci. Rep.*, 5, 12179, <https://doi.org/10.1038/srep12179>, 2015.
- 1025 Gerhardt, P.: *Methods for General and Molecular Bacteriology*, American Society for Microbiology, Washington DC, 1994.
- Hansen, T. A.: *Sulfide als Electronendonor voor Rhodospirillaceae*, Proefschrift, University of Groningen, 1974.
- 1030 Hedrich, S., and Johnson, D. B.: *Acidithiobacillus ferridurans* sp. nov., an acidophilic iron-, sulfur- and hydrogen-metabolizing chemolithotrophic gammaproteobacterium, *Int. J. Syst. Evol. Microbiol.*, 63, 4018-4025, <https://doi.org/10.1099/ijs.0.049759-0>, 2013.
- 1035 Hensel, M., Hinsley, A. P., Nikolaus, T., Sawers, G., and Berks, B. C.: The genetic basis of tetrathionate respiration in *Salmonella typhimurium*, *Mol. Microbiol.*, 32, 275-287, <https://doi.org/10.1046/j.1365-2958.1999.01345.x>, 1999.

- Hensen, D., Sperling, D., Trüper, H. G., Brune, D. C., and Dahl, C.: Thiosulphate oxidation in the phototrophic sulphur bacterium *Allochromatium vinosum*, *Mol. Microbiol.*, 62, 794-810, <https://doi.org/10.1111/j.1365-2958.2006.05408.x>, 2006.
- 1040 Holmkvist, L., Ferdelman, T. G., and Jørgensen, B. B.: A cryptic sulfur cycle driven by iron in the methane zone of marine sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta.*, 75, 3581-3599, <https://doi.org/10.1016/j.gca.2011.03.033>, 2011.
- 1045 Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M. C., Rattei, T., Mende, D. R., Sunagawa, S., Kuhn, M., and Jensen, L. J.: eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences, *Nucleic Acids Res.*, 44, D286-D293, <https://doi.org/10.1093/nar/gkv1248>, 2016.
- 1050 Hyatt, D., Chen, G. L., LoCascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J.: Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11, 119, <https://doi.org/10.1186/1471-2105-11-119>, 2010.
- Jessen, G.L., Lichtschlag, A., Ramette, A., Pantoja, S., Rossel, P.E., Schubert, C.J., Struck, U. and Boetius, A.: Hypoxia causes preservation of labile organic matter and changes seafloor microbial community composition (Black Sea). *Sci. Adv.*, 3, p.e1601897, <https://doi.org/10.1126/sciadv.1601897>, 2017.
- 1055 Jørgensen, B. B.: A thiosulfate shunt in the sulfur cycle of marine sediments, *Science*, 249, 152-154, <https://doi.org/10.1126/science.249.4965.152>, 1990.
- Jørgensen, B. B., and Bak, F.: Pathways and microbiology of thiosulfate transformations and sulfate reduction in a marine sediment (Kattegat, Denmark), *Appl. Environ. Microbiol.*, 57, 847-856, 1991.
- 1060 Jørgensen, B. B., Findlay, A. J., and Pellerin, A.: The Biogeochemical Sulfur Cycle of Marine Sediments, *Front. Microbiol.*, 10, 849, <https://doi.org/10.3389/fmicb.2019.00849>, 2019.
- Kamyshny Jr, A.: Improved cyanolysis protocol for detection of zero-valent sulfur in natural aquatic systems. *Limnol. Oceanogr.: Methods*, 7, 442-448, <https://doi.org/10.4319/lom.2009.7.442>, 2009.
- 1065 Kaprálek, F.: The physiological role of tetrathionate respiration in growing *Citrobacter*, *Microbiology*, 71, 133-139, <https://doi.org/10.1099/00221287-71-1-133>, 1972.
- Kanao, T., Kamimura, K., and Sugio, T.: Identification of a gene encoding a tetrathionate hydrolase in *Acidithiobacillus ferrooxidans*, *J. Biotechnol.*, 132, 16-22, <https://doi.org/10.1016/j.jbiotec.2007.08.030>, 2007.
- 1070 Kanao, T., Kosaka, M., Yoshida, K., Nakayama, H., Tamada, T., Kuroki, R., Yamada, H., Takada, J., and Kamimura, K.: Crystallization and preliminary X-ray diffraction analysis of tetrathionate hydrolase from *Acidithiobacillus ferrooxidans*, *Acta Cryst. F.*, 69, 692-694, <https://doi.org/10.1107/S1744309113013419>, 2013.
- 1075 Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M.: KEGG as a reference resource for gene and protein annotation, *Nucleic Acids Res.*, 44, 457-462, <https://doi.org/10.1093/nar/gkv1070>, 2016.
- Kelly, D. P., and Wood, A. P.: Synthesis and determination of thiosulfate and polythionates, *Methods. Enzymol.*, 243, 475-501, [https://doi.org/10.1016/0076-6879\(94\)43037-3](https://doi.org/10.1016/0076-6879(94)43037-3), 1994.
- 1080 Kikumoto, M., Nogami, S., Kanao, T., Takada, J., and Kamimura, K.: Tetrathionate-forming thiosulfate dehydrogenase from the acidophilic, chemolithoautotrophic bacterium *Acidithiobacillus ferrooxidans*, *Appl. Environ. Microbiol.*, 79, 113-120, <https://doi.org/10.1128/AEM.02251-12>, 2013.
- Lahiri, C., Mandal, S., Ghosh, W., Dam, B., and Roy, P.: A novel gene cluster *soxSRT* is essential for the chemolithotrophic oxidation of thiosulfate and tetrathionate by

- 1085 *Pseudaminobacter salicylatoxidans* KCT001, Curr. Microbiol., 52, 267-273,  
<https://doi.org/10.1007/s00284-005-0176-x>, 2006.
- Langmead, B., and Salzberg, S.L.: Fast gapped-read alignment with Bowtie 2, Nat. Methods., 9,  
357–359, <https://doi.org/10.1038/nmeth.1923>, 2012.
- Li, D., Liu, C. M., Luo, R., Sadakane, K., and Lam, T. W.: MEGAHIT: an ultra-fast single-node  
1090 solution for large and complex metagenomics assembly via succinct de Bruijn graph,  
Bioinformatics., 31, 1674-1676, <https://doi.org/10.1093/bioinformatics/btv033>, 2015.
- Mason, J., and Kelly, D. P.: Thiosulfate oxidation by obligately heterotrophic bacteria, Microb.  
Ecol., 15, 123-134, <https://doi.org/10.1007/BF02011707>, 1988.
- 1095 Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E. M., Kubal, M., Paczian, T.,  
Rodriguez, A., Stevens, R., Wilke, A., and Wilkening, J.: The metagenomics RAST server—a  
public resource for the automatic phylogenetic and functional analysis of metagenomes,  
BMC Bioinformatics, 9, 386, <https://doi.org/10.1186/1471-2105-9-386>, 2008.
- Müller, F. H., Bandejas, T. M., Urich, T., Teixeira, M., Gomes, C. M. and Kletzin, A.: Coupling of  
1100 the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel  
membrane-bound thiosulphate: quinone Oxidoreductase, Mol. Microbiol., 53, 1147-1160,  
<https://doi.org/10.1111/j.1365-2958.2004.04193.x>, 2004.
- Nurk, S., Bankevich, A., Antipov, D., Gurevich, A. A., Korobeynikov, A., Lapidus, A., Prjibelski, A.  
D., Pyshkin, A., Sirotkin, A., Sirotkin, Y., and Stepanauskas, R.: Assembling single-cell  
1105 genomes and mini-metagenomes from chimeric MDA products, J. Comput. Biol., 20, 714-  
737, <https://doi.org/10.1089/cmb.2013.0084>, 2013.
- Oltmann, L. F., and Stouthamer, A. H.: Reduction of tetrathionate, trithionate and thiosulphate,  
and oxidation of sulphide in *Proteus mirabilis*, Arch. Microbiol., 105, 135-142,  
<https://doi.org/10.1007/BF00447128>, 1975.
- 1110 Parte, A. C.: LPSN—list of prokaryotic names with standing in nomenclature, Nucleic Acids Res.,  
42, 613-616, <https://doi.org/10.1093/nar/gkt1111>, 2013.
- Patwardhan, S., Foustoukos, D. I., Giovannelli, D., Yucel, M. and Vetriani, C.: Ecological  
succession of sulfur-oxidizing Epsilon-and Gammaproteobacteria during colonization of a  
shallow-water gas vent, Front. Microbiol., 9, 2970,  
<https://doi.org/10.3389/fmicb.2018.02970>, 2018.
- 1115 Podgorsek, L., and Imhoff, J. F.: Tetrathionate production by sulfur oxidizing bacteria and the role  
of tetrathionate in the sulfur cycle of Baltic Sea sediments, Aquat. Microb. Ecol., 17, 255-  
265, <https://doi.org/10.3354/ame017255>, 1999.
- Price-Carter, M., Tingey, J., Bobik, T. A., and Roth, J. R.: The Alternative Electron Acceptor  
Tetrathionate Supports B12 Dependent Anaerobic Growth of *Salmonella* enteric Serovar  
1120 *Typhimurium* on Ethanolamine or 1, 2-Propanediol, J. Bacteriol., 183, 2463–2475,  
<https://doi.org/10.1128/JB.183.8.2463-2475.2001>, 2001.
- Pyne, P., Alam, M., and Ghosh, W.: A novel *soxO* gene, encoding a glutathione disulfide  
reductase, is essential for tetrathionate oxidation in *Advenella kashmirensis*, Microbiol.  
Res., 205, 1-7, <https://doi.org/10.1016/j.micres.2017.08.002>, 2017.
- 1125 Pyne, P., Alam, M., Rameez, M. J., Mandal, S., Sar, A., Mondal, N., Debnath, U., Mathew, B.,  
Misra, A. K., Mandal, A. K., and Ghosh, W.: Homologs from sulfur oxidation (Sox) and  
methanol dehydrogenation (Xox) enzyme systems collaborate to give rise to a novel  
pathway of chemolithotrophic tetrathionate oxidation, Mol. Microbiol., 109, 169-191,  
<https://doi.org/10.1111/mmi.13972>, 2018.
- 1130 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner,  
F. O.: The SILVA ribosomal RNA gene database project: improved data processing and

- web-based tools, *Nucleic Acids Res.*, 41, 590-596, <https://doi.org/10.1093/nar/gks1219>, 2012.
- 1135 Quatrini, R., Appia-Ayme, C., Denis, Y., Jedlicki, E., Holmes, D. S., and Bonnefoy, V.: Extending the models for iron and sulfur oxidation in the extreme acidophile *Acidithiobacillus ferrooxidans*, *BMC Genomics*, 10, 394, <https://doi.org/10.1186/1471-2164-10-394>, 2009.
- 1140 Rameez, M. J., Pyne, P., Mandal, S., Chatterjee, S., Alam, M., Bhattacharya, S., Mondal, N., Sarkar, J. and Ghosh, W.: Two pathways for thiosulfate oxidation in the alphaproteobacterial chemolithotroph *Paracoccus thiocyanatus* SST, *Microbiol. Res.*, 230, 126345, <https://doi.org/10.1016/j.micres.2019.126345>, 2019.
- Rowe Annette, R., Chellamuthu, P., Lam, B., Okamoto, A., and Nealson, K. H.: Marine sediments microbes capable of electrode oxidation as a surrogate for lithotrophic insoluble substrate metabolism, *Front. Microbiol.*, 5, 784, <https://doi.org/10.3389/fmicb.2014.00784>, 2015.
- 1145 Roy, C., Alam, M., Mandal, S., Haldar, P.K., Bhattacharya, S., Mukherjee, T., Roy, R., Rameez, M. J., Misra, A. K., Chakraborty, R. and Nanda, A. K., Mukhopadhyay, S. K., and Ghosh, W.: Global association between thermophilicity and vancomycin susceptibility in bacteria, *Front. Microbiol.*, 7, 412, <https://doi.org/10.3389/fmicb.2016.00412>, 2016.
- 1150 Roy, C., Rameez, M. J., Haldar, P. K., Peketi, A., Mondal, N., Bakshi, U., Mapder, T., Pyne, P., Fernandes, S., Bhattacharya, S., Roy, R., Mandal, S., O'Neill, W. K., Mazumdar, A., Mukhopadhyay, S. K., Mukherjee, A., Chakraborty, R., Hallsworth, J. E., and Ghosh, W.: Microbiome and ecology of a hot spring-microbialite system on the Trans-Himalayan Plateau, *Sci. Rep.*, 10, 5917, <https://doi.org/10.1038/s41598-020-62797-z>, 2020.
- Roy, A. B., and Trudinger, P. A.: *The Biochemistry of Inorganic Compounds of Sulfur*, Cambridge University Press, Cambridge, 1970.
- 1155 Rudnicki, M. D., Elderfield, H., and Spiro, B.: Fractionation of sulfur isotopes during bacterial sulfate reduction in deep ocean sediments at elevated temperatures, *Geochim. Cosmochim. Acta.*, 65, 777-789, [https://doi.org/10.1016/S0016-7037\(00\)00579-2](https://doi.org/10.1016/S0016-7037(00)00579-2), 2001.
- 1160 Rzhepishevskaya, O. I., Valdés, J., Marcinkeviciene, L., Gallardo, C. A., Meskys, R., Bonnefoy, V., Holmes, D. S., and Dopson, M.: Regulation of a novel *Acidithiobacillus caldus* gene cluster involved in metabolism of reduced inorganic sulfur compounds, *Appl. Environ. Microbiol.*, 73, 7367-7372, <https://doi.org/10.1128/AEM.01497-07>, 2007.
- Schippers, A., Rohwerder, T., and Sand, W.: Intermediary sulfur compounds in pyrite oxidation: implications for bioleaching and biodepyritization of coal, *Appl. Microbiol. Biotechnol.*, 52, 104-110, <https://doi.org/10.1007/s002530051495>, 1999.
- 1165 Schippers, A., and Jørgensen, B. B.: Oxidation of pyrite and iron sulfide by manganese dioxide in marine sediments, *Geochim. Cosmochim. Acta.*, 65, 915-922, [https://doi.org/10.1016/S0016-7037\(00\)00589-5](https://doi.org/10.1016/S0016-7037(00)00589-5), 2001.
- 1170 Sorokin, D. Y., Teske, A., Robertson, L. A., and Kuenen, J. G.: Anaerobic oxidation of thiosulfate to tetrathionate by obligately heterotrophic bacteria, belonging to the *Pseudomonas stutzeri* group, *FEMS Microbiol. Ecol.*, 30, 113-123, <https://doi.org/10.1111/j.1574-6941.1999.tb00640.x>, 1999.
- Sorokin, D.: Oxidation of inorganic sulfur compounds by obligatory organotrophic bacteria, *Mikrobiologiya*, 72, 725-739, <https://doi.org/10.1023/B:MIC1.0000008363.24128.e52003>, 2003.
- 1175 Sorokin, D. Y., Robertson, L. A., and Kuenen J. G.: Sulfur cycling in *Catenococcus thiocyclus*, *FEMS Microbiol. Ecol.*, 19, 117-125, <https://doi.org/10.1111/j.1574-6941.1996.tb00204.x>, 1996.

- 1180 Stoffels, L., Krehenbrink, M., Berks, B. C., and Uden, G.: Thiosulfate Reduction in *Salmonella enterica* Is driven by the proton motive force, *Appl. Environ. Microbiol.*, 194, 475–485. <https://doi.org/10.1128/JB.06014-11>, 2011.
- Sujith, P. P., and Bharathi, P. L.: Manganese oxidation by bacteria: biogeochemical aspects. In *Molecular Biomineralization*, 52, 49-76, <https://doi.org/10.1007/978-3-642-21230-73>, 2011.
- 1185 Tebo, B. M., Johnson, H. A., McCarthy, J. K., and Templeton, A. S.: Geomicrobiology of manganese (II) oxidation, *Trends Microbiol.*, 13, 421-428, <https://doi.org/10.1016/j.tim.2005.07.009>, 2005.
- Tringe, S. G., von Mering, C., Kobayashi, A., Salamov, A. A., Chen, K., Chang, H. W., Podar, M., Short, J. M., Mathur, E. J., Detter, J. C., Bork, P., Hugenholtz, P., and Rubin, E. M.: Comparative metagenomics of microbial communities. *Science*, 308, 554-557, <https://doi.org/10.1126/science.1107851>, 2005.
- 1190 Tuttle, J. H.: Thiosulfate oxidation and tetrathionate reduction by intact cells of marine pseudomonad strain 16B, *Appl. Environ. Microbiol.*, 39, 1159-1166, 1980.
- 1195 Thamdrup, B., Finster, K., Fossing, H., Hansen, J. W., and Jørgensen, B. B.: Thiosulfate and sulfite distributions in porewater of marine sediments related to manganese, iron, and sulfur geochemistry, *Geochim. Cosmochim. Acta.*, 58, 67-73, [https://doi.org/10.1016/0016-7037\(94\)90446-4](https://doi.org/10.1016/0016-7037(94)90446-4), 1994.
- Tostevin, R., Turchyn, A. V., Farquhar, J., Johnston, D. T., Eldridge, D. L., Bishop, J. K., and McIlvin, M.: Multiple sulfur isotope constraints on the modern sulfur cycle, *Earth Planet. Sci. Lett.*, 396, 14-21, <https://doi.org/10.1016/j.epsl.2014.03.057>, 2014.
- 1200 Troelsen, H. and Jørgensen, B. B.: Seasonal dynamics of elemental sulfur in two coastal sediments. *Estuar. Coast Shelf S.*, 15, 255-266, [https://doi.org/10.1016/0272-7714\(82\)90062-2](https://doi.org/10.1016/0272-7714(82)90062-2), 1982.
- Vassiliadis, P.: The Rappaport-Vassiliadis (RV) enrichment medium for the isolation of salmonellas: An overview, *J. Appl. Bacteriol.*, 54, 69-76, <https://doi.org/10.1111/j.1365-2672.1983.tb01302.x>, 1983.
- 1205 van Zyl, L. J., van Munster, J. M., and Rawlings, D. E.: Construction of *arsB* and *tetH* mutants of the sulfur-oxidizing bacterium *Acidithiobacillus caldus* by marker exchange, *Appl. Environ. Microbiol.*, 74, 5686-5694, <https://doi.org/10.1128/AEM.01235-08>, 2008.
- Vishniac, W., and Santer, M.: The *Thiobacilli*. *Bacteriol. Rev.*, 21, 195-213, 1957.
- 1210 Wasmund, K., Mußmann, M., and Loy, A.: The life sulfuric: microbial ecology of sulfur cycling in marine sediments, *Environ. Microbiol. Rep.*, 9, 323-344, <https://doi.org/10.1111/1758-2229.12538>, 2017.
- 1215 Watsuji, T. O., Had, E., Miyazaki, M., Ichimura, M., and Takai, K.: *Thiomicrospira hydrogeniphila* sp. nov., an aerobic, hydrogen-and sulfur-oxidizing chemolithoautotroph isolated from a seawater tank containing a block of beef tallow, *Int. J. Syst. Evol. Microbiol.*, 66, 3688-3693, <https://doi.org/10.1099/ijsem.0.001250>, 2016.
- Zhu, W., Lomsadze, A., and Borodovsky, M.: *Ab initio* gene identification in metagenomic sequences, *Nucleic Acids Res.*, 38, e132, <https://doi.org/10.1093/nar/gkq275>, 2010.
- 1220 Zopfi, J., Ferdelman, T. G., and Fossing, H.: Distribution and fate of sulfur intermediates - sulfite, tetrathionate, thiosulfate, and elemental sulfur - in marine sediments, in: *Sulfur Biogeochemistry - Past and Present*, edited by: Amend, J. P., Edwards, K. J., and Lyons, T. W., Geological Society of America Special Paper 379, Boulder, Colorado, 97-116, 2004.



## Legend of figures

1225 **Figure 1. Relative abundance of the tetrathionate-metabolizing bacterial groups plotted**  
1230 **pair-wise against each other or against sediment-depth, along (A) SSK42/5 and (B)**  
**SSK42/6.** Relative abundances are expressed as the percentages of metagenomic reads  
matching protein-coding genomic sequences from the genera of tetrathionate-forming  
(*Pseudomonas* and *Halomonas*), tetrathionate-oxidizing (*Acidithiobacillus*, *Guyparkeria*,  
1235 *Halothiobacillus* and *Thiomicrospira*) and tetrathionate-reducing (*Citrobacter*, *Proteus* and  
*Salmonella*) bacteria. Presence of these genera in the various sediment-samples of SSK42/5 and  
SSK42/6 was corroborated via manual scrutiny of the amplified 16S rRNA gene sequence-based  
diversity data reported previously for these cores (Fernandes et al., 2018) as well as by  
searching the individual metagenomic sequence datasets against the 16S rRNA gene sequence  
1240 database of the Ribosomal Database Project (using BlastN with minimum alignment length 50  
bp, minimum identity cut-off 90% and maximum e-value cut-off  $1e^{-5}$ ). Plots corroborated by  
Pearson correlation coefficient (CC) and/or Spearman rank correlation coefficient (RCC) values  $\geq$   
 $+ 0.8$  with  $P < 0.05$  are shown in blue. Whereas none of the plots were corroborated by negative  
CC or RCC values numerically  $\geq 0.8$  with  $P < 0.05$ , those corroborated by positive/negative CC  
and/or RCC values numerically  $\leq 0.8$  are shown in black, irrespective of whether  $P$  is  $< 0.05$ . All  
CC and RCC values pertaining to SSK42/5 and SSK42/6 are given in Table S6 and Table S7  
respectively.

1245 **Figure 2. Time course formation and transformations of tetrathionate by the representative**  
**strains of the various species-level clusters isolated from 275 cmbsf of SSK42/6. (A**  
through **D)** thiosulfate to tetrathionate conversion by *Halomonas* sp. 15WGF, *Methylophaga* sp.  
SBPC3, *Pseudomonas* sp. SBBC and *Stenotrophomonas* sp. SBPC3 respectively. **(E and F)**  
oxidation of thiosulfate to sulfate via tetrathionate, and oxidation of tetrathionate to sulfate, by  
*Pusillimonas ginsengisoli* MTCC 12558 respectively. **(G)** oxidation of tetrathionate to sulfate by  
1250 *Guyparkeria* sp. SB14A. **(H)** reduction of tetrathionate to thiosulfate by *Enterobacter* sp.  
RVSM5a. —●—, —▲— and —▼— indicate the concentrations of sulfur (mM S) present in the  
medium at a given time-point of incubation in the form of thiosulfate, sulfate and tetrathionate  
respectively. —■— denotes the pH of a culture at a given time-point of incubation.

1255 **Figure 3. Rates of formation and transformations of tetrathionate in slurry-cultures of the**  
**sediment-samples, and concentrations of different sulfur species recorded along the two**  
**sediment-cores. (A and B)** *in vitro* rates of tetrathionate formation and its downstream oxidation  
(if any) in ASWT medium, for the different sedimentary communities of SSK42/5 and SSK42/6  
respectively. **(C and D)** *in vitro* rates of tetrathionate oxidation in ASWTr medium, for the different

1260 sedimentary communities of SSK42/5 and SSK42/6 respectively. (E and F) *in vitro* rates of  
tetrathionate reduction in RVTr medium, for the different sedimentary communities of SSK42/5  
and SSK42/6 respectively. (G and H) concentrations of sulfur species measured along SSK42/5  
and SSK42/6 respectively. In A through F, brown, violet and green circles represent tetrathionate  
1265 formation, oxidation and reduction rates respectively. In G and H, orange, light blue and purple  
circles represent the concentrations of sulfide, thiosulfate and sulfate respectively (sulfide and  
sulfate data were re-plotted from Fernandes et al., 2018).

**Figure 4. Bubble plot showing the percentages of metagenomic reads from individual  
sediment-samples of SSK42/5 that matched genomic sequences of the six tetrathionate-  
metabolizing bacterial isolates:** (*Ent*) *Enterobacter* sp. RVSM5a, (*Gpk*) *Guyparkeria* sp.  
1270 SB14A, (*Hlm*) *Halomonas* sp. MCC 3301, (*Mtp*) *Methylophaga* sp. MTCC 12599, (*Psb*)  
*Pseudomonas bauzanensis* MTCC 12600, and (*Psg*) *Pusillimonas ginsengisoli* MTCC 12558.  
Scales for sediment-depth (plotted in y-axis) and percentage of metagenomic reads from a given  
depth matching genomic sequences of an isolate (represented by bubble size) are both linear;  
1275 the individual isolates are placed along the x-axis.

**Figure 5. Bubble plot showing the percentages of metagenomic reads from individual  
sediment-samples of SSK42/6 that matched genomic sequences of the six tetrathionate-  
metabolizing bacterial isolates:** (*Ent*) *Enterobacter* sp. RVSM5a, (*Gpk*) *Guyparkeria* sp.  
1280 SB14A, (*Hlm*) *Halomonas* sp. MCC 3301, (*Mtp*) *Methylophaga* sp. MTCC 12599, (*Psb*)  
*Pseudomonas bauzanensis* MTCC 12600, and (*Psg*) *Pusillimonas ginsengisoli* MTCC 12558.  
Scales for sediment-depth (plotted in y-axis) and percentage of metagenomic reads from a given  
depth matching genomic sequences of an isolate (represented by bubble size) are both linear;  
the individual isolates are placed along the x-axis.

1285 **Figure 6.** Schematic diagram showing the network of biotic and abiotic process that are  
apparently involved in the formation and transformations of tetrathionate in the Arabian Sea  
sediments explored in this study.

**Table 1.** Bacteria isolated from 275 cmbsf of SSK42/6, and their tetrathionate-metabolizing properties.

	Bacteria isolated in ASWT		Bacteria isolated in ASWTY	Bacteria isolated in MSTY			Bacteria isolated in RVTr
	<i>Guyparkeria</i> sp.	<i>Methylophaga</i> sp.	<i>Halomonas</i> sp.	<i>Pseudomonas</i> sp.	<i>Stenotrophomonas</i> sp.	<i>Pusillimonas ginsengisoli</i>	<i>Enterobacter</i> sp.
Identification up to lowest taxonomic level possible	<i>Guyparkeria</i> sp.	<i>Methylophaga</i> sp.	<i>Halomonas</i> sp.	<i>Pseudomonas</i> sp.	<i>Stenotrophomonas</i> sp.	<i>Pusillimonas ginsengisoli</i>	<i>Enterobacter</i> sp.
Total number of strains isolated for the species-level cluster	6	2	2	2	1	2	4
Name of the representative strain	SB14A	SB9B = MTCC 12599	15WGF = MCC 3301	SBBB = MTCC 12600	SBPC3	SBSA = MTCC 12558	RVSM5a
16S rRNA gene sequence accession number of the representative strain	LN999387	LN999390	LT607031	LN999396	LN999400	LN999398	MH593841
Tetrathionate-metabolizing phenotype (medium in which phenotype was tested)	(i) Tetrathionate to sulfate (ASWTr)	(i) Thiosulfate to tetrathionate (ASWTM)	(i) Thiosulfate to tetrathionate (ASWTY)	(i) Thiosulfate to tetrathionate (MSTY)	(i) Thiosulfate to tetrathionate (MSTY)	(i) Thiosulfate to tetrathionate (MSTY) (ii) Tetrathionate to sulfate (MSTrY)	(i) Tetrathionate to thiosulfate (RVTr)
NCBI accession number for whole genome sequence	SWAW01000000	SSXS01000000	SSXT01000000	SWAV01000000	-	RAPG00000000	VTPT00000000

Figure 1





Figure 2



1310

1315

1320

1325

Figure 3




Figure 4




Figure 5

### SSK42/6








Figure 6



1330