

Carbon dioxide dynamics in an agricultural headwater stream driven by hydrology and primary production

Marcus B. Wallin^{1,2*}, Joachim Audet³, Mike Peacock¹, Erik Sahlée², Mattias Winterdahl^{2,4,5}

5

¹Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, Sweden

²Department of Earth Sciences, Uppsala University, Uppsala, Sweden

³Department of Bioscience, Aarhus University, Silkeborg, Denmark

⁴Department of Physical Geography, Stockholm University, Stockholm, Sweden

10 ⁵Bolin Centre for Climate Research, Stockholm, Sweden

Correspondence to: Marcus B. Wallin (marcus.wallin@slu.se)

Abstract. Headwater streams are known to be hotspots for carbon dioxide (CO₂) emissions to the atmosphere and are hence important components in landscape carbon balances. However, surprisingly little is known about stream CO₂ dynamics and emissions in agricultural settings, a land use type that globally cover ca 40% of the continental area. Here we present hourly measured in-situ stream CO₂ concentration data from a 11.3 km² temperate agricultural headwater catchment covering more than one year (in total 339 days excluding periods of ice and snow cover). The stream CO₂ concentrations during the entire study period were generally high (median 3.44 mg C L⁻¹, corresponding to partial pressures (*p*CO₂) of 4778 µatm) but were also highly variable (IQR = 3.26 mg C L⁻¹). The CO₂ concentration dynamics covered a variety of different time-scales from seasonal to hourly, and with an interplay of hydrological and biological controls. The hydrological control was strong (although with both positive as well as negative influences dependent on season) and CO₂ concentrations changed rapidly in response to rainfall and snowmelt events. However, during growing-season baseflow and receding flow conditions, aquatic primary production seemed to control the stream CO₂ dynamics resulting in elevated diel patterns. During the dry summer period, rapid rewetting following precipitation events generated high CO₂ pulses exceeding the overall median level of stream CO₂ (up to 3 times higher) observed during the whole study period. This finding highlights the importance of stream intermittency and its effect on stream CO₂ dynamics. Given the observed high levels of CO₂ and its temporally variable nature, agricultural streams clearly need more attention in order to understand and incorporate these considerable dynamics in large scale extrapolations.

15
20
25

1. Introduction

Fluvial systems (streams and rivers) are estimated to dominate the inland water CO₂ source globally, surpassing CO₂ emissions by lakes and reservoirs by a factor of six (Raymond et al. 2013). However, this estimate relies on a number of assumptions

30 and the scarcity of empirical data makes it uncertain. One of the critical gaps in the global upscaling is the lack of direct measurements from agriculture dominated areas (Osborne et al. 2010). Globally, agricultural land covers about 40% of the total continental area (Ramankutty et al., 2008) but there are few studies specifically focusing on the magnitude and dynamics of CO₂ emissions from agricultural streams. The few studies that do exist have shown e that agricultural stream CO₂ concentrations are generally high and up to 5 times greater than those in streams draining forested areas which are more extensively studied (Borges et al. 2018; Bodmer et al. 2016; Wallin et al. 2018). For example, Bodmer et al. (2016) measured partial pressure of CO₂ ($p\text{CO}_2$) in German and Polish streams and examined differences between forested and agricultural catchments. They found that $p\text{CO}_2$ was generally 2-3 times higher in agricultural streams compared to streams draining forested areas. Similarly, Borges et al. (2018) found high CO₂ concentrations in streams and rivers dominated by agriculture in the river system Meuse, Belgium. They linked the higher $p\text{CO}_2$ in agricultural streams to elevated levels of dissolved organic carbon (DOC), particulate organic carbon (POC) and inorganic nitrogen. On the other hand, Deirmendjian et al. (2019) showed that there was no difference in $p\text{CO}_2$ between forest and cropland streams in south-west France despite higher $p\text{CO}_2$ in forest groundwater compared to cropland groundwater. They explained the similar stream $p\text{CO}_2$ by more efficient gas exchange in the forest streams compared to the low-gradient cropland streams.

45 There are numerous factors influencing CO₂ patterns in stream systems and site-specific controls often dominate. Hence, large scale generalizations are difficult to make (Crawford et al. 2017). Based on high-frequency data, CO₂ concentrations in streams draining nutrient-poor forest and peatlands, as well as tropical forests, are often found related to variations in stream discharge but with site-specific response patterns, with CO₂ found either positively or negatively related to stream discharge (Crawford et al. 2017; Dinsmore et al. 2013; Johnson et al. 2007). These response patterns have often been connected to the catchment characteristics and changes in hydrological pathways, which in turn control the dominant source areas (both from a vertical and lateral point of view) of CO₂ in the catchment soils (Campeau et al. 2018; Leith et al. 2015; Dinsmore and Billett 2008). In contrast, other catchments lack a strong hydrological control and instead display clear diel cycles in stream CO₂ concentration indicating a metabolic control (Crawford et al. 2017). Here the interplay of photosynthesis and respiration (in-stream or terrestrial) could result in large day to night time differences in stream CO₂.

These recent findings concerning dynamics and controls on stream CO₂ concentrations have been possible due to the development of cost-effective CO₂ sensors (e.g. Johnson et al. 2010; Bastviken et al. 2015) which have enabled continuous data collection covering relevant time-scales (<hourly resolution). However, very little information about stream CO₂ dynamics exists from agricultural areas, a land use type that is heavily managed by humans from multiple aspects including hydrological drainage, nutrient additions, soil cultivation etc. As a consequence, CO₂ patterns in agricultural streams could potentially be very different than in other land use types with amplified diel CO₂ dynamics due to high metabolism and/or quicker response to hydrological events due to effective drainage systems.

In addition to the concentration gradient between the stream water and the above air, gas exchange is also highly dependent on the physical conditions at the air-water interface. For stream systems, the gas transfer velocity (often the variable given to describe the efficiency of the air-water gas exchange) is related to a combination of hydrological and morphological conditions of the stream channel, often including slope, velocity and water depth (Raymond et al. 2012; Wallin et al. 2011). All these variables are proxies for describing the turbulence of the stream water, which controls the gas exchange but that is rarely directly measured (Kokic et al. 2018). Agricultural areas are often located in flat landscapes resulting in drainage systems that are low-gradient and slow-flowing (Rhoads et al. 2003; Hughes et al. 2010), conditions that prevent effective air-water gas exchange (Hall & Ulseth, 2019). However, whether the elevated *p*CO₂ observed in agricultural streams is an effect of land use specific hydro-morphological stream conditions preventing efficient gas exchange or an effect of high internal (aquatic) or external (terrestrial) CO₂ production is currently unknown.

Although recent studies have identified agricultural streams as high *p*CO₂ systems, there are still large knowledge gaps to be filled in order to improve our understanding concerning the influence of these waterbodies in landscape C cycling. Here we present high-resolution (hourly) CO₂ concentration measurements in a Swedish agricultural headwater stream during more than a year (in total 339 days excluding periods of ice and snow cover). The study aimed to 1) quantify CO₂ concentration

levels in an agricultural stream and explore its temporal dynamics, 2) identify the main drivers causing temporal variability in
80 stream CO₂ concentration and how they might vary with season.

2. Methods

2.1. Study area

The study was conducted within the 11.3 km² Sundbromark (SBM) catchment (59°55'N, 17°32'E), located 5 km NW of the city of Uppsala, Sweden (Figure 1B). The catchment is a part of the hydro-meteorological observatory Marsta that was
85 established in the late 1940s (Halldin et al. 1999). The 30 year (1960-1991) mean annual temperature for the area is 5.3°C (mean January and July temperatures are -4.5 and 16.0°C, respectively) and with a mean annual precipitation of 535 mm. The length of the growing season is on average ca 210 days from early April to the end of October (SMHI). The catchment is dominated by agricultural land (86%) mainly used for cereal production and pasture, and with minor influence of forest (8%) and urban areas (6%). The area is flat with only 28 m elevation difference from 41 m.a.s.l. at the highest point to 13 m.a.s.l. at
90 the catchment outlet (Table 1). The bedrock consists of gneissic granites and the soils are dominated by post glacial clay at lower elevations and with some influence of glacial clay and silt at higher elevations. Although the bedrock does not contain any known carbonates, the soils are alkaline due to glacial carbonate containing deposits resulting in a stream pH ranging between 7.4 and 8.4 (Table 2), and with high electrical conductivity (EC, ranging 791-1908 μS cm⁻¹) (Osterman 2018). The nutrient and DOC levels of the stream water (Table 1) are at the lower end (within the 25th percentile) of monitored agricultural
95 catchments in Sweden (Linefur et al. 2018; Kyllmar et al. 2014). The oxygen conditions are mainly undersaturated (median D.O. = 53%) during the growing season. The arable fields are to a large extent artificially drained with extensive tile drainage pipe systems connected to the stream network.

To explore how representative the SBM catchment is for streams draining agricultural areas in the region, a snapshot sampling survey was performed across 10 streams (denoted region UPP 2 in the study by Audet et al. (2019)) of various sizes (catchment
100 area 8.5-740 km²) and agricultural influences (30-86%) distributed within a radius of 10 km from the city center of Uppsala (Figure 1A, Table S1).

2.2. Field sampling and analysis

The measurements were conducted from September 26, 2017 to December 12, 2018 (in total 339 days of measurements excluding periods of ice, and snow cover). Stream CO₂ concentration was monitored using an EosGP sensor (Eosense, Dartmouth, Canada). The sensor was covered by copper tape in order to avoid biofouling. Sensor accuracy is <1% of the calibrated range (0-2% CO₂) + 1% of the reading corresponding to a maximum error of ca 0.3 mg C L⁻¹ based on the maximum CO₂ measured in the current study. The CO₂ sensor was calibrated against known gas standards (400, 1000, 5000 and 20,000 ppm) before and after deployment. No significant drift (exceeding the above given uncertainty) in the instrument was observed during the period. Volume fraction outputs from the sensor were corrected for variations in temperature and pressure (atmospheric and water depth) using the method described in Johnson et al. (2010) and expressed in the unit of mg C L⁻¹.

Water level, water temperature and EC were measured together with CO₂ concentration at a V-notch weir (Figure S1). Water level was measured using a pressure transducer (1400, MJK Automation, Sweden) mounted in a stilling well representing the stream water level at the V-notch weir. Discharge was calculated from a stage-discharge rating curve based on a series of manual measurements and according to a rating curve presented in Holmqvist (1998). Water temperature and EC were monitored using a thermocouple (Type T) and a CS547A-L conductivity sensor (Campbell, UK), respectively. The sensors (except for the pressure transducer) were deployed under the water surface attached to a wooden rod in the center of the stream just upstream of the weir. All sensors were connected to a CR1000X data logger (Campbell, UK) measuring at a 1 min interval and storing average values at a temporal resolution of 30 (in 2017) or 60 (in 2018) minutes.

Stable isotopic analysis of the dissolved inorganic carbon (DIC) ($\delta^{13}\text{C}$ -DIC) was performed on six occasions during the falling limb of the snowmelt discharge peak in 2018 in order to explore the temporal variability in DIC source. At each sampling occasion a sample for analysis of $\delta^{13}\text{C}$ -DIC was taken in a 60 mL glass vial completely filled with stream water and closed airtight with a rubber septum below the water surface. In order to preserve the sample, 1 mL of highly concentrated ZnCl₂ solution was injected in each sample (with subsequent release of 1 ml of sample in order to keep atmospheric pressure) directly after sample collection. Samples were kept cold and dark until analysis. Prior to analysis, 2 mL of sample was injected into 12

mL septum-sealed pre-combusted glass vials (Labco Limited) pre-filled with He gas, and pre-injected with 0.1 mL of concentrated phosphoric acid in order to convert all DIC species to CO₂(g) (Campeau et al. 2017a). The samples were analyzed using an isotope ratio mass spectrometer (DeltaV Plus, Thermo Fisher Scientific, Bremen, Germany) Gasbench II (Thermo
130 Fisher Scientific, Bremen, Germany) measuring the CO₂ in the headspace. Each sample was analyzed seven times (sample volume; 100 µL per sample) and the first two injections for each sample were discarded to avoid memory effects, and the mean was taken of the other five to give the final result. The δ¹³C-DIC values are given in terms of deviation from known carbonate standards in per mille where R is the isotopic ratio of [¹³C]/[¹²C]:

$$\delta^{13}\text{C-DIC (‰)} = (R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$$

135 Precipitation, air temperature and incoming shortwave (global) radiation data (Figure 2) were obtained from the Marsta meteorological observatory located within the catchment ca 2.5 km from the stream sampling station (Halldin et al. 1999). In the absence of direct measurements of photosynthetically active radiation (PAR) shortwave incoming radiation was used as a proxy for available photosynthetic light.

140 A spatial sampling campaign for CO₂ concentration, pH, EC and water temperature was conducted on June 21, 2018 across ten agricultural streams (including the SBM stream) located in different catchments around the city of Uppsala (Figure 1A). The sampling was performed between 10.00 and 14.00 during the day. Samples for CO₂ analysis were collected using the headspace method (Hope et al., 2004; Kokic et al. 2015). Briefly, 30 mL bubble-free water were collected in 60 mL polypropylene syringes and equilibrated with a known volume of ambient air by shaking vigorously for 1 min. The equilibrated
145 headspace (15-20 mL) was recovered and analysed on an Ultraportable Greenhouse Gas Analyzer (UGGA) (Los Gatos Research, USA) equipped with a soda lime filter and manual injection port. In situ CO₂ concentration was calculated from the UGGA-determined ppm values using Henry's law considering stream temperature (Weiss 1974), atmospheric pressure, the added ambient air, as well as the water-air volume ratio in the syringe. pH, EC and water temperature were measured in-situ in the streams with handheld instruments, for pH with a pH110 pH-meter (VWR, USA), and for EC and temperature with a
150 HI 99300 (Hanna Instr., USA).

2.3. Delineation of the stream network and catchment characteristics

Catchment area and characteristics were calculated in QGIS 3.8 based on a high resolution (2×2 m) digital elevation model (DEM) derived from LIDAR data (GSD Elevation data, grid 2+, Swedish Land Survey). Land use distribution within the catchment was derived from the CORINE Land Cover 2018 product (European Environment Agency), and soil and bedrock characteristics were based on digital versions of the Quaternary deposits (1:25,000 – 1:100,000) and bedrock (1:50,000 – 1:250,000) maps (Swedish Geological Survey).

2.4. Data analysis

Out of the total data set (339 days) from the SBM catchment, only data measured at discharge rates $> 0 \text{ L s}^{-1}$ (i.e. excluding standing water or completely dry conditions) were used in the analysis of the stream CO_2 data (Figure S1). For further evaluation of the control on stream CO_2 concentration, the data set was divided into four periods (Autumn (49 days), Snowmelt (17 days), Spring (91 days) and Dry period (138 days)) according to distinct phases in the hydrograph (Figure 3, Table S2). The stream CO_2 dynamics observed among the different periods were examined visually and any hydrological controls on the CO_2 were identified by the presence and direction of CO_2 -discharge hysteresis loops (Evans and Davies, 1998). Similar hysteresis analysis was used to investigate diel patterns in the CO_2 concentration data. Spearman's rank correlation coefficient was used to test for monotonic relationships between the diel amplitude in stream CO_2 concentration and potential drivers. Correlations were considered significant if $p < 0.05$. The software JMP 14.2.0 (SAS Institute Inc., Cary, NC, USA) was used for all statistical calculations.

3. Results

The mean air temperature and total precipitation for the entire period (September 26, 2017-December 12, 2018) were $6.8 \text{ }^\circ\text{C}$ and 704 mm, respectively. The summer and autumn of 2018 were dry with generally low precipitation, the exception was on July 29 with 82 mm rain within 24 hours (Figure 2). Mean and median stream discharge for study period were 30.6 and 0.9 L s^{-1} , respectively, and with a total range from 0 to 668 L s^{-1} (corresponding to a range from 0 to 5.0 mm day^{-1}) (Figure 3). However, due to high water table exceeding the range of the pressure transducer the absolute peak discharge occurring during April 5 to April 7 was missed in the measurements. The large skewness between mean and median discharge was an effect of

175 the large number of days without waterflow over the weir during the summer and autumn 2018, 128 days (38%) out of the study period. According to frequency analysis, 67% of the days had a mean daily discharge $<5 \text{ L s}^{-1}$. Despite the few days with discharge $>100 \text{ L s}^{-1}$ (7% of the entire period), those days accounted for 69% of the accumulated discharge. The majority (84%) of these high discharge days occurred during the snowmelt in April.

3.1. General CO₂ patterns

180 The stream CO₂ concentrations during the entire study period (median and mean 3.44 mg C L^{-1} and 3.94 mg C L^{-1} , respectively, corresponding to a $p\text{CO}_2$ of $4778 \text{ } \mu\text{atm}$ and $5324 \text{ } \mu\text{atm}$) were highly variable (IQR = 3.26 mg C L^{-1}) (Figure 3) and displayed a bimodal distribution with frequency peaks at $\sim 2.7 \text{ mg C L}^{-1}$ and $\sim 6.1 \text{ mg C L}^{-1}$ (Figure S2). The lower peak was associated with the snowmelt and spring period, whereas the higher peak was attributed to the autumn period 2017 and to rain events during the dry period of summer/autumn 2018. In addition to the bimodal shape a very distinct peak in frequently measured
185 concentrations was observed at $\sim 1.6 \text{ mg C L}^{-1}$. This peak was attributed to the minimum concentrations values for the diel cycles observed during the spring period.

3.2. Controls on stream CO₂ concentration

The autumn period started dry with low discharge ($<3 \text{ L s}^{-1}$) for the initial month of measurements. The CO₂ concentrations were at the same time highly dynamic but unrelated to variations in discharge. The CO₂ concentration reached the maximum
190 for the autumn ($10.89 \text{ mg C L}^{-1}$, which was also the maximum for the entire study period) during late October followed by a decline in CO₂ to ca 2 mg C L^{-1} in early November. During November and December four main rain events were identified which all displayed an increasing stream CO₂ concentration with increasing discharge. In three of these events a positive clockwise hysteresis loop was observed (Figure 4) where the CO₂ concentration reached its maximum before the discharge did. At the last event during the autumn 2017, the relationship between CO₂ concentration and discharge was close to linear, but still
195 positive. During the snowmelt period the hydrograph was characterized by a diel cycle with melting during day-time resulting in daily discharge peaks which were suppressed during night-time freezing. In contrast to the autumn events the daily discharge peaks were negatively related to the stream CO₂ concentration, and with an anti-clockwise hysteresis loop where the minimum CO₂ concentration was reached before the highest discharge of the event (Figure 5). After the snowmelt discharge peak the

spring and early summer periods (late April to early July) were dry with limited precipitation and with a steady decline in runoff (Figure 3). During this period the CO₂ concentration displayed a pronounced diel cycle with daily maximum and minimum CO₂ concentrations reached during early mornings (06:00) and late afternoons (18:00), respectively (Figure 6). The medium amplitude of the diel CO₂ cycle for this period was 2.03 mg C L⁻¹, corresponding to pCO₂ = 2974 μatm (IQR = 1.23 mg C L⁻¹, corresponding to pCO₂ = 2212 μatm), and with the size of the diel CO₂ concentration amplitude being related to both the daily mean water temperature and the shortwave radiation (Figure 7). The diel pattern displayed a clear negative anti-clockwise CO₂-streamwater temperature hysteresis loop, where the median CO₂ concentration could differ up to 75% between day and night-time although being measured at the same stream water temperature (Figure 8).

From early July the stream dried out and hence no runoff over the V-notch weir was generated. During this period the CO₂ sensor was mostly recording an atmospheric signal. However, for five rain events during the summer and early autumn runoff was generated which allowed stream CO₂ determination for shorter periods (Figure 9). During these runoff events (< 2 days long) high CO₂ concentration pulses were recorded (up to 11 mg C L⁻¹). At all events CO₂ was recorded for a longer period than the discharge as the small dam above the v-notch weir was still water-filled for some time after runoff over the weir ceased. Also, common for all events was that the stream CO₂ concentration continued to increase although the discharge peak had passed. During July 29 a heavy rain storm occurred with 82 mm precipitation in 24 hours. Although more than 15% of the long-term annual mean precipitation fell during one day, low discharge was generated (maximum discharge 6.1 L s⁻¹) due to high evapotranspiration and dry soils (Figures 3 and 9). However, the rainstorm event resulted in close to the highest stream CO₂ concentration (10.81 mg C L⁻¹) being observed during the studied period. As soon as the stream was more permanently refilled in early December and with discharge generated over the weir, the stream CO₂ concentration was back to similarly high levels (typically 5-8 mg C L⁻¹) as observed in the autumn of 2017.

220 3.3. Sources of DIC

The δ¹³C-DIC data collected during the falling limb of the spring discharge peak (discharge range 130-9.6 L s⁻¹) were ranging from -13.8 to -12.2‰. This narrow range suggests a relatively constant source of inorganic C during the spring period.

Although there was a tendency towards more negative $\delta^{13}\text{C-DIC}$ values at higher discharge, no significant relationship was found (Figure 10). $\delta^{13}\text{C-DIC}$ was also unrelated to the stream CO_2 concentration (data not shown).

225 **3.4. Spatial representativeness**

The ten streams manually sampled around Uppsala displayed a wide range in CO_2 concentrations (1.8-4.6 mg C L⁻¹) on the day of sampling (June-21 2018), and with the SBM stream (site 3 in Table S1) being close to the overall median (SBM, 2.7 mg C L⁻¹; overall median, 3.0 mg C L⁻¹) (Table S1). Furthermore, the CO_2 concentration manually sampled at SBM was close to the sensor recorded CO_2 (2.59 mg C L⁻¹) at the hour of sampling. The SBM stream was also close to the spatial median
230 DOC concentration but slightly elevated in NO_3 and PO_4 . The CO_2 concentration was on a spatial scale related to pH but unrelated to catchment area or land use distribution within the catchment. Furthermore, the CO_2 concentration was on a spatial scale unrelated to mean stream concentrations of DOC, PO_4 and NO_3 , although these variables were sampled during a different period than the CO_2 .

4. Discussion

235 In order to produce large scale estimates of the exchange of GHGs between inland surface waters and the atmosphere, a basic requirement is to know the aqueous concentrations of the gases of interest and how they might vary over time. Headwater streams have been identified as “hotspots” for CO_2 emissions (Raymond et al. 2013; Wallin et al. 2018), but there is limited data capturing the temporal resolution, specifically from streams draining agricultural regions, making large scale generalizations uncertain. Due to effective drainage, high nutrient conditions and often high sun-light exposure (due to limited
240 tree cover), agricultural streams could potentially be very different in their CO_2 dynamics compared with streams draining other environments. Here we continuously measured stream CO_2 concentration in a headwater catchment dominated by agricultural land use (86%) covering more than one year of the snow-free period. In line with findings from similar studies from other environments (arctic tundra, boreal forest, temperate peatlands, alpine) (e.g. Rocher-Ros et al. 2019; Riml et al. 2019; Crawford et al. 2017; Peter et al. 2014; Dinsmore et al. 2013) we found a mixture of controls on stream CO_2 operating
245 at different time-scales generating a highly dynamic stream CO_2 concentration pattern. These time-scales covers seasonal patterns to diel cycles, or even shorter scales associated with discharge events. Both the magnitude of CO_2 concentrations, and

their associated temporal dynamics were found to be high in the current agricultural stream when compared with the literature. The mean CO₂ concentration (3.94 mg C L⁻¹ corresponding to a pCO₂ of 5324 µatm) is at the high end when compared with other high-frequency CO₂ data sets covering low-order (<3rd stream order) catchments draining multiple environments, including arctic tundra, boreal forest, hemi-boreal forest, temperate forest, temperate peatlands and alpine areas (typically ranging from ca 0.2 to 6 mg C L⁻¹) (Crawford et al. 2017; Natchimuthu et al. 2017; Peter et al. 2014; Dinsmore et al. 2013). Still, CO₂ concentrations in SBM do not seem to be exceptionally high compared to snapshot-based data from other agricultural streams.

The spatial variability seen in this study, although only based on snapshot samples, and previous studies indicate that CO₂ concentrations in agricultural streams are comparably high (Borges et al. 2018; Bodmer et al., 2016; Sand-Jensen & Staehr, 2012). In addition, the observed temporal dynamics presented here are, to our knowledge, among the most pronounced in the literature, although the number of high-frequency stream CO₂ data sets are limited. For example, the rapid decrease in stream CO₂ during the autumn of 2017, the strong diel cycle (diel amplitude up to almost 5.0 mg C L⁻¹) during the spring/early summer period, or the rapid and high CO₂ pulses (up to 11.0 mg C L⁻¹) occurring in accordance to rain events during the dry late summer/autumn period. These high CO₂ dynamics clearly illustrate the need for continuous high frequency CO₂ concentration measurements in streams in general, and in agricultural streams more specifically. Without such high-frequency data, representative estimates of agricultural stream CO₂ will be associated with high uncertainty. Although based on measurements from a single stream, these findings in turn indicate that current large-scale stream CO₂ emission estimates (e.g. Raymond et al. 2013; Humborg et al. 2010), which are largely based on snapshot concentration data with low (or no) resolution in time, might be specifically uncertain for agricultural regions.

According to our continuous data the highly dynamic pattern in stream CO₂ concentration is driven by a complex interplay of hydrology and biology. The high autumn concentrations observed both in 2017 and 2018 are likely an effect of high respiration of organic matter in the stream channel and/or in the adjacent soil water (Figure 3D). This is supported by efficient aquatic microbial DOC degradation (<800 µg C L⁻¹ d⁻¹) observed during the autumn period across the ten streams (agricultural land

use, 30-86%) included in the spatial sampling campaign (Peacock et al. unpublished 2020). This could be compared with organic C degradation rates determined in boreal forest and mire streams displaying typically lower rates ($<300 \mu\text{g C L}^{-1} \text{d}^{-1}$, Berggren et al. 2009). The positive CO_2 -discharge relationships indicated that event flow pathways were in contact with soils with higher concentrations of CO_2 compared to flow pathways during base flow (Evans & Davies, 1998; Seibert et al., 2009). Also, the clock-wise shape of the hysteresis loop suggests that there is a buildup of CO_2 in the catchment that is flushed out during rain events (Figure 4). The CO_2 pool seems to be limited as the CO_2 concentration drops before the maximum discharge peak occurs, or that vertical patterns in the CO_2 soil profile control the stream CO_2 dependent on dominating flow paths (Evans and Davies, 1998; Öquist et al. 2009). This could explain that the stream CO_2 increase did not reach any source limitation at rain events of lower magnitude (Figure 4D). Similar positive CO_2 concentration-discharge patterns have been observed across different low-order streams (e.g. Crawford et al. 2017; Dinsmore et al. 2013), but the absolute patterns are often concluded to be highly site-specific and even event-specific. Here we suggest, by exploring the hysteresis loops, that such positive relationships are influenced by the size of the available catchment CO_2 pool or the hydrological connectivity to it. In a highly drained low-elevation agricultural landscape where much of the stream runoff is generated through drainage pipes (Castellano et al. 2019), the extent and spatial distribution of these terrestrial source areas and connections between ground- and surface water are central for the CO_2 patterns observed in the stream. Strong hydrological control has been found for DOC in agricultural streams in USA and France, where high discharge events flush allochthonous DOC, via subsurface drainage pipes, into streams (Morel et al. 2009; Royer & David, 2005). In contrast to the seasonally variable CO_2 -discharge response patterns observed in the current study, Morel et al. (2009) suggested that stream DOC is non-limited and would continue to rise until the maximum discharge peak is reached. Whether this discrepancy in source limitation between CO_2 and DOC (although based on different studies and environments) indicate differences in the source areas of the different carbon components require further investigation.

In contrast to the patterns observed during the autumn, during the snow melt period the stream CO_2 was diluted when discharge increased following a diel pattern (Figure 5). The melting and freezing between day and night-time suggests that melt-water from the surface snowpack during day time to a larger extent reached the stream without picking up an elevated CO_2 signal.

Similar dilution patterns in conjunction with snowmelt have been observed in catchments of various land use but specifically in peatland catchments with limited forest cover (e.g. Wallin et al. 2013). The similarity between this agricultural catchment and open peatlands could potentially be the effect of an efficient melting of the snowpack. Both non-forested peatlands and agricultural fields are open areas subject to direct sunlight, and wind and rain exposure, while the soil under the snow remains frozen. As a result, a large share of the melt-water will never infiltrate the soil but instead reach the surface drainage system as overland flow (Laudon et al. 2007). This is further accompanied by the low hydraulic conductivity of clay soils, which are dominating the catchment of the current study. Although we did not capture the 2-3 days of peak spring flood (due to a water level out of the range of the pressure transducer) it was evident that the stream CO₂ concentration was diluted from ca 6.0 mg C L⁻¹ to ca 2.0 mg C L⁻¹ during these days, something that is further supported by the similar drop in EC during the peak spring flood from ca 900 to ca 150 µS cm⁻¹. However, as soon as the discharge peak passed, the stream CO₂ concentration recovered rapidly to the pre-peak levels suggesting a shift to hydrological pathways that mobilize a high CO₂ pool, again supported by the concurrent increase in EC. April and May 2018 were characterized by warm and clear weather with an average 4.2°C higher air temperature and 255 more sun hours than the 30-year mean (1961-1990, SMHI). Altogether, this stimulates a kick-start of the aquatic primary production upon snowmelt, which likely explains the steady decline in CO₂ that occurred during late April/early May. During the spring and early summer, a strong diel pattern in CO₂ concentration further developed, likely driven by aquatic primary production consuming CO₂ during day-time. Such diel CO₂ patterns are commonly observed in stream CO₂ time series at base-flow or during receding flow conditions (e.g. Riml et al. 2019; Peter et al. 2014) and are especially pronounced in amplitude in nutrient-rich streams or in streams without canopy shading (Alberts et al. 2017; Crawford et al. 2017; Rocher-Ros et al. 2019). Initial evaluation of the δ¹³C-DIC data collected during the spring period suggests a relatively steady mixture of geogenic and biogenic DIC although somehow related to variations in discharge (Figure 10). However, given the suppressed stream CO₂ during the spring period, together with the strong diel cycle caused by aquatic primary production, fractionation of a strict biogenic DIC pool (with a δ¹³C-DIC from -28 to -20‰) could theoretically push the δ¹³C-DIC towards the less negative values observed in the current study (from -13.8 to -12.2‰) (Campeau et al. 2017b). Combined studies on aquatic metabolism, C dynamics and stable isotopic composition would further be recommended to disentangle the dynamic CO₂ source patterns in this type of agricultural system.

The spring and early summer of 2018 were generally dry leading to the stream channel drying out during long periods. The rapid rewetting periods (< 2 days) that occurred following larger precipitation events resulted in high CO₂ pulses (3-11 mg C L⁻¹) generally exceeding the overall median level of stream CO₂ (3.44 mg C L⁻¹) observed during the study period. The intermittent nature of streams, with distinct drying and rewetting episodes, is known to generate high CO₂ concentration pulses and subsequent emissions (Marcé et al. 2019). Such rapid pulses are generally suggested to be a result of intense respiration in the stream bed sediments upon rewetting, or due to a rapid mobilization of terrestrial C, both organic (DOC) and inorganic (CO₂) in connection to precipitation events. However, the findings of high CO₂ pulses upon rewetting have mostly been done in areas that display pronounced dry and wet seasons e.g. Mediterranean areas or Australia (e.g. Gomez-Gener et al. 2015; Looman et al. 2017). Here we show that such stream intermittency can also cause high and rapid CO₂ pulses in a Swedish agricultural setting, highlighting the need for expanding the geographical coverage of studies that investigate stream intermittency in relation to GHG dynamics and emissions. Areas that display stream intermittency will likely also increase in the future given the predicted changes in temperature and precipitation patterns. An obvious tool in this work is the use of continuous sensor-based measurements which allow capturing the episodic and unpredictable nature of these phenomena.

5. Conclusions

It is evident from the current study that the stream CO₂ dynamics in an agricultural headwater catchment are highly variable across a variety of different time-scales and with an interplay of hydrological and biological controls. The hydrological control was strong (although with both positive as well as negative influences dependent on season) and rapid in response to rainfall and snowmelt events. However, during growing-season baseflow and receding flow conditions, the aquatic primary production seems to control the stream CO₂ dynamics, which in turn sets the basis for atmospheric emissions. During the dry summer period, rapid rewetting following precipitation events generated high CO₂ pulses exceeding the overall median level of stream CO₂ (up to 3 times higher). This finding thus highlights the importance of stream intermittency in agricultural areas and its effect on stream CO₂ dynamics. Given the observed high levels of CO₂ and its temporally variable nature, agricultural streams clearly need more attention in order to understand and incorporate these considerable dynamics in large scale extrapolations.

6. Data availability

Data is available from the Uppsala University data repository, LINK WILL BE ADDED

7. Author contribution

MBW and MW brought the idea and designed the study. MBW funded and instrumented the catchment and analysed the data.
350 MW conducted the GIS analysis. JA, MP and ES provided ideas and data. MBW wrote the manuscript with great support from all co-authors.

8. Competing interests

The authors declare that they have no conflict of interest

9. Acknowledgements

355 Financial support to MBW from the King Carl-Gustaf XVI award for environmental science and from the Finn Malmgren foundation is acknowledged. JA was supported by FORMAS (grant 2015-1559). Jacob Smeds, My Osterman, Philip Johansson and Maud Oger are acknowledged for invaluable support in field and lab.

References

- 360 Alberts, J.M., Beaulieu, J.J. and Buffam, I., 2017, Watershed land use and seasonal variation constrain the influence of riparian canopy cover on stream ecosystem metabolism. *Ecosystems* 20(3), 553-567.
- Audet, J., Bastviken, D., Bundschuh, M., Buffam, I., Feckler, A., Klemedtsson, L., Laudon, H., Löfgren, S., Natchimuthu, S., Öquist, M., Peacock, M. and Wallin, M.B. 2019, Forest streams are important sources for nitrous oxide emissions. *Global Change Biology*, 26(2) doi:10.1111/gcb.14812
- 365 Berggren M., Laudon H., Jansson M., 2009, Hydrological control of organic carbon support for bacterial growth in boreal headwater streams. *Microbial Ecology*, 57, 170-178. doi:10.1007/s00248-008-9423-6
- Bodmer, P., Heinz, M., Pusch, M., Singer, G. and Premke, K., 2016, Carbon dynamics and their link to dissolved organic matter quality across contrasting stream ecosystems. *Science of The Total Environment*, 553, 574–586.
- Borges, A.V., Darchambeau, F., Lambert, T., Bouillon, S., Morana, C., Brouyère, S., Hakoun, V., Jurado, A., Tseng, H.-C., Descy, J.-P. and Roland, F.A.E., 2018, Effects of agricultural land use on fluvial carbon dioxide, methane and nitrous oxide concentrations in a large European river, the Meuse (Belgium). *Science of The Total Environment*, 610–611, 342–355.
- 370 Campeau, A., Bishop, K., Nilsson, M. B., Klemedtsson, L., Laudon, H., Leith, F. I., Öquist, M. G., Wallin, M. B., 2018, Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater catchments, *Journal of Geophysical Research – Biogeosciences*, 123 (1), 149-167, doi:10.1002/2017JG004083
- 375 Campeau, A., Bishop K., Billett, M. F., Garnett, M. H., Laudon, H., Leach, J. A., Nilsson, M. B., Öquist, M. G., Wallin, M. B., 2017a, Aquatic export of young dissolved and gaseous carbon from a pristine boreal fen: implications for peat carbon stock stability, *Global Change Biology*, 23 (12), 5523-5536, doi:10.1111/gcb.13815
- Campeau, A., Wallin, M. B., Giesler, R., Löfgren, S., Mörth, C-M., Schiff, S. L., Venkiteswaran, J. J., Bishop, K., 2017b, Multiple sources and sinks of dissolved inorganic carbon across Swedish streams, refocusing the lens of stable C isotopes.
380 *Scientific Reports*, 7, 9158, doi:10.1038/s41598-017-09049-9

- Castellano, M.J., Archontoulis, S.V., Helmers, M.J., Poffenbarger, H.J. and Six, J., 2019, Sustainable intensification of agricultural drainage. *Nature Sustainability*, 2(10), 914-921.
- Crawford, J. T., Stanley, E. H., Dornblaser, M. M., & Striegl, R. G., 2017, CO₂ time series patterns in contrasting headwater streams of North America. *Aquatic Sciences*, 79(3), 473-486. doi:10.1007/s00027-016-0511-2
- 385 Deirmendjian, L., Anschutz, P., Morel, C., Mollier, A., Augusto, L., Loustau, D., Cotovicz, L.C., Buquet, D., Lajaunie, K., Chaillou, G., Voltz, B., Charbonnier, C., Poirier, D. and Abril, G., 2019, Importance of the vegetation-groundwater-stream continuum to understand transformation of biogenic carbon in aquatic systems – A case study based on a pine-maize comparison in a lowland sandy watershed (Landes de Gascogne, SW France). *Science of the Total Environment* 661, 613-629. doi:10.1016/j.scitotenv.2019.01.152
- 390 Dinsmore, K. J., M. B. Wallin, M. S. Johnson, M. F. Billett, K. Bishop, J. Pumpanen & A. Ojala, 2013. Contrasting CO₂ concentration discharge dynamics in headwater streams: A multi-catchment comparison. *Journal of Geophysical Research: Biogeosciences*, 118, 445-461.
- Dinsmore, K. J. & M. F. Billett, 2008, Continuous measurement and modeling of CO₂ losses from a peatland stream during stormflow events. *Water Resources Research*, 44, 11.
- 395 Evans, C., Davies, T.D., 1998, Causes of Concentration/Discharge Hysteresis and its Potential as a Tool for Analysis of Episode Hydrochemistry. *Water Resources Research* 34, 129–137.
- Gómez-Gener, L., Obrador, B., von Schiller, D., Marcé, R., Casas-Ruiz, J.P., Proia, L., Acuña, V., Catalán, N., Muñoz, I., Koschorreck, M., 2015. Hot spots for carbon emissions from Mediterranean fluvial networks during summer drought. *Biogeochemistry* 125, 409–426.
- 400 Hall Jr, R.O. and Ulseth, A.J., 2019, Gas exchange in streams and rivers. *Wiley Interdisciplinary Reviews: Water*, p.e1391
- Halldin, S., Bergström, H., Gustafsson, D., Dahlgren, L., Hjelm, P., Lundin, L.C., Mellander, P.E., Nord, T., Jansson, P.E., Seibert, J., Stähli, M., Szilágyi Kishné, A. and Smedman, A.S. (1999) Continuous long-term measurements of soil–plant–atmosphere variables at an agricultural site. *Agricultural and Forest Meteorology*, 98-99, 75-102.
- Holmqvist, M., 1998, Avrinningsdynamik i fem små områden. *Vattenbalans, recession, magasin-koefficient och dynamiskt vattenmagasin*. MSc thesis, Uppsala University, 54 pp
- 405 Hope, D., S. M. Palmer, M. F. Billett, and J. J. Dawson, 2004, Variations in dissolved CO₂ and CH₄ in a first-order stream and catchment: an investigation of soil–stream linkages, *Hydrol. processes*, 18, 3255-3275.
- Hughes, R.M., Herlihy, A.T. and Kaufmann, P.R., 2010, An Evaluation of Qualitative Indexes of Physical Habitat Applied to Agricultural Streams in Ten US States 1. *JAWRA Journal of the American Water Resources Association*, 46(4), 792-806.
- 410 Johnson, M. S., M. F. Billett, K. J. Dinsmore, M. Wallin, K. E. Dyson, and R. S. Jassal, 2010, Direct and continuous measurement of dissolved carbon dioxide in freshwater aquatic systems - methods and applications, *Ecology*, 3, 68-78, doi:10.1002/eco.95.
- Johnson, M. S., Weiler, M., Couto, E. G., Riha, S. J., & Lehmann, J., 2007, Storm pulses of dissolved CO₂ in a forested headwater Amazonian stream explored using hydrograph separation. *Water Resources Research*, 43(11), w11201..
- 415 Kokic J, Wallin MB, Chmiel HE, Denfeld BA, Sobek S., 2015, Carbon dioxide evasion from headwater systems strongly contributes to the total export of carbon from a small boreal lake catchment. *J Geophys Res-Biogeo.* 120:13–28. doi:10.1002/2014JG002706
- Kokic J, Sahlée E, Sobek S, Vachon D, Wallin MB, 2018, High spatial variability of gas transfer velocity in streams revealed by turbulence measurements, *Inland Waters*, 8:4, 461-473, doi:10.1080/20442041.2018.1500228
- 420 Kyllmar, K., Forsberg, L. S., Andersson, S., & Martensson, K., 2014, Small agricultural monitoring catchments in Sweden representing environmental impact. *Agriculture Ecosystems & Environment*, 198, 25– 35. doi:10.1016/j.agee.2014.05.016
- Laudon, H., V. Sjöblom, I. Buffam, J. Seibert, and M. Mörth, 2007, The role of catchment scale and landscape characteristics for runoff generation of boreal streams, *J. Hydrol.*, 344(3-4), 198-209,
- 425 Leith, F. I., Dinsmore, K. J., Wallin, M. B., Billett, M. F., Heal, K. V., Laudon, H., Öquist, M. G., Bishop, K., 2015, Carbon dioxide transport across the hillslope–riparian–stream continuum in a boreal headwater catchment, *Biogeosciences*, 12, 1881-1892, doi:10.5194/bg-12-1-2015
- Linefur, H., Norberg, L., Kyllmar, K., Andersson, S. och Blomberg, M., 2018, Växtnäringsförluster i små jordbruksdominerade avrinningsområden 2016/2017. Uppsala: Sveriges lantbruksuniversitet. (Ekohydrologi, 155).
- Looman, A., Maher, D.T., Pendall, E., Bass, A., Santos, I.R., 2017. The carbon dioxide evasion cycle of an intermittent first-order stream: contrasting water–air and soil–air exchange. *Biogeochemistry*, 132, 87–102.
- 430

- Marcé, R., Obrador, B., Gómez-Gener, L., Catalán, N., Koschorreck, M., Arce, M.I., Singer, G. and von Schiller, D., 2019, Emissions from dry inland waters are a blind spot in the global carbon cycle. *Earth-Science Reviews* 188, 240-248.
- Morel, B., Durand, P., Jaffrezic, A., Gruau, G., & Molenat, J., 2009, Sources of dissolved organic carbon during stormflow in a headwater agricultural catchment. *Hydrological Processes*, 23(20), 2888-2901.
- 435 Natchimuthu, S., Wallin, M. B., Klemetsson, L., Bastviken, D., 2017, Spatio-temporal patterns of stream methane and carbon dioxide emissions in a hemiboreal catchment in Southwest Sweden, *Scientific Reports*, 7, 39729, doi:10.1038/srep39729
- Osborne, B., Saunders, M., Walmsley, D., Jones, M., Smith, P., 2010, Key questions and uncertainties associated with the assessment of the cropland greenhouse gas balance. *Agriculture, Ecosystems & Environment* 139, 293–301, doi:10.1016/j.agee.2010.05.009
- 440 Osterman, M., 2018, Carbon dioxide in agricultural streams - magnitude and patterns of an understudied atmospheric carbon source, MSc thesis, Uppsala University, 58 pp
- Öquist, M. G., M. Wallin, J. Seibert, K. Bishop, and H. Laudon, 2009, Dissolved inorganic carbon export across the soil/stream interface and its fate in a boreal headwater stream, *Environmental Science & Technology*, 43(19), 7364-7369, doi:10.1021/es900416h.
- 445 Peter, H., Singer, G. A., Preiler, C., Chiffard, P., Steniczka, G., & Battin, T. J., 2014, Scales and drivers of temporal pCO₂ dynamics in an Alpine stream. *Journal of Geophysical Research: Biogeosciences*, 119(6), 1078-1091. doi:10.1002/2013JG002552
- Ramankutty, N., Evan, A.T., Monfreda, C. and Foley, J.A., 2008, Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles*, 22(1).
- 450 Raymond PA, et al. 2013, Global carbon dioxide emissions from inland waters. *Nature*, 503 (7476), 355-359.
- Raymond, P. A., C. J. Zappa, D. Butman, T. L. Bott, J. Potter, P. Mulholland, A. E. Laursen, W. H. McDowell & D. Newbold 2012, Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers. *Limnology and Oceanography - Fluids and Environments*, 2, 41-53.
- Rhoads, B.L., Schwartz, J.S. and Porter, S., 2003, Stream geomorphology, bank vegetation, and three-dimensional habitat hydraulics for fish in midwestern agricultural streams. *Water Resources Research*, 39(8).
- 455 Riml, J., Campeau, A., Bishop, K., Wallin, M. B., 2019, Spectral decomposition of high-frequency CO₂ concentrations reveals soil-stream linkages, *Journal of Geophysical Research – Biogeosciences*, doi:10.1029/2018JG004981
- Rocher-Ros, G., Sponseller, R.A., Bergström, A.-K., Myrstener, M. and Giesler, R., 2019, Stream metabolism controls diel patterns and evasion of CO₂ in Arctic streams. *Global Change Biology*, doi:10.1111/gcb.14895
- 460 Royer, T. V., & David, M. B., 2005, Export of dissolved organic carbon from agricultural streams in Illinois, USA. *Aquatic Sciences*, 67(4), 465-471.
- Sand-Jensen, K., Staehr, P.A., 2012, CO₂ dynamics along Danish lowland streams: water–air gradients, piston velocities and evasion rates. *Biogeochemistry* 111, 615–628. doi:10.1007/s10533-011-9696-6
- Seibert, J., Grabs, T., Köhler, S., Laudon, H., Winterdahl, M., Bishop, K., 2009, Linking soil- and stream-water chemistry based on a Riparian Flow-Concentration Integration Model. *Hydrology and Earth System Sciences* 13, 2287–2297. <https://doi.org/10.5194/hess-13-2287-2009>
- 465 Wallin, M. B., Campeau, A., Audet, J., Bastviken, D., Bishop, K., Kokic, J., Laudon, H., Lundin, E., Löfgren, S., Natchimuthu, S., Sobek, S., Teutschbein, C., Weyhenmeyer, G., Grabs, T., 2018, Carbon dioxide and methane emissions of Swedish low-order streams – a national estimate and lessons learnt from more than a decade of observations, *Limnology and Oceanography – Letters*, 3 (3), 156-167, doi:10.1002/lol2.10061
- 470 Wallin, M. B., Grabs, T., Buffam, I., Laudon, H., Ågren, A., Öquist, M. G., Bishop, K., 2013, Evasion of CO₂ from streams – The dominant component of the carbon export through the aquatic conduit in a boreal catchment, *Global Change Biology*, 19(3), 785-797, doi:10.1111/gcb.12083.
- 475 Wallin, M. B., Öquist, M. G., Buffam, I., Billett, M. F., Nisell, J., Bishop, K. H., 2011, Spatiotemporal variability in the gas transfer coefficient (KCO₂) of boreal streams; implications for large scale estimates of CO₂ evasion, *Global Biogeochemical Cycles*, 25, GB3025, doi:10:1029/2010GB003975

Table 1. Catchment characteristics of the Sundbromark (SBM) catchment

Catchment area (km ²)	11.3
Elevation range (masl)	13-41
<i>Land use distribution (%)</i>	
Agricultural land	86
Forest	8
Urban	6
<i>Main Soil type distribution (%)</i>	
Post glacial clay	48
Glacial silt	22
Glacial clay	14
Sandy till	12
<i>Main bedrock distribution (%)</i>	
Granodiorite granite	89
Tonalite granodiorite	6
Dacite rhyolite	3
Granite	2

480

Table 2. Water chemistry at the outlet of the SBM catchment collected during June-November 2017 (n = 8) (Osterman 2018).

	Median	Mean	Min-Max
pH	7.7	7.8	7.4-8.4
EC ($\mu\text{S cm}^{-1}$)	1082	1273	791-1908
NH ₄ -N (mg L^{-1})	0.10	0.08	0.01-0.1
NO ₃ -N (mg L^{-1})	0.7	1.9	0.09-6.5
PO ₄ -P (mg L^{-1})	0.07	0.09	0.01-0.2
DOC (mg L^{-1})	10.0	9.6	4.2-13.1
D.O. (%)	53	62	31-119




Figure 1. Location of the study with A) sampled sites of the spatial survey, and B) The Sundbromark (SBM) catchment. Catchment delineation and land use distributions are given according to GSD elevation data, grid 2+ (©Swedish Land Survey) and CORINE Land Cover 2018 (European Environment Agency).






Figure 2. (A) Daily mean air temperature, and (B) daily precipitation during the study period (Sep 26, 2017-Dec 12, 2018) at the Marsta Observatory. Due to malfunctioning sensor the precipitation data for July 29 2018 is collected from the nearby (3 km) SMHI station, Årna. The dotted lines refer to the hydrological periods displayed in figure 3.

495



500 **Figure 3. Time series of A) stream discharge (Q) with sampling days for $\delta^{13}C$ -DIC highlighted by red dots, B) stream water temperature, C) electrical conductivity (EC), and D) CO_2 concentration for the study period September 26, 2017-December 12, 2018, with break for the ice- and snow-covered period December-March. The CO_2 data include periods when the sensor was above the water surface during dry periods in summer/autumn of 2018.**



505 **Figure 4. Stream CO₂ concentration (black) and discharge (red) for the autumn 2017 period with CO₂-Q hysteresis plots for four rain events.**




Figure 5. Stream CO₂ concentration (black) and discharge (red) for the snowmelt period 2018 with CO₂-Q hysteresis plots for four discharge events.




Figure 6. Time series of (A) Stream CO₂ concentration (black) and discharge (red), and (B) water temperature (black) and shortwave incoming radiation (SR, red) covering the period April-July 2018. Note the reverse axis for shortwave incoming radiation.




Figure 7. Diel amplitude in stream CO_2 concentration in relation to A) daily mean stream water temperature, and B) daily mean shortwave radiation (SR), covering the period April-July 2018. Statistics are given according to Spearman's rank correlation.

520





Figure 8. CO₂-Water temperature hysteresis loop based on the median daily values presented in figure 7 covering the period April-July 2018.



525 **Figure 9.** Stream CO₂ concentration (black) and discharge (red) for the dry period (July-September 2018). Periods when the CO₂ sensor was above the water table capturing an atmospheric signal (i.e. with concentrations <0.5 mg C L⁻¹) are highlighted by the lower box.




Figure 10. $\delta^{13}\text{C-DIC}$ as a function of stream discharge. The six sampling occasions covered the falling limb of the snowmelt peak April-June 2018.