

Is there warming in the pipeline? A multi-model analysis of the zero emission commitment from CO₂

Andrew H. MacDougall¹, Thomas L. Frölicher^{2,3}, Chris D. Jones⁴, Joeri Rogelj^{5,6}, H. Damon Matthews⁷, Kirsten Zickfeld⁸, Vivek K. Arora⁹, Noah J. Barrett¹, Victor Brovkin^{10,11}, Friedrich A. Burger^{2,3}, Micheal Eby¹², Alexey V. Eliseev^{13,14}, Tomohiro Hajima¹⁵, Philip B. Holden¹⁶, Aurich Jeltsch-Thömmes^{2,3}, Charles Koven¹⁷, Nadine Mengis¹⁸, Laurie Menviel¹⁹, Martine Michou²⁰, Igor I. Mokhov^{13,14}, Akira Oka²¹, Jörg Schwinger²², Roland Séférian²⁰, Gary Shaffer^{23,24}, Andrei Sokolov²⁵, Kaoru Tachiiri¹⁵, Jerry Tjiputra²², Andrew Wiltshire⁴, and Tilo Ziehn²⁶

¹St. Francis Xavier University, Antigonish, B2G 2W5, Canada

²Climate and Environmental Physics, Physics Institute, University of Bern, Switzerland

³Oeschger Centre for Climate Change Research, University of Bern, Switzerland

⁴Met Office Hadley Centre, Exeter, EX1 3PB, UK

⁵Grantham Institute for Climate Change and the Environment, Imperial College London, London, UK

⁶International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

⁷Concordia University, Montreal, Canada

⁸Department of Geography, Simon Fraser University, Burnaby, Canada

⁹Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC, Canada

¹⁰Max Planck Institute for Meteorology, Hamburg, Germany

¹¹CEN, University of Hamburg, Germany

¹²University of Victoria, Victoria, BC, Canada

¹³Faculty of Physics, Lomonosov Moscow State University, Moscow, Russia

¹⁴A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, Moscow, Russia

¹⁵Research Center for Environmental Modeling and Application, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

¹⁶School of Environment, Earth and Ecosystem Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

¹⁷Climate and Ecosystem Sciences Division, Lawrence Berkeley National Lab, Berkeley, CA, USA

¹⁸Biogeochemical Modelling Department, GEOMAR – Helmholtz Centre for Ocean Research, Kiel, Germany

¹⁹Climate Change Research Centre, PANGAEA, The University of New South Wales, Sydney, Australia

²⁰CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

²¹Atmosphere and Ocean Research Institute, The University of Tokyo

²²NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Bergen, Norway

²³Research Center GAIA Antarctica, University of Magallanes, Punta Arenas, Chile

²⁴Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

²⁵Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, USA

²⁶Commonwealth Scientific and Industrial Research Organisation, Oceans and Atmosphere, Aspendale, VIC, Australia

Correspondence: AH MacDougall (amacdoug@stfx.ca)

Abstract. The Zero Emissions Commitment (ZEC) is the change in global mean temperature expected to occur following the cessation of net CO₂ emissions, and as such is a critical parameter for calculating the remaining carbon budget. The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) was established to gain a better understanding of the potential magnitude and sign of ZEC, in addition to the processes that underlie this metric. Eighteen Earth system models

5 of both full and intermediate complexity participated in ZECMIP. All models conducted an experiment where atmospheric
CO₂ concentration increases exponentially until 1000 PgC has been emitted. Thereafter emissions are set to zero and models
are configured to allow free evolution of atmospheric CO₂ concentration. Many models conducted additional second priority
simulations with different cumulative emissions totals and an alternative idealized emissions pathway with a gradual transition
to zero emissions. The inter-model range of ZEC 50 years after emissions cease for the 1000 PgC experiment is -0.36 to
10 0.29 °C with a model ensemble mean of -0.07°C, median of -0.05°C and standard deviation of 0.19 °C. Models exhibit a
wide variety of behaviours after emissions cease, with some models continuing to warm for decades to millennia and others
cooling substantially. Analysis shows that both the carbon uptake by the ocean and the terrestrial biosphere are important for
counteracting the warming effect from the reduction in ocean heat uptake in the decades after emissions cease. This warming
effect is difficult to constrain due to high uncertainty in the efficacy of ocean heat uptake. Overall, the most likely value of ZEC
15 on multi-decadal timescales is close to zero, consistent with previous model experiments and simple theory.

1 Introduction

The long-term temperature goal of the Paris Agreement is to hold global warming well below 2°C, and to endeavour to
keep warming to no more than 1.5°C (United Nations, 2015). An important metric to assess the feasibility of this target is
the ‘remaining carbon budget’ (e.g. Rogelj et al., 2018), which represents the total quantity of CO₂ that can still be emitted
20 without causing a climate warming that exceeds the temperature limits of the Paris Agreement (e.g. Rogelj et al., 2019a). The
remaining carbon budget can be estimated from five factors: 1) historical human induced warming to date; 2) the Transient
Climate Response to cumulative CO₂ emissions (TCRE); 3) the estimated contribution of non-CO₂ climate forcings to future
warming; 4) a correction for the feedback processes presently unrepresented by Earth System Models (ESMs); and 5) the
unrealized warming from past CO₂ emissions, called Zero Emissions Commitment (ZEC) (e.g. Rogelj et al., 2019a). Of these
25 five factors, ZEC is the only quantity whose uncertainty was not formally assessed in the recent Intergovernmental Panel
on Climate Change (IPCC) Special Report on 1.5°C. Here we present the results of a multi-model analysis that uses the
output of dedicated model experiments that were submitted to the Zero Emission Commitment Model Intercomparison Project
(ZECMIP). This intercomparison project explicitly aims at quantifying the ZEC and identifying the processes that affect its
magnitude and sign across models (Jones et al., 2019).

30 ZEC is the change in global temperature that is projected to occur following a complete cessation of net CO₂ emissions
(Matthews and Weaver, 2010). After emissions of CO₂ cease, carbon is expected to be redistributed between the atmosphere,
ocean, and land carbon pools, such that the atmospheric CO₂ concentration continues to evolve over centuries to millennia (e.g.
Maier-Reimer and Hasselmann, 1987; Cao et al., 2009; Siegenthaler and Joos, 1992; Sarmiento et al., 1992; Enting et al., 1994;
Archer and Brovkin, 2008; Archer et al., 2009; Eby et al., 2009; Joos et al., 2013). In parallel, ocean heat uptake is expected
35 to decline as the ocean comes into thermal equilibrium with the elevated radiative forcing (Matthews and Caldeira, 2008). In
previous simulations of ZEC, the carbon cycle has acted to remove carbon from the atmosphere and counteract the warming
effect from the reduction in ocean heat uptake, leading to values of ZEC that are close to zero (e.g. Plattner et al., 2008;

Matthews and Caldeira, 2008; Solomon et al., 2009; Frölicher and Joos, 2010; Gillett et al., 2011). In the recent assessment of ZEC in the IPCC Special Report on Global Warming of 1.5 °C, the combined available evidence indicated that past CO₂ emissions do not commit to substantial further global warming (Allen et al., 2018). A ZEC of zero was therefore applied for the computation of the remaining carbon budget for the IPCC 1.5 °C Special Report (Rogelj et al., 2018). However, the evidence available at that time consisted of simulations from only a relatively small number of models using a variety of experimental designs. Furthermore, some recent simulations have shown a more complex evolution of temperature following cessation of emissions (e.g. Frölicher et al., 2014; Frölicher and Paynter, 2015). Thus a need to assess ZEC across a wider spectrum of climate models using a unified experimental protocol has been articulated (Jones et al., 2019).

ZEC was one of the metrics that emerged from the development of ESMs at the turn of the 21st century (Hare and Meinshausen, 2006). The concept was first conceptualized by Hare and Meinshausen (2006) who used the Model for the Assessment of Greenhouse gas Induced Climate Change (MAGICC), a climate model emulator, to explore temperature evolution following a complete cessation of all anthropogenic emissions. Matthews and Caldeira (2008) introduced the CO₂-only concept of ZEC which is used here. Their experiments used the intermediate complexity University of Victoria Earth System Climate Model (UVic ESCM) to show that stabilizing global temperature would require near zero CO₂ emissions. Plattner et al. (2008) used a wide range of different Earth System Models of Intermediate Complexity (EMICs) following a similar experiment and found that ZEC is close to (or less than) zero. These initial results with intermediate complexity models were subsequently supported by emission-driven ESM simulations (Lowe et al., 2009; Frölicher and Joos, 2010; Gillett et al., 2011). Zickfeld et al. (2013) quantified the ZEC under different scenarios and for a range of EMICs, but the resulting range is biased towards negative values as slightly negative instead of zero emissions were prescribed in some models. Some recent ESM simulations indicate that climate warming may continue after CO₂ emissions cease. For example, Frölicher and Paynter (2015) performed a simulation with the full ESM GFDL-ESM2M where emissions cease after 2°C of warming is reached. The simulations show some decades of cooling followed by a multi-centennial period of renewed warming resulting in an additional 0.5°C of warming 1000 years after emissions cease.

Two studies have examined in detail the underlying physical and biogeochemical factors that generate ZEC. Ehlert and Zickfeld (2017) examine ZEC with a set of idealized experiments conducted with the UVic ESCM. The study partitioned ZEC into a thermal equilibrium component represented by the ratio of global mean surface air temperature anomaly to unrealized warming, and a biogeochemical equilibrium component represented by the ratio of airborne fraction of carbon to equilibrium airborne fraction of carbon. The study found that the thermal equilibrium component of ZEC is much greater than the biogeochemical equilibrium component, implying a positive warming commitment. Williams et al. (2017) examine ZEC using the theoretical framework developed by Goodwin et al. (2007). The framework allows for the calculation of equilibrium atmospheric CO₂ concentration if the cumulative effect of the land carbon sink is known. The framework was applied to the same simulation conducted for Frölicher and Paynter (2015). The analysis showed that ZEC emerges from two competing contributions: 1) a decline in the fraction of heat taken up by the ocean interior leading to radiative forcing driving more surface warming; 2) uptake of carbon by the terrestrial biosphere and ocean system removing carbon from the atmosphere, causing a

cooling effect. Both studies focused on the long-term value of ZEC after multiple centuries and thus neither study examined what drives ZEC in the policy relevant timeframe of a few decades following cessation of emissions.

While we focus here on the ZEC from CO₂ emissions only, the ZEC concept has also been applied to the climate commitment resulting from other greenhouse gas emissions and aerosols (Frölicher and Joos, 2010; Matthews and Zickfeld, 2012; Mauritsen and Pincus, 2017; Allen et al., 2018; Smith et al., 2019), wherein the ZEC is characterized by an initial warming due to the removal of aerosol forcing, followed by a more gradual cooling from the decline in non-CO₂ greenhouse gas forcing. The ZEC from all emissions over multiple centuries is generally consistent with ZEC from only CO₂ emissions, for moderate future scenarios (Matthews and Zickfeld, 2012).

In addition to the ZEC, other definitions of warming commitment have also been used in the literature. The ‘constant composition commitment’ is defined as the unrealized warming that results from constant atmospheric greenhouse gas and aerosol concentrations (Wigley, 2005; Meehl et al., 2005; Hare and Meinshausen, 2006). This variety of warming commitment was highlighted prominently in the 2007 IPCC report (Meehl et al., 2007), leading to a widespread misunderstanding that this additional ‘warming in the pipeline’ was the result of past greenhouse gas emissions. However, the constant composition commitment rather results primarily from the future CO₂ and other emissions that are required to maintain stable atmospheric concentrations over time (Matthews and Weaver, 2010; Matthews and Solomon, 2013). Another related concept is the future ‘emissions commitment’ which quantifies the committed future CO₂ (and other) emissions that will occur as a result of the continued operation of existing fossil fuel infrastructure (Davis et al., 2010; Davis and Socolow, 2014; Smith et al., 2019; Tong et al., 2019). This concept is also distinct from the ZEC, as it quantifies an aspect of socioeconomic inertia (rather than climate inertia), which has been argued to be an important driver of potentially unavoidable future climate warming (Matthews and Solomon, 2013; Matthews, 2014).

When considering climate targets in the range of 1.5 to 2.0°C and accounting for the approximately 1°C of historical warming to date (Allen et al., 2018; Rogelj et al., 2018), a ZEC on the order of ± 0.1°C can make a large difference in the remaining carbon budget. Hence there is a need for a precise quantification and in-depth understanding of this value. This can be achieved by a systematic assessment of ZEC across the range of available models and a dedicated analysis of the factors that control the value of ZEC in these simulations. Thus the goals of this study based on the simulations of the Zero Emissions Commitment Model Intercomparison Project (ZECMIP) are: 1) to estimate the value of ZEC in the decades following cessation of emissions in order to facilitate an estimate of the remaining carbon budget; 2) to test if ZEC is sensitive to the pathway of emissions; 3) to establish whether ZEC is dependent on the cumulative total CO₂ that are emitted before emissions cease; 4) to identify which physical and biogeochemical factors control the sign and magnitude of ZEC in models.

The most policy-relevant question related to ZEC is: will global temperature continue to increase following complete cessation of greenhouse gas and aerosol emissions? The present iteration of ZECMIP aims to answer part of this question by examining the temperature response in idealized CO₂-only climate model experiments. To answer the question in full, the behaviour of non-CO₂ greenhouse gases, aerosols, and land-use-change must be accounted for in a consistent way. Such efforts will be the focus of future iterations of ZECMIP.

2 Methods

2.1 Protocol and Simulations

Here we summarize the ZECMIP protocol, the full protocol for ZECMIP is described in Jones et al. (2019). The ZECMIP protocol requested modelling groups to conduct three idealized simulations of two different types each - A and B. Type A simulations are initialized from one of the standard climate model benchmark experiments in which specified atmospheric CO₂ concentration increases at a rate of 1% per year from its pre-industrial value of around 285 ppm until quadrupling, referred to as the 1pctCO₂ simulation in the Climate Model Intercomparison Project (CMIP) framework (Eyring et al., 2016). The three type A simulations are initialized from the 1pctCO₂ simulation when diagnosed cumulative emissions of CO₂ reach 750, 1000, and 2000 Pg C. After the desired cumulative emission is reached the models are set to freely evolving atmospheric CO₂ mode, with zero further CO₂ emissions. Since net anthropogenic emissions are specified to be zero in type A simulations, atmospheric CO₂ concentration is expected to decline in these simulations in response to carbon uptake by the ocean and land. A consequence of the protocol is that for the type A simulations each model branches from the 1pctCO₂ simulations in a different year, contingent on when a model reaches the target cumulative emissions, which in turn depends on each model's representation of the carbon cycle and feedbacks. An example of emissions for the type A experiments is shown in Figure 1a. The three type B simulations are initialized from pre-industrial conditions and are emissions driven from the beginning of the simulation. Emissions follow bell-shaped pathways wherein all emissions occur within a 100 year window (Figure 1b). In all experiments land-use-change and non-CO₂ forcings are held at their pre-industrial levels.

Due to the late addition of ZECMIP to the CMIP Phase 6 (CMIP6) (Eyring et al., 2016), only the 1000 Pg C type A experiment (esm-1pct-brch-1000PgC) was designated as top priority ZECMIP simulation. The other simulations were designated as second priority simulations and were meant to be conducted if participating modelling groups had the resources and time. Both full ESMs and EMICs were invited to participate in ZECMIP. ESMs were requested to perform the top priority simulation for 100 years after CO₂ emissions cease, and more years and more experiments as resources allowed. EMICs were requested to conduct all experiments for at least 1000 years of simulations following cessation of emissions. Table 1 shows the experiments and experimental codes for ZECMIP.

Table 1. Experiments designed for ZECMIP

Name	Code	Cumulative Emissions (PgC)	Priority
A1	esm-1pct-brch-1000PgC	1000	1
A2	esm-1pct-brch-750PgC	750	2
A3	esm-1pct-brch-2000PgC	2000	2
B1	esm-bell-1000PgC	1000	2
B2	esm-bell-750PgC	750	2
B3	esm-bell-2000PgC	2000	2




Figure 1. (a) Example of diagnosed emission from the UVic ESCM for the type A experiments. Emissions are diagnosed from the 1pctCO₂ experiment which has prescribed atmospheric CO₂ concentrations. The target cumulative emissions total is reached part-way through the final year of emissions, thus that final year has a lower average emission rate than the previous year. (b) Time series of global CO₂ emissions for bell curve pathways B1 to B3. The numbers in the legend indicate the cumulative amount of CO₂ emissions for each simulation.

130 2.2 Model Descriptions

Eighteen models participated in ZECMIP: nine comprehensive ESMs and nine EMICs. The primary features of each model are summarized in Table A1 and A2 for ESMs, and Table A3 and A4 for EMICs. The ESMs in alphabetical order are: 1) CSIRO Australian Community Climate and Earth System Simulator, ESM version 1.5 – ACCESS-ESM1.5, 2) Canadian Centre for Climate Modelling and Analysis (CCCma) – CanESM5, 3) Community Earth System Model 2 – CESM2, 4) Centre National de Recherches Météorologiques (CNRM) – CNRM-ESM2-1, 5) Geophysical Fluid Dynamics Laboratory (GFDL) – GFDL-ESM2M, 6) Japan Agency for Marine-Earth Science and Technology (JAMSTEC/team MIROC) – MIROC-ES2L, 7) Max Planck Institute Earth System model, version 1.2, low resolution – MPI-ESM1.2-LR, 8) Norwegian Earth System Model 2 – NorESM2 and 9) UK (Met Office Hadley Centre and NERC) – UKESM1-0-LL. The nine EMICs in alphabetical order are: 1) Bern three dimensional Earth System Model – Bern3D-LPX, 2) Climate-Biosphere model, version 2 - CLIMBER-2, 3) Danish Centre for Earth System Science Earth System Model – DCESS1.0, 4) A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences – IAPRAS, 5) Loch-Vecode-ECbilt-Clio Model – LOVECLIM 1.2, 6) Massachusetts Institute of Technology Earth System Model – MESM, 7) Model for Interdisciplinary Research on Climate-lite/Japan Uncertainty Modelling Project-Loosely Coupled Model – MIROC-lite, 8) Planet Simulator - Grid-Enabled Integrated Earth system model – PLASIM-GENIE, and 9) University of Victoria Earth System Climate Model – UVic ESCM 2.10. For brevity models are referred to by their short names in the remainder of the manuscript. Table 2 shows the ZECMIP experiments that each modelling group submitted.

Table 3 shows three benchmark climate metrics for each model, Equilibrium Climate Sensitivity (ECS), Transient Climate Response (TCR), and TCRE. ECS is the climate warming expected if atmospheric CO₂ concentration was doubled from the pre-industrial value and maintained indefinitely while the climate system is allowed to come into equilibrium with the elevated radiative forcing (e.g. Planton, 2013; Charney et al., 1979). There are a variety of methods to compute ECS from climate model outputs (e.g. Knutti et al., 2017). Here we use ECS values computed using the method of Gregory et al. (2004), called “effective climate sensitivity”. The method of Gregory et al. (2004) computes ECS from the slope of the scatter plot between change in global temperature and planetary heat uptake, with values from the benchmark experiment where atmospheric CO₂ concentration is instantaneously quadrupled (4×CO₂ experiment). TCR is the atmospheric surface temperature change (relative to the preindustrial temperature) when atmospheric CO₂ is doubled in year 70 of the 1pctCO₂ experiment, computed using a 20 year averaging window centred about year 70 of the experiment (e.g. Planton, 2013). TCRE is described in the introduction and is computed from year 70 of the 1% experiment (e.g. Planton, 2013).

Bern and UVic submitted three versions of their models with three different ECSs. For Bern ECSs of 2.0 °C, 3.0 °C, and 5.0 °C, and for UVic ECSs of 2.0 °C, 3.8 °C, and 5.0 °C. These ECS values are true equilibrium climate sensitivities computed by allowing each model to come fully into equilibrium with the changed radiative forcing. For each model the central ECS value was used for the main analysis, 3.0 °C for Bern and 3.8 °C for UVic. The remaining experiments were used to explore the relationship between ECS and ZEC.

Table 2. Experiments conducted for ZECMIP by model. Full ESMs are listed on top followed by EMICs.

Model	A1	A2	A3	B1	B2	B3
ACCESS	X	X	X	–	–	–
CanESM5	X	–	X	–	–	–
CESM2	X	–	–	–	–	–
CNRM	X	–	–	–	–	–
GFDL	X	X	X	X	X	X
MIROC-ES2L	X	X	X	–	–	–
MPI-ESM	X	–	–	–	–	–
NorESM2	X	–	–	–	–	X
UKESM	X	X	X	–	–	–
Bern	X	X	X	X	X	X
CLIMBER	X	–	–	–	–	–
DCESS	X	X	X	X	X	X
IAPRAS	X	X	X	X	X	X
LOVECLIM	X	X	–	X	–	–
MESM	X	X	X	X	X	X
MIROC-lite	X	X	X	X	X	X
P. GENIE	X	X	X	X	X	X
UVic	X	X	X	X	X	X

Table 3. Benchmark climate model characteristics for each model: Equilibrium Climate Sensitivity (ECS), Transient Climate Response (TCR), and Transient Climate Response to Cumulative CO₂ Emissions (TCRE). UKESM reported a maximum to minimum range for TCR and TCRE based on four ensemble members.

Model	ECS (°C)	TCR (°C)	TCRE (K EgC ⁻¹)
ACCESS	3.9	1.9	1.9
CanESM5	5.7	2.8	2.3
CESM2	5.1	2.0	2.0
CNRM	4.8	1.9	1.9
GFDL	2.4	1.4	1.2
MIROC-ES2L	2.7	1.5	1.3
MPI-ESM	2.8	1.8	1.6
NorESM2	2.6	1.5	1.5
UKESM	5.4	2.68 to 2.85	2.49 to 2.66
Bern	2.6	1.6	1.5
CLIMBER	2.8	1.8	1.7
DCESS	3.0	2.0	2.0
IAPRAS	2.2	1.5	1.5
LOVECLIM	2.8	1.5	1.4
MESM	2.9	1.8	1.7
MIROC-lite	1.7	1.2	1.1
P. GENIE	3.4	1.7	1.6
UVic	3.5	1.8	1.8

2.3 Quantifying ZEC

ZEC is the change in global average surface air temperature following the cessation of CO₂ emissions. Thus ZEC must be calculated relative to the global temperature when emissions cease. Typically such a value would be computed from a 20 year window centred on the year when emissions cease. However, for the ZECMIP type A experiments such a calculation underestimates the temperature of cessation, due to the abrupt change in forcing when emissions suddenly cease, leading to an overestimation of ZEC values. That is, a roughly linear increase in temperature pathway abruptly changes to a close to stable temperature pathway. Therefore we define the temperature of cessation to be the global mean surface air temperature from benchmark 1pctCO2 experiment averaged over a 20 year window centred on the year emissions cease in the respective ZECMIP type-A experiment (year the ZECMIP experiment branches from the 1% experiment). For the EMICs which lack internal variability this method provides an unbiased estimate of the temperature of cessation.

Earlier studies have examined ZEC from decadal (Matthews and Zickfeld, 2012; Mauritsen and Pincus, 2017; Williams et al., 2017; Allen et al., 2018; Smith et al., 2019) to multi-centennial timescales (Frölicher and Paynter, 2015; Ehlert and Zickfeld, 2017). One of the main motivations of this present study is to inform the impact of the ZEC on the remaining carbon budget. This remaining carbon budget is typically used to assess the consistency of societal emissions pathways with the international temperature target of the Paris Agreement (UNEP, 2018). In this emission pathway and policy context, the ZEC within a few decades of emissions cessation is more pertinent than the evolution of the Earth system hundreds or thousands of years into the future. Therefore, we define values of ZEC_X as a 20 year average temperature anomaly centred at year *X* after emissions cease. Thus, 50-year ZEC (ZEC₅₀) is the global mean temperature relative to the temperature of cessation averaged from year 40 to year 59 after emissions cease. We similarly define 25-year ZEC (ZEC₂₅) and 90-year ZEC (ZEC₉₀).

2.4 Analysis Framework

A key question of the present study is why some models have positive ZEC and some models have negative or close to zero ZEC. From elementary theory we understand that the sign of ZEC will depend on the pathway of atmospheric CO₂ concentration and ocean heat uptake following cessation of emissions (Wigley and Schlesinger, 1985). Complicating this dynamic is that atmospheric CO₂ change has contributions both from the net carbon flux from the ocean and the terrestrial biosphere. Using the forcing-response equation (Wigley and Schlesinger, 1985) and the common logarithmic approximation for the radiative forcing from CO₂ (Myhre et al., 1998), we can partition ZEC into contributions from ocean heat uptake, ocean carbon uptake, and net carbon flux into the terrestrial biosphere. The full derivation of the relationship is shown in Appendix B and the summary equations are shown below:

$$\lambda T_{ZEC} = -R \int_{t=ze}^{\infty} \frac{f_O}{C_A} dt - R \int_{t=ze}^{\infty} \frac{f_L}{C_A} dt - \epsilon(N - N_{ze}), \quad (1)$$

where λ (W m⁻²K⁻¹) is the climate feedback parameter, T_{ZEC} (K) is ZEC, R (W m⁻²) is the radiative forcing from an e-fold increase in atmospheric CO₂ burden, t is time (a), ze is the time that emissions cease, f_O (PgC a⁻¹) is ocean carbon uptake,

195 f_L (PgC a⁻¹) is carbon uptake by land, C_A is atmospheric CO₂ content (PgC), N is planetary heat uptake (W m⁻²), N_{ze} is planetary heat uptake at the time emissions cease, and ϵ is the efficacy of planetary heat uptake. The equation states that ZEC is proportional to the sum of three energy balance terms: 1) the change in radiative forcing from carbon taken up by the ocean; 2) the change in radiative forcing from carbon taken up or given off by land; and 3) the change in effective ocean heat uptake. The two integral terms can be evaluated numerically from the ZECMIP model output, and thus can be simplified into two energy forcing terms F_{ocean} and F_{land} :

$$200 \quad F_{ocean} = R \int_{t=ze}^{\infty} \frac{f_O}{C_A} dt, \quad (2)$$

and,

$$F_{land} = R \int_{t=ze}^{\infty} \frac{f_L}{C_A} dt, \quad (3)$$

and thus:

$$\lambda T_{ZEC} = -F_{ocean} - F_{land} - \epsilon(N - N_{ze}). \quad (4)$$

205 Values for R were computed from the effective radiative forcing value for the models that simulate internal variability, with effective radiative forcing provided by each modelling group. Bern, DCESS, and UVic prescribe exact values for R and thus these values were used for calculations with these models. Effective radiative forcing for a doubling of CO₂ is $\frac{1}{2}$ the y-intercept of a 4×CO₂ Gregory plot (Gregory et al., 2004). R values and the effective climate sensitivities were used to calculate λ for each model. Efficacy (Winton et al., 2010) was calculated from:

$$210 \quad \epsilon = \frac{\lambda T - R \ln\left(\frac{C_A}{C_{Ao}}\right)}{N}, \quad (5)$$

where C_{Ao} (PgC) is the pre-industrial CO₂ burden, and T (K) is the global mean temperature anomaly relative to pre-industrial temperature. T , N , and C_A values were taken from the benchmark 1pctCO₂ for each model, as an average value from year 10 to year 140 of that experiment. Computed ϵ values are shown in Table 4. Effective climate sensitivity is here calculated as a time average fit and hence is assumed to be a constant, while efficacy values are expected to change in time. In CLIMBER 215 planetary or ocean heat uptake is not included into standard output and hence is not analyzed using this framework. CESM2 and NorESM2 are also excluded as the 4×CO₂ experiment results for these models are not yet available.

Efficacy has been shown to arise from spatial patterns in ocean heat uptake (Winton et al., 2010, 2013; Rose et al., 2014), with ocean heat uptake in the high latitudes being more effective at cooling the atmosphere than ocean heat uptake in low latitudes (Rose et al., 2014). This spatial structure in the effectiveness of ocean heat uptake in turn is suspected to originate

220 from shortwave radiation cloud feedbacks (Andrews et al., 2015). The method we have used to calculate efficacy folds state-dependent feedbacks and temporal change in the climate feedback parameter (Rugenstein et al., 2016) into the efficacy parameter.

Notably, calculated effective climate sensitivities and effective radiative forcings vary slightly within models due to the internal variability (Gregory et al., 2015), hence the efficacy values calculated here are associated with some uncertainty. Efficacy values are known to evolve in time (Winton et al., 2010), thus the efficacy value from the 1pctCO₂ experiment may be different than efficacy 50 years after emission cease in the ZECMIP experiments. To test this effect yearly efficacy values were calculated for the four EMICs without internal variability (Bern, DCESS, MESM and UVic). These tests showed that efficacy was 3.5% to 25% away from the values for the 1pctCO₂ experiment 50 years after emission cease (Figure A1). Thus we have assigned efficacy a $\pm 30\%$ uncertainty. Notably efficacy declines in three of the four models, consistent with previous work showing strong trends in efficacy over time (Williams et al., 2017). Radiative forcing from CO₂ is not precisely logarithmic (Gregory et al., 2015; Byrne and Goldblatt, 2014; Etminan et al., 2016) and therefore the calculated F_{ocean} and F_{land} values will be slightly different than the changes in radiative forcing experienced within each model, except for the three models that prescribe CO₂ radiative forcing. Also accounting for the uncertainty in recovering R values from model output, we assign a $\pm 10\%$ uncertainty to radiative forcing values.

Table 4. Efficacy ϵ and radiative forcing for $2\times\text{CO}_2$ R values for each model. Efficacy values are calculated from the 1pctCO₂ experiment.

Model	Efficacy	Radiative forcing $2\times\text{CO}_2$ (W m^{-2})
ACCESS	1.2	2.9
CanESM5	1.0	3.4
CNRM	0.9	3.2
GFDL	1.3	3.6
MIROC-ES2L	1.0	4.1
MPI-ESM	1.1	4.1
UKESM	1.0	4.0
Bern	1.0	3.7
DCESS	1.1	3.7
IAPRAS	1.1	3.7
LOVECLIM	1.0	3.7
MESM	0.8	4.1
MIROC-lite	1.0	3.0
P. GENIE	0.9	4.2
UVic	1.0	4.1

3.1 A1 Experiment results

Figure 2 shows the evolution of atmospheric CO₂ concentration and temperature for the 100 years after emissions cease for the A1 experiment (1% branched at 1000 PgC). In all simulations atmospheric CO₂ concentration declines after emissions cease, with a rapid decline in the first few decades followed by a slower decline thereafter. The rates of decline vary across the models. By 50 years after emissions cease in the A1 experiment the change in atmospheric CO₂ concentration ranged from -91 to -52 ppm, with a mean of -76 ppm and median of -80 ppm. Temperature evolution in the 100 years following cessation of emissions varies strongly by model, with some models showing declining temperature, others having ZEC close to zero, and others showing continued warming following cessation of emissions. Some models such as UKESM, CNRM, and UVic exhibit continued warming in the centuries following cessation of emissions. Other models such as IAPRAS and DCESS exhibit a temperature peak, then decline. Still other models show ZECs that hold close to zero (e.g. MPI-ESM), while some models show continuous decline in temperature following cessation of emissions (e.g. P. GENIE). Table 5 shows the ZEC₂₅, ZEC₅₀, and ZEC₉₀ values for the A1 experiment. The table shows values of ZEC₅₀ ranging from -0.36 to 0.29 °C with a model ensemble mean of -0.06°C, median of -0.05°C and a standard deviation of 0.19°C. Tables C1 and C2 show ZEC₂₅, ZEC₅₀, and ZEC₉₀ for the A2 and A3 experiment. Figure 3 shows the evolution of atmospheric CO₂ concentration, temperature anomalies (relative to the year emissions cease), and ocean heat uptake for 1000 years following cessation of emissions in the A1 experiment. All models show continued decline in atmospheric CO₂ concentration for centuries after emissions cease. One model (P. GENIE) shows a renewed growth in atmospheric CO₂ concentration beginning about 600 years after emissions cease, resulting from release of carbon from soils overwhelming the residual ocean carbon sink. Of the nine models that extended simulations beyond 150 years, seven show temperature on a long-term decline (Bern, MESM, DCESS, IAPRAS, LOVECLIM, P. GENIE, and MIROC-ES2L), GFDL shows temperature declining and then increasing within 200 years after cessation but ultimately remaining close to the temperature at cessation, and the UVic model shows slow, warming. Most models show continuous decline in ocean heat uptake with values approaching zero. Three models (GFDL, LOVECLIM and IAPRAS) show the ocean transition from a heat sink to a heat source.

Table 5. Temperature anomaly relative to the year emissions cease averaged over a 20 year time window centred about the 25th, 50th, and 90th year following cessation of anthropogenic CO₂ emissions (ZEC₂₅, ZEC₅₀, and ZEC₉₀ respectively) for the A1 (1% to 1000 PgC experiment).

Model	ZEC ₂₅ (°C)	ZEC ₅₀ (°C)	ZEC ₉₀ (°C)
ACCESS	0.04	0.01	-0.03
CanESM5	-0.04	-0.13	-0.17
CESM2	-0.11	-0.31	-0.34
CNRM	0.11	0.06	0.25
GFDL	-0.18	-0.27	-0.19
MIROC-ES2L	-0.02	-0.08	-0.21
MPI-ESM	-0.22	-0.27	-0.37
NorESM	-0.27	-0.33	-0.32
UKESM	0.21	0.28	0.33
Bern	0.05	0.01	-0.08
DCESS	0.11	0.06	-0.04
CLIMBER	0.0	-0.07	-0.14
IAPRAS	0.34	0.29	0.03
LOVECLIM	-0.02	-0.04	-0.03
MESM	0.05	0.01	-0.06
MIROC-lite	-0.02	-0.06	-0.09
P. GENIE	-0.19	-0.36	-0.71
UVic	0.04	0.03	0.02
Mean	-0.01	-0.07	-0.12
Median	-0.01	-0.05	-0.08
Standard Deviation	0.15	0.19	0.23




Figure 2. (a,c) Atmospheric CO₂ concentration anomaly and (b,d) Zero Emissions Commitment following cessation of emissions under the experiment where 1000 PgC was emitted following the 1% experiment (A1). ZEC is the temperature anomaly relative to the estimated temperature at the year of cessation. Top row shows output for ESMs and bottom row for EMICs.




Figure 3. (a) Change in atmospheric CO₂ concentration, (b) change in temperature, and (c) ocean heat uptake following cessation of emissions for the A1 experiment (1000 PgC following 1%) for 1000 years following cessation of emissions.

3.2 Effect of emissions rate, 1% vs. Bell

260 The bell experiments were designed to test whether temperature evolution following cessation of emissions depends on the pathway of emissions before emissions cease. These experiments also illustrate model behaviour during a gradual transition to zero emissions (e.g. MacDougall, 2019), a pathway that is consistent with most future scenarios (Eyring et al., 2016). Nine of the participating models conducted both the A1 and B1 experiments, GFDL and eight of the EMICs (CLIMBER is the EMIC which did not conduct the B1 experiment). Figure 4 shows the temperature evolution (relative to pre-industrial temperature) 265 for both experiments. All models show that by the 100th year of the experiments, when emissions cease in the bell experiment, the temperature evolution is very close in the two experiments. For seven of the models, GFDL, Bern, DCESS, LOVECLIM, MESM, MIROC-lite, and UVic, the temperature evolution in the A1 and B1 experiments is indistinguishable after emissions cease in the Bell experiment. That is, models suggest that in the long term the past pathway of CO₂ is largely irrelevant to total temperature change, and determined only by the total amount of cumulative emissions.

270 Figure 5 shows ZEC for both experiments. There is no sharp discontinuity in forcing in the Bell experiments and thus the temperature of cessation for these experiments is simply calculated relative to a temperature average from a 20 year window centred about the year 100 when emissions cease. Despite the long-term temperature evolution being the same for both experiments, the change in temperature relative to time of cessation is different in most models. This feature is not unexpected as theoretical work on the TCRE relationship suggest that direct proportionality between cumulative emissions of 275 CO₂ and temperature change should break down when emission rates are very low (MacDougall, 2017), as emissions are near the end of the Bell experiments. That is, in the type-B experiments emissions decline gradually and hence the Earth system is closer to thermal and carbon cycle equilibrium when emissions cease. These results support using the type-A experiment (1% followed by sudden transition to zero emissions) to calculate ZEC for providing a correction to the remaining carbon budget, as the experiment provides a clear separation between TCRE and ZEC, while for a gradual transition to zero emissions scenario 280 the two effects are mixed as emissions approach zero.

The B2 experiment (750 PgC) was designed to assess ZEC for an emissions total that would imply a climate warming of close to 1.5°C (Jones et al., 2019). The mean change in emission rate for the B2 experiment during the ramp-down phase of the experiment (year 50 to 100) is -0.39 PgC a⁻². This rate is similar to the rate of -0.29 [-0.05 to -0.64] PgC a⁻² for stringent mitigation scenarios from the IPCC Special Report on 1.5° for the period from year 2020 to 2050 CE (Rogelj et al., 2018). 285 Therefore we would expect similar behaviour in the stringent mitigation scenarios and the type-B experiments. That is, for the effect of ZEC to manifest while emissions are ramping down. The A2 experiment (1% 750 PgC) branches from the 1pctCO2 experiment between year 51 and 60 in the models that performed that experiment. Emission in the A2 experiment cease in year 100. Thus the temperature correction expected by time emission cease for the stringent mitigation scenarios would be in the range of ZEC₄₀ to ZEC₅₀ for the B2 experiment.




Figure 4. Temperature evolution of A1 (1% to 1000 PgC) and B1 (Bell shaped emissions of 1000 PgC over 100 years) experiments relative to pre-industrial temperature. Solid lines are the A1 experiment and dashed lines are the B1 experiment. Vertical blue line shows when emissions cease in the A1 experiment and vertical red line shows where emissions cease in the B1 experiment.




Figure 5. ZEC for the A1 (1% to 1000 PgC) and B1 (Bell shaped emissions of 1000 PgC over 100 years) experiments. Solid lines are the A1 experiment and dashed lines are the B1 experiment.

Twelve models conducted at least two type-A (1%) experiments such that ZEC could be calculated for 750 PgC, 1000 PgC, and 2000 PgC of cumulative emissions, five ESMs (ACCESS, CanESM5, GFDL, MIROC-ES2L, and UKESM), and all of the EMICs except CLIMBER. Two of the models conducted only two of the type-A experiments, CanESM5 conducted the A1 and A3 experiments, while LOVECLIM conducted the A1 and A2 experiments. Figure 6 shows the ZEC_{50} for each model for the three experiments. All of the full ESMs exhibit higher ZEC_{50} with higher cumulative emissions. The EMICs have a more mixed response with Bern, MESM, LOVECLIM and UVic showing increased ZEC_{50} with higher cumulative emissions, DCESS and IAPRAS showing slightly declining ZEC_{50} with higher cumulative emissions, and P. GENIE showing a strongly declining ZEC_{50} with higher emissions. The inter-model range for the ZEC_{50} of the A2 (750 PgC) experiment is -0.31 to 0.30 °C with a mean value of -0.03 °C, a median of -0.06 °C and standard deviation of 0.15 °C. The inter-model range A3 (2000 PgC) experiment -0.40 to 0.52 °C with a mean of 0.10 °C, a median of 0.10 °C and standard deviation of 0.26 °C. Note that different subsets of models conducted each experiment, such that ranges between experiments are not fully comparable.




Figure 6. Values of ZEC_{50} for the 750, 1000 and 2000 PgC experiments branching from the 1% experiment (type-A). Panel (a) shows results for full ESMs and panel (b) for EMICs

3.4 Analysis of Results

The framework introduced in Section 2.4 was applied to the ZECMIP output to partition the energy balance components of ZEC into contributions from the warming effect of the reduction in ocean heat uptake ($-\Delta N$), and the effect of the change in radiative forcing from the ocean (F_{ocean}) and terrestrial carbon fluxes (F_{land}). Figure 7 shows the results of this analysis for each model averaged over the period 40 to 59 years after emissions cease for the 1000 PgC 1% (A1) experiment (the same time interval as ZEC_{50}). The components of the bars in Figure 7a are the terms of the right hand side of Equation 4. The results suggest that both ocean carbon uptake and terrestrial carbon uptake are critical for determining the sign of ZEC in the decades following cessation of emissions. Previous efforts to examine ZEC, while acknowledging the terrestrial carbon sink, have emphasized the role of ocean heat and carbon uptake (Ehlert and Zickfeld, 2017; Williams et al., 2017). These studies also focused on ZEC on timescales of centuries, not decades. In CanESM5 and CNRM the terrestrial carbon sink dominates the reduction in radiative forcing, while in ACCESS, IAPRAS, MESM, P. GENIE, and UVic the ocean carbon uptake dominates the reduction in radiative forcing. The remaining models have substantial contributions from both carbon sinks. In all models the reduction in forcing from ocean carbon uptake is smaller than the reduction in ocean heat uptake, suggesting that the post-cessation net land carbon sink is critical to determining ZEC values. The ocean carbon uptake itself varies substantially between models, with some of the EMICs (P. GENIE, MESM, and IAPRAS) having very high ocean carbon uptake, and two of the ESMs (CanESM5 and CNRM) having very low ocean carbon uptake. Given that the behaviour of the terrestrial carbon cycle varies strongly between models (Friedlingstein et al., 2006; Arora et al., 2013, 2019) and that many models lack feedbacks related to nutrient limitation and permafrost carbon pools, the strong dependence of ZEC_{50} on terrestrial carbon uptake is concerning for the robustness of ZEC_{50} estimates. Notably, the three ESMs, with the weakest terrestrial carbon sink response (ACCESS, MIROC-ES2L, and UKESM), include terrestrial nutrient limitations (Table A1, A2). However, despite including terrestrial nutrient limitation Bern and MPI-ESM simulate a terrestrial carbon uptake in the middle and upper parts of the inter-model range, respectively. The UVic model includes permafrost carbon and has a relatively weak terrestrial carbon uptake (Table A4). IAPRAS does not account for either nutrient limitations or permafrost carbon and has the weakest terrestrial carbon uptake among all models studied here (Table A3).

To further investigate the effect of nutrient limitation on ZEC we have compared models with and without terrestrial nutrient limitations. Eight of the models that participated in ZECMIP included a representation of the terrestrial nitrogen cycle, ACCESS, CESM2, MIROC-ES2L, MPI-ESM, NorESM, UKESM, Bern and MESM. One model (ACCESS) includes a representation of the terrestrial phosphorous cycle. Figure 8 shows behaviour of the terrestrial carbon cycle before and after emissions cease for models with and without terrestrial nutrient limitations. Figure 8a shows that consistent with Arora et al. (2019) models with a terrestrial nitrogen cycle have on average a lower carbon uptake than those without. However, after emissions cease there is little difference in the terrestrial uptake of carbon between models with and without nutrient limitations. For both sets of model the median uptake is almost the same at 68 PgC and 63 PgC respectively, and the range for models without nutrient limitation fully envelops the range for those with nutrient limitations. Thus, while nutrient limitations do not appear to have a controlling influence on the *magnitude* of the post cessation terrestrial carbon uptake they have a marked impact on its

uncertainty. As with carbon cycle feedbacks (Arora et al., 2019) those models including terrestrial nitrogen limitation exhibit substantially smaller spread than those which do not. This offers hope for future reductions in ZEC uncertainty as more models begin to include nitrogen - and thereafter phosphorus - limitations on the land carbon sink.

340 Figure 9a shows the relationship between ocean heat uptake (Figure 9b), cumulative ocean carbon uptake (Figure 9c), and the cumulative terrestrial carbon uptake when emissions cease and 50 years after emissions cease. Excluding the clear outlier of IAPRAS Figure 9 shows a clear negative relationship ($R=-0.80$) between ocean heat uptake before emissions cease and the change in ocean heat uptake 50 years after emissions cease in the A1 experiment. That is, models with high ocean uptake before emissions cease tend to have a strong reduction in ocean heat uptake after emissions cease. Similarly there is a strong ($R=0.88$) positive relationship between ocean carbon uptake before emissions cease and uptake in the 50 years after emissions cease. The relationship between uptake (or in one case net release) of carbon by the terrestrial biosphere before and after emissions cease is weaker ($R=0.72$) but clear. Therefore explaining why the energy balance components illustrated by Figure 7 vary between 345 models would seem to relate strongly to why models have varying ocean heat, ocean carbon uptake, and terrestrial carbon cycle behaviour before emissions cease.

It has long been suggested that the reason that long-term ZEC was close to zero is compensation between ocean heat 350 and ocean carbon uptake (Matthews and Caldeira, 2008; Solomon et al., 2009; Frölicher and Paynter, 2015), which are both dominated by the ventilation of the thermocline (Sabine et al., 2004; Banks and Gregory, 2006; Xie and Vallis, 2012; Frölicher et al., 2015; Goodwin et al., 2015; Zanna et al., 2019). However, Figure 7 shows that this generalization is not true for decadal time-scales. The two quantities do compensate one-another but in general the effect from reduction in ocean heat uptake is larger than the change in radiative forcing from the continued ocean carbon uptake. Thus going forward additional emphasis 355 should be placed on examining the role of the terrestrial carbon sink in ZEC for policy relevant timescales. Also notable is the large uncertainty in effective ocean heat uptake, which originates from the uncertainty in efficacy. As efficacy is related to spatial patterns in ocean heat uptake and coupled shortwave cloud feedbacks (Rose et al., 2014; Andrews et al., 2015), shifts in these patterns in time thus likely affect the values of ZEC and hence represents an important avenue for further investigation.

Figure 10 compares the energy fluxes for the ten models that conducted all of the type-A (1%) experiments. All three energy 360 balance components seem to be affected by the cumulative emissions leading up to cessation of emissions, however there is no universal pattern. Most models show a larger reduction in ocean heat uptake with higher cumulative emissions, but UKESM has the largest reduction for the 1000 PgC experiment. Variations in the reduction in radiative forcing from ocean carbon uptake tend to be small between simulations within each model but show no consistent patterns between models. Most of the models show a smaller terrestrial carbon sink for the 2000 PgC experiment than the other two experiments, the exception being 365 IAPRAS which shows the opposite pattern. Examining in detail why these factors change in each model could be a productive avenue for future research.

3.5 Relationship to other climate metrics

Figure 11 shows the relationship between ECS, TCR, TCRE, Realized Warming and ZEC_{50} for the A1 (1000 PgC) experiment. Realized warming is the ratio of TCR to ECS. TCR is transient warming when CO_2 is doubled and ECS is warming at




Figure 7. (a) Energy fluxes following cessation of CO₂ emissions for the 1000 PgC 1% (A1) experiment. ΔN is the change in ocean heat uptake relative to the time emissions ceased. A reduction in ocean heat uptake will cause climate warming, hence $-\Delta N$ is displayed. F_{ocean} is the change in radiative forcing caused by ocean carbon uptake, and F_{land} is the change in radiative forcing caused by terrestrial carbon uptake. Vertical black lines are estimated uncertainty ranges. (b) ZEC₅₀ values for each model. Models are arranged in ascending order of ZEC₅₀

370 equilibrium following doubling of CO₂, their ratio is the fraction of warming from CO₂ that has been realized, hence ‘Realized Warming’ (e.g. Frölicher et al., 2014). ECS shows virtually no correlation with ZEC₅₀ (R=0.04) and thus ZEC and ECS appear to be independent. Both TCR and Realized Warming show weak positive correlations with ZEC₅₀ (R= 0.25 and 0.30 respectively). TCRE shows the strongest relationship to ZEC₅₀ with a correlation coefficient of 0.34. However, these relationships may not be robust due to small number of non-independent models. The poor correlation between ZEC and other climate

375 metrics is not unexpected as ZEC is determined by the difference in warming caused by reduction in ocean heat uptake and cooling caused by continued land and ocean carbon uptake after the cessation of emissions. Small differences between large quantities are not expected to correlated well to the quantities used to calculate them.

Bern and UVic both conducted the ZECMIP experiments with three versions of their models with different equilibrium climate sensitivities, allowing for examination of the effect of ECS on ZEC. Figure 12 shows the ZEC₅₀ for these simulations.

380 The figure shows that for both Bern and UVic higher ECS corresponds to higher ZEC. For Bern for the A1 (1% 1000 PgC) experiment ZEC₅₀ is 0.01, 0.03, and 0.18 $^{\circ}\text{C}$ for ECSs of 2.0, 3.0 and 5.0 $^{\circ}\text{C}$ respectively. For UVic for the A1 (1% 1000 PgC)




Figure 8. Terrestrial carbon uptake for models with and without a nitrogen cycle, before emissions cease and after emissions cease. Eight models have a representation of nutrient limitations and ten do not. Circles indicate data points for ESMs and triangle indicate data points for EMICs.




Figure 9. Relationship between variables before emissions cease and 50 years after emissions cease. (a) Ocean Heat Uptake (OHU) is computed for 20 year windows with the value at cessation taken from the 1pctCO₂ experiment analogous to the how temperature of cessation is computed. (b) Cumulative ocean carbon uptake, (c) Cumulative land carbon uptake. Each marker represents value from a single model. Line of best fit excludes the outlier model IAPRAS which is marked with a magenta square.




Figure 10. Energy fluxes following cessation of CO₂ emissions for the type-A experiments (1%) for each model. ΔN is the change in ocean heat uptake relative to the time emissions ceased. A reduction in ocean heat uptake will cause climate warming, hence $-\Delta N$ is displayed. F_{ocean} is the change in radiative forcing caused by ocean carbon uptake, and F_{land} is the change in radiative forcing caused by terrestrial carbon uptake. All fluxes are computed for averages from 40 to 59 years after emissions cease.

experiment ZEC₅₀ is -0.15, 0.01, and 0.22 for ECSs of 2.0, 3.8 and 5.0 °C respectively. Note that the ECS values given here are for true equilibrium climate sensitivity, not effective climate sensitivities as used in the remainder of this study. Figure 13 compares the energy fluxes for the three versions of Bern and UVic. For Bern ocean carbon uptake is unaffected by climate sensitivity, while for UVic there is a small decline in ocean carbon uptake for an ECS of 5.0°C. For Bern the reduction in ocean heat uptake is larger at higher climate sensitivity, while for UVic this quantity is almost the same for ECSs of 3.8 and 5.0 °C. In Bern the terrestrial carbon sink is weaker in versions with higher climate sensitivity. In UVic the terrestrial carbon sink is weakest for a climate sensitivity of 3.8 °C. Overall the results suggest a relationship between higher ECS and higher ZEC within these models.




Figure 11. Relationship between: ECS (a), TCR (b), TCRE (c), Realized Warming (d) and ZEC₅₀. Line of best fit is shown in red. Correlation coefficients are displayed in each panel.




Figure 12. Values of 50 year ZEC for the 750, 1000 and 2000 PgC experiments branching from the type-A (1%), for versions of Bern and UVic with varying Equilibrium Climate Sensitivity. Note that the ECS values given here are for true equilibrium climate sensitivity, not effective climate sensitivities as used in the remainder of the study.




Figure 13. Energy fluxes following cessation of CO₂ emissions for the type-A experiments (1%) for versions of Bern and UVic with varying Equilibrium Climate Sensitivity. ΔN is the change in ocean heat uptake relative to the time emissions ceased. A reduction in ocean heat uptake will cause climate warming, hence $-\Delta N$ is displayed. F_{ocean} is the change in radiative forcing caused by ocean carbon uptake, and F_{land} is the change in radiative forcing caused by terrestrial carbon uptake. All fluxes are computed for averaged from 40 to 59 years after emissions cease.

4.1 Drivers of ZEC

The analysis here has shown that across models decadal-scale ZEC is poorly correlated to other metrics of climate warming, such as TCR and ECS, though relationships may exist within model frameworks (Figure 12). However, the three factors that drive ZEC, ocean heat uptake, ocean carbon uptake and net land carbon flux correlate relatively well to their states before emissions cease. Thus it may be useful to conceptualize ZEC as a function of these three components each evolving in their own way in reaction to the cessation of emissions. Ocean heat uptake evolves due to changes in ocean dynamics (e.g. Frölicher et al., 2015) as well as the complex feedbacks that give rise to changes in ocean heat uptake efficacy (Winton et al., 2010). Ocean carbon uptake evolution is affected by ocean dynamics, changes to ocean biogeochemistry, and changes in atmosphere-ocean CO₂ chemical disequilibrium, where the latter is also influenced by land carbon fluxes (e.g. Sarmiento and Gruber, 2006). The response of the land biosphere to cessation of emissions is expected to be complex with contributions from the response of photosynthesis to declining atmospheric CO₂ concentration, a continuation of enhanced soil respiration (e.g. Jenkinson et al., 1991), and release of carbon from permafrost soils (Schuur et al., 2015), among other factors. Investigating the evolution of the three components in detail may be a valuable avenue of future analysis. Similarly, given their clearer relationships to the state of the Earth system before emissions cease, focusing on the three components independently may prove useful for building a framework to place emergent constraints on ZEC. Future work will explore evaluation opportunities by assessing relationships between these quantities in the idealized 1% simulation and values at the end of the historical simulations up to present day.

Our analysis has suggested that the efficacy of ocean heat uptake is crucial for determining the temperature effect from ocean heat uptake following cessation of emissions. Efficacy itself is generated by spatial patterns in ocean heat uptake and shortwave cloud feedback processes (Rose et al., 2014; Andrews et al., 2015). Thus, evaluating how these processes and feedbacks evolve after emissions cease is crucial for better understanding ZEC. As the spatially resolved outputs for ZECMIP are now available (see Section 5), evaluating such feedbacks presents a promising avenue for future research.

4.2 Policy Implications

One of the main motivations to explore ZEC are its implications for policy and society's ability to limit global warming to acceptable levels. Climate policy is currently aiming at limiting global mean temperature increase to well below 2 °C and pursuing to limit it to 1.5 °C (United Nations, 2015). To stay within these temperature limits, emission reduction targets are being put forward. These targets can take the form of emissions reductions in specific years, like the Paris Agreement nationally determined contributions for the years 2025 or 2030 (Rogelj et al., 2016), but also of net zero emissions targets that cap the cumulative CO₂ emissions a country is contributing to the atmosphere (Haites et al., 2013; Rogelj et al., 2015; Geden, 2016). Because ZEC does affect the required stringency of emissions reductions or of the maximum warming one can expect, it is important to clearly understand its implications within a wider policy context. First, for policy analysts and scientists, the quantification of ZEC₅₀ will help inform better estimates of the remaining carbon budget compatible with limiting warming to 1.5 °C or well below 2 °C over the course of this century. Analysts need to be clear, however, that ZEC₅₀ is only then an

adequate adjustment for TCRE-based carbon budget estimates if the TCRE values are based on 1% CO₂ increase simulations. In contrast, however, our results also show that when CO₂ emissions ramp down gradually (c.f. the B-series of ZECMIP experiments) ZEC₅₀ is generally much smaller, because part of it is already realized during the emissions ramp down. Hence this means that in a situation in which society successfully gradually reduces its global CO₂ emissions to net zero at rates comparable to the B2 experiment (see Section 3.2), the expected additional warming on time scales of decades to maximum a century is small. Finally, over multiple centuries, warming might still further increase or decrease. In the former case, a certain level of carbon-dioxide removal would be required over the coming centuries. The level implied by the long-term ZEC, however, represents much less a challenge than the urgent drastic emissions cuts required to limit warming to either 1.5 °C or 2 °C over the next decades (Rogelj et al., 2019b).

4.3 Moving towards ZECMIP-II

For the first iteration of ZECMIP the experimental protocol has focused solely on the response of the Earth system to zero emission of CO₂. However, many other non-CO₂ greenhouse gases, aerosol, and land-use-changes all affect global climate (e.g. IPCC, 2013). To truly explore the question whether global temperature will continue to increase following complete cessation of greenhouse gas and aerosol emissions the effect of each anthropogenic forcing agent must be accounted for (e.g. Frölicher and Joos, 2010; Matthews and Zickfeld, 2012; Mauritsen and Pincus, 2017; Allen et al., 2018; Smith et al., 2019). We envision a second iteration of ZECMIP accounting for these effects with a set of self-consistent idealized experiments, as part of the formal CMIP7 process.

440 5 Conclusions

Here we have analysed model output from the 18 models that participated in ZECMIP. We have found that the inter-model range of ZEC 50 years after emissions cease for the A1 (1% to 1000 PgC) experiment is -0.36 to 0.29 °C with a model ensemble mean of -0.07 °C, median of -0.05 °C, and standard deviation of 0.19 °C. Models show a range of temperature evolution after emissions cease from continued warming for centuries to substantial cooling. All models agree that following cessation of
445 CO₂ emissions, the atmospheric CO₂ concentration will decline. Comparison between experiments with a sudden cessation of emissions and a gradual reduction in emissions show that long term temperature change is independent of the pathway of emissions. However, in experiments with a gradual reduction in emissions a mixture of TCRE and ZEC effects occur as the rate of emissions declines. As the rate of emission reduction in these idealized experiments is similar to that in stringent mitigation scenarios, a similar pattern may emerge if deep emission cuts commence.

450 ESM simulations agree that higher cumulative emissions lead to a higher ZEC, though some EMICs show the opposite relationship. Analysis of the model output shows that both ocean carbon uptake and the terrestrial carbon uptake are critical for reducing atmospheric CO₂ concentration following the cessation of CO₂, and thus counteracting the warming effect of reduction in ocean heat uptake. The three factors that contribute to ZEC (ocean heat uptake, ocean carbon uptake and net land carbon flux) correlate well to their states prior to the cessation of emissions.

455 The results of the ZECMIP experiments are broadly consistent with previous work on ZEC, with a most likely value of ZEC close to zero and a range of possible model behaviours after emissions cease. In our analysis of ZEC we have shown that terrestrial uptake of carbon plays a more important role in determining that value of ZEC on decadal timescales than has been previously suggested. However our analysis is consistent with previous results from Ehlert and Zickfeld (2017) and Williams et al. (2017) in terms of ZEC arising from balance of physical and biogeochemical factors.

460 Overall, the most likely value of ZEC on decadal time-scales is assessed to be close to zero, consistent with prior work. However substantial continued warming for decades or centuries following cessation of emission is a feature of a minority of the assessed models and thus cannot be ruled out purely on the basis of models.

Data availability. ESM data will be published and freely available as per CMIP6 data policy on the Earth System Grid Federation (<https://esgf-node.llnl.gov/projects/cmip6/>). EMIC data will be published and freely available on a dedicated server (terra.seos.uvic.ca/ZEC). The annual
465 global mean variables used for the present analysis will also be made available on the server.

Appendix A: Model Description Tables

Table A1. Model descriptions of the atmospheric, oceanic, and carbon cycle components for the full Earth System Models (ESMs) that participated in this study.

Model	ACCESS-ESM1.5	CanESM5	CESM2	CNRM-ESM2-1	GFDL-ESM2M
Short Name	ACCESS	CanESM5	CESM2	CNRM	GFDL
Model Expansion	CSIRO Australian Community Climate and Earth System Simulator, ESM version 1.5	Canadian Earth System Model, version 5	Community Earth System Model 2	CNRM-CERFACS Earth System Model, version 2	Geophysical Fluid Dynamics Laboratory Earth system model version 2
Atmosphere	Unified Model (UM) 7.3, 1.875°x1.25°, L38	CanAM5, 2.81° x 2.81°, L49	CAM6, 0.9°x1.25°	ARPEGE-Climat T127 (1.4°), 91 levels	AM2, 2° x 2.5°, L24
Ocean	MOM5, 1° tripolar grid, finer 10S-10N and S. Ocean, L50	NEMO, 1° finer 20°N - 20°S, L45	POP2	NEMO, 1° tripolar grid, L75	MOM4p1, 1° tripolar grid finer at equator, L50
Z-coordinate or Isopycnal Sea Ice	Z-coordinate	Z-coordinate	Z-coordinate	Z-coordinate	Z-coordinate
Land Carbon Cycle	CICE4.1	LIM2	POP2	GELATOv6	SIS
Model Name	CABLE	CLASS-CTEM	CLM5	SURFEX (ISBA-CTRIIP)	LM3.0
Dynamic Vegetation	No	No	No	No	Yes
Nitrogen Cycle	Yes	No	Yes	No	No
Phosphorus Cycle	Yes	No	No	No	No
Permafrost Carbon	No	No	Yes	No	No
Ocean Carbon Cycle					
Model Name	WOMBAT	CMOC	MARBL	PISCESv2-gas	TOPAZ2
Explicit Nutrients	Yes	Yes	yes	yes	Yes
If Yes List	P, Fe	N	N, P, Si, Fe	N, P, Si, Fe	N, P, Si, Fe
Reference	(Law et al., 2017; Ziehn et al., 2020)	(Swart et al., 2019)	(Danabasoglu and Oth-ers, 2019; Lawrence et al., 2019)	(Séférian et al., 2019; Decharme et al., 2019; Delire et al., 2020)	(Dunne et al., 2012, 2013; Burger et al., 2020)

Table A2. Model descriptions of the atmospheric, oceanic, and carbon cycle components for the full Earth System Models (ESMs) that participated in this study.

Model	MIROC-ES2L	MPI-ESM1.2-LR	NorESM2-LM	UKESM1-0-LL
Short Name	MIROC-ES2L	MPI-ESM	NorESM2	UKESM
Model Expansion	Model for Interdisciplinary Research on Climate, Earth System version2 for Long-term simulations	Max-Planck-Institute Earth System model, version 1.2, low resolution	Norwegian Earth System Model 2	United Kingdom Earth System Model, vn1
Atmosphere	CCSR-NIES AGCM, T42, L40	ECHAM6, T63 (ca.1.8°x1.8°), L47	CAM6, 0.9°x2.5°, L32	HadGAM3. N96 (1.25° x 1.875°), L85
Ocean	CCSR Ocean Component model (COCO), 360x256 grids with tripolar grid, L62	MPIOM1.6, GR1.5 (1.5°x1.5°)	Bergen Layered Ocean Model, 1° finer near Equator, L53	NEMO, 1° tripolar grid, L75
Z-coordinate or Isopycnal Sea Ice	Z-coordinate COCO	Z-coordinate Thermodynamic – Dynamic	Isopycnal Community Sea-ice model	Z-coordinate CICE sea ice model
Land Carbon Cycle				
Model Name	MATSIRO/VISIT-e	JSBACH 3.2	CLM5	JULES
Dynamic Vegetation	No	Yes	No	Yes
Nitrogen Cycle	Yes	Yes	Yes	Yes
Phosphorus Cycle	No	No	No	No
Permafrost Carbon	No	No	Yes	No
Ocean Carbon Cycle				
Model Name	OECO2	HAMOCC6	iHAMOCC	MEDUSA-2
Explicit Nutrients	Yes	Yes	Yes	Yes
If Yes List	N, P, Fe	N, P, Si, Fe	P, N, Fe	N, Si, Fe
Reference	(Hajima et al., 2019)	(Ilyina et al., 2013; Mauritsen et al., 2019; Goll et al., 2017)	(Tjiputra et al., 2020)	(Sellar et al., 2019; Best et al., 2011; Clark et al., 2011; Yool et al., 2013)

Table A3. Model descriptions of the atmospheric, oceanic, and carbon cycle components for Earth system models of intermediate complexity (EMICs) that participated in this study.

Model	Bern3D-LPX	CLIMBER-2	DCESS	IAPRAS	LOVECLIM 1.2
Short Name	Bern	CLIMBER	DCESS	IAPRAS	LOVECLIM
Model Expansion	Bern3D-LPX	Climate-Biosphere model, version 2	Danish Center for Earth System Science Earth System Model version 1.0	A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences	LOVECLIM v1.2
Atmosphere	2D Energy-Moisture Balance, 4.5° 9° (<i>onaverage</i>)	Statistical-Dynamical, 51° x 10°	Energy-Moisture Balance model	Statistical-Dynamical model, 4.5° 6.0° ,L11	ECBilt, 5.625° x 5.625°, L3
Ocean	4.5° x 9° (on average) , L32	2D, 3-basin zonally-averaged, 2.5° lat, L21	2 box in lat, 100 m Z resolution	Statistical-Dynamical model 4.5° 6.0°,L3	CLIO, 3° ,L20
Z-coordinate or isopycnal Sea Ice	Z-coordinate 4.5°,9° (on average)	Z-coordinate Thermodynamic-Dynamic	Z-coordinate Mean Surface Temperature Parameterization	Z-coordinate Mean Surface Temperature Parameterization	Z-coordinate Thermodynamic-dynamic
Land Carbon Cycle					
Model Name	LPX v1.4	VECODE			VECODE
Dynamic Vegetation	Yes	Yes	Yes	No	Yes
Nitrogen Cycle	Yes	No	No	No	No
Phosphorus Cycle	No	No	No	No	No
Permafrost Carbon	No	No	No	No	No
Ocean Carbon Cycle					
Model Name	Bern3D v2.0s				LOCH
Explicit Nutrients	Yes	Yes	Yes	No	Yes
If Yes List	P, Si, Fe	P	P		P, Si
Reference	(Ritz et al., 2011; Roth et al., 2014; Jeltsch-Thömmes et al., 2019; Lienert and Joos, 2018)	(Brovkin et al., 2002; Ganopolski et al., 2001)	(Shaffer et al., 2008)	(Eliseev, 2011; Eliseev and Mokhov, 2011; Mokhov and Eliseev, 2012; Eliseev et al., 2014; Mokhov et al., 2020)	(Menviel et al., 2008; Goosse et al., 2010; Mouchet, 2011)

Table A4. Model descriptions of the atmospheric, oceanic, and carbon cycle components for Earth system models of intermediate complexity (EMICs) that participated in this study.

Model	MESM	MIROC- lite/JUMP- LCM	PLASIM- GENIE	UVic ESCM 2.10
Short Name	MESM	MIROC-lite	P. GENIE	UVic
Model Expansion	MIT Earth System Model	Model for In- terdisciplinary Research on Climate- lite/Japan Uncertainty Modelling Project- Loosely Coupled Model	Planet Sim- ulator - Grid-ENabled Integrated Earth system model	University of Victoria Earth System Cli- mate Model version 2.10
Atmosphere	Zonally aver- aged 4° lat., L11	2D Energy- Moisture Balance, 6° x 6°	PLASIM, T21, L10	2D Energy- Moisture Balance, 3.6°1.8°,L1
Ocean	4°x5° mixed layer, anomaly diffusing model	CCSR Ocean Compo- nent model (COCO), 6° x 6°, L15	GOLDSTEIN, T21, L16	MOM2, 3.6°1.8°,L19
Z-coordinate isopycnal Sea Ice	or NA Thermodynamic- Dynamic	Z-coordinate COCO	Z-coordinate GOLDSTEIN	Z-coordinate Thermodynamic- Dynamic
Land Carbon Cycle				
Model Name	TEM	Sim-CYCLE	ENTS	TRIFFID
Dynamic Vegetation	No	No	Yes	Yes
Nitrogen Cycle	Yes	No	No	No
Phosphorus Cycle	No	No	No	No
Permafrost Carbon	No	No	No	Yes
Ocean Carbon Cycle				
Model Name	OCM		BIOGEM	2NZPD
Explicit Nutrients	No	Yes	Yes	Yes
If Yes List		N	P	N, P
Reference	(Sokolov et al., 2018)	(Tachiiri et al., 2010; Oka et al., 2011)	(Holden et al., 2018, 2019)	(Mengis et al., 2020)

Appendix B: Analytical framework

A key question for this study is explaining why some models have positive ZECs and some models have negative or close to zero ZECs. From elementary theory we understand that the sign of ZEC will depend on the pathway of atmospheric CO₂ concentration and ocean heat uptake following cessation of emissions. Complicating this dynamic is that atmospheric CO₂ change has contributions both from ocean carbon uptake and the net flux from the terrestrial biosphere. Here we devise a simple method for partitioning the contribution to ZEC from the ocean carbon flux, net land carbon flux, and the ocean heat uptake.

We begin with the forcing response equation (Wigley and Schlesinger, 1985):

$$F = \lambda T + \epsilon N, \quad (B1)$$

where F (W m⁻²) is radiative forcing, N (W m⁻²) is planetary heat uptake, ϵ (dimensionless) is the efficacy of planetary heat uptake, λ (W m⁻²K⁻¹) is the climate feedback parameter, and T (K) is the change in global temperature (relative to pre-industrial). This equation can be re-written as:

$$\lambda T = F - \epsilon N. \quad (B2)$$

To compute the rate of change of temperature we take the derivative of equation B2 in time giving:

$$\lambda \frac{dT}{dt} = \frac{dF}{dt} - \epsilon \frac{dN}{dt}. \quad (B3)$$

Radiative forcing from CO₂ can be approximated using the classical logarithmic relationship (Myhre et al., 1998):

$$F = R \ln \left(\frac{C_A}{C_{Ao}} \right), \quad (B4)$$

where R (W m⁻²) is the radiative forcing from an e-fold increase in atmospheric CO₂ concentration, C_A (PgC) is atmospheric CO₂ burden and C_{Ao} (PgC) is the original atmospheric CO₂ burden. Recalling that the derivative of $\frac{d \ln(x)}{dx} = \frac{1}{x}$ the derivative of equation B4 is:

$$\frac{dF}{dt} = R \left(\frac{C_{Ao}}{C_A} \right) \left(\frac{dC_A}{dt} \right) \frac{1}{C_{Ao}}, \quad (B5)$$

which simplifies to:

$$\frac{dF}{dt} = \frac{R}{C_A} \left(\frac{dC_A}{dt} \right). \quad (B6)$$

490 After emissions cease atmospheric CO₂ concentration can be expressed as:

$$C_A = C_{ze} - (C_O - C_{Oze}) - (C_L - C_{Lze}), \quad (\text{B7})$$

Where C_{ze} (PgC) is atmospheric CO₂ burden at the time emissions reach zero, C_O (PgC) is the carbon content of the ocean, and C_L (PgC) is the carbon content of land. C_{Oze} (PgC) is the carbon content of the ocean at the time emissions reach zero and C_{Lze} (PgC) is the carbon content of land when emissions reach zero. Thus the derivative of C_A is:

$$495 \quad \frac{dC_A}{dt} = -\frac{dC_O}{dt} - \frac{dC_L}{dt} \quad (\text{B8})$$

$\frac{dC_O}{dt}$ is the flux of carbon into the ocean f_O , and $\frac{dC_L}{dt}$ is the flux of carbon into land f_L .

$$\frac{dC_A}{dt} = -f_O - f_L. \quad (\text{B9})$$

Substituting equation B9 in equation B6 we find:

$$\frac{dF}{dt} = -R \frac{f_O + f_L}{C_A}, \quad (\text{B10})$$

500 which can be split into:

$$\frac{dF}{dt} = -R \frac{f_O}{C_A} - R \frac{f_L}{C_A}, \quad (\text{B11})$$

which can be substituted into equation B3:

$$\lambda \frac{dT}{dt} = -R \frac{f_O}{C_A} - R \frac{f_L}{C_A} - \epsilon \frac{dN}{dt}. \quad (\text{B12})$$

If we integrate equation B12 from time emissions reach zero we get:

$$505 \quad \lambda T_{ZEC} = -R \int_{t=ze}^{\infty} \frac{f_O}{C_A} dt - R \int_{t=ze}^{\infty} \frac{f_L}{C_A} dt - \epsilon(N - N_{ze}), \quad (\text{B13})$$

where T_{ZEC} (K) is ZEC, N_{ze} (W m⁻²) is the planetary heat uptake when emissions cease. The integrals $\int_{t=ze}^{\infty} \frac{f_O}{C_A} dt$ and $\int_{t=ze}^{\infty} \frac{f_L}{C_A} dt$ can be computed numerically from ZECMIP output. Therefore we define:

$$F_{ocean} = R \int_{t=ze}^{\infty} \frac{f_O}{C_A} dt, \quad (\text{B14})$$

and,

$$510 \quad F_{land} = R \int_{t=ze}^{\infty} \frac{f_L}{C_A} dt; \quad (\text{B15})$$

and thus:

$$\lambda T_{ZEC} = -F_{ocean} - F_{land} - \epsilon(N - N_{ze}), \quad (\text{B16})$$

Appendix C: ZEC values for A2 and A3 experiments

Table C1. Temperature anomaly relative to the year emissions cease averaged over a 20 year time window centred about the 25th, 50th, and 90th year following cessation of anthropogenic CO₂ emissions (ZEC₂₅, ZEC₅₀, and ZEC₉₀ respectively) for the A2 (750 PgC 1% experiment).

Model	ZEC ₂₅ (°C)	ZEC ₅₀ (°C)	ZEC ₉₀ (°C)
ACCESS	-0.03	-0.06	-0.12
GFDL	-0.26	-0.31	-0.26
MIROC-ES2L	-0.04	-0.17	-0.20
UKESM	0.13	0.11	0.08
Bern	0.03	-0.02	-0.09
DCESS	0.12	0.07	-0.02
IAPRAS	0.34	0.30	0.08
LOVECLIM	-0.06	-0.06	0.05
MESM	0.0	-0.07	-0.13
MIROC-lite	0.0	-0.06	-0.09
P. GENIE	0.03	-0.13	-0.12
UVic	0.02	-0.03	-0.06
Mean	0.02	-0.03	-0.07
Median	0.0	-0.06	-0.09
Standard Deviation	0.13	0.15	0.10

Table C2. Temperature anomaly relative to the year emissions cease averaged over a 20 year time window centred about the 25th, 50th, and 90th year following cessation of anthropogenic CO₂ emissions (ZEC₂₅, ZEC₅₀, and ZEC₉₀ respectively) for the A3 (2000 PgC 1% experiment).

Model	ZEC ₂₅ (°C)	ZEC ₅₀ (°C)	ZEC ₉₀ (°C)
ACCESS	0.21	0.52	0.65
CanESM5	0.01	-0.06	-0.05
GFDL	-0.12	-0.14	-0.06
MIROC-ES2L	0.03	-0.07	-0.12
UKESM	0.37	0.51	0.75
Bern	0.17	0.17	0.15
DCESS	0.09	0.06	-0.01
IAPRAS	0.34	0.28	0.0
MESM	0.12	0.13	0.11
MIROC-lite	0.08	0.03	-0.02
P. GENIE	-0.15	-0.40	-0.48
UVic	0.11	0.21	0.33
Mean	0.11	0.10	0.10
Median	0.10	0.10	-0.01
Standard Deviation	0.16	0.26	0.34

Appendix D: Efficacy Evolution




Figure A1. Evolution of efficacy for the four EMICs without substantial internal variability. Red horizontal line is the efficacy value estimated from the 1pctCO₂ experiment and vertical blue line marks 50 years after emissions cease.

515 *Author contributions.* CDJ, TLF, AHMD, JR, HDM, and KZ initiated the research. CDJ and AHMD coordinated the project. AHMD, VKA, VB, FAB, AVE, TH, PBH, AJT, CK, NM, LM, MM, IIM, RS, GS, JT, JS, TZ, KT, and AO carried out model simulations. AHMD, CDJ, TLF, JR, HDM, and KZ contributed to the analysis. ME and NJB contributed critical technical support and programming. AHMD, HDM, VKA, and JR contributed significantly to the writing of the paper.

Competing interests. AVE is a member of the journal Editorial Board. All other authors declare no competing interests

520 *Acknowledgements.* AHMD, ME, HDM, and KZ are each grateful for support from respective National Science and Engineering Research
Council of Canada Discovery Grants. AHMD is thankful to Compute Canada. TLF and FAB acknowledge support from the Swiss National
Science Foundation under grant PP00P2 - 170687 and from the European Union's Horizon 2020 Research and Innovative Programme
under grant agreement number 821003 (CCiCC). GFDL ESM2M simulations were performed at the Swiss National Supercomputing Centre
(CSCS). CDJ and AW acknowledge the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101) and the EU
525 Horizon 2020 CRESCENDO project, grant number 641816. JR, RS and MM acknowledge support from the European Union's Horizon
2020 Research and Innovation Programme under grant agreement number 820829 (CONSTRAIN). VB is grateful to Thomas Raddatz and
Veronika Gayler for assistance with the MPI-ESM experimental setup and acknowledges support from the EU Horizon 2020 CRESCENDO
project. AVE was supported by the Russian Foundation for Basic Research (grant 18-05-00087). TH and KT acknowledge the support from
"Integrated Research Program for Advancing Climate Models" by the Ministry of Education, Culture, Sports, Science, and Technology of
530 Japan. PBH was funded by LC3M, a Leverhulme Trust Research Centre Award (RC-2015-029). AJT acknowledges support by the Swiss
National Science Foundation (# 200020_172476) and the Oeschger Centre for Climate Change Research. IIM was supported by the Russian
Science Foundation (grant 19-17-00240). GS was supported by FONDECYT (Chile) grant 1190230. LM acknowledges support from the
Australian Research Council FT180100606. RS and MM acknowledge the EU Horizon 2020 CRESCENDO project, grant number 641816
and Supercomputing time provided by the Météo-France/DSI supercomputing center. JT and JS acknowledge Research Council of Norway
535 funded projects KeyClim (295046) and IMPOSE (294930).

The authors thank all participants of the expert workshop on carbon budgets, co-organized with the support of the Global Carbon Project, the CRESCENDO project, the University of Melbourne, and Simon Fraser University, for discussions that initiated the authors' thinking about ZECMIP.

We thank R. Williams and an anonymous reviewer for their helpful comments.

540 **References**

- Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J., Mahowald, N., Mulugetta, Y., Perez, R., Wairiu, M., and Zickfeld, K.: Framing and Context, in: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, World Meteorological Organization, 2018.
- 545 Andrews, T., Gregory, J. M., and Webb, M. J.: The dependence of radiative forcing and feedback on evolving patterns of surface temperature change in climate models, *Journal of Climate*, 28, 1630–1648, 2015.
- Archer, D. and Brovkin, V.: The millennial atmospheric lifetime of anthropogenic CO₂, *Climatic Change*, 90, 283–297, 2008.
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven, G., Montenegro, A., et al.: Atmospheric lifetime of fossil fuel carbon dioxide, *Annual review of earth and planetary sciences*, 37, 117–134, 2009.
- 550 Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., and Tatiana Ilyina, T. H., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon-Concentration and Carbon-Climate Feedbacks in CMIP5 Earth System Models., *Journal of Climate*, 26, 2013.
- Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C., Krasting, J., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T.: Carbon-concentration and carbon-climate feedbacks in CMIP6 models, and their comparison to CMIP5 models, *Biogeosciences Discussions*, 2019, 1–124, <https://doi.org/10.5194/bg-2019-473>, <https://www.biogeosciences-discuss.net/bg-2019-473/>, 2019.
- 555 Banks, H. T. and Gregory, J. M.: Mechanisms of ocean heat uptake in a coupled climate model and the implications for tracer based predictions of ocean heat uptake, *Geophysical Research Letters*, 33, 2006.
- Best, M., Pryor, M., Clark, D., Rooney, G., Essery, R., Ménard, C., Edwards, J., Hendry, M., Porson, A., Gedney, N., et al.: The Joint UK Land Environment Simulator (JULES), model description—Part 1: energy and water fluxes, *Geoscientific Model Development*, 4, 677–699, 2011.
- 565 Brovkin, V., Bendtsen, J., Claussen, M., Ganopolski, A., Kubatzki, C., Petoukhov, V., and Andreev, A.: Carbon cycle, vegetation, and climate dynamics in the Holocene: Experiments with the CLIMBER-2 model, *Global Biogeochemical Cycles*, 16, 86–1, 2002.
- Burger, F. A., Frölicher, T. L., and John, J. G.: Increase in ocean acidity variability and extremes under increasing atmospheric CO₂, *Biogeosciences Discussions*, 2020, 1–32, <https://doi.org/10.5194/bg-2020-22>, <https://www.biogeosciences-discuss.net/bg-2020-22/>, 2020.
- Byrne, B. and Goldblatt, C.: Radiative forcing at high concentrations of well-mixed greenhouse gases, *Geophysical Research Letters*, 41, 570 152–160, 2014.
- Cao, L., Eby, M., Ridgwell, A., Caldeira, K., Archer, D., Ishida, A., Joos, F., Matsumoto, K., Mikolajewicz, U., Mouchet, A., et al.: The role of ocean transport in the uptake of anthropogenic CO₂, *Biogeosciences*, 6, 375–390, 2009.
- Charney, J., Arakawa, A., Baker, D., Bolin, B., Dickinson, R., Goody, R., Leith, C., Stommel, H., and Wunsch, C.: Carbon dioxide and climate: a scientific assessment: report of an ad hoc study group on carbon dioxide and climate, woods hole, Massachusetts, July 23–27, 1979 to the climate research board, assembly of mathematical and physical sciences, National Research Council, National Academy of Sciences. Climate Research Board. <http://books.google.com/books>, 1979.
- 575

- Clark, D., Mercado, L., Sitch, S., Jones, C., Gedney, N., Best, M., Pryor, M., Rooney, G., Essery, R., Blyth, E., et al.: The Joint UK Land Environment Simulator (JULES), model description–Part 2: carbon fluxes and vegetation dynamics, *Geoscientific Model Development*, 4, 701–722, 2011.
- 580 Danabasoglu, G. and Others: The Community Earth System Model 2 (CESM2), Submitted to *Journal of Advances in Modeling the Earth System*, 2019.
- Davis, S. J. and Socolow, R. H.: Commitment accounting of CO₂ emissions, *Environmental Research Letters*, 9, 084 018, 2014.
- Davis, S. J., Caldeira, K., and Matthews, H. D.: Future CO₂ emissions and climate change from existing energy infrastructure, *Science*, 329, 1330–1333, 2010.
- 585 Decharme, B., Delire, C., Minvielle, M., Colin, J., Vergnes, J.-P., Alias, A., Saint-Martin, D., Sférian, R., Sénési, S., and Voltaire, A.: Recent Changes in the ISBA-CTRIP Land Surface System for Use in the CNRM-CM6 Climate Model and in Global Off-Line Hydrological Applications, *Journal of Advances in Modeling Earth Systems*, 2019.
- Delire, C., Sférian, R., Decharme, B., Alkama, R., Carrer, D., Joetzjer, E., Morel, X., and Rocher, M.: The global land carbon cycle simulated with ISBA, *Journal of Advances in Modeling Earth Systems*, <https://doi.org/10.1029/2019MS001886>, 2020.
- 590 Dunne, J. P., John, J. G., Adcroft, A. J., Griffies, S. M., Hallberg, R. W., Shevliakova, E., Stouffer, R. J., Cooke, W., Dunne, K. A., Harrison, M. J., et al.: GFDL’s ESM2 global coupled climate-carbon earth system models. Part I: Physical formulation and baseline simulation characteristics, *Journal of Climate*, 25, 6646–6665, 2012.
- Dunne, J. P., John, J. G., Shevliakova, E., Stouffer, R. J., Krasting, J. P., Malyshev, S. L., Milly, P., Sentman, L. T., Adcroft, A. J., Cooke, W., et al.: GFDL’s ESM2 global coupled climate-carbon earth system models. Part II: carbon system formulation and baseline simulation
- 595 characteristics, *Journal of Climate*, 26, 2247–2267, 2013.
- Eby, M., Zickfeld, K., Montenegro, A., Archer, D., Meissner, K. J., and Weaver, A. J.: Lifetime of Anthropogenic Climate Change: Millennial Time Scales of Potential CO₂ and Surface Temperature Perturbations, *Journal of Climate*, 22, 2501–2511, <https://doi.org/10.1175/2008JCLI2554.1>, 2009.
- Ehlert, D. and Zickfeld, K.: What determines the warming commitment after cessation of CO₂ emissions?, *Environmental Research Letters*, 12, 015 002, 2017.
- 600 Eliseev, A.: Estimation of changes in characteristics of the climate and carbon cycle in the 21st century accounting for the uncertainty of terrestrial biota parameter values, *Izvestiya, Atmospheric and Oceanic Physics*, 47, 131, 2011.
- Eliseev, A., Mokhov, I., and Chernokulsky, A.: An ensemble approach to simulate CO₂ emissions from natural fires, *Biogeosciences*, 11, 3205–3223, 2014.
- 605 Eliseev, A. V. and Mokhov, I. I.: Uncertainty of climate response to natural and anthropogenic forcings due to different land use scenarios, *Advances in Atmospheric Sciences*, 28, 1215, 2011.
- Enting, I. G., Wigley, T., and Heimann, M.: Future emissions and concentrations of carbon dioxide: key ocean/atmosphere/land analyses, 1994.
- Etminan, M., Myhre, G., Highwood, E., and Shine, K.: Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision
- 610 of the methane radiative forcing, *Geophysical Research Letters*, 43, 12–614, 2016.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, 2016.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiy, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler,

- 615 K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate–Carbon Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison, *Journal of Climate*, 19, 3337–3353, 2006.
- Frölicher, T. L. and Joos, F.: Reversible and irreversible impacts of greenhouse gas emissions in multi-century projections with the NCAR global coupled carbon cycle-climate model, *Climate Dynamics*, 35, 1439–1459, 2010.
- Frölicher, T. L. and Paynter, D. J.: Extending the relationship between global warming and cumulative carbon emissions to multi-millennial
620 timescales, *Environmental Research Letters*, 10, 075 002, 2015.
- Frölicher, T. L., Winton, M., and Sarmiento, J. L.: Continued global warming after CO₂ emissions stoppage, *Nature Climate Change*, 4, 40–44, 2014.
- Frölicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., and Winton, M.: Dominance of the Southern Ocean in anthropogenic carbon and heat uptake in CMIP5 models, *Journal of Climate*, 28, 862–886, 2015.
- 625 Ganopolski, A., Petoukhov, V., Rahmstorf, S., Brovkin, V., Claussen, M., Eliseev, A., and Kubatzki, C.: CLIMBER-2: a climate system model of intermediate complexity. Part II: model sensitivity, *Climate Dynamics*, 17, 735–751, 2001.
- Geden, O.: An actionable climate target, *Nature Geoscience*, 9, 340, <https://doi.org/10.1038/ngeo2699>, <http://dx.doi.org/10.1038/ngeo2699>, 2016.
- Gillett, N., Arora, V., Zickfeld, K., Marshall, S., and Merryfield, W.: Ongoing climate change following a complete cessation of carbon
630 dioxide emissions, *Nature Geoscience*, 4, 83–87, 2011.
- Goll, D. S., Winkler, A. J., Raddatz, T., Dong, N., Prentice, I. C., Ciais, P., and Brovkin, V.: Carbon–nitrogen interactions in idealized simulations with JSBACH (version 3.10), *Geoscientific Model Development*, 10, 2009–2030, <https://doi.org/10.5194/gmd-10-2009-2017>, <https://www.geosci-model-dev.net/10/2009/2017/>, 2017.
- Goodwin, P., Williams, R. G., Follows, M. J., and Dutkiewicz, S.: Ocean–atmosphere partitioning of anthropogenic carbon dioxide on
635 centennial timescales, *Global Biogeochemical Cycles*, 21, 2007.
- Goodwin, P., Williams, R. G., and Ridgwell, A.: Sensitivity of climate to cumulative carbon emissions due to compensation of ocean heat and carbon uptake, *Nature Geoscience*, 8, 29–34, 2015.
- Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Selten, F., Barriat, P.-Y., Campin, J.-M., Deleersnijder, E., Driesschaert, E., Goelzer, H., Janssens, I., Loutre, M.-F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P.-P.,
640 Munhoven, G., Pettersson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinville, B., Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate complexity LOVECLIM version 1.2, *Geoscientific Model Development*, 3, 603–633, <https://doi.org/10.5194/gmd-3-603-2010>, <https://www.geosci-model-dev.net/3/603/2010/>, 2010.
- Gregory, J., Ingram, W., Palmer, M., Jones, G., Stott, P., Thorpe, R., Lowe, J., Johns, T., and Williams, K.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophysical Research Letters*, 31, 2004.
- 645 Gregory, J. M., Andrews, T., and Good, P.: The inconstancy of the transient climate response parameter under increasing CO₂, *Phil. Trans. R. Soc. A*, 373, 20140 417, 2015.
- Haites, E., Yamin, F., and Höhne, N.: Possible Elements of a 2015 Legal Agreement on Climate Change, IDDRI SciencesPo Working Paper, pp. 1–24, <http://theredddesk.org/sites/default/files/resources/pdf/Possible%20Elements%20of%20a%202015%20Legal%20Agreement%20on%20Climate%20Change.pdf>, 2013.
- 650 Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Description of the MIROC-ES2L Earth system model and evaluation of its cli-

- mate–biogeochemical processes and feedbacks, *Geoscientific Model Development Discussions*, 2019, 1–73, <https://doi.org/10.5194/gmd-2019-275>, <https://www.geosci-model-dev-discuss.net/gmd-2019-275/>, 2019.
- 655 Hare, B. and Meinshausen, M.: How much warming are we committed to and how much can be avoided?, *Climatic Change*, 75, 111–149, 2006.
- Holden, P. B., Edwards, N. R., Ridgwell, A., Wilkinson, R., Fraedrich, K., Lunkeit, F., Pollitt, H., Mercure, J.-F., Salas, P., Lam, A., et al.: Climate–carbon cycle uncertainties and the Paris Agreement, *Nature Climate Change*, 8, 609, 2018.
- Holden, P. B., Edwards, N. R., Rangel, T. F., Pereira, E. B., Tran, G. T., and Wilkinson, R. D.: PALEO-PGEM v1.0: A statistical emulator of Pliocene-Pleistocene climate, *Geoscientific Model Development*, 12, 5137–5155, <https://doi.org/10.5194/gmd-12-5137-2019>, <https://www.geosci-model-dev.net/12/5137/2019/>, 2019.
- 660 Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287–315, 2013.
- IPCC: Summary for Policymakers, in: Working Group I Contribution to the Intergovernmental Panel on Climate Change Fifth Assessment Report Climate Change 2013: The Physical Science Basis, edited by Alexander, L., Allen, S., Bindoff, N. L., Bréon, F.-M., Church, J., Cubasch, U., Emori, S., Forster, P., Friedlingstein, P., Gillett, N., Gregory, J., Hartmann, D., Jansen, E., Kirtman, B., Knutti, R., Kani-
665 kicharla, K. K., Lemke, P., Marotzke, J., Masson-Delmotte, V., Meehl, G., Mokhov, I., Piao, S., Plattner, G.-K., Dahe, Q., Ramaswamy, V., Randall, D., Rhein, M., Rojas, M., Sabine, C., Shindell, D., Stocker, T. F., Talley, L., Vaughan, D., and Xie, S.-P., Cambridge University Press, 2013.
- 670 Jeltsch-Thömmes, A. T. D., Battaglia, G., Cartapanis, O., Jaccard, S., and Joos, F.: Low terrestrial carbon storage at the Last Glacial Maximum: constraints from multi-proxy data, *Climate of the Past*, 15, 849–879, 2019.
- Jenkinson, D., Adams, D., and Wild, A.: Model estimates of CO₂ emissions from soil in response to global warming, *Nature*, 351, 304–306, 1991.
- Jones, C. D., Frölicher, T. L., Koven, C., MacDougall, A. H., Matthews, H. D., Zickfeld, K., Rogelj, J., Tokarska, K. B., Gillett, N., Ilyina, T., Meinshausen, M., Mengis, N., Séférian, R., and Eby, M.: The Zero Emission Commitment Model Intercomparison Project (ZECMIP) contribution to CMIP6: Quantifying committed climate changes following zero carbon emissions, In Review *Geoscientific Model Development*, 2019.
- 675 Joos, F., Roth, R., Fuglestedt, J., Peters, G., Enting, I., Bloh, W. v., Brovkin, V., Burke, E., Eby, M., Edwards, N., et al.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmospheric Chemistry and Physics*, 13, 2793–2825, 2013.
- 680 Knutti, R., Rugenstein, M. A., and Hegerl, G. C.: Beyond equilibrium climate sensitivity, *Nature Geoscience*, 10, ngeo3017, 2017.
- Law, R. M., Ziehn, T., Matear, R. J., Lenton, A., Chamberlain, M. A., Stevens, L. E., Wang, Y.-P., Srbinovsky, J., Bi, D., Yan, H., et al.: The carbon cycle in the Australian Community Climate and Earth System Simulator (ACCESS-ESM1)–Part 1: Model description and pre-industrial simulation, *Geoscientific Model Development*, 10, 2567–2590, 2017.
- 685 Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., van Kampenhout, L., Kennedy, D., et al.: The Community Land Model version 5: Description of new features, benchmarking, and impact of forcing uncertainty, *Journal of Advances in Modeling Earth Systems*, 2019.
- Lienert, S. and Joos, F.: A Bayesian ensemble data assimilation to constrain model parameters and land-use carbon emissions, *Biogeosciences*, 15, 2909–2930, 2018.

- 690 Lowe, J. A., Huntingford, C., Raper, S. C. B., Jones, C. D., Liddicoat, S. K., and Gohar, L. K.: How difficult is it to recover from dangerous levels of global warming?, *Environmental Research Letters*, 4, 014 012, 2009.
- MacDougall, A. H.: The oceanic origin of path-independent carbon budgets, *Scientific Reports*, 7, 10 373, <https://doi.org/10.1038/s41598-017-10557-x>, 2017.
- MacDougall, A. H.: Limitations of the 1% experiment as the benchmark idealized experiment for carbon cycle intercomparison in C⁴MIP, 695 *Geoscientific Model Development*, 12, 597–611, 2019.
- Maier-Reimer, E. and Hasselmann, K.: Transport and storage of CO₂ in the ocean: an inorganic ocean-circulation carbon cycle model, *Climate dynamics*, 2, 63–90, 1987.
- Matthews, H. and Weaver, A.: Committed climate warming, *Nature Geoscience*, 3, 142–143, 2010.
- Matthews, H. and Zickfeld, K.: Climate response to zeroed emissions of greenhouse gases and aerosols, *Nature Climate Change*, 2, 338–341, 700 2012.
- Matthews, H. D.: A growing commitment to future CO₂ emissions, *Environmental Research Letters*, 9, 111 001, 2014.
- Matthews, H. D. and Caldeira, K.: Stabilizing climate requires near-zero emissions, *Geophysical Research Letters*, 35, L04 705, <https://doi.org/10.1029/2007GL032388>, 2008.
- Matthews, H. D. and Solomon, S.: Irreversible does not mean unavoidable, *Science*, 340, 438–439, 2013.
- 705 Mauritsen, T. and Pincus, R.: Committed warming inferred from observations, *Nature Climate Change*, 7, 652, 2017.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., et al.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, 2019.
- Meehl, G. A., Washington, W. M., Collins, W. D., Arblaster, J. M., Hu, A., Buja, L. E., Strand, W. G., and Teng, H.: How much more global 710 warming and sea level rise?, *science*, 307, 1769–1772, 2005.
- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S., Watterson, I., Weaver, A., and Zhao, Z.-C.: Global climate projections, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M., and Miller, H., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- 715 Mengis, N., Keller, D. P., MacDougall, A., Eby, M., Wright, N., Meissner, K. J., Oschlies, A., Schmittner, A., Matthews, H. D., and Zickfeld, K.: Evaluation of the University of Victoria Earth System Climate Model version 2.10 (UVic ESCM 2.10), *Geoscientific Model Development Discussions*, 2020, 1–28, <https://doi.org/10.5194/gmd-2019-373>, <https://www.geosci-model-dev-discuss.net/gmd-2019-373/>, 2020.
- 720 Menviel, L., Timmermann, A., Mouchet, A., and Timm, O.: Meridional reorganizations of marine and terrestrial productivity during Heinrich events, *Paleoceanography*, 23, PA1203, <https://doi.org/doi:10.1029/2007PA001445>, 2008.
- Mokhov, I. and Eliseev, A.: Modeling of global climate variations in the 20th-23rd centuries with new RCP scenarios of anthropogenic forcing, in: *Doklady Earth Sciences*, vol. 443, p. 532, Springer Science & Business Media, 2012.
- Mokhov, I., Eliseev, A., and Guryanov, V.: Model estimates of global and regional climate changes in the Holocene, In Press: *Doklady Earth 725 Sci.*, 2020.
- Mouchet, A.: A 3D model of ocean biogeochemical cycles and climate sensitivity studies, Ph.D. thesis, Université de Liège, Liège, Belgium, <http://hdl.handle.net/2268/98995>, 2011.

- Myhre, G., Highwood, E. J., Shine, K. P., and Stordal, F.: New estimates of radiative forcing due to well mixed greenhouse gases, *Geophysical research letters*, 25, 2715–2718, 1998.
- 730 Oka, A., Tajika, E., Abe-Ouchi, A., and Kubota, K.: Role of the ocean in controlling atmospheric CO₂ concentration in the course of global glaciations, *Climate dynamics*, 37, 1755–1770, 2011.
- Planton, S.: Annex III: Glossary, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2013.
- 735 Plattner, G.-K., Knutti, R., Joos, F., Stocker, T., Von Bloh, W., Brovkin, V., Cameron, D., Driesschaert, E., Dutkiewicz, S., Eby, M., Edwards, N. R., Fichetfet, T., Hargreaves, C., Jones, C. D., Loutre, M. F., Matthews, H. D., Mouchet, A., Möller, S. A., Nawrath, S., Price, A., Sokolov, A., Strassmann, K. M., and Weaver, A. J.: Long-term climate commitments projected with climate–carbon cycle models, *Journal of Climate*, 21, 2721–2751, 2008.
- Ritz, S. P., Stocker, T. F., and Joos, F.: A coupled dynamical ocean–energy balance atmosphere model for paleoclimate studies, *Journal of Climate*, 24, 349–375, 2011.
- 740 Rogelj, J., Schaeffer, M., Meinshausen, M., Knutti, R., Alcamo, J., Riahi, K., and Hare, W.: Zero emission targets as long-term global goals for climate protection, *Environmental Research Letters*, 10, 105 007, <https://doi.org/10.1088/1748-9326/10/10/105007>, <http://stacks.iop.org/1748-9326/10/i=10/a=105007>, 2015.
- Rogelj, J., den Elzen, M., Höhne, N., Fransen, T., Fekete, H., Winkler, H., Schaeffer, R., Sha, F., Riahi, K., and Meinshausen, M.: Paris Agreement climate proposals need a boost to keep warming well below 2°C, *Nature*, 534, 631–639, <https://doi.org/10.1038/nature18307>,
745 <http://dx.doi.org/10.1038/nature18307>, 2016.
- Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., Mundaca, L., Seférian, R., and Vilarino, M. V.: Mitigation pathways compatible with 1.5°C in the context of sustainable development, in: *Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and*
750 *efforts to eradicate poverty*, World Meteorological Organization, 2018.
- Rogelj, J., Forster, P. M., Kriegler, E., Smith, C. J., and Sférian, R.: Estimating and tracking the remaining carbon budget for stringent climate targets, *Nature*, 571, 335, 2019a.
- Rogelj, J., Huppmann, D., Krey, V., Riahi, K., Clarke, L., Gidden, M., Nicholls, Z., and Meinshausen, M.: A new scenario logic for the Paris Agreement long-term temperature goal, *Nature*, 573, 357–363, <https://doi.org/10.1038/s41586-019-1541-4>, <https://doi.org/10.1038/s41586-019-1541-4>, 2019b.
755
- Rose, B. E., Armour, K. C., Battisti, D. S., Feldl, N., and Koll, D. D.: The dependence of transient climate sensitivity and radiative feedbacks on the spatial pattern of ocean heat uptake, *Geophysical Research Letters*, 41, 1071–1078, 2014.
- Roth, R., Ritz, S., and Joos, F.: Burial-nutrient feedbacks amplify the sensitivity of atmospheric carbon dioxide to changes in organic matter remineralisation, *Earth system dynamics*, 5, 321–343, 2014.
- 760 Rugenstein, M. A., Gregory, J. M., Schaller, N., Sedláček, J., and Knutti, R.: Multiannual ocean–atmosphere adjustments to radiative forcing, *Journal of Climate*, 29, 5643–5659, 2016.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C., Wallace, D. W., Tilbrook, B., et al.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sarmiento, J. L. and Gruber, N.: *Ocean biogeochemical dynamics*, Princeton University Press, 2006.

- 765 Sarmiento, J. L., Orr, J. C., and Siegenthaler, U.: A perturbation simulation of CO₂ uptake in an ocean general circulation model, *Journal of Geophysical Research: Oceans*, 97, 3621–3645, 1992.
- Schuur, E., McGuire, A., Schädel, C., Grosse, G., Harden, J., Hayes, D., Hugelius, G., Koven, C., Kuhry, P., Lawrence, D., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, 2015.
- 770 Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., et al.: Evaluation of CNRM Earth-System model, CNRM-ESM 2-1: role of Earth system processes in present-day and future climate, *Journal of Advances in Modeling Earth Systems*, 2019.
- Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., O’Connor, F. M., Stringer, M., Hill, R., Palmieri, J., et al.: UKESM1: Description and evaluation of the UK Earth System Model, *Journal of Advances in Modeling Earth Systems*, 2019.
- 775 Shaffer, G., Malskær Olsen, S., and Pepke Pedersen, J. O.: Presentation, calibration and validation of the low-order, DCESS Earth System Model (Version 1), *Geoscientific Model Development*, 1, 17–51, <https://doi.org/10.5194/gmd-1-17-2008>, <https://www.geosci-model-dev.net/1/17/2008/>, 2008.
- Siegenthaler, U. and Joos, F.: Use of a simple model for studying oceanic tracer distributions and the global carbon cycle, *Tellus B*, 44, 186–207, 1992.
- 780 Smith, C. J., Forster, P. M., Allen, M., Fuglestedt, J., Millar, R. J., Rogelj, J., and Zickfeld, K.: Current fossil fuel infrastructure does not yet commit us to 1.5°C warming, *Nature communications*, 10, 101, 2019.
- Sokolov, A., Kicklighter, D., Schlosser, A., Wang, C., Monier, E., Brown-Steiner, B., Prinn, R., Forest, C., Gao, X., Libardoni, A., et al.: Description and evaluation of the MIT Earth System Model (MESM), *Journal of Advances in Modeling Earth Systems*, 10, 1759–1789, 2018.
- 785 Solomon, S., Plattner, G.-K., Knutti, R., and Friedlingstein, P.: Irreversible climate change due to carbon dioxide emissions, *Proceedings of the national academy of sciences*, 106, 1704–1709, 2009.
- Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., et al.: The Canadian Earth System Model version 5 (CanESM5. 0.3), *Geoscientific Model Development*, 12, 4823–4873, 2019.
- Tachiiri, K., Hargreaves, J. C., Annan, J. D., Oka, A., Abe-Ouchi, A., and Kawamiya, M.: Development of a system emulating the global carbon cycle in Earth system models, *Geoscientific Model Development*, 3, 365–376, <https://doi.org/10.5194/gmd-3-365-2010>, <https://www.geosci-model-dev.net/3/365/2010/>, 2010.
- 790 Tjiputra, J. F., Schwinger, J., Bentsen, M., Moree, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y., Olivie, D., Seland, O., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), Submitted to *Geoscientific Model Development*, 2020.
- 795 Tong, D., Zhang, Q., Zheng, Y., Caldeira, K., Shearer, C., Hong, C., Qin, Y., and Davis, S. J.: Committed emissions from existing energy infrastructure jeopardize 1.5°C climate target, *Nature*, p. 1, 2019.
- UNEP: The Emissions Gap Report 2018, Report, UNEP, <https://doi.org/ISBN:978-92-807-3726-4>, 2018.
- United Nations: Paris Agreement: Twenty-first conference of the parties of the United Nations Framework Convention on Climate Change, United Nations, 2015.
- 800 Wigley, T. M.: The climate change commitment, *Science*, 307, 1766–1769, 2005.
- Wigley, T. M. and Schlesinger, M. E.: Analytical solution for the effect of increasing CO₂ on global mean temperature, *Nature*, 315, 649–652, 1985.

- Williams, R. G., Roussenov, V., Frölicher, T. L., and Goodwin, P.: Drivers of continued surface warming after cessation of carbon emissions, *Geophysical Research Letters*, 44, 10–633, 2017.
- 805 Winton, M., Takahashi, K., and Held, I. M.: Importance of ocean heat uptake efficacy to transient climate change, *Journal of Climate*, 23, 2333–2344, 2010.
- Winton, M., Adcroft, A., Griffies, S. M., Hallberg, R. W., Horowitz, L. W., and Stouffer, R. J.: Influence of ocean and atmosphere components on simulated climate sensitivities, *Journal of climate*, 26, 231–245, 2013.
- Xie, P. and Vallis, G. K.: The passive and active nature of ocean heat uptake in idealized climate change experiments, *Climate Dynamics*, 38, 810 667–684, 2012.
- Yool, A., Popova, E., and Anderson, T.: MEDUSA-2.0: an intermediate complexity biogeochemical model of the marine carbon cycle for climate change and ocean acidification studies, *Geoscientific Model Development*, 6, 1767–1811, 2013.
- Zanna, L., Khatiwala, S., Gregory, J. M., Ison, J., and Heimbach, P.: Global reconstruction of historical ocean heat storage and transport, *Proceedings of the National Academy of Sciences*, 116, 1126–1131, 2019.
- 815 Zickfeld, K., Eby, M., Weaver, A. J., Alexander, K., Crespin, E., Edwards, N. R., Eliseev, A. V., Feulner, G., Fichefet, T., Forest, C. E., et al.: Long-Term Climate Change Commitment and Reversibility: An EMIC Intercomparison., *Journal of Climate*, 26, 2013.
- Ziehn, T., Chamberlain, M., Law, R., Lenton, A., Bodman, R., Dix, M., Stevens, L., Wang, Y.-P., and Srbinovsky, J.: The Australian Earth System Model: ACCESS-ESM1.5, In Review: *Journal of Southern Hemisphere Earth Systems Science (JSHESS)*, 2020.