

The soil organic carbon stabilization potential of old and new wheat cultivars: a $^{13}\text{CO}_2$ labelling study

Marijn Van de Broek^{1*}, Shiva Ghiasi^{2*}, Charlotte Decock^{1,3}, Andreas Hund⁴, Samuel Abiven⁵, Cordula Friedli^{4,5}, Roland A. Werner², Johan Six¹

¹Sustainable Agroecosystems group, Department of Environmental Systems Science, Swiss Federal Institute of Technology, ETH Zürich, Zürich, Switzerland

²Grassland Sciences group, Department of Environmental Systems Science, Swiss Federal Institute of Technology, ETH Zürich, Zürich, Switzerland

³California State University, San Luis Obispo, CA, USA

⁴Group of Crop Science, Department of Environmental Systems Science, Swiss Federal Institute of Technology, ETH Zürich, Zürich, Switzerland

⁵Department of Geography, University of Zürich, Zürich, Switzerland

*Marijn Van de Broek and Shiva Ghiasi contributed equally to this study

Correspondence to: Marijn Van de Broek (Marijn.vandebroek@usys.ethz.ch)

Abstract. Over the past decades, average global wheat yields have increased by about 250 %, mainly due to the cultivation of high-yielding wheat cultivars. This selection process not only affected aboveground parts of plants, but in some cases also reduced root biomass, with potentially large consequences for the amount of organic carbon (OC) transferred to the soil. To study the effect of wheat breeding for high-yielding cultivars on subsoil OC dynamics, two old and two new wheat cultivars from the Swiss wheat breeding program were grown for one growing season in 1.5 m-deep lysimeters and pulse-labelled with $^{13}\text{CO}_2$, to quantify the amount of assimilated carbon that was transferred belowground and can potentially be stabilized in the soil. The results show that although the old wheat cultivars with higher root biomass transferred more assimilated carbon belowground compared to more recent cultivars, no significant differences in net rhizodeposition were found between the different cultivars. As a consequence, the long-term effect of wheat cultivar selection on SOC stocks will depend on the amount of root biomass that is stabilized in the soil. Our results suggest that the process of wheat selection for high-yielding cultivars resulted in lower amounts of belowground carbon translocation, with potentially important effects on SOC stocks. Further research is necessary to quantify the long-term importance of this effect.

1 Introduction

Soil management has a large influence on the size of the soil organic carbon (SOC) stock in managed arable soils. This is evident from the large decrease in SOC that is generally observed after soils under natural vegetation are converted to arable

land (Don et al., 2011; Guo and Gifford, 2002; Poeplau et al., 2011). As a consequence, the mineralization of SOC and the loss of forest caused by land use change has contributed about 30 % to the increase in atmospheric CO₂ concentration since the onset of the industrial revolution (Le Quéré et al., 2018). Current contributions of the agricultural sector to global warming have been estimated to be about 11 %, but are mostly in the form of N₂O and CH₄ and not anymore as CO₂ (Tubiello et al., 2015).

The rising awareness that there is potentially an opportunity to increase subsoil organic carbon (OC) stocks (Chen et al., 2018) has led to the proposal that agricultural soils can be a sink of atmospheric CO₂ by applying appropriate climate-smart agricultural practices (Chenu et al., 2018; Minasny et al., 2017; Paustian et al., 2016). Multiple management practices have been shown to increase the OC content of cultivated soils, including the application of organic amendments to soils (Sandén et al., 2018), increasing the amount of crop residues returned to the field (Lehtinen et al., 2014) and planting of cover crops (Kong and Six, 2010; Poeplau and Don, 2015)..

In addition, growing crops with deeper roots and/or higher root biomass has been put forward as a strategy to increase OC sequestration in arable soils (Kell, 2011), while deep rooting can also decrease the effect of drought in climates where deep soil water is available during the main cropping season (Wasson et al., 2012). However, a direct or marker-assisted selection for root traits is very rare in conventional breeding programs. Accordingly, we have very limited knowledge if and how breeders alter the root system and potentially affect belowground carbon cycling. One way to evaluate the effect of breeder's selection on root characteristics and subsoil carbon cycling is to compare old and new varieties of the same breeding programme. For the Swiss wheat breeding programmes, the selection process reduced the mass and depth of roots under well-watered conditions (Friedli et al., 2019), as has been found for other breeding programs (Aziz et al., 2017), but modern genotypes enhanced root allocation to deep soil layers under drought. However, this pattern has not been observed consistently (Cholick et al., 1977; Feil, 1992; Lupton et al., 1974). To the best of our knowledge, there is no information about the effect of breeding on changes in subsoil OC dynamics and root respiration.

One reason for the lack of quantitative data about the effects of rooting depth on SOC sequestration is related to difficulties in measuring the amount of carbon transferred from roots to the soil (gross rhizodeposition) and the proportion of carbon that is eventually stabilized there (net rhizodeposition), after a portion of gross rhizodeposits are lost from the soil through microbial mineralization or leaching. The fact that rhizodeposition occurs below the soil surface greatly prevents direct observations of this *'hidden half of the hidden half'* of the SOC cycle (Pausch and Kuzyakov, 2018). First of all, direct measurements of root exudation rates are hampered by the fact that rhizodeposits are used by rhizosphere microorganisms within a couple of hours after they are released, resulting in very low concentrations of root carbon exudates in the soil (Kuzyakov, 2006). Second, the release of carbon exudates by agricultural crops is not equally divided throughout the growing season, but mainly occurs in the first 1 – 2 months of the growing period and decreases sharply thereafter (Gregory and Atwell, 1991; Keith et al., 1986; Kuzyakov and Domanski, 2000; Pausch and Kuzyakov, 2018; Swinnen et al., 1994). Third, measurements of the effects of rhizodeposits on changes in SOC stocks are further complicated by the priming effect, i.e. their positive effect on the mineralization of native SOC (Fontaine et al., 2007; de Graaff et al., 2009).

To overcome these difficulties, rates of C rhizodeposition can be measured by labelling plants with $^{13}\text{CO}_2$ or $^{14}\text{CO}_2$ (Jones et al., 2009; Kuzyakov and Domanski, 2000) and subsequently tracing the amount of photosynthetically assimilated ^{13}C or ^{14}C label in the soil at the end of the growing season (Kong and Six, 2010, 2012). However, the continuous application of $^{13}\text{CO}_2$ or $^{14}\text{CO}_2$ during the course of an entire growing season to plants is often not feasible, as this requires the set-up of open-top chambers while continuously supplying the crops with the isotopic label, which comes at a high financial cost. Therefore, plants are commonly labelled at fixed time intervals during the growing season (repeated pulse-labelling). This results in reliable estimates of the partitioning of assimilated carbon to different plant compartments, as well as into the soil (Kong and Six, 2010; Kuzyakov and Domanski, 2000; Sun et al., 2018).

In addition, assessing the magnitude of the carbon transfer from roots to the soil is not straightforward, particularly under field conditions. While carbon inputs from crops to the soil are often derived from yield measurements (Keel et al., 2017; Kong et al., 2005; Taghizadeh-Toosi et al., 2016), these quantities are often poorly related to root biomass or the magnitude of root exudates (Hirte et al., 2018; Hu et al., 2018). A better understanding of the factors controlling the rates of carbon rhizodeposition by different agricultural crops is thus necessary to assess how different crops affect SOC cycling and to provide SOC models with reliable rates of carbon inputs to the soil.

The present study addresses the following research question: do wheat cultivars with shallow roots and lower root biomass lead to less net carbon rhizodeposition compared to wheat cultivars with deeper roots and higher root biomass? To address this question, four different bread wheat cultivars from a century of Swiss wheat breeding (Fossati and Brabant, 2003; Friedli et al., 2019) were grown in large mesocosms, which allowed to study the plant-soil system under controlled conditions that closely resemble a field situation. We hypothesized that wheat cultivars with shallow roots and lower root biomass would result in less net carbon rhizodeposition over the course of a growing season, compared to cultivars with deeper roots and higher root biomass.

2. Materials and Methods

2.1 Experimental set-up

2.1.1 ETH mesocosm platform

To assess the effect of wheat root characteristics on net rhizodeposition in a realistic soil environment under controlled conditions, an experiment was set up at the mesocosm platform of the Sustainable Agroecosystems Group at the Research Station for Plant Sciences Lindau (ETH Zürich, Switzerland). The platform was located inside a greenhouse and consisted of 12 cylindrical lysimeters with a diameter of 0.5 m and a height of 1.5 m, constructed using 10 mm wide polyethylene (Figure S1). The lysimeters were equipped with probes installed at five different depths (0.075, 0.30, 0.60, 0.90 and 1.20 m below the surface) to measure the volumetric moisture content at a temporal resolution of 30 min (ECH₂O EC-5, Decagon Devices, US) and to sample soil pore water (Prenart, Frederiksberg, Denmark) and soil pore air (Membrana, Wuppertal, Germany). The

lysimeters were filled with mechanically homogenized soil, collected from an agricultural field in Estavayer-Le-Lac, Switzerland. The upper 0.15 m of the lysimeters were filled with topsoil, collected from the A-horizon, while the remainder (0.15 – 1.35 m depth) was filled with subsoil. The bottom 0.15 m of the lysimeters (1.35 – 1.50 m depth) consisted of a layer of gravel (Blähton, Erik Schweizer, Switzerland), to facilitate drainage of soil water through the bottom of the lysimeters. The top and subsoil had a sandy clay loam texture with 21 % silt, 21 % clay, 58 % sand, and top- and subsoil pH values were 7.8 and 7.5, respectively. The OC concentration of the top- and subsoil was 0.77 ± 0.01 % and 0.40 ± 0.01 %, respectively, with a C:N ratio of 6.9 and 5.0, respectively. No carbonates were detected in the soil.

At the top of each lysimeter, pneumatically activated chambers were placed, that were automatically closed when applying the $^{13}\text{CO}_2$ label (see section 2.1.3). These chambers were made of stainless steel with fitted Plexiglas panes and covered a rectangular area of 0.5 x 0.5 m with an initial height of 0.1 m. Chamber heights were extended with increasing plant height, using one or two height extensions of 0.5 m each (Figure S1).

2.1.2 Wheat cultivars and growth conditions

Four wheat (*Triticum aestivum* L.) cultivars from the Swiss wheat breeding program (Fossati and Brabant, 2003; Friedli et al., 2019) with different breeding ages were selected: Mont-Calme 268 (introduced in 1926), Probus (1948), Zinal (2003) and CH Claro (2007). Generally, more recent cultivars of this program on average have more shallow roots and lower root biomass under well-watered conditions compared to the older cultivars (Friedli et al., 2019). CH Claro was selected as a modern variety with relatively deep rooting.

Before the wheat plants were transplanted to the lysimeters, wheat seeds were germinated in a greenhouse for 2 – 3 days on perforated anti-algae foil laid over 2-mm moistened fleece at a warm temperature (20 °C during day and 18 °C during night) and good light conditions. Next, the seedlings were planted in containers filled with the same topsoil used to fill the lysimeters and transferred to a climate chamber for vernalization for 52 days (Baloch et al., 2003). First, the seedlings were kept 45 days at 4 °C, with 8 hours of light per day and a light intensity of 10 kilolux. During the 3 subsequent days, daylight intensity was increased to 36 kilolux, daytime temperature was increased to 12 °C and night temperature to 10 °C. During the last 4 days, daytime temperature was increased to 16 °C, and night temperature to 12 °C. The relative humidity was maintained at 60 ± 10 % during the entire vernalization period. After vernalization, 70 seedlings were transplanted to every lysimeter, corresponding to a plant density of 387 plants m^{-2} . At the timing of transplanting, the plants were at the onset of tillering.

The experimental set-up consisted of a randomized complete block design. Each of the four wheat cultivars was planted in three lysimeters, i.e. 3 replicates per cultivar, resulting in a total of 12 lysimeters. These were placed in 3 blocks of 4 rows, where each wheat cultivar was planted in one lysimeter in each block. The plants were grown in the greenhouse for about 5 months, between 24 August 2015 and 1 February 2016. Despite uneven maturing of plants within and between the lysimeters, all plants had reached flowering stage at the time of harvest. Fertilizer was applied to the soil lysimeters a first time on 5 October 2015, at a rate of 84 kg N ha^{-1} , 36 kg P_2O_5 ha^{-1} , 48 kg K_2O ha^{-1} and 9 kg Mg ha^{-1} , and a second time on 4 December 2015, at a rate of 56 kg N ha^{-1} , 24 kg P_2O_5 ha^{-1} , 32 kg K_2O ha^{-1} and 6 kg Mg ha^{-1} . The lysimeters were watered manually twice

per week with a similar amount of water, to keep soil moisture close to field capacity. Differences in the amount of water used by the different cultivars resulted in differences in the soil water content between the cultivars (Figure S2). The temperature in the greenhouse was set to 20 °C during the day and 15 °C during the night. During the experiment, the average temperature in the greenhouse was 16.9 °C, with a minimum and maximum of 9.3 °C and 29.8 °C respectively. The average humidity was 63.7 %, with a minimum and maximum of 35.3 % and 86.4% respectively.

2.1.3 Repeated ^{13}C pulse-labelling

In order to study carbon allocation within the atmosphere-plant-soil system, a ^{13}C pulse-labelling approach was used. 99% $^{13}\text{CO}_2$ (Euriso-top, Saint-Aubin, France) was applied once per week (2 pm on Thursdays) by injecting 15, 56 or 98 mL CO_2 into each chamber depending on the chamber extension used, in order to yield a target $^{13}\text{CO}_2$ content of 58 atom%. A weekly labelling frequency has been shown to ensure a sufficient abundance of root-derived ^{13}C in the soil at the end of the experiment (Bromand et al., 2001; Kong and Six, 2010). After chamber closure, CO_2 concentration in one chamber was monitored using a CO_2 analyzer (Li-820, LICOR, Lincoln, US). Throughout the experiment, CO_2 concentrations were measured in the same chamber. After the CO_2 concentration dropped below 200 ppm, another $^{13}\text{CO}_2$ pulse was injected to yield a post-label CO_2 concentration of 570 ppm in the chamber headspace. The chamber lids were kept closed for two hours after label injection to achieve sufficient uptake and then re-opened to avoid condensation. On the same day of pulse-labelling, all chambers were closed overnight to recuperate ^{13}C lost through night respiration and allowed to be taken up by the plants in the morning before reopening the chambers.

2.2 Measurements

2.2.1 Belowground CO_2 concentration and $\delta^{13}\text{CO}_2$

Soil gas sampling was performed once per week (Wednesdays) by attaching a pre-evacuated 110 mL crimp serum vial to a sampling port at each depth, leaving it equilibrating overnight. For each sample, a 20 mL subsample was transferred to a pre-evacuated Labco exetainer (12 mL), and used to determine the CO_2 concentration. The CO_2 concentration of each sample was determined using a gas chromatograph equipped with a thermal conductivity detector (Bruker 456-GC, Germany). In addition, the $\delta^{13}\text{C}$ value of CO_2 was measured for CO_2 samples collected along the depth profiles on the last sampling date, using a Gasbench II modified as described by Zeeman et al. (2008) coupled to a Delta^{plus}XP isotope ratio mass spectrometer (IRMS, ThermoFisher, Germany). The standard deviation of the measurements was < 0.15 ‰.

2.2.2 Sampling and general soil analyses

At the end of the experiment, the aboveground biomass of the wheat plants was harvested separately for each lysimeter and separated into leaves, ears and stems. Soil from the lysimeters was collected by destructive sampling to analyze bulk density, root biomass and other soil properties. The sampling was done layer by layer. After a soil layer had been sampled, it was

removed completely from the lysimeter and the next layer was sampled. From each depth increment (0 – 0.15, 0.15 – 0.45, 0.45 – 0.75, 0.75 – 1.05, 1.05 – 1.35 m depth), five soil cores were collected per lysimeter using a soil core sampler (5.08 cm diameter, Giddings Machine Company Inc., Windsor, CO, US). Three of the five cores per lysimeter and depth increment were used for the determination of root biomass based on a combination of buoyancy and sieving through a 530 µm sieve, using a custom-built root washing station. The remaining two soil cores were sieved at 8 mm, air-dried and stored for further analysis. Prior to air drying, the fresh weight and volume for each core was determined, and a subsample was taken for the determination of gravimetric soil moisture content. Bulk density was calculated based on fresh weight, gravimetric moisture content, and core volume. Soil texture was measured using a particle size analyzer (LS 13 320, Beckman Coulter, Indianapolis, USA). Prior to analysis 0.1g of soil was shaken for 4 h with 4 ml of 10 % Na-hexametaphosphate and sonicated for 1 min.

2.2.3 Soil microbial biomass

Soil microbial biomass was extracted from soil samples that had been frozen at -20 °C for 6 months immediately after sampling. Two subsamples of 40 g were taken from each sieved soil sample. One set was fumigated for 24 hours using chloroform. Next, total dissolved OC was extracted from each fumigated and non-fumigated subsample by shaking it in 200 mL 0.05 M K₂SO₄ for one hour, prior to filtering through a Whatman 42 filter paper. Total OC concentrations in K₂SO₄ extracts were determined using a CN analyzer (multi N/C 2100 S analyser, Analytik Jena, Germany). To determine microbial biomass carbon per unit of dry soil, the gravimetric soil water content was determined by drying about 10 g of each soil sample at 105 °C and subtracting the weights before and after drying. The carbon content of the soil microbial biomass was calculated according to Vance et al. (1987) as:

$$TOC_{MB} = \frac{TOC_F - TOC_{NF}}{0.45} \quad (\text{Eq. 1})$$

Where TOC_F and TOC_{NF} are the total OC in fumigated and non-fumigated samples, respectively. The remainder of the filtered samples was freeze dried in order to analyze the δ¹³C value. The δ¹³C value of soil microbial biomass was calculated using mass balance according to Ruehr et al. (2009):

$$\delta^{13}C_{MB} = \frac{(\delta^{13}C_F \cdot C_F - \delta^{13}C_{NF} \cdot C_{NF})}{C_F - C_{NF}} \quad (\text{Eq. 2})$$

Where C_F and C_{NF} represent total carbon content of the fumigated and non-fumigated samples, respectively.

2.2.4 Organic carbon concentration and isotopic composition of plant material, soil organic carbon and microbes

The OC concentration and isotopic composition (δ¹³C) of above- and belowground plant material, fumigated and non-fumigated soil (for the determination of microbial biomass C and δ¹³C) and bulk soil were measured by weighing 2, 4, 80, 80

and 100 mg, respectively, of each sample into Sn capsules (9 x 5 mm, Saentis, CH) for analysis with a Flash EA 1112 Series elemental analyzer (ThermoFisher, Germany) coupled to a Delta^{plus} XP IRMS via a ConFlo III (Brooks et al., 2003; Werner et al., 1999; Werner and Brand, 2001). The measurement precision (SD) of the quality control standards (tyrosine Tyr-Z1, caffeine Caf-Z1), was 0.37 (‰) for above and belowground plant material, microbes and the soil samples.

2.3 Data processing

2.3.1 Excess ¹³C calculations

The mass of ¹³C label that was recovered in (i) the aboveground vegetation, (ii) roots of wheat plants and (iii) the soil was calculated following Studer et al. (2014):

$$m^E(^{13}\text{C}) = \frac{\chi^E(^{13}\text{C}) \cdot m(\text{C}) \cdot M(^{13}\text{C})}{\chi(^{12}\text{C}) \cdot M(^{12}\text{C}) + \chi(^{13}\text{C}) \cdot M(^{13}\text{C})} \quad (\text{Eq. 3})$$

Where $m^E(^{13}\text{C})$ is the mass of recovered ¹³C label (g m⁻²), $\chi^E(^{13}\text{C})$ is the excess atom fraction (unitless, calculated following Coplen (2011)), $m(\text{C})$ is the total mass (g m⁻²) of C, $M(^{12}\text{C})$ and $M(^{13}\text{C})$ are the molar weight of ¹²C and ¹³C (g mol⁻¹), respectively, and $\chi(^{12}\text{C})$ and $\chi(^{13}\text{C})$ are the ¹²C and ¹³C atom fraction (unitless), respectively.

To calculate the excess atom fraction ($\chi^E(^{13}\text{C})$) of the soil compartment, the isotopic composition of the soil at the start of the experiment was used as the reference value (-26.45 ± 0.04 ‰ for the topsoil, -25.01 ± 0.13 ‰ for the subsoil). As all lysimeters were labelled with ¹³CO₂, no control treatment for the wheat plants was present. Therefore, a δ¹³C reference value of -28 ‰ was assumed for the aboveground parts and roots of all wheat plants. The calculation of excess ¹³C is very sensitive to variability in input parameter values, including the δ¹³C value of plant biomass and soil. Therefore, a sensitivity analysis was used to show that varying the initial δ¹³C value of the wheat plants with +/- 3 ‰, a typical range over which δ¹³C values can vary in the field because of e.g. precipitation (Kohn, 2010), led to changes in calculated $m^E(^{13}\text{C})$ in the order of +/- 1 % for aboveground biomass and +/- 1 – 5 % for belowground biomass. The effect of the initial δ¹³C value of the biomass on the calculated amount of recovered ¹³C label in the wheat plants was thus limited. Calculations of the effect of wheat cultivar on belowground excess ¹³C were only performed for the upper 0.45 m of the lysimeters, as missing data for deeper soil layers prevented including these layers in the statistical analyses.

2.3.2 Net carbon rhizodeposition

The absolute amount of carbon rhizodeposition for the different depth segments in the lysimeters was calculated following Janzen and Bruinsma (1989):

$$Rhizodeposition\ C = \frac{\chi^E(^{13}C)_{soil}}{\chi^E(^{13}C)_{root}} \cdot C_{soil} \quad (\text{Eq. 4})$$

Where rhizodeposition C is expressed in g kg⁻¹ for the considered layer, $\chi^E(^{13}C)_{soil}$ and $\chi^E(^{13}C)_{root}$ are the excess ¹³C atom fraction in the soil and roots respectively, calculated as described in section 2.3.1, and C_{soil} is the OC concentration of the considered soil layer (g kg⁻¹). This approach assumes that the isotopic enrichment of rhizodeposits and roots are equal. The absolute amount of carbon rhizodeposition for each soil layer was calculated by multiplying rhizodeposition C (g kg⁻¹) with the carbon content (kg) present in each of the respective layers. We note that data on the C concentration and $\delta^{13}C$ value of root biomass could not be obtained from a number of soil layer below 0.45 m depth for certain cultivars, due to the limited root biomass that could be retrieved. Therefore, net C rhizodeposition was only calculated for the two uppermost soil layers (0 – 0.45 m depth), as only for these layers all necessary data to calculate net C rhizodeposition was available for the three replicates of every cultivar.

2.3.3 Subsoil CO₂ production

Depth profiles of subsoil CO₂ production in the lysimeters were calculated using the weekly measured depth profiles of CO₂ concentration throughout the experiment. To assess the variability among the different lysimeters, these calculations were performed separately for every lysimeter and average CO₂ production depth profiles were calculated for each cultivar. Measurements of CO₂ concentration, soil temperature and soil moisture content were performed at discrete depths (0.075, 0.30, 0.60, 0.90 and 1.20 m depth). Continuous depth profiles of these variables at a vertical resolution of 0.05 m were obtained using linear interpolation. Depth profiles of CO₂ production were calculated using the discretized form of the mass balance equation of CO₂ in a diffusive one-dimensional medium, following Goffin et al. (2014):

$$P(z)_i = \frac{\Delta(\varepsilon_i[CO_2]_i)}{\Delta t} + \frac{F_{top_i} - F_{bot_i}}{\Delta z} \quad (\text{Eq. 5})$$

Where $P(z)$ is the CO₂ production in layer i (μmol CO₂ m⁻³ s⁻¹) over timespan Δt , t is the time (s), ε_i is the air-filled porosity in layer i (m³ m⁻³), $[CO_2]_i$ is the CO₂ concentration of layer i (μmol CO₂ m⁻³), F_{top_i} and F_{bot_i} are the CO₂ fluxes transported through the upper and lower boundaries of layer i (μmol CO₂ m⁻² s⁻¹) during timespan Δt , respectively, and z is the depth (m). The vertical CO₂ fluxes are calculated as (Goffin et al., 2014):

$$F_{top_i} = -\bar{D}_{s;i-1,i} \frac{[CO_2]_{i-1} - [CO_2]_i}{\Delta z} \quad (\text{Eq. 6})$$

$$F_{bot_i} = -\bar{D}_{s;i,i+1} \frac{[CO_2]_i - [CO_2]_{i+1}}{\Delta z} \quad (\text{Eq. 7})$$

Where $\bar{D}_{s,i,j}$ is the harmonic average of the effective diffusivity coefficient (D_s) between layers i and j , and Δz is the layer thickness. The effective diffusivity coefficient is calculated using a formula appropriate for repacked soils (Moldrup et al., 2000):

$$D_{s,i} = D_{0,t} \frac{\varepsilon_{i,t}^{2.5}}{\Phi_i} \quad (\text{Eq. 8})$$

Where D_0 is the gas diffusion coefficient of CO_2 in free air over timespan Δt ($\text{m}^2 \text{s}^{-1}$), ε_i is the air-filled porosity of layer i over timespan Δt ($\text{m}^3 \text{m}^{-3}$) and Φ_i is the total soil porosity of layer i ($\text{m}^3 \text{m}^{-3}$). The total soil porosity was calculated as $\Phi_i = 1 - \rho_i/\rho_p$ where ρ_i is the soil bulk density (ton m^{-3}) and ρ_p is the particle density (2.65 ton m^{-3}). Due to the large vertical variability in measured bulk density depth profiles, a constant bulk density profile was assumed for the subsoil (below 0.15 m depth), calculated as the average of the measured bulk density values for these layers. The air-filled porosity over timespan Δt was calculated as the difference between the total porosity ($\text{m}^3 \text{m}^{-3}$) and the average measured water-filled pore space over timespan Δt ($\text{m}^3 \text{m}^{-3}$). The latter was measured throughout the experiment (see section 2.1.1) and corrected based on differences between these measurements at the end of the experiment and the measured volumetric water content of the sampled soil at the end of the experiment. For this purpose, different correction equations were used for (i) the upper soil layer (0 – 15 cm) and (ii) all deeper layers combined.

The gas diffusion coefficient in free air was corrected for the individual lysimeters for variations in temperature and soil moisture throughout the experiment (Massman, 1998), as:

$$D_0 = D_{0,stp} \frac{p_0}{p} \left(\frac{T}{T_0} \right)^\alpha \quad (\text{Eq. 9})$$

Where $D_{0,stp}$ is the gas diffusion coefficient for CO_2 in free air under standard temperature (0°C) and pressure (1 atm) ($1.385 \cdot 10^{-5} \text{ m}^2 \text{s}^{-1}$ (Massman, 1998)) and α is a coefficient (1.81; Massman, 1998)). Semi-continuous measurements of soil temperature in every lysimeter were used to calculate D_0 values throughout the experiment, while a constant atmospheric pressure of 1 atm throughout the experiment was assumed.

To obtain depth profiles of the total amount of CO_2 produced by the different wheat cultivars during the experiment (expressed as $\text{g CO}_2 \text{m}^{-2}$), the calculated CO_2 production rates between all measurement days ($P(z)$) were summed for the timespan of the experiment and converted to $\text{g CO}_2 \text{m}^{-2}$ using the molecular mass of CO_2 (44.01 g mol^{-1}). We applied the boundary condition of the absence of a flux of CO_2 at the bottom of the lysimeters. It is noted that these calculations do not make a distinction between the source of CO_2 of production, thereby combining both autotrophic and heterotrophic CO_2 production (total soil respiration). For more information about these methods, reference is made to Goffin et al. (2014).

2.4 Statistics

To account for the three blocks in the randomized complete block design, statistically significant differences between aboveground characteristics of different cultivars were checked using a two-way analysis of variance (anova) without interactions (Dean et al., 2015), followed by a Tukey's test, based on the values obtained for the individual replicates ($n = 3$ for every cultivar) using a significance level of 0.05. This was done after checking for homogeneity of variance (Levene's test) and normality (Shapiro-Wilk test) using a confidence level of 0.05. These analyses were performed in Matlab®. The effect of cultivar and depth on soil bulk density, belowground biomass, belowground C allocation and net C rhizodeposition was assessed using a linear mixed-effects model, with cultivar and depth being fixed effects and blocks being treated as a random effect (*lmer* function in R (R Core Team (2019))). Next, a pair-wise comparison was used to check for statistically significant differences between the cultivars (*emmeans* package in R). Belowground biomass was log-transformed to increase normality and homogeneity of variances for the latter analysis. Uncertainties on reported variables are expressed as standard errors ($n = 3$).

3. Results

3.1 Aboveground biomass

The aboveground biomass produced at the end of the experiment was significantly different between Zinal and Probus, while the aboveground biomass of CH Claro and Mont-Calme 268 was not significantly different from any other cultivar (Figure 1, Table 1). The biomass of the ears was significantly higher for Zinal, compared to CH Claro, Probus and Mont-Calme (Figure 1, Table S1). It is noted that these data should be interpreted with care, since not all plants reached maturity at the time of harvest, and is potentially not representative for the biomass of the ears of full-grown plants. No significant differences were found between the $\delta^{13}\text{C}$ values of aboveground biomass of the different cultivars (Figure 2). The high $\delta^{13}\text{C}$ values of the aboveground biomass of all wheat cultivars (266 ‰ on average) showed that a substantial amount of the $^{13}\text{CO}_2$ tracer was incorporated by all wheat plants (Figure 2).

3.2 Belowground biomass

The average root biomass was highest in the topsoil and significantly lower in the subsoil layers of all four wheat cultivars (Figure 1B). Root biomass of Zinal was significantly lower compared to the root biomass of Probus and Mont-Calme 268, while the root biomass of CH Claro was not significantly different from any of the other cultivars (Figure 1B). These differences were mostly present in the two uppermost soil layers, while root biomass was not significantly different between different cultivars at any depth, except for Zinal and Mont-Calme 268 between 0.45 – 0.75 m depth (Figure 1). The root:shoot ratio varied between 0.10 ± 0.02 and 0.19 ± 0.08 , and was not significantly different between the different cultivars (Table 1).

The depth profiles of the $\delta^{13}\text{C}$ of root biomass were different between the old and more recent wheat cultivars (Figure 2). In the two uppermost soil layers, no significant differences were detected between the $\delta^{13}\text{C}$ values of root biomass of the different cultivars. These differences could not be checked for statistical significance in deeper soil layers due to a lack of sufficient recovered root biomass in each lysimeter. The $\delta^{13}\text{C}$ values of the roots of the old wheat cultivars showed only limited variation with depth, with values between ca. 150 and 200 ‰. In contrast, the $\delta^{13}\text{C}$ values of the roots of the more recent wheat cultivars were highest in the two uppermost soil layers (0 – 45 cm) and showed an abrupt decrease with depth in deeper soil layers.

3.3 Soil and soil organic carbon characteristics

The SOC concentration in the lysimeters was similar to the OC concentration of the initial soil (Figure 3A). A direct comparison between the SOC concentration before and after the experiment could not be made, as no measurements of the OC concentration of the soil in the lysimeters before the start of the experiment could be made. However, the SOC concentration measured at the different depths in the lysimeters was similar to the OC concentration measured on the soil that was used to fill the lysimeters (Figure 3A). No statistically significant differences in SOC concentration were found between the different cultivars at any depth.

The SOC in the two uppermost soil layers (0 – 45 cm) of all wheat cultivars was enriched in ^{13}C compared to the soil that was used to fill the lysimeters (Figure 3B). Although the $\delta^{13}\text{C}$ value of SOC was not significantly different at any depth between any of the cultivars, the largest increase in the $\delta^{13}\text{C}$ value of topsoil OC was observed for Probus and Mont-Calme 268 (Figure 3B), indicating that the soil under the old cultivars incorporated more of the ^{13}C label, compared to the more recent cultivars. The limited difference between (i) the $\delta^{13}\text{C}$ values of the soil used to fill the lysimeters and (ii) the measurements at the end of the experiment below a depth of 0.45 m, indicates a lower amount of incorporated ^{13}C label in the subsoil. Similarly, the $\delta^{13}\text{C}$ value of topsoil microbial biomass was more positive compared to deeper soil layers for all cultivars, indicating that microbes utilized more substrate enriched in ^{13}C in the two uppermost soil layers, compared to deeper soil layers (Figure 3C). Statistically significant differences were only detected in the layer between 0.15 and 0.45 m depth, where the $\delta^{13}\text{C}$ value of microbial biomass under Zinal was significantly lower compared to Mont-Calme 268. However, as the microbial biomass under Zinal was substantially higher compared to under Mont-Calme 268 in this layer (Figure S3), this does not necessarily imply that microbes under Mont-Calme 268 incorporated more excess ^{13}C compared to under Zinal. Depth profiles of microbial biomass carbon were relatively constant (200 – 500 $\mu\text{g C g soil}^{-1}$) with no consistent differences between different cultivars (Figure S3). The $\delta^{13}\text{C}$ values of soil CO_2 ($\delta^{13}\text{CO}_2$) at the end of the experiment were similar for all wheat cultivars for the two uppermost layers (0 – 0.45 m) (Figure 3D). Deeper down the profile, the CO_2 under the old wheat cultivars was more enriched in ^{13}C compared to the more recent cultivars, by an average of ca. 30 ‰. The only statistically significant differences were detected in the lowermost layer, where the $\delta^{13}\text{C}$ value of CO_2 of Zinal and CH Claro were significantly lower compared to Mont-Calme 268.

There was no significant effect of cultivar on the bulk density of the soil at the end of the experiment ($F_{3,59} = 1.9$, $p = 0.23$), while there was a significant effect of depth on bulk density ($F_{4,59} = 19.4$, $p < 0.0005$). The average bulk density of all lysimeters

was highest in the topsoil ($1.67 \pm 0.12 \text{ Mg m}^{-3}$) and varied with depth (Figure S4A). The gravimetric moisture content in the
345 lysimeters at the end of the experiment increased with depth for all cultivars, from ca 0.1 g g^{-1} in the top layer to ca. 0.15 g g^{-1} in the bottom layer (Figure S4B), and was only significantly different between Mont-Calme 268 and Zinal in the uppermost soil layer. The soil moisture content changed relatively little throughout the experiment for all lysimeters, after an initial phase of decreasing soil moisture content at the onset of the experiment (Figure S2).

3.4 Excess ^{13}C and carbon rhizodeposition

350 The total amount of ^{13}C label that was present in the plant-soil system at the end of the experiment, expressed as excess ^{13}C , differed significantly between different wheat cultivars (Figure 4A). When accounting for excess ^{13}C in aboveground biomass and in the soil and roots down to a depth of 0.45 m, the lowest amount of ^{13}C label was found in the Zinal lysimeters ($1.19 \pm 0.11 \text{ g m}^{-2}$), followed by CH Claro ($1.64 \pm 0.06 \text{ g m}^{-2}$) and the older wheat cultivars ($2.05 \pm 0.09 \text{ g m}^{-2}$ for Mont-Calme 268 and $2.01 \pm 0.19 \text{ g m}^{-2}$ for Probus), with the majority of ^{13}C tracer in the above-ground biomass (Figure 4A). Despite these
355 differences, the relative distribution of the assimilated ^{13}C between aboveground biomass, roots and soil was similar between the different wheat cultivars (Figure 4B). On average, $80.7 \pm 1.7 \%$ of the assimilated tracer ended up in aboveground biomass, $8.4 \pm 1.5 \%$ in root biomass and $10.9 \pm 1.4 \%$ in the soil. It is noted that root-respired ^{13}C label is not included in this analysis, which may lead to an underestimation of the fraction of ^{13}C label that was allocated belowground.

The total amount of net carbon rhizodeposition measured at the end of the experiment down to 0.45 m decreased with depth
360 for all wheat cultivars (Figure 4C), with this difference only being statistically significant for Zinal. The highest amount of net carbon rhizodeposition was observed for Probus ($108 \pm 34 \text{ g C m}^{-2}$), followed by CH Claro ($97 \pm 24 \text{ g C m}^{-2}$), Mont-Calme ($83 \pm 29 \text{ g C m}^{-2}$) and Zinal ($62 \pm 11 \text{ g C m}^{-2}$). There was thus no clear relationship between the amount of net carbon rhizodeposition and year of release of the wheat cultivars.

3.5 CO_2 concentration and production

365 Throughout the experiment, the change in the CO_2 concentration of the two uppermost soil layers was limited, with average values for the topsoil between 470 and 761 ppm for all cultivars (Figure 5). Deeper down the lysimeters, relatively constant CO_2 concentrations were observed during the first 3 weeks of the experiment, ca. 5.000 – 10.000 ppm. After 3 weeks, subsoil CO_2 concentrations abruptly increased and remained high throughout the experiment. These were substantially larger for the older cultivars (with maximum values of ca. 30.000 ppm) compared to the younger cultivars (with maximum values ca. 24.000
370 ppm).

Despite these high CO_2 concentrations in the subsoil, CO_2 production was mainly taking place in the topsoil, with the highest rates of CO_2 production between 0.10 and 0.20 m depth for all cultivars (Figure 6). For the young cultivars (Zinal and CH Claro), 95 % of CO_2 was produced above a depth of 0.3 m. In contrast, in older cultivars (Probus and Mont-Calme 268) 95 % of CO_2 was produced above a depth of 0.55 and 0.6 m, respectively. Despite these observations, neither the calculated total

375 amount of subsoil CO₂ production or the depth above which 95 % of CO₂ was produced were significantly different between
any of the cultivars.

4. Discussion

The aim of the present study was to assess differences in belowground carbon transfer and net rhizodeposition by wheat
cultivars with different root biomass and rooting depth. Our results show that although there are marked differences in both
380 the amount of carbon transferred belowground and the timing of belowground carbon transfer, there is no clear relationship
between root characteristics and the amount of net rhizodeposition. Therefore, the fate of root biomass might determine the
total amount of subsoil carbon stabilization in the long-term.

4.1 Plant biomass carbon dynamics and CO₂ production

No consistent differences in total aboveground biomass between old and new wheat cultivars were observed. The aboveground
385 biomass values were at the high end of reported values for wheat plants in the field (Mathew et al., 2017), while the lack of
consistent differences in the biomass of wheat cultivars released over a time span of multiple decades has generally been
observed (Brancourt-Hulmel et al., 2003; Feil, 1992; Lupton et al., 1974; Wacker et al., 2002).

The fraction of biomass in the grain-bearing ears was, however, much larger for the modern wheat cultivars (on average 9 and
47 % of total aboveground biomass for CH Claro and Zinal respectively) compared to the old wheat cultivars (on average 1
390 and 2 % for Mont-Calme 268 and Probus respectively). While an increase in the fraction of biomass allocated to grains is
generally observed in old versus modern wheat cultivars (Brancourt-Hulmel et al., 2003; Feil, 1992; Shearman et al., 2005),
mostly as a consequence of the introduction of reduced height genes (Tester and Langridge, 2010), the harvest index reported
here for the old cultivars might have been underestimated because older cultivars were not yet fully mature at plant harvest.
The total root biomass of the older wheat cultivars was substantially larger compared to the more recent cultivars, although
395 these differences were not consistently significant between all modern and old varieties (Table 1). These differences were
mostly apparent in the top 0.45 m of the lysimeters (Figure 1). It is not clear if the lack of statistically significant differences
in the root biomass within the deeper soil layers was due to (i) inability to collect all fine roots from the soil or (ii) actual
differences in root biomass. These results are in line with a recent study on the biomass of roots of different wheat cultivars of
the Swiss wheat breeding program, including the cultivars used in our experiment (Friedli et al., 2019). This study showed that
400 under well-watered conditions, older wheat cultivars had a substantially higher root biomass compared to the more recently
released wheat cultivars. Similar results have been obtained for wheat cultivars released in e.g. Australia (Aziz et al., 2017)
and other countries around the world (Waines and Ehdaie, 2007). The root:shoot ratio of the wheat cultivars in our study ($0.10 \pm 0.2 - 0.19 \pm 0.08$, Table 1) were at the low end of reported values for wheat plants globally (Mathew et al., 2017), but in line
with reported values for wheat cultivars of the Swiss wheat breeding program, including the cultivars used in our study (an
405 average value of 0.14 for all cultivars studied by Friedli et al. (2019)).

The maximum rooting depth was similar between the old and recent wheat cultivars (Figure 1B). This is in contrast with the results from Friedli et al. (2019), who found that the older wheat cultivars had deeper roots (the depth above which 95 % of roots were found (D_{95}) was on average 101 cm) compared to the more recent cultivars included in the present study (average D_{95} of 85 cm). These differences might partly arise from the different set-up used in both studies. Both experiments were
410 carried out in a controlled greenhouse environment, but Friedli et al. (2019) used soil columns with a diameter of 0.11 m, while in our study lysimeters with a diameter of 0.5 m were used. Additional information about subsoil root dynamics could be obtained from the measured depth profiles of the CO_2 concentration and $^{13}\text{CO}_2$, with the latter only being measured in the last phase of the experiment. The calculated depth profiles of CO_2 production showed that CO_2 was being produced down to greater depths under the old wheat cultivars (Figure 6). Combined with the higher $\delta^{13}\text{C}$ values of subsoil CO_2 of the lysimeters under
415 the old wheat cultivars at the end of the experiment (Figure 3D), this suggests that the roots of the old wheat cultivars respired CO_2 at greater depths compared to the recent wheat cultivars.

The $\delta^{13}\text{C}$ values of root biomass suggest that the temporal root carbon dynamics of the old and recent wheat cultivars differed substantially (Figure 2B). The root biomass of the old wheat cultivars had a high $\delta^{13}\text{C}$ value at all measured depths, indicating that the $^{13}\text{CO}_2$ label was allocated to the roots at all depths throughout the experiment. In contrast, the root biomass of the
420 recent wheat cultivars was greatly enriched in ^{13}C in the top 0.45 m, while deeper roots were much less enriched in ^{13}C . This suggests that both old and more recent wheat cultivars grew roots down to depths of > 1 m in the beginning of the experiment (when the total amount of ^{13}C assimilated by the plants was limited), while only the old cultivars kept on allocating carbon down to deep roots (> 0.45 m) throughout the experiment (thus having assimilated more ^{13}C over the period of root growth compared to the more recent cultivars). The similar $\delta^{13}\text{C}$ value of the aboveground biomass of all wheat cultivars (Figure 2A)
425 suggests that the differences in $\delta^{13}\text{C}$ values of the root biomass are unlikely to be caused by differences in the relative amount of $^{13}\text{CO}_2$ assimilated by the plants, relative to unlabeled CO_2 . Thus, these results suggest that old wheat cultivars allocate photosynthates down to their roots throughout a substantial part of the plant growth phase, while this is not the case for more recent cultivars.

4.2 Carbon allocation by wheat plants

430 The partitioning of the ^{13}C label was very similar between the different wheat cultivars (Figure 4B). It is noted that the amount of rhizosphere-respired $^{13}\text{CO}_2$ could not be included in these calculations, although this typically accounts for ca. 7 - 14 % of assimilated carbon in crops, or 40 % of total belowground C allocation (Kuzyakov and Domanski, 2000; Pausch and Kuzyakov, 2018). The fraction of assimilated carbon that is transferred belowground reported here is therefore underestimated. The belowground transfer of ca. 20 % of assimilated C for all cultivars is in line with previous studies, which have reported fractions
435 of similar magnitude for wheat plants, when not accounting for rhizosphere CO_2 respiration: 18 – 25 % (Hirte et al., 2018), 18 % (as reviewed by Kuzyakov and Domanski (2000)), 15 % (Keith et al., 1986), 17 % (Gregory and Atwell, 1991) and 31 % (Sun et al., 2018). In contrast, reported values of the partitioning of belowground translocated carbon by wheat plants to (i) roots and/or (ii) net rhizodeposition are much more variable, with reported net rhizodeposition carbon as a percentage of total

belowground carbon (root carbon and net rhizodeposition carbon combined) for wheat plants between 23 % (as summarized by Kuzyakov and Domanski (2000)) and 72 % (Sun et al., 2018). The results obtained here (68 %) are thus at the high end of reported values. However, they were similar to results from a field study in Switzerland which used two modern Swiss wheat cultivars, among which CH Claro (58 %; Hirte et al. (2018)).

4.3 Rates of net carbon rhizodeposition

The total amount of carbon assimilated by the wheat cultivars that was transferred to roots and soil in the top 0.45 m at the end of the experiment ranged between $97 \pm 14 \text{ g m}^{-2}$ (Zinal) and $164 \pm 38 \text{ g m}^{-2}$ (Probus) (Table 2). It is noted that the total amount of belowground carbon translocation by the wheat plants is underestimated, as rhizosphere respiration could not be included in our calculations. These numbers are in the range of reported values for wheat plants of $94 - 295 \text{ g m}^{-2}$ (as summarized by Keith et al. (1986)) and the value reported by Kuzyakov and Domanski (2000) (150 g m^{-2}), as well as the reported amount for two recent wheat cultivars of the Swiss wheat breeding program (including CH Claro) of $110 - 134 \text{ g m}^{-2}$ (Hirte et al., 2018). In contrast to the total amount of carbon translocated belowground, the amount of net carbon rhizodeposition was not consistently different between the old and more recent wheat cultivars ($62 \pm 11 - 108 \pm 34 \text{ g m}^{-2}$) (Figure 4, Table 2). These values are higher compared to values calculated by Pausch and Kuzyakov (2018) ($18 - 34 \text{ g m}^{-2}$, depth unknown) and Hirte et al. (2018) ($63 - 73 \text{ g m}^{-2}$; down to 0.75 m depth).

A large uncertainty associated with calculated values of subsoil carbon sequestration using isotopic labelling approaches is related to the assumption that the isotopic enrichment of roots and rhizodeposits are similar (eq. 4). This simplification is made because of the difficulties in measuring quantitative characteristics of rhizodeposits in a soil medium (Oburger and Jones, 2018), but leads to erroneous calculations of the amount of carbon rhizodeposition when this assumption is violated (Stevenel et al., 2019). To assess the uncertainty of calculated values of subsoil carbon sequestration, we calculated how these values differ when the value of root $\delta^{13}\text{C}$ is varied with $\pm 25 \%$ (Figure S5). This results in calculated values of total carbon rhizodeposition, down to a depth of 0.45 m, of $69 - 105 \text{ g m}^{-2}$ for Mont-Calme 268, $88 - 138 \text{ g m}^{-2}$ for Probus, $51 - 78 \text{ g m}^{-2}$ for Zinal and $81 - 121 \text{ g m}^{-2}$ for CH Claro, or uncertainties in the amount of carbon rhizodeposits between -18 and $+28 \%$. Further research on the effect of the assumption of using root $\delta^{13}\text{C}$ values as a proxy for carbon rhizodeposits is thus necessary to better quantify the effect on estimates of carbon sequestration.

4.4 The effect of old and recent wheat cultivars on net carbon rhizodeposition

Our results indicate that the old wheat cultivars, with deeper active roots throughout the experiment and larger root biomass, allocated more assimilated carbon belowground, although the differences were not statistically significant (Figure 4C, Table 2). However, we found no evidence that wheat cultivars with larger root biomass lead to higher net carbon rhizodeposition (Table 2). Our hypothesis, which stated that wheat cultivars with larger root biomass and deeper roots would lead to larger amounts of net carbon rhizodeposition, could therefore not be confirmed.

470 The total amount of OC that will be stabilized in the soil by the studied wheat cultivars will therefore depend on the long-term fate of the root biomass. The root biomass was higher for the old wheat cultivars, although these differences were mainly limited to the upper 0.45 m of the soil. Due to the destructive sampling of vegetation and soil at the end of the experiment, the fate of root biomass after harvest could not be assessed. Based on the results, one could therefore hypothesize that the higher root biomass of old wheat cultivars would lead to larger rates of carbon sequestration in the long-term. Similarly, Mathew et al. (2017) suggested that growing grasses and maize plants would lead to larger SOC stocks because these plants have the highest total and root biomass, compared to growing crops with a lower biomass. However, it is not straightforward to make predictions about the amount of root biomass that will be stabilized in the soil in the long term, as this depends on the efficiency with which plant-derived biomass is incorporated in microbial biomass (Cotrufo et al., 2013) and interactions between soil depth, the microbial community composition and its substrate preference (e.g. Kramer and Gleixner, 2008), among other factors. During the past century, there has been a continuing increase in the importance of wheat cultivars with smaller root biomass (Fossati and Brabant, 2003; Friedli et al., 2019; Waines and Ehdaie, 2007). This can have profound implications for OC stocks of soils under wheat cultivation, as rhizodeposition and root-derived carbon are the most important inputs of OC to the soil (Kong and Six, 2010). Testing the long-term effect of the gradual change in wheat cultivars on OC inputs to the soil would thus require experiments that run over multiple growing seasons, and allow the quantification of the amount of root carbon that is eventually stabilized in the soil.

Correct knowledge on the amount of OC that is transferred belowground by plants is necessary to reliably model SOC dynamics. However, this knowledge is currently limited and changes in belowground carbon allocation due to the cultivation of different cultivars is generally not considered in SOC models. Moreover, it has recently been shown that accounting for changes in belowground carbon allocation by relating this to changes in aboveground biomass does not improve model results (Taghizadeh-Toosi et al., 2016). Rather, it has been suggested that more reliable model results are obtained when crop-specific amounts of belowground carbon allocation are used, independent of aboveground biomass production (Taghizadeh-Toosi et al., 2016). Since model results are very sensitive to the amount of carbon inputs (Keel et al., 2017), and cereal crops are grown on ca. 20 % of croplands globally (Leff et al., 2004) (covering ca. 12 % of global land mass and storing ca. 10 % of global SOC in the upper meter of soil (Govers et al., 2013)), a correct assessment of a potential decrease in belowground carbon inputs by wheat plants over the past century through the cultivation of different cultivars will have important implications for the simulation of changes in SOC on the global scale.

Assessing the overall impact of the past evolution of wheat cultivars on SOC stocks also requires taking into account the amount of land needed to produce sufficient food. For example, if future research would show that more recent wheat cultivars lead to less SOC stabilization compared to older cultivars, this does not necessarily imply a net loss of SOC as a consequence of the historical shift to planting recently developed wheat cultivars. If the aim is to increase overall SOC stocks, it might be more favorable to grow high-yield wheat cultivars that sequester less OC per unit area compared to a low-yielding cultivar, if this results in a larger area of arable land that can be taken out of cultivation. This land can be put under native vegetation, such as forest or grassland, which stores substantially more SOC compared to arable land (Jobbágy and Jackson, 2000).

5. Conclusion

505 In this study, four different wheat cultivars were grown in lysimeters and labeled with $^{13}\text{CO}_2$ using repeated pulse-labelling to quantify the effect of rooting depth and root biomass on net carbon rhizodeposition. Our results show that there is no clear trend between the time of cultivar development and the amount of net carbon rhizodeposition with large variabilities being observed between replicates of the same cultivars. Based on these results, the hypothesis that wheat cultivars with a larger root biomass and deeper roots would promote carbon net carbon rhizodeposition, was rejected. An important remaining uncertainty
510 is related to the fate of root biomass after harvest, which might contribute to the stabilized SOC pool over the long-term.

Data availability

Additional figures and tables can be found in the supplementary information. The data associated with this manuscript is available in the online version of this manuscript.

Author contribution

515 CDK, SA, AH, CF and JS conceived the idea for the study
CDK set up the lysimeter and labelling experiments and collected the data
SG, CDK, SA and RW performed lab analyses
MVdB, SG, CDK and JS analysed and interpreted the data and performed the statistics
MVdB and SG wrote the manuscript, with contributions from CDK, AH, SA, JS, CF and RW
520 MVdB and SG contributed equally to this manuscript

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgements

The lysimeter infrastructure for this study was funded by ETH core start-up funds provided to Johan Six. Charlotte Decock is
525 grateful for funding from the Plant Fellows, a postdoctoral fellowship administered by the Zurich Basel Plant Science center and funded under the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement no GA-2010-267243 – PLANT FELLOWS. Additional funding for this project came from the Swiss National Science Foundation (Project number 200021_160232). The authors are very grateful to Matti Barthel, Benjamin Wild and Christopher Mikita, for their help with setting up the lysimeters, data collection and interpretation. The

530 authors appreciate the grassland sciences group of ETH Zürich for providing the laboratory facility to perform part of the analyses performed in this study. We thank Brigitta Herzog and Hansueli Zellweger for their help with greenhouse management and plant protection at the ETH Research Station for Plant Sciences in Lindau.

References

- 535 Aziz, M. M., Palta, J. A., Siddique, K. H. M. and Sadras, V. O.: Five decades of selection for yield reduced root length density and increased nitrogen uptake per unit root length in Australian wheat varieties, *Plant Soil*, 413, 181–192, doi:10.1007/s11104-016-3059-y, 2017.
- Baloch, D. M., Karow, R. S., Marx, E., Kling, J. G. and Witt, M. D.: Vernalization Studies with Pacific Northwest Wheat, *Agron. J.*, 95, 1201–1208, doi:10.2134/agronj2003.1201, 2003.
- 540 Brancourt-Hulmel, M., Doussinault, G., Lecomte, C., Bérard, P., Le Buanec, B. and Trottet, M.: Genetic improvement of agronomic traits of winter wheat cultivars released in France from 1946 to 1992, *Crop Sci.*, 43, 37–45, 2003.
- Bromand, S., Whalen, J. K., Janzen, H. H., Schjoerring, J. K. and Ellert, B. H.: A pulse-labelling method to generate ¹³C-enriched plant materials, *Plant Soil*, 235, 253–257, doi:10.1023/A:1011922103323, 2001.
- 545 Brooks, P. D., Geilmann, H., Werner, R. A. and Brand, W. A.: Improved precision of coupled $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements from single samples using an elemental analyzer/isotope ratio mass spectrometer combination with a post-column six-port valve and selective CO₂ trapping; improved halide robustness of the combustion, *Rapid Commun. Mass Spectrom.*, 17, 1924–1926, doi:10.1002/rcm.1134, 2003.
- Chen, S., Martin, M. P., Saby, N. P. A., Walter, C., Angers, D. A. and Arrouays, D.: Fine resolution map of top- and subsoil carbon sequestration potential in France, *Sci. Total Environ.*, 630, 389–400, doi:10.1016/j.scitotenv.2018.02.209, 2018.
- 550 Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D. and Balesdent, J.: Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations, *Soil Tillage Res.*, doi:10.1016/j.still.2018.04.011, 2018.
- Cholick, F. A., Welsh, J. R. and Cole, C. V.: Rooting Patterns of Semi-dwarf and Tall Winter Wheat Cultivars under Dryland Field Conditions, *Crop Sci.*, 17, 637–639, 1977.
- Coplen, T. B.: Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results, *Rapid Commun. Mass Spectrom.*, 25, 2538–2560, doi:10.1002/rcm.5129, 2011.
- 555 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K. and Paul, E.: The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?, *Glob. Chang. Biol.*, 19, 988–995, doi:10.1111/gcb.12113, 2013.
- Dean, A., Morris, M., Stufken, J. and Bingham, D., Eds.: *Handbook of design and analysis of experiments*, CRC Press., 2015.
- 560 Don, A., Schu, J. and Freibauer, A.: Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis, *Glob. Chang. Biol.*, 17(4), 1658–1670, doi:10.1111/j.1365-2486.2010.02336.x, 2011.
- Feil, B.: Breeding Progress in Small Grain Cereals — A Comparison of Old and Modern Cultivars, *Plant Breed.*, 108, 1–11, 1992.
- Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B. and Rumpel, C.: Stability of organic carbon in deep soil layers controlled by fresh carbon supply, *Nature*, 450, 277–280, doi:10.1038/nature06275, 2007.
- 565 Fossati, D. and Brabant, C.: La sélection du blé en Suisse Le programme des stations fédérales Une progression impressionnante, *Rev. Suisse Agric.*, 35(4), 169–180, 2003.
- Friedli, C. N., Abiven, S., Fossati, D. and Hund, A.: Modern wheat semi-dwarfs root deep on demand: response of rooting

- depth to drought in a set of Swiss era wheats covering 100 years of breeding, *Euphytica*, 215(85), 1–15, doi:10.1007/s10681-019-2404-7, 2019.
- 570 Goffin, S., Aubinet, M., Maier, M., Plain, C., Schack-Kirchner, H. and Longdoz, B.: Characterization of the soil CO₂ production and its carbon isotope composition in forest soil layers using the flux-gradient approach, *Agric. For. Meteorol.*, 188, 45–57, doi:10.1016/j.agrformet.2013.11.005, 2014.
- Govers, G., Merckx, R., Van Oost, K. and van Wesemael, B.: Managing Soil Organic Carbon for Global Benefits: A STAP Technical Report, Global Environmental Facility, Washington, D.C., 2013.
- 575 de Graaff, M.-A., Van Kessel, C. and Six, J.: Rhizodeposition-induced decomposition increases N availability to wild and cultivated wheat genotypes under elevated CO₂, *Soil Biol. Biochem.*, 41, 1094–1103, doi:10.1016/j.soilbio.2009.02.015, 2009.
- Gregory, P. J. and Atwell, B. J.: The fate of carbon in pulse-labelled crops of barley and wheat, *Plant Soil*, 136, 205–213, doi:10.1007/BF02150051, 1991.
- 580 Guo, L. B. and Gifford, R. M.: Soil carbon stocks and land use change: a meta analysis, *Glob. Chang. Biol.*, 8, 345–360, doi:10.1046/j.1354-1013.2002.00486.x, 2002.
- Hirte, J., Leifeld, J., Abiven, S., Oberholzer, H.-R. and Mayer, J.: Below ground carbon inputs to soil via root biomass and rhizodeposition of field-grown maize and wheat at harvest are independent of net primary productivity, *Agric. Ecosyst. Environ.*, 265, 556–566, doi:10.1016/j.agee.2018.07.010, 2018.
- 585 Hu, T., Sørensen, P., Wahlström, E. M., Chirinda, N., Sharif, B., Li, X. and Olesen, J. E.: Root biomass in cereals, catch crops and weeds can be reliably estimated without considering aboveground biomass, *Agric. Ecosyst. Environ.*, 251, 141–148, doi:10.1016/j.agee.2017.09.024, 2018.
- Janzen, H. H. and Bruinsma, Y.: Methodology for the quantification of root and rhizosphere nitrogen dynamics by exposure of shoots to ¹⁵N-labelled ammonia, *Soil Biol. Biochem.*, 21(2), 189–196, doi:10.1016/0038-0717(89)90094-1, 1989.
- 590 Jobbágy, E. G. and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to climate and vegetation, *Ecol. Appl.*, 10(2), 423–436, doi:10.1890/1051-0761(2000)010[0423:TVDOSO]2.0.CO;2, 2000.
- Jones, D. L., Nguyen, C. and Finlay, R. D.: Carbon flow in the rhizosphere: carbon trading at the soil–root interface, *Plant Soil*, 321, 5–33, doi:10.1007/s11104-009-9925-0, 2009.
- 595 Keel, S. G., Leifeld, J., Mayer, J., Taghizadeh-Toosi, A. and Olesen, J. E.: Large uncertainty in soil carbon modelling related to method of calculation of plant carbon input in agricultural systems, *Eur. J. Soil Sci.*, 68(6), 953–963, doi:10.1111/ejss.12454, 2017.
- Keith, H., Oades, J. M. and Martin, J. K.: Input of carbon to soil from wheat plants, *Soil Biol. Biochem.*, 18(4), 445–449, 1986.
- Kell, D. B.: Breeding crop plants with deep roots: their role in sustainable carbon, nutrient and water sequestration, *Ann. Bot.*, 108, 407–418, doi:10.1093/aob/mcr175, 2011.
- 600 Kohn, M. J.: Carbon isotope compositions of terrestrial C₃ plants as indicators of (paleo)ecology and (paleo)climate, *Proc. Natl. Acad. Sci. U. S. A.*, 107(46), 19691–19695, doi:10.1073/pnas.1004933107, 2010.
- Kong, A. Y. Y. and Six, J.: Tracing Root vs. Residue Carbon into Soils from Conventional and Alternative Cropping Systems, *Soil Sci. Soc. Am. J.*, 74(4), 1201, doi:10.2136/sssaj2009.0346, 2010.
- Kong, A. Y. Y. and Six, J.: Microbial community assimilation of cover crop rhizodeposition within soil microenvironments in alternative and conventional cropping systems, *Plant Soil*, 356, 315–330, doi:10.1007/s11104-011-1120-4, 2012.
- 605 Kong, A. Y. Y., Six, J., Bryant, D. C., Denison, R. F. and van Kessel, C.: The Relationship between Carbon Input, Aggregation, and Soil Organic Carbon Stabilization in Sustainable Cropping Systems, *Soil Sci. Soc. Am. J.*, 69, 1078–1085, doi:10.2136/sssaj2004.0215, 2005.
- Kramer, C. and Gleixner, G.: Soil organic matter in soil depth profiles: Distinct carbon preferences of microbial groups during

carbon transformation, *Soil Biol. Biochem.*, 40, 425–433, doi:10.1016/j.soilbio.2007.09.016, 2008.

- 610 Kuzyakov, Y.: Sources of CO₂ efflux from soil and review of partitioning methods, *Soil Biol. Biochem.*, 38, 425–448, doi:10.1016/j.soilbio.2005.08.020, 2006.
- Kuzyakov, Y. and Domanski, G.: Carbon Input By Plants in the Soil. Review, *J. Plant Nutr. Soil Sci.*, 163, 421–431, doi:10.1002/1522-2624(200008)163:4<421::AID-JPLN421>3.0.CO;2-R, 2000.
- 615 Leff, B., Ramankutty, N. and Foley, J. A.: Geographic distribution of major crops across the world, *Global Biogeochem. Cycles*, 18, 1–27, doi:10.1029/2003GB002108, 2004.
- Lehtinen, T., Schlatter, N., Baumgarten, A., Bechini, L., Krüger, J., Grignani, C., Zavattaro, L., Costamagna, C. and Spiegel, H.: Effect of crop residue incorporation on soil organic carbon and greenhouse gas emissions in European agricultural soils, *Soil Use Manag.*, 30(4), 524–538, doi:10.1111/sum.12151, 2014.
- 620 Lupton, F. G. H., Oliver, R. H., Ellis, F. B., Barnes, B. T., Howse, K. R., Welbank, P. J. and Taylor, P. J.: Root and shoot growth of semi-dwarf and taller winter wheats, *Ann. Appl. Biol.*, 77, 129–144, 1974.
- Massman, W. J.: A review of the molecular diffusivities of H₂O, CO₂, CH₄, CO, O₃, SO₂, NH₃, N₂O, NO, and NO₂ in air, O₂ and N₂ near STP, *Atmos. Environ.*, 32(6), 1111–1127, doi:10.1016/S1352-2310(97)00391-9, 1998.
- Mathew, I., Shimelis, H., Mutema, M. and Chaplot, V.: What crop type for atmospheric carbon sequestration: Results from a global data analysis, *Agric. Ecosyst. Environ.*, 243, 34–46, doi:10.1016/j.agee.2017.04.008, 2017.
- 625 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B. and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, doi:10.1016/j.geoderma.2017.01.002, 2017.
- 630 Moldrup, P., Olesen, T., Gamst, J., Schjønning, P., Yamaguchi, T. and Rolston, D. E.: Predicting the Gas Diffusion Coefficient in Repacked Soil, *Soil Sci. Soc. Am. J.*, 64, 1588–1594, doi:10.2136/sssaj2000.6451588x, 2000.
- Oburger, E. and Jones, D. L.: Sampling root exudates – Mission impossible?, *Rhizosphere*, 6, 116–133, doi:10.1016/j.rhisph.2018.06.004, 2018.
- 635 Pausch, J. and Kuzyakov, Y.: Carbon input by roots into the soil: Quantification of rhizodeposition from root to ecosystem scale, *Glob. Chang. Biol.*, 24, 1–12, doi:10.1111/gcb.13850, 2018.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P. and Smith, P.: Climate-smart soils, *Nature*, 532, 49–57, doi:10.1038/nature17174, 2016.
- Poeplau, C. and Don, A.: Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-analysis, *Agric. Ecosyst. Environ.*, 200, 33–41, doi:10.1016/j.agee.2014.10.024, 2015.
- 640 Poeplau, C., Don, A., Vesterdal, L., Le, J., van Wesemael, B., Schumacher, J. and Gensior, A.: Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach, *Glob. Chang. Biol.*, 17(7), 2415–2427, doi:10.1111/j.1365-2486.2011.02408.x, 2011.
- 645 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., Boden, T. A., Tans, P. P., Andrews, O. D., Arora, V. K., Bakker, D. C. E., Barbero, L., Becker, M., Betts, R. A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Cosca, C. E., Cross, J., Currie, K., Gasser, T., Harris, I., Hauck, J., Haverd, V., Houghton, R. A., Hunt, C. W., Hurtt, G., Ilyina, T., Jain, A. K., Kato, E., Kautz, M., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lima, I., Lombardozzi, D., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., Padin, X. A., Peregon, A., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Reimer, J., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Viovy, N., Vuichard, N., Walker, A. P., Watson, A. J., Wiltshire, A. J., Zaehle, S. and Zhu, D.: Global Carbon Budget 2017, *Earth Syst. Sci. Data*,

10, 405–448, doi:10.5194/essd-10-405-2018, 2018.

R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2019.

- 655 Ruehr, N. K., Offermann, C. A., Gessler, A., Winkler, J. B., Ferrio, J. P., Buchmann, N. and Barnard, R. L.: Drought effects on allocation of recent carbon: from beech leaves to soil CO₂ efflux, *New Phytol.*, 184, 950–961, doi:10.1111/j.1469-8137.2009.03044.x, 2009.
- Sandén, T., Spiegel, H., Stüger, H.-P., Schlatter, N., Haslmayr, H.-P., Zavattaro, L., Grignani, C., Bechini, L., D'Hose, T., Molendijk, L., Pecio, A., Jarosz, Z., Guzmán, G., Vanderlinden, K., Giráldez, J. V., Mallast, J. and ten Berge, H.: European long-term field experiments: knowledge gained about alternative management practices, edited by M. Aitkenhead, *Soil Use Manag.*, 34, 167–176, doi:10.1111/sum.12421, 2018.
- 660 Shearman, V. J., Sylvester-bradley, R., Scott, R. K. and Foulkes, M. J.: Physiological Processes Associated with Wheat Yield Progress in the UK, *Crop Sci.*, 45, 175–185, 2005.
- Stevenel, P., Frossard, E., Abiven, S., Rao, I. M., Tamburini, F. and Oberson, A.: Using a Tri-Isotope (¹³C, ¹⁵N, ³³P) Labelling Method to Quantify Rhizodeposition, in *Methods in Rhizosphere Biology Research*, edited by D. Reinhardt and A. K. Sharma, pp. 169–195, Springer, Singapore., 2019.
- 665 Studer, M. S., Siegwolf, R. T. W. and Abiven, S.: Carbon transfer, partitioning and residence time in the plant-soil system: a comparison of two $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ labelling techniques, *Biogeosciences*, 11, 1637–1648, doi:10.5194/bg-11-1637-2014, 2014.
- 670 Sun, Z., Chen, Q., Han, X., Bol, R., Qu, B. and Meng, F.: Allocation of photosynthesized carbon in an intensively farmed winter wheat–soil system as revealed by ¹⁴CO₂ pulse labelling, *Sci. Rep.*, 8(3160), 1–10, doi:10.1038/s41598-018-21547-y, 2018.
- Swinnen, J., van Veen, J. A. and Merckx, R.: ¹⁴C pulse-labeling of field-grown spring wheat: an evaluation of its use in rhizosphere carbon budget estimations, *Soil Biol. Biochem.*, 26(2), 161–170, 1994.
- 675 Taghizadeh-Toosi, A., Christensen, B. T., Glendining, M. and Olesen, J. E.: Consolidating soil carbon turnover models by improved estimates of belowground carbon input, *Sci. Rep.*, 6(32568), 1–7, doi:10.1038/srep32568, 2016.
- Tester, M. and Langridge, P.: Breeding technologies to increase crop production in a changing world, *Science* (80-.), 327, 818–822, doi:10.1126/science.1183700, 2010.
- 680 Tubiello, F. N., Salvatore, M., Ferrara, A. F., House, J., Federici, S., Rossi, S., Biancalani, R., Condor Golec, R. D., Jacobs, H., Flammini, A., Prosperi, P., Cardenas-Galindo, P., Schmidhuber, J., Sanz Sanchez, M. J., Srivastava, N. and Smith, P.: The Contribution of Agriculture, Forestry and other Land Use activities to Global Warming, 1990-2012, *Glob. Chang. Biol.*, 21(7), 2655–2660, doi:10.1111/gcb.12865, 2015.
- Vance, E. D., Brookes, P. C. and Jenkinson, D. S.: An extraction method for measuring soil microbial biomass C, *Soil Biol. Biochem.*, 19(6), 703–707, doi:10.1016/0038-0717(87)90052-6, 1987.
- 685 Voss-Fels, K. P., Stahl, A., Wittkop, B., Lichthardt, C., Nagler, S., Rose, T., Chen, T.-W., Zetzsche, H., Seddig, S., Majid Baig, M., Ballvora, A., Frisch, M., Ross, E., Hayes, B. J., Hayden, M. J., Ordon, F., Leon, J., Kage, H., Friedt, W., Stützel, H. and Snowdon, R. J.: Breeding improves wheat productivity under contrasting agrochemical input levels, *Nat. Plants*, 5, 706–714, doi:10.1038/s41477-019-0445-5, 2019.
- 690 Wacker, L., Jacomet, S. and Körner, C.: Trends in Biomass Fractionation in Wheat and Barley from Wild Ancestors to Modern Cultivars, *Plant Biol.*, 4, 258–265, doi:10.1055/s-2002-25735, 2002.
- Waines, J. G. and Ehdaie, B.: Domestication and Crop Physiology: Roots of Green-Revolution Wheat, *Ann. Bot.*, 100, 991–998, doi:10.1093/aob/mcm180, 2007.
- Wasson, A. P., Richards, R. A., Chatrath, R., Misra, S. C., Prasad, S. V. S., Rebetzke, G. J., Kirkegaard, J. A., Christopher, J.

- 695 and Watt, M.: Traits and selection strategies to improve root systems and water uptake in water-limited wheat crops, *J. Exp. Bot.*, 63(9), 3485–3498, doi:10.1093/jxb/ers111, 2012.
- Werner, R. A. and Brand, W. A.: Referencing strategies and techniques in stable isotope ratio analysis, *Rapid Commun. Mass Spectrom.*, 15, 501–519, doi:10.1002/rcm.258, 2001.
- 700 Werner, R. A., Bruch, B. A. and Brand, W. A.: ConFlo III - an interface for high precision $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis with an extended dynamic range, *Rapid Commun. Mass Spectrom.*, 13(13), 1237–1241, doi:10.1002/(SICI)1097-0231(19990715)13:13<1237::AID-RCM633>3.0.CO;2-C, 1999.
- Zeeman, M. J., Werner, R. A., Eugster, W., Siegwolf, R. T. W., Wehrle, G., Mohn, J. and Buchmann, N.: Optimization of automated gas sample collection and isotope ratio mass spectrometric analysis of $\delta^{13}\text{C}$ of CO_2 in air, *Rapid Commun. Mass Spectrom.*, 22, 3883–3892, doi:10.1002/rcm.3772, 2008.




Figure 1 Aboveground (A) and root (B) biomass of the different wheat cultivars at the end of the experiment. Bars represent the average per wheat cultivar, error bars show the standard error ($n = 3$) and circles show the individual data points. The inset in (B) shows a detail of the subsoil root biomass. If statistically significant differences were present, these are indicated with letters, with variables sharing a letter not being significantly different. For root biomass, statistical analyses were performed using a three-way analysis of variance, in contrast to the two-way analysis used to calculate significant differences for the total root biomass summed over all depths as reported in Table 1 (see section 2.4).




Figure 2 $\delta^{13}\text{C}$ values of aboveground (A) and belowground (B) biomass for the different wheat cultivars at the end of the experiment. Bars (A) and symbols (B) represent the average per wheat cultivar based on 3 replicates, error bars show the standard error ($n = 3$), symbols without error bars indicate samples for which no 3 replicates were available. Circles show the individual data points. If statistically significant differences were present for root biomass at the same depth, these are indicated with letters, with variables sharing a letter not being significantly different and data points without error bars being left out of the analyses.




Figure 3 Depth profiles of organic carbon concentration (A), the $\delta^{13}\text{C}$ value of organic carbon (B), the $\delta^{13}\text{C}$ value of microbial biomass (C) and the $\delta^{13}\text{C}$ value of soil CO_2 , averaged per wheat cultivar at the end of the experiment. Error bars represent the standard error ($n = 3$). *The initial soil indicates measurements performed on the soil that was used to fill the lysimeters prior to the experiments.




Figure 4 Absolute (A) and relative (B) distribution of excess ^{13}C between aboveground biomass, root biomass and soil for the different wheat cultivars. Soil compartments are calculated down to 0.45 m depth. (C) shows the total carbon rhizodeposition and root carbon for the upper two soil layers for the different wheat cultivars (0 – 0.45 m depth). Error bars represent the standard error ($n = 3$). In (A), letter indicate significant differences between the total amount of excess ^{13}C of the different cultivars. In (C), letters are provided when the total belowground C allocation differed between the different depth layers, which was only the case for Zinal.




Figure 5 Changes in the CO₂ concentration (ppm) in the lysimeters throughout the experiment for the four wheat cultivars. The average CO₂ concentration of three replicates are shown (n = 3). Dots indicate the measured data points.




Figure 6 Depth profiles of calculated cumulative CO₂ production (g CO₂ m⁻² per 0.05 m depth layer). Dots show the calculated production rates, thin lines show the calculated CO₂ production for the individual lysimeters, while the thick lines show the average based on three replicates (two for CH Claro).

745

Table 1 Characteristics (\pm standard error, $n = 3$) of the biomass of the different wheat cultivars at the end of the experiment. Values that share a letter in the same column are not significantly different.

Cultivar (year of release)	Aboveground biomass		Root biomass		Root:shoot ratio
	Biomass (g m^{-2})	OC %	Biomass (g m^{-2})	OC %	
CH Claro (2007)	$1064 \pm 207^{\text{ab}}$	$40.5 \pm 0.3^{\text{ab}}$	$107 \pm 28^{\text{ab}}$	$38.7 \pm 1.9^{\text{a}}$	$0.10 \pm 0.02^{\text{a}}$
Zinal (2003)	$710 \pm 114^{\text{b}}$	$40.0 \pm 0.14^{\text{a}}$	$97 \pm 20^{\text{b}}$	$39.1 \pm 1.1^*$	$0.14 \pm 0.01^{\text{a}}$
Probus (1948)	$1154 \pm 220^{\text{a}}$	$41.9 \pm 0.2^{\text{b}}$	$161 \pm 54^{\text{a}}$	$38.1 \pm 0.5^{\text{a}}$	$0.13 \pm 0.03^{\text{a}}$
Mont-Calme 268 (1926)	$1119 \pm 174^{\text{ab}}$	$40.8 \pm 0.4^{\text{ab}}$	$205 \pm 67^{\text{a}}$	$36.8 \pm 1.3^{\text{a}}$	$0.19 \pm 0.08^{\text{a}}$

* Was excluded from statistical analysis due to missing data

750

Table 2 Average belowground carbon allocation (net rhizodeposition and root biomass combined) and net carbon rhizodeposition by the different wheat cultivars, calculated down to a depth of 0.45 m (variation is reported as the standard error, $n = 3$). No statistically significant differences in belowground C allocation of net C rhizodeposition between any of the cultivars were detected.

	Belowground C allocation (g m^{-2})	Net C rhizodeposition (g m^{-2})
CH Claro	131 ± 26	97 ± 24
Zinal	97 ± 14	62 ± 11
Probus	164 ± 38	108 ± 34
Mont-Calme 268	154 ± 39	83 ± 29