

Interactive comment on “What was the source of the atmospheric CO₂ increase during the Holocene?” by Victor Brovkin et al.

Peter Köhler

peter.koehler@awi.de

Received and published: 4 May 2019

In the figures of my previous comment the labels of the top x-axes (time in yr CE) are placed wrong, they need to get shifted by 50 years. Furthermore, the x-axes started at -100 yr BP, that has now been changed to -66 yr BP = 2016 CE, which is the first year with data.

I apologize for any confusion, the corrected figures are attached below, including their full captions.

Fig. 1: Atmospheric CO₂ spline and underlying data (2016 CE to -6050 CE or -66 BP to 8,000 BP). Black spline as published in Köhler *et al.* (2017) against time series (gold) used in Brovkin *et al.* (2019). Error bars around the ice core data points are $\pm 2\sigma$. WDC

Printer-friendly version

Discussion paper



data have been adjusted to reduce offsets, see *Köhler et al. (2017)* for details.

Fig. 2: Atmospheric CH₄ spline and underlying data (2016 CE to -6050 CE or -66 BP to 8,000 BP). Black spline as published in *Köhler et al. (2017)* against time series (gold) used in *Brovkin et al. (2019)*. Details on plotted data are explained in *Köhler et al. (2017)*. The maximum ice core data uncertainty ($\pm 2\sigma$) is sketched in the lower left corner. Latitudinal origin of data is indicated by NH and SH, implying northern and southern hemisphere, respectively.

Fig 3: Atmospheric N₂O spline and underlying data (2016 CE to -6050 CE or -66 BP to 8,000 BP). Black spline as published in *Köhler et al. (2017)* against time series (gold) used in *Brovkin et al. (2019)*. Details on plotted data are explained in *Köhler et al. (2017)*. The maximum ice core data uncertainty ($\pm 2\sigma$) is sketched in the upper right corner. Filled symbols: data taken for spline; open symbols: data not taken for spline.

References

- Brovkin, V., S. Lorenz, T. Raddatz, T. Ilyina, I. Stemmler, M. Toohey, and M. Claussen (2019), What was the source of the atmospheric CO₂ increase during the Holocene?, *Biogeosciences Discussions*, 2019, 1–25, doi:10.5194/bg-2019-64.
- Köhler, P., C. Nehrbass-Ahles, J. Schmitt, T. F. Stocker, and H. Fischer (2017), A 156 kyr smoothed history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363–387, doi:10.5194/essd-9-363-2017.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2019-64>, 2019.

BGD

Interactive
comment

Printer-friendly version

Discussion paper






Fig. 1. Corrected Fig. 1 on CO₂ of previous comment

[Printer-friendly version](#)[Discussion paper](#)




Fig. 2. Corrected Fig. 2 on CH₄ of previous comment

Printer-friendly version

Discussion paper






Fig. 3. Corrected Fig. 3 on N₂O of previous comment

Printer-friendly version

Discussion paper

