

Interactive comment on “Can ocean community production and respiration be determined by measuring high-frequency oxygen profiles from autonomous floats?” by Christopher Gordon et al.

David Nicholson

dnicholson@whoi.edu

Received and published: 30 April 2020

General comments: This study seeks to apply dissolved oxygen measurements from profiling floats to estimate primary production and respiration from diel oxygen cycles. The study region in the shelf break region of the northern Gulf of Mexico is a challenging environment for this approach because it is a region of low productivity, but high physical variability and thus is a good testbed to evaluate the limits of diel approaches. Further, near 30 N the Coriolis frequency is approximately 24-hours and near-inertial oscillations can confound biologically-driven diurnal cycles. In general, this study found that physical variability was too great to allow for robust estimates of biological rates in

[Printer-friendly version](#)

[Discussion paper](#)



this challenging environment. This contribution is valuable as it is important to recognize the limitations of such methods.

Given the strong vertical O₂ gradient and slow response time of the optode sensors, a significant portion of the manuscript is dedicated to optimizing methods for deconvolving the oxygen time series and estimate sensor response time in situ. Indeed, this is such a major part of the manuscript, I would recommend changing the title of the paper to in some way reflect the time response part of the manuscript. Although dynamic corrections to optode oxygen sensors on floats has been explored in depth before (Bittig et al., 2014; Bittig and Körtzinger, 2017) the dataset here provides a valuable addition particularly because both ascent and descent profiles were logged, time-stepped and recorded in full resolution without binning.

Overall, I think this is a valuable contribution that will spur improved methodologies for correcting dissolved oxygen in biogeochemical Argo applications. The results should be further applicable to other platforms such as gliders and profiling moorings. I have several concerns about the analysis and some suggestions to extend interpretation that I think would be worthwhile for the authors to consider.

Specific comments:

1. It seems a 7-pt moving mean smoothing is performed prior to deconvolution. With the stated 5 m resolution and 12 cm s⁻¹ average vertical velocity that works out to averaging over about a 40 sec period. A moving average also is a filter (and one with a messy response in the frequency domain). I am concerned that this step would alter the calculated sensor response time that is determined by deconvolution after this averaging. Does the moving average operator slow down and/or complicating the sensor response before the deconvolution is even applied? An easy test is to report if the same median time responses are recovered without the moving mean step.

2. Bittig and Kortzinger (2017) outlined a detailed approach for scaling tau as a function of temperature and flow speed. What is the implication of using a constant tau here

BGD

Interactive
comment

Printer-friendly version

Discussion paper



instead of the temperature and boundary-layer dependent τ . Is it possible to apply the Bittig approach as well for comparison? There is a significant vertical temperature gradient in the study region and thus an expectation that response time would be slower in deeper water than near the surface.

3. A recent publication by Barone et al. (2019) quantifies GPP and R from diel cycles with uncertainty and fit statistics. Applying this approach would provide a more quantitative assessment of how good (or bad) daily diel fits are.

4. Are any corrections made for air-sea O₂ flux? It sounds like there was significant atmospheric forcing. Barone et al. 2019 outlines how diel O₂ inventories can be corrected for air-sea flux prior to fitting a diel cycle.

Technical suggestions:

L21: should specify that 12-24 hour incubations approximate NPP (cite Marra 2009). Other short-term incubation approaches also are fairly commonly used and measure something closer to GPP.

P2-L42: add (Barone et al., 2019) and (Johnson, 2010)

P3-L67: add salinity to list of corrections

P6: L67: There is temperature dependence both to molecular diffusivity and kinematic viscosity

P8 L84: since τ is a function of environment it is also a function of time but treated as a constant. How does that impact interpretation?

Fig 2: The label 'Scatter' in B seems odd. Maybe use 'difference' instead?

P9 L29: change 'listen' to 'listed'

P13 L53: It seems possible that depending on sensor orientation there could be a big difference in up vs. down response time. Was this tested at all? Barone et al. (2019)

Printer-friendly version

Discussion paper



found ~ 35 sec tau for the same sensor on Seagliders (see supplemental info).

Barone, B., Nicholson, D., Ferrón, S., Firing, E. and Karl, D.: The estimation of gross oxygen production and community respiration from autonomous time-series measurements in the oligotrophic ocean, *Limnol. Oceanogr. Methods*, 17(12), 650–664, doi:10.1002/lom3.10340, 2019.

Bittig, H. C. and Körtzinger, A.: Technical note: Update on response times, in-air measurements, and in situ drift for oxygen optodes on profiling platforms, *Ocean Sci*, 13(1), 1–11, doi:10.5194/os-13-1-2017, 2017.

Bittig, H. C., Fiedler, B., Scholz, R., Krahnemann, G. and Körtzinger, A.: Time response of oxygen optodes on profiling platforms and its dependence on flow speed and temperature, *Limnol. Oceanogr. Methods*, 12(8), 617–636, doi:10.4319/lom.2014.12.617, 2014. Johnson, K. S.: Simultaneous measurements of nitrate, oxygen, and carbon dioxide on oceanographic moorings: Observing the Redfield ratio in real time, *Limnol. Oceanogr.*, 55(2), 615–627, doi:10.4319/lo.2010.55.2.0615, 2010.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-119>, 2020.

BGD

Interactive
comment

Printer-friendly version

Discussion paper

