Winter atmospheric nutrients and pollutants deposition on West

Sayan mountain lakes (Siberia)

Daniel Diaz-de-Quijano¹; Aleksander Vladimirovich Ageev¹; Elena Anatolevna Ivanova¹; Olesia Valerevna Anishchenko^{1,2}

¹Siberian Federal University, 79, Svobondyi prospekt, Krasnoyarsk, 660041, Krasnoyarskii Krai, Russian Federation.
²Institute of Biophysics, Siberian Branch, Russian Academy of Sciences, 50/50, Akademgorodok, Krasnoyarsk, 660036, Krasnoyarskii Krai, Russian Federation.

Correspondence to: Diaz-de-Quijano, D. (diasdekikhanobarbero@sfu-kras.ru, daniquijano@gmail.com)

Код поля изменен Код поля изменен

	Abstract. The world map of anthropogenic atmospheric nitrogen deposition and its effects on natural ecosystems is not
10	described with equal precision everywhere. In this paper, we report atmospheric nutrient, sulphate and spheroidal
	carbonaceous particles (SCPs) deposition rates, based on snowpack analyses, of a formerly unexplored Siberian mountain
	region. Then, we discuss their potential effects on lake phytoplankton biomass limitation.

We estimate that the nutrient depositions observed in the late season snowpack ($40\pm16 \text{ mg NO}_3\text{-N}_{c}\text{m}^{-2}$ and $0.58\pm0.13 \text{ mg}$ TP-P₂m⁻²) would correspond to yearly depositions lower than $119\pm71 \text{ mg NO}_3\text{-N}_{c}\text{m}^{-2}\text{s}\text{y}^{-1}$ and higher than $1.71\pm0.91 \text{ mg}$ TP-

- P_xm²_xy¹. These yearly deposition estimates would approximately fit the predictions of global deposition models and correspond to the very low nutrient deposition range although they are still higher than world background values. In spite of the fact that such low atmospheric nitrogen deposition rate would be enough to induce nitrogen limitation in unproductive mountain lakes_x phosphorus deposition was also extremely low and the lake water N:P ratio resulted to be unaffected by atmospheric nutrient deposition. In the end, the studied lakes phytoplankton appeared to be hanging on the
- 20 fence between phosphorus and nitrogen limitation. We conclude that these pristine lakes are fragile sensitive systems exposed to the predicted climate warming, increased winter precipitation, enhanced forest fires and shifts in anthropogenic nitrogen emissions that could finally couple their water chemistry to that of atmospheric nutrient deposition and unlock temperature-inhibited responses of phytoplankton to nutrient shifts.

25 1 Introduction

Worldwide nitrogen cycle perturbation is the second most important global environmental concern, just after massive extinction of species and even more important than global warming (Rockström et al., 2009; Steffen et al., 2015). The

1

6	удалено: ×
6	удалено:
6	удалено:
(удалено: ·
6	удалено: ·
9	удалено: ·
6	удалено: the extremely low
6	отформатировано: Английский (США)
1 H	удалено: would have made the bioavailable N:P deposition ratio to be frankly high. In the end,
(1	удалено: , with a trend towards nitrogen limitation
7	

удалено: slight imbalances in the nutrient deposition might have important effects on the ecology of these lakes under the expected scenario of cli

anthropogenic mobilization of formerly inaccessible nitrogen compartments has more than doubled natural nitrogenasemediated inputs of reactive nitrogen forms into the global nitrogen cycle (Vitousek et al., 1997). Massive fossil fuel combustion since the industrial revolution, chemical fixation of atmospheric diatomic nitrogen to produce fertilizers since the Second World War and the wide extension of leguminous crops are the most important human sources of nitrogen cycle

45 perturbation (Vitousek et al., 1997). A substantive part of this anthropogenic reactive nitrogen is then spread, air-transported and deposited all over the world with a diverse impact on different ecosystems. The effects of atmospheric nitrogen deposition on primary production have been documented in the usually nitrogen-limited

terrestrial ecosystems (Bobbink et al., 2010; DeForest et al., 2004; Güsewell, 2004; LeBauer and Treseder, 2008), as well as in commonly phosphorus-limited lakes (Bergström et al., 2005). A series of studies all over Sweden and abroad showed

- 50 atmospheric nitrogen deposition to have turned unproductive lake phytoplankton from natural nitrogen to induced phosphorus limitation (Bergström and Jansson, 2006; Bergström et al., 2005; Elser et al., 2009) when temperature was not a limiting factor (Bergström et al., 2013). Of course, these changes do not only concern primary production limitation, but also primary producer species composition, cascade effects over the food web, secondary production, species interactions, etc. Likewise, these studies showed that it was reasonable to study the relationship between atmospheric nutrient deposition and
- 55 lake phytoplankton growth limitation independently from biogeochemical processes occurring at the levels of the watershed, runoff and river transport, lake sediments, etc.

Nevertheless, ecological processes are not homogeneous around the World. There is a particular and dynamic geography of reactive nitrogen sources, an atmospheric conveyor belt with a conspicuous structure, an evolving climate with patchy temperature and precipitation changes, and a multiplicity of lake districts with distinct individual lakes in them. If it is true

- 60 that climatic and atmospheric nutrient deposition models have helped a lot to describe this geography, the latter ones lack empirical measurements for some regions of the world, which might undermine their regional spatial reliability in comparison to climate models (Fagerli et al., 2019; Gauss et al., 2019; Lamarque et al., 2013; Mahowald et al., 2008), Moreover, not all lake districts of the world have been studied with the same intensity, so certain processes might be overlooked and the limnological paradigms might be site-biased (Marcé et al., 2015). In this study, we analysed the
- 65 snowpack in the West Sayan mountains (south central Siberia) in order to gauge atmospheric nitrogen, phosphorus, sulphate and spheroidal carbonaceous particles (SCPs) deposition rates. As far as we know, no such measurements had been pursued in this site before, so they might be useful to contrast and inform world deposition models. Besides, we have also assessed lake phytoplankton nutrient limitation regime and discussed the potential influence of nutrient deposition on it.

According to published global models (IPCC, 2013; Lamarque et al., 2013), the West Sayan mountains, in south central 70 Siberia, correspond to a low atmospheric nitrogen deposition area with a cold but increasingly warming climate in the last

decades. Our aim was to corroborate it because in case it was confirmed, it would be an adequate site to study the effects of global warming on ecosystems with a minimal interference of atmospheric nitrogen deposition. In other words, identifying and studying such areas could help disentangle warming and nitrogen fertilization as drivers of ecological change. It could also contribute to assess the worthiness to implement global nitrogen cycle policies, besides climate ones.

удалено: not only

удалено: paradigmatically

удалено: but also in the paradigmatically

(удалено: world
~(удалено: у
\mathcal{I}	удалено:

УДАЛЕНО: "does not occur on a needle tip", as Ramon Margalef used to say (Bascompte and Solé, 2005; Margalef, 1986)...

Код поля изменен

отформатировано: Испанский отформатировано: Испанский

удалено: cold удалено: but notably warmed and relatively

85 2 Methods

90

100

2.1 Study site and sampling

West Sayan mountain range is located in south central Siberia (fig. 1). It has a central position in the Altay-Sayan mountain system, in between the Altay mountains (to the west) and East Sayan mountains (to the east), which are constituents of the Sayan-Baikal mobile fold belt south the Siberian craton (Logatchev, 1993). West Sayan orogeny occurred in the ancient Paleozoic, by folding Paleozoic and Precambrian deposits, during the Baikal tectogenesis and in the Cenozoic era (namely

during the Pliocene-Pleistocene Epochs) (Chernov et al., 1988). With a north-west orientation and heights from 400 to 2700 m.a.s.l., West Sayan mountains combine old eroded with typical glacial reliefs, carved during the Pleistocene glaciation in the highest ridges. The source of Yenisei river, the first Siberian river in terms of discharge, is located in West Sayan mountains and its headwater tributaries are also Sayanic.



Figure 1. Distribution of sampling points in the Ergaki Natural Park. Snow cores: Tsirkovoe (A), Oiskoe (B) and Tushkan (C);-Lakes: Tsirkovoe (1), Oiskoe (2), Raduzhnoe (3), Karovoe (4), Svetloe (5); Olenva Rechka metereological station (M). Ergaki location in Eurasia. Mountain ridge (grev), open spaces (white), forest and bushes (green), three lane federal road (vellow). Source: http://www.shandl.narod.ru/map.htm and Wikipedia CC BY-SA 3.0

The present study was performed in the Ergaki Natural Park, in the West Sayan mountains. With an altitude range from 700 to 2466 m.a.s.l., this park is well known for the glacial landscapes of both Ergaki and Aradan ridges embedded in a boreal mountainous taiga matrix, that extends far to the north. The landscape is spattered with monumental and pictoric granite-syenite rocks, and the general geology is rich in granitoids (Voskresenskii, 1962). South from the park, sub-boreal larch

3

отформатировано: Шрифт: 12 пт

Отформатировано: Название объекта

105 taigas and central Asian steppes develop. The closest gardens and agricultural fields are located downhill more than 35 Km north from the northernmost sampling point and constitute a modest patch within the taiga matrix. <u>Meteorological conditions</u> in the Ergaki Natural Park are characterized by high precipitation (1243 mm) and extreme temperatures, ranging from -36.8 to +33.3°C (fig. <u>2</u> a). From a geobotanical point of view the park is located in the holarctic kingdom, circumboreal region and Altair Sayan province (Takhtadzhyan, 1978). An exhaustive floristic description is available at (Stepanov, 2016), where 110 our study area corresponds to the L3 district.



Figure 2., Records, at Olenya Rechka meteorological station. a: Climograph (1 February 2005 - 30 April 2019). The median yearly-precipitation was 1242.975 mm, with a winter (2nd October to 5th April) and summer precipitations of 464.95 mm and 778.025 mm, respectively. Temperature: median (red), interquartile range (yellow band), 5th to 25th percentiles and 75th to 95th percentiles (lower and upper golden bands), below the 5th percentile and above the 95th percentile (lower and upper dark grav bands), below the 5th percentile and above the 95th percentile (lower and upper dark grav bands), below the 5th percentile and above the 95th percentile (lower and upper dark grav bands). The red line corresponds to the 2016-2017 snowpack thickness record until the snow sampling date. c: Air temperatures measured all the 3 hours during the time period when analysed snowpacks were laying on their respective locations (2016-17).

Snowpack cores were sampled at three sites of the Ergaki Natural Park: next to lake Tsirkovoe (Цирковое) next to lake Oiskoe (Ойское) and on a forest glade close to Tushkan stream (Тушкан) (fig. 1, table 1). Snow sampling was conducted the 5th of April 2017, integrating a snowfall period of 6 months and 5 days according to precipitation data recorded in the closeby Olenya Rechka metereological station (http://rp5.ru). The 1st and 24th September 2016 snowfalls thinner than 0.5 cm and 2 cm respectively were registered but they melted the following day. The first important snowfall occurred the 1st October 2016 evening and left a 22 cm pack that was not significantly reduced anymore until the sampling day (fig. 2,b). Air

удалено: Climate
удалено: ru
удалено: is
удалено: 1
удалено: province
удалено: у
Удалено: Lowland deciduous birch (<i>Betula pendula</i> Roth.) and aspen (<i>Populus tremula</i> L.) forests are succeeded by pine (<i>Pinus</i> sylvestris L. 1753) and larch (<i>Larix sibirca</i> Ledeb. 1833) light taiga which extends from the lowlands to an altitude of about 1300-140011
удалено: Interesting facts are remarked in this monography, such as the presence of southern fabaceae species originary from arid[2]
отформатировано: Шрифт: 12 пт
Отформатировано: Название объекта
удалено: Climatic
удалено: characteristics
удалено:[3]
отформатировано: Английский (США)
отформатировано: Английский (США)
удалено: (Ойское)
отформатировано: Английский (США)
отформатировано: Английский (США)

temperatures recorded in the mentioned meteorological station during this time window were mostly below zero, with positive temperatures only for some hours around midday during the first and last weeks (fig. 2,c).

÷ .						• •			
	A				Distance in m to local pollution sources				
	Symbol in the	latitude	longitude	Altitude	road	cottage	inflow from houses		
	map			(m)					
Tsirkovoe	<u>1, A</u>	52°52'28.3"N	93°14'53.1"E	1428	466	-	- 4		
Oiskoe,	2, B	<u>52°50'28.3"N</u>	93°14'46.0"E	1418	251	229	288		
<u>Tushkan</u>	<u>C</u>	<u>52°46'16.9"N</u>	<u>93°21'17.0"E</u>	1125	725	471			
Raduzhnoe	<u>3</u>	52°50'08.4"N	93°20'44.5"E	1462	4600	3000	-		
Karovoe	<u>4</u>	52°49'57.4"N	93°21'41.6"E	1632	5265	4000	-		
Svetloe	<u>5</u>	52°48'02.2"N	93°25'05.4"E	1511	5647	5470	-		

 Table 1 Sampling sites and distance to local perturbations. Temporary summer camps are present on Svetloe lake shore and used

 185
 to be on Raduzhnoe's. See Table S1 for lake and watershed cover characteristics.

The snow core sampling was conducted following a modified version of the MOLAR project protocols for atmospheric deposition assessment (Mosello et al., 1997). Sampling areas were chosen on a map to be accessible but as far as possible from local sources of air pollution. Definitive locations were also chosen to represent average snowpack thicknesses by checking it across the sampling areas using a snow probe. Hence, wind- and orography-induced secondary modifications of

- 190 the snowpack were minimised. An aluminium tube and piston (1m x 2.5 cm inner diameter), plastic shovel, plastic containers and rubber gloves were soaked in ~4% HCl and MQ water rinsed before being used to pick up snow cores. A protective mask and synthetic clothes were worn during sampling. The snow was stored in the plastic containers and kept at 20°C until further analyses. Two cores divided in three segments (0-40, 40-80 and 80-115 cm) were sampled at Tsirkovoe, whereas three cores divided in two segments (0-60 and 60-115 cm) were sampled at Oiskoe and Tushkan.
- 195 A selection of five accessible Ergaki mountain lakes was chosen to represent a variety in dimensions, altitude, and watershed vegetation covers (fig. 1, table 1 and S1). Oiskoe and Svetloe (Светлос) lakes are relatively large forest lakes (0.57 and 0.37 Km², 21 and 24 m maximum depth) with low water transparency (4 and 8 m average Secchi disk, respectively). Tsirkovoe, Raduzhnoe (Радужное) and Karovoe (Каровое) lakes are located at an alpine landscape and are smaller and shallower (0.02, 0.03, and 0.08 Km², respectively; 15, 4 and 7m maximum depth, respectively). Secchi disk was not tested at Tsirkovoe but
- 200 Raduzhnoe and Karovoe lake beds were visible. Karovoe and Svetloe lakes represent a 7% of the watershed area, Tsirkovoe and Oiskoe, a 5%, and Raduzhnoe is only a 1.4% of its watershed area. As for vegetation cover, Oiskoe and Svetloe watersheds have a 25 and 28% forest cover, respectively, whereas the other lakes have less than 10% forest covers. These two watersheds are quite similar in terms of land cover: they have quite equilibrated percentages of forests, shrubs, meadows and scree. Oiskoe is also the watershed with higher peatland cover
- 205 (6%), followed by Svetloe (3%) and Radushnoe (1.6%). Karovoe and Raduzhnoe watersheds are dominated by scree (73%)
 - 5



and 52%, respectively) and meadows (14% and 24%, respectively), whereas Tsirkovoe watershed is dominated by shrubs (56%) and scree (37%).

Lake water was sampled at different depths (from 1 to 5 separate depths, or composite sample from that depths, depending on the sampling expedition, see table S2) and consequently analysed for chlorophyll and nutrient content in early September 2015 (Tsirkovoe, Oiskoe and Raduzhnoe) and in late August 2017 (Tsirkovoe, Oiskoe and Karovoe). Data from a previously published study integrating June and August samplings 2011-12 (Oiskoe, Raduzhnoe, Karovoe and Svetloe) was also used (Anishchenko et al., 2015). Water samples were sieved *in situ* to remove zooplankton, transported to the field laboratory at 4-10°C in the dark, filtered for chlorophyll analyses and frozen at -20°C for further chemical analyses.

265 2.2 Chemical analyses

Snow and lake water samples were fully thaw<u>ed</u> and stirred before analyses. Snow water equivalent (SWE) was calculated by multiplying the snowpack depth and the ratio of melted water volume to sampled snow volume. <u>Samples were gently</u> vacuum filtered for NO_2^- , NO_3^- , soluble reactive phosphorus –SRP– and $SO_4^2^-$ analyses and filtered with single-use plastic syringes for NH₄⁺ analyses. Cellulose filters were used in 2011 and 2012 surveys according to the Russian National

- 270 Standards (Gladyshev et al., 2015; Tolomeev et al., 2014) and 0.45 μm membrane filters "Porafil" (Macherey-Nagel, Germany) were used in 2015 and 2017. Nitrate was reduced to NO_2^- by the cadmium reduction method. Nitrite was determined by the colorimetric method after reacting with sulphanilamide and α-naphthylamine. Detection limits were 0.001 $\underline{mg} N - NO_3^-$; t^{-1} , 0,0002 mg $N - NO_2^-$; t^{-1} , and 0.011 mg N-NH4· t^{-1} . Lake water dissolved inorganic nitrogen (DIN) was calculated as the sum of nitrate, nitrite and ammonium. Soluble reactive phosphorus was assessed using the ascorbic acid and
- 275 ammonium molybdate method. Total phosphorus (<u>TP</u>) was measured the same way after persulfate digestion of unfiltered samples. In the 2015 survey, lake water TP was calculated as the sum of soluble reactive phosphorus and seston particulate phosphorus (PP). The latter was collected on the above-mentioned membrane filters and measured after persulfate digestion. All these analyses were made according to the Russian National Standards (Gladyshev et al., 2015; Tolomeev et al., 2014), which generally coincide to those from APHA (APHA, 1989). In 2015 nutrients were measured using a Flow Injection
- 280 Analyser Lachat Quickchem 8500 autoanalyzer Series 2 FIA System (Hach Ltd, Loveland, CO, U.S.). The method detection limits (MDL) were 0.0263 mg N-NO₃·l⁻¹, 0.0057 mg N-NO₂·l⁻¹, 0.0479 mg N-NH₃·l⁻¹ and 0.0011 mg P-PO₄·l⁻¹₋₋As for SO₄⁻⁻ analysis, snow and lake water samples were concentrated by heating, HClO₄ and HNO₃ mixture was added and evaporated, then an ion-exchange column was used to remove interferences of cations. Samples were titrated with BaCl₂ solution in the presence of nitrochromazo until blue color appearance (Kalacheva et al., 2002). Finally, total nitrogen (TN) was digested
- 285 from total snow and lake water samples using persulfate and boric acids and subsequently transformed into NO_3^- (Grasshoff et al., 1983). The natural light absorption of this nitrate at 210 nm was determined using a Spekol 1300 photometer (Analytik Jena, Germany) and corrected for organic matter interference by subtracting absorption at 275 nm (Slanina et al., 1976).

•(удалено: ; Радужное
•(удалено: ; Каровое
•(удалено: ; Светлое

удалено: Water and snow

удалено: n

удалено: Ammonium was determined by nesslerization in samples filtered through 0.45 µm membrane filters "Porafil" (Macherey-Nagel, Germany) with single-use plastic syringes. Detection limit for N-NH₄ ammonium was 0.014 011 mg⁻¹. The other dissolved chemical species (NO_2 , NO_3 , soluble reactive phosphorus –SRP– and SO_4^{2-7}) were analysed on gently vacuum filtered aliquots also using the same mentioned filters in 2015 and 2017, and. cellulose filters in 2011 and 2012 according to the Russian National Standards (Gladyshev et al., 2015; Tolomeev et al., 2014). . . .

2.3 Chlorophyll and SCPs analyses

A known fraction of melted snow samples was filtered through GF/C filters to collect SCPs (Mosello et al., 1997). Nitric, hydrofluoric and chlorhydric acids were used to remove organic, siliceous and carbonate material, respectively (Rose, 1994;

315 Yang et al., 2001). Determinate fractions of the samples were mounted on NAPHRAX and counted at 400X under an Axiostar plus microscope (Zeiss). Negative controls and a sediment reference standard were likewise processed to correct final counts for <u>eventual</u> experimental bias (Rose, 2008).

Phytoplankton chlorophyll was assessed according to the UNESCO standard protocols (VA, 1997). Samples had been filtered in the field laboratory through BaSO₄-covered 0.45 µm membrane filters "Porafil" (Macherey-Nagel, Germany),

320 folded inwards and frozen. They were then let thaw, dried in the dark, and scraped along with BaSO₄ into centrifuge tubes. Pigments were extracted in 100% acetone for 9h in the dark at +4°C. After filtration through 0.2 µm polycarbonate filters, MQ water was added to get pigments dissolved in a 90% acetone solution, f.c. Photometric measurements were used to calculate chlorophyll concentrations (Jeffrey and Humphrey, 1975).

2.4 Air mass retrotrajectory analysis, maps and statistics

- 325 The retrotrajectories of air masses flowing on the three snow sampling sites were obtained using the Hybrid Single-Particle Lagrangian Integrated Trajectories (HYSPLIT) model for archive trajectories (Rolph et al., 2017; Stein et al., 2015) of the National Oceanic and Atmospheric Administration Air Resources Laboratory (NOAA, USA). A total of 187 daily retrotrajectories embracing the sampled period of atmospheric deposition were reconstructed as the snowpack bulk deposition airshed. Each trajectory started three days back in the past. It recorded a per hour latitude, longitude and altitude
- 330 coordinates and ended up at the snow sampling coordinates, at 0 m above model ground level at 24h of consecutive days. All the analyses were performed within the R environment (R Development Core Team, 2017). Total retrotrajectory length and average wind speed the hour before getting to the sampling point were calculated using the Vincenty (ellipsoid) distance method within the geosphere package (Hijmans, 2017). The openair package was used to determine wind direction and to draw wind roses (Carslaw and Ropkins, 2012). The number of per hour coordinates at 0 m above model ground level was
- 335 calculated to characterise the direct interaction of each air mass with the Earth crust. Retrotrajectories were mapped using the ggmap package (Kahle and Wickham, 2013). One-way ANOVA comparing sites and Pearson correlation analyses of chemical and wind variables in the seven analysed snow core sections were performed using built-in functions of the R statistical environment.

Detailed watershed land cover/land use maps were manually defined for each lake. Polygons were defined using QGIS 340 3.14.16-Pi on the basis of Google Satellite and Open Street Map XYZ tiles at a resolution that distinguished single trees. Lake, whole watershed and different vegetation and land cover areas were calculated using ellipsoidal project. удалено: any

3 Results and discussion

345 3.1 Potential fragmentation of nutrients by snow melting

The three sampled snow cores were 115 cm deep, but had different <u>snow water equivalent (SWE)</u>: 25±1 cm in Tsirkovoe, 27±1 cm in Oiskoe and 12±0.3 cm in Tushkan. First of all, the snowpack temperature profile was measured to determine if snow melting could have occurred before sampling. Major snow thawing can be discarded in any of the 3 sampling sites because snow temperature was not around 0°C but always lower. Nevertheless, the deepest snowpack layers fall within the

350 range between -2 and 0°C: Oiskoe at 110 cm deep, Tsirkovoe from 90 to 110 cm, and namely Tushkan from 60 to 110 cm deep. This indicates that snow melt was either about to occur or could have even started in these particular layers, triggering a sequential elution of solutes (Mosello et al., 1997). In that hypothetical case, snowpack-based atmospheric deposition estimates would be biased. In order to discard such a case, solute concentrations in the upper and colder snow layers were compared to those in the deeper and warmer ones (fig. 3, table S3).

8

удалено: 1



	The hypothesis was that deeper and warmer layers, suspect of possible melting, would show lower solute concentration in
	case of important melting, preferentially in those solutes that elute firstly during snow melting. Yet, because the first
	centimetres of snowpack were formed much faster than the rest of the snowpack, it is conceivable that the deep layers were
365	originally poorer in airborne chemicals and particles, which would bother the initial hypothesis. Indeed, the first third of
	snowpack thickness at Olenya Rechka meteorological station deposited in only 19 days (from 10/01 to 10/19), whereas it
	took 39 days (from 10/01 to 11/08) to attain half of its thickness at sampling date (i.e. 187 days after initial snowpack
	formation). It is likely that the deepest Tsirkovoe, Oiskoe and Tushkan snow core segments (80-115 cm in the first case and
	60-115 cm in the others) would have formed in about 19 and 39 days, respectively. Nevertheless, if precipitation rate had, a
370	determinant effect on the vertical distribution of solutes and particles content, the lower values in deeper layers should be
	expectable in all the measured variables, and it was not the case (fig. 3).
	Thus, no significant differences were found between the upper and deeper layers in any of the measured variables except for
	nitrate, with lower values in the deep layers (ANOVA, p-value= 7.75×10-5). The other significant differences between upper
1	and deeper snow layers (TN in Tsirkovoe, SRP in Tushkan, TP in Oiskoe and SCPs in Tushkan; t-tests) were not consistent
375	across sampling sites (fig. 3). Besides, sulphate also had slightly lower concentrations in the deep snow layers but this
	difference was not statistically significant. This is especially explanatory because preferential elution of ions during snow
	melt occurs either in the sequence $SO_4^2 > NO_3 > NH_4^+$ (Kuhn, 2001) or $SO_4^2 > NH_4^+ > NO_3^-$ (Wang et al., 2018a), but sulphate
	always elutes preferentially to inorganic nitrogen species, according to the literature (Cragin et al., 1996; Kuhn, 2001;
	Stottlemyer and Rutkowski, 1990; Williams and Melack, 1991). In other words, higher proportions of sulphate are released
380	during early snow melting steps as compared to nitrate or ammonium. As a result, only significantly lower values of sulphate
1	should be observable in incipient thawing snow layers whereas both sulphate and nitrate would be significantly leaked at a
	more advanced thawing stage. Therefore, we suggest that the only observed differences in nitrate concentrations between

Figure 3 Chemical composition of upper (dark grey) and lower (light grey) layers of the 2016-17 snowpack in Ergaki mountains. All values are in µg[1¹ except SCPs (counts[1¹). Bars represent mean values and whiskers, standard deviation. The two upper

layers were averaged in Tsirkovoe, where the snow core was divided into three layers. Column pairs with "a" and "b" letters are

significantly different (one-way ANOVA, p-v<0.05 in the case of nitrate; t-tests in the other cases; n=3 except in Tsirkovoe, where

360

upper layer n=4 and lower layer n=2).

fact of being non-significant allows us to discard thawing as a cause, and entails sulphate load estimates would not be thaw-385 biased nor any of the other solutes, which should elute at a later stage. As a conclusion, snowpack-based estimates of atmospheric deposition should always be cautiously considered, but major elution of solutes due to snow melting was not detected in the present study, probably thanks to the consistently negative temperatures along almost the whole integrated time period.

удалено: удалено: •

удалено: had

удалено:

удалено: n'

10

layers might not be due to snow melting. Even if it is true that sulphate also tends to be lower at deep warm snow layers, the

3.2 Snow nutrients and pollutants composition

Nutrient concentrations in Ergaki snowpack (table 2, table S³) generally take intermediate positions in comparison with other snowpack studies around the world. For instance the average 191±35 μg NO₃-N₆I⁻¹ in Ergaki is comparable to an old record in the Pyrenees (115±106 μg NO₃-N₆I⁻¹, Catalan 1989) but lower than a later one in the same mountains (280 μg NO₃-N₆I⁻¹, Felip et al. 1995). It also takes an intermediate position relative to the Alps: lower than in Tyrolean Alps in the early 1990s (308 μg NO₃-N₆I⁻¹, Felip et al. 1995) but higher than most sampling points in the French Alps in a more recent study
(Dambrine et al., 2018). Finally, nitrate concentration in Ergaki snow was in between that of the Bothnian Bay of the Baltic sea (480 ±130 μg NO₃-N₆I⁻¹, Rahm et al. 1995) and the lake Tahoe basin in Sierra Nevada (14-138 μg NO₃-N₆I⁻¹, Pearson et al. 2015). Note that, paradoxically, the former is considered a low atmospheric nitrogen deposition region (Bergström and Jansson, 2006) whereas the latter has been reckoned as an airborne nutrient enriched area (Sickman et al., 2003) where atmospheric nitrogen deposition has shifted phytoplankton limitation from N and P colimitation to persistent P limitation

удалено: 1		
удалено:		
удалено: higher		
удалено: than		
удалено:		
удалено: bit more r	ecent	
удалено:		

		NH4-N	NO2-N	NO3-N	Т	Р	TP	SO4-S	3
					Ν	0			С
						4			Р
									s
						Р			
Average concentration	in	n.d.	n.d.	191	4	2	3.33	864	8
snow <u>(μg·l⁻¹)</u>				(34)	8		(2.42)	(106)	0
					3	5			5
					(5			
					1	(<u>S</u>
					6	2			<u>C</u>
					5				<u>P</u>
)	1			<u>s</u>
						3			2
)			<u>1</u>
									2
									<u>1</u>
									(
									2
									7
									5

11

отформатировано: без подчеркивания, Цвет шрифта: Авто, подстрочные Отформатированная таблица отформатировано: без подчеркивания, Цвет шрифта: Авто, подстрочные отформатировано: без подчеркивания, Цвет шрифта: Авто, подстрочные отформатировано: без подчеркивания, Цвет шрифта: Авто, подстрочные отформатировано: без подчеркивания, Цвет шрифта: Авто, подстрочные

Отформатировано: По левому краю

)		
Half year deposition	n.d.	n.d.	40	9 0 0.58	190	ŀ	Отформатировано: По левому краю	
<u>(mg·m⁻²)</u>			(16)	7 . (0.13)	(91)	5		
				4		9		
				(3				
				5 (×		
				6 0				
) .		1		
				1		<u>0</u>		
				5		3		
)		<u>s</u>		
						<u>C</u>		
						<u>P</u>		
						<u>s</u>		
						±		
						<u>m</u>		
						=		
						2		
						(
						4		
						8		
)		
Deposition rate (~time)	n.d.	n.d.	79	1 0 1.13	372	3	Отформатировано: По левому краю	
$(mg \cdot m^{-2} \cdot y^{-2})$			(47)	9 . (0.60)	(236)	1		
				1 8		2		
				(4				
				1 (×		
				3 0				
				2 .		<u>1</u>		
) 4		<u>0</u>		
				8		<u>3</u>		
)		<u>S</u>		

						C P s : m 2 : Y : Y (1 (1 7 4)	
Deposition (~precipitation) (mg·m ⁻² ·y ⁻²)	rate n.d.	n.d.	119 (71)	2 1 1.71 8 . (0.91) 8 2 (6 1 (9 0 8 .) 7 3)	560 (356)	4 7 0 × 1 0 3 S C P 8 ∴ m 2	Отформатировано: По левому краю

				1	
				Y	
				÷	
				1	
				(
				2	
				6	
				2	
				2	
		100	0.000)	-
Modelled	For year	<u>c. 100</u>	$\underline{0}$ <u>c. 2 (Mahowald et</u>	100-200	
deposition rates	2000	(Lamarque et	<u>. al., 2008)</u>	(Lamarque et	отформатировано: Английский (США)
$(\text{mg} \cdot \text{m}^{-2} \cdot \text{y}^{-2})$		al., 2013)	1	al., 2013)	отформатировано: Английский (США)
			Ξ		отформатировано: Шрифт: не полужирный, Цвет шрифта: Авто, Английский (США)
			<u>0</u>		отформатировано: Шрифт: не полужирный, Цвет шрифта: Авто, Английский (США)
			4 		отформатировано: Испанский
			<u>5</u>		отформатировано: Английский (США)
			(
			М		
			a		
			h		
			0		
			W		
			a		
			1		
			d		
			e		
			t		
			a 1		
			1		
			,		
		14			

	8				
1)			отформатировано: Испанский	
	For year	<u>50-200</u>	<u>50-200</u>		
	<u>2030</u>	(Lamarque et	(Lamarque et		
		<u>al., 2013)</u>	<u>al., 2013)</u>		
ļ	Table 2. Average concentrations, half year depositions and	l estimated yearly deposition rates	as averaged by the 3 sampled sites		
	(μg·l ⁻¹ , mg·m ⁻² and mg·m ⁻² ·y ⁻¹ , respectively) (SCPs in cours	nts;1 ⁻¹ , 10 ³ counts;m ⁻² and 10 ³ count red. Vearly deposition rates were	ts; m ⁻² ; y ⁻¹). Mean values are shown,	~	удалено: ·
	winter depositions and either assuming a constant deposit	ion rate (time weighted estimate,	row 3) or a precipitation-dependent	M	удалено: ·
42	20 <u>deposition rate (precipitation-weighted estimate, row 4). Se</u>	e section 3.3 for further discussion.			удалено: ·
1	Total nitrogen and total phosphorus in Ergaki snowpack	(table 2) were higher than in the	first mentioned Pyrenean study but		удалено: ·
ĺ.	lower than in the Baltic: 194 \pm 135 and 1054 \pm 363 µg TI	N-N-1-1, and 2.38±0.59 and 9.3±5	.1 μg TP-P ₂ 1 ⁻¹ , respectively. Total		удалено:
	phosphorus concentration also was within the lowest rar	ge of that measured around lake	Tahoe (3-109 µg TP-P-1-1, Pearson	11 1	удалено:
I	et al. 2015). Nevertheless, ammonium and nitrite were u	ndetectable in the present study b	ut detected in most of the previous	177	удалено:
42	25 studies in the snowpack (e.g. Catalan 1989: Pearson et a	l. 2015). Ammonium was also de	tected in the snow surrounding the	$\langle \rangle \rangle$	удалено:
I	city of Krasnovarsk by our own lab using the same an	alvtical method as here (unpubl)	It is very likely that ammonium	- \}	удалено:
	concentrations in the present study were under the deter	ation limit as nitrate values were	more than five-fold lower than in	Ì	удалено: -
ī	the Kreeneversk eity enew complex, where amonium	had haan dataatad. Finally, nutria	nt bioavailability is an attribute of		
	the Krashoyarsk city show samples, where animonium	had been detected. Finally, hutrie		1	УЛЭЛОНО: 7
	the Ergaki snowpack as $73\pm12\%$ IP was in the form of p	hosphate and about $44\pm19\%$ TN	vas nitrate.		удалено: ?
43	30 Besides ammonium and nitrite, sulphate concentration	s in Ergaki snowpack were also	unusual. Sulphate was the most		VIII VIII VIII VIII VIII VIII VIII VII
	abundant of the measured ions. It doubled that in the Py	renees in the late eighties (401±10	06 μg SO ₄ -S ₂ 1 ⁻¹ , Catalan 1989) and		удалено:
-	quadrupled that on lake Tahoe (Pearson et al., 2015). S	Sulphate concentration in Ergaki	snowpack was only similar to the	Å	удалено:
	highest values in the literature for non-urban areas, such	as on the south coast of lake Su	perior in the eighties (828 \pm 216 µg	//	отформатировано: Английский (США)
	SO4-S ₄ 1 ⁻¹ in average, Stottlemyer and Rutkowski 199	00). Altogether, nitrogen and pl	nosphorus concentrations reached	//	отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто
43	35 intermediate-low values but sulphate concentration was r	emarkably high in Ergaki snowpa	ck.	/)	удалено: as well as
	The organic nitrogen in Ergaki snowpack represented a 5	6±19% of the total nitrogen, which	h is a high but reasonable value as		отформатировано: Шрифт: не полужирный, без
	compared to the literature. The relative share of organic	to total nitrogen in the snowpack	s of the Pyrenees, Alps and Sierra		подчеркивания, цвет шрифта: Авто
	Nevada (USA) mountains, and on the Baltic sea were 1	0±9%, 41±13%, 49±17% and 21±	12%, respectively (Catalan, 1989;		подчеркивания, Цвет шрифта: Авто
44	Clement et al., 2012; Pearson et al., 2015; Rahm et al., 2015; Rahm et al., 2010; Snowpack records than in wet deposition because dry	(1995), Organic nitrogen has been deposition of organic nitrogen i	reported elsewhere to be higher in s integrated in the snowpack and		отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто
ſ	because microbial uptake and assimilation of inorganic r	hitrogen might occur in the snowp	ack (Clement et al., 2012; Pearson	_	отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто
	et al., 2015). In the case of microbial conversion from in	organic to organic nitrogen, it cou	ld be hypothesised that deeper and		отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто
		15			

460 older snow layers should have higher organic nitrogen shares. Such a pattern was only observed at Tushkan (ANOVA, p-value= 0.0178) whereas no significant differences were found in the percentatge of organic nitrogen between upper and lower snow layers at Oiskoe and Tsirkovoe sites, We, therefore, hypothesise that a different combination of phenomena might be responsible for organic nitrogen dominance at different sites,

3.3 Atmospheric deposition load

- 465 Roughly half year cumulative deposition corresponding to the snow season -187 days- is summarized in table 2 (second row; see table S3 for original measurements). Unfortunately, snow-free season depositions were not measured in the present study and, consequently, yearly deposition rates could not be determined. Nevertheless, preliminary estimations were conducted assuming either a constant deposition rate along the year –time-weighted estimate– or a precipitation-weighted deposition rate (table 2, 3rd and 4th rows, respectively). These assumptions entail different simplifications concerning the seasonal
- 470 pattern of emission, transport and deposition of the different chemical species in this particular part of the world. The precipitation-weighted estimate should be, *a priori*, a more accurate estimate because wet deposition is known to be the main contributor to total deposition<u>in wet climates like that in West Sayan mountains</u>. Indeed, the accumulated precipitation registered in Olenya Rechka meteorological station during the studied snow season was 419 mm, whereas almost the double (819 mm) were registered during the following months up to complete a year. Note that the 2016-17 seasonality was <u>slightly</u>
- 475 more prominent than the median 2005-2019 seasonal precipitation (fig. 2 a). Nevertheless, both estimations neglect the emission seasonality, which might turn the constant deposition estimate into the most credible <u>option in some cases</u>. To evaluate our different estimates, we compared them to seasonal depositions in the literature and discussed their likely seasonal emissions.

The precipitation-dependent estimate of atmospheric phosphorus deposition (table 2, 4th row) was probably the closest to the actual yearly load, although it might be an underestimate. That is because two thirds of the yearly precipitation and biogenic aerosols -mainly pollen- co-occur during the snow-free season and pollen represents an important share of atmospheric phosphorus sources in low-atmospheric-phosphorus taiga landscapes like Ergaki (Banks and Nighswander, 2000; Doskey and Ugoagwu, 1992; Mahowald et al., 2008; Wang et al., 2015). In order to evaluate the magnitude of our underestimation, we checked similar studies with seasonal resolution in cold forest landscapes including the lake of Bays (ON, Canada)

- 485 (Eimers et al., 2018), lake Simcoe (ON, Canada) (Brown et al., 2011), and a Tibetan forest (Wang et al., 2018b). Phosphorus Joads <u>during the snow period</u> were about 9, 11 and 18 times larger than in Ergaki, snow-free season atmospheric phosphorus depositions, 2.4, 5 and 7.4 times higher than in the snow season, and snow-free season precipitations, 0.88, 0.91 and 9.1 times that of snow season, respectively. <u>Thus, in the hypothetical case that phosphorus deposition seasonality strictly</u> depended on the yearly phosphorus load, the seasonal factor for Ergaki should be much lower than 2.4. On the other hand, if
- 490 <u>it</u> depended on precipitation seasonality <u>alone</u>, Ergaki precipitation seasonality (1.95) would correspond to a phosphorus deposition seasonality of about 5.5. If we applied <u>these factors (2.4 and 5.5</u>) to estimate snow-free season phosphorus deposition and added it to measured snow season deposition, <u>the</u> yearly deposition rates they would be: 1.972 mg TP-P; m⁻

отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто

удалено: n

отформатировано: Шрифт: не полужирный, без подчеркивания, Цвет шрифта: Авто

отформатировано: Шрифт: не полужирный

(удалено:

удалено: 1

удалено: a bit

удалено: 1

удалено: one

удалено: 🕤

отформатировано: без подчеркивания, Цвет шрифта: Авто, надстрочные

YRANEHO: As a rule of thumb, weak seasonality is observed for chemical species with low deposition loads. This was clearly the case of atmospheric phosphorus deposition. Atmospheric phosphorus depositions are particularly low in taiga landscapes like Ergaki, where spring and summer biogenic aerosols – mainly pollen– represent the largest share of atmospheric phosphorus sources

Yganeno: Additionally, our study site has a much higher precipitation during the snow-free season. The co-ocurrence of biogenic acrosols and almost two thirds of the precipitation during the snow-free season implies that even our precipitation-dependent estimates (table 2, 4th row) might be underestimates, as biogenic acrosols are not taken into account. In order to evaluate the

(удалено:

удалено: and with seasonal resolution were checked. Although we didn't find any study with seasonal resolution and snow season atmospheric phosphorus loads as low as those measured in Ergaki, some clues were given by a bunch of sampling sites around ...

удалено: snow period

yganeho: In conclusion, the magnitude of snow season phosphorus load, and the seasonalities of precipitation and phosphorus deposition were positively correlated, and the snow-free to snow season phosphorus deposition factor could be guessed, in principle, for Ergaki. Nevertheless, the available data is too scarce to make any formal prediction based on a multiple nonlinear regression. If...

удалено: phosphorus deposition seasonality just

YRANEHO: The latter factor (5.5) would definitely provide an overestimation of snow-free season phosphorus deposition, whereas it is not certain if the former one (2.4) would either over- or underestimate it ...

удалено: 2.4 and 5.5

удалено: in order to calculate

удалено:

² y² and 1.462 mg PO4-P₂ m² y², and 3.77 mg TP-P₂ m² y² and 2.795 mg PO4-P₂ m² y², respectively. <u>The latter factor</u> (5.5) would definitely provide an overestimation of snow-free season phosphorus deposition. The former one (2.4) resulted in slightly higher values than our precipitation-dependent estimate and it is not certain if that would either over- or

- 535 underestimate the actual value The best atmospheric nitrate deposition estimate was the precipitation-dependent estimate (table 2, 4th row), according to its seasonal distribution in other natural forested sites in the literature. The Ergaki autumn-winter nitrate deposition was four and five times lower than in a Chinese "background" and a south Bohemian forest site, respectively (Kopáček et al. 2011b; Xu et al. 2018). Nevertheless, HNO3 and total inorganic nitrogen species atmospheric concentrations were stable along the
- 540 year in the Chinese background site, and only particulate NO3 was slightly higher in autumn-winter than spring-summer. In the south Bohemian forest, the April to September precipitation and nitrate deposition were only 15% and 5% higher than in October-March, respectively. Therefore, atmospheric nitrogen concentration was also fairly stable at this site, with just slightly higher atmospheric nitrate concentrations in the autumn-winter semester that would be mainly counterbalanced by a lower precipitation. In the light of these observations, it is likely that atmospheric nitrate concentration in Ergaki during the
- 545 snow season, was either similar to or slightly higher than in the snow-free season. In case of invariable atmosphere nitrate concentrations along the year, higher precipitation during the snow-free season would trigger also a higher nitrate deposition, and our precipitation-dependent estimate (table 2, 4th row) would be our best estimate. In the hypothetical case where the proportion between snow and snow-free season atmospheric nitrate concentrations was the same as in the Chinese site (1.24), and taking into account the higher snow-free season precipitation in Ergaki (1.95), the resulting snow-free nitrate
- 550 deposition would be about 1.6 times higher than that measured in the snow season and the yearly load would be about 104±62 mg NO₃-N₂m⁻²y⁻¹. In conclusion, the actual yearly nitrate deposition should be somewhere in-between the timedependent and the precipitation-dependent estimates, but closer to the latter estimate,

Time-based sulphate atmospheric deposition estimate was better than the precipitation-dependent estimate. Winter atmospheric sulphate depositions were higher than in summer both in the pristine Canadian Rocky Mountains (2-5 times)

- 555 and in a Japanese site receiving sulphate from Chinese coal combustion (3.5-4 times) (Ohizumi et al., 2016; Wasiuta et al., 2015). Unexpectedly high seasonality in the pristine location was due to much higher precipitation in winter but the atmospheric sulphate concentration was relatively constant along the year. On the other hand, sulphate deposition seasonality in the polluted site was attributed to higher coal burning and emissions in winter. As winter atmospheric sulphate deposition in Ergaki (190±91 mg SO4-S·m²) was about four times higher than the background values in the Canadian Rocky mountains
- 560 (\leq 50 mg SO₄-S·m²) and 11-12 times lower than in a Japanese site, it is likely that winter atmospheric sulphate concentrations in Ergaki were also higher than in summer. In this case, the precipitation-weighted yearly deposition estimate would be an overestimation but we cannot rigorously determine if the actual value would Jay above or below the timeweighted estimate. At most, we could assume a linear relationship between yearly sulphate deposition load and its seasonality. Then, a seasonality 11-12 times lower than in the Japanese site form the literature would imply a yearly 565 deposition load of 251 mg SO4-Scm² in Ergaki. Accordingly, our time-dependent estimate would be our best estimate but

	удалено.				
	удалено:				
$\langle \rangle$	удалено: ·				
$\langle \rangle$	удалено:				
()	удалено: ·				
1	удалено:				
$\langle \rangle$	удалено:				
	удалено: For all the above-mentioned reasons, it is very likely that the actual yearly load in Ergaki was safely below the latter estimates, whereas the former ones are not far above the precipitation-dependent estimates (table 2, 4 th row) and, possibly [11]				
\mathbb{N}	удалено: ¶ [12]				
	удалено: As for atmospheric nitrate deposition, no significantly 13				
	перемещено вниз [1]: (Кора́ček et al. 2011b; Xu et al. 2018)				
	удалено: A simplistic conclusion would be to think that this [14]				
$\langle \rangle$	перемещено (вставка) [1]				
//	удалено: А				
()	удалено: in the Bohemian forest				
()	удалено: , which denotes a				
$\langle \rangle$	удалено: fairly stable				
()	удалено: Similarly, stable atmospheric HNO3 and total inorgani5]				
1	удалено: ,				
)	удалено: ,				
	удалено: that				
2	удалено: in the				
$\left\langle \right\rangle$	удалено: ·				
//	удалено: ·				
$\langle \rangle$	удалено: it is reasonable to think that				
$\langle \rangle \rangle$	удалено: was				
	удалено: . Indeed, in case of a nitrate emissions seasonality as [h6]				
	удалено: precipitation-dependent				
	удалено: would be closer to the actual value				
	удалено: In the case of sulphate, w				
	удалено: in Ergaki (190±91 mg SO ₄ -S·m ⁻²), was about four times				
1	удалено: can be				
	удалено: According to these observations, an area with an [18]				
Ç	удалено: to have a somewhat higher				
	удалено: than summer				
	удалено: be				
1	удалено: orientatively				
	удалено:				

still an overestimate. In conclusion, for the sake of a simpler discussion, we will only consider the time-weighted estimate of yearly sulphate deposition and precipitation-weighted estimates of phosphorus and nitrate depositions. Nevertheless, these estimates must be interpreted cautiously: Whereas nitrate and sulphate deposition estimates might be slightly overestimated,

685 phosphorus would be underestimated. The selected yearly deposition rate estimates (table 2, 3rd and 4th rows) were compared to global model predictions from the literature (table 2 5th row). A global deposition model predicted c. 100 mg NO₃-N₂m², y⁻¹ and 100-200 mg SO₄-S m², y⁻¹ удалено: ··m⁻²·...y⁻¹ and 100-200 mg SO₄-S ·...m⁻²·...y⁻¹ loads рад loads on West Savan mountains for year 2000, whereas it forecasted ranges from 50-200 mg NO₃-N₂-y⁻¹ and 50-200 mg отформатировано: без подчеркивания, Цвет шрифта: Авто, надстрочные SO4-Scm²y¹ in 2030, according to different scenarios (Lamarque et al., 2013). The EMEP wet deposition model reported similar values (50-100 mg oxidized N m²-y¹ and 50-200 mg oxidized S m²-y¹, including SO₂ and SO₄, and HNO₃, HONO 690 and NO₃, respectively) for year 2017 on the limits of their geographical prediction, at a location about 200 Km west of our sampling site (Fagerli et al., 2019; Gauss et al., 2019). Therefore, our 2016-17 nitrate deposition estimate roughly fitted the models, whereas sulphate deposition was clearly higher than expected. In the case of phosphorus deposition, our TP estimate was slightly lower than predicted (c. 2 mg TP-P_e $m^{-2}y^{-1}$), although uncertainties linked to pollen contribution could make the 695 actual TP-P deposition match or even surpass the modelled values. On the other hand, the phosphate fraction would be clearly higher than expected (0.1-0.5 mg PO₄-P₂m⁻²y⁻¹, Mahowald et al. 2008). In conclusion, the atmospheric nitrate deposition in Ergaki mountain ridge is at the very low range and is between 5 to 20 times lower than in polluted areas of the world. Nevertheless, it is clearly above the background deposition of 0-50 mg NO₃-Nm⁻².y⁻¹, as it used to be the case of most Siberia in 1850 or the Antarctica and unpolluted parts of the oceans in 2000 удалено: ··m⁻²·...y⁻¹, as it used to be the case of most Siberia ippon 700 (Lamarque et al., 2013). As for total phosphorus deposition, the uncertainty linked to non-measured spring-summer biogenic and wildfire contributions, makes it hard to position the studied site within a World ranking. Our estimate excluding these important biogenic and wildfire contributions (1.71 mg TP-Pt m²ty²) and our literature-based estimate including them $(1.972 \text{ mg TP-P}_{e} \text{ m}^{2} \text{y}^{2})$ would be lower than any terrestrial measurement, according to a worldwide review ($\geq 3 \text{ mg TP-P}_{e}$ $m^2 y^2$) (Tipping et al., 2014). Nevertheless, it is also possible that pollen and wildfires accounted for a larger contribution 705 and that the present study site exceeded the latter value. In any case, atmospheric phosphorus deposition in Ergaki would be above the background values corresponding to the poles and the oceans ($\leq 1-2 \text{ mg TP-P}, \text{m}^2, y^2$) and $\leq 0.5 \text{ mg PO}_4$ -P, m^2, y^2), excluding the Atlantic strip downwind of the Sahara (Mahowald et al., 2008). Finally, our yearly sulphate deposition estimate should be cautiously considered, as it could be overestimated due to regionally widespread coal combustion for **удалено:** expectably lower deposition during summer heating during winter (see section 3.5). In any case, it would positively exceed the background values of 0-50 and 50-100 mg 510 SO4m²y¹ typical in the polar areas and southern hemisphere oceans, respectively (Lamarque et al., 2013). удалено: ··m⁻²· ... [21] 3.4 SCPs deposition rate

[22]

The calculated SCPs deposition rate in Ergaki Natural Park ($312\pm174\chi10^3$ SCPs;m²;y⁻¹) was high above the background rates recorded in Baikal middle basin ($57\chi10^3$ SCPs;m²;y⁻¹), Svalbard islands ($13\chi10^3$ SCPs;m²;y⁻¹) and Nevada Rocky mountains ($1.3\pm0.8\chi10^3$ SCPs;m²;y⁻¹) (Reinemann et al., 2014; Rose et al., 1998). Indeed, it is also far below the records in

more polluted areas such as the case of lake Paione Superiore in western Alps as it used to be during the more polluted times of 1980s and early 1990s (40900×10^3 SCPs; m²; y⁻¹) or a set of north African lakes ($1098-23694 \times 10^3$ SCPs; m²; y⁻¹), where production of electricity by thermal means has increased in the last years (Rose et al., 1999b, 2003). In comparison to a couple of lakes sampled in 1992 in the Khamar-Daban mountains (Southern Siberia) (262 and 780×10^3 SCPs; m²; y⁻¹), the

- SCP deposition rate in Ergaki was more similar to the lake that was relatively farther from Irkutsk <u>city</u> pollution source (Rose et al., 1998). Our data also falls in the lower range of Tatra mountains (225-5240×10³ SCPs_xm²_xy⁻¹) and the Pyrenees (229-630×10³ SCPs_xm²_xy⁻¹) in the mid-1990s (Rose et al., 1998, 1999a; Šporka et al., 2002). An interesting and paradoxical case to compare with is lake Grânästjärn in 1980, with a very similar SCP deposition (300×10³ SCPs_xm²_xy⁻¹) but sulphate and nitrate deposition rates 2.6 and 1.8 times higher than in Ergaki (Bergström et al., 2005; Wik and Renberg, 1996). This may lead to think that our SCP, sulphate and nitrate data didn't match. Nevertheless, at least sulphate depositions differing up to c. 40% have been observed at a particulate low SCP deposition rate (Rose and Monteith, 2005). Additionally, sulphate measurements in Sweden might include a higher percentage of marine sulphate than in the heart of Eurasia. Finally, apart
- 780 **3.5 Spatial distribution and origin of atmospheric depositions**

The spatial distribution of deposited chemical species showed two different patterns. On the one hand, phosphorus forms and SCPs showed even distribution between sampling sites. On the other hand, NO₃, TN and SO₄ depositions were significantly higher on Tsirkovoe and Oiskoe than on Tushkan (fig. 4). Even distribution of phosphorus deposition suggests a common source of atmospheric phosphorus for all Ergaki sites. Under very low atmospheric phosphorus deposition, like in Ergaki

from this single Swedish lake where the proportion of nitrate to SCPs differs so much from ours, both SCP and nitrate

deposition measured in this study are generally comparable to the low range of values in the literature.

- 785 mountain ridge, biogenic and combustion origins are more important than mineral (Mahowald et al., 2008). This is supported by several evidences in our case. Firstly, even if the predominant western air mass retrotrajectories partly traverse Kazakhstan steppe a percentage of days, they hardly ever cross central Asian deserts (fig. 5 a and b). Moreover, direct contact between air mass retrotrajectories and the Earth crust occurs more often in the taiga ecoregion (yellow dots, fig. 5 a and b). Finally, the percentage of TP which is soluble in our snow samples (79%) is comparable to European aerosols (50-
- 790 100%), but much higher than Saharan dust (8-25%) (Mahowald et al., 2008), so it suggests that phosphorus aerosols in Ergaki were not of desert origin. Thus, Ergaki mountain lakes differ from Baikal lake, which can be influenced by dust originating at the Gobi desert (Jambers and Van Grieken, 1997).

удалено	×
удалено	
удалено	
удалено	×
удалено	- -
удалено	
удалено	
удалено	• ×
удалено	
удалено	
удалено	×
удалено	. ·
удалено	: ·
удалено	:
удалено	×
удалено	: ·
удалено	· ·
удалено	×
удалено	· ·
удалено	



815 Figure 4. Half year deposition of some airborne chemical species along with snow at 3 different sites in Ergaki mountain ridge. Bars are average values and whiskers represent standard deviation. Sites with different letters on error bars belong to different groups defined by post-hoc Tukey HSD analysis (one-way ANOVA p-values<0.01).



Figure 5. Daily three-day long air mass retrotrajectories flowing onto lake Oiskoe from 1st October 2016 to 5th April 2017 (a). Only retrotrajectories of air masses causing precipitation onto Oiskoe at different zooms (b, c and e) and to Tsirkovoe (d) and Tushkan (f). Yellow dots represent hourly records where the air mass retrotrajectories contacted the Earth crust. Water bodies are in black. Grey lines are either political borders or the road. Sources: Map tiles by <u>Stamen Design</u>, under <u>CC BY 3.0</u>. Data by <u>OpenStreetMap</u>, under <u>ODbL</u>; Airmass retrotrajectories by http://www.ready.noaa.gov.

Nitrate and sulphate are tracers of fossil fuel or biomass combustion (Mahowald et al., 2008). Their spatial correlation (lower

- 825 in Tushkan than in the other two sites, fig. 4) suggests that alternative nitrate origins like chemical fertilizers or secondary transformations like nitrification/denitrification by microbes in the snowpack might be unimportant. In the same vein, nitrate and sulphate had a relatively high, although not significant, Pearson correlation coefficient (0.63) in the seven analysed snow core sections. The reason for lower nitrate and sulphate values in Tushkan is still unclear. This site is a forest glade located at a 300 m Jower altitude than the other two. We argue that this altitude difference is too limited to trigger any differences in
- 830 nitrate deposition and, in any case, the higher the site, the lower the expected deposition (Dambrine et al., 2018). Nonetheless, shorter distance between Tsirkovoe and Oiskoe and their wind regimes might be more decisive, even if the former is located on the north face and the latter on the south <u>face</u>. The wind speeds there were 3.4 m_es⁻¹ and 3.3 m_es⁻¹, respectively, whereas in Tushkan, farther from the ridge, it was 3.0 m_es⁻¹. Despite wind rose circular correlation between Tsirkovoe and Oiskoe was not significant, their local air mass retrotrajectories were quite more similar than to Tushkan (fig.
- 5 d, e and f). Perhaps, the two more northern sites might be also more exposed to the dominant snow-forming westerlies that flow between latitudes 52° and 53° and above (fig. 5 c). As for SCPs, which are also originated by combustion, the nitrate and sulphate spatial distribution is not followed. We speculate that the particulate character of SCPs might impose different atmospheric transport properties, so even if SCPs are good tracers of air pollution, slight mismatches between SCPs and chemical pollutants might occur (Rose and Ruppel, 2015; Wik and Renberg, 1996). Anyway, a trend to higher values in the 840 northernmost sites is also observed (fig. 4).

Finally, local wind speed differences between sites might be a tracer of air mass origin. Generally, faster winds would be capable to deliver chemical species from longer distances. The average retrotrajectory length had a weak, non significant, but still positive Pearson correlation with nitrate (0.47) and negative with phosphate (-0.48), TP (-0.25) and TN (-0.20). This suggests that nitrate deposited on Ergaki mountains might originate at farther distances than phosphorus and particulate

- 845 nitrogen. To sum up, we hypothesise that northern cities might contribute more to the nitrate deposition than southern ones. These combustion-produced chemicals would be uploaded to the northernmost half of the dominant westerlies conveyor belt flowing directly onto Ergaki mountains, rather than on the southernmost half of the westerlies flow, which mainly traverse Kazakhstan and turn northwards to Ergaki mountain ridge just before reaching the city of Kyzyl (fig. 5 c). These urban sources might not only include those at the local scale (Abakan, Minusinsk, Chernogorsk, rather than Kyzyl in the south) but
- also those at a regional one (central and eastern south Siberian rather than northern Kazakhstan cities).

удалено: altitude

удалено:	
удалено:	
удалено:	

855 3.6 Nutrient deposition and lake water stoichiometry

The molar stoichiometry of volume weighted mean concentrations in the snowpack, i.e., the concentrations that would be measured if we had sampled the whole snow core at once, were 169±76 NO3-N/TP-P, 251±134 NO3-N/PO4-P, and 486±357 TN-N/TP-P (mol/mol). The molar stoichiometry of estimated yearly atmospheric nutrient deposition in Ergaki were 169±105 and 128±79 NO3-N/TP-P (mol/mol) according to the precipitation-dependent yearly load estimation and to

- 860 the abovementioned literature-based corrections, respectively. As no other DIN form but nitrate was detected in our samples, NO3-N/TP-P ratio can be compared to DIN-N/TP-P ratios in the literature. The latter ratio has been previously used to assess nitrogen and phosphorus bioavailability both from deposition and in mountain lake water (Brahney et al., 2015). Total phosphorus was used instead of PO4 because: (1) otherwise, the fast biological uptake of deposited PO4 would lead to underestimate the actual atmospheric load, and (2) organic phosphorus sources are bioavailable to many mountain lake
- 865 phytoplankton species via extracellular enzyme activity, phagotrophy and osmotrophy (Diaz-de-Quijano, 2014; Medina-Sanchez et al., 2004). Thus, our estimated NO3-N/TP-P deposition molar ratios in Ergaki would belong to the higher quartile, as compared to a set of alpine regions of the world (Brahney et al., 2015) (fig. 6 a). The closest nutrient deposition stoichiometries to that of Ergaki (precipitation-dependent estimates) were recorded at the Sant Nicolau valley in the Pyrenees during the second half of the 1990s (170.5 DIN-N/TP-P, molar) and the Tatra mountains (165.8; calculation based on DIN
- 870 wet deposition during 1990-94 and average TP wet deposition during 1998-2009) (Brahney et al., 2015; Kopáček et al., 2000, 2011b; Ventura et al., 2000). In the case of the alternative 128±79 NO3-N/TP-P (mol/mol) deposition estimate for Ergaki, southern Sweden lake district was the closest value (125.6). In any case, Ergaki atmospheric nutrient deposition stoichiometry was by far larger than that of northern Sweden (20.1 molar DIN-N/TP-P), the paradigm of pristine areas with very low anthropogenic atmospheric nitrogen deposition and naturally nitrogen limited lakes. This was due to the
- 875 <u>extraordinary low phosphorus deposition as compared to nitrogen, which was also relatively low but not that much (see section 3.3).</u>

The molar stoichiometry of Ergaki lakes water was on average above the Redfield ratio (30.7±34.2 DIN-N/TP-P, fig. 6 a, table S2), which suggests lake phytoplankton would generally be P limited. Nevertheless, values below the Redfield ratio (16:1), i.e., corresponding *sensu lato* to N limitation, were also frequently observed. All the lakes had DIN-N/TP-P ratios

- both above and below the Redfield ratio at different surveys with the exception of lake Tsirkovoe, which was always above the Redfield ratio (P limitation) (table 3). Most lakes had DIN-N/TP-P ratios below the Redfield ratio in June and August 2011 and early September 2015, and above it in June and August 2012 and late August 2017. Therefore, no temporal trend can be drawn from our lake water chemistry data set (see table 3). Besides to the Redfield criterion, a previous study determined that DIN-N/TP-P molar ratios of 3.3, 4.9 and 7.5 would correspond to 75%, 50% and 25% probabilities of chlorophyll increase under a N enrichment experiment (Bergström, 2010). According to this more restrictive criterion, only lake Oiskoe would approach the 75% probability of having a positive increase of chlorophyll under N enrichment in the
 - 24

2015 survey. Nevertheless, it is very likely that this case was due to watershed-level processes. Forest and peat cover (25%

	and 6%, respectively) in Oiskoe watershed are more important than in other lakes watersheds and, more importantly, the lake	
	receives an inflow passing by nearby houses (table 1 and S1). A local phosphorus input from nearby houses could have	
890	occurred at Oiskoe in 2015 pointing to mesotrophic status of the lake. The high TP values could be also partly due to having	
	calculated TP as the sum of SRP and PP instead of measuring it directly in 2015 but only lake Oiskoe recorded	
	unprecedentedly high values, which might be due to the mentioned human activities in its watershed. In conclusion, lake	
	water nutrient stoichiometry suggests a variable, but slightly P-dominated, nutrient limitation regime	отфор
	In conclusion, a decoupling between atmospheric deposition and lake water nutrient ratio was observed in Ergaki as	
895	compared to many other mountain regions of the world (fig. 6 a). We argue that this was due to the extraordinarily low	
	absolute values of nutrient, and especially phosphorus, atmospheric deposition on Ergaki mountain ridge. Accordingly, the	
	sum of other biogeochemical processes occurring at the watershed soils, run-off, rivers, water column, sediments, etc. would	

be sufficient to modify the original stoichiometry of air-borne nutrients.

отформатировано: Английский (США)

отформатировано: Английский (Соединенное Королевство)

Отформатировано: Обычный



удалено: The relationship between atmospheric deposition and lake water concentration of nutrients is modulated by an array of processes occurring in the watershed soils, run-off, rivers, water column, sediments, etc. A non-exhaustive list would include: nitrification and denitrification in streams and wetlands, phosphorus sorption by sediments and soils, bedrock weathering, deposition of particulate nutrients on lake sediments and redissolution of nutrients back to the water column, mineralization of organic nutrients or avoidance of such mineralization via the mycoloop, etc. (Chróst and Siuda, 2002; Grossart et al., 2016; Schlesinger, 2000). In this section and the following one, we consider all these processes as a black box and focus only on the eventual relationship between atmospheric nutrient deposition and phytoplankton biomass and/or nutrient chemistry in lake water, as other studies did before (Bergström and Jansson, 2006; Bergström et al., 2005; Elser et al., 2009). The molar stoichiometry of precipitation-dependent yearly deposition estimations on Ergaki mountain ridge were 169±105 NO3-N/TP-P, 251±167 NO3-N/PO4-P, and 424±310 TN-N/TP-P (mol/mol). Nevertheless, as already mentioned, the precipitation dependent nitrate estimation might be overestimated and the phosphorus ones are definitely underestimated. If we took the abovementioned speculative guesses based on our autumn-winter measurements and its relationship to spring-summer measurements found in the literature, the molar ratios would be 128±79 NO3-N/TP-P and 190±126 NO3-N/PO4-P (mol/mol). Of course, any discussion based on these estimates is preliminary and needs to be contrasted with future year-long measurements of atmospheric nutrient deposition. including snow-free season too. Nevertheless, in both speculative cases, the NO3-N/TP-P deposition molar ratio in Ergaki would belong to the higher quartile, as compared to a set of alpine regions of the world (Brahney et al., 2015) (fig. 6 a). Briefly, it would be due to the extraordinary low phosphorus deposition as compared to nitrogen, which was also relatively low but not that much. As no other DIN form but nitrate was detected in our samples NO3-N/TP-P ratio can be compared to DIN-N/TP-P ratios in the literature. The closest nutrient deposition stoichiometries to that of Ergaki (precipitation-dependent estimates) were at Sant Nicolau valley in the Pyrenees (170,5 DIN-N/TP-P, molar) and the Tatra mountains (165.8) (Brahney et al., 2015; Kopáček et al., 2000, 2011b; Ventura et al., 2000). The Tatra and Pyrenees lakes used to be P-limited at the turn of the millennium, when they had these atmospheric nutrient deposition stoichiometries. However, a more comprehensive metaanalysis in the Pyrenees, covering the 1998-2010 period, showed a lower average DIN-N/TP-P (c. 116) and a trend to shift into potential nitrogen limitation (Camarero and Catalan, 2012). Also the Tatra mountains value in Brahney's review should be considered cautiously because it is based on DIN wet deposition during 1990-94 and average TP wet deposition during 1998-2009. Average values for the latter time interval in both, Tatra mountains and Bohemian forest, also had a lower DIN-N/TP-P ratio (98). On the other hand, the alternative 128±79 NO3-N/TP-P (mol/mol) deposition estimate for Ergaki would relocate our lake district closer to the other dots in the graph (fig. 6 a), with southern Sweden lake district as the closest one (125.6). In any case, Ergaki atmospheric nutrient deposition stoichiometry is by far larger than that of northern Sweden (20.1 molar DIN-N/TP-P), the paradigm of pristine areas with very low anthropogenic atmospheric nitrogen deposition and naturally nitrogen limited lakes. ... [23] отформатировано: Шрифт: 12 пт, без подчеркивания, Цвет шрифта: Авто

отформатировано: Испанский

Figure 6. The relationship between atmospheric nutrient deposition, nutrient bioavailability and phytoplankton biomass in Ergaki lakes (black square) in the context of different world data sets. a: Atmospheric deposition versus lake DIN-N/TP-P molar ratios of several alpine regions of the world, as reproduced from (Brahney et al., 2015), including Ergaki means and standard deviations: precipitation-dependent estimates (rhombus and dashed lines) and literature corrected estimate (square and solid lines). Vertical dashed line represents the referential north Sweden value (20.1). Horizontal dashed line represents the Redfield ratio (16). b: Yearly atmospheric DIN-N deposition versus Chl a/TP-P ratio of 13 Swedish regions, as reproduced from (Bergström and Jansson, 2006; Bergström et al., 2005), including Ergaki means and standard deviations (as before). Dashed lines represent approximate limits of N, N-P and P limitation areas, according to the authors (250 and 500 mg DIN-N_m²;y⁻¹). c: Lake DIN-N concentration versus Chl a/TP-P ratio of different lake districts in the world (black circles) and its segmented regression fit as

concentration versus Cm at 1r-r ratio of ultrerent take districts in the world (black circles) and its segmented regression fit as reproduced from (Camarero and Catalan, 2012), including Ergaki medians (black square) and particular observations from the following lakes (triangles): Tsirkovoe (yellow), Oiskoe (red), Raduzhnee (blue), Karovoe (grey) and Svetloe (green). Dashed line
 970 represents limit between N and P limitation conditions, according to the authors.

Year	month	Lake	DIN-N/TP-P	TN/TP	Limiting
			(mol/mol)	(mol/mol)	nutrient
2011	early June and	Oiskoe	<u>4.3</u>		N
	August	Svetloe	4.3		N
		Raduzhnoe	16.2		<u>N-P</u>
		Karovoe	<u>6.5</u>		N
		mean	<u>7.9</u>		N
2012	early June and	Oiskoe	109.2		Р
	August	Svetloe	<u>49.5</u>		<u>P</u>
		Raduzhnoe	<u>42.1</u>		<u>P</u>
		Karovoe	<u>29.6</u>		<u>P</u>
		mean	<u>57.6</u>		<u>P</u>
2015	early September	Oiskoe	3.7	11.6	Ν
		Raduzhnoe	10.5	30.9	Ν
		Tsirkovoe	23.6	82.8	Р
		mean	12.6	41.8	Ν
2017	late August	Oiskoe	6	<u>56.6</u>	N
		Karovoe	25.8	<u>61.9</u>	P
		Tsirkovoe	<u>97.9</u>	<u>164.1</u>	Р
		mean	43.3	<u>94.2</u>	Р

Table 3. Lake water N:P ratios along the studied years and limiting nutrient according to nutrient availability (DIN-N/TP-P) and the Redfield ratio.

(удалено: FIGURE

Отформатировано: Название объекта

удалено:

удалено:

удалено: In the same vein, the average molar stoichiometry of lake water samples was widely above the Redfield ratio (50±128 DIN-N/TP-P, fig. 6 a, table S2), which suggests lake phytoplankton could rather be P limited. Nonetheless, it is important to pay attention to the high dispersion of our data. The ratios in the 2012 survey were outliers one order of magnitude higher than in the other years, stretching the mean upwards (table 3). Indeed, there was a trend to turn from potential P (early June and August 2011-12) to N or N-P colimitation (early September 2015 and late August 2017) in all the studied lakes but in Tsirkovoe, if we compare the available nutrient ratio to the Redfield ratio (16:1, table 3). Besides to the Redfield criterion, a previous study determined that DIN-N/TP-P molar ratios of 3.3, 4.9 and 7.5 would correspond to 75%, 50% and 25% probabilities of chlorophyll increase under a N enrichment experiment (Bergström, 2010). According to this more restrictive criterion, only lake Oiskoe would approach the 75% probability of having a positive increase of chlorophyll under N enrichment in the last two surveys (2015, 2017). The effects of atmospheric nutrient deposition could be blurred by watershed processes at this lake. Forest and peat cover (25% and 6%, respectively) are important in comparison to the other lakes and, more importantly, the lake receives an inflow passing by nearby houses (table 1). Neverthelessa

Ν	удалено: 19.1
Ŋ	удалено: Р
N	удалено: 18.7
$\langle \rangle$	удалено: 71.3
$\langle \rangle$	удалено: 28.1
$\langle \rangle$	удалено: 34.3
$\langle \rangle$	удалено: 422.9
$\langle \rangle$	удалено: 194.4
$\langle \rangle$	удалено: 176.9
()	удалено: 38.1
Ì	удалено: 208.1
	удалено: Karovoe
1	удалено: 3.8
	удалено: 35.8
	удалено: 16.3
1	удалено: 39.1
Ì	удалено: N-
Ċ	удалено: 61.9
()	удалено: 103.7
$\langle \rangle$	удалено: 27.3
Ì	удалено: 59.5

3.7 Atmospheric input and lake phytoplankton biomass limitation

055

Atmospheric nitrogen deposition rate was lower in West Sayan mountains than in the most pristine areas in Sweden during the period 1995-2001, and two orders of magnitude lower than the most impacted Swedish region (Bergström et al., 2013, 2005; Bergström and Jansson, 2006) (fig. 6 b). The Swedish atmospheric deposition gradient was used to establish the new

- paradigm, according to which the natural state of phytoplankton growth in many unproductive lakes ($\leq 25 \mu g \text{ TP-P}_{e} \text{I}^{-1}$) would be nitrogen limitation, when atmospheric deposition is below c. 250 mg DIN-Nem⁻²-y⁻¹. According to that, the West Sayan mountain lakes (79-119 mg DIN-N₂m⁻²y⁻¹) would clearly belong to the potential atmospherically induced N limitation domain, where slight increases in atmospheric nitrogen deposition trigger larger phytoplankton biomass shifts. In a further study, Camarero and Catalan reckoned the upper threshold of N limitation of phytoplankton growth regime as ~100 µg DIN-
- 060 N_il⁻¹ in lake water (Camarero and Catalan, 2012) (fig. 6 c). Despite some of our measurements fell above the threshold, the median of 54, µg DIN-N_il⁻¹ would confirm, potential N limitation in most of the sampled lakes (table S2). Nevertheless, the potential character of that nitrogen limitation should be highlighted. If we combine all the mentioned pieces of evidence, it is understandable that atmospheric nutrient deposition was so low (fig. 6 b) that it led to a decoupling between atmospheric deposition and lake water nutrient stoichiometries (fig. 6 a). In the end, the lake water DIN concentration was low enough to
- allow the occurrence of phytoplankton growth N limitation events (fig. 6 c), but their occurrence was alternated with phosphorus limitation periods (table 3).

The future dynamics of DIN and TP concentrations in Ergaki lakes are likely to determine changes in phytoplankton biomass and species composition because the observed stoichiometries suggested a switching regime between N and P limitation events. That would promote alternatively low-P- and low-N-adapted species. Moreover, in the case of N, we found that shifts

- 070 resulting in DIN concentrations ranging from zero to the double of the average registered values would trigger important phytoplankton biomass responses (fig. 6 c). During our survey, nutrient concentrations in lakes were not majorly connected to atmospheric deposition (fig. 6 a), but future dynamics of atmospheric nutrients deposition in West Sayan mountain lakes are uncertain. On the one hand, the observed high snow sulphate concentrations and the wind analysis, made us suggest coal combustion in cities of central and eastern south Siberia to be the main winter atmospheric nitrate source. Additionally,
- 1075 predicted winter precipitation in this region under the RCP4.5 scenario (IPCC Representative Concentration Pathway scenario assuming 4.5 W₂m² radiative forcing by 2100) would increase 10-30% in 2016-2035 and up to 20-30% in 2081-2100 (IPCC, 2013). Therefore, it is likely that atmospheric nitrogen deposition increased even in a scenario where actual emissions did not change. On the other hand, wild forest fire events in south central Siberia have multiplied and intensified during the last decades and are expected to follow this trend in the 21st century as well (Brazhnik et al., 2017; Malevsky-
- 1080 Malevich et al., 2008). Their effect on atmospheric nutrient dynamics will be complex. Apart from modifying the sources of phosphorus rich biogenic aerosol particles, wildfires themselves used to be considered as a nitrogen volatilization pulse that left phosphorus on the burnt land (Hungate et al., 2003; Raison, 1979). Nevertheless, a recent study unveiled their relevance as a source of atmospheric phosphorus too (Wang et al., 2015). To sum up, uncertainties on the magnitude and timing of

удалено: The molar stoichiometry of volume weighted mean concentrations, i.e. the concentrations that would be measured if we had sampled the whole snow core at once, were 169±76 NO3-N/TP-P, 251±134 NO3-N/PO4-P, and 486±357 TN-N/TP-P (mol/mol). To pick up two cases from the literature, the nutrient ratios in the Bothnian Bay (northern Sweden) snowpack were 164±121 NO3-N/TP-P and 326±179 TN-N/TP-P (mol/mol), and on lake Redon (Catalan Pyrenees), 111±106 NO3-N/TP-P, 198±107 DIN-N/TP-P, 118+116 NO3-N/SRP-P and 216+104 TN-N/TP-P (mol/mol) (Catalan, 1989; Rahm et al., 1995), Therefore, NO3-N/TP-P or DIN-N/TP-P in the snowpacks were similar but TN/TP was higher in Ergaki.

удалено: by

удалено:

удалено: ed

удалено:

удалено:

удалено: According to these authors, unproductive N-limited lakes would shift into N-P colimitation regime under atmospheric depositions of 250-500 mg DIN-N·m-2·y-1, and into P limitation above 500. Therefore.

удалено:

удалено:

удалено:

According to Bergström and colleagues, the Chla/TP-P ratio of Nlimited lakes would increase with atmospheric N deposition up to the threshold where they would become P-limited. Above this threshold, extra nitrogen inputs would not change the Chla/TP-P ratio, and eventual phosphorus inputs would also trigger chlorophyll increases, so no change in the ratio would be observed. Later onIn a further

удалено: this

удалено:

удалено: 81

удалено:

удалено: s

удалено: The non normal distribution of lake DIN in our data e median a better estimate of the central value. Thus, the vicinity of West Sayan lakes to the threshold suggests that an eventual moderate increase in DIN could lead to a 2-3 fold phytoplankton biomass increase and a state change into P-limitation regime and consequent phytoplankton community shifts.

удалено: and phytoplankton limitation regime

удалено: is

удалено:

удалено: This would probably push lake DIN above the mentioned threshold and would trigger a phytoplankton shift.

future fossil fuel combustion, precipitation and wildfire regimes make it difficult to predict <u>the details of future atmospheric</u> nutrient deposition on West Sayan mountains. Nevertheless, it is very likely that deposition rates will increase and, perhaps, represent a sufficient input to finally couple the nutrient ratio of deposition inputs and lake water and/or to promote an increase in phytoplankton biomass₇

- 135 Finally, the sensitiveness of lake phytoplankton to nutrient availability might likely depend on temperature too as it was the case in the pristine Swedish north (Bergström et al., 2013). Low values of atmospheric nitrogen deposition in that region triggered a low DIN availability and a regional N limitation of lake phytoplankton growth. Nevertheless, the authors found clear phytoplankton responses to experimental NH4NO3 additions only in warm enough and N-limited lakes. As summer lake water temperatures were roughly 5-18 °C there and 5-14 °C here in West Sayan lakes, it is conceivable that the
- 140 phytoplankton response to nutrient availability shifts was temperature-mediated in Ergaki mountain lakes. Of course, the predicted warming in the region (IPCC, 2013) would decrease the system resilience, as eventual changes in nutrients availability could not be temperature mitigated and would be directly reflected in phytoplankton biomass and species composition changes.

YAAACHO: the status of West Sayan mountain lakes, but it is
credible that they were pushed into a higher trophic state and
phosphorus limitation regime, with its consequent phytoplankton
community shift.

(удалено: such

удалено: changes

удалено: The atmospheric nitrogen deposition in the Swedish

УДАЛЕНО: region declined to $< 100 \text{ mg DIN-N} \cdot \text{m}^{-2} \cdot \text{y}^{-1}$ in 2011, virtually as deposition in West Sayan lakes, but...

удалено: They found clear phytoplankton responses to experimental NH₄NO₃ additions only in warm enough and N-limited lakes. We speculate that relatively lower temperatures in West Sayan lakes might have impeded the Chl a/TP-P ratio to increase so far (fig. 6 c), but...

удалено: expected

1145 4. Conclusions

The Ergaki Natural Park in the West Sayan mountains was reckoned as an above-background but low atmospheric nutrient deposition area. Our atmospheric total phosphorus and nitrate deposition estimates reasonably fitted those predicted by global deposition models, whereas sulphate and phosphate proved to be higher than expected. While nitrogen values were comparable to the lowest records in other mountain areas of the World, phosphorus deposition was likely at the very lowest range ever measured on terrestrial ecosystems before. Nevertheless, any conclusions regarding yearly atmospheric phosphorus deposition loads from this study should be contrasted with further year-long measurements in West Sayan mountains, including the presumably important biogenic and wildfire contributions during spring and summer seasons. The atmospheric reactive nitrogen deposition load on Ergaki Natural Park was similar to that in northern Sweden, so low enough to potentially induce nitrogen limitation in lake phytoplankton. However, the strongly mismatching nutrient ratios in

- 155 atmospheric deposition and lake water suggested a decoupling between the two compartments. The stoichiometry of atmospheric nutrient deposition are highly determined by the extremely low phosphorus deposition and the biogeochemical processes occurring at the watershed level would easily modify the nutrient ratio of the tiny and unbalanced atmospheric input. Under such conditions, a succession of N and P limitation events was observed rather than the expected N limitation. Thus, we hypothesise that the recently suggested paradigm of nitrogen limitation of lake phytoplankton growth in natural
- 160 pre-industrial conditions might apply for Sweden and other regions with a minimal atmospheric phosphorus deposition but maybe not for large areas in Siberia. A larger number of lakes distributed over Siberia and whole year long atmospheric nutrient deposition, including spring and summer seasons, should be studied to contrast that hypothesis.

29

удалено:	W		

удалено: In conclusion, t

удалено: However, according to our best estimate of phosphorus deposition, tT

удалено: could be

отформатировано: без подчеркивания, Цвет шрифта: Авто, Английский (США) 185 Finally, the West Sayan mountain range constitutes an excellent pristine area to study the effects of past global warming with a minimal influence of atmospheric nitrogen and phosphorus deposition. The studied mountain lakes are also good sensitive sentinels for upcoming climate warming and atmospheric nutrient deposition dynamics. The expected increase in precipitation and wildfires and future fossil fuel combustion, could finally couple these lakes to the atmospheric nutrient deposition and, if climate warming unlocked the phytoplankton response to it, promote a phytoplankton biomass increase and species composition change.

Data availability

The data sets presented in this study are available as supplementary tables S1 and S2 at the repository of the library of the library of the Siberian Federal University: http://elib.sfu-kras.ru/handle/2311/135098.

Team list

Diaz-de-Quijano, Daniel; Ageev, Aleksander Vladimirovich; Ivanova, Elena Anatolevna; Anishchenko, Olesia Valerevna

Author contribution

DD and EAI designed the snow sampling, DD and AVA performed it and DD and OVA analysed snow samples. All the authors participated in sampling and sample analyses of lake water. DD prepared the manuscript with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

1205 The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model and READY website (http://www.ready.noaa.gov) used in this publication. We also would like to thank the direction of the Ergaki Natural Park for their support and willingness to collaborate.

30

отформатировано: без подчеркивания, Цвет шрифта: Авто, Английский (США)

отформатировано: без подчеркивания, Цвет шрифта: Авто, Английский (США)

отформатировано: без подчеркивания, Цвет шрифта: Авто, Английский (США)

УДАЛЕНО: As a result, the deposited DIN-N/TP-P ratio would be at the upper quartile of alpine regions worldwide and similar to southern Sweden, where phytoplankton in low productive lakes was generally phosphorus limited. As a synthesis of these contradictory these...

YGAJEHO: s, the observed lake nutrient stoichiometry and nutrient limitation regime resulted to be variable, with an apparent trend over the years and/or along the seasonal succession from phosphorus to nitrogen limitation.[¶]

This frontier status between phosphorus and nitrogen limitation and the low atmospheric deposition rates make West Sayan mountain lakes very sensitive to any shift in nutrient deposition. For example, the expected precipitation and wildfires increase could push these lakes to the phosphorus limitation realm, increased phytoplankton biomass and community change, in case that fossil fuel combustion was not limited. Nevertheless, such changes could be temperature limited in a way that lakes were on the eve of a temperature unlocking of primary production and a consequent change of state in the ecosystems. Therefore both, phytoplankton biomass and species composition in these lakes might serve as helpful early warnings of global and regional environmental change.[¶]

Lastly, further studies on the effect of atmospheric nutrient deposition on Siberian mountain lakes, namely including the whole year long deposition, are strongly recommended. The pre-industrial lake phytoplankton nitrogen limitation paradigm was based on a particular region of the world: first in Sweden and then Norway, Colorado, etc. (Bergström et al., 2013, 2005; Bergström and Jansson, 2006; Elser et al., 2009). Nevertheless, increasing atmospheric phosphorus deposition from Saharan origin on the Pyrenees was able to reduce lake DIN, despite increasign atmospheric nitrogen deposition (Camarero and Catalan, 2012). This, along with the extremely low snow-season atmospheric phosphorus deposition measured in this study opens the possibility to hypothesise variations in the general paradigm, there where atmospheric phosphorus depositions were particularly high or low. Thus, in case that the whole year long atmospheric nitrogen to phosphorus deposition ratio kept high enough due to low phosphorus deposition, phytoplankton in Siberian mountain lakes at pre-industrial times could have been either phosphorus limited or, simply, insignificantly affected by atmospheric inputs.

Financial Support

The reported study was funded by the Russian Foundation for Basic Research (RFBR), project number 20-04-00960 and the program 5-100 of the Ministry of Science and Higher Education of the Russian Federation.

1250 References

Anishchenko, O. V., Glushchenko, L. A., Dubovskaya, O. P., Zuev, I. V., Ageev, A. V. and Ivanova, E. A.: Morphometry and metal concentrations in water and bottom sediments of mountain lakes in Ergaki Natural Park, Western Sayan Mountains, Water Resour., 42(5), 670–682, doi:10.1134/S0097807815050036, 2015.

APHA: Standard methods for the examination of water and wastewater, 17th ed., edited by American Public Health 1255 Association., 1989.

Banks, H. H. and Nighswander, J. E.: Relative Contribution of Hemlock Pollen to the Phosphorus Loading of the Clear Lake Ecosystem Near Minden, Ontario, in Symposium on Sustainable Management of Hemlock Ecosystems in Eastern North America, Proceedings, pp. 168–174., 2000.

Bergström, A.-K. and Jansson, M.: Atmospheric nitrogen deposition has caused nitrogen enrichment and eutrophication of lakes in the northern hemisphere, Glob. Chang. Biol., 12(4), 635–643, doi:10.1111/j.1365-2486.2006.01129.x, 2006.

Bergström, A.-K., Faithfull, C., Karlsson, D. and Karlsson, J.: Nitrogen deposition and warming – effects on phytoplankton nutrient limitation in subarctic lakes, Glob. Chang. Biol., 19, 2557–2568, doi:10.1111/gcb.12234, 2013.

Bergström, A. K.: The use of TN:TP and DIN:TP ratios as indicators for phytoplankton nutrient limitation in oligotrophic lakes affected by N deposition, Aquat. Sci., 72(3), 277–281, doi:10.1007/s00027-010-0132-0, 2010.

1265 Bergström, A. K., Blomqvist, P. and Jansson, M.: Effects of atmospheric nitrogen deposition on nutrient limitation and phytoplankton biomass in unproductive Swedish lakes, Limnol. Oceanogr., 50(3), 987–994, doi:10.4319/lo.2005.50.3.0987, 2005.

Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J. W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L. and De Vries, W.: Global

1270 assessment of nitrogen deposition effects on terrestrial plant diversity: A synthesis, Ecol. Appl., 20(1), 30–59, doi:10.1890/08-1140.1, 2010.
 Brahney, J., Mahowald, N., Ward, D. S., Ballantyne, A. P. and Neff, J. C.: Is atmospheric phosphorus pollution altering

global alpine lake stoichiometry?, Global Biogeochem. Cycles, 29(9), 1369–1383, 2015.

Brazhnik, K., Hanley, C. and Shugart, H. H.: Simulating Changes in Fires and Ecology of the 21st Century Eurasian Boreal
 Forests of Siberia, Forests, 8(49), 1–27, doi:10.3390/f8020049, 2017.

Brown, L. J., Taleban, V., Gharabaghi, B. and Weiss, L.: Seasonal and spatial distribution patterns of atmospheric phosphorus deposition to Lake Simcoe, ON, J. Great Lakes Res., 37(SUPPL. 3), 15–25, doi:10.1016/j.jglr.2011.01.004,

2011.

Camarero, L. and Catalan, J.: Atmospheric phosphorus deposition may cause lakes to revert from phosphorus limitation back to nitrogen limitation, Nat. Commun., 3, 1118, doi:10.1038/ncomms2125, 2012.

Carslaw, D. C. and Ropkins, K.: openair --- an R package for air quality data analysis, Environ. Model. Softw., 27–28, 52–61, 2012.

Catalan, J.: The Winter Cover of a High-Mountain Mediterranean Lake (Estany Redó, Pyrenees), Water Resour. Res., 25(3), 519–527, 1989.

1285 Chernov, G., Vdovin, V., Okishev, P., Petkevich, M., Mistryukov, A., Zyatkova, L. and Milyaeva, L.: Relief Altae-Sayanskoi Gornoi Oblasti, edited by V. Nikolaev and G. Chernov, Nauka Sibirskoe Otdelenie., 1988. Clement, J. C., Robson, T. M., Guillemin, R., Saccone, P., Lochet, J., Aubert, S. and Lavorel, S.: The effects of snow-N

deposition and snowmelt dynamics on soil-N cycling in marginal terraced grasslands in the French Alps, Biogeochemistry, 108(1–3), 297–315, doi:10.1007/s10533-011-9601-3, 2012.

1290 Cragin, J. H., Hewitt, A. D. and Colbeck, S. C.: Grain-scale mechanisms influencing the elution of ions from snow, Atmos. Environ., 30(1), 119–127, doi:10.1016/1352-2310(95)00232-N, 1996.

Dambrine, E., Arthaud, F., Avrillier, J.-N., Millery, A. and Nicolas, M.: Altitudinal and Latitudinal Variations of Snowpack N Concentration over the French Alps, Water Air Soil Pollut, 229(250), 1–12, 2018.

DeForest, J. L., Zak, D. R., Pregitzer, K. S. and Burton, A. J.: Atmospheric Nitrate Deposition, Microbial Community 1295 Composition, and Enzyme Activity in Northern Hardwood Forests, Soil Sci. Soc. Am. J., 68(1), 132,

doi:10.2136/sssaj2004.1320, 2004.

Diaz-de-Quijano, D.: On the track of cellular ecology. Methodological improvements and contributions of single-cell phosphatase activity on the ecology of phytoplankton in Pyrenean lakes., 2014.

Doskey, P. V and Ugoagwu, B. J.: Macronutrient chemistry of pollen, Commun. Soil Sci. Plant Anal., 23(1&2), 15–23, doi:10.1080/00103629209368566, 1992.

Eimers, M. C., Hillis, N. P. and Watmough, S. A.: Phosphorus Deposition in a Low-Phosphorus Landscape: Sources, Accuracy and Contribution to Declines in Surface Water P, Ecosystems, 21(4), 782–794, doi:10.1007/s10021-017-0184-2, 2018.

Elser, J. J., Andersen, T., Baron, J. S., Bergström, A.-K., Jansson, M., Kyle, M., Nydick, K. R., Steger, L. and Hessen, D. O.:

1305 Shifts in lake N:P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition., Science (80-.)., 326(5954), 835–837, doi:10.1126/science.1176199, 2009.

Fagerli, H., Tsyro, S., Jonson, J. E., Nyíri, Á., Gauss, M., Simpson, D., Wind, P., Benedictow, A., Klein, H., Mortier, A., Aas, W., Hjellbrekke, A.-G., Solberg, S., Platt, S. M., Yttri, K. E., Tørseth, K., Gaisbauer, S., Mareckova, K., Matthews, B., Schindlbacher, S., Sosa, C., Tista, M., Ullrich, B., Wankmüller, R., Scheuschner, T., Bergström, R., Johanson, L., Jalkanen,

1310 J.-P., Metzger, S., Denier van der Gon, H. D. C., Kuenen, J. J. P., Visschedjik, A. J. H., Barregård, L., Molnár, P. and Stockfelt, L.: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Oslo., 2019.

Felip, M., Sattler, B., Psenner, R. and Catalan, J.: Highly Active Microbial Communities in the Ice and Snow Cover of High Mountain Lakes, Appl. Environ. Microbiol., 61(6), 2394–2401 [online] Available from: http://aem.asm.org/cgi/content/abstract/61/6/2394, 1995.

1315 Gauss, M., Tsyro, S., Benedictow, A., Fagerli, H., Hjellbrekke, A.-G., Aas, W. and Solberg, S.: EMEP / MSC-W model performance for acidifying and eutrophying components and photo-oxidants in 2017. Supplementary material to EMEP Status Report 1/2019, Oslo., 2019.

Gladyshev, M. I., Kolmakova, O. V., Tolomeev, A. P., Anishchenko, O. V., Makhutova, O. N., Kolmakova, A. A., Kravchuk, E. S., Glushchenko, L. A., Kolmakov, V. I. and Sushchik, N. N.: Differences in organic matter and

1320 bacterioplankton between sections of the largest arctic river: Mosaic or continuum?, Limnol. Oceanogr., 60(4), 1314–1331, doi:10.1002/lno.10097, 2015.

Grasshoff, K., Ehrhamdt, M. and Kremling, K.: Methods of sea water analysis, 2nd Ed., Wiley Verlag-Chemie, Weinheim., 1983.

Güsewell, S.: N: P ratios in terrestrial plants: variation and functional significanc, New Phytol., 164, 243–266, doi:10.1111/j.1469-8137.2004.01192.x, 2004.

Hijmans, R. J.: geosphere: Spherical Trigonometry, [online] Available from: https://cran.r-project.org/package=geosphere, 2017.

Hungate, B. A., Cole, J., Vitousek, P. M., Stewart, J., Victoria, R., Kenichi, S. and Naiman, R. J.: Disturbance and element interactions, in SCOPE 61: Interactions Among the Major Biogeochemical Cycles, edited by J. M. Melillo, C. B. Field, and

 B. Moldan, pp. 47–62, Island Press, Washington DC., 2003.
 IPCC: Climate Change 2013. The Physical Science Basis. Annex I: Atlas of Global and Regional Climate Projections., 2013.

Jambers, W. and Van Grieken, R.: Single particle characterization of inorganic suspension in Lake Baikal, Siberia, Environ. Sci. Technol., 31(5), 1525–1533, doi:10.1021/es9608003, 1997.

1335 Jassby, A. D., Reuter, J. E., Axler, R. P., Goldman, C. R. and Hackley, S. H.: Atmospheric Deposition of Nitrogen and Phosphorus in theAnnual Nutrient Load of Lake Tahoe (California‐Nevada), Water Resour. Res., 30(7), 2207– 2216, 1994.

Jeffrey, S. W. and Humphrey, G. F.: New spectrophotometric equations for determining chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton, Biochem. und Physiol. der Pflanz., 167(2), 191–194, 1975.

 Kahle, D. and Wickham, H.: ggmap: Spatial Visualization with ggplot2, R J., 5(1), 144–161 [online] Available from: http://journal.r-project.org/archive/2013-1/kahle-wickham.pdf, 2013.
 Kalacheva, G. S., Gubanov, V. G., Gribovskaya, I. V., Gladchenko, I. A., Zinenko, G. K. and Savitsky, S. V.: Chemical

analysis of Lake Shira water (1997-2000), Aquat. Ecol., 36(2), 123-141, doi:10.1023/A:1015695813280, 2002.

Kopáček, J., Stuchlík, E., Straškrabová, V. and Pšenáková, P.: Factors governing nutrient status of mountain lakes in the 1345 Tatra Mountains, Freshw. Biol., 43, 369–384, doi:10.1046/j.1365-2427.2000.00569.x, 2000.

Kopáček, J., Turek, J., Hejzlar, J. and Porcal, P.: Bulk deposition and throughfall fluxes of elements in the Bohemian Forest (central Europe) from 1998 to 2009, Boreal Environ. Res., 16, 495–508, 2011a.

Kopáček, J., Hejzlar, J., Vrba, J. and Stuchlík, E.: Phosphorus loading of mountain lakes: Terrestrial export and atmospheric deposition, Limnol. Oceanogr., 56(4), 1343–1354, doi:10.4319/lo.2011.56.4.1343, 2011b.

- 1350 Kuhn, M.: The nutrient cycle through snow and ice, a review, Aquat. Sci., 63(2), 150–167, doi:10.1007/PL00001348, 2001. Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C.-U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., MacKenzie, I. a., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D. and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP):
- 1355 evaluation of historical and projected future changes, Atmos. Chem. Phys., 13(16), 7997-8018, doi:10.5194/acp-13-7997-2013, 2013.

LeBauer, D. S. and Treseder, K. K.: Nitrogen Limitation of Net Primary Productivity in Terrestrial Ecosystems is Globally Distributed, Ecology, 2, 371–379, doi:10.1007/BF00153104, 2008.

Logatchev, N. A.: History and geodynamics of the Lake Baikal rift in the context of the Eastern Siberia rift system: a review, 1360 Bull. Centres Rech. Explor. Elf Aquitaine, 17(2), 353–370, 1993.

- Mahowald, N., Jickells, T. D., Baker, A. R., Artaxo, P., Benitez-Nelson, C. R., Bergametti, G., Bond, T. C., Chen, Y., Cohen, D. D., Herut, B., Kubilay, N., Losno, R., Luo, C., Maenhaut, W., McGee, K. A., Okin, G. S., Siefert, R. L. and Tsukuda, S.: Global distribution of atmospheric phosphorus sources, concentrations and deposition rates, and anthropogenic impacts, Global Biogeochem. Cycles, 22(4), 1–19, doi:10.1029/2008GB003240, 2008.
- 1365 Malevsky-Malevich, S. P., Molkentin, E. K., Nadyozhina, E. D. and Shklyarevich, O. B.: An assessment of potential change in wildfire activity in the Russian boreal forest zone induced by climate warming during the twenty-first century, Clim. Change, 86, 463–474, doi:10.1007/s10584-007-9295-7, 2008.

Marcé, R., Obrador, B., Morguí, J.-A., Riera, J., López, P. and Armengol, J.: Carbonate weathering as a driver of CO2 supersaturation in lakes, Nat. Geosci., 8, 107–111, doi:doi:10.1038/ngeo2341, 2015.

1370 Medina-Sanchez, J. M., Villar-Argaiz, M. and Carrillo, P.: Neither with nor without you: A complex algal control on bacterioplankton in a high mountain lake, Limnol. Oceanogr., 49, 1722–1733 [online] Available from: Dani DEA Mixotrofia, 2004.

Mosello, R., Berg, T., Nicuks, U. and Appleby, P.: Atmospheric Deposition. Sampling and Analysis of Direct Deposition, Snow Pack and Soil Cores, in Measuring and Modelling the Dynamic Response of Remote Mountian Lake Ecosystems to Environmental Change, pp. 1–16., 1997.

Ohizumi, T., Take, N., Inomata, Y., Yagoh, H., Endo, T., Takahashi, M., Yanahara, K. and Kusakabe, M.: Long-term variation of the source of sulfate deposition in a leeward area of Asian continent in view of sulfur isotopic composition, Atmos. Environ., 140, 42–51, doi:10.1016/j.atmosenv.2016.05.057, 2016.

1375

Pearson, C., Schumer, R., Trustman, B. D., Rittger, K., Johnson, D. W. and Obrist, D.: Nutrient and mercury deposition and



1380 storage in an alpine snowpack of the Sierra Nevada, USA, Biogeosciences, 12(12), 3665–3680, doi:10.5194/bg-12-3665-2015, 2015.

R Development Core Team: R: A Language and Environment for Statistical Computing, R Found. Stat. Comput., doi:10.1007/978-3-540-74686-7, 2017.

Rahm, L., Hakansson, B., Larsson, P., Fogelqvist, E., Bremle, G. and Valderrama, J.: Nutrient and Persistent Pollutant 1385 Deposition on the Bothnian Bay Ice and Snow fields, Water Air Soil Pollut, 84, 187–201, 1995.

Raison, R. J.: Modification of the Soil Environment by Vegetation Fires, with Particular Reference to Nitrogen Transformations: a Review, Plant Soil, 51, 73–108, 1979.

Reinemann, S. A., Porinchu, D. F., Gustin, M. A. and Mark, B. G.: Historical trends of mercury and spheroidal carbonaceous particle deposition in sub-alpine lakes in the Great Basin, United States, J. Paleolimnol., 52, 405–418, doi:10.1007/s10933-1390
 014-9801-7, 2014.

Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F. S., Lambin, E. F., Lenton, T. M., Scheffer, M., Folke, C., Schellnhuber, H. J., Nykvist, B., de Wit, C. A., Hughes, T., van der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P. K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R. W., Fabry, V. J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P. and Foley, J. A.: A safe operating space for humanity., Nature, 461(7263), 472–475, doi:10.1038/461472a, 2009.

Rolph, G., Stein, A. and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, Environ. Model. Softw., 95, 210–228, doi:10.1016/j.envsoft.2017.06.025, 2017.

Rose, N. L.: A note on further refinements to a procedure for the extraction of carbonaceous fly-ash particles from sediments, J. Paleolimnol., 11(2), 201–204, doi:10.1007/BF00686866, 1994.

1400 Rose, N. L.: Quality control in the analysis of lake sediments for spheroidal carbonaceous particles, Limnol. Oceanogr., 6, 172–179, doi:10.4319/lom.2008.6.172, 2008.

Rose, N. L. and Monteith, D. T.: Temporal trends in spheroidal carbonaceous particle deposition derived from annual sediment traps and lake sediment cores and their relationship with non-marine sulphate, Environ. Pollut., 137, 151–163, doi:10.1016/j.envpol.2004.12.022, 2005.

1405 Rose, N. L. and Ruppel, M.: Environmental Archives of Contaminant Particles, in Environmental Contaminants, vol. 18, pp. 187–221., 2015.

Rose, N. L., Appleby, P. G., Boyle, J. F., Mackay, A. W. and Flower, R. J.: The spatial and temporal distribution of fossilfuel derived pollutants in the sediment record of Lake Baikal, eastern Siberia, J. Paleolimnol., 20, 151–162, 1998.

Rose, N. L., Harlock, S. and Appleby, P. G.: The spatial and temporal distributions of spheroidal carbonaceous fly-ash
particles (scp) in the sediment records of european mountain lakes, Water. Air. Soil Pollut., 1–32, 1999a.

Rose, N. L., Harlock, S. and Appleby, P. G.: The spatial and temporal distributions of spheroidal carbonaceous fly-ash particles (SCP) in the sediment records of european mountain lakes, Water. Air. Soil Pollut., 113, 1–32, 1999b.

Rose, N. L., Flower, R. J. and Appleby, P. G.: Spheroidal carbonaceous particles (SCPs) as indicators of atmospherically

deposited pollutants in North African wetlands of conservation importance, Atmos. Environ., 37, 1655–1663, 1415 doi:10.1016/S1352-2310(03)00012-8, 2003.

Sickman, J. O., Melack, J. M. and Clow, D. W.: Evidence for nutrient enrichment of high-elevation lakes in the Sierra Nevada, California, Limnol. Oceanogr., 48(5), 1885–1892, doi:10.4319/lo.2003.48.5.1885, 2003.

Slanina, J., Lingerak, W. A. and Bergman, L.: A fast determination of nitrate in rain and surface waters by means of UV spectrophotometry, Zeitschrift für Anal. Chemie, 280(5), 365–368, 1976.

1420 Šporka, F., Štefková, E., Bitušík, P., Thompson, A. R., Agustí-Panareda, A., Appleby, P. G., Grytnes, J. A., Kamenik, C., Krno, I., Lami, A., Rose, N. and Shilland, N. E.: The paleolimnological analysis of sediments from high mountain lake Nižné Terianske pleso in the High Tatras (Slovakia), J. Paleolimnol., 28, 95–109, 2002.

Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R., Carpenter, S. R., Vries, W. De, Wit, C. A. De, Folke, C., Gerten, D., Heinke, J., Mace, G. M., Persson, L. M., Ramanathan, V., Reyers, B. and Sörlin,

1425 S.: Planetary boundaries: Guiding human development on a changing planet, Science (80-.)., 347(6223), 1259855-1–10, doi:10.1126/science.1259855, 2015.

Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: Noaa's hysplit atmospheric transport and dispersion modeling system, Bull. Am. Meteorol. Soc., 96(12), 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.

Stepanov, N. V.: Sosudistye Rasteniya Priyeniseiskikh Sayan, Institut Fundamentalnoy Biologii i Biotekhnologii. Siberian 1430 Federal University, Krasnoyarsk., 2016.

Stottlemyer, R. and Rutkowski, D.: Multiyear Trends in Snowpack Ion Accumulation and Loss, Northern Michigan with from, Water Resour. Res., 26(4), 721–737, 1990.

Takhtadzhyan, A. L.: Floristicheskie Oblasti Zemli, Nauka., 1978.

Tipping, E., Benham, S., Boyle, J. F., Crow, P., Davies, J., Fischer, U., Guyatt, H., Helliwell, R., Jackson-Blake, L., Lawlor,
1435 A. J., Monteith, D. T., Rowe, E. C. and Toberman, H.: Atmospheric deposition of phosphorus to land and freshwater,
Environ. Sci. Process. Impacts, 16(7), 1608–1617, doi:10.1039/c3em00641g, 2014.

Tolomeev, A. P., Anishchenko, O. V., Kravchuk, E. S., Kolmakova, O. V., Glushchenko, L. A., Makhutova, O. N., Kolmakova, A. A., Kolmakov, V. I., Trusova, M. Y., Sushchik, N. N. and Gladyshev, M. I.: Component elements of the carbon cycle in the middle and lower Yenisei River, Contemp. Probl. Ecol., 7(4), 489–500, doi:10.1134/S1995425514040118, 2014.

VA: Phytoplankton pigments in oceanography: guidelines to modern methods, edited by S. W. Jeffrey, R. F. C. Mantoura, and S. W. Wright, Paris., 1997.

Ventura, M., Camarero, L., Buchaca, T., Bartumeus, F., Livingstone, D. M. and Catalan, J.: The main features of seasonal variability in the external forcing and dynamics of a deep mountain lake (Redó. Pyrenees), J. Limnol., 59(SUPPL. 1), 97–108, doi:10.4081/jlimnol.2000.s1.97, 2000.

Vitousek, P. M., Aber, J. D., Howarth, R. H., Likens, G. E., Matson, P. A., Schindler, D. W., Schlesinger, W. H. and Tilman,
 D. G.: Human alteration of the global nitrogen cycle: Source and consequences, Ecol Appl, 7(3), 737–750,

doi:10.1038/nn1891, 1997.

Voskresenskii, S.: Geomorfologia Sibirii, Izdatelstvo MGU., 1962.

1450 Wang, R., Balkanski, Y., Boucher, O., Ciais, P. and Tao, S.: Significant contribution of combustion-related emissions to the atmospheric phosphorus budget, Nat. Geosci., 8(January), 4–10, doi:10.1038/NGEO2324, 2015.

Wang, S., Shi, X., Cao, W. and Pu, T.: Seasonal Variability and Evolution of Glaciochemistry at An Alpine Temperate Glacier on the Southeastern Tibetan Plateau, Water, 10(114), 1–14, doi:10.3390/w10020114, 2018a.

Wang, W., Liu, X., Xu, J., Dore, A. J. and Xu, W.: Imbalanced nitrogen and phosphorus deposition in the urban and forest l455 environments in southeast Tibet, Atmos. Pollut. Res., 9(4), 774–782, doi:10.1016/j.apr.2018.02.002, 2018b.

Wasiuta, V., Norman, A. L., Lafrenière, M. J. and Hastings, M. G.: Background atmospheric sulfate deposition at a remote alpine site in the Southern Canadian Rocky Mountains, J. Geophys. Res. Atmos., 120(21), 11,352-11,367, doi:10.1002/2015JD023835, 2015.

Wik, M. and Renberg: Environmental records of carbonaceous fly-ash particles from fossil-fuel combustion, J. Paleolimnol.,1460 15(ii), 193–206, 1996.

Williams, M. W. and Melack, J. M.: Solute Chemistry of Snowmelt and Runoff in an Alpine Basin , Sierra Nevada, Water Resour. Res., 27(7), 1575–1588, doi:10.1029/90WR02774, 1991.

Xu, W., Liu, L., Cheng, M., Zhao, Y., Zhang, L., Pan, Y., Zhang, X., Gu, B., Li, Y., Zhang, X., Shen, J., Lu, L., Luo, X., Zho, Y., Feng, Z., Collett, J. L., Zhang, F. and Liu, X.: Spatial-temporal patterns of inorganic nitrogen air concentrations and deposition in eastern China, Atmos. Chem. Phys., 18(15), 10931–10954, doi:10.5194/acp-18-10931-2018, 2018.

l465 deposition in eastern China, Atmos. Chem. Phys., 18(15), 10931–10954, doi:10.5194/acp-18-10931-2018, 2018.
 Yang, H., Rose, N. L. and Battarbee, R. W.: Dating of recent catchment peats using spheroidal carbonaceous particle (SCP) concentration profiles with particular reference to Lochnagar, Scotland, Holocene, 11(5), 593–597, doi:10.1191/095968301680223549, 2001.

Страница 4: [1] удалено	Daniel Diaz de Quijano	17.12.2020 19:10:00
Страница 4: [2] удалено	Daniel Diaz de Quijano	17.12.2020 19:10:00
Страница 4: [3] удалено	danila	30.10.2020 16:49:00
Страница 5: [4] удалено	danila	30.10.2020 16:48:00
Страница 5: [5] отформатировано	Unknown	
Шрифт: 12 пт, без подчеркивания, Цвет	г шрифта: Авто	
Страница 5: [6] отформатировано	Daniel Diaz de Quijano	24.11.2020 18:26:00
Шрифт: полужирный, Цвет шрифта: Ф	он 1, Английский (США)	
Страница 5: [7] отформатировано	Daniel Diaz de Quijano	22.11.2020 19:17:00
без подчеркивания, Цвет шрифта: Фон	1	
Страница 5: [7] отформатировано	Daniel Diaz de Quijano	22.11.2020 19:17:00
оез подчеркивания, Цвет шрифта: Фон	1	
	Daniel Diaz de Quijano	21 11 2020 21.14.00
		21.11.2020 21.14.00
осз подчеркивания, цвет шрифта. Авте		
Страница 5: [8] отформатировано	Daniel Diaz de Quijano	21.11.2020 21:14:00
без подчеркивания, Цвет шрифта: Авто	, Английский (США)	
.		
Страница 5: [9] отформатировано	Daniel Diaz de Quijano	21.11.2020 20:46:00
без подчеркивания, Английский (США)	
		21.11.2020.20.46.00
страница 5: [9] отформатировано	Daniel Diaz de Quijano	21.11.2020 20:46:00
зез подчеркивания, Английский (США)	
Страница 5: [9] отформатировано	Daniel Diaz de Ouijano	21.11.2020 20:46:00
без полчеркивания. Английский (США		
	7	
Страница 5: [9] отформатировано	Daniel Diaz de Quijano	21.11.2020 20:46:00
без подчеркивания, Английский (США)	
- ``	·	
Страница 5: [9] отформатировано	Daniel Diaz de Quijano	21.11.2020 20:46:00
без подчеркивания, Английский (США)	
Страница 5: [9] отформатировано	Daniel Diaz de Quijano	21.11.2020 20:46:00
без подчеркивания, Английский (США)	
	de a lla	20 10 2020 10 57.00
траница 5: [10] Отформатировано	aaniia	30.10.2020 16:57:00

восточноазиатскими буквами и цифрами

▼		
A		
Страница 17: [14] удалено	Daniel Diaz de Quijano	20.12.2020 00:41:00
Χ		
		20.42.2020.00.40.00
Страница 17: [15] удалено	Daniel Diaz de Quijano	20.12.2020 00:48:00
▼		
		20.42.2020.00.50.00
Страница 17: [16] удалено	Daniel Diaz de Quijano	20.12.2020 00:59:00
Υ		
	Daniel Dias de Onitana	20 12 2020 01.27.00
Страница 17: [17] удалено	Daniel Diaz de Quijano	20.12.2020 01:27:00
Υ		
	Daniel Diaz de Ouijane	20 12 2020 01.25.00
Страница 17: [10] удалено	Daniel Diaz de Quijano	20.12.2020 01:25:00
Υ		
	Daniel Diaz de Quijano	24 11 2020 18:26:00
Страница 10. [19] удалено		24.11.2020 18.20.00
▼		
	Daniel Diaz de Quijano	24 11 2020 18:26:00
страница 10. [19] удалено		24.11.2020 10.20.00
Страница 18: [19] удадено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
Страница 18: [19] удалено	Daniel Diaz de Ouijano	24.11.2020 18:26:00
- Страница 18: [19] удалено	Daniel Diaz de Ouijano	24.11.2020 18:26:00
-		
×		
- Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
×		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
· · · · · · · · · · · · · · · · · · ·		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
.		
_		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
v		
<u> </u>		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00

Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
A		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
v		
A		
Страница 18: [19] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
A		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
	Daniel Diaz de Quijano	24 11 2020 18:26:00
страница 10. [20] удалено	Damer Diaz de Quijano	24.11.2020 10.20.00
×		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
•		
▲		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
V		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
A		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
Χ		
A		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
▼		
	Daniel Diaz de Quijane	24 11 2020 18:26:00
Страница 10. [20] удалено	Damer Diaz de Quijano	24.11.2020 18:20:00
	Daniel Diaz de Quijano	24 11 2020 18:26:00
страница 10. [20] удалено	Damer Diaz de Quijano	24.11.2020 10.20.00
V		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
τ		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
T		
Страница 18: [20] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
↓		

A		
Страница 18: [21] удалено	Daniel Diaz de Quijano	24.11.2020 18:26:00
τ		
A		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
▼		
A		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
▼		
A		24.44.2020.40.20.00
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
	Daniel Diaz de Quijano	24 11 2020 18:30:00
erpanniqu Ior [22] yaurene		
Υ		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
V		
<u>ــــــــــــــــــــــــــــــــــــ</u>		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
▼		
<u> </u>		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
▼		
A		
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
▼		
A	N N	
Страница 18: [22] удалено	Daniel Diaz de Quijano	24.11.2020 18:30:00
Υ		
	Daniel Diaz de Ouijano	24 11 2020 18:30:00
Страница 10. [22] удалено	Daniel Diaz de Quijano	24.11.2020 10.50.00
V		
Страница 18: [22] удалено	Daniel Diaz de Ouijano	24.11.2020 18:30:00
T		
Страница 26: [23] удалено	Daniel Diaz de Quijano	16.12.2020 22:22:00
I		
A		
Страница 27: [24] удалено	Daniel Diaz de Quijano	16.12.2020 22:21:00
V		
_		

l