1   **Wetter environment and increased grazing reduced the area burned in northern Eurasia: 2002 – 2016**

2   Wei Min Hao[1], Matthew C. Reeves[2], L. Scott Baggett[3], Yves Balkanski[4], Philippe Ciais[4], Bryce L. Nordgren[1], Alexander Petkov[1],
3   Rachel E. Corley[1], Florent Mouillot[5], Shawn P. Urbanski[1], Chao Yue[6]
4
5   [1]United States Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory, 5775 Highway 10 West, Missoula, MT
6   59808 USA.
7   [2]United States Forest Service, Rocky Mountain Research Station, Forestry Sciences Laboratory, 800 E. Beckwith, Missoula, MT
8   59801, USA.
9   [3]United States Forest Service, Rocky Mountain Research Station, 240 West Prospect, Fort Collins, CO 80526, USA.
10 [4]Laboratoire des Sciences du Climat et de l'Environnement, LSCE CEA CNRS UVSQ, 91191 Gif Sur Yvette, France.
11 [5]UMR CEFE 5175, Centre National de la Recherche Scientifique (CNRS), Université de Montpellier, Université Paul-Valéry
12 Montpellier, Ecole Pratique des Hautes Etudes (EPHE), Institut de Recherche pour le Développement, 34293 Montpellier CEDEX 5,
13 France.
14 [6]Institute of Soil and Water Conservation, Northwest A&F University, Yangling, Shaanxi 712100, P.R. China.
15
16 **Correspondence:** Wei Min Hao (weimin.hao@usda.gov)
17

18 Supporting Information

19

Fig. S1.1. Effects of year and percent of grass area on the area burned.

Fig. S1.2. Effects of PDSI and percent of grass area on the area burned.

23
24
25
26
27

**Fig. S1.3.** Effects of livestock density and year on the area burned.

Fig. S1.4. Effects of PDSI and livestock density on the area burned.

38
39
40
41
42
43

44
45 **Fig. S2**: Number of sheep and goats in Kazakhstan from 1993 to 2014 (Food and Agriculture Organization, 2016).
46
47
48

49
50 **Fig. S3**. Number of cattle in Kazakhstan from 1993 to 2014 (Food and Agriculture Organization, 2016).
51
52
53