

Warming increases soil respiration in a carbon-rich soil without changing microbial respiratory potential

Marion Nyberg<sup>1</sup>, Mark J. Hovenden<sup>1</sup>

<sup>1</sup> School of Biological Sciences, University of Tasmania, Private Bag 55, Hobart, Tasmania, 7001, Australia

Corresponding author: Marion Nyberg (current affiliation University of British Columbia)

Mnybel@mail.ubc.ca

**Table Supplementary 1 Above ground biomass removal quantities at the Silver Plains Warming Experiment for all ambient dominant removal (ADR), warmed dominant removal (WDR), ambient random removal (ARR), and warmed random removal (WRR) plots.**

Treatment	Block	Above ground biomass dry per plot (g)
ADR	1	202.34
WDR	1	170.95
ARR	1	189.47
WRR	1	215.1
ADR	2	112.11
WDR	2	296.1
ADR	3	174.43
WDR	3	137.72
ARR	3	144.6
WRR	3	153.51
ADR	4	48.15
WDR	4	219.88
ADR	5	169.61
WDR	5	201.65
ARR	5	183.57
WRR	5	177.2
ADR	6	77.99
WDR	6	156.56
ADR	7	247.39

WDR	7	242.16
ARR	7	238.38
WRR	7	221.59
ADR	8	277.9
WDR	8	213.94

---