



## Warming increases soil respiration in a carbon-rich soil without changing microbial respiratory potential

Marion Nyberg<sup>1</sup>, Mark J. Hovenden<sup>1</sup>

<sup>1</sup>School of Natural Sciences, University of Tasmania, Hobart, 7001, Australia

5 *Correspondence to:* Marion Nyberg (current affiliation University of British Columbia) (mnybe1@mail.ubc.ca)

**Abstract.** Increases in global temperatures due to climate change threaten to tip the balance between carbon (C) fluxes, liberating large amounts of C from soils. Evidence of warming-induced increases in CO<sub>2</sub> efflux from soils has led to suggestions that this response of soil respiration (R<sub>s</sub>) will trigger a positive land C–climate feedback cycle, ultimately warming the earth further. Currently, there is little consensus about the mechanisms driving the warming-induced R<sub>s</sub> response, and there are relatively few studies from ecosystems with large soil C stores. Here, we investigate the impacts of experimental warming on R<sub>s</sub> in the C-rich soils of a Tasmanian grassy sedgeland, and whether alterations of plant community composition or differences in microbial respiratory potential could contribute to any effects. *In situ*, warming increased R<sub>s</sub> on average by 28% and this effect was consistent over time and across plant community composition treatments. In contrast, warming had no impact on microbial respiration in incubation experiments. Plant community composition manipulations did not influence R<sub>s</sub> or the R<sub>s</sub> response to warming. Processes driving the R<sub>s</sub> response in this experiment were, therefore, not due plant community effects and are more likely due to increases in belowground autotrophic respiration and the supply of labile substrate through rhizodeposition and root exudates. CO<sub>2</sub> efflux from this high-C soil increased by more than a quarter in response to warming, suggesting inputs need to increase by at least this amount if soil C stocks are to be maintained. These results indicate the need for comprehensive investigations of both C inputs and losses from C-rich soils if efforts to model net ecosystem C exchange of these crucial, C-dense systems are to be successful.

### 25 **1 Introduction**

Globally, more carbon (C) is stored in soils than the amount of C in the atmosphere and in plants combined (Canadell et al., 2007). Simple physiology suggests that soil respiration (R<sub>s</sub>) rates will increase as soil temperatures rise (Gillooly et al., 2001), stimulating CO<sub>2</sub> emissions from the soil – a response that has the potential to outweigh plant productivity responses to global warming and lead to a net loss of C from soils (Melillo et al., 2017). Recently, numerous studies have suggested that global warming is indeed disturbing the balance between ecosystem C inputs and outputs (Melillo et al., 2017). This suggests the possibility of a positive feedback whereby warming increases C efflux from soils, which accelerates climate change leading to further C losses and so on (Bridgman et al., 2008; Melillo et al., 2017; Bond-Lamberty et al., 2018). Importantly, it is possible that warming-induced C losses increase with soil C content such that it has been suggested that soils storing the most C could shift from C sinks to C sources (Cai et al., 2010; Wilson et al., 2016; Jasey et al., 2018).

Increases in respiration of soil organic carbon (SOC) as an effect of experimental warming occur almost universally (Rustad et al., 2001) however, increasing soil temperatures stimulate not only soil microbes and



enzyme activity, but also net primary productivity (NPP) and fresh C input from litterfall, and root exudations (Rustad et al., 2001), enhancing substrate availability for microbial respiration (Lu et al., 2013; Wang et al., 2017).  
40 Warming effects have also been demonstrated to drive microbial priming, whereby decomposition is enhanced through increased input of labile C compounds (van der Wal and de Boer, 2017). Despite this, greater above ground plant biomass is not directly linked to immediate or long-term increases in the storage of SOC and hence the mechanisms driving the response of Rs to warming are uncertain (Jackson et al., 2017).

Effects of temperature on environmental factors such as soil moisture, substrate availability and  
45 evapotranspiration also influence and mediate rates of decomposition of SOM, and efflux of CO<sub>2</sub> (Davidson et al., 2000; Eliasson et al., 2005; Lu et al., 2013). These effects include extension of growing seasons and shifts in species composition and community structure (Chen et al., 2016). Considering this, changes in plant community composition, and subsequent shifts in functional traits have the potential to influence the quantity and quality of organic matter in the soil, as well as the physical soil structure (Metcalf et al., 2011). This suggests that there is  
50 potential for the response of soil C dynamics to warming to be partially or even wholly dependent upon changes to plant community composition (Jackson et al., 2017).

Both experimental and global warming have impacts on soil water availability, which is itself a primary determinant of Rs (Schimel et al., 1994). Following a unimodal relationship, respiration is highest at an intermediate (35-50% by volume) soil water content (SWC), which stimulates microbial activity and enhances  
55 above and below ground labile C inputs (Chou et al., 2008; Zhou et al., 2010; Wang et al., 2017). Anaerobic conditions in wet and flooded soils suppress microbial activity, slowing decomposition of SOM (Davidson and Janssens, 2006). Similarly, low SWC can have a similar effect by reducing microbial activity, restricting soil respiration (Carey et al., 2016). As warming generally leads to lower soil water content (Zhang et al., 2013; Li et al., 2017), the impact on Rs depends upon the underlying soil water content, increasing respiration of wet soils  
60 but reducing respiration in drier soils (Almagro et al., 2009). Essentially, the effect of warming on SWC could either offset or exacerbate direct warming effects on soil respiration, potentially disturbing the entire global C balance.

Substrate availability is another factor that is affected by warming, and thus has the potential to shift the temperature sensitivity of SOM decomposition (Davidson and Janssens, 2006). Largely, increased temperatures  
65 lead to the loss of physical or chemical protection of SOM, and thus enhanced microbial respiration of soil organic carbon (SOC) (Davidson and Janssens, 2006). Partitioning SOC into pools as a function of recalcitrance and residence time assists with analysing effects of environmental manipulations on long-term C storage (Pendall et al., 2011). As C inputs to the soil and consequently into these various pools occur in response to the interplay between rates of NPP, decomposition, climatic conditions and soil characteristics (Ontl and Schulte, 2012), the  
70 fate of SOC is either transformation into highly recalcitrant humus, important for the stabilisation and long-term storage of SOC, or loss to the atmosphere as CO<sub>2</sub>. Thus, factors such as oxygen availability, substrate quantity and quality, nutrient limitation and activity of extracellular enzymes are key to the soil respiration response. Carefully controlled laboratory incubations are necessary to eliminate confounding factors and pinpoint the mechanisms driving responses observed in the field (Davidson and Janssens, 2006). Ultimately, distinguishing  
75 between the potential driving factors is vital for our ability to model future C fluxes and to extend the observations from field experiments more widely.



To understand the consequences of warming on soil C dynamics and particularly  $R_s$ , it is necessary to distinguish between warming-related increases in  $R_s$  that are simply due to an increase in the biochemical response of  $R_s$  to temperature, and potential alterations of the temperature sensitivity of  $R_s$  caused by climate warming. The increase in  $R_s$  with rising temperature has been widely documented (Luo et al., 2001; Rustad et al., 2001), however, the temperature sensitivity of  $R_s$  in soils that have undergone experimental warming is much more variable (Song et al., 2014; Carey et al., 2016). Shifts in the temperature sensitivity under warming are likely to be driven by both changes in microbial community composition and changes in the physical and chemical properties of the soil (Davidson and Janssens, 2006). Additionally, effects of warming such as soil drying affect various ecosystem processes and thus might shift the temperature response of  $R_s$  (Carey et al., 2016; Moinet et al., 2018). The effect of temperature on  $R_s$  is thus complex, and there are a number of biotic and abiotic factors influencing the response of SOM decomposition to warming. Until these various influences are characterised accurately, projecting future soil C emissions will remain problematic.

Although measurements of soil respiration *in situ* often demonstrate warming-related increases, the mechanisms behind this response cannot be revealed by simple field observations (Davidson and Janssens, 2006). In particular, it is difficult to distinguish changes in microbial community composition and functioning in response to warming from *in situ* measurements alone. These changes include acclimation (Luo et al., 2001) or adaptation (Bradford et al., 2008), encompassing both physiological and genetic changes within individuals and species, changes in community structure (Sheik et al., 2011) and a shift towards microbial use of slowly decomposing C (Bracho et al., 2016). Hence, a shift in temperature sensitivity of SOM decomposition is likely to be driven by warming through a change in microbial respiratory potential, expressed as the  $\text{CO}_2$  mineralisation rate.

Here, we use a manipulative experiment to examine the potential influences of climate change, specifically warming and plant community composition, on soil C dynamics. We examine soil respiration responses both *in situ* and in laboratory incubation experiments to disentangle the mechanisms involved in the response of soil respiration to both warming and manipulation of the plant community. Specifically, we ask the following questions:

- 1.) Does warming increase soil respiration in a Tasmanian C-rich soil?
- 2.) If so, is this due to changes in microbial respiratory potential?
- 3.) Does altering plant community composition change the response of soil respiration to warming?

105

## 2 Materials and Methods

### 2.1 Study site

All field measurements and soil samples were taken at the Silver Plains warming experiment in the Tasmanian central plateau, Australia (42°09'S, 147°08'E; 890 m a.s.l.). The site is a natural grassy sedge-land with an average summer temperature of 16°C, average winter temperature of 6°C and average annual rainfall of 720 mm (BOM, 2018). Soil at the site is peaty, being an organosol containing on average 8 kg C m<sup>-2</sup> in the top 10 cm. The vegetation at the site is heavily grazed year-round by a range of native vertebrate herbivores, including wallabies,

110



pademelon and wombats, as well as by feral fallow deer, resulting in an extremely low vegetation stature of a few cm, with the exception of inflorescences which can extend up to 30 cm above the ground.

115

## 2.2 Experimental design

The experiment was set up in the 2014 austral winter as a fully orthogonal, two-factor random block design, with warming and species removal as fixed factors across eight replicate blocks (Fig. 1). The experiment consists of forty 2 x 2 metre plots, with 3 metres between each plot, of which 20 were warmed using hexagonal polycarbonate open-top chambers (OTC) with an internal diameter of 1.5 m, with the remainder of being unwarmed, ambient plots. To investigate the impact of altering plant community composition, the dominant species, *Poa gunnii*, was removed by plucking in one warmed and one ambient plot (henceforth “dominant removal” plots) in each block. One warmed and one ambient plot in each block was left untouched (henceforth “no removal” plots). To control for possible effects of removing biomass during the dominant species removal treatment, we also removed biomass from one additional warmed and unwarmed plot in every second block, removing the same amount of biomass as was removed from the “dominant removal” plots in the same block but with this biomass removed randomly from across the plot, rather than from a single species (henceforth “random removal” plots). Plant biomass was removed in the spring and summer of 2014/15 by gently removing small plants by hand and by repeatedly clipping larger plants to ground level until green shoots no longer emerged. The amount of biomass removed in each plot is presented in Table S1. After the initial removal treatment, all plots were left undisturbed until the following spring, at which time all plots were surveyed to determine whether removed plants had re-established. As removed plants had not re-established at this time, no further removal occurred. Plant biomass was not measured directly in the plots in order to reduce disturbance. However, measures of vegetation cover and height indicated that the vegetation in removed plots had recovered completely within two years and were very similar to untouched control plots by this time, except in terms of species composition.

120  
125  
130  
135




Figure 1. Conceptual diagram for the experimental design of the Silver Plains warming experiment. Each block contains a warmed and unwarmed plot with no species removed (WNR) and (ANR) respectively; a warmed and unwarmed plot with the dominant species removed (WDR) and (ADR) respectively; and in every second block, i.e. in four blocks, there is a warmed and unwarmed plot with random biomass removal (WRR) and (ARR) respectively.

Air temperature at 5 cm height and soil temperature at 5 cm depth in each plot was logged continuously with iButton dataloggers. Over the entire five year period, the warming treatment increased air temperature 5 cm above the soil surface by 1.56°C ( $P < 0.004$ ) and soil temperature at 5 cm depth by 1.29°C ( $P < 0.001$ ).

### 2.3 *In situ* methods

A 50 mm length of 100 mm diameter PVC pipe was inserted into the soil to a depth of 2 cm, extending 3 cm above ground height, within the centre 0.25 m<sup>2</sup> of each plot for soil respiration measurements. Soil respiration was measured with a CO<sub>2</sub>/H<sub>2</sub>O infrared gas analyser (IRGA) (Licor, model LI-6400) with attachment of a Licor 6400-09 soil chamber, which attached to PVC pipes. Soil respiration was measured *in situ* monthly from August 2017 to June 2018. On each occasion, three measurements of *in situ* soil respiration, defined as the CO<sub>2</sub> efflux rate, were made in each plot. The average value of these three measurements was used in subsequent analyses. Soil temperature and moisture in each plot were measured at the exact same time as the soil respiration measurements on each occasion. Soil temperature was measured with a soil thermocouple probe (LiCor 6000-09TC) attached to the LI-6400. Volumetric soil water content (SWC) was estimated at 5 locations in each plot using a hand held TDR probe at 0-5cm depth. The 5 separate measurements of SWC were then averaged to obtain one SWC value per plot on each measuring occasion.



Six randomly placed soil samples, amounting to a total of approximately 25-30 g fresh weight, were collected from each plot using a 1.5 cm diameter hand corer to a depth of 5 cm below ground level, twice throughout the year. Samples were collected on the 02/03/18, representing the end of summer, or growing season soil, and on the 25/06/18, representing winter soils.

#### 2.4 Laboratory incubations

Soil cores collected *in situ* were immediately placed on ice for return to the laboratory, where they were refrigerated (4°C) overnight. The following day, the samples were composited at the plot level and sieved through a 4 mm sieve for one minute to remove leaves and large roots. A 10 g fresh-weight sub-sample was removed and oven dried from each composite sample for the determination of total soil C. Each subsample was ground to a powder in a Retsch Mixer Mill (MM200, Retsch GmbH, Haan) and then C content was analysed by combustion in a Perkin Elmer 2400 Series II Elemental Analyser (Perkin Elmer Australia, Melbourne). The remaining soil was used immediately for laboratory incubations to determine microbial respiration, as detailed below.

Microbial respiration as a function of temperature was determined by incubation using soils sampled in the Silver Plains warming experiment at the end of summer and in mid-winter 2018. For each plot, three replicate samples weighing four to eight grams from the composite sample were placed in 100 mL specimen jars, each of which was incubated at a different temperature. Each sample was wetted to bring them to 90% of field capacity for winter soils and 60% of field capacity for summer soils to represent prevailing soil moisture conditions in each respective season. Once water was added to all soil samples, specimen jars were placed in 500 ml preserving jars with tightly fitting lids containing a septum to allow gas headspace samples to be collected by syringe. Jars were stored in dark incubation cabinets at temperatures at one of 10, 17 or 25°C, with one sample from each plot at each temperature. Headspace gas of jars were sampled (20 ml) using a syringe on days 1, 2, 4, 5, 7, 9, 12, 15, 19, 23, 29, 35, 49, 56, 63. After extracting samples from each jar, headspace samples were analysed for CO<sub>2</sub> concentration, representing soil respiration, and microbial respiratory potential was thus defined as the rate of CO<sub>2</sub> release. To analyse headspace gas, samples were injected directly into an infrared gas analyser (LI-6262, Li-Cor, Lincoln, NE). After measurements were taken and analysed, jars were ventilated for 20 minutes and headspace gas equilibrated with atmospheric air. Following this, lids were replaced and headspace gas was sampled and analysed again to obtain starting CO<sub>2</sub> concentration for each jar. C mineralisation over the sample period was calculated from the increase in headspace CO<sub>2</sub> concentration.

Total C mineralisation over the entire incubation period was simply the sum of the amount of C mineralised over each sample period. Daily C mineralization results (dC/dt) were analysed using non-linear curve fitting routines in R (version 3.4.3, R Core Team, 2017), with a single pool plus constant model (Pendall et al., 2011) to estimate the size of the labile C pool ( $C_0$ ), the intrinsic decay constant of the labile pool ( $k$ ), and the intrinsic decay constant of the stable C pool ( $Y_0$ ):

$$\frac{dC}{dT} = C_0 k e^{-kt} + Y_0$$

(1)



## 2.5 Data analysis

195 Field soil respiration rates were analysed using a 2-factor repeated measures ANOVA with warming and removal  
as the fixed factors. Since soil temperature (Ts) and SWC are known controllers of Rs and varied substantially  
over the year, we also analysed field Rs with a 2-factor ANCOVA with Ts and SWC and the interaction between  
Ts and SWC as covariates. Treatment means were calculated as least-squares means using the lsmeans package  
to account for the influences of covariates (Russel V. Lenth, 2016). Treatment effects on SWC and Ts were  
200 analysed using 2-factor repeated measures ANOVA exactly as for Rs.

Because there was a significant influence of warming on Rs, we created a separate model of the influence of SWC  
and Ts on *in situ* Rs for warmed and unwarmed plots. Since the respiration temperature relationship is best  
described by an Arrhenius-type function (Fang & Moncrieff 2001), we used multiple regression techniques to fit  
an exponential relationship to Rs and SWC, Ts and the interaction between Ts and SWC. Such a non-linear  
205 relationship fitted the observed data far better than a linear model, as compared by the Akaike information criterion  
corrected for finite sample size.

Total cumulative CO<sub>2</sub> emitted in laboratory incubations, C<sub>a</sub>, k, and Y<sub>0</sub> for each season were compared using three-  
factor analysis of variance ANOVA for both summer and winter soils with incubation temperature, warming and  
species removal as fixed factors, including all interactions. Seasonal differences were also analysed using four-  
210 factor ANOVA, with season also included as a fixed factor along with warming effect, removal and incubation  
temperature.

All statistical analyses were carried out in R (version 3.4.3). Data were checked for heteroscedasticity and  
normality and the required transformations were made using the Box Cox power and logarithmic transformations.  
Significant treatment effects were further analysed using Tukey's HSD *post hoc* comparisons.

215

## 3 Results

### 3.1 *In situ* soil respiration

#### 3.1.1 CO<sub>2</sub> efflux

Experimental warming drove a significant increase in soil respiration over the course of the year ( $F_{1,12}=58.48$ ,  
220  $P<0.001$ ; Table 1) but there was no significant influence of the species removal treatment, so neither the dominant  
nor random removal treatments were different to the untouched plots ( $F_{2,12}=1.1$ ,  $P=0.36$ ), nor was there a warming  
 $\times$  removal interaction effect on CO<sub>2</sub> efflux ( $F_{2,12}=0.14$ ,  $P=0.87$ ). As expected, time of year had a strong effect of  
CO<sub>2</sub> efflux ( $F_{6,12}=11.84$ ,  $P<0.001$ ), with the highest rates,  $13.23 \pm 0.37 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ , in summer, decreasing  
through to  $1.4 \pm 0.06 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  in winter (Table 1). Despite the strong variation in C efflux rates across  
225 the year, there was no significant interaction between month and warming ( $F_{5,12}=1.17$ ,  $P=0.38$ ), indicating that  
the warming effect was consistent across the year.



**Table 1. The impact of experimental warming on soil CO<sub>2</sub> efflux, soil temperature and soil water content in the Silver Plains Warming Experiment from August 2017 to June 2018. Values shown are means with standard errors in parentheses (n=20).**

Month	Treatment	CO <sub>2</sub> efflux	Soil temperature	SWC
		μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup>	°C	%
August	Ambient	1.8 (0.1)	5.1 (0.1)	34.3 (0.6)
	Warmed	2.3 (0.1)	5.3 (0.1)	27.4 (0.5)
November	Ambient	8.0 (0.3)	16.0 (0.3)	19.8 (1.0)
	Warmed	12.2 (0.4)	16.9 (0.3)	16.5 (0.7)
January	Ambient	11.6 (0.3)	17.1 (0.2)	19.3 (0.4)
	Warmed	13.2 (0.4)	17.8 (0.2)	17.5 (0.6)
February	Ambient	7.6 (0.2)	18.9 (0.1)	13.3 (0.2)
	Warmed	12.9 (0.5)	18.8 (0.1)	12.6 (0.3)
March	Ambient	6.1 (0.1)	13.3 (0.3)	20.6 (0.3)
	Warmed	9.0 (0.2)	13.5 (0.2)	15.8 (0.3)
April	Ambient	4.7 (0.1)	12.6 (0.2)	13.7 (0.3)
	Warmed	7.3 (0.1)	13.2 (0.1)	11.1 (0.3)
May	Ambient	4.0 (0.1)	10.9 (0.3)	1225 (0.3)
	Warmed	5.7 (0.1)	12.3 (0.2)	9.8 (0.2)
June	Ambient	1.4 (0.1)	3.3 (0.2)	52.0 (1.7)
	Warmed	1.7 (0.1)	4.1 (0.1)	45.3 (1.2)

230

### 3.1.2 Soil temperature



Time of year had a strong impact on soil temperature ( $F_{6,12}=27.61$ ,  $P<0.001$ ), which varied from  $3.33 \pm 0.18$  °C to  $18.89 \pm 0.14$ °C over the study period. Experimental warming had a significant impact on soil temperature increasing soil temperature at 5 cm depth by  $0.55$ °C on average ( $F_{1,12}=7.31$ ,  $P=0.02$ ). This impact was sustained over the year with no significant month x warming interaction ( $F_{5,12}=0.88$ ,  $P=0.52$ ), indicating that the warming chambers had a similar effect on soil temperature across the year. Neither removal treatment, i.e. neither dominant nor random biomass removal ( $F_{2,12}=1.99$ ,  $P=0.18$ ), nor warming x removal interactions ( $F_{2,12}=0.45$ ,  $P=0.65$ ) affected soil temperature. Thus, the warming treatment increased soil temperatures consistently over the year and across the species removal treatments.

### 3.1.3 Soil water content

Soil water content (SWC) also varied over the year ( $F_{6,12}=6.21$ ,  $P=0.003$ ) reflecting precipitation patterns at Silver Plains (BOM, 2018). Over the course of the year, SWC ranged from  $9.83 \pm 0.17\%$  to  $52 \pm 1.69\%$ , with moisture levels decreasing from winter 2017 through to autumn 2018, and then steeply increasing again in winter 2018 (Table 1). Experimental warming significantly decreased SWC throughout the year by 3% on average ( $P<0.001$ ), which is expected considering the drying effect of warming. However, the impact of warming on SWC depended upon the month, as indicated by a significant sampling month x warming effect ( $F_{5,12}=6.09$ ,  $P=0.005$ ). Warming had the greatest effect on SWC in August 2017 and June 2018, when SWC was highest and soil temperature was lowest. SWC in these winter months was substantially higher than during the rest of the year, with SWC on average 34 and 58% higher in August and June respectively, than the overall mean value ( $21.3 \pm 0.5\%$ ). The proportional reduction in SWC due to warming in these months was nearly two-times the yearly average. Otherwise, the warming effect was similar between sampling months. There was no significant influence of the removal treatment, i.e. neither the dominant nor random removal treatments were different to the untouched plots ( $F_{2,12}=0.23$ ,  $P=0.8$ ), nor was there a warming x removal interaction effect on SWC ( $F_{2,12}=0.52$ ,  $P=0.61$ ), again indicating that plant species removal did not alter the influence of the warming treatment.

### 3.1.4 Relationships between environmental factors and CO<sub>2</sub> efflux

Both soil temperature ( $F_{1,33}= 33.62$ ,  $P<0.001$ ) and SWC ( $F_{1,33}= 5.95$ ,  $P=0.02$ ) were strong controllers of soil CO<sub>2</sub> efflux over the year at Silver Plains (Fig. 2). However, treatment effects on these abiotic factors alone were insufficient to explain the higher C efflux in warmed plots, as ANCOVA indicated that the warming treatment still induced significant increases in CO<sub>2</sub> efflux when variation in soil T and SWC were accounted for ( $F_{1,33}= 44.83$ ,  $P<0.001$ ). Thus, the warming treatment increased soil CO<sub>2</sub> efflux independently of its effects on soil temperature and SWC (Fig. 2). Across the whole year LS mean CO<sub>2</sub> efflux rates for ambient soils was  $6.07$  ( $C.I.=5.69,6.45$ )  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  but  $8.48$   $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  ( $C.I.=8.09,8.86$ ) for warmed soils, amounting to a warming-induced increase of 28% at a common soil temperature and SWC. As CO<sub>2</sub> efflux measurements spanned a large variation in both soil T and SWC, it was possible to discern a trend whereby the stimulation of C efflux by warming became more pronounced as soil temperature increased (Fig. 2). Neither removal, i.e. neither dominant nor random biomass removal ( $F_{2,33}=0.89$ ,  $P=0.42$ ), nor a warming x removal interaction ( $F_{2,33}=0.57$ ,  $P=0.57$ ) affected CO<sub>2</sub> efflux, as indicated by ANCOVA.




Figure 2. CO<sub>2</sub> efflux as a function of soil temperature and soil water content for warmed (W) and ambient (A) plots at Silver Plains from August 2017 to June 2018. Size of the point represents SWC %, with larger points corresponding to higher SWC. The regression lines indicate the relationship between CO<sub>2</sub> efflux and soil temperature at median SWC in ambient plots (solid line) and warmed plots (dashed line).  
 275

### 3.1.5 Models of CO<sub>2</sub> efflux

As ANCOVA indicated that soil CO<sub>2</sub> efflux at Silver Plains was significantly influenced by soil temperature, SWC and a strong warming effect, the relationship between these covariates and CO<sub>2</sub> efflux could be estimated separately for ambient and warmed treatments. First a general regression model of CO<sub>2</sub> efflux was fit and selected using model selection based on AICc. The most parsimonious and accurate model was one which included soil temperature (T<sub>s</sub>), SWC, and a SWC x T<sub>s</sub> interaction term (Int.term).  
 280

This model was then fit independently to ambient and warmed plots using the relative coefficient values, with 89% of the variance in CO<sub>2</sub> efflux explained in warmed plots Eq. (2) and 82% in ambient plots Eq. (3).

285 
$$\text{CO}_2 \text{ efflux}_{\text{ambient}} = e^{(-0.8+0.359 \log(\text{SWC})+0.115 (T_s)+0.003(\text{Int.term}))}$$

$$R_2 = 0.82 \tag{2}$$

$$\text{CO}_2 \text{ efflux}_{\text{warmed}} = e^{(-0.06+0.148 \log(\text{SWC})+0.124 (T_s)+0.002(\text{Int.term}))}$$


$$R_2 = 0.89 \tag{3}$$

290


Thus, it is possible to model CO<sub>2</sub> efflux across a range of soil temperature and SWC values in both ambient, unwarmed (Fig. 3A) and warmed conditions (Fig. 3B). From these plots, it is possible to determine that while the CO<sub>2</sub> efflux rate increases more steeply with rising temperature in warmed plots than in unwarmed plots, the way



in which it does so is also dependent upon the SWC (Fig. 3A and B). Thus, the impact of experimental warming  
295 on soil CO<sub>2</sub> efflux was greatest in warm ( $T > 15^{\circ}\text{C}$ ) relatively dry conditions (SWC < 30%; Fig. 4).



**Figure 3.** CO<sub>2</sub> efflux ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) modelled as a function of soil temperature and SWC in A.) ambient plots and B.) warmed plots. Colour indicates predicted CO<sub>2</sub> efflux values and field observations are shown as individual points. Regions beyond the observed range of CO<sub>2</sub> efflux rates are shown in grey.



300

Figure 4. Delta CO<sub>2</sub> efflux ( $\mu \text{ mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ). The amount of extra CO<sub>2</sub> that is likely to be released due to warming as a function of soil temperature ( $^{\circ}\text{C}$ ) and SWC (%). Data points represent actual measurements and colour indicates predicted CO<sub>2</sub> efflux. Points on the contour graphs are field observations and regions beyond field observations, and thus where CO<sub>2</sub> cannot be predicted, are greyed out.

305

### 3.2 Laboratory incubations


#### 3.2.1 Total C mineralisation

To determine whether experimental treatments altered potential microbial respiration, soil samples were collected in summer and winter for laboratory incubations. These incubations allowed the temperature sensitivity of soil respiration, the size of the labile C pool ( $C_a$ ) and its decay constant ( $k$ ) to be assessed, as well as estimating the decay constant of the more resilient stable C pool ( $Y_0$ ) to be assessed in constant, optimal conditions. From soils collected in summer, the total amount of C mineralised increased substantially as an effect of incubation temperature, however there were no effects of either the warming or removal treatments. On average, soil incubated at  $17^{\circ}\text{C}$  for two months emitted 48% more C than at  $10^{\circ}\text{C}$ , and a further 22% at  $25^{\circ}\text{C}$  ( $F_{2,82}=80.9$ ,

310



315  $P < 0.001$ ; Fig. 5). From soils collected in winter, total C mineralised again only increased significantly as an  
 effect of incubation temperature, with on average a 26% increase in C emitted at 17°C from 10°C, and a further  
 27% increase at 25°C ( $F_{2,112}=49.56$ ,  $P < 0.001$ ; Fig. 5). Just like the situation with soil collected in summer, there  
 were no treatment effects on the total amount of C mineralised from winter soils ( $F_{1,112}=0.04$ ,  $P=0.84$ ). Between  
 seasons, winter soils emitted on average 24% less C than summer soils ( $F_{1,196}=33.66$ ,  $P < 0.001$ ), most likely  
 320 because of the higher SWC used for the winter soils, and neither removal treatment, i.e. neither dominant nor  
 random biomass removal ( $F_{2,196}=0.67$ ,  $P=0.51$ ), nor warming significantly affected total C mineralised overall  
 ( $F_{1,196}=0.01$ ,  $P=0.92$ ).



325 **Figure 5. Total C mineralised in summer and winter from soils in no removal (NR), (random removal (RR), and dominant removal (DR) plots at incubation temperatures, 10, 17 and 25°C for warmed (W) and ambient (A) treatments.**

### 3.2.2 Labile C

In summer soil, incubation temperature significantly increased the size of  $C_a$  on average by 50% from 10°C to 17°C, and by a further 18% at 25°C ( $P < 0.001$ ) (Fig. 6A). There were no treatment effects on the size of  $C_a$ . Winter  
 330 soil incubations reflect similar results to those for summer soils, with a 27% increase in  $C_a$  pool size from 10°C to 17°C, and a further 27% increase to 25°C ( $P=0.001$ ). As with summer soil there were no treatment effects. Overall, season had no effect on  $C_a$ , however incubation temperature increased  $C_a$  across the two seasons of 36% from 10°C to 17°C and a further 24% at 25°C ( $P < 0.001$ ).



The intrinsic decay constant of the labile pool ( $k$ ) in summer soil was not affected by incubation temperature  
335 ( $F_{2,82}=0.39$ ,  $P=0.68$ ), the warming ( $F_{1,82}=0.06$ ,  $P=0.8$ ), or removal treatments, i.e. neither dominant nor random  
biomass removal ( $F_{2,82}=0.31$ ,  $P=0.73$ ), was significantly influenced by an interaction between warming and  
species removal ( $F_{2,82}=3.14$ ,  $P=0.05$ ) (Fig. 6C). In ambient plots, removing the dominant species tended to  
increase  $k$ , however, in warmed plots, the opposite occurred. Post hoc analysis revealed the greatest differences  
in  $k$  were observed between warmed x no removal and warmed x dominant removal plots, and warmed x dominant  
340 removal and ambient x dominant removal plots. In winter, there were no treatment or incubation temperature  
effects on  $k$ , however  $k$  was on average 42% greater in summer ( $F_{1,196}=201.09$ ,  $P<0.001$ ).

### 3.2.3 Intrinsic decay constant of the stable C pool

From summer soil, the size of the stable C pool ( $Y_0$ ) also increased significantly ( $F_{2,82}=78.01$ ,  $P<2^{-16}$ ) as a function  
345 of incubation temperature with an average increase of 47% from 10°C to 17°C, and a further 20% at 25°C (Fig.  
6C). There were no treatment effects on the  $Y_0$  of summer soil. For winter soils, responses to treatments were  
similar to those of summer soils. There were no treatment effects, but incubation temperature increased  $Y_0$  on  
average by 27% from 10°C to 17°C, and a further 28% at 25°C ( $F_{2,112}=45.9$ ,  $P<0$ ). Overall  $Y_0$  was 39% higher in  
summer than in winter ( $F_{1,196}=137.61$ ,  $P<0.001$ ), and incubation temperature also significantly increased  $Y_0$   
350 overall, with on average a 38% increase from 10°C to 17°C, and a further 23% at 25 °C ( $F_{1,196}=107.28$ ,  $P<0.001$ ),  
however there were no treatment effects.

### 3.2.4 Proportion of total C that was labile

From summer soil, the proportion of total C that was from  $C_a$  was only affected by incubation temperature with  
355 on average a 49% increase from 10°C to 17°C, and a further 22% increase when incubated at 25°C ( $F_{2,82}=77.73$ ,  
 $P<0.001$ ; Fig. 6D). There were no treatment effects. Similarly, in winter, the proportion of total C that was  $C_a$   
increased only as a function of increasing incubation temperature, with on average a 24% increase from 10°C to  
17°C and a further 27% at 25°C ( $F_{2,112}=22.19$ ,  $P<0.001$ ). Overall, the proportion of total C that was  $C_a$ , increased  
substantially as a function of incubation temperature ( $F_{2,196}=67.94$ ,  $P<0.001$ ) with a 35% increase from 10°C to  
360 17°C, and a further 25% increase at 25°C, however there were no overall treatment effects.




Figure 6. A.) Labile C pool size ( $C_a$ ) in summer and winter soils in no removal (NR), random removal (RR) and dominant removal (DR) plots at incubation temperature, 10, 17 and 25°C for warmed (black) and ambient (white) treatments. B.) Intrinsic decay constant of the labile C pool ( $k$ ) both summer and winter soils in no removal (NR), random removal (RR), and dominant removal (DR) plots at incubation temperature, 10, 17 and 25°C for warmed (black) and ambient (white) treatments. C.) The intrinsic decay constant of the stable C pool ( $Y_0$ ) in summer and winter soils in no removal (NR), random removal (RR), and dominant removal (DR) plots at incubation temperature, 10, 17 and 25°C for warmed (black) and ambient (white) treatments. D.) Proportion of total C that is from the labile C pool ( $C_a$ ) in both summer and winter soils in no removal (NR), random removal (RR), and dominant removal (DR) plots at incubation temperatures 10, 17 and 25°C for warmed (black) and ambient (white) treatments.

### 3.3 Total soil C content

Overall, irrespective of removal treatment, total soil C % averaged  $19.2 \pm 0.4$  ( $P < 0.001$ ). C % was  $18.7 \pm 0.7$  and  $19.7 \pm 0.6$  in ambient and warmed soils respectively, however there were no significant treatment effects.

All incubation results were also analysed per gram of soil C but results were essentially identical to those expressed per gram of soil dry weight, above.

## 4 Discussion

The main aim of this study was to investigate whether warming increases  $R_s$  *in situ*, and whether any observed treatment effects were due to an increased ability of the soil microbial community to mineralise SOC. Additionally, we investigated whether manipulating plant community composition affected the  $R_s$  response to warming. Results



demonstrated strong warming-related increases in  $R_s$  *in situ*, however there were no warming effects on microbial respiratory potential. Additionally, the warming treatment increased soil temperature and decreased soil water content significantly *in situ*, however the warming effect on  $R_s$  was greater than expected considering the impact it had on these abiotic factors. Thus, warming increased  $R_s$  more than simply by increasing soil temperature and reducing soil water content. Manipulating the plant community by removing the dominant species or removing biomass had no impact on  $R_s$ , nor did these treatments influence the impact of warming on  $R_s$ . This suggests that the warming-induced increase in  $R_s$  was independent of any influences on plant community composition. Similarly, removal treatments did not affect microbial respiratory potential, however there was a complex warming and removal interaction that influenced the decay constant of the soil labile C pool ( $k$ ). Overall, the results from this study suggest that as there was no change in microbial respiratory potential, the observed increase in soil respiration *in situ* was largely an effect of altered plant activity in warmed plots.

#### 395 **4.1 Possible mechanisms leading to the warming-induced increase in soil respiration**

Warming increased  $R_s$  *in situ* over the course of the sampling period from November 2017 to June 2018. This increase in  $\text{CO}_2$  efflux observed from soils *in situ* encompasses the response of both microbial (heterotrophic) respiration, and plant root (autotrophic) respiration, and amounted to an average increase in soil C efflux of 28%. The observed increase of  $R_s$  in response to warming is in line with multiple other studies, although most of these focus on soils in the low- to mid-range of soil C stocks and in northern hemisphere locations (Lu et al., 2013; van Gestel et al., 2018). There are 4 possible mechanisms whereby could have increased in  $R_s$ : 1.) Increased temperature sensitivity of  $R_s$ ; 2.) Influence through change in plant community composition; 3.) Enhanced substrate supply through SOM and 4.) Plant induced alteration to soil microhabitat. The substantial  $R_s$  response to warming could be due to one or a combination of these processes and determining which were likely to be involved has significant ramifications for our ability to predict future soil C dynamics.

##### 4.1.1 Increased temperature sensitivity of $R_s$

One of the proposed mechanisms behind the increased  $R_s$  response to warming, and subsequent loss of soil C stores is an increase in the temperature sensitivity of  $R_s$ , i.e. increased decomposition of SOM (Kirschbaum, 1995). This response, mainly attributed to an increase in enzyme kinetics with temperature, is linked strongly to substrate availability (Davidson and Janssens, 2006). At Silver Plains, the overall significant increase in  $R_s$  rates from warmed plots *in situ* implied that the temperature sensitivity of  $R_s$  was higher under warming. The highest  $R_s$  rates were recorded during the growing season in spring and summer, suggesting primary productivity, microbial activity and environmental factors such as precipitation are likely to substantially influence respiration rates (Almagro et al., 2009). However, despite the strong dependence of  $R_s$  on soil water content and soil temperature, warmed plots had higher rates of C efflux from the soil under particular combinations of soil temperature and moisture (Section 3.1.5, Fig. 3). The restrictive effect of high soil water content and low soil temperature on  $R_s$  observed in this study is widely documented and due to the creation of anoxic conditions limiting microbial access to substrate (Schimel et al., 1994; Syed et al., 2006; Sierra et al., 2015). Hence the observed effect of soil water content and soil temperature on  $R_s$  was anticipated, however the degree to which



warming enhanced the response of  $R_s$  to temperature, was greater than expected. This observation could be explained by the greater effect of warming on air temperature than soil temperature at 5cm, thus considering most microbial activity occurs in the surface soil layers, it is possible the soil was warmed more than the amount measured, partially accounting for the large increase in  $R_s$ . Interestingly, the degree of stimulation by warming  
425 increased as soil temperature increased, i.e., there was a greater warming-induced stimulation of C efflux when soil was warm than when it was cold. This contrasts previous findings that indicate a greater warming effect on  $R_s$  at lower temperatures (Wang et al., 2014), and results from the incubation experiment in this study that reveal warming did not affect microbial respiration at any incubation temperature.

The large apparent increase in  $R_s$  observed *in situ* implied that warming possibly triggered an increase in microbial  
430 respiratory potential. Partitioning of  $R_s$  in incubation experiments allows the response of microbial respiration to warming to be observed under optimal conditions, controlling for soil water content and soil temperature. In stark contrast to the *in situ* observations, soil incubations revealed no differences in the temperature sensitivity of microbial respiration between warmed and ambient soil. There were no differences among treatments in the total C mineralisation rate measured under laboratory conditions, which would indicate that the ability of the soil  
435 microbial community to mineralise soil C was unchanged. This lack of any treatment effect was similar in winter and summer-collected soils, even though there was a strong seasonal effect on the  $\text{CO}_2$  respiration rate in incubated soils. Winter soils emitted significantly less  $\text{CO}_2$  than soils collected at the end of summer, a response attributed to decreased access to substrate as an effect of limited enzyme activity in cold temperatures (Suseela et al., 2012). Additionally, despite claims that the warming-induced increase in  $R_s$  is due to a strong, positive relationship  
440 between the average turnover time of labile C pools and mean annual air temperature (Trumbore et al., 1996), there was no difference in the size of the respired labile C pool ( $C_a$ ) between warmed and ambient soils, or between seasons. Therefore, this suggests that the warming treatment did not increase the temperature sensitivity of labile C decomposition. Additionally, there were no warming or removal treatment effects on the decay constant of the stable C pool ( $Y_0$ ), calculated from  $\text{CO}_2$  emission rates late in the incubation period. This implies that stable C,  
445 which is chemically and physically protected (Schlesinger, 1997), was not sensitive to warming, a response that contrasts results obtained elsewhere (Leifeld and Fuhrer, 2005; Hartley and Ineson, 2008). Thus, lack of a warming effect indicates that warming-induced increases in labile, or stable C temperature sensitivity are not driving the  $R_s$  response to warming observed *in situ*.

Essentially, the incubation studies revealed that four years of experimental warming had not altered either the  
450 potential for microbial respiration or its inherent temperature response, as soils incubated at the same temperature respired more or less at the same rate, regardless of whether they were collected from warmed or ambient plots. These results indicate that the warming-induced stimulation of  $R_s$  *in situ* was not due to changes in the inherent temperature response of microbial respiration. Considering the soil incubation experiment decoupled microbial respiration from plant activity and particularly C inputs, the lack of a warming treatment effect on C emissions in  
455 the incubation experiments is evidence that plants play a large role in the respiration response.

#### 4.1.3 Influence through change in plant species community composition



Plant community compositional change drives ecosystem responses to global changes, particularly when it involves shifts in the dominance or abundance of plant functional types (Bret-Harte et al., 2008). This is particularly true with  $R_s$  and global warming, as warming-related changes in plant functional types, and hence the resources they input to the soil, are highly likely to occur (Saleska et al., 2002). By investigating the effects of dominant species removal, random biomass removal, and warming on the response of respiration, there is scope to gain insight into future ecosystem dynamics under a changing climate. Removal of a dominant species from an ecosystem has promoted species diversity and altered ecosystem function, implying dominants reduce the establishment of other species (Wardle et al., 1999). Metcalfe et al., (2011) highlight the significant role that functional traits of the dominant species hold on many soil processes, including decomposition and respiration and hence one would expect to observe these effects in this study. Plants modify local soil conditions through root exudations of hormones, sugars, phenolics and amino acids, essentially structuring the rhizosphere microbial community composition. This means that changes in plant community composition have the potential to affect  $R_s$  and thus ecosystem functioning (Van Nuland et al., 2016), and hence the motive to investigate how  $R_s$  responds to the combination of warming and manipulated plant community composition. Results from Silver Plains demonstrate neither removal of the dominant plant species nor random removal of biomass had any effect on  $R_s$  or the temperature response of  $R_s$  *in situ*, and very little effect on microbial respiratory potential. This suggests that, in line with previous studies, temperature had greater control on  $R_s$  than variation in plant community composition (Duval and Radu, 2018). This result is surprising because despite removing the dominant species, there did not appear to be any functional shifts within the community, suggesting it may have been replaced by a functionally-similar species, or there was a compensatory response by functionally-different species (Bret-Harte et al., 2008). Differences in plant chemistry, morphology and physiology affect the quantity and quality of root and leaf litter, leading to changes in SOM decomposition rates, shifts in microbial respiratory potential and community structure (Van Nuland et al., 2016). However, the response of  $R_s$  *in situ* indicates that despite removal of the dominant species or random removal of biomass, community function was maintained.

Despite the absence of an effect on  $R_s$  *in situ* from manipulating species composition, microbial respiration dynamics indicated that warming and biomass removal (both random and dominant removal treatments) reduced the intrinsic decay constant of the labile C pool ( $k$ ) in summer soil. Interestingly, biomass removal had the opposite effect in ambient plots. In a previous clipping experiment, which is representative of biomass removal, a decrease in  $R_s$  due to clipping was explained as relocation of assimilates to shoots, reduction in the supply of photosynthates to roots, and thus decreased root respiration (Zhou et al., 2010). Hence there is likely to be less available substrate under warming and biomass removal scenarios, and  $k$  is therefore lower. Considering this, the interactive effect of warming and biomass removal on  $k$  is complex and requires further investigation to explore the mechanistic basis behind the response. The absence of an influence on  $R_s$  through variation in plant community composition suggests this mechanism is not driving the warming-induced  $R_s$  response to warming.

#### 4.1.4 Enhanced substrate supply

##### SOM



495 As SOM forms mainly from plant litter, warming related increases in both above and belowground primary  
productivity suggest supply of SOM will be greater under warming (Rustad et al., 2001; Lin et al., 2010; Wu et  
al., 2011), at least in systems that are not water limited. Additionally, experimental warming often increases leaf-  
drop, root-turnover and the subsequent decomposition of leaf and root litter (Lu et al., 2013), with the combined  
500 effects of warming and higher C inputs on respiration rates reported to be greater than the impact of either factor  
in isolation (Hopkins et al., 2014). Root and leaf litter have fast turnover times, implying they represent a major  
source of C for microbial decomposition. Therefore, an increase in the input of easily degradable C would promote  
microbial activity (Wan et al., 2005; Hogberg and Read, 2006), stimulating soil C efflux. Considering this, an  
increase in substrate supply seems like a conceivable explanation for the increase in  $R_s$  observed *in situ*. However,  
incubation experiments indicated no influence of warming on the total amount of C between warmed and ambient  
505 plots, or on the size of the labile pool ( $C_a$ ), or total C respired. This indicates that substrate supply and availability  
from plant biomass is similar in warmed and ambient plots. Previous investigations suggest that despite warming-  
related increases in litter quantity, enhanced respiration due to increased labile C concentration in soils is likely  
to offset additional C inputs (Lu et al., 2013), meaning changes to both inputs and losses of soil C could balance  
each other. Interestingly neither *in situ*  $R_s$ , nor total soil C or  $C_a$  was affected by plant community composition  
510 manipulations, suggesting substrate supply and availability was similar regardless of warming and removal  
treatments. This result contrasts with those from previous clipping experiments that demonstrated that biomass  
removal limits substrate supply (Wan and Luo, 2003; Xue et al., 2016). Overall this suggests that increased  
substrate supply through SOM is not a driving mechanism behind the warming-induced increase in  $R_s$  observed  
*in situ* although specific tests of this mechanism, such as through the use of stable isotope tracing, would be  
515 required to be confident.

#### 4.1.6 Plant induced alteration to soil microhabitat

The final mechanism that could be driving the warming-induced increases in  $R_s$  are plant induced alterations to  
the soil microhabitat. In this study, as in most, soil for incubations was not analysed as intact soil cores, rather  
520 being sieved and homogenised, altering the microhabitat conditions. This is potentially problematic, as it is  
assuming that rhizosphere processes are not influencing the overall  $R_s$  response. Previous studies have  
demonstrated the important role roots play in stabilising SOM (Hinsinger et al., 2009), with disturbed soils having  
a lower capacity to protect SOM due to mechanical disruption of macroaggregates, and hence C is more readily  
decomposed by microorganisms (Beare et al., 1994). Additionally, macrofauna such as earthworms and  
525 nematodes play an important role in the early stages of SOM decomposition (Wardle et al., 2004), and therefore  
the absence of these species from the incubations could have influenced the rates of C efflux. Considering this, it  
is possible that through homogenisation of soil in incubation studies, soil C dynamics and decomposition rates  
are confounded by disturbances to the soil microhabitat.

530

#### 4.3 Implications



Large C stocks within this type of peaty habitat imply they are globally important for the C cycle, thus understanding potential losses are immensely important for the global C budget. Previous studies on the response of Rs to warming have been largely centred around northern hemisphere sites, and with that there has been large unexplained variability in the response (van Gestel et al., 2018). This implies that the mechanisms behind the Rs response to warming are poorly characterised. Our results indicate that warming-related increases in CO<sub>2</sub> efflux from C-rich soils in grassy peatlands are expected in the future, however microbial respiratory potential is not the driving factor, and thus there is a strong link to plant activity and C inputs. Moreover, the results indicate that the impact of warming on soil CO<sub>2</sub> efflux is strongly dependent upon both soil temperature and moisture conditions, improving the confidence that current and future soil CO<sub>2</sub> efflux can be modelled from these variables. However, the lack of observations in certain combinations of soil temperature and moisture mean that predictions using the models presented here should be limited to the observed range. Future work should test the generality of these models in previously unobserved combinations of soil moisture and temperature.

Predictions regarding future climate conditions require a more comprehensive mechanistic understanding of temperature and decomposition relationships, especially considering the global variation in these relationships. Further investigation into the role of inputs is required, as warming could be driving increases in inputs, thereby balancing the accelerated C efflux and preventing net loss of C from soils. Alternatively, warming could lead to depletion of huge stores of C. This effect is no doubt subject to great variation depending on the ecosystem and hence the necessity to examine the response, accounting for heterogeneity in soil and vegetation types worldwide. Most importantly, this study revealed that C inputs through root exudates and root respiration were the two mechanisms most likely to be driving the Rs response to warming. Thus, more research into the influence of root exudates and root respiration on Rs, particularly under warming, will provide a more comprehensive insight to the Rs response. Ultimately, thorough investigations into the whole ecosystem C exchange is required to advance understanding into how warming will affect rates of inputs and outputs.

The increase in Rs in response to warming observed here is in line with previous experimental warming studies, although few have been conducted in C-rich soils. Thus, the results from this study contribute directly to a field of knowledge that is currently underrepresented. Despite a strong warming effect, there appear to be no significant effects of plant community manipulation, suggesting that warming exerts more of an influence on CO<sub>2</sub> efflux from soils than differences in plant communities. Additionally, results suggest that the microbial respiratory potential in this system is not altered by experimental warming and hence cannot be decoupled from plant activity if we are to enhance our ability to predict C cycling dynamics in a warmer climate. Current findings suggest warming is likely to trigger a positive feedback cycle whereby increases in global temperatures will enhance CO<sub>2</sub> efflux from soils, subsequently warming the earth further. As the huge C stocks in the soil have the potential to either amplify or attenuate global warming, the impacts of climate change on soil C dynamics require urgent investigation. A more comprehensive representation of ecosystem C exchange is needed, as well as the mechanisms involved, if we want to decrease CO<sub>2</sub> efflux from soils and ensure these huge C sinks are stabilised, or potentially even increased such that the biosphere can sequester more atmospheric CO<sub>2</sub> and help to stabilise the climate.



**Author contribution:** Marion Nyberg and Mark Hovenden were responsible for conceptualization and project  
570 administration. Marion Nyberg performed data acquisition and curation, formal analysis and writing. Mark  
Hovenden developed the methodology and provided supervision.

#### Competing interest

The authors declare that they have no conflict of interest

575

#### Acknowledgements

We would like to thank Meagan Porter and Rose Brinkhoff for help setting up the experiment and the Tasmanian  
Land Conservancy for permission to use the land for the experiment. This research did not receive any specific  
grant from funding agencies in the public, commercial, or not-for-profit sectors.

580

#### References

- Almagro, M., López, J., Querejeta, J. I., and Martínez-Mena, M.: Temperature dependence of soil CO<sub>2</sub> efflux is  
strongly modulated by seasonal patterns of moisture availability in a Mediterranean ecosystem, *Soil Biology  
and Biochemistry*, 41, 594-605, <https://doi.org/10.1016/j.soilbio.2008.12.021>, 2009.
- 585 Beare, M. H., Hendrix, P. F., Cabrera, M. L., and Coleman, D. C.: Aggregate-protected and unprotected organic  
matter pools in conventional- and no-tillage soils, *Soil Science Society of America Journal*, 58,  
[10.2136/sssaj1994.03615995005800030021x](https://doi.org/10.2136/sssaj1994.03615995005800030021x), 1994.
- Bengtson, P., Barker, J., and Grayston, S. J.: Evidence of a strong coupling between root exudation, C and N  
availability, and stimulated SOM decomposition caused by rhizosphere priming effects, *Ecology and Evolution*,  
590 2, 1843-1852, [10.1002/ece3.311](https://doi.org/10.1002/ece3.311), 2012.
- Bond-Lamberty, B., Bailey, V. L., Chen, M., Gough, C. M., and Vargas, R.: Globally rising soil heterotrophic  
respiration over recent decades, *Nature*, 560, 80-83, [10.1038/s41586-018-0358-x](https://doi.org/10.1038/s41586-018-0358-x), 2018.
- Bond-Lamberty, B., and Thomson, A.: Temperature-associated increases in the global soil respiration record,  
*Nature*, 464, 579-582, [10.1038/nature08930](https://doi.org/10.1038/nature08930), 2010.
- 595 Boone, R. D., Nadelhoffer, K. J., Canary, J. D., and Kaye, J. P.: Roots exert a strong influence on the  
temperature sensitivity of soil respiration, *Nature*, 396, 570-572, [10.1038/25119](https://doi.org/10.1038/25119), 1998.
- Bracho, R., Natali, S., Pegoraro, E., Crummer, K. G., Schädel, C., Celis, G., Hale, L., Wu, L., Yin, H., Tiedje, J.  
M., Konstantinidis, K. T., Luo, Y., Zhou, J., and Schuur, E. A. G.: Temperature sensitivity of organic matter  
decomposition of permafrost-region soils during laboratory incubations, *Soil Biology and Biochemistry*, 97, 1-  
600 14, [10.1016/j.soilbio.2016.02.008](https://doi.org/10.1016/j.soilbio.2016.02.008), 2016.



- Bradford, M. A., Davies, C. A., Frey, S. D., Maddox, T. R., Melillo, J. M., Mohan, J. E., Reynolds, J. F., Treseder, K. K., and Wallenstein, M. D.: Thermal adaptation of soil microbial respiration to elevated temperature, *Ecology Letters*, 11, 1316-1327, 10.1111/j.1461-0248.2008.01251.x, 2008.
- 605 Bret-Harte, M. S., Mack, M. C., Goldsmith, G. R., Sloan, D. B., Demarco, J., Shaver, G. R., Ray, P. M., Biesinger, Z., and Chapin, F. S.: Plant functional types do not predict biomass responses to removal and fertilization in Alaskan tussock tundra, *Journal of Ecology*, 96, 713-726, 10.1111/j.1365-2745.2008.01378.x, 2008.
- Bridgman, S. D., Pastor, J., Dewey, B., Weltzin, J. F., and Updegraff, K.: Rapid carbon response of peatlands to climate change, *Ecology*, 89, 3041-3048, 10.1890/08-0279.1, 2008.
- 610 Cai, T., Flanagan, L. B., and Syed, K. H.: Warmer and drier conditions stimulate respiration more than photosynthesis in a boreal peatland ecosystem: analysis of automatic chambers and eddy covariance measurements, *Plant, Cell & Environment*, 33, 394-407, 10.1111/j.1365-3040.2009.02089.x, 2010.
- Canadell, J. G., Kirschbaum, M. U. F., Kurz, W. A., Sanz, M.-J., Schlamadinger, B., and Yamagata, Y.: Factoring out natural and indirect human effects on terrestrial carbon sources and sinks, *Environmental Science & Policy*, 10, 370-384, <https://doi.org/10.1016/j.envsci.2007.01.009>, 2007.
- 615 Carey, J. C., Tang, J., Templer, P. H., Kroeger, K. D., Crowther, T. W., Burton, A. J., Dukes, J. S., Emmett, B., Frey, S. D., Heskell, M. A., Jiang, L., Machmuller, M. B., Mohan, J., Panetta, A. M., Reich, P. B., Reinsch, S., Wang, X., Allison, S. D., Bamminger, C., Bridgman, S., Collins, S. L., de Dato, G., Eddy, W. C., Enquist, B. J., Estiarte, M., Harte, J., Henderson, A., Johnson, B. R., Larsen, K. S., Luo, Y., Marhan, S., Melillo, J. M., Penuelas, J., Pfeifer-Meister, L., Poll, C., Rastetter, E., Reinmann, A. B., Reynolds, L. L., Schmidt, I. K., Shaver, G. R., Strong, A. L., Suseela, V., and Tietema, A.: Temperature response of soil respiration largely unaltered with experimental warming, *Proceedings of the National Academy of Science U S A*, 113, 13797-13802, 10.1073/pnas.1605365113, 2016.
- 625 Chen, J., Luo, Y., Xia, J., Wilcox, K. R., Cao, J., Zhou, X., Jiang, L., Niu, S., Estera, K. Y., Huang, R., Wu, F., Hu, T., Liang, J., Shi, Z., Guo, J., and Wang, R.-W.: Warming effects on ecosystem carbon fluxes are modulated by plant functional types, *Ecosystems*, 20, 515-526, 10.1007/s10021-016-0035-6, 2016.
- Chou, W. W., Silver, W. L., Jackson, R. D., Thompson, A. W., and Allen-Diaz, B.: The sensitivity of annual grassland carbon cycling to the quantity and timing of rainfall, *Global Change Biology*, 14, 1382-1394, 10.1111/j.1365-2486.2008.01572.x, 2008.
- 630 Davidson, E. A., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165-173, 10.1038/nature04514, 2006.
- Davidson, E. A., Trumbore, S. E., and Amundson, R.: Soil warming and organic carbon content, *Nature*, 408, 789-790, 10.1038/35048672, 2000.
- 635 Deyn, G. B. D., Cornelissen, J. H. C., and Bardgett, R. D.: Plant functional traits and soil carbon sequestration in contrasting biomes, *Ecology Letters*, 11, 516-531, 10.1111/j.1461-0248.2008.01164.x, 2008.



- Duval, T. P., and Radu, D. D.: Effect of temperature and soil organic matter quality on greenhouse-gas production from temperate poor and rich fen soils, *Ecological Engineering*, 114, 66-75, 10.1016/j.ecoleng.2017.05.011, 2018.
- 640 Eliasson, P. E., McMurtrie, Pepper, D. A., Stromgren, M., Linder, S., and Agren, G. I.: The response of heterotrophic CO<sub>2</sub> flux to soil warming, *Global Change Biology*, 11, 167-181, 2005.
- Fischer, H., Eckhardt, K.-U., Meyer, A., Neumann, G., Leinweber, P., Fischer, K., and Kuzyakov, Y.: Rhizodeposition of maize: short-term carbon budget and composition, *Journal of Plant Nutrition and Soil Science*, 173, 67-79, 10.1002/jpln.200800293, 2010.
- 645 Fontaine, S., Bardoux, G., Abbadie, L., and Mariotti, A.: Carbon input to soil may decrease soil carbon content, *Ecology Letters*, 7, 314-320, 10.1111/j.1461-0248.2004.00579.x, 2004.
- Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., and Charnov, E. L.: Effects of size and temperature on metabolic rate, *Science*, 293, 2248, 2001.
- Hartley, I. P., and Ineson, P.: Substrate quality and the temperature sensitivity of soil organic matter decomposition, *Soil Biology and Biochemistry*, 40, 1567-1574, 10.1016/j.soilbio.2008.01.007, 2008.
- 650 Heimann, M., and Reichstein, M.: Terrestrial ecosystem carbon dynamics and climate feedbacks, *Nature*, 451, 289-292, 10.1038/nature06591, 2008.
- Hinsinger, P., Bengough, A. G., Vetterlein, D., and Young, I. M.: Rhizosphere: biophysics, biogeochemistry and ecological relevance, *Plant and Soil*, 321, 117-152, 10.1007/s11104-008-9885-9, 2009.
- Hogberg, P., and Read, D. J.: Towards a more plant physiological perspective on soil ecology, *Trends in Ecology and Evolution*, 21, 548-554, 2006.
- 655 Hopkins, F. M., Filley, T. R., Gleixner, G., Lange, M., Top, S. M., and Trumbore, S. E.: Increased belowground carbon inputs and warming promote loss of soil organic carbon through complementary microbial responses, *Soil Biology and Biochemistry*, 76, 57-69, 10.1016/j.soilbio.2014.04.028, 2014.
- Jackson, R. B., Lajtha, K., Crow, S. E., Hugelius, G., Kramer, M. G., and Piñeiro, G.: The ecology of soil carbon: pools, vulnerabilities, and biotic and abiotic controls, *Annual Review of Ecology, Evolution, and Systematics*, 48, 419-445, 10.1146/annurev-ecolsys-112414-054234, 2017.
- 660 Jassey, V. E. J., Reczuga, M. K., Zielińska, M., Słowińska, S., Robroek, B. J. M., Mariotte, P., Seppely, C. V. W., Lara, E., Barabach, J., Słowiński, M., Bragazza, L., Chojnicki, B. H., Lamentowicz, M., Mitchell, E. A. D., and Buttler, A.: Tipping point in plant–fungal interactions under severe drought causes abrupt rise in peatland ecosystem respiration, *Global Change Biology*, 24, 972-986, 10.1111/gcb.13928, 2018.
- Karhu, K., Auffret, M. D., Dungait, J. A., Hopkins, D. W., Prosser, J. I., Singh, B. K., Subke, J. A., Wookey, P. A., Agren, G. I., Sebastia, M. T., Gouriveau, F., Bergkvist, G., Meir, P., Nottingham, A. T., Salinas, N., and Hartley, I. P.: Temperature sensitivity of soil respiration rates enhanced by microbial community response, *Nature*, 513, 81-84, 10.1038/nature13604, 2014.



- 670 Kirschbaum, M. U. F.: The temperature dependence of soil organic matter decomposition, and the effect of global warming on soil organic C storage, *Soil Biology and Biochemistry*, 27, 753-760, [https://doi.org/10.1016/0038-0717\(94\)00242-S](https://doi.org/10.1016/0038-0717(94)00242-S), 1995.
- Leifeld, J., and Fuhrer, J.: The temperature response of CO<sub>2</sub> production from bulk soils and soil fractions is related to soil organic matter quality, *Biogeochemistry*, 75, 433-453, [10.1007/s10533-005-2237-4](https://doi.org/10.1007/s10533-005-2237-4), 2005.
- 675 Li, G., Kim, S., Han, S., Chang, H., and Son, Y.: Effect of soil moisture on the response of soil respiration to open-field experimental warming and precipitation manipulation, *Forests*, 8, 56, [10.3390/f8030056](https://doi.org/10.3390/f8030056), 2017.
- Lin, D., Xia, J., and Wan, S.: Climate warming and biomass accumulation of terrestrial plants: a meta-analysis, *New Phytologist*, 188, 187-198, [10.1111/j.1469-8137.2010.03347.x](https://doi.org/10.1111/j.1469-8137.2010.03347.x), 2010.
- Lu, M., Zhou, X., Yang, Q., Li, H., Luo, Y., Fang, C., Chen, J., Yang, X., and Li, B.: Responses of ecosystem carbon cycle to experimental warming: a meta-analysis, *Ecology*, 94, 726-738, [10.1890/12-0279.1](https://doi.org/10.1890/12-0279.1), 2013.
- Luo, Y.: Terrestrial carbon-cycle feedback to climate warming, *Annual Review of Ecology, Evolution, and Systematics*, 38, 683-712, [10.1146/annurev.ecolsys.38.091206.095808](https://doi.org/10.1146/annurev.ecolsys.38.091206.095808), 2007.
- Luo, Y., Wan, S. Q., Hui, D. F., and Wallace, L. L.: Acclimatization of soil respiration to warming in a tall grass prairie, *Nature*, 413, 622-625, [10.1038/35098065](https://doi.org/10.1038/35098065), 2001.
- 685 Melillo, J. M., Frey, S. D., DeAngelis, K. M., Werner, W. J., Bernard, M. J., Bowles, F. P., Pold, G., Knorr, M. A., and Grandy, A. S.: Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world, *Science*, 358, 101, 2017.
- Metcalfe, D. B., Fisher, R. A., and Wardle, D. A.: Plant communities as drivers of soil respiration: pathways, mechanisms, and significance for global change, *Biogeosciences*, 8, 2047-2061, [10.5194/bg-8-2047-2011](https://doi.org/10.5194/bg-8-2047-2011), 2011.
- 690 Millar, D. J., Cooper, D. J., Dwire, K. A., Hubbard, R. M., and von Fischer, J.: Mountain peatlands range from CO<sub>2</sub> sinks at high elevations to sources at low elevations: Implications for a changing climate, *Ecosystems*, 20, 416-432, [10.1007/s10021-016-0034-7](https://doi.org/10.1007/s10021-016-0034-7), 2016.
- Moinet, G. Y. K., Hunt, J. E., Kirschbaum, M. U. F., Morcom, C. P., Midwood, A. J., and Millard, P.: The temperature sensitivity of soil organic matter decomposition is constrained by microbial access to substrates, *Soil Biology and Biochemistry*, 116, 333-339, [10.1016/j.soilbio.2017.10.031](https://doi.org/10.1016/j.soilbio.2017.10.031), 2018.
- 695 Ontl, T. A., and Schulte, L. A.: Soil carbon storage *Nature Education Knowledge*, 3, 2012.
- Pendall, E., Osana, Y. U. I., Williams, A. L., and Hovenden, M. J.: Soil carbon storage under simulated climate change is mediated by plant functional type, *Global Change Biology*, 17, 505-514, [10.1111/j.1365-2486.2010.02296.x](https://doi.org/10.1111/j.1365-2486.2010.02296.x), 2011.
- 700 Phillips, R. P., Finzi, A. C., and Bernhardt, E. S.: Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO<sub>2</sub> fumigation, *Ecology Letters*, 14, 187-194, [10.1111/j.1461-0248.2010.01570.x](https://doi.org/10.1111/j.1461-0248.2010.01570.x), 2011.



- 705 Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., Gurevitch, J., and Gcte, N.: A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming, *Oecologia*, 126, 543-562, 10.1007/s004420000544, 2001.
- Saleska, S. R., Shaw, M. R., Fischer, M. L., Dunne, J. A., Still, C. J., Holman, M. L., and Harte, J.: Plant community composition mediates both large transient decline and predicted long-term recovery of soil carbon under climate warming, *Global Biogeochemical Cycles*, 16, 2002.
- 710 Schimel, D. S., Braswell, B. H., Holland, E. A., McKeown, R., Ojima, D. S., Painter, T. H., Parton, W. J., and Townsend, A. R.: Climatic, edaphic, and biotic controls over storage and turnover of carbon in soils, *Global Biogeochemical Cycles*, 8, 279-293, 10.1029/94GB00993, 1994.
- Schlesinger, W. H.: *Biogeochemistry: an analysis of global change*, Second edition ed., Academic Press, California, 1997.
- 715 Schlesinger, W. H., and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7-20, 2000.
- Shahzad, T., Chenu, C., Genet, P., Barot, S., Perveen, N., Mougin, C., and Fontaine, S.: Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the rhizosphere priming effect induced by grassland species, *Soil Biology and Biochemistry*, 80, 146-155, 10.1016/j.soilbio.2014.09.023, 2015.
- 720 Sheik, C. S., Beasley, W. H., Elshahed, M. S., Zhou, X., Luo, Y., and Krumholz, L. R.: Effect of warming and drought on grassland microbial communities, *The ISME Journal*, 5, 1692-1700, 10.1038/ismej.2011.32, 2011.
- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture, *Journal of Advances in Modeling Earth Systems*, 7, 335-356, 10.1002/2014ms000358, 2015.
- 725 Song, B., Niu, S., Luo, R., Luo, Y., Chen, J., Yu, G., Olejnik, J., Wohlfahrt, G., Kiely, G., Noormets, A., Montagnani, L., Cescatti, A., Magliulo, V., Law, B. E., Lund, M., Varlagin, A., Raschi, A., Peichl, M., Nilsson, M. B., and Merbold, L.: Divergent apparent temperature sensitivity of terrestrial ecosystem respiration, *Journal of Plant Ecology*, 7, 419-428, 10.1093/jpe/rtu014, 2014.
- 730 Suseela, V., Conant, R. T., Wallenstein, M. D., and Dukes, J. S.: Effects of soil moisture on the temperature sensitivity of heterotrophic respiration vary seasonally in an old-field climate change experiment, *Global Change Biology*, 18, 336-348, 10.1111/j.1365-2486.2011.02516.x, 2012.
- Syed, K. H., Flanagan, L. B., Carlson, P. J., Glenn, A. J., and Van Gaalen, K. E.: Environmental control of net ecosystem CO<sub>2</sub> exchange in a treed, moderately rich fen in northern Alberta, *Agricultural and Forest Meteorology*, 140, 97-114, 10.1016/j.agrformet.2006.03.022, 2006.
- 735 Trumbore, S. E., Chadwick, O. A., and Amundson, R.: Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393, 1996.
- van der Wal, A., and de Boer, W.: Dinner in the dark: Illuminating drivers of soil organic matter decomposition, *Soil Biology and Biochemistry*, 105, 45-48, 10.1016/j.soilbio.2016.11.006, 2017.



- van Gestel, N., Shi, Z., van Groenigen, K. J., Osenberg, C. W., Andresen, L. C., Dukes, J. S., Hovenden, M. J., Luo, Y., Michelsen, A., Pendall, E., Reich, P. B., Schuur, E. A. G., and Hungate, B. A.: Predicting soil carbon  
740 loss with warming, *Nature*, 554, E4-E5, 10.1038/nature25745, 2018.
- Van Nuland, M. E., Wooliver, R. C., Pfennigwerth, A. A., Read, Q. D., Ware, I. M., Mueller, L., Fordyce, J. A., Schweitzer, J. A., Bailey, J. K., and Fox, C.: Plant-soil feedbacks: connecting ecosystem ecology and evolution, *Functional Ecology*, 30, 1032-1042, 10.1111/1365-2435.12690, 2016.
- Wan, S., Hui, D., Wallace, L., and Luo, Y.: Direct and indirect effects of experimental warming on ecosystem  
745 carbon processes in a tallgrass prairie, *Global Biogeochemical Cycles*, 19, n/a-n/a, 10.1029/2004gb002315, 2005.
- Wan, S., and Luo, Y.: Substrate regulation of soil respiration in a tallgrass prairie: results of a clipping and shading experiment, *Global Biogeochemical Cycles*, 17, 10.1029/2002GB001971, 2003.
- Wang, J., Sun, J., Xia, J., He, N., Li, M., and Niu, S.: Soil and vegetation carbon turnover times from tropical to  
750 boreal forests, *Functional Ecology*, 32, 71-82, 10.1111/1365-2435.12914, 2017.
- Wang, X., Liu, L., Piao, S., Janssens, I. A., Tang, J., Liu, W., Chi, Y., Wang, J., and Xu, S.: Soil respiration under climate warming: differential response of heterotrophic and autotrophic respiration, *Global Change Biology*, 20, 3229-3237, 10.1111/gcb.12620, 2014.
- Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., van der Putten, W. H., and Wal, D. H.: Ecological  
755 linkages between aboveground and belowground biota, *Science*, 304, 1629-1633, 10.1126/science.1094875, 2004.
- Wardle, D. A., Bonner, K. I., Barker, G. M., Yeates, G. W., Nicholson, K. S., Bardgett, R. D., Watson, R. N., and Ghani, A.: Plant removals in perennial grassland: vegetation dynamics, decomposers, soil biodiversity, and ecosystem properties, *Ecological Monographs*, 69, 535-568, 1999.
- 760 Wilson, D., Farrell Catherine, A., Fallon, D., Moser, G., Müller, C., and Renou-Wilson, F.: Multiyear greenhouse gas balances at a rewetted temperate peatland, *Global Change Biology*, 22, 4080-4095, 10.1111/gcb.13325, 2016.
- Wu, Z., Dijkstra, P., Koch, G. W., Peñuelas, J., and Hungate, B. A.: Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation, *Global Change Biology*,  
765 17, 927-942, 10.1111/j.1365-2486.2010.02302.x, 2011.
- Xue, K., Yuan, M. M., Xie, J., Li, D., Qin, Y., Hale, L. E., Wu, L., Deng, Y., He, Z., Van Nostrand, J. D., Luo, Y., Tiedje, J. M., and Zhou, J.: Annual removal of aboveground plant biomass alters soil microbial responses to warming, *MBio*, 7, 10.1128/mBio.00976-16, 2016.
- Yin, H., Li, Y., Xiao, J., Xu, Z., Cheng, X., and Liu, Q.: Enhanced root exudation stimulates soil nitrogen  
770 transformations in a subalpine coniferous forest under experimental warming, *Global Change Biology*, 19, 2158-2167, 10.1111/gcb.12161, 2013.



Zhang, N., Liu, W., Yang, H., Yu, X., Gutknecht, J. L., Zhang, Z., Wan, S., and Ma, K.: Soil microbial responses to warming and increased precipitation and their implications for ecosystem C cycling, *Oecologia*, 173, 1125-1142, 10.1007/s00442-013-2685-9, 2013.

775 Zhou, X., Sherry, R. A., An, Y., Wallace, L. L., and Luo, Y.: Main and interactive effects of warming, clipping, and doubled precipitation on soil CO<sub>2</sub> efflux in a grassland ecosystem, *Global Biogeochemical Cycles*, 20, 10.1029/2005GB002526, 2006.

Zhou, Y., Li, M.-H., Cheng, X.-B., Wang, C.-G., Fan, A. N., Shi, L.-X., Wang, X.-X., and Han, S.: Soil respiration in relation to photosynthesis of *Quercus mongolica* trees at elevated CO<sub>2</sub>, *PLOS ONE*, 5, e15134, 780 10.1371/journal.pone.0015134, 2010.

Zhu, B., and Cheng, W.: Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition, *Global Change Biology*, 17, 2172-2183, 10.1111/j.1365-2486.2010.02354.x, 2011.

785

790

795

800

805