

Dear Dr. Sara Vicca,

Thank you for accepting our revised manuscript with technical corrections.

Based on your suggestion, we thoroughly revised the manuscript and improved the formulations in many places to improve the readability.

Even if the new version has many changes, such changes are just textual improvements and the contents and interpretations are the same as the previous version.

We hope that our revisions are satisfactory.

Sincerely yours,

Greta Formaglio

1 **Herbicide weed control increases nutrient leaching compared to mechanical**
2 **weeding in a large-scale oil palm plantation**

3

4 Greta Formaglio¹, Edzo Veldkamp¹, Xiaohong Duan ², Aiyen Tjoa³, Marife D. Corre¹

5

6 ¹Soil Science of Tropical and Subtropical Ecosystems, University of Göttingen, Göttingen,
7 37073, Germany

8 ²Institute of Biochemical Plant Pathology, Helmholtz Zentrum Munich, 85764, Germany

9 ³Faculty of Agriculture, Tadulako University, Palu, 94118, Indonesia

10

11 *Correspondence to:* Greta Formaglio (gformag@gwdg.de)

12

13 **Abstract**

14 Nutrient leaching in intensively managed oil palm plantations can diminish soil fertility and
15 water quality. There is a need to reduce this environmental footprint without sacrificing yield.

16 ~~In a large-scale oil palm plantation on Acrisol soil, w~~We quantified nutrient leaching ~~in a large-~~
17 ~~scale oil palm plantation on Acrisol soil with using a full~~ factorial ~~treatment combinations~~
18 ~~of experiment with~~ two fertilization rates (260 N, 50 P, 220 K kg ha⁻¹ yr⁻¹ as conventional
19 practice, and 136 N, 17 P, 187 K kg ha⁻¹ yr⁻¹, equal to harvest export, as reduced management)
20 and two weeding methods (conventional herbicide application, and mechanical weeding as
21 reduced management). ~~Each of the four treatment combinations was represented by a 2500 m²~~
22 ~~plot, replicated in four blocks. In each plot~~Over the course of one year, we collected monthly
23 soil-pore water ~~was collected monthly~~ at 1.5 m depth ~~for one year~~ in three distinct management
24 zones: palm circle, inter-row, and frond-stacked area. Nutrient leaching ~~In in~~ the palm circle,
25 ~~nutrient leaching~~ was low due to low solute concentrations and small drainage fluxes, probably
26 resulting from large plant uptake. ~~Conversely~~In contrast, nitrate and aluminum leaching losses
27 were high in the inter-row, ~~nitrate and aluminum leaching losses were high~~ due to their high
28 concentrations and, large drainage fluxes, ~~possibly~~probably resulting from low plant uptake,
29 and acidic lower pH. In the frond-stacked area, base cation leaching was high, presumably from
30 frond litter decomposition, but N leaching was low. Mechanical weeding reduced leaching
31 losses of all nutrients compared to the conventional herbicide weeding, probably because
32 herbicides decreased ground vegetation, and ~~thereby thus~~ reduced soil nutrient retention. The
33 leaching of total N-nitrogen in our the mechanical weeding with reduced fertilization treatment
34 (32 ± 6 kg N ha⁻¹ yr⁻¹) was less than half of the the highest with conventional management (73
35 ± 20 kg N ha⁻¹ yr⁻¹) and the lowest in mechanical weeding with reduced fertilization (32 ± 6 kg
36 N ha⁻¹ yr⁻¹) whereas ~~its yields remained comparable among all~~ were not affected by our these
37 treatments. Our findings suggest that mechanical weeding and reduced fertilization should be
38 included in the program by the Indonesian Ministry of Agriculture ~~program~~ for precision

39 farming (e.g. variable rates with plantation age), particularly for large-scale oil palm
40 plantations. We furthermore suggest to include mechanical weeding and reduced fertilization,
41 ~~and in the~~ science-based policy recommendations, such as those endorsed by the Roundtable
42 for Sustainable Palm Oil association.

43 **1 Introduction**

44 Agricultural expansion is a major driver of tropical deforestation (Geist and Lambin, 2002),
45 which has global impacts on ~~reducing~~ carbon sequestration (Asner et al., 2010; van Straaten et
46 al., 2015; Veldkamp et al. *in press*), greenhouse gas regulation (e.g. Murdiyarso et al., 2010;
47 Meijide et al., 2020; Veldkamp et al. *in press*) ~~and~~ biodiversity (e.g. Clough et al., 2016) ~~and~~
48 ~~increasing profit gains at the expense of ecosystem multifunctionality (Grass et al., 2020)~~. Oil
49 palm is the ~~most important rapidly expanding~~ dominant tree-cash crop that replaces tropical
50 forest in Southeast Asia (Gibbs et al., 2010; Carlson et al., 2013) due to its high yields, ~~with~~
51 low production costs and rising global demand (Carter et al., 2007; Corley, 2009; ~~Grass et al.,~~
52 2020). Currently, Indonesia ~~produces~~ contributes 57% of the global palm oil production
53 ~~worldwide~~ (FAO, 2018), ~~which and this production~~ is projected to further expand in the future,
54 threatening ~~the~~ the remaining tropical forests (Pirker et al., 2016; Vijay et al., 2016). ~~Forest~~
55 ~~Forest-to-to-~~oil palm conversion is associated with a decrease in soil fertility because of high
56 nutrient export via harvest, reduced rates of soil-N cycling, and decreases in soil organic carbon
57 (SOC) and nutrient stocks (van Straaten et al., 2015; Allen et al., 2015; Allen et al., 2016). ~~D~~The
58 ~~declines~~ in soil fertility ~~reinforces~~ promote the dependency on fertilizer inputs and threatens the
59 long-term productivity ~~of the area~~ (Syers 1997), which could further ~~exacerbate~~ stimulate
60 expansion of oil palm production in new areas ~~land use conversion~~. Leaching ~~can~~ contributes to
61 the ~~impoverishment~~ reduction of soil nutrient ~~stocks as well as reduction in~~ and negatively
62 affects water quality, ~~and potentially leading to~~ eutrophication of water bodies. ~~Increased~~ High
63 ~~nutrient~~ loads of nutrients ~~to in~~ water bodies due to agricultural expansion and intensification,

Formatted: Font: Italic

Formatted: Font: Italic

64 common in temperate areas (Carpenter et al., 1998), are increasingly reported ~~for in humid~~
65 tropical regions (Figueiredo et al., 2010; Teklu et al., 2018). ~~Given the~~ ~~Because of the~~ ~~typically~~
66 high precipitation rates, leaching losses can be ~~large-substantial~~ in intensively managed
67 plantations in the tropics, although deeply weathered tropical soils also have the capacity to
68 ~~store-retain~~ large quantities of N and P (Neill et al., 2013; Jankowski et al., 2018). Indeed, nitrate
69 (NO_3^-), ~~the most leachable form of N~~, can be ~~retained-adsorbed in the subsoil~~ by ~~the~~ anion
70 exchange capacity ~~in the subsoil~~ of highly weathered acidic soils (Wong et al., 1990), whereas
71 P can be fixed to Fe and Al (hydr)oxides, ~~of common in heavily weathered~~ tropical soils (Roy
72 et al., 2016). Nevertheless, ~~there are some evidences of~~ ~~reductions in~~ stream water quality
73 ~~reductions due to~~ ~~have been reported in~~ oil palm cultivation in Malaysia (Luke et al., 2017;
74 Tokuchi et al., 2019). ~~This illustrates, signifying~~ the importance of quantifying nutrient leaching
75 losses in ~~other~~ areas with expansive oil palm plantations, ~~especially in such as~~ Jambi, Indonesia,
76 one of the hotspots of forest conversion to oil palm in Indonesia (Drescher et al., 2016).

77 ~~Nutrient leaching losses in oil palm plantations are calculated from water drainage~~
78 ~~fluxes and solute nutrient concentrations in the soil water (Kurniawan et al., 2018 Tarigan et al.,~~
79 ~~2020). Despite their relatively low drainage fluxes (as a consequence of high~~
80 ~~evapotranspiration; Röhl et al., 2019; a result from higher evapotranspiration and deeper rooting~~
81 ~~depth than annual crops Tarigan et al., 2020). Can have low leaching losses, as a consequence~~
82 ~~of high evapotranspiration and thus low drainage fluxes (Tarigan et al., 2020). However, most~~
83 ~~large-scale oil palm plantations are large-scale enterprises that are with their characterized by~~
84 ~~intensive management with typically have high fertilization rates, that may result in typically~~
85 ~~have high nitrate (NO_3^-) concentrations in the soil water and, resulting in large and sustained,~~
86 ~~nitrate (NO_3^-) leaching losses and herbicide application. Intensive agriculture in the tropics is~~
87 ~~associated with high N leaching losses (Huddell et al., 2020). Even in tree cash or perennial~~
88 ~~crop plantations, despite their generally higher evapotranspiration and deeper rooting depth than~~

89 ~~annual crops, high fertilization rates can result in sustained, large nutrient leaching losses~~ (e.g.
90 Wakelin et al., 2011; Cannavo et al., 2013). ~~In the leachate, Large NO₃⁻ leaching from high N~~
91 ~~fertilization is always~~ accompanied by ~~leaching of~~ cations ~~(normally bases) because of its~~
92 ~~negative charge~~ (Cusack et al., 2009; Dubos et al., 2017), ~~further~~ impoverishing highly
93 weathered tropical soils that are inherently low in base cations (Allen et al., 2016; Kurniawan
94 et al., 2018). ~~High fertilization rates are typically applied is necessary~~ to support ~~the~~ high
95 yields of oil palm plantations; ~~but however, reduction in better balanced well-adjusted~~
96 fertilization rates, e.g. to levels that compensate for nutrient export through harvest, may ~~create~~
97 ~~opportunities to~~ reduce nutrient leaching losses while maintaining high productivity.

98 ~~Herbicides are commonly used for weed control. On the other hand, the use of herbicides~~
99 ~~for weed control can exacerbate nutrient leaching losses, as prolonged absence of ground~~
100 ~~vegetation reduces uptake of redistributed nutrients from applied fertilizers (Abdalla et al.,~~
101 ~~2019). Chemical weeding with herbicides is commonly practiced in large-scale oil palm~~
102 ~~plantations. Here, the herbicides is are applied placed in the area where the fertilizers are~~
103 ~~applied, close to the palm stems to reduce competition by weeds for nutrients and water with~~
104 ~~ground vegetation, and in the inter-rows, to facilitate access during harvest (Corley and Tinker,~~
105 ~~2016). Herbicides do not only eradicate aboveground vegetative parts but also removes roots,~~
106 ~~slowing down weed regeneration. Consequently, the use of herbicides for weed control can~~
107 ~~exacerbate nutrient leaching losses, because the absence of ground vegetation reduces the~~
108 ~~uptake and thus retention of nutrients from applied fertilizers (Abdalla et al., 2019). In contrast~~
109 ~~to herbicide application, mechanical weeding does not kill eradicate the roots and only removes~~
110 ~~aboveground part, allowing for~~ relatively fast regeneration of ground vegetation, which could
111 take up redistributed nutrients and ~~could thus~~ reduce leaching losses.

112 ~~In oil palm plantations, different management zones can be distinguished, which have~~
113 ~~to be taken into account. To when investigate investigating nutrient leaching losses, in an oil~~

114 ~~palm plantation, the spatial structure created by the planting design and by the management~~
115 ~~practices must be taken into account, which is only partly considered in the sampling designs~~
116 ~~of previous studies. Typically, we can identify t~~Three contrasting management zones in oil
117 palm plantations ~~can be identified~~: (1) the palm circle, an area of 2 m radius around the palm's
118 stem where the fertilizers are applied and weeded; (2) the inter-row, which is unfertilized and
119 weed control is ed less frequently than the palm circle ~~but unfertilized~~; and (3) the frond-stacked
120 area, usually every second inter-row, where the ~~cut-pruned~~ senesced fronds are piled up and no
121 weeding or fertilization is done. In ~~these each~~ management zones, the extent of nutrient leaching
122 losses depend on the interplay of water fluxes, root uptake and soil nutrient ~~contents~~
123 ~~determine concentrations the extent of nutrient leaching losses~~. Root uptake, which is related to
124 root density, ~~which~~ is high inside the palm circle and lower in the inter-row (Lamade et al.
125 1996; Jourdan and Rey, 1997). ~~In t~~The palm circle, ~~receives the fertilizer-s are applied directly,~~
126 but also ~~takes up high amounts~~ uptake of water and nutrients is highest (Nelson et al., 2006).
127 Hence, large leaching losses may only occur shortly following ~~pulse high~~ fertilization if and
128 during high drainage fluxes occur, e.g. directly following intensive rain showers (from high
129 precipitation) events (Banabas et al., 2008a). The inter-row ~~experiences has~~ higher water input
130 from precipitation than the palm circle because of the lower interception by the canopy
131 ~~intereception~~ (Banabas et al., 2008b). Here, root density and thus root uptake is low, resulting in
132 a_ and_ large water fluxes_ within the soil because of low root uptake, stimulating nutrient
133 transport to lower depths. However, nutrient leaching may be low in the inter-row because there
134 is no direct fertilizer application. The frond-stacked area receives nutrients from decomposition
135 of nutrient-rich fronds (Kotowska et al., 2016). ~~Furthermore, and such~~ mulching with senesced
136 fronds prevents runoff and promotes water infiltration as a consequence of enhanced owing to
137 the high macroporosity, a result of high_ by increased_ organic matter and biological activity
138 (Moradi et al., 2015). Low canopy interception and h High water infiltration may generate

139 high water drainage fluxes, resulting in intermediate nutrient leaching losses in ~~the front-~~
140 ~~stacked~~ this area management zone.

141 In this study, we aimed to quantify nutrient leaching losses in our experiment that was
142 established in an intensively managed, large-scale oil palm plantation; in order ~~and~~ to assess ~~if~~
143 ~~whether reduced-lower intensity of~~ management intensity (i.e. reduced fertilization rates equal
144 to harvest export and mechanical weeding) can reduce leaching losses without affecting yield
145 ~~in oil palm plantations~~. We tested the following hypotheses: (1) leaching losses in the palm
146 circle are larger than in ~~the~~ other management zones because of direct fertilizer application; (2)
147 leaching losses under herbicide application are higher than mechanical weeding because of
148 ~~lower regeneration of ground vegetation that can augment~~ the reduced nutrient retention owing
149 to reduced weed growth; (3) nutrient leaching fluxes under reduced ~~under conventional high~~
150 fertilization rates are substantial-lower compared to ~~reduced-conventional, high rates-~~ but yield
151 ~~not affected~~ ~~because of excessive nutrient inputs~~. Our study provides the first a-systematic
152 quantification of leaching losses, an important environmental footprint of oil palm production,
153 taking into consideration ~~its spatial variation in~~ the different management zones, and evaluates
154 the effectiveness of alternative management practices ~~for~~ to reduce on leaching and yield
155 reduction.

157 2 Materials and methods

158 2.1 Study area and experimental design

159 ~~This~~ Our study was conducted in a state-owned oil palm plantation in Jambi province, Indonesia
160 (1° 43' 8" S, 103° 23' 53" E, 73 m above sea level). Mean annual air temperature is 26.7 ± 1.0
161 °C and mean annual precipitation is 2235 ± 385 mm (1991–2011; data from Sultan Thaha
162 airport, Jambi). During our study period (March 2017–February 2018), the mean daily air

163 temperature was 26.3 °C and annual precipitation was 2772 mm, with a dry period between
164 July and October (precipitation < 140 mm month⁻¹). The soil is highly weathered, loam Acrisol
165 soil (Allen et al., 2015) and nutrient inputs from bulk precipitation in the area, measured in
166 2013, were 12.9 kg N, 0.4 kg P, 5.5 kg K ha⁻¹ yr⁻¹ (Kurniawan et al., 2018).

167 This oil palm plantation was established between 1998 and 2002, and ~~so~~the palms were
168 16–20 years old during our study period. The plantation ~~has~~ is mostly located on flat terrain,
169 and it encompassed 2025 ha, with a planting density of approximately 142 palms ha⁻¹, spaced
170 8 m apart ~~on rows~~. The rows between palms are used alternately for harvesting operations and
171 to pile-up senesced fronds, which are regularly ~~cut~~ pruned to facilitate harvesting of fruits; this
172 frond-stacked area covers approximately 15% of the plantation area. The palm circle, a -2 m
173 radius from the stem, where ~~in both~~ fertilizers ~~are applied~~ and ~~chemically weeded~~ herbicides are
174 applied four times a year, covers 18% of the plantation. The remaining 67% ~~can be~~ classified
175 classified as inter-row, which is not fertilized but weeded ~~two times~~ twice a year.

176 In November 2016, a ~~two (fertilization rates) by two (weeding methods)~~ factorial
177 management experiment was established with two fertilization rates and two weeding methods
178 (in this plantation as part of the framework of the EFForTS project, described in detail by Darras
179 et al., (2019). For ~~fertilization~~ fertilizer treatments, the conventional rates were 260 N, 50 P,
180 220 K kg ha⁻¹ yr⁻¹, whereas the reduced rates were 136 N, 17 P, 187 K kg ha⁻¹ yr⁻¹. Reduced
181 fertilization rates were ~~determined~~ established to compensate for nutrient exports via fruit
182 harvest and were ~~based~~ assessed on by multiplying the nutrient concentrations measured in the
183 fruit bunches ~~multiplied by~~ with the annual yield. The fertilizer sources were urea (CH₄N₂O),
184 triple superphosphate (Ca(H₂PO₄)₂·H₂O) and muriate of potash (KCl), in granular forms. ~~These~~
185 Fertilizers were applied ~~according to~~ following the plantation's standard practices: split in two
186 applications per year (in April and October), spread in a band ~~within~~ at approximately 2 m
187 radius from the palm, ~~and this area~~ which that was raked before fertilizer application. For both

188 ~~fertilization fertilizer~~ treatments, we also applied lime (426 kg dolomite ha⁻¹ yr⁻¹; CaMg(CO₃)₂)
189 and micronutrients (142 kg Micro-Mag ha⁻¹ yr⁻¹ with 0.5% B₂O₃, 0.5% CuO, 0.25% Fe₂O₃,
190 0.15% ZnO, 0.1% MnO and 18% MgO). ~~were also applied besides the N, P and K fertilizers,~~
191 as commonly practiced in large-scale plantations on acidic Acrisol soils (Pahan, 2010). ~~For~~
192 ~~weeding treatments, the~~ conventional ~~method weed control~~ was ~~the use of~~ done using a
193 herbicides (glyphosate), whereas the ~~reduced alternative~~ method was mechanical weeding using
194 a brush cutter; ~~the~~ cut plant materials were left on the ground. Herbicide Glyphosate was
195 applied following plantation's standard practice: 1.5 L glyphosate ha⁻¹ yr⁻¹ to the palm circle,
196 split four times a year, and 0.75 L glyphosate ha⁻¹ yr⁻¹ to the inter-row, split two times a year.
197 ~~The mechanical~~ weeding was carried out in the same areas and frequencies as herbicide
198 application. This management experiment comprised of four replicate blocks, ~~and each had~~
199 with four plots (50 m x 50 m each) assigned to four treatment combinations: conventional rate–
200 herbicide, conventional rate–mechanical weeding, reduced rate–herbicide, and reduced rate–
201 mechanical weeding.

202

203 2.2 Soil water sampling

204 Over the course of one year, we collected monthly soil-pore water samples ~~over one year,~~
205 using suction cup lysimeters (P80 ceramic, maximum pore size 1 µm; CeramTec AG,
206 Marktredwitz, Germany). We installed the lysimeters in January 2017, randomly choosing two
207 palms per plot and sampling in the three management zones: 1) within in the palm circle, at 1
208 m from the palm stem, 2) in the frond-stacked area, at about 4 m from the palm stem, and 3) in
209 the inter-row, at approximately 4 m from the palm stem (Fig. A1). In total, we installed, 96
210 lysimeters ~~were installed~~ (4 treatments x 4 replicates x 2 subplots x 3 management zones). The
211 lysimeters were inserted into the soil to 1.5 m depth, so that the soil-pore water was collected
212 well below the rooting depth of 1 m, which is common ~~to for~~ oil palm plantations on loam

213 Acrisol soils near our study site (Kurniawan et al., 2018). Starting in March 2017, we sampled
214 soil water ~~was sampled~~ by applying 40 kPa vacuum (Kurniawan et al., 2018; Dechert et al.,
215 2005) to the lysimeters. Water samples were ~~and~~ collected in dark glass bottles, which were
216 stored in a bucket buried in the field. We consider ~~Although there was only~~ the two-month
217 acclimatization of lysimeters ~~between their installation and the beginning of~~ before sampling ~~;~~
218 ~~we considered this to be~~ sufficient, because ~~as~~ soil disturbance was minimized and biochemical
219 processes are rapid in tropical soils. During sampling, ~~Once a week~~, we transferred once a week
220 the collected water into plastic bottles ~~and which were~~ transported ~~them~~ to the field station,
221 where they were ~~stored~~ frozen for storage. ~~The Soil water~~ collection continued ~~over~~ during a
222 month until a volume of 100 mL was collected from each lysimeter, or until the end of the
223 month. The frozen water samples were transported by air ~~freight~~ to the University of
224 Goettingen, Germany, where element concentrations were determined. We measured the
225 concentrations of mineral N (NH_4^+ and NO_3^-), total dissolved N (TDN) and Cl⁻ by using
226 continuous flow injection colorimetry (SEAL Analytical AA3, SEAL Analytical GmbH,
227 Norderstadt, Germany), as described in details ~~s~~ by Kurniawan et al. (2018). We calculated
228 ~~d~~ Dissolved organic N (DON) ~~was calculated~~ as the difference between TDN and mineral N.
229 We measured the concentrations of base cations (Na, K, Ca, Mg), total Al, total Fe, total Mn,
230 total S, and total P with using an inductively coupled plasma–atomic emission spectrometer
231 (iCAP 6300; Thermo Fischer Scientific GmbH, Dreieich, Germany).

232 We determined a partial cation-anion charge balance of the major elements
233 (concentrations $> 0.03 \text{ mg L}^{-1}$) in soil-pore water by converting the concentrations to $\mu\text{mol}_{\text{charge}}$
234 L^{-1} . For this, we ~~We~~ assumed S to be in the form of sulfate (SO_4^{2-}) and total Al to have a charge
235 of 3⁺. We calculated the combined contribution of organic acids (RCOO^-) and bicarbonate
236 (HCO_3^-) as the difference between the measured cations and anions (Kurniawan et al., 2018).

237

238 2.3 Modeling water drainage

239 The water balance was modeled using the water sub-model of the Expert-N software, version
240 5.0 (Priesack, 2005), which was successfully used in previous research to estimate drainage
241 fluxes from different land uses in Indonesia (Dechert et al., 2005; Kurniawan et al., 2018). The
242 model inputs were climate data (solar radiation, temperature, precipitation, relative humidity,
243 and wind speed), and soil (texture, bulk density, and hydraulic functions) and vegetation
244 characteristics (biomass, leaf area index, and root distribution). The climate data were ~~taken~~
245 collected from the climatological station in the plantation (described in detail by Meijide et al.,
246 2017), ~~whereas and the for the~~ oil palm biomass ~~was used taken used published data~~ from ~~a~~
247 ~~study on~~ oil palm plantations near our study site (Kotowska et al., 2015). We measured sSoil
248 bulk density and porosity in the top 10 cm ~~were measured in of~~ each management zone at our
249 study site, whereas for the 10–50–cm depth these were measured in the inter-row. ~~Data for~~
250 soil bulk density and porosity for the 50–200–cm depth, as well as soil texture, soil hydraulic
251 parameters (i.e. water retention curve, saturated hydraulic conductivity and Van Genuchten
252 parameters ~~for the water retention curve~~), and root distribution ~~were taken~~ we used published
253 data from Allen et al. (2015) and Kurniawan et al. (2018), choosing their studied oil palm
254 plantations closest to our study site. The Expert-N water sub-model calculates daily water
255 drainage based on precipitation, evapotranspiration, canopy interception, runoff, and change in
256 soil water storage. Evapotranspiration is calculated using the Penman-Monteith method (Allen,
257 1998), applying a plant factor of 1.06 (Meijide et al., 2017), with plant transpiration based on
258 leaf area index (LAI), plant biomass, and maximum rooting depth. The canopy interception is
259 calculated from the percentage of throughfall and the maximum water storage capacity of the
260 canopy. Runoff is calculated from soil texture and bulk density, which determine the water
261 infiltration rate, and from the slope, which was 5% (Röll et al., 2019). The vertical water

262 movement is calculated using Richards equation based on soil hydraulic functions (Hillel,
263 1982).

264 To model the drainage in the different management zones, we used the measured soil
265 bulk density and porosity in the top 10 cm and adjusted other input parameters to simulate
266 differences in water balance in each management zone (Table A1). For the palm circle, we set
267 the LAI to 3.65, which is the maximum LAI measured at our site (Fan et al., 2015), to simulate
268 high water uptake in the palm circle (Nelson et al., 2006) and maximum rooting depth to 1 m,
269 which is reported for oil palm plantations near our site (Kurniawan et al., 2018). The percentage
270 throughfall in the palm circle was set ~~to~~at 10% and the water storage capacity of oil palm stem
271 was set to 8.4 mm (Tarigan et al., 2018). For the inter-row, we set the LAI and the maximum
272 rooting depth ~~as~~at half ~~the values~~ of the palm circle (1.8 LAI, 50 cm rooting depth), as roots
273 are shallower between palms (Nelson et al., 2006); ~~the~~ throughfall was set ~~to~~at 50%, and the
274 palm stem's water storage capacity was set ~~to~~at 4.7 mm (based on canopy storage capacity
275 reported by Tarigan et al., 2018). For the frond-stacked area, the LAI was set to 0.75, which is
276 half of the minimum measured in the studied plantation (Darras et al., 2019), ~~as~~because
277 understory vegetation is absent at this zone. Values for interception in the frond-stacked area
278 was set to the same values as the inter-row, whereas the runoff was set to 0 (no overland runoff),
279 ~~as~~because mulching with senesced fronds ~~increases water infiltration and slows down~~prevents
280 runoff (Tarigan et al., 2016).

281 For validation of the Expert-N water sub-model outputs, we measured weekly soil water
282 matric potential at ~~depths of~~ 30 cm and 60 cm depths over the study period and compared the
283 measured values with the modeled matric potential. Matric potential was measured by installing
284 a tensiometer (with a P80 ceramic, maximum pore size 1 μm ; CeramTec AG, Marktredwitz,
285 Germany) at each depth in each management zone near ~~to~~ two palms in two treatments (i.e.
286 conventional rate-herbicide, and reduced rate-mechanical weeding), for a total of 12

287 tensiometers. We summed the modeled daily drainage at 1.5 m depth to get the monthly
288 drainage fluxes, which we then multiplied with the element concentrations in soil water to get
289 the monthly nutrient leaching fluxes.

290

291 **2.4 Soil biochemical characteristics and nutrient retention efficiency**

292 We measured soil biochemical properties in the same sampling locations (Figure A1) at the
293 following four depth intervals: 0–5 cm, 5–10 cm, 10–30 cm, and 30–50 cm. In each plot, sSoil
294 samples from the same management zone ~~in each plot~~ were pooled to make one composite
295 sample, totaling 192 soil samples (4 treatments x 4 replicates x 3 management zones x 4 depths).

296 The samples were air-dried and sieved (2 mm). We and measured ~~for~~ pH (on a 1:4 soil-to-water
297 ratio) and ~~for~~ effective cation exchange capacity (ECEC), by percolating the soils with
298 unbuffered 1 mol L⁻¹ NH₄Cl and measuring-analyzing the cations (Ca, Mg, K, Na, Al, Fe, Mn)
299 in percolates using ICP-AES. A subsample was finely ground and analyzed for organic C and
300 total N using a CN analyzer (Vario EL Cube, Elementar Analysis Systems GmbH, Hanau,
301 Germany), and for ¹⁵N natural abundance signature using isotope ratio mass spectrometer
302 (IRMS; Delta Plus, Finnigan MAT, Bremen, Germany). We calculated the soil element stocks
303 for each depth by multiplying the element concentration with the measured bulk density and
304 summed-adding them for the top 50 cm; other soil characteristics (e.g. pH, ECEC, base
305 saturation) in the top 50 cm soil were calculated as the depth-weighted average of the sampled
306 depths.

307 In addition, we calculated the N and base cation retention efficiency in the soil for each
308 experimental treatment and management zone following the formula: nutrient retention
309 efficiency = 1 – (nutrient leaching loss / soil-available nutrient) (Kurniawan et al., 2018). We
310 used the gross N mineralization rates in the top 5 cm soildepth (Table A2) as an index of soil-

311 available N whereas soil-available base cations was the sum of the stocks of K, Na, Mg and Ca
312 in the top 10 cm ~~depth-soil~~, expressed in mol_{charge} m⁻².

313

314 2.5 Statistical analyses

315 For soil biochemical properties measured once, we tested for differences among management
316 zones and experimental treatments for the entire 50 cm depth, using the analysis of variances
317 (ANOVA) with Tukey HSD as a post hoc test. The soil variables that showed non-normal
318 distribution or unequal variances ~~were log-transformed prior to the analysis~~, ~~tested~~ with
319 Shapiro–Wilk and Levene’s tests, respectively, ~~were log-transformed prior to the analysis~~. Base
320 cation and N retention efficiency were also tested for differences between experimental
321 treatments in the same way. For repeatedly measured variables, i.e. soil-pore water solute
322 concentrations and leaching fluxes, we used linear mixed-effects models (LME; Bates et al.,
323 2015) to assess the differences among management zones and treatments. For testing
324 ~~management zone~~ differences among management zones, we conducted the LME with
325 management zone as fixed effect and random effects for sampling months and experimental
326 treatments nested with replicate plots, which were also nested with subplots. For testing
327 treatment differences, we calculated for each replicate plot on each sampling month the area-
328 weighted average of the three management zones (i.e. palm circle accounts for 18% of the
329 plantation area, the frond-stacked area 15%, and the inter-row 67%), and LME was carried out
330 with treatment as fixed effect and random effects for sampling months and replicate plots nested
331 with subplots. If the residuals of the LME models were not normally distributed, we applied
332 either logarithmic or square root transformation. Differences were assessed with ANOVA
333 (Kuznetsova et al., 2017) followed by Tukey HSD (Hothorn et al., 2008). We also used LME
334 to assess differences in soil water matric potential among management zones, with management
335 zone as fixed effect and measurement days and depth nested with treatment as random effects.

336 Comparability between modeled and measured soil water matric potential for each depth in
337 each management zone ($n = 50$ field measurements) was assessed using Pearson correlation
338 test. All tests were considered significant at $P \leq 0.05$, except for soil pH, ~~for~~ which ~~we~~
339 ~~considered~~ ~~accepted~~ ~~there was~~ a marginal significance at $P = 0.06$. All statistical analyses were
340 performed with R version 3.6.1 (R Core Team, 2019).

341

342 **3 Results**

343 **3.1 Soil biochemical properties and water balance**

344 Soil biochemical properties in the top 50 cm did not differ between experimental treatments (all
345 $P > 0.05$) but strongly differed among management zones (Table 1). The frond-stacked area,
346 where senesced fronds were regularly piled like mulch material, had higher SOC and total N
347 stocks ($P < 0.01$) compared to the other management zones. The inter-row, with regular
348 weeding but without direct fertilizer and lime inputs, showed lower exchangeable base cation
349 contents (i.e. Ca, Mg, K) compared to the other management zones ($P \leq 0.02$) and higher
350 exchangeable Al content than the palm circle ($P = 0.01$). This was reflected in the lower base
351 saturation and higher Al saturation in the inter-row compared to the other zones ($P < 0.01$).
352 Also, inter-row had the lowest ECEC ($P < 0.01$) and marginally lower pH than the palm circle
353 ($P = 0.06$). The palm circle, where fertilizers and lime were applied, had generally comparable
354 exchangeable element contents with the frond-stacked area, except for K, which was higher in
355 the palm circle ($P < 0.01$), and for Mn, which was higher in the frond-stacked area ($P < 0.01$).

356 There were strong positive correlations between field-measured and modeled soil water
357 matric potential (Fig. 1). The matric potential was generally lowest in the palm circle,
358 intermediate in the inter-row, and highest in the frond-stacked area ($P < 0.01$). This pattern was
359 also reflected in the low drainage flux in the palm circle and high drainage flux in the frond-

360 stacked area (Table 2; Fig. 2). In the palm circle, the low drainage flux had resulted from high
361 plant transpiration and interception whereas the high drainage flux in the frond-stacked area
362 was due to low evapotranspiration and runoff with the senesced frond mulch (Table 2).
363 Compared to annual precipitation, the calculated annual evapotranspiration was 51%, 31%, and
364 38% in the palm circle, frond-stacked area, and inter-row, respectively; annual drainage fluxes
365 at 1.5-m depth were 20% of precipitation in the palm circle, 65% in the frond-stacked area, and
366 43% in the inter-row. Over the course of one year, the monthly drainage fluxes displayed two
367 peaks, in May and November, which occurred following several ~~Seasonally, the monthly~~
368 ~~drainage fluxes had two peak periods, May and November, after consecutive~~ days of moderate
369 rainfall. Lowest drainage fluxes were measured, and were lowest during the end of the dry
370 season ~~towards the start of the wet season~~ (Fig. 2).

371

372 3.2 Differences in leaching losses among management zones and treatments

373 We detected clear treatment differences ~~F~~or element concentrations in soil-pore water at 1.5 m
374 depth, ~~treatment differences were clear in~~ between the palm circle and inter-row (Fig. 3), with
375 the herbicide treatment showing higher element concentrations than the mechanical weeding
376 ($P \leq 0.02$). The frond-stacked area had generally lower ionic charge concentrations compared
377 to the other management zones (Fig. 3). ~~The~~ dominant cations in leachate were Al^{3+} , Ca^{2+} ,
378 Mg^{2+} , K^+ , and Na^+ across experimental treatments and management zones. Dissolved ~~Among~~
379 ~~the management zones,~~ Al^{3+} concentrations were highest in the inter-row, intermediate in the
380 palm circle, and lowest in the frond-stacked area ($P < 0.01$). The Ca^{2+} concentrations ~~of Ca^{2+}~~
381 were similar in the palm circle and frond-stacked area ($P = 0.42$), and ~~these both~~ were higher
382 than in the inter-row ($P < 0.01$). The concentrations of Mg^{2+} and K^+ were higher in the palm
383 circle than in the other two management zones ($P < 0.01$). The Na^+ concentrations were higher
384 in the palm circle and inter-row than in the frond-stacked area ($P < 0.01$). As for N, NH_4^+

385 concentrations were lowest in the frond-stacked area, followed by the palm circle, and highest
386 in the inter-row ($P = 0.01$). Across treatments, NH_4^+ was 4-18% of TDN whereas DON was
387 ~~only~~ 1-7% of TDN. Thus, NO_3^- was ~~thus~~ the main form of dissolved N, ~~and-which was this was~~
388 highest in the inter-row, followed by the frond-stacked area, and lowest in the palm circle ($P <$
389 0.01). The dominant anion was Cl^- with higher concentrations in the palm circle than in the
390 other zones ($P < 0.01$).

391 Monthly leaching fluxes showed a common pattern among the major solutes (Fig. 4):
392 ~~there were~~ two peaks of leaching losses (May and November) ~~that~~ followed fertilizer
393 applications, ~~and-whereas~~ lower leaching losses occurred during the dry season from July to
394 October. Leaching fluxes of NO_3^- ~~showed~~ followed a similar spatial pattern as ~~NO_3^- 's~~
395 concentrations: higher in the inter-row, followed by the frond-stacked area, and lowest in the
396 palm circle ($P < 0.01$; Fig. 4). Total Al leaching fluxes were also higher in the inter-row than
397 the other zones ($P < 0.01$; Fig. 4). ~~On the other hand~~In contrast, base cation leaching fluxes
398 ~~showed~~ displayed opposite spatial patterns ~~than their~~ compared to their concentrations: Ca, K,
399 and Mg leaching were higher in the frond-stacked area than the palm circle and inter-row (all
400 $P < 0.01$; Fig. 4). Leaching of Na was higher in both the frond-stacked area and inter-row than
401 the palm circle ($P < 0.01$; Fig. 4).

402 Reduced intensity of management ~~clearly~~ strongly influenced nutrient leaching losses
403 (Fig. 5; Table 3). ~~Specifically,~~ mechanical weeding reduced NO_3^- and cation leaching
404 compared to herbicide weed control ($P \leq 0.03$; Fig. 5; Table 3). Leaching of NO_3^- was highest
405 in the conventional fertilization-herbicide treatment and lowest in reduced management
406 treatments ($P \leq 0.02$; Fig. 5). This was also reflected in the leaching fluxes of accompanying
407 cations; specifically, total Al and Ca leaching were higher in conventional fertilization-
408 herbicide treatment than the reduced management treatments (all $P \leq 0.02$; Fig. 5). For the other
409 base cations, mechanical weeding ~~clearly~~ lowered leaching losses compared to herbicide

410 weeding, in particular K and Na leaching in both fertilization rates and Mg leaching in
411 conventional fertilization (all $P \leq 0.03$; Fig. 5).

412

413 3.3 Annual leaching losses and nutrient retention efficiency

414 In proportion to the applied fertilizer, annual leaching losses of TDN (Table 3) were 28% of the
415 applied N in the herbicide treatment for both conventional and reduced fertilization rates, 24%
416 in the mechanical weeding with conventional fertilization, and only 19% in the mechanical
417 weeding with reduced fertilization. The annual leaching of K (Table 3) was 4% of the applied
418 K fertilizer in the herbicide treatment and 3% in the mechanical weeding for both fertilization
419 rates. In this highly weathered Acrisol soils with high capacity for P fixation by Fe and Al
420 (hydr)oxides, ~~there was~~ we detected no leaching of dissolved P (Table 3).

421 Both N and base cation retention efficiencies were generally lower in the inter-row
422 compared to the other management zones ($P \leq 0.03$), except for reduced fertilization–
423 mechanical weeding where there were no differences among management zones (Table 4). The
424 area-weighted average N retention efficiency was comparable among experimental treatments
425 ($P = 0.89$) but there was a trend of increasing efficiency with decreasing management intensity
426 (Table 4). Base cation retention efficiency showed ~~clear~~ strong differences among experimental
427 treatments for each management zones: in the palm circle, it was highest in mechanical weeding
428 and lowest in the herbicide treatment ($P = 0.04$); in the frond-staked area and inter-row, it was
429 lowest in the most intensive management treatment (conventional fertilization–herbicide) and
430 highest in either mechanical weeding or reduced fertilization ($P \leq 0.05$; Table 4). The area-
431 weighted average d base cation retention efficiency was also ~~clearly~~ influenced by weeding
432 method, being lowest in herbicide treatment and highest in mechanical weeding both with
433 conventional fertilization ($P = 0.03$; Table 4).

434

435 4 Discussion

436 4.1 Water model and temporal pattern of nutrient leaching losses

437 To our knowledge, ~~this our~~ study is the first ~~attempt that to model~~ has modeled water drainage
438 fluxes from the different management zones of an oil palm plantation, which makes comparison
439 with other published values challenging~~limiting our comparison with literature values. Our~~
440 ~~modeled~~ annual transpiration rates in the palm circle (Table 2) ~~was were~~ remarkably similar to
441 the values estimated with the eddy covariance technique in the same oil palm plantation (827–
442 829 mm yr⁻¹; Meijide et al., 2017; Röhl et al., 2019). ~~Furthermore, and,~~ our average daily
443 transpiration rate (2.3 mm d⁻¹) was within the range of ~~that rates~~ measured with drone-based
444 photogrammetry (3 ± 1 mm d⁻¹; Ahongshangbam et al., 2019), also in the same plantation~~in~~
445 ~~the same oil palm plantation. Also, t~~The modeled annual runoff in the palm circle and inter-row
446 (Table 2) ~~was were also~~ within the range of runoff estimates in oil palm plantations in Jambi
447 province (10–20% of rainfall; Tarigan et al., 2016) and in Papua New Guinea (1.4–6% of
448 rainfall; Banabas et al., 2008b). Considering the areal proportions of the three management
449 zones, the weighted-average drainage flux (1161 mm yr⁻¹) was lower than ~~that the~~ estimated for
450 smallholder oil palm plantations near our study site (1614 mm drainage flux with 3418 mm
451 precipitation measured in 2013; Kurniawan et al., 2018). However, higher evapotranspiration
452 rates in large-scale compared to smallholder oil palm plantations in our study area (Röhl et al.,
453 2019) ~~may explain these differences. Nevertheless,~~ ~~However,~~ ratios of drainage flux to annual
454 precipitation were comparable between our study and ~~that by the study by~~ Kurniawan et al. (
455 ~~2018)~~. We conclude from ~~Also, evapotranspiration rate is higher in large-scale than~~
456 ~~smallholder oil palm plantations in our study area (Röhl et al., 2019), which could have led to a~~
457 ~~lower drainage flux in large-scale plantations. Based on~~ these comparisons with literature

458 values and on the good agreement between modeled and measured soil water matric potential
459 (Fig. 1), ~~we conclude~~ that our modeled water drainage fluxes were reliable. ~~Moreover,~~ the
460 frond-stacked areas had larger drainage fluxes, ~~caused by as a result of a combined combination~~
461 ~~of with~~ low evapotranspiration and runoff (Table 2) and enhanced porosity (~~indicated by lower~~
462 ~~bulk density;~~ Table 1) from organic matter that facilitates water infiltration (Moradi et al.,
463 2015), ~~as indirectly indicated by its low soil bulk density (Table 1)~~. This suggests that piling
464 senesced fronds may amend groundwater recharge, which, ~~in turn,~~ could moderate discharge
465 fluctuations in water catchments of oil palm-converted areas (Tarigan et al., 2020).

466 The temporal peaks of nutrient leaching fluxes (May and November; Fig. 4) likely
467 resulted from the combined effect of high drainage flux and fertilizer application. ~~High-Large~~
468 drainage fluxes might have stimulated the downward transport of nutrients and decreased their
469 residence time in the soil, and thus their adsorption onto the soil exchange sites (Lohse and
470 Matson, 2005). ~~Although~~ large drainage fluxes usually dilute the nutrient concentrations in
471 the soil-pore water; ~~;~~ ~~however,~~ the combined fertilizer and lime applications were able to
472 maintained high nutrient concentrations as manifested by the parallel peaks of drainage and
473 nutrient leaching fluxes (Figs. 2 and 4). The high NO₃⁻ leaching following urea-N fertilization
474 (Fig. 4) suggests ~~increased-rapid~~ nitrification (Silver et al., 2005), fast NO₃⁻ transport through
475 the soil column, and ~~reduced-limited~~ anion adsorption capacity (Wong et al., 1990). The latter
476 was possibly ~~aggravated-affected~~ by the ~~additional-added~~ Cl⁻ from fertilization with KCl (Fig.
477 3), which ~~could-may have~~ saturated the soil anion exchange sites, particularly ~~at in~~ this mature
478 plantation, ~~with already which has been intensively fertilized for~~ 16–20 years ~~of high~~
479 ~~fertilization rates~~. ~~Due to its negative charge,~~ Large-NO₃⁻-leaching fluxes ~~is-are~~ always
480 accompanied by comparable large-leaching fluxes of ~~buffering-positive~~ cations (Dubos et al.,
481 2017; Kurniawan et al., 2018), resulting in ~~their~~ similar temporal leaching patterns (Fig. 4).
482 ~~These-Our~~ findings ~~showed-illustrate~~ that fertilization should be avoided during periods of high

483 drainage fluxes, ~~which were~~. Generally, the high drainage was a consequence of ~~related to~~
484 ~~protracted~~ extended period of moderate rainfall (Fig. 2). However, it is expected that reliable
485 ~~p~~Prediction of periods ~~of with~~ high ~~precipitation rainfall~~ and drainage ~~will become even more~~
486 ~~difficult with~~ ~~will further be confounded by~~ climate change, which is ~~widening the range~~
487 ~~between wet and dry seasons and~~ increasing ~~the~~ uncertainties in rainfall intensity and
488 distribution (Chou et al., 2013; Feng et al., 2013). Fertilization during the dry ~~period season~~ is
489 also not advisable ~~given the high volatilization of applied~~ because plant uptake is low during
490 ~~this period (Corley and Tinker, 2016) and application of urea together with lime will cause urea~~
491 ~~to volatilize easily~~, even in ~~these~~ acidic soils ~~as this is always accompanied by liming~~ (Goh et
492 al., 2003; Pardon et al., 2016) ~~and the low palm uptake during the dry season (Corley and~~
493 ~~Tinker, 2016)~~.

494 Our results suggest that there are several viable options to reduce leaching losses without
495 sacrificing production. Thus, ~~sp~~Spreading ~~the fertilization or applications~~ over a longer period
496 of time and reducing fertilization rates, e.g. at compensatory level equal to harvest export, as
497 we tested in our experiment, are recommendable alternatives to present practices, ~~seem viable~~
498 options to reduce leaching losses without sacrificing production. In addition, ~~t~~One other option
499 is the use of organic amendments, such as empty fruit bunches, compost, palm oil mill effluent,
500 or slow-release fertilizers, which have been shown to reduce N leaching in tropical cropping
501 systems (Nyamangara et al., 2003; Steiner et al., 2008; Mohanty et al., 2018), will also reduce
502 leaching losses. ~~On addition,~~ organic fertilizer have the additional advantage that of ~~can~~
503 improve ~~improving~~ soil fertility ~~can be improved~~ in oil palm plantations (Comte et al., 2013;
504 Boafo et al., 2020), as was also ~~evident with~~shown by mulching of senesced oil palm fronds
505 (i.e. high SOC, total N, ECEC and base saturation in the frond-stacked area; Table 1).

506

507 **4.2 Leaching losses in the different management zones**

508 A surprising result, in contrast ~~Contrary~~ to our first hypothesis, was that nutrient leaching losses
509 among management zones were generally large in the inter-row, especially for mineral N
510 (largely NO₃; Fig. 3), and lower in the palm circle (Fig. 4). ~~This strikingly large mineral N~~
511 ~~leaching losses in the inter row were surprising~~ We did not expect this given that ~~because the~~
512 ~~inter-row this area~~ did not receive direct fertilizer inputs (see section 2.1). ~~This~~ Our results
513 suggests that mineral N was transported via surface and/or subsurface lateral flows from the
514 ~~directly~~ fertilized palm circle to the inter-row via surface and subsurface lateral flow as these
515 ~~two zones which~~ were just only 3 m apart (Fig. A1). We expect that the contribution of sSurface
516 transport of mineral N was ~~probably a~~ minor process at our site, because of the low runoff
517 (Table 2) ~~;-). Also~~ in an oil palm plantation in Papua New Guinea, the loss of N fertilizer via
518 surface runoff was only 0.3–2.2 kg N ha⁻¹ yr⁻¹ (Banabas et al., 2008b). The dominant form of
519 transport of mineral N in our experiment ~~Mineral N~~ was probably likely predominantly
520 transported to the inter row via subsurface lateral flow. Acrisol soils are characterized by
521 clay translocation to a subsurface soil horizon from upper to lower depths that ~~could can~~ create
522 ~~an impeding stagnating~~ layer above which ~~conducive to~~ lateral water flow can occur (Elsenbeer,
523 2001). Indeed, the clay contents of the Acrisol soils at our study area increase with depth, and
524 soil bulk density at 100–150 cm is was larger ~~at 100–150 cm~~ than at 150–200 cm depth (Allen
525 et al., 2016). In addition, ~~the~~ palm roots spreading from the palm circle to the inter-row may
526 create channels for subsurface lateral flow of dissolved ions such as like NO₃⁻ (Li and Ghodrati,
527 1994). Higher mineral N leaching in the inter-row than palm circle ~~was had~~ also been observed
528 in a study in Brazil ~~and where~~ it was attributed to lower root density and higher N mineralization
529 at increasing distance from the palm's stem (Schroth et al., 2000). Hence, a combination of
530 lower root uptake, higher N mineralization, and subsurface lateral transport (particularly for
531 NO₃⁻) all may all have contributed to higher mineral N leaching losses in the inter-row than the
532 palm circle. In the inter-row, tThe main ~~accompanying~~ cation that accompanied the ~~of~~ leached
533 NO₃⁻ ~~in the inter row~~ was Al³⁺ (Figs. 3 and 4). This is because this zone's soil pH (Table 1) was

534 within the Al-buffering range (pH 3–5; van Breemen et al., 1983) ~~as because~~ this zone ~~had~~
535 ~~receives had~~ no direct lime application and ~~consequently had~~ a low base saturation (Table 1).
536 Our findings ~~also showed~~ that if leaching is measured only within the palm circle, this could
537 ~~lead to a substantially underestimate-underestimation of~~ mineral N and Al leaching losses.

538 ~~Despite the direct application of fertilizer. The~~ palm circle had relatively low N leaching
539 losses (Figs. 3 and 4), ~~which despite the direct application of fertilizer. This~~ was probably due
540 to the large root density, ~~in this zone that facilitates~~ an efficient nutrient uptake (Edy et al.,
541 2020; Nelson et al., 2006). ~~Hence, the~~ dominant anion in soil-pore water in the palm circle
542 was Cl⁻ (Fig. 3), ~~which was~~ enhanced by the applied KCl fertilizer, which was accompanied by
543 high base cation concentrations relative to dissolved Al (Fig. 3). The former was due to the
544 applied micromag fertilizer and dolomite (section 2.1), which increased pH and exchangeable
545 bases and rendered Al in insoluble form (~~i.e. lower exchangeable Al;~~ Table 1; Schlesinger and
546 Bernhardt, 2013). Despite their high concentrations, ~~base cations the~~ leaching fluxes of ~~base~~
547 ~~cations~~ in the palm circle (Fig. 4) were constrained by the low water drainage flux ~~due to high~~
548 ~~evapotranspiration~~ (Table 2).

549 ~~Although the~~ frond-stacked area was at the same distance from the palm circle as the
550 inter-row (Fig. A1), ~~but had substantially lower~~ mineral N leaching losses ~~were substantially~~
551 ~~lower~~ (Figs. 3 and 4). Decomposition of nutrient-rich fronds (Kotowska et al., 2016) resulted
552 in high SOC and N stocks (Table 1), which can support ~~a~~ large microbial biomass in this zone
553 (Haron et al., 1998). ~~Immobilization of mineral N by the large microbial biomass, converting~~
554 ~~mobile NO₃⁻ to less mobile organic N, may have caused~~ ~~Thus,~~ the low mineral N leaching in the
555 frond-stacked area ~~may be attributed to immobilization of mineral N by large microbial~~
556 ~~biomass, converting mobile NO₃⁻ to less mobile organic N~~ (e.g. Corre et al., 2010). In addition,
557 ~~it could be possible that~~ palm root uptake of nutrients (including mineral N) ~~was~~ ~~may have been~~
558 higher in the frond-stacked area ~~compared to~~ ~~than in~~ the inter-row ~~as because~~ roots ~~tend to~~

559 proliferate in nutrient-rich zones (Table 1; Hodge, 2004). ~~Indeed, This is supported by~~ studies
560 ~~that showed~~ have shown higher root density and higher water uptake under the frond piles
561 compared to the inter-row (Nelson et al., 2006; Rüegg et al., 2019). ~~The larger base cation~~
562 ~~leaching in the frond-stacked area compared to the inter-row (Fig. 4) were probably a reflection~~
563 ~~of (The high ECEC, base saturation and pH in frond-stacked area (Table 1). These favorable~~
564 ~~soil characteristics, despite having no direct lime application, were due to~~ were probably caused
565 by the release of nutrients from decomposition of frond litter, which contain high ~~levels of~~ base
566 cation concentrations (Kotowska et al., 2016). ~~Thus, the larger base cations leaching in the~~
567 ~~frond stacked area compared to the inter row (Fig. 4) merely mirrored their high exchangeable~~
568 ~~concentrations (Table 1).~~ Finally, the low Al leaching ~~of Al was low~~ in the frond-stacked area
569 (Figs. 3 and 4) ~~can be explained by the higher soil pH, because Al becomes insoluble and~~
570 ~~exchangeable Al decreases as pH increases (i.e. lower exchangeable Al; (Table 1). Altogether,~~
571 ~~these~~ Our results highlighted the benefits of piling senesced fronds ~~on to~~ the soil to reduce
572 leaching of mineral N and Al, which could otherwise ~~can potentially diminish~~ affect ground
573 water quality. ~~In other areas such as Borneo, oil palm plantations in other areas (e.g. Borneo;~~
574 ~~Rahman et al., 2018)~~ were reported to practice piling of senesced fronds on every inter-row
575 ~~(Rahman et al., 2018). I, which we did not observed in our study region this is~~ rarely
576 practiced as that is claimed because it to hinders access to palms during harvest. Maybe
577 chopping up senesced leaves with a shredder before spreading them on the soil can both
578 improve access and at the same time ~~Nonetheless, our findings implied that increase in the~~
579 ~~frond stacked area can contribute to sustainable~~ enhance nutrient management of oil palm
580 plantations.

581

582 4.3 Leaching losses under different intensity of management

583 ~~There was a clear influence of m~~Management intensity treatments ~~on strongly affected~~ nutrient
584 leaching losses with generally lower leaching fluxes ~~in under less intensive~~~~reduced~~
585 management ~~intensity~~ (Fig. 5; Table 3). In line with our second hypothesis, ~~the weeding~~
586 ~~methods clearly influenced leaching losses: the~~ mechanical weeding ~~treatment~~ had lower
587 nutrient leaching fluxes than the herbicide ~~treatment application~~ (Fig. 5; Table 5). Plots with
588 ~~m~~Mechanical weeding ~~was associated with had a better~~~~more~~higher ground vegetation cover
589 (Darras et al., 2019) and higher nutrient retention efficiency than herbicide weeding (Table 4).
590 ~~Leaching losses were probably retained better by, suggesting that~~ faster regrowth of understory
591 vegetation ~~by under~~ mechanical weeding ~~has additionally contributed to the uptake of nutrients~~
592 ~~and thus reducing leaching losses~~. This is in line with ~~some~~ studies in temperate forests and in
593 a cedar plantation ~~that showed~~ed that understory vegetation can take up excess NO₃⁻ in the soil
594 (Olsson and Falkengren-Grerup, 2003) and reduce NO₃⁻ leaching and the mobilization of Ca
595 and Mg (Fukuzawa et al., 2006; Baba et al., 2011). ~~Enhanced Denser~~ understory vegetation in
596 oil palm plantations may also positively impact biodiversity by increasing plant species richness
597 and soil macrofauna diversity and abundance (Ashton-Butt et al., 2018; Luke et al., 2019),
598 which may facilitate nutrient uptake and recycling ~~of nutrients~~. In addition, sincrease in soil
599 macrofauna ~~might may~~ have contributed to lower Na leaching ~~of Na~~ with mechanical weeding
600 (Fig. 5), ~~since because~~ herbivores and decomposers can take up a large substantial amounts of
601 Na (Kaspari et al., 2009). Following the first three years after establishment of the experiment,
602 oil palm y~~The yield in the first three years following the experiment establishment was on~~
603 ~~average~~approximately 30 Mg of fresh fruit bunches ha⁻¹ yr⁻¹ and did it was comparable~~not~~
604 ~~different~~ among experimental treatments (Figure A2₃; Darras et al. 2019). This ~~indicated~~
605 ~~shows~~attests that during the first three years the reduced management intensity did not affect
606 productivity. However, at least during the first three years, but the long-term monitoring of
607 yield is essential as it may take a longer time period before the yield to ~~responds~~ to our
608 experimental ~~the t~~reatments (e.g. Tao et al. 2017). ~~Also, the costs~~ of the two weeding

609 treatments (i.e. herbicide vs mechanical) ~~was comparable~~were not different because it is a
610 common practice to combine the use of herbicide with the periodic mechanical cutting of
611 resistant ground vegetation (Pahan, 2010; Darras et al., 2019). In addition, the use of glyphosate
612 ~~is has been~~ associated with possible health risks to workers and the environment (van Bruggen
613 et al., 2018). ~~Therefore, these~~In summary, our results ~~altogether~~ advocate for a higher-more
614 sustainable management sustainability of with mechanical weeding ~~over compared to~~ herbicide
615 application.

616 The decrease in NO₃⁻ leaching with ~~reduction of~~ N fertilization rates ~~decreased NO₃⁻~~
617 leaching, without affecting yield, supporting ~~ing~~ our third hypothesis. Our results suggest that
618 excess N applied with the conventional fertilization rate (above harvest export; section 2.1)
619 was largely lost through leaching (Table 3), as ~~Comparing conventional and reduced~~
620 ~~fertilization rates,~~ there were no differences in total N stocks (section 3.1), mineral N levels
621 (Darras et al., 2019), N retention efficiency (Table 4) and oil palm yield (Darras et al., 2019).
622 We attribute ~~suggesting that excess N (above harvest export; section 2.1) from high N~~
623 fertilization was largely lost through leaching (Table 3). The ~~decreased declines in~~ Al and Ca
624 leaching with reduced fertilization ~~can be attributed to the lowered~~ NO₃⁻ leaching, ~~since because~~
625 Al and Ca cations accompanied the leached NO₃⁻; these were the accompanying cations (Figs. 4
626 and 5). ~~The~~ Also, a reduction of Ca leaching may also be also related to ~~could have resulted~~
627 ~~from~~ the lower application rate of triple superphosphate fertilizer, which contains 16% of Ca.
628 The reduced K fertilization ~~had no effect on~~ did not affect K leaching (Fig. 5) probably because
629 K fertilization rates ~~was were~~ only reduced by 15% of the conventional rate due owing to high
630 K ~~requirements of export with harvested~~ oil palm fruits (section 2.1). ~~We conclude~~ Our study
631 provides evidence that this mature (16–20 years old) plantation with conventional management
632 was ~~overly over-~~ fertilized with N, and ~~that we suggest that inclusion of lower a reduction in~~ N
633 fertilization rates (related to N export with fruit bunches) may be included in the Indonesian

Formatted: Subscript

Formatted: Superscript

634 program for precision farming (Ministry of Agriculture of Indonesia, 2016) will substantially
635 and quickly improve the ~~to reduce~~ environmental footprint of oil palm production.

636 ~~Comparing the N leaching losses in the studied plantation with~~ Compared to other
637 fertilized tropical plantations (Table A3), our plantation had similar N leaching estimates
638 reported in another oil palm study using a model validated with field data from Suamatra
639 (Pardon et al. 2020). In contrast, higher-lower N leaching losses were ~~than~~ reported in other
640 large-scale oil palm plantations on similar soils with comparable fertilization rates (Omoti et
641 al., 1983; Tung et al., 2009). However, in these studies, ~~the~~ leaching losses were exclusively
642 measured in the palm circle (Omoti et al., 1983) or the sampling location was not specified
643 (Tung et al., 2009). Both studies may thus have ~~such that N leaching may be~~ underestimated
644 N leaching, because as our results showed the highest contribution ~~of the inter-row~~ to leaching
645 losses from the inter-row (Figs. 3 and 4). ~~Last, our values are in the same range as the N leaching~~
646 ~~estimated in another oil palm plantation study using a model that was validated with field data~~
647 ~~from Sumatra (Pardon et al. 2020). The~~ N leaching fluxes in our plantation were also higher
648 than fluxes reported from ~~in~~ smallholder oil palm plantations in the same area, which typically
649 ~~had much~~ owing to their lower fertilization rates (Kurniawan et al., 2018). In contrast, N
650 leaching ~~On the other hand, in~~ our plantation ~~had was~~ lower N leaching losses than from an an
651 oil palm plantation ~~and or~~ coffee agroforestry systems on volcanic soils (Banabas et al., 2008b;
652 Tully et al., 2012; Cannavo et al., 2013). ~~This may be caused by the, which have high~~ inherently
653 higher nutrient contents, and highly porosity of these volcanic soils ~~which that facilitates~~ ~~soils~~
654 ~~and~~ high infiltration rates. ~~The~~ N leaching losses from our plantation were also lower ~~than~~
655 ~~in~~ compared to banana plantations, which had substantially higher ~~characterized by very high~~
656 fertilization rates (Wakelin et al., 2011; Armour et al., 2013). ~~Last, our values are in the same~~
657 ~~range as the N leaching estimated in another oil palm plantation study using a model that was~~
658 ~~validated with field data from Sumatra (Pardon et al. 2020).~~

659 The ~~high fluxes of NO₃⁻ and Al nutrients leached~~ at 1.5 m depth ~~should be considered~~
660 ~~lost from uptake of oil palm roots, as the majority of the root mass and the highest root density~~
661 ~~are in the top 0.5 m depth (Nelson et al., 2006; Schroth et al., 2000; Kurniawan et al., 2018).~~
662 ~~The high leaching fluxes of NO₃⁻ and Al implies~~ied a substantial risk of groundwater pollution.
663 During the period of high drainage fluxes following fertilization, NO₃⁻ concentrations in soil-
664 pore water reached ~~to concentrations of~~ 20–40 mg NO₃⁻ L⁻¹ in the inter-row (covering 67% of
665 the plantation area), which ~~was is~~ close to the upper limit of 50 mg NO₃⁻ L⁻¹ ~~limit~~ for drinking
666 water (WHO, 2011), ~~and~~ Al concentrations in soil-pore water even exceeded the limit of 0.2
667 mg Al L⁻¹ in 60% of the samples. ~~Nevertheless, before reaching streams and rivers, This does~~
668 ~~not automatically mean that surface water will be contaminated, as these~~ NO₃⁻ and Al
669 concentrations can be diluted ~~by surface flow and~~ partially retained in the soil ~~along flow paths;~~
670 ~~NO₃⁻ can be temporarily adsorbed in the deeper layers of highly weathered soils by its~~
671 ~~inherently high anion exchange capacity (Harmand et al., 2010; Jankowski et al., 2018) and can~~
672 ~~be consumed by or~~ denitrifi~~ed~~ation (Wakelin et al., 2011). Such processes are especially
673 effective in r-Riparian buffers, which can mitigate the transport of these agricultural pollutants
674 to streams (Luke et al., 2017; Chellaiah and Yule, 2018). Our results thus also support the
675 importance of restoring ~~Restoring~~ riparian buffers in areas former forests converted to oil palm
676 plantations, which is also an important ~~have been listed as one~~ sustainability ~~eriteria~~ criterion ;
677 endorsed by the Roundtable for Sustainable Palm Oil association (RSPO, 2018), ~~and that~~ may
678 also provide additional regulation services (Woodham et al., 2019).

679 5 Conclusions

680 Our findings show that nutrient leaching losses in an oil palm plantation differed among
681 management zones, as a result of fertilization, liming, mulching and of different drainage
682 fluxes. ~~The reduction of management intensity, i.e. Implementation of-~~ mechanical weeding
683 with reduced fertilization rates, was effective in reducing nutrient leaching losses without

684 ~~reduction in affecting~~ yield ~~at least~~ during the first three years of this experiment. Long-term
685 investigation of this management experiment is important and planned in order to get a reliable
686 response of yield and to make a more holistic economic analysis ~~that,~~ including valuation of
687 regulation services. Greenhouse gas emissions should also be quantified, as another important
688 parameter of the environmental footprint of oil palm production. ~~Our~~ Our ultimate goal is that
689 our present and future findings ~~and these further investigations should~~ will be incorporated into
690 science-based policy recommendations such as those endorsed by the RSPO.

691 **Data availability**

692 All data of this study are deposited at the EFForTS-IS data repository ([https://efforts-is.uni-](https://efforts-is.uni-goettingen.de)
693 [goettingen.de](https://efforts-is.uni-goettingen.de)), an internal data-exchange platform, which is accessible to all members of the
694 Collaborative Research Center (CRC) 990. Based on the data sharing agreement within the
695 CRC 990, these data are currently not publicly accessible but will be made available through a
696 written request to the senior author.

697 **Author contribution**

698 GF performed the [experiments/field measurements](#), analysed the data and wrote the
699 manuscript in consultation with MDC. EV and MDC conceived and planned the experiment.
700 XD helped carry out the water model simulations. AT aided in [organizing the](#) field activities
701 [organization](#) and [facilitating the granting](#) collaborations [agreements among partners](#). All
702 authors contributed to the final version of the manuscript.

703 **Competing interests**

704 No conflict of interest to declare

705 **Acknowledgments**

706 This study was part of the project A05 in the CRC990-EFForTS, funded by the German
707 Research Foundation (DFG, Project ID: 192626868 – SFB 990). We acknowledge the
708 collaborations with PTPN VI plantation, Kevin Darras, and project Z01, for the
709 implementation and maintenance of this field experiment. We thank Christian Stiegler, with
710 project A03, for the climate data, and Eckart Priesack for the Expert-N water sub-model. We
711 especially thank our field and laboratory assistants for their valuable dedications in all field
712 and laboratory activities. This research was conducted under the research permit of Ministry
713 of Research and Technology of Indonesia, 539351/SIP/FRP/E5/Dit.KI/X/2016.

714 **References**

- 715 Abdalla, M., Hastings, A., Cheng, K., Yue, Q., Chadwick, D., Espenberg, M., Truu, J., Rees,
716 R. M., and Smith, P.: A critical review of the impacts of cover crops on nitrogen leaching,
717 net greenhouse gas balance and crop productivity, *Global Change Biology*, 25, 2530–
718 2543, doi:10.1111/gcb.14644, 2019.
- 719 Ahongshangbam, J., Khokthong, W., Ellsäßer, F., Hendrayanto, H., Hölscher, D., and Röhl,
720 A.: Drone-based photogrammetry-derived crown metrics for predicting tree and oil palm
721 water use, *Ecohydrology*, 12, 1–18, doi:10.1002/eco.2115, 2019.
- 722 Allen, K., Corre, M. D., Kurniawan, S., Utami, S. R., and Veldkamp, E.: Spatial variability
723 surpasses land-use change effects on soil biochemical properties of converted lowland
724 landscapes in Sumatra, Indonesia, *Geoderma*, 284, 42–50,
725 doi:10.1016/j.geoderma.2016.08.010, 2016.
- 726 Allen, K., Corre, M. D., Tjoa, A., and Veldkamp, E.: Soil nitrogen-cycling responses to
727 conversion of lowland forests to oil palm and rubber plantations in Sumatra, Indonesia,
728 *PloS One*, 10, e0133325, doi:10.1371/journal.pone.0133325, 2015.
- 729 Allen, R. G.: Crop evapotranspiration: Guidelines for computing crop water requirements,
730 *FAO irrigation and drainage paper*, 56, Rome, 300 pp., 1998.
- 731 Armour, J. D., Nelson, P. N., Daniells, J. W., Rasiyah, V., and Inman-Bamber, N. G.: Nitrogen
732 leaching from the root zone of sugarcane and bananas in the humid tropics of Australia,
733 *Agriculture, Ecosystems & Environment*, 180, 68–78, doi:10.1016/j.agee.2012.05.007,
734 2013.
- 735 Ashton-Butt, A., Aryawan, A. A. K., Hood, A. S. C., Naim, M., Purnomo, D., Suhardi,
736 Wahyuningsih, R., Willcock, S., Poppy, G. M., Caliman, J.-P., Turner, E. C., Foster, W.
737 A., Peh, K. S. H., and Snaddon, J. L.: Understory vegetation in oil palm plantations
738 benefits soil biodiversity and decomposition rates, *Frontiers in Forests and Global Change*,
739 1, 1–13, doi:10.3389/ffgc.2018.00010, 2018.

740 Asner, G. P., Powell, G. V. N., Mascaro, J., Knapp, D. E., Clark, J. K., Jacobson, J., Kennedy-
741 Bowdoin, T., Balaji, A., Paez-Acosta, G., Victoria, E., Secada, L., Valqui, M., and
742 Hughes, F. R.: High-resolution forest carbon stocks and emissions in the Amazon,
743 Proceedings of the National Academy of Sciences of the United States of America, 107,
744 16738–16742, doi:10.1073/pnas.1004875107, 2010.

745 Baba, M., Abe, S., Kasai, M., Sugiura, T., and Kobayashi, H.: Contribution of understory
746 vegetation to minimizing nitrate leaching in a Japanese cedar plantation, Journal of Forest
747 Research, 16, 446–455, doi:10.1007/s10310-010-0244-3, 2011.

748 Banabas, M., Scotter, D. R., and Turner, M. A.: Losses of nitrogen fertiliser under oil palm in
749 Papua New Guinea: 2. Nitrogen transformations and leaching, and a residence time model,
750 Soil Research, 46, 340–347, doi:10.1071/SR07174, 2008a.

751 Banabas, M., Turner, M. A., Scotter, D. R., and Nelson, P. N.: Losses of nitrogen fertiliser
752 under oil palm in Papua New Guinea: 1. Water balance, and nitrogen in soil solution and
753 runoff, Australian Journal of Soil Research, 46, 332–339, doi:10.1071/SR07171, 2008b.

754 Bates, D., Maechler, M., Bolker, B., and Walker, S.: Fitting linear mixed-effect models using
755 lme4, Journal of Statistical Software, 67, 1–48, 2015.

756 Boafo, D. K., Kraisornpornson, B., Panphon, S., Owusu, B. E., and Amaniampong, P. N.:
757 Effect of organic soil amendments on soil quality in oil palm production, Applied Soil
758 Ecology, 147, 103358, doi:10.1016/j.apsoil.2019.09.008, 2020.

759 Cannavo, P., Harmand, J.-M., Zeller, B., Vaast, P., Ramírez, J. E., and Dambrine, E.: Low
760 nitrogen use efficiency and high nitrate leaching in a highly fertilized *Coffea arabica*–*Inga*
761 *densiflora* agroforestry system: a 15N labeled fertilizer study, Nutrient Cycling in
762 Agroecosystems, 95, 377–394, doi:10.1007/s10705-013-9571-z, 2013.

763 Carlson, K. M., Curran, L. M., Asner, G. P., Pittman, A. M., Trigg, S. N., and Marion
764 Adeney, J.: Carbon emissions from forest conversion by Kalimantan oil palm plantations,
765 Nature Climate Change, 3, 283–287, doi:10.1038/nclimate1702, 2013.

766 Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and Smith,
767 V. H.: Nonpoint pollution of surface waters with phosphorus and nitrogen, *Ecological*
768 *Applications*, 8, 559–568, doi:10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2,
769 1998.

770 Carter, C., Finley, W., Fry, J., Jackson, D., and Willis, L.: Palm oil markets and future supply,
771 *European Journal of Lipid Science and Technology*, 109, 307–314,
772 doi:10.1002/ejlt.200600256, 2007.

773 Chellaiah, D. and Yule, C. M.: Effect of riparian management on stream morphometry and
774 water quality in oil palm plantations in Borneo, *Limnologica*, 69, 72–80,
775 doi:10.1016/j.limno.2017.11.007, 2018.

776 Chou, C., Chiang, J. C. H., Lan, C.-W., Chung, C.-H., Liao, Y.-C., and Lee, C.-J.: Increase in
777 the range between wet and dry season precipitation, *Nature Geoscience*, 6, 263–267,
778 doi:10.1038/ngeo1744, 2013.

779 Clough, Y., Krishna, V. V., Corre, M. D., Darras, K., Denmead, L. H., Mejjide, A., Moser, S.,
780 Musshoff, O., Steinebach, S., Veldkamp, E., Allen, K., Barnes, A. D., Breidenbach, N.,
781 Brose, U., Buchori, D., Daniel, R., Finkeldey, R., Harahap, I., Hertel, D., Holtkamp, A.
782 M., Hörandl, E., Irawan, B., Jaya, I. N. S., Jochum, M., Klärner, B., Knohl, A., Kotowska,
783 M. M., Krashevskaya, V., Kreft, H., Kurniawan, S., Leuschner, C., Maraun, M., Melati, D.
784 N., Opfermann, N., Pérez-Cruzado, C., Prabowo, W. E., Rembold, K., Rizali, A., Rubiana,
785 R., Schneider, D., Tjitrosoedirdjo, S. S., Tjoa, A., Tschardtke, T., and Scheu, S.: Land-use
786 choices follow profitability at the expense of ecological functions in Indonesian
787 smallholder landscapes, *Nature Communications*, 7, 13137, doi:10.1038/ncomms13137,
788 2016.

789 Comte, I., Colin, F., Grünberger, O., Follain, S., Whalen, J. K., and Caliman, J.-P.:
790 Landscape-scale assessment of soil response to long-term organic and mineral fertilizer

791 application in an industrial oil palm plantation, Indonesia, *Agriculture, Ecosystems &*
792 *Environment*, 169, 58–68, doi:10.1016/j.agee.2013.02.010, 2013.

793 Corley, R. H. V. and Tinker, P. B.: *The oil palm*, 5th ed., John Wiley & Sons, Ltd, Hoboken,
794 NJ, 2016.

795 Corley, R.H.V.: How much palm oil do we need?, *Environmental Science & Policy*, 12, 134–
796 139, doi:10.1016/j.envsci.2008.10.011, 2009.

797 Corre, M. D., Veldkamp, E., Arnold, J., and Wright, S. J.: Impact of elevated N input on soil
798 N cycling and losses in old-growth lowland and montane forests in Panama, *Ecology*, 91,
799 1715–1729, doi:10.1890/09-0274.1, 2010.

800 Cusack, D. F., Silver, W., and McDowell, W. H.: Biological nitrogen fixation in two tropical
801 forests: ecosystem-level patterns and effects of nitrogen fertilization, *Ecosystems*, 12,
802 1299–1315, doi:10.1007/s10021-009-9290-0, 2009.

803 Darras, K. F. A., Corre, M. D., Formaglio, G., Tjoa, A., Potapov, A., Brambach, F., Sibhatu,
804 K. T., Grass, I., Rubiano, A. A., Buchori, D., Drescher, J., Fardiansah, R., Hölscher, D.,
805 Irawan, B., Kneib, T., Krashevskaya, V., Krause, A., Kreft, H., Li, K., Maraun, M., Polle,
806 A., Ryadin, A. R., Rembold, K., Stiegler, C., Scheu, S., Tarigan, S., Valdés-Uribe, A.,
807 Yadi, S., Tschardtke, T., and Veldkamp, E.: Reducing fertilizer and avoiding herbicides in
808 O oil palm plantations—ecological and economic valuations, *Frontiers in Forests and*
809 *Global Change*, 2, doi:10.3389/ffgc.2019.00065, 2019.

810 Dechert, G., Veldkamp, E., and Brumme, R.: Are partial nutrient balances suitable to evaluate
811 nutrient sustainability of land use systems? Results from a case study in central Sulawesi,
812 Indonesia, *Nutrient Cycling in Agroecosystems*, 72, 201–212, doi:10.1007/s10705-005-
813 1546-2, 2005.

814 Drescher, J., Rembold, K., Allen, K., Beckschäfer, P., Buchori, D., Clough, Y., Faust, H.,
815 Fauzi, A. M., Gunawan, D., Hertel, D., Irawan, B., Jaya, I. N. S., Klarner, B., Kleinn, C.,
816 Knohl, A., Kotowska, M. M., Krashevskaya, V., Krishna, V., Leuschner, C., Lorenz, W.,

817 Meijide, A., Melati, D., Nomura, M., Pérez-Cruzado, C., Qaim, M., Siregar, I. Z.,
818 Steinebach, S., Tjoa, A., Tschardtke, T., Wick, B., Wiegand, K., Kreft, H., and Scheu, S.:
819 Ecological and socio-economic functions across tropical land use systems after rainforest
820 conversion, *Philosophical Transactions of the Royal Society of London. Series B,*
821 *Biological sciences*, 371, doi:10.1098/rstb.2015.0275, 2016.

822 Dubos, B., Snoeck, D., and Flori, A.: Excessive use of fertilizer can increase leaching
823 processes and modify soil reserves in two Ecuadorian oil palm plantations, *Experimental*
824 *Agriculture*, 53, 255–268, doi:10.1017/S0014479716000363, 2017.

825 Edy, N., Yelianti, U., Irawan, B., Polle, A., and Pena, R.: Differences in root nitrogen uptake
826 between tropical lowland rainforests and oil palm plantations, *Frontiers in Plant Science*,
827 11, 92, doi:10.3389/fpls.2020.00092, 2020.

828 Elsenbeer, H.: Hydrologic flowpaths in tropical rainforest soilsclapes - a review, *Hydrological*
829 *Processes*, 15, 1751–1759, doi:10.1002/hyp.237, 2001.

830 Fan, Y., Roupsard, O., Bernoux, M., Le Maire, G., Panferov, O., Kotowska, M. M., and
831 Knohl, A.: A sub-canopy structure for simulating oil palm in the Community Land Model
832 (CLM-Palm): phenology, allocation and yield, *Geoscientific Model Development*, 8,
833 3785–3800, doi:10.5194/gmd-8-3785-2015, 2015.

834 Feng, X., Porporato, A., and Rodriguez-Iturbe, I.: Changes in rainfall seasonality in the
835 tropics, *Nature Climate Change*, 3, 811–815, doi:10.1038/nclimate1907, 2013.

836 Figueiredo, R. O., Markewitz, D., Davidson, E. A., Schuler, A. E., Watrin, O. d. S., and Silva,
837 P. d. S.: Land-use effects on the chemical attributes of low-order streams in the eastern
838 Amazon, *Journal of Geophysical Research*, 115, doi:10.1029/2009JG001200, 2010.

839 Food and Agriculture Organization, FAOSTAT: <http://faostat.fao.org/site/339/default.aspx>,
840 last access: 30 March 2020.

841 Fukuzawa, K., Shibata, H., Takagi, K., Nomura, M., Kurima, N., Fukazawa, T., Satoh, F., and
842 Sasa, K.: Effects of clear-cutting on nitrogen leaching and fine root dynamics in a cool-

843 temperate forested watershed in northern Japan, *Forest Ecology and Management*, 225,
844 257–261, doi:10.1016/j.foreco.2006.01.001, 2006.

845 Geist, H. J. and Lambin, E. F.: Proximate causes and underlying driving forces of tropical
846 deforestation, *BioScience*, 52, 143–150, doi:10.1641/0006-
847 3568(2002)052[0143:PCAUDF]2.0.CO;2, 2002.

848 Gibbs, H. K., Ruesch, A. S., Achard, F., Clayton, M. K., Holmgren, P., Ramankutty, N., and
849 Foley, J. A.: Tropical forests were the primary sources of new agricultural land in the
850 1980s and 1990s, *Proceedings of the National Academy of Sciences of the United States*
851 *of America*, 107, 16732–16737, doi:10.1073/pnas.0910275107, 2010.

852 Goh, K.-J., Härdter, R., and Fairhurst, T.: Fertilizing for maximum return, *Oil Palm*
853 *Management for Large and Sustainable Yields*, 279–306, 2003.

854 Grass, I., Kubitzka, C., Krishna, V. V., Corre, M. D., Mußhoff, O., Pütz, P., Drescher, J.,
855 Rembold, K., Ariyanti, E. S., Barnes, A. D., Brinkmann, N., Brose, U., Brümmer, B.,
856 Buchori, D., Daniel, R., Darras, K. F. A., Faust, H., Fehrmann, L., Hein, J., Hennings, N.,
857 Hidayat, P., Hölscher, D., Jochum, M., Knohl, A., Kotowska, M. M., Krashevskaya, V.,
858 Kreft, H., Leuschner, C., Lobite, N. J. S., Panjaitan, R., Polle, A., Potapov, A. M.,
859 Purnama, E., Qaim, M., Röhl, A., Scheu, S., Schneider, D., Tjoa, A., Tschardtke, T.,
860 Veldkamp, E., and Wollni, M.: Trade-offs between multifunctionality and profit in
861 tropical smallholder landscapes, *Nature Communications*, 11, 1186, doi:10.1038/s41467-
862 020-15013-5, 2020.

863 Harmand, J.-M., Ávila, H., Oliver, R., Saint-André, L., and Dambrine, E.: The impact of
864 kaolinite and oxi-hydroxides on nitrate adsorption in deep layers of a Costarican Acrisol
865 under coffee cultivation, *Geoderma*, 158, 216–224, doi:10.1016/j.geoderma.2010.04.032,
866 2010.

867 Haron, K., Brookes, P. C., Anderson, J. M., and Zakaria, Z. Z.: Microbial biomass and soil
868 organic matter dynamics in oil palm (*Elaeis Guineensis* Jacq.) plantations, West Malaysia,
869 Soil Biology and Biochemistry, 30, 547–552, doi:10.1016/S0038-0717(97)00217-4, 1998.

870 Hillel, D: Introduction to Soil Physics, 107–114, Academic Press, California, USA, 1982.

871 Hodge, A.: The plastic plant: root responses to heterogeneous supplies of nutrients, New
872 Phytologist, 162, 9–24, doi:10.1111/j.1469-8137.2004.01015.x, 2004.

873 Hothorn, T., Bretz, F., and Westfall, P.: Simultaneous inference in general parametric models,
874 Biometrical Journal, 50, 346–363, 2008.

875 ~~Huddell, A. M., Galford, G. L., Tully, K. L., Crowley, C., Palm, C. A., Neill, C., Hickman, J.
876 E., and Menge, D. N. L.: Meta analysis on the potential for increasing nitrogen losses
877 from intensifying tropical agriculture, Global Change Biology, 26, 1–13,
878 doi:10.1111/gcb.14951, 2020.~~

879 Jankowski, K., Neill, C., Davidson, E. A., Macedo, M. N., Costa, C. J., Galford, G. L.,
880 Santos, L. M., Lefebvre, P., Nunes, D., Cerri, C. E. P., McHorney, R., O’Connell, C., and
881 Coe, M. T.: Deep soils modify environmental consequences of increased nitrogen fertilizer
882 use in intensifying Amazon agriculture, Scientific Reports, 13478, doi:10.1038/s41598-
883 018-31175-1, 2018.

884 Jourdan, C. and Rey, H.: Modelling and simulation of the architecture and development of the
885 oil-palm (*Elaeis guineensis* Jacq.) root system, Plant and Soil, 190, 235–246,
886 doi:10.1023/A:1004270014678, 1997.

887 Kaspari, M., Yanoviak, S. P., Dudley, R., Yuan, M., and Clay, N. A.: Sodium shortage as a
888 constraint on the carbon cycle in an inland tropical rainforest, Proceedings of the National
889 Academy of Sciences of the United States of America, 106, 19405–19409,
890 doi:10.1073/pnas.0906448106, 2009.

891

892 Kotowska, M. M., Leuschner, C., Triadiati, T., and Hertel, D.: Conversion of tropical lowland
893 forest reduces nutrient return through litterfall, and alters nutrient use efficiency and
894 seasonality of net primary production, *Oecologia*, 180, 601–618, doi:10.1007/s00442-015-
895 3481-5, 2016.

896 Kotowska, M. M., Leuschner, C., Triadiati, T., Meriem, S., and Hertel, D.: Quantifying
897 above- and belowground biomass carbon loss with forest conversion in tropical lowlands
898 of Sumatra (Indonesia), *Global Change Biology*, 21, 3620–3634, doi:10.1111/gcb.12979,
899 2015.

900 Kurniawan, S., Corre, M. D., Matson, A. L., Schulte-Bisping, H., Utami, S. R., van Straaten,
901 O., and Veldkamp, E.: Conversion of tropical forests to smallholder rubber and oil palm
902 plantations impacts nutrient leaching losses and nutrient retention efficiency in highly
903 weathered soils, *Biogeosciences*, 15, 5131–5154, doi:10.5194/bg-15-5131-2018, 2018.

904 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: LmerTest package: tests in linear
905 mixed effects models, *Journal of Statistical Software*, 82, doi:10.18637/jss.v082.i13, 2017.

906 Lamade, E., Djegui, N., and Leterme, P.: Estimation of carbon allocation to the roots from
907 soil respiration measurements of oil palm, *Plant and Soil*, 181, 329–339,
908 doi:10.1007/BF00012067, 1996.

909 Li, Y. and Ghodrati, M.: Preferential transport of nitrate through soil columns containing root
910 channels, *Soil Science Society of America Journal*, 58, 653–659,
911 doi:10.2136/sssaj1994.03615995005800030003x, 1994.

912 Lohse, K. A. and Matson, P.: Consequences of nitrogen additions for soil losses from wet
913 tropical forests, *Ecological Applications*, 15, 1629–1648, doi:10.1890/03-5421, 2005.

914 Luke, S. H., Barclay, H., Bidin, K., Chey, V. K., Ewers, R. M., Foster, W. A., Nainar, A.,
915 Pfeifer, M., Reynolds, G., Turner, E. C., Walsh, R. P. D., and Aldridge, D. C.: The effects
916 of catchment and riparian forest quality on stream environmental conditions across a
917 tropical rainforest and oil palm landscape in Malaysian Borneo, *Ecohydrology*:

918 ecosystems, land and water process interactions, *ecohydrogeomorphology*, 10, 1-17,
919 doi:10.1002/eco.1827, 2017.

920 Luke, S. H., Purnomo, D., Advento, A. D., Aryawan, A. A. K., Naim, M., Pikstein, R. N., Ps,
921 S., Rambe, T. D. S., Soeprpto, Caliman, J.-P., Snaddon, J. L., Foster, W. A., and Turner,
922 E. C.: Effects of understory vegetation management on plant communities in oil palm
923 plantations in Sumatra, Indonesia, *Frontiers in Forests and Global Change*, 2, 1–13,
924 doi:10.3389/ffgc.2019.00033, 2019.

925 Meijide, A., La Rua, C. de, Guillaume, T., Röhl, A., Hassler, E., Stiegler, C., Tjoa, A., June,
926 T., Corre, M. D., Veldkamp, E., and Knohl, A.: Measured greenhouse gas budgets
927 challenge emission savings from palm-oil biodiesel, *Nature Communications*, 11, 1089,
928 doi:10.1038/s41467-020-14852-6, 2020.

929 Meijide, A., Röhl, A., Fan, Y., Herbst, M., Niu, F., Tiedemann, F., June, T., Rauf, A.,
930 Hölscher, D., and Knohl, A.: Controls of water and energy fluxes in oil palm plantations:
931 environmental variables and oil palm age, *Agricultural and Forest Meteorology*, 239, 71–
932 85, doi:10.1016/j.agrformet.2017.02.034, 2017.

933 Ministry of Agriculture of Indonesia: Oil palm replanting guideline, Regulation No:
934 18/Permentan/KB.330/5/2016, 2016.

935 Mohanty, S., Swain, C. K., Tripathi, R., Sethi, S. K., Bhattacharyya, P., Kumar, A., Raja, R.,
936 Shahid, M., Panda, B. B., Lal, B., Gautam, P., Munda, S., and Nayak, A. K.: Nitrate
937 leaching, nitrous oxide emission and N use efficiency of aerobic rice under different N
938 application strategy, *Archives of Agronomy and Soil Science*, 64, 465–479,
939 doi:10.1080/03650340.2017.1359414, 2018.

940 Moradi, A., Teh, C. B. S., Goh, K. J., Husni, A. M. H., and Ishak, C. F.: Effect of four soil
941 and water conservation practices on soil physical processes in a non-terraced oil palm
942 plantation, *Soil and Tillage Research*, 145, 62–71, doi:10.1016/j.still.2014.08.005, 2015.

943 Murdiyarso, D., Hergoualc'h, K., and Verchot, L. V.: Opportunities for reducing greenhouse
944 gas emissions in tropical peatlands, *Proceedings of the National Academy of Sciences of*
945 *the United States of America*, 107, 19655–19660, doi:10.1073/pnas.0911966107, 2010.

946 Neill, C., Coe, M. T., Riskin, S. H., Krusche, A. V., Elsenbeer, H., Macedo, M. N.,
947 McHorney, R., Lefebvre, P., Davidson, E. A., Scheffler, R., Figueira, A. M. e. S., Porder,
948 S., and Deegan, L. A.: Watershed responses to Amazon soya bean cropland expansion and
949 intensification, *Philosophical Transactions of the Royal Society of London. Series B,*
950 *Biological sciences*, 368, 20120425, doi:10.1098/rstb.2012.0425, 2013.

951 Nelson, P. N., Banabas, M., Scotter, D. R., and Webb, M. J.: Using soil water depletion to
952 measure spatial distribution of root activity in oil palm (*Elaeis Guineensis* Jacq.)
953 plantations, *Plant and Soil*, 286, 109–121, doi:10.1007/s11104-006-9030-6, 2006.

954 Nyamangara, J., Bergström, L. F., Piha, M. I., and Giller, K. E.: Fertilizer use efficiency and
955 nitrate leaching in a tropical sandy soil, *Journal of environmental quality*, 32, 599–606,
956 doi:10.2134/jeq2003.5990, 2003.

957 Olsson, M. O. and Falkengren-Grerup, U.: Partitioning of nitrate uptake between trees and
958 understory in oak forests, *Forest Ecology and Management*, 179, 311–320,
959 doi:10.1016/S0378-1127(02)00544-3, 2003.

960 Omoti, U., Ataga, D. O., and Isenmila, A. E.: Leaching losses of nutrients in oil palm
961 plantations determined by tension lysimeters, *Plant and Soil*, 73, 365–376,
962 doi:10.1007/BF02184313, 1983.

963 Pahan, I.: *Complete guide to oil palm*, 8th ed., Penebar Swadaya, Jakarta, Indonesia, 160-164,
964 2010.

965 Pardon, L., Bessou, C., Nelson, P. N., Dubos, B., Ollivier, J., Marichal, R., Caliman, J.-P., and
966 Gabrielle, B.: Key unknowns in nitrogen budget for oil palm plantations. A review,
967 *Agronomy for Sustainable Development*, 36, 20, doi:10.1007/s13593-016-0353-2, 2016.

968 Pardon, L., Bockstaller, C., Marichal, R., Sionita, R., Nelson, P. N., Gabrielle, B., Laclau, J.-
969 P., Pujianto, Caliman, J.-P., and Bessou, C.: IN-Palm: An agri-environmental indicator to
970 assess nitrogen losses in oil palm plantations, *Agronomy Journal*, 112, 786–800,
971 doi:10.1002/agj2.20109, 2020.

972 Pirker, J., Mosnier, A., Kraxner, F., Havlík, P., and Obersteiner, M.: What are the limits to oil
973 palm expansion?, *Global Environmental Change*, 40, 73–81,
974 doi:10.1016/j.gloenvcha.2016.06.007, 2016.

975 Priesack, E.: Expert-N model library documentation, Institute of Soil Ecology, 2005.

976 R Core Team: R: A language and environment for statistical computing, Foundation for
977 Statistical Computing, Vienna, Austria, 2019.

978 Rahman, N., Neergaard, A. de, Magid, J., van de Ven, G. W. J., Giller, K. E., and Bruun, T.
979 B.: Changes in soil organic carbon stocks after conversion from forest to oil palm
980 plantations in Malaysian Borneo, *Environmental Research Letters*, 13, 105001,
981 doi:10.1088/1748-9326/aade0f, 2018.

982

983 Röhl, A., Niu, F., Meijide, A., Ahongshangbam, J., Ehbrecht, M., Guillaume, T., Gunawan,
984 D., Hardanto, A., Hendrayanto, Hertel, D., Kotowska, M. M., Kreft, H., Kuzyakov, Y.,
985 Leuschner, C., Nomura, M., Polle, A., Rembold, K., Sahner, J., Seidel, D., Zemp, D. C.,
986 Knohl, A., and Hölscher, D.: Transpiration on the rebound in lowland Sumatra,
987 *Agricultural and Forest Meteorology*, 274, 160–171, doi:10.1016/j.agrformet.2019.04.017,
988 2019.

989 Roy, E. D., Richards, P. D., Martinelli, L. A., Della Coletta, L., Lins, S. R. M., Vazquez, F.
990 F., Willig, E., Spera, S. A., VanWey, L. K., and Porder, S.: The phosphorus cost of
991 agricultural intensification in the tropics, *Nature plants*, 2, 16043,
992 doi:10.1038/nplants.2016.43, 2016.

993 RSPO: Principles and criteria: For the production of sustainable palm oil, 2018.

994 Rüegg, J., Quezada, J. C., Santonja, M., Ghazoul, J., Kuzyakov, Y., Buttler, A., and
995 Guillaume, T.: Drivers of soil carbon stabilization in oil palm plantations, *Land*
996 *Degradation & Development*, 30, 1–12, doi:10.1002/ldr.3380, 2019.

997 Schlesinger, W. H. and Bernhardt, E. S.: *Biogeochemistry: An analysis of global change*, 3rd
998 ed., Academic Press, Waltham, Mass, 672 pp., 2013.

999 Schroth, G., Rodrigues, M.R.L., and D'Angelo, S. A.: Spatial patterns of nitrogen
1000 mineralization, fertilizer distribution and roots explain nitrate leaching from mature
1001 Amazonian oil palm plantation, *Soil Use and Management*, 16, 222–229,
1002 doi:10.1111/j.1475-2743.2000.tb00197.x, 2000.

1003 Silver, W. L., Thompson, A. W., Reich, A., Ewel, J. J., and Firestone, M. K.: Nitrogen
1004 cycling in tropical plantation forests: potential controls on nitrogen retention, *Ecological*
1005 *Applications*, 15, 1604–1614, doi:10.1890/04-1322, 2005.

1006 Steiner, C., Glaser, B., Gerdal Teixeira, W., Lehmann, J., Blum, W. E.H., and Zech, W.:
1007 Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol
1008 amended with compost and charcoal, *Journal of Plant Nutrition and Soil Science*, 171,
1009 893–899, doi:10.1002/jpln.200625199, 2008.

1010 Syers, J. K.: Managing soils for long-term productivity, *Phil. Trans. R. Soc. Lond. B*, 352,
1011 1011–1021, doi:10.1098/rstb.1997.0079, 1997.

1012 Tao, H.-H., Snaddon, J. L., Slade, E. M., Caliman, J.-P., Widodo, R. H., Suhardi, and Willis,
1013 K. J.: Long-term crop residue application maintains oil palm yield and temporal stability
1014 of production, *Agronomy for Sustainable Development*, 37, 33, doi:10.1007/s13593-017-
1015 0439-5, 2017.

1016 Tarigan, S., Stiegler, C., Wiegand, K., Knohl, A., and Murti Laksono, K.: Relative contribution
1017 of evapotranspiration and soil compaction to the fluctuation of catchment discharge: case
1018 study from a plantation landscape, *Hydrological Sciences Journal*, 58, 1–10,
1019 doi:10.1080/02626667.2020.1739287, 2020.

1020 Tarigan, S., Wiegand, K., Sunarti, and Slamet, B.: Minimum forest cover required for
1021 sustainable water flow regulation of a watershed: a case study in Jambi Province,
1022 Indonesia, *Hydrology and Earth System Sciences*, 22, 581–594, doi:10.5194/hess-22-581-
1023 2018, 2018.

1024 Tarigan, S. D., Sunarti, Wiegand, K., Dislich, C., Slamet, B., Heinonen, J., and Meyer, K.:
1025 Mitigation options for improving the ecosystem function of water flow regulation in a
1026 watershed with rapid expansion of oil palm plantations, *Sustainability of Water Quality
1027 and Ecology*, 8, 4–13, doi:10.1016/j.swaqe.2016.05.001, 2016.

1028 Teklu, B. M., Hailu, A., Wiegand, D. A., Scholten, B. S., and van den Brink, P. J.: Impacts of
1029 nutrients and pesticides from small- and large-scale agriculture on the water quality of
1030 Lake Ziway, Ethiopia, *Environmental Science and Pollution Research*, 25, 13207–13216,
1031 doi:10.1007/s11356-016-6714-1, 2018.

1032 Tokuchi, N., Samejima, H., Hon, J., and Fukushima, K.: Influence of herbicide use in oil palm
1033 plantations on stream water chemistry in Sarawak, *Anthropogenic Tropical Forests*, 209–
1034 216, doi:10.1007/978-981-13-7513-2_11, 2019.

1035 Tully, K. L., Lawrence, D., and Scanlon, T. M.: More trees less loss: nitrogen leaching losses
1036 decrease with increasing biomass in coffee agroforests, *Agriculture, Ecosystems &
1037 Environment*, 161, 137–144, doi:10.1016/j.agee.2012.08.002, 2012.

1038 Tung, P. G. A., Yusoff, M. K., Majid, N. M., Joo, G. K., and Huang, G. H.: Effect of N and K
1039 fertilizers on nutrient leaching and groundwater quality under mature oil palm in Sabah
1040 during the monsoon period, *American Journal of Applied Sciences*, 6, 1788–1799, 2009.

1041 van Breemen, N., Mulder, J., and Driscoll, C. T.: Acidification and alkalization of soils, *Plant
1042 and Soil*, 283–308, 1983.

1043 van Bruggen, A.H.C., He, M. M., Shin, K., Mai, V., Jeong, K. C., Finckh, M. R., and Morris,
1044 J. G.: Environmental and health effects of the herbicide glyphosate, *Science of the Total
1045 Environment*, 616-617, 255–268, doi:10.1016/j.scitotenv.2017.10.309, 2018.

1046 van Straaten, O., Corre, M. D., Wolf, K., Tchienkoua, M., Cuellar, E., Matthews, R. B., and
1047 Veldkamp, E.: Conversion of lowland tropical forests to tree cash crop plantations loses
1048 up to one-half of stored soil organic carbon, Proceedings of the National Academy of
1049 Sciences of the United States of America, 112, 9956–9960, doi:10.1073/pnas.1504628112,
1050 2015.

1051 Veldkamp, E., Schmidt, M., Powers, J. S., and Corre, M. D.: Deforestation and reforestation
1052 impacts on soils in the tropics, Nature Reviews Earth & Environment, *in press*.

1053 Vijay, V., Pimm, S. L., Jenkins, C. N., and Smith, S. J.: The impacts of oil palm on recent
1054 deforestation and biodiversity loss, PloS One, 11, e0159668,
1055 doi:10.1371/journal.pone.0159668, 2016.

1056 Wakelin, S. A., Nelson, P. N., Armour, J. D., Rasiah, V., and Colloff, M. J.: Bacterial
1057 community structure and denitrifier (*nir-gene*) abundance in soil water and groundwater
1058 beneath agricultural land in tropical North Queensland, Australia, Soil Research, 49, 65–
1059 76, doi:10.1071/SR10055, 2011.

1060 WHO: Guidelines for drinking-water quality, 4th ed., World Health Organization, Geneva,
1061 2011.

1062 Wong, M. T.F., Hughes, R., and Rowell, D. L.: Retarded leaching of nitrate in acid soils from
1063 the tropics: measurement of the effective anion exchange capacity, Journal of Soil
1064 Science, 655–663, 1990.

1065 Woodham, C. R., Aryawan, A. A. K., Luke, S. H., Manning, P., Caliman, J.-P., Naim, M.,
1066 Turner, E. C., and Slade, E. M.: Effects of replanting and retention of mature oil palm
1067 riparian buffers on Ecosystem functioning in oil palm plantations, Frontiers in Forests and
1068 Global Change, 2, 29, doi:10.3389/ffgc.2019.00029, 2019.

1069

Formatted: Font: Italic

1070 **Tables and figures**

1071 **Table 1** Soil physical and biochemical characteristics (mean \pm standard errors, $n = 4$ plots) in
 1072 the top 50 cm depth for each management zone, averaged across experimental treatments.
 1073 Means within a row followed by different letters indicate significant differences among
 1074 management zones (one-way ANOVA with Tukey HSD or Kruskal–Wallis H test with multiple
 1075 comparisons extension at $P \leq 0.05$). Bulk density measured in the top 10 cm of soil, whereas
 1076 all the other parameters are for the 0–50 cm soil depth: element stocks are the sum of the
 1077 sampled soil depths (0–5 cm, 5–10 cm, 10–30 cm and 30–50 cm) and the rest are depth-
 1078 weighted averages, calculated for each replicate plot. ECEC, effective cation exchange capacity

Soil properties		Palm circle	FronD-stacked area	Inter-row
Bulk density	g cm^{-3}	1.37 ± 0.01^a	0.89 ± 0.01^b	1.36 ± 0.01^b
Soil organic C	kg m^{-2}	6.2 ± 0.6^b	9.1 ± 0.8^a	6.4 ± 0.2^b
Total N	g m^{-2}	402 ± 31^b	571 ± 39^a	426 ± 15^{ab}
soil C:N ratio		15.5 ± 0.5^a	15.7 ± 0.3^a	15.0 ± 0.5^a
^{15}N natural abundance	‰	5.9 ± 0.1^a	5.3 ± 0.2^a	5.7 ± 0.2^a
pH	1:4 (H ₂ O)	5.05 ± 0.08^a	5.00 ± 0.08^{ab}	4.81 ± 0.05^b
ECEC	$\text{mmol}_c \text{kg}^{-1}$	35 ± 2^a	28 ± 2^a	18 ± 1^b
Base saturation	%	48 ± 3^a	46 ± 4^a	20 ± 2^b
Aluminum saturation	%	52 ± 4^b	50 ± 2^b	78 ± 2^a
Mg	g m^{-2}	32 ± 3^a	28 ± 6^a	9 ± 1^b
Ca	g m^{-2}	169 ± 21^a	157 ± 15^a	37 ± 5^b
K	g m^{-2}	39 ± 13^a	13 ± 1^b	6 ± 1^b
Na	g m^{-2}	1.5 ± 0.4^a	0.7 ± 0.2^a	0.6 ± 0.2^a
Al	g m^{-2}	66 ± 4^b	71 ± 4^{ab}	87 ± 3^a

Fe	g m ⁻²	1.4 ± 0.2 ^a	1.8 ± 0.4 ^a	1.8 ± 0.5 ^a
Mn	g m ⁻²	0.7 ± 0.1 ^b	1.8 ± 0.3 ^a	0.6 ± 0.2 ^b
H	g m ⁻²	0.2 ± 0.0 ^a	0.2 ± 0.0 ^a	0.2 ± 0.1 ^a

1079

1080 **Table 2** Annual water balance simulated from March 2017 to February 2018 for each
 1081 management zone.

Water flux (mm yr ⁻¹)	Palm circle	FronD-stacked area	Inter-row
Precipitation	2772	2772	2772
Transpiration	828	448	401
Evaporation	228	214	434
Interception	351	209	209
Runoff	338	0	216
Drainage (at 1.5 m depth)	556	1806	1179

1082

1083 **Table 3** Annual leaching losses at 1.5 m depth for each experimental treatment from March
1084 2017 to February 2018. Values are area-weighted averages of leaching losses in each
1085 management zone (mean \pm standard error, $n = 4$ plots). Means followed by different letters
1086 indicate differences among experimental treatments (linear-mixed effect models on monthly
1087 values followed by Tukey HSD test for multiple comparisons at $P \leq 0.05$). Treatments: ch =
1088 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
1089 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding. DON =
1090 dissolved organic N; TDN = total dissolved N.

Element leaching (kg ha ⁻¹ yr ⁻¹)	ch	cw	rh	rw
NO ₃ ⁻ -N	71.5 \pm 20.1 ^a	48.2 \pm 13.0 ^{ab}	36.3 \pm 20.1 ^b	30.0 \pm 5.7 ^b
NH ₄ ⁺ -N	1.7 \pm 0.2 ^a	1.7 \pm 0.1 ^a	1.8 \pm 0.1 ^a	1.7 \pm 0.2 ^a
DON	0.5 \pm 0.5 ^a	0.6 \pm 0.3 ^a	0.4 \pm 0.1 ^a	0.3 \pm 0.0 ^a
TDN	73.6 \pm 20.2 ^a	50.4 \pm 13.1 ^{ab}	38.4 \pm 8.9 ^b	32.0 \pm 5.8 ^b
Ca	26.6 \pm 4.3 ^a	19.4 \pm 4.4 ^b	18.2 \pm 1.8 ^b	17.0 \pm 2.1 ^b
Mg	11.6 \pm 2.5 ^a	7.7 \pm 0.8 ^b	9.1 \pm 0.7 ^{ab}	10.8 \pm 3.6 ^{ab}
K	8.1 \pm 1.3 ^a	6.2 \pm 0.7 ^b	8.9 \pm 0.6 ^a	5.7 \pm 1.1 ^b
Na	15.9 \pm 3.5 ^{ab}	13.6 \pm 2.4 ^b	18.9 \pm 3.1 ^a	13.1 \pm 1.2 ^b
Mn	0.3 \pm 0.1 ^a	0.2 \pm 0.0 ^b	0.2 \pm 0.0 ^{bc}	0.1 \pm 0.0 ^c
Total Al	40.8 \pm 11.5 ^a	20.8 \pm 7.6 ^b	19.9 \pm 6.8 ^b	21.8 \pm 3.1 ^b
Total S	2.4 \pm 0.5 ^a	1.8 \pm 0.4 ^a	2.1 \pm 0.6 ^a	4.9 \pm 3.3 ^a
Total Fe	0.2 \pm 0.0 ^a	0.5 \pm 0.3 ^a	0.2 \pm 0.0 ^a	0.5 \pm 0.3 ^a
Total P	0.0 \pm 0.0 ^a	0.1 \pm 0.0 ^a	0.0 \pm 0.0 ^a	0.0 \pm 0.0 ^a

CI 79.7 ± 15.8^a 36.9 ± 8.3^b 67.7 ± 8.7^a 78.3 ± 7.5^a

1091

1092


1093 **Table 4** N and base cation retention efficiencies in the soil for each management zone and
 1094 experimental treatment (means \pm standard error, $n = 4$ plots). Means followed by different
 1095 lowercase letters indicate differences among experimental treatments for each management
 1096 zone, whereas different uppercase letters indicate differences among management zones for
 1097 each experimental treatment (one-way ANOVA with Tukey HSD or Kruskal–Wallis H test
 1098 with multiple comparisons extension at $P \leq 0.05$). Weighted-average is based on the areal
 1099 coverage of each management zone: 18% for palm circle, 15% for frond-stacked area, and 67%
 1100 for inter-row. Treatments: ch = conventional fertilization–herbicide; cw = conventional
 1101 fertilization–mechanical weeding; rh = reduced fertilization–herbicide; rw = reduced
 1102 fertilization–mechanical weeding. See section 2.4 for calculations of N and base cation
 1103 retention efficiency.

	ch	cw	rh	rw
N retention efficiency ($\text{mg N m}^{-2} \text{d}^{-1} / \text{mg N m}^{-2} \text{d}^{-1}$)				
Palm circle	0.987 ± 0.002^{aA}	0.982 ± 0.007^{aAB}	0.986 ± 0.003^{aAB}	0.997 ± 0.000^{aA}
Frond-stacked area	0.984 ± 0.004^{aA}	0.989 ± 0.004^{aA}	0.993 ± 0.001^{aA}	0.987 ± 0.002^{aA}
Inter-row	0.877 ± 0.025^{aB}	0.870 ± 0.022^{aB}	0.900 ± 0.018^{aB}	0.906 ± 0.039^{aA}
Weighted-average	0.925 ± 0.022^a	0.934 ± 0.020^a	0.945 ± 0.012^a	0.946 ± 0.018^a
Base cation retention efficiency ($\text{mol}_c \text{m}^{-2} \text{yr}^{-1} / \text{mol}_c \text{m}^{-2} \text{yr}^{-1}$)				

Palm circle	$0.967 \pm 0.008^{ab A}$	$0.982 \pm 0.002^{a A}$	$0.937 \pm 0.013^{b A}$	$0.974 \pm 0.010^{ab A}$
FronD-stacked area	$0.884 \pm 0.013^{b A}$	$0.950 \pm 0.004^{a A}$	$0.960 \pm 0.002^{a A}$	$0.928 \pm 0.016^{ab A}$
Inter-row	$0.588 \pm 0.086^{b B}$	$0.875 \pm 0.022^{a B}$	$0.704 \pm 0.048^{ab B}$	$0.822 \pm 0.063^{ab A}$
Weighted-average	0.876 ± 0.009^b	0.945 ± 0.007^a	0.902 ± 0.019^{ab}	0.934 ± 0.012^{ab}


1104

1105 **Figure 1** Pearson correlation test between modeled (red line) and field-measured soil water
 1106 matric potential (black points) ($n = 50$ field measurements over one year) for each management
 1107 zone at 30 cm and 60 cm depths.
 1108




1109

1110 **Figure 2** Monthly water drainage at 1.5 m depth, simulated in each management zone, and
1111 daily rainfall from March 2017 to February 2018. The gray shaded area represent the dry season
1112 (precipitation < 140 mm month⁻¹)




1113

1114 **Figure 3.** Partial cation-anion charge balance of the major solutes (with concentrations > 0.03
 1115 mg L⁻¹) in soil water at 1.5 m depth for each experimental treatment in the different
 1116 management zones. The combined concentrations of organic acids (RCOO⁻) and carbonates
 1117 (HCO₃⁻) are calculated as the difference between the measured cations and anions. Treatments:
 1118 ch = conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding;
 1119 rh = reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding.




1120
 1121 **Figure 4** Monthly leaching losses at 1.5 m depth (mean \pm standard errors, $n = 4$ plots) for each
 1122 management zone. Black arrows indicate fertilizer applications and the gray shaded area
 1123 represents the dry season (precipitation < 140 mm month⁻¹).

1124



1125

1126 **Figure 5** Average monthly leaching losses at 1.5 m depth for each experimental treatment from
 1127 March 2017 to February 2018. Values are area-weighted averages of leaching losses in each
 1128 management zone (means \pm standard errors, $n = 4$ plots). For each parameter, different letters
 1129 indicate significant differences among treatments (linear-mixed effect models on monthly
 1130 values followed by Tukey HSD test for multiple comparisons at $P \leq 0.05$). Treatments: ch =
 1131 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
 1132 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding
 1133



1134

1135 **Appendices**

1136 **Table A1** Parameters used ~~for~~ in the Expert-N water sub-model for each management zone.

<u>Parameters</u>	Depth (cm)	Palm circle	Inter-row	Fron- stacked area
Interception				
Saturation capacity (mm d ⁻¹)		8.4	4.7	4.7
Throughfall (%)		50	10	10
Plant water uptake				
Plant height (cm)		874	874	874
Leaf area index		3.64	1.8	0.75
Leaf number		40	40	40
Aboveground biomass (kg ha ⁻¹)		47400	47400	47400
Maximum rooting depth (cm)		100	50	50
Crop cover		0.8	0.6	0.6
Root biomass (kg ha ⁻¹)		15600	15600	15600
Root partition (%)	0–10	29	29	29
	10–30	31	31	31
	30–50	18	18	18
	50–100	15	15	15
	100–150	5	5	5
	150–200	2	2	2
Soil properties				
Bulk density (g cm ⁻³)	0–10	1.37	1.36	0.8
	10–30	1.36	1.36	1.26
	30–50	1.52	1.52	1.52
	50–100	1.50	1.50	1.50
	100–150	1.58	1.58	1.58
	150–200	1.46	1.46	1.46
Texture – Clay (%)	0–10	15.8	15.8	15.8
	10–30	24.5	24.5	24.5
	30–50	37.5	37.5	37.5
	50–100	41.0	41.0	41.0
	100–150	43.3	43.3	43.3
	150–200	47.6	47.6	47.6
Texture – Sand (%)	0–10	53.3	53.3	53.3
	10–30	47.6	47.6	47.6
	30–50	35.9	35.9	35.9
	50–100	34.4	34.4	34.4
	100–150	31.7	31.7	31.7

Organic matter (%)	150–200	29.8	29.8	29.8
	0–10	3.2	2.9	8.7
	10–30	2.8	2.6	3.7
	30–50	2.0	1.6	2.0
	50–100	2.5	2.5	2.5
	100–150	2.0	2.0	2.0
Porosity (Vol %)	150–200	1.2	1.2	1.2
	0–10	48.8	48.8	70.0
	10–30	45.7	45.7	45.7
	30–50	41.9	41.9	41.9
	50–100	43.3	43.3	43.3
	100–150	40.3	40.3	40.3
Field capacity (Vol %)	150–200	45.0	45.0	45.0
	0–10	27.2	27.2	27.2
	10–30	27.4	27.4	27.4
	30–50	21.3	21.3	21.3
	50–100	23.1	23.1	23.1
	100–150	24.5	24.5	24.5
Wilting point (Vol %)	150–200	28.1	28.1	28.1
	0–10	18.3	18.3	18.3
	10–30	17.3	17.3	17.3
	30–50	17.9	17.9	17.9
	50–100	17.3	17.3	17.3
	100–150	20.4	20.4	20.4
Saturated hydraulic conductivity (mm d ⁻¹)	150–200	24.5	24.5	24.5
	0–10	400	400	200
	10–30	200	200	400
	30–50	200	200	300
	50–100	150	150	150
	100–150	260	260	260
Van Genuchten α (cm ⁻¹)	150–200	260	260	260
	0–10	0.059	0.059	0.059
	10–30	0.025	0.025	0.035
	30–50	0.010	0.010	0.020
	50–100	0.008	0.008	0.015
	100–150	0.021	0.021	0.021
Van Genuchten n	150–200	0.021	0.021	0.021
	0–10	1.70	1.70	1.70
	10–30	1.71	1.71	1.81
	30–50	1.12	1.12	1.25
	50–100	1.09	1.09	1.15
	100–150	1.21	1.21	1.21
	150–200	1.23	1.23	1.23

1137 **Table A2** Gross N mineralization rates (means \pm SE, $n = 4$ plots) in the top 5 cm soil for each
 1138 treatment and management zone in a large-scale plantation in Jambi, Indonesia. Measurements
 1139 were done on intact soil cores in February 2018 using the ^{15}N pool dilution technique, as
 1140 described in details by Allen et al. (2015). Treatments: ch = conventional fertilization–
 1141 herbicide; cw = conventional fertilization–mechanical weeding; rh = reduced fertilization–
 1142 herbicide; rw = reduced fertilization–mechanical weeding

Gross N mineralization ($\text{mg N m}^{-2} \text{d}^{-1}$)

	ch	cw	rh	rw
palm circle	135 \pm 39	115 \pm 25	111 \pm 34	210 \pm 13
frond-stacked area	584 \pm 100	845 \pm 207	581 \pm 188	430 \pm 134
inter-row	288 \pm 64	239 \pm 39	227 \pm 51	262 \pm 56

1143 *Note:* These data are not included in the main manuscript to avoid redundant publication as they
 1144 were already included in another manuscript presently in review.


1145 **Table A3** Literature comparison of annual N fertilization and total N leaching losses across
 1146 tropical plantations.

Author	Soil type	rainfall (mm yr ⁻¹)	Type of plantation management	N applied (kg ha ⁻¹ yr ⁻¹)	Total N leaching (kg ha ⁻¹ yr ⁻¹)	Percentage N leached (%)
Present study	loam Acrisol	2772	intensive oil palm	260	74	28
Present study	loam Acrisol	2772	intensive oil palm	130	38	28
Omoti et al. 1983	sandy clay Acrisol	2000	intensive oil palm	150	9	6
Kurniawan et al. 2018	loam Acrisol	3418	smallholder oil palm	88	11	12.5
Tung et al. 2009	Acrisol	-	intensive oil palm	128	3 (150 days)	2
Tung et al. 2009	Acrisol	-	intensive oil palm	251	3 (150 days)	1
Banabas et al. 2008	clay loam Andosol	2398	intensive oil palm	100	37	37
Banabas et al. 2008	sandy loam Andosol	3657	intensive oil palm	100	103	103
Cannavo et al. 2013	clay loam Andosol	2678	coffee agroforestry	250	157	63


Tully et al., 2012	clay loam	2700	coffee	120	119	99
	Andosol		agroforestry			
Armour et al. 2013	clay Acrisol	1958	intensive	476	164	34
			banana			
Wakelin et al. 2011	loam	2685	intensive	305	116	38
	Acrisol		banana			

1147
1148

1149 **Figure A1** Lysimeter locations at each treatment plot, with two subplots (blue rectangles) that
1150 each included the three management zones (blue crosses): 1) lysimeters in the palm circle were
1151 at 1 m from the palm stem, 2) in the frond-stacked area, at about 4 m from the palm stem, and
1152 3) in the inter-row, at approximately 4 m from the palm stem.



1153
1154 **Figure A2** Annual yield of each experimental treatment from 2017 to 2019. Treatments: ch =
1155 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
1156 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding.



1157

1158 *Note:* yield was measured by weighing the harvested fresh fruit bunches from each palm in
 1159 the inner 30 m x 30 m area of each plot.

1160