



1 **Herbicide weed control increases nutrient leaching as compared to**
2 **mechanical weeding in a large-scale oil palm plantation**

3

4 Greta Formaglio¹, Edzo Veldkamp¹, Xiaohong Duan², Aiyen Tjoa³, Marife D. Corre¹

5

6 ¹Soil Science of Tropical and Subtropical Ecosystems, University of Göttingen, Göttingen,

7 37073, Germany

8 ²Institute of Biochemical Plant Pathology, Helmholtz Zentrum Munich, 85764, Germany

9 ³Faculty of Agriculture, Tadulako University, Palu, 94118, Indonesia

10

11 *Correspondence to:* Greta Formaglio (gformag@gwdg.de)

12



13 **Abstract**

14 Nutrient leaching in intensively managed oil palm plantations can diminish soil fertility and
15 water quality. There is a need to reduce this environmental footprint without sacrificing yield.
16 We quantified nutrient leaching in a large-scale oil palm plantation on Acrisol soil with factorial
17 treatment combinations of two fertilization rates (260 N, 50 P, 220 K kg ha⁻¹ yr⁻¹ as conventional
18 practice, and 136 N, 17 P, 187 K kg ha⁻¹ yr⁻¹, equal to harvest export, as reduced management)
19 and two weeding methods (conventional herbicide, and mechanical weeding as reduced
20 management). Each of the four treatment combinations was represented by a 2500 m² plot,
21 replicated in four blocks. In each plot, soil-pore water was collected monthly at 1.5 m depth for
22 one year in three management zones: palm circle, inter-row, and frond-stacked area. In the palm
23 circle, nutrient leaching was low due to low solute concentrations and small drainage fluxes,
24 resulting from large plant uptake. Conversely, in the inter-row, nitrate and aluminum leaching
25 losses were high due to their high concentrations, large drainage fluxes, low plant uptake, and
26 acidic pH. In the frond-stacked area, base cation leaching was high, presumably from frond
27 litter decomposition, but N leaching was low. Mechanical weeding, even with conventional
28 high fertilization rates, reduced leaching losses of all nutrients. Mechanical weeding with
29 reduced fertilization had the lowest N and base cation leaching whereas its yield and economic
30 gross margin remain comparable with the conventional management practices. Herbicide weed
31 control decreased ground vegetation, and thereby reduced efficiency of soil nutrient retention.
32 Our findings signified that mechanical weeding and reduced fertilization should be included in
33 the Indonesian Ministry of Agriculture program for precision farming (e.g. variable rates with
34 plantation age), particularly for large-scale plantations, and in the science-based policy
35 recommendations, such as those endorsed by the Roundtable for Sustainable Palm Oil
36 association.



37 **1 Introduction**

38 Agricultural expansion is a major driver of tropical deforestation (Geist and Lambin, 2002),
39 which have global impacts on reducing carbon sequestration (Asner et al., 2010; van Straaten
40 et al., 2015), greenhouse gas regulation (e.g. Meijide et al., 2020; Murdiyarso et al., 2010), and
41 biodiversity (e.g Clough et al., 2016) and increasing profit gains at the expense of ecosystem
42 multifunctionality (Grass et al., 2020). Oil palm is the most important rapidly expanding tree-
43 cash crop that replaces tropical forest in Southeast Asia (Gibbs et al., 2010; Carlson et al., 2013)
44 due to its high yield with low production costs and rising global demand (Carter et al., 2007;
45 Corley, 2009). Currently, Indonesia produces 57 % of palm oil worldwide (FAO, 2018) and
46 this production is projected to expand in the future, threatening the remaining tropical forest
47 (Vijay et al., 2016; Pirker et al., 2016). Forest to oil palm conversion is associated with a
48 decrease in soil fertility, because of high nutrient export via harvest, reduced rates of soil-N
49 cycling, and decreases in soil organic carbon (SOC) and nutrient stocks (Allen et al., 2015;
50 Allen et al., 2016; van Straaten et al., 2015). The decline in soil fertility reinforces the
51 dependency on fertilizer inputs, and a severe decline can lead to abandonment of the area with
52 further expansion of oil palm plantations in another, exacerbating land-use change. Leaching
53 can contribute to the impoverishment of soil nutrients as well as reduction in water quality and
54 eutrophication of water bodies. Increased nutrient loads to water bodies due to agricultural
55 expansion and intensification, common in temperate areas (Carpenter et al., 1998), are
56 increasingly reported for tropical regions (Figueiredo et al., 2010; Teklu et al., 2018). Given the
57 typically high precipitation rates, leaching losses can possibly be large in intensively managed
58 plantations in the tropics, although deeply weathered tropical soils also have the capacity to
59 store large quantities of N and P (Jankowski et al., 2018; Neill et al., 2013). Indeed, NO_3^- , the
60 most leachable form of N, can be retained in the subsoil by anion exchange capacity of highly
61 weathered acidic soils (Wong et al., 1990) whereas P can be fixed to Fe and Al (hydr)oxides of
62 tropical soils (Roy et al., 2016). Nevertheless, there are some evidences of streamwater quality



63 reductions due to oil palm cultivation in Malaysia (Luke et al., 2017; Tokuchi et al., 2019),
64 signifying the importance of quantifying nutrient leaching losses in other areas with expansive
65 oil palm plantations, especially in Jambi, Indonesia, one of the hotspots of forest conversion to
66 oil palm in Indonesia (Drescher et al., 2016).

67 Although oil palm plantations can possibly have low leaching losses, as a consequence
68 of high evapotranspiration and thus low drainage fluxes (Tarigan et al., 2020), most of oil palm
69 plantations are large-scale enterprises that are characterized by intensive management with high
70 fertilization rates and herbicide application. Intensive agriculture in the tropics is associated
71 with high N leaching losses (Huddell et al., 2020). Even in tree-cash or perennial crop
72 plantations, despite their generally higher evapotranspiration and deeper rooting depth than
73 annual crops, high fertilization rates result in sustained, large nutrient leaching losses (e.g.
74 Cannavo et al., 2013; Wakelin et al., 2011). Large NO_3^- leaching from high N fertilization is
75 always accompanied by leaching of cations (Cusack et al., 2009; Dubos et al., 2017),
76 impoverishing highly weathered tropical soils that are inherently low in base cations (Allen et
77 al., 2016; Kurniawan et al., 2018). Fertilization is necessary to support high yields of oil palm
78 plantations, but reduction in fertilization rates, e.g. to levels that compensate for nutrient export
79 through harvest, may reduce nutrient leaching losses while maintaining high productivity. On
80 the other hand, the use of herbicide for weed control can exacerbate nutrient leaching losses, as
81 prolonged absence of ground vegetation reduces uptake of redistributed nutrients from applied
82 fertilizers far from reach of crop roots (Abdalla et al., 2019). Herbicide weeding, common in
83 large-scale oil palm plantation, is practiced in the area where the fertilizers are applied, to reduce
84 competition for nutrients and water with ground vegetation, and in the inter-rows, to facilitate
85 access during harvest (Corley and Tinker, 2016). However, herbicide not only eradicates
86 aboveground vegetative parts but also removes roots slowing down regeneration. In contrast,



87 mechanical weeding only removes aboveground part, allowing relatively fast regeneration of
88 ground vegetation, which could take up redistributed nutrients and could reduce leaching losses.

89 To investigate nutrient leaching losses in an oil palm plantation, the spatial structure
90 created by the planting design and by the management practices must be taken into account,
91 which is only partly considered in the sampling designs of previous studies. Three management
92 zones in oil palm plantations can be identified: (1) the palm circle, an area around the palm's
93 trunk where the fertilizers are applied and weeded; (2) the inter-row, weeded less frequently
94 than the palm circle but unfertilized; and (3) the frond-stacked area, usually every second inter-
95 row, where the cut senesced fronds are piled up. In these management zones, the interplay of
96 water fluxes, root uptake and soil nutrient contents determine the extent of nutrient leaching
97 losses. The palm circle despite having direct fertilization have also large water and nutrient
98 uptake (Nelson et al., 2006) because of high root density (Lamade et al., 1996) such that large
99 leaching losses may only occur following pulse high fertilization and during high drainage
100 (from high precipitation) events (Banabas et al., 2008a). The inter-row experiences higher water
101 input from precipitation than the palm circle because of lower canopy interception (Banabas et
102 al., 2008b), and large water flux within the soil because of low root uptake, stimulating nutrient
103 transport to lower depths. However, as there is no direct fertilizer application on the inter-row,
104 nutrient leaching may be low. The frond-stacked area receives nutrients from decomposition of
105 nutrient-rich fronds (Kotowska et al., 2016) and such mulching with senesced fronds prevents
106 runoff and promotes water infiltration as a consequence of enhanced macroporosity by
107 increased organic matter (Moradi et al., 2015). High water infiltration may generate high water
108 drainage fluxes, resulting in intermediate nutrient leaching losses in the frond-stacked area.

109 In this study, we aimed to quantify nutrient leaching losses in an intensively managed,
110 large-scale oil palm plantation, and to assess if reduced intensity of management (i.e. reduced
111 fertilization rates equal to harvest export and mechanical weeding) can reduce leaching losses



112 in oil palm plantations. We tested these hypotheses: (1) leaching losses in the palm circle will
113 be larger than in the other management zones because of direct fertilizer application; (2)
114 leaching losses under herbicide application will be higher than mechanical weeding because of
115 slower regeneration of ground vegetation that can augment nutrient retention; (3) nutrient
116 leaching fluxes under conventional high fertilization rates will be substantial compared to
117 reduced rates because of excessive nutrient inputs. Our study provides a systematic
118 quantification of an important environmental footprint of oil palm production, taking into
119 consideration its spatial variation in management zones, and evaluates the effectiveness of
120 alternative management practices for leaching reduction.

121 **2 Materials and methods**

122 **2.1 Study area and experimental design**

123 This study was conducted in a state-owned oil palm plantation in Jambi province, Indonesia (1°
124 43' 8" S, 103° 23' 53" E, 73 m above sea level). Mean annual air temperature is 26.7 ± 1.0 °C
125 and mean annual precipitation is 2235 ± 385 mm (1991–2011; data from Sultan Thaha airport,
126 Jambi). During our study period (March 2017–February 2018), the mean daily air temperature
127 was 26.3 °C and annual precipitation was 2772 mm, with a dry period between July and October
128 (precipitation < 140 mm month⁻¹). The soil is highly weathered, loam Acrisol soil (Allen et al.,
129 2015) and nutrient inputs from bulk precipitation in the area, measured in 2013, were 12.9 kg
130 N, 0.4 kg P, 5.5 kg K ha⁻¹ yr⁻¹ (Kurniawan et al., 2018).

131 This oil palm plantation was established between 1998 and 2002, and so the palms were
132 16–20 years old during our study period. The plantation encompassed 2025 ha, with a planting
133 density of approximately 142 palms ha⁻¹, spaced 8 m apart on rows. The rows between palms
134 are used alternately for harvesting operations and to pile-up senesced fronds, which are
135 regularly cut to facilitate harvesting of fruits; this frond-stacked area covers 15 % of the



136 plantation. The palm circle, 2 m radius from the trunk, wherein fertilizers are applied and
137 weeded four times a year, covers 18 % of the plantation. The remaining 67 % can be classified
138 as inter-row, which is not fertilized but weeded two times a year.

139 In November 2016, a two (fertilization rates) by two (weeding methods) factorial
140 management experiment was established in this plantation as part of the framework of the
141 EFForTS project, described in detail by Darras et al. (2019). For fertilization treatments, the
142 conventional rates were 260 N, 50 P, 220 K kg ha⁻¹ yr⁻¹, whereas the reduced rates were 136 N,
143 17 P, 187 K kg ha⁻¹ yr⁻¹. Reduced fertilization rates were determined to compensate for nutrient
144 exports via fruit harvest and were based on the nutrient concentrations measured in the fruit
145 bunches multiplied by the annual yield. The fertilizer sources were urea (CH₄N₂O), triple
146 superphosphate (Ca(H₂PO₄)₂·H₂O) and muriate of potash (KCl), and these were applied
147 according to the plantation's standard practices: split in two applications per year (in April and
148 October), applied in a band within a 2 m radius from the palm, and this area was raked before
149 fertilizer application. For both fertilization treatments, lime (426 kg dolomite ha⁻¹ yr⁻¹;
150 CaMg(CO₃)₂) and micronutrients (142 kg Micro-Mag ha⁻¹ yr⁻¹ with 0.5 % B₂O₃, 0.5 % CuO,
151 0.25 % Fe₂O₃, 0.15 % ZnO, 0.1 % MnO and 18 % MgO) were also applied besides the N, P
152 and K fertilizers, as commonly practiced in large-scale plantations on acidic Acrisol soils
153 (Pahan, 2010). For weeding treatments, the conventional method was the use of herbicide
154 (glyphosate), whereas the reduced method was mechanical weeding using a brush cutter.
155 Glyphosate was applied following plantation's standard practice: 1.5 L ha⁻¹ yr⁻¹ to the palm
156 circle, split four times a year, and 0.75 L ha⁻¹ yr⁻¹ to the inter-row, split two times a year. The
157 mechanical weeding was carried out in the same areas and frequencies as herbicide application.
158 This management experiment comprised of four replicate blocks and each had four plots (50 m
159 x 50 m each) assigned to four treatment combinations: conventional rate–herbicide,



160 conventional rate–mechanical weeding, reduced rate–herbicide, and reduced rate–mechanical
161 weeding.

162

163 **2.2 Soil water sampling**

164 We collected monthly soil-pore water samples over one year, using suction cup lysimeters (P80
165 ceramic, maximum pore size 1 μm ; CeramTec AG, Marktredwitz, Germany). We installed the
166 lysimeters in January 2017, choosing two palms per plot and sampling in the three management
167 zones: 1) in the palm circle, at 1 m from the palm trunk, 2) in the frond-stacked area, at about
168 4 m from the palm trunk, and 3) in the inter-row, at approximately 4 m from the palm trunk
169 (Fig. A1). In total, 96 lysimeters were installed (4 treatment plots x 4 replicates x 2 subplots x
170 3 management zones). The lysimeters were inserted into the soil till 1.5 m depth, so that the
171 soil-pore water was collected well below the rooting depth of 1 m which is common to oil palm
172 plantations on loam Acrisol soils near our study site (Kurniawan et al., 2018). Starting in March
173 2017, soil water was sampled by applying 40 kPa vacuum (Kurniawan et al., 2018; Dechert et
174 al., 2005) to the lysimeters and collected in dark glass bottles, which were stored in a bucket
175 buried in the field. Once a week, we transferred the collected water into plastic bottles and
176 transported them to the field station, where they were stored frozen. The collection continued
177 over a month until a volume of 100 mL was collected from each lysimeter, or until the end of
178 the month. The frozen water samples were transported by air freight to the University of
179 Goettingen, Germany, where element concentrations were determined. We measured the
180 concentrations of mineral N (NH_4^+ and NO_3^-), total dissolved N (TDN) and Cl by continuous
181 flow injection colorimetry (SEAL Analytical AA3, SEAL Analytical GmbH, Norderstadt,
182 Germany), as described in details by Kurniawan et al. (Kurniawan et al., 2018). Dissolved
183 organic N (DON) was calculated as the difference between TDN and mineral N. We measured
184 the concentrations of base cations (Na, K, Ca, Mg), total Al, total Fe, total Mn, total S, and total



185 P with an inductively coupled plasma–atomic emission spectrometer (iCAP 6300; Thermo
186 Fischer Scientific GmbH, Dreieich, Germany).

187 We determined a partial cation-anion charge balance of the major elements
188 (concentrations $> 0.03 \text{ mg L}^{-1}$) in soil-pore water by converting the concentrations to $\mu\text{mol}_{\text{charge}}$
189 L^{-1} . We assumed S to be in the form of sulfate (SO_4^{2-}) and total Al to have a charge of 3^+ . We
190 calculated the contribution of organic acids (RCOO^-) and bicarbonate (HCO_3^-) as the difference
191 between the measured cations and anions (2018).

192

193 **2.3 Modeling water drainage**

194 The water balance was modeled using the water sub-model of the Expert-N software, version
195 5.0 (Priesack, 2005), which was successfully used to estimate drainage fluxes from different
196 land uses in Indonesia (Dechert et al., 2005; Kurniawan et al., 2018). The model inputs were
197 climate data (solar radiation, temperature, precipitation, relative humidity, and wind speed), and
198 soil (texture, bulk density, and hydraulic functions) and vegetation characteristics (biomass,
199 leaf area index, and root distribution). The climate data were taken from the climatological
200 station in the plantation (described in detail by Mejjide et al., 2017), and the oil palm biomass
201 was taken from a study on oil palm plantations near our study site (Kotowska et al., 2015). Soil
202 bulk density and porosity in the top 10 cm were measured in each management zone at our
203 study site, whereas for the 10–50 cm depth these were measured in the inter-row, assuming that
204 the differences in soil bulk density among management zones would be minimal below the
205 topsoil. Data for soil bulk density and porosity for the 50–200 cm depth, as well as soil texture,
206 soil hydraulic parameters (i.e. water retention curve, saturated hydraulic conductivity and Van
207 Genuchten parameters for the water retention curve), and root distribution were taken from
208 Allen et al. (2015) and Kurniawan et al. (2018), choosing their studied oil palm plantations



209 closest to our study site. Expert-N water sub-model calculates daily water drainage based on
210 precipitation, evapotranspiration, canopy interception, runoff, and change in soil water storage.
211 Evapotranspiration is calculated using Penman-Monteith method (Allen, 1998), applying a
212 plant factor of 1.06 (Meijide et al., 2017), with plant transpiration based on leaf area index
213 (LAI), plant biomass, and maximum rooting depth. The canopy interception is calculated from
214 the percentage of throughfall and the maximum water storage capacity of the canopy. Runoff
215 is calculated from soil texture and bulk density, which determine the water infiltration rate, and
216 from the slope, which was 5 % (Röll et al., 2019). The vertical water movement is calculated
217 using Richard's equation based on soil hydraulic functions.

218 To model the drainage in the different management zones, we used the measured soil
219 bulk density and porosity in the top 10 cm and adjusted other input parameters to simulate
220 differences in water balance in each management zone. For the palm circle, we set the LAI to
221 3.65, which is the maximum LAI measured at our site (Fan et al., 2015), to simulate high water
222 uptake in the palm circle (Nelson et al., 2006) and maximum rooting depth to 1 m, which is
223 reported for oil palm plantations near our site (Kurniawan et al., 2018). The percentage
224 throughfall in the palm circle was set to 50 % and the water storage capacity of oil palm trunk
225 was set to 8.4 mm (Tarigan et al., 2018). For the inter-row, we set the LAI and the maximum
226 rooting depth as half of the palm circle (1.8 LAI, 50 cm rooting depth), as roots are shallower
227 between palms (Nelson et al., 2006); the throughfall was set to 10 %, and the palm trunk's water
228 storage capacity was set to 4.7 mm (based on canopy storage capacity reported by Tarigan et
229 al., 2018). For the frond-stacked area, the LAI was set to 0.75, which is half of the minimum
230 measured in the studied plantation (Darras et al., 2019), as understory vegetation is absent at
231 this zone. Values for interception in the frond-stacked area was set to the same values as the
232 inter-row, whereas the runoff was set to 0, as mulching with senesced fronds slows down runoff
233 (Tarigan et al., 2016).



234 For validation of the Expert-N water sub-model outputs, we measured soil water matric
235 potential at depths of 30 cm and 60 cm over the study period and compared the measured values
236 with the modeled matric potential. Matric potential was measured by installing a tensiometer
237 (P80 ceramic, maximum pore size 1 μm ; CeramTec AG, Marktrechwitz, Germany) at each depth
238 in each management zone near to two palms in two treatments (i.e. conventional rate–herbicide,
239 and reduced rate–mechanical weeding), for a total of 12 tensiometers. We summed the modeled
240 daily drainage at 1.5 m depth to get the monthly drainage fluxes, which we then multiplied with
241 the element concentrations in soil water to get the monthly nutrient leaching fluxes.

242

243 **2.4 Soil biochemical characteristics and nutrient retention efficiency**

244 We measured soil biochemical properties in the same sampling locations (Figure S1) at four
245 depth intervals: 0–5 cm, 5–10 cm, 10–30 cm, and 30–50 cm. Soil samples from the same
246 management zone in each plot were pooled to make one composite sample, totaling to 192 soil
247 samples (4 treatments plots x 4 replicates x 3 management zones x 4 depths). The samples were
248 air-dried and sieved (2 mm) and measured for pH (1:4 soil-to-water ratio) and for effective
249 cation exchange capacity (ECEC), by percolating the soils with unbuffered 1 mol L⁻¹ NH₄Cl
250 and measuring the cations (Ca, Mg, K, Na, Al, Fe, Mn) in percolates using ICP-AES. A
251 subsample was finely ground and analyzed for organic C and total N using a CN analyzer (Vario
252 EL Cube, Elementar Analysis Systems GmbH, Hanau, Germany), and for ¹⁵N natural
253 abundance signature using isotope ratio mass spectrometer (IRMS; Delta Plus, Finnigan MAT,
254 Bremen, Germany). We calculated the soil element stocks for each depth by multiplying the
255 element concentration with the measured bulk density and summed for the top 50 cm; other soil
256 characteristics (e.g. pH, ECEC, base saturation) in the top 50 cm soil were calculated as the
257 depth-weighted average of the sampled depths.



258 In addition, we calculated the N and base cation retention efficiency in the soil for each
259 experimental treatment and management zone following the formula: nutrient retention
260 efficiency = $1 - (\text{nutrient leaching loss} / \text{soil-available nutrient})$ (Kurniawan et al., 2018). We
261 used the gross N mineralization rates in the top 5 cm soil (Table A1) as an index of soil-available
262 N whereas soil-available base cations was the sum of the stocks of K, Na, Mg and Ca in the top
263 10 cm soil, expressed in $\text{mol}_{\text{charge}} \text{m}^{-2}$.

264

265 **2.5 Statistical analyses**

266 For soil biochemical properties measured once, we tested for differences among management
267 zones as well as among experimental treatments for the entire 50 cm depth, using the analysis
268 of variances (ANOVA) with Tukey HSD as a post hoc test. The soil variables that showed non-
269 normal distribution or unequal variances, tested with Shapiro–Wilk and Levene’s tests,
270 respectively, were log-transformed prior to the analysis. Base cation and N retention efficiency
271 were also tested for differences between experimental treatments in the same way. For
272 repeatedly measured variables, i.e. soil-pore water solute concentrations and leaching fluxes,
273 we used linear mixed-effects models (LME; Bates et al., 2015) to assess the differences among
274 management zones and treatments. For testing management zone differences, we conducted the
275 LME with management zone as fixed effect and random effects for sampling months and
276 experimental treatments nested with replicate plots, which were also nested with subplots. For
277 testing treatment differences, we calculated for each replicate plot on each sampling month the
278 area-weighted average of the three management zones (i.e. palm circle accounts for 18 % of the
279 plantation area, the frond-stacked area 15 %, and the inter-row 67 %), and LME was carried
280 out with treatment as fixed effect and random effects for sampling months and replicate plots
281 nested with subplots. If the residuals of the LME models were not normally distributed, we
282 applied either logarithmic or square root transformation. Differences were assessed with



283 ANOVA (Kuznetsova et al., 2017) followed by Tukey HSD (Hothorn et al., 2008). We also
284 used LME to assess differences in soil water matric potential among management zones, with
285 management zone as fixed effect and measurement days and depth nested with treatment as
286 random effects. Comparability between modeled and measured soil water matric potential for
287 each depth in each management zone ($n = 50$ field measurements) was assessed using Pearson
288 correlation test. All tests were considered significant at $P \leq 0.05$, except for soil pH which we
289 considered a marginal significance at $P = 0.06$. All statistical analyses were performed with R
290 version 3.6.1 (R Core Team, 2019).

291

292 **3 Results**

293 **3.1 Soil biochemical properties and water balance**

294 Soil biochemical properties in the top 50 cm did not differ between experimental treatments (all
295 $P > 0.05$) but strongly differed among management zones (Table 1). The frond-stacked area,
296 where senesced fronds were regularly piled like mulch material, had higher SOC and total N
297 stocks ($P < 0.01$) compared to the other management zones. The inter-row, with regular
298 weeding but without direct fertilizer and lime inputs, showed lower exchangeable base cation
299 contents (i.e. Ca, Mg, K) compared to the other management zones ($P \leq 0.02$) and higher
300 exchangeable Al content than the palm circle ($P = 0.01$). This was reflected in the lower base
301 saturation and higher Al saturation in the inter-row compared to the other zones ($P < 0.01$).
302 Also, inter-row had the lowest ECEC ($P < 0.01$) and marginally lower pH than the palm circle
303 ($P = 0.06$). The palm circle, where fertilizers and lime were applied, had generally comparable
304 exchangeable element contents with the frond-stacked area, except for K, which was higher in
305 the palm circle ($P < 0.01$), and for Mn, which was higher in the frond-stacked area ($P < 0.01$).



306 There were high positive correlations between field-measured and modeled soil water
307 matric potential (Fig. 1). The matric potential was generally lowest in the palm circle,
308 intermediate in the inter-row, and highest in the frond-stacked area ($P < 0.01$). This pattern was
309 also reflected in the low drainage flux in the palm circle and high drainage flux in the frond-
310 stacked area (Table 2; Fig. 2). In the palm circle, the low drainage flux had resulted from high
311 plant transpiration and interception whereas the high drainage flux in the frond-stacked area
312 was due to low evapotranspiration and runoff with the senesced frond mulch (Table 2). In ratio
313 to annual precipitation, the calculated annual evapotranspiration was 51 %, 31 %, and 38 % in
314 the palm circle, frond-stacked area, and inter-row, respectively; annual drainage fluxes at 1.5
315 m depth were 20 % of precipitation in the palm circle, 65 % in the frond-stacked area, and 43
316 % in the inter-row. Seasonally, the monthly drainage fluxes had two peak periods, May and
317 November, after consecutive days of moderate rainfall, and were lowest during the end of the
318 dry season towards the start of the wet season (Fig. 2).

319

320 **3.2 Differences in leaching losses among management zones and treatments**

321 For element concentrations in soil-pore water at 1.5 m depth, treatment differences were
322 exhibited clearly in the palm circle and inter-row (Fig. 3), with the herbicide treatment showing
323 higher element concentrations than the mechanical weeding ($P \leq 0.02$). The frond-stacked area
324 had generally lower ionic charge concentrations compared to the other management zones (Fig.
325 3). The dominant cations were Al^{3+} , Ca^{2+} , Mg^{2+} , K^+ , and Na^+ across experimental treatments
326 and management zones. Among the management zones, Al^{3+} concentrations were highest in the
327 inter-row, intermediate in the palm circle, and lowest in the frond-stacked area ($P < 0.01$). The
328 concentrations of Ca^{2+} were similar in the palm circle and frond-stacked area ($P = 0.42$), and
329 these were higher than the inter-row ($P < 0.01$). The concentrations of Mg^{2+} and K^+ were higher
330 in the palm circle than in the other two management zones ($P < 0.01$). The Na^+ concentrations



331 were higher in the palm circle and inter-row than in the frond-stacked area ($P < 0.01$). As for
332 dissolved N, NH_4^+ concentrations were lowest in the frond-stacked area, followed by the palm
333 circle, and highest in the inter-row ($P = 0.01$). Across treatments, NH_4^+ was 4–18 % of TDN
334 whereas DON was 1–7 % of TDN. Thus, NO_3^- was the main form of dissolved N, and this was
335 highest in the inter-row, followed by the frond-stacked area, and lowest in the palm circle ($P <$
336 0.01). The dominant anion was Cl^- with higher concentrations in the palm circle than in the
337 other zones ($P < 0.01$).

338 Monthly leaching fluxes showed a common pattern among the major solutes (Fig. 4):
339 there were two peaks of leaching losses (May and November) that followed fertilizer
340 applications, and lower leaching losses during the dry season from July to October. Leaching
341 fluxes of NO_3^- showed similar pattern as its concentrations: higher in the inter-row, followed
342 by the frond-stacked area, and lowest in the palm circle ($P < 0.01$; Fig. 4). Total Al leaching
343 fluxes were also higher in the inter-row than the other zones ($P < 0.01$; Fig. 4). On the other
344 hand, although base cation concentrations were large in the palm circle (Fig. 3), the low
345 drainage fluxes in this zone (Fig. 2; Table 2) resulted in opposite patterns of base cation leaching
346 fluxes among management zones; Ca, K, and Mg leaching were higher in the frond-stacked area
347 than the palm circle and inter-row (all $P < 0.01$; Fig. 4). Leaching of Na was higher in both the
348 frond-stacked area and inter-row than the palm circle ($P < 0.01$; Fig. 4).

349 Reduced intensity of management clearly influenced nutrient leaching losses (Fig. 5;
350 Table 3). Specifically, mechanical weeding reduced NO_3^- and cation leaching compared to
351 herbicide weed control ($P \leq 0.03$; Fig. 5; Table 3). Leaching of NO_3^- was highest in the
352 conventional fertilization–herbicide treatment and lowest in reduced management treatments
353 ($P \leq 0.02$; Fig. 5). This was also reflected in the leaching fluxes of accompanying cations;
354 specifically, total Al and Ca leaching were higher in conventional fertilization–herbicide
355 treatment than the reduced management treatments (all $P \leq 0.02$; Fig. 5). For the other base



356 cations, mechanical weeding clearly lowered leaching losses compared to herbicide weeding,
357 in particular K and Na leaching in both fertilization rates and Mg leaching in conventional
358 fertilization (all $P \leq 0.03$; Fig. 5).

359

360 **3.3 Annual leaching losses and nutrient retention efficiency**

361 In proportion to the applied fertilizer, annual leaching losses of TDN (Table 3) were 28 % of
362 the applied N in the herbicide treatment for both conventional and reduced fertilization rates,
363 24 % in the mechanical weeding with conventional fertilization, and only 19 % in the
364 mechanical weeding with reduced fertilization. The annual leaching of K (Table 3) was 4 % of
365 the applied K fertilizer in the herbicide treatment and 3 % in the mechanical weeding for both
366 fertilization rates. In this highly weathered Acrisol soils with high capacity for P fixation by Fe
367 and Al (hydr)oxides, there was no leaching of dissolved P (Table 3).

368 Both N and base cation retention efficiencies were generally lower in the inter-row
369 compared to the other management zones ($P \leq 0.03$), except for reduced fertilization–
370 mechanical weeding where there were no differences among management zones (Table 4). The
371 area-weighted average N retention efficiency was comparable among experimental treatments
372 ($P = 0.89$) but there was a trend of increasing efficiency with decreasing management intensity
373 (Table 4). Base cation retention efficiency showed clear differences among experimental
374 treatments for each management zones: in the palm circle, it was highest in mechanical weeding
375 and lowest in the herbicide treatment ($P = 0.04$); in the frond-staked area and inter-row, it was
376 lowest in the most intensive management treatment (conventional fertilization–herbicide) and
377 highest in either mechanical weeding or reduced fertilization ($P \leq 0.05$; Table 4). The area-
378 weighted average base cation retention efficiency was also clearly influenced by weeding



379 method, being lowest in herbicide treatment and highest in mechanical weeding both with
380 conventional fertilization ($P = 0.03$; Table 4).

381

382 **4 Discussion**

383 **4.1 Water model and temporal pattern of nutrient leaching losses**

384 To our knowledge, this study is the first attempt to model drainage fluxes from the different
385 management zones of an oil palm plantation, making our comparisons with literature values
386 limited. Our modeled annual transpiration rate in the palm circle (Table 2) was remarkably
387 similar to the values estimated with the same Penman–Monteith method ($827\text{--}829\text{ mm yr}^{-1}$;
388 Mejjide et al., 2017; Röhl et al., 2019), and our average daily transpiration rate (2.3 mm d^{-1}) was
389 within the range of that measured with drone-based photogrammetry ($3 \pm 1\text{ mm d}^{-1}$;
390 Ahongshangbam et al., 2019), all in the same oil palm plantation. Also, the modeled annual
391 runoff in the palm circle and inter-row (Table 2) was within the range of runoff estimates in oil
392 palm plantations in Jambi province (10–20 % of rainfall; Tarigan et al., 2016) and in Papua
393 New Guinea (1.4–6 % of rainfall; Banabas et al., 2008b). Considering the areal proportions of
394 the three management zones, the weighted-average drainage flux (1161 mm yr^{-1}) was lower
395 than that estimated for smallholder oil palm plantations near our study site (1614 mm drainage
396 $\text{flux with } 3418\text{ mm precipitation measured in } 2013$; Kurniawan et al., 2018), although their
397 ratios to annual precipitation were comparable. Aside from the difference in precipitation
398 during our study period compared to the relatively wet year of 2013, evapotranspiration rate is
399 higher in large-scale than smallholder oil palm plantations in our study area (Röhl et al., 2019),
400 which would lead to lower drainage flux in large-scale plantation. Moreover, in the frond-
401 stacked area, enhanced porosity from organic matter that facilitates water infiltration (Moradi
402 et al., 2015), as indirectly indicated by its low soil bulk density (Table 1), combined with low



403 evapotranspiration and runoff, resulted in large drainage flux (Table 2). This suggests that piling
404 senesced fronds may amend groundwater recharge, which could moderate discharge
405 fluctuations in water catchments of oil palm converted areas (Tarigan et al., 2020). Based on
406 these comparisons with literature values and on the good agreement between modeled and
407 measured soil water matric potential (Fig. 1), we conclude that our modeled drainage fluxes
408 were reliable.

409 The temporal peaks of nutrient leaching fluxes (May and November; Fig. 4) had resulted
410 from the combined effect of high drainage flux and fertilizer application. The high drainage
411 fluxes in May and November (Fig. 2) might have stimulated the downward transport of
412 elements and decreased their residence time in the soil, and thus their adsorption onto the soil
413 exchange sites (Lohse and Matson, 2005). These high water fluxes usually dilute the element
414 concentrations in the soil-pore water; however, high concentrations were maintained because
415 of fertilizer and lime applications in the same periods, resulting in parallel peaks of drainage
416 and leaching fluxes (Figs. 2 and 4). The high NO_3^- leaching following urea-N fertilization (Fig.
417 4) suggests increased nitrification (Silver et al., 2005), fast NO_3^- transport through the soil
418 column, and reduced anion adsorption capacity, which otherwise would have delayed anion
419 leaching (Wong et al., 1990). The latter was possibly aggravated by the additional Cl^- from
420 fertilization with KCl (Fig. 3), which could saturate the soil anion exchange sites, particularly
421 at this mature plantation with already 16–20 years of high fertilization rates. Large NO_3^-
422 leaching is always accompanied by large leaching of buffering cations (Dubos et al., 2017;
423 Kurniawan et al., 2018), resulting in their similar temporal patterns (Fig. 4). These findings
424 showed that fertilization should be avoided during periods of high drainage fluxes. Generally,
425 the high drainage was a consequence of a protracted period of moderate rainfall (Fig. 2).
426 Prediction of periods of high precipitation and drainage will further be confounded by climate
427 change, which is widening the range between wet and dry seasons and increasing the



428 uncertainties in rainfall intensity and distribution (Chou et al., 2013; Feng et al., 2013).
429 Fertilization during the dry period is also not advisable given the high volatilization of applied
430 urea even in acidic soil as this is always accompanied by liming (Goh et al., 2003; Pardon et
431 al., 2016) and the low palm uptake during the dry season (Corley and Tinker, 2016). Thus,
432 reduction of fertilization rates, e.g. at compensatory level equal to harvest export, seems a viable
433 option to reduce leaching losses without sacrificing production. One other option is the use of
434 organic amendments and slow-release fertilizers, which have been shown to reduce N leaching
435 in tropical cropping systems (Nyamangara et al., 2003; Mohanty et al., 2018; Steiner et al.,
436 2008) and to improve soil fertility in oil palm plantations (Comte et al., 2013; Boafo et al.,
437 2020), as was also evident with mulching of senesced oil palm fronds (i.e. high SOC, total N,
438 ECEC and base saturation in the frond-stacked area; Table 1).

439

440 **4.2 Leaching losses in the different management zones**

441 Contrary to our first hypothesis, leaching losses were generally higher in the inter-row,
442 especially for mineral N (largely NO_3^- ; Fig. 3), compared to the other zones, whereas the palm
443 circle had the lowest leaching (Fig. 4). This strikingly large mineral N leaching losses in the
444 inter-row were surprising given that this area did not receive direct fertilizer inputs (see section
445 2.1). This result suggests that mineral N was transported from the directly fertilized palm circle
446 to the inter-row via surface and subsurface lateral flow as these two zones were just 3 m apart
447 (Fig. A1). Surface transport of mineral N was probably a minor process at our site because of
448 the low runoff (Table 2); in an oil palm plantation in Papua New Guinea, the loss of N fertilizer
449 via surface runoff is only $0.3\text{--}2.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Banabas et al., 2008b). Mineral N was probably
450 predominantly transported to the inter-row via subsurface lateral flow. Acrisol soils are
451 characterized by clay translocation from upper to lower depths that could create an impeding
452 layer conducive to lateral water flow (Elsenbeer, 2001). Indeed, the clay contents of the Acrisol



453 soils at our study area increase with depth, and soil bulk density is highest at 100–150 cm than
454 at 150–200 cm depth (Allen et al., 2016). In addition, the palm roots spreading from the palm
455 circle to the inter-row may create channels for subsurface lateral flow of dissolved ions like
456 NO_3^- (Li and Ghodrati, 1994). Higher mineral N leaching in the inter-row than palm circle was
457 also observed in Brazil and it was attributed to lower root density and higher N mineralization
458 at increasing distance from the palm's trunk (Schroth et al., 2000). Hence, a combination of
459 lower root uptake, higher N mineralization, and subsurface lateral transport (particularly for
460 NO_3^-) may all have contributed to higher mineral N leaching losses in the inter-row than the
461 palm circle. The main accompanying cation for NO_3^- leaching in the inter-rows was Al^{3+} (Figs.
462 3 and 4), as this zone's soil pH (Table 1) was within the Al-buffering range (pH 3–5; van
463 Breemen et al., 1983), having no direct lime application and thus low base saturation (Table 1).
464 Our findings showed that if leaching is measured only within the palm circle, this largely
465 underestimates mineral N and Al leaching losses.

466 The palm circle had relatively low N leaching losses (Figs. 3 and 4) despite the direct
467 application of fertilizer. This was probably due to the large root density in this zone that
468 facilitates an efficient nutrient uptake (Edy et al., 2020; Nelson et al., 2006). Hence, the
469 dominant anion in soil-pore water in the palm circle was Cl^- (Fig. 3), enhanced by the applied
470 KCl fertilizer, which was accompanied by high base cation concentrations relative to dissolved
471 Al (Fig. 3). The former was due to the applied micromag fertilizer and dolomite (section 2.1),
472 which increased pH and exchangeable bases and rendered Al in insoluble form (i.e. lower
473 exchangeable Al; Table 1; Schlesinger and Bernhardt, 2013). Despite their high concentrations,
474 the leaching fluxes of base cations in the palm circle (Fig. 4) were constrained by the low water
475 drainage flux due to high evapotranspiration (Table 2).

476 The frond-stacked area was at the same distance from the palm circle as the inter-row
477 (Fig. A1) but had substantially lower mineral N leaching losses (Figs. 3 and 4). Decomposition



478 of nutrient-rich fronds (Kotowska et al., 2016) resulted in high SOC and N stocks (Table 1),
479 which can support large microbial biomass in this zone (Haron et al., 1998). Thus, the low
480 mineral N leaching in the frond-stacked area may be attributed to immobilization of mineral N
481 by large microbial biomass, converting mobile NO_3^- to less mobile organic N (e.g. Corre et al.,
482 2010). In addition, it could be possible that palm root uptake of nutrients (including mineral N)
483 was higher in the frond-stacked area compared to the inter-row as roots proliferate in nutrient-
484 rich zones (Table 1; Hodge, 2004). This is supported by studies that showed higher root density
485 and higher water uptake under the frond piles compared to the inter-row (Rüegg et al., 2019;
486 Nelson et al., 2006). The high ECEC, base saturation and pH in frond-stacked area (Table 1),
487 despite having no direct lime application, were due to the release of nutrients from
488 decomposition of frond litter, which contain high levels of base cations (Kotowska et al., 2016).
489 Thus, although leaching of base cations were larger in the frond-stacked area than in the inter-
490 row (Fig. 4), these losses merely mirrored their high exchangeable levels (Table 1). Finally, the
491 leaching of Al was low in the frond-stacked area (Figs. 3 and 4) because Al becomes insoluble
492 as pH increased (i.e. lower exchangeable Al; Table 1). Altogether, these results highlighted the
493 benefits of piling senesced fronds onto the soil to reduce leaching of mineral N and Al, which
494 otherwise can potentially diminish ground water quality, and to amend soil fertility (Table 1).
495 Oil palm plantations in other areas (e.g. Borneo; Rahman et al., 2018) were reported to practice
496 piling of senesced fronds on every inter-row, which we did not observed in our study region as
497 that is claimed to hinder access to palms during harvest; nonetheless, our findings implied that
498 increase in the frond-stacked area can contribute to sustainable management practices of oil
499 palm plantations.

500

501 **4.3 Leaching losses under different intensity of management**



502 There was a clear influence of management intensity treatments on nutrient leaching losses with
503 a general reduction of leaching in reduced management intensity (Fig. 5; Table 3). In line with
504 our second hypothesis, the weeding methods clearly influenced leaching losses with a common
505 pattern of lower leaching fluxes in mechanical weeding than herbicide treatment (Fig. 5; Table
506 3). Mechanical weeding was associated with more ground vegetation cover (Darras et al., 2019)
507 and higher nutrient retention efficiency than herbicide weeding (Table 4), suggesting that faster
508 regrowth of understory vegetation by mechanical weeding have additionally contributed to the
509 uptake of nutrients and thus reducing leaching losses. This is in line with some studies in
510 temperate forests and a cedar plantation, which showed that understory vegetation can take up
511 excess NO_3^- in the soil (Olsson and Falkengren-Grerup, 2003) and reduce NO_3^- leaching and
512 the mobilization of Ca and Mg (Baba et al., 2011; Fukuzawa et al., 2006). Enhanced understory
513 vegetation in oil palm plantations may also positively impact biodiversity by increasing plant
514 species richness and soil macrofauna diversity and abundance (Luke et al., 2019; Ashton-Butt
515 et al., 2018), which may facilitate uptake and recycling of nutrients. Increase in soil macrofauna
516 might have contributed to lower leaching of Na with mechanical weeding (Fig. 5), since
517 herbivores and decomposers take up a large amount of Na (Kaspari et al., 2009). In addition,
518 the use of glyphosate is associated with possible health risks to workers and the environment
519 (van Bruggen et al., 2018); also, the economic gross margin (i.e. revenues minus costs) is
520 comparable between mechanical weeding and herbicide treatment because of needed labor for
521 periodic mechanical cutting of resistant ground vegetation in oil palm plantations with herbicide
522 weeding (Darras et al., 2019; Pahan, 2010). Altogether, these results advocate for the higher
523 sustainability of mechanical weeding over herbicide application.

524 The reduction of N fertilization rates decreased NO_3^- leaching, supporting our third
525 hypothesis. Comparing conventional and reduced fertilization rates, there were no differences
526 in total N stocks (section 3.1), mineral N levels (Darras et al., 2019), N retention efficiency



527 (Table 4) and oil palm yield (Darras et al., 2019), suggesting that excess N (above harvest
528 export; section 2.1) from high N fertilization was largely lost through leaching (Table 3). The
529 decreased Al and Ca leaching with reduced fertilization can be attributed to the lowered NO_3^-
530 leaching, since these were the accompanying cations (Figs. 4 and 5). Also, a reduction of Ca
531 leaching could have resulted from the lower application rate of triple superphosphate fertilizer,
532 which contains 16 % of Ca. The reduced K fertilization had no effect on K leaching (Fig. 5)
533 because K fertilization rate was only reduced by 15 % of the conventional rate due to high K
534 requirements of oil palm fruits (section 2.1). We conclude that this mature (16–20 years old)
535 plantation with conventional management was overly fertilized for N, and that a reduction in N
536 fertilization rate may be included in the Indonesian program for precision farming (Ministry of
537 Agriculture of Indonesia, 2016) to reduce environmental footprint of oil palm production.

538 Comparing the N leaching losses in the studied plantation with other fertilized tropical
539 plantations (Table A2), our plantation had higher N leaching than other large-scale oil palm
540 plantations on similar soils with comparable fertilization rates (Omoti et al., 1983; Tung et al.,
541 2009). However, in these studies the leaching losses were measured in the palm circle (Omoti
542 et al., 1983) or the sampling location was not specified (Tung et al., 2009), such that N leaching
543 may be underestimated as our results showed the high contribution of the inter-row to leaching
544 losses (Figs. 3 and 4). The N leaching fluxes in our plantation were also higher than in
545 smallholder oil palm plantations in the same area, which typically had much lower fertilization
546 rates (Kurniawan et al., 2018). On the other hand, our plantation had lower N leaching losses
547 than an oil palm plantation and coffee agroforestry systems on volcanic soils (Banabas et al.,
548 2008b; Cannavo et al., 2013; Tully et al., 2012), which have high inherent nutrient contents,
549 highly porous soils and high infiltration rates. The N leaching losses from our plantation were
550 also lower than in banana plantations, characterized by very high fertilization rates (Wakelin et
551 al., 2011; Armour et al., 2013).



552 The nutrients leached at 1.5 m depth should be considered lost from uptake of oil palm
553 roots, as majority of the root mass and the highest root density are in the top 0.5 m depth (Nelson
554 et al., 2006; Schroth et al., 2000; Kurniawan et al., 2018). The high leaching fluxes of NO_3^- and
555 Al implied a risk of groundwater pollution. During the high drainage fluxes following
556 fertilization, NO_3^- concentrations in soil-pore water reached to 20–40 mg L^{-1} in the inter-row
557 (covering 67 % of the plantation area), which was close to the 50 mg L^{-1} limit for drinking water
558 (WHO, 2011), and Al concentrations in soil-pore water exceeded the limit of 0.2 mg L^{-1} in 60
559 % of the samples. Nevertheless, before reaching to streams and rivers, these NO_3^- and Al
560 concentrations can be diluted by surface flow and retained in the soil along flow paths: NO_3^-
561 can be temporarily adsorbed in the deeper layers of highly weathered soils by its inherently
562 high anion exchange capacity (Harmand et al., 2010; Jankowski et al., 2018) and can be
563 consumed by denitrification (Wakelin et al., 2011). Riparian buffers can mitigate the transport
564 of these agricultural pollutants to streams (Luke et al., 2017; Chellaiah and Yule, 2018).
565 Restoring riparian buffers in former forests converted to oil palm plantations have been listed
566 as one sustainability criteria, endorsed by the Roundtable for Sustainable Palm Oil association
567 (RSPO, 2018), and may provide additional regulation services (Woodham et al., 2019).

568 **5 Conclusions**

569 Our findings show that nutrient leaching losses in an oil palm plantation differed among
570 management zones, as a result of fertilization, liming, mulching and of different drainage
571 fluxes. The reduction of management intensity, i.e. mechanical weeding with reduced
572 fertilization rates, was effective in reducing nutrient leaching losses without reduction in yield
573 at least during the first two years of this experiment (Darras et al., 2019). Long-term
574 investigation of this management experiment is important to get a reliable response of yield and
575 a holistic economic analysis, including valuation of regulation services. Greenhouse gas
576 emissions should also be quantified, as another important parameter of environmental footprint



577 of oil palm production. Our findings and these further investigations should be incorporated
578 into science-based policy recommendations such as those endorsed by the RSPO.



579 **Data availability**

580 All data of this study are deposited at the EFForTS-IS data repository ([https://efforts-is.uni-](https://efforts-is.uni-goettingen.de)
581 [goettingen.de](https://efforts-is.uni-goettingen.de)), an internal data-exchange platform, which is accessible to all members of the
582 Collaborative Research Center (CRC) 990. Based on the data sharing agreement within the
583 CRC 990, these data are currently not publicly accessible but will be made available through a
584 written request to the senior author.

585 **Author contribution**

586 GF performed the experiments, analysed the data and wrote the manuscript in consultation
587 with MDC. EV and MDC conceived and planned the experiment. XD helped carry out the
588 water model simulations. AT aided in field activities organization and granting collaborations
589 agreements. All authors contributed to the final version of the manuscript.

590 **Competing interests**

591 No conflict of interest to declare

592 **Aknowledgments**

593 This study was part of the project A05 in the CRC990-EFForTS, funded by the German
594 Research Foundation (DFG, Project ID: 192626868 – SFB 990). We acknowledge the
595 collaborations with PTPN VI plantation, Kevin Darras, and project Z01, for the
596 implementation and maintenance of this field experiment. We thank Christian Stiegler, with
597 project A03, for the climate data, and Eckart Priesack for the Expert-N water sub-model. We
598 especially thank our field and laboratory assistants for their valuable dedications in all field
599 and laboratory activities. This research was conducted under the research permit of Ministry
600 of Research and Technology of Indonesia, 539351/SIP/FRP/E5/Dit.KI/X/2016.



601 **References**

- 602 Abdalla, M., Hastings, A., Cheng, K., Yue, Q., Chadwick, D., Espenberg, M., Truu, J., Rees,
603 R. M., and Smith, P.: A critical review of the impacts of cover crops on nitrogen leaching,
604 net greenhouse gas balance and crop productivity, *Global Change Biology*, 25, 2530–
605 2543, doi:10.1111/gcb.14644, 2019.
- 606 Ahongshangbam, J., Khokthong, W., Ellsäßer, F., Hendrayanto, H., Hölscher, D., and Röhl,
607 A.: Drone-based photogrammetry-derived crown metrics for predicting tree and oil palm
608 water use, *Ecohydrology*, 12, 1–18, doi:10.1002/eco.2115, 2019.
- 609 Allen, K., Corre, M. D., Kurniawan, S., Utami, S. R., and Veldkamp, E.: Spatial variability
610 surpasses land-use change effects on soil biochemical properties of converted lowland
611 landscapes in Sumatra, Indonesia, *Geoderma*, 284, 42–50,
612 doi:10.1016/j.geoderma.2016.08.010, 2016.
- 613 Allen, K., Corre, M. D., Tjoa, A., and Veldkamp, E.: Soil nitrogen-cycling responses to
614 conversion of lowland forests to oil palm and rubber plantations in Sumatra, Indonesia,
615 *PloS One*, 10, e0133325, doi:10.1371/journal.pone.0133325, 2015.
- 616 Allen, R. G.: Crop evapotranspiration: Guidelines for computing crop water requirements,
617 *FAO irrigation and drainage paper*, 56, Rome, 300 pp., 1998.
- 618 Armour, J. D., Nelson, P. N., Daniells, J. W., Rasiah, V., and Inman-Bamber, N. G.: Nitrogen
619 leaching from the root zone of sugarcane and bananas in the humid tropics of Australia,
620 *Agriculture, Ecosystems & Environment*, 180, 68–78, doi:10.1016/j.agee.2012.05.007,
621 2013.
- 622 Ashton-Butt, A., Aryawan, A. A. K., Hood, A. S. C., Naim, M., Purnomo, D., Suhardi,
623 Wahyuningsih, R., Willcock, S., Poppy, G. M., Caliman, J.-P., Turner, E. C., Foster, W.
624 A., Peh, K. S. H., and Snaddon, J. L.: Understory vegetation in oil palm plantations
625 benefits soil biodiversity and decomposition rates, *Frontiers in Forests and Global Change*,
626 1, 1–13, doi:10.3389/ffgc.2018.00010, 2018.



- 627 Asner, G. P., Powell, G. V. N., Mascaro, J., Knapp, D. E., Clark, J. K., Jacobson, J., Kennedy-
628 Bowdoin, T., Balaji, A., Paez-Acosta, G., Victoria, E., Secada, L., Valqui, M., and
629 Hughes, F. R.: High-resolution forest carbon stocks and emissions in the Amazon,
630 Proceedings of the National Academy of Sciences of the United States of America, 107,
631 16738–16742, doi:10.1073/pnas.1004875107, 2010.
- 632 Baba, M., Abe, S., Kasai, M., Sugiura, T., and Kobayashi, H.: Contribution of understory
633 vegetation to minimizing nitrate leaching in a Japanese cedar plantation, Journal of Forest
634 Research, 16, 446–455, doi:10.1007/s10310-010-0244-3, 2011.
- 635 Banabas, M., Scotter, D. R., and Turner, M. A.: Losses of nitrogen fertiliser under oil palm in
636 Papua New Guinea: 2. Nitrogen transformations and leaching, and a residence time model,
637 Soil Research, 46, 340–347, doi:10.1071/SR07174, 2008a.
- 638 Banabas, M., Turner, M. A., Scotter, D. R., and Nelson, P. N.: Losses of nitrogen fertiliser
639 under oil palm in Papua New Guinea: 1. Water balance, and nitrogen in soil solution and
640 runoff, Australian Journal of Soil Research, 46, 332–339, doi:10.1071/SR07171, 2008b.
- 641 Bates, D., Maechler, M., Bolker, B., and Walker, S.: Fitting linear mixed-effect models using
642 lme4, Journal of Statistical Software, 67, 1–48, 2015.
- 643 Boafo, D. K., Kraisornpornson, B., Panphon, S., Owusu, B. E., and Amaniampong, P. N.:
644 Effect of organic soil amendments on soil quality in oil palm production, Applied Soil
645 Ecology, 147, 103358, doi:10.1016/j.apsoil.2019.09.008, 2020.
- 646 Cannavo, P., Harmand, J.-M., Zeller, B., Vaast, P., Ramírez, J. E., and Dambrine, E.: Low
647 nitrogen use efficiency and high nitrate leaching in a highly fertilized Coffea arabica–Inga
648 densiflora agroforestry system: a 15N labeled fertilizer study, Nutrient Cycling in
649 Agroecosystems, 95, 377–394, doi:10.1007/s10705-013-9571-z, 2013.
- 650 Carlson, K. M., Curran, L. M., Asner, G. P., Pittman, A. M., Trigg, S. N., and Marion
651 Adeney, J.: Carbon emissions from forest conversion by Kalimantan oil palm plantations,
652 Nature Climate Change, 3, 283–287, doi:10.1038/nclimate1702, 2013.



- 653 Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and Smith,
654 V. H.: Nonpoint pollution of surface waters with phosphorus and nitrogen, *Ecological*
655 *Applications*, 8, 559–568, doi:10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2,
656 1998.
- 657 Carter, C., Finley, W., Fry, J., Jackson, D., and Willis, L.: Palm oil markets and future supply,
658 *European Journal of Lipid Science and Technology*, 109, 307–314,
659 doi:10.1002/ejlt.200600256, 2007.
- 660 Chellaiah, D. and Yule, C. M.: Effect of riparian management on stream morphometry and
661 water quality in oil palm plantations in Borneo, *Limnologica*, 69, 72–80,
662 doi:10.1016/j.limno.2017.11.007, 2018.
- 663 Chou, C., Chiang, J. C. H., Lan, C.-W., Chung, C.-H., Liao, Y.-C., and Lee, C.-J.: Increase in
664 the range between wet and dry season precipitation, *Nature Geoscience*, 6, 263–267,
665 doi:10.1038/ngeo1744, 2013.
- 666 Clough, Y., Krishna, V. V., Corre, M. D., Darras, K., Denmead, L. H., Mejjide, A., Moser, S.,
667 Musshoff, O., Steinebach, S., Veldkamp, E., Allen, K., Barnes, A. D., Breidenbach, N.,
668 Brose, U., Buchori, D., Daniel, R., Finkeldey, R., Harahap, I., Hertel, D., Holtkamp, A.
669 M., Hörandl, E., Irawan, B., Jaya, I. N. S., Jochum, M., Klarner, B., Knohl, A., Kotowska,
670 M. M., Krashevskaya, V., Kreft, H., Kurniawan, S., Leuschner, C., Maraun, M., Melati, D.
671 N., Opfermann, N., Pérez-Cruzado, C., Prabowo, W. E., Rembold, K., Rizali, A., Rubiana,
672 R., Schneider, D., Tjitrosoedirdjo, S. S., Tjoa, A., Tschardtke, T., and Scheu, S.: Land-use
673 choices follow profitability at the expense of ecological functions in Indonesian
674 smallholder landscapes, *Nature Communications*, 7, 13137, doi:10.1038/ncomms13137,
675 2016.
- 676 Comte, I., Colin, F., Grünberger, O., Follain, S., Whalen, J. K., and Caliman, J.-P.:
677 Landscape-scale assessment of soil response to long-term organic and mineral fertilizer



- 678 application in an industrial oil palm plantation, Indonesia, *Agriculture, Ecosystems &*
679 *Environment*, 169, 58–68, doi:10.1016/j.agee.2013.02.010, 2013.
- 680 Corley, R. H. V. and Tinker, P. B.: The oil palm, 5th ed., John Wiley & Sons, Ltd, Hoboken,
681 NJ, 2016.
- 682 Corley, R.H.V.: How much palm oil do we need?, *Environmental Science & Policy*, 12, 134–
683 139, doi:10.1016/j.envsci.2008.10.011, 2009.
- 684 Corre, M. D., Veldkamp, E., Arnold, J., and Wright, S. J.: Impact of elevated N input on soil
685 N cycling and losses in old-growth lowland and montane forests in Panama, *Ecology*, 91,
686 1715–1729, doi:10.1890/09-0274.1, 2010.
- 687 Cusack, D. F., Silver, W., and McDowell, W. H.: Biological nitrogen fixation in two tropical
688 forests: ecosystem-level patterns and effects of nitrogen fertilization, *Ecosystems*, 12,
689 1299–1315, doi:10.1007/s10021-009-9290-0, 2009.
- 690 Darras, K. F. A., Corre, M. D., Formaglio, G., Tjoa, A., Potapov, A., Brambach, F., Sibhatu,
691 K. T., Grass, I., Rubiano, A. A., Buchori, D., Drescher, J., Fardiansah, R., Hölscher, D.,
692 Irawan, B., Kneib, T., Krashevskaya, V., Krause, A., Kreft, H., Li, K., Maraun, M., Polle,
693 A., Ryadin, A. R., Rembold, K., Stiegler, C., Scheu, S., Tarigan, S., Valdés-Urbe, A.,
694 Yadi, S., Tschardtke, T., and Veldkamp, E.: Reducing fertilizer and avoiding herbicides in
695 O oil palm plantations—ecological and economic valuations, *Frontiers in Forests and*
696 *Global Change*, 2, doi:10.3389/ffgc.2019.00065, 2019.
- 697 Dechert, G., Veldkamp, E., and Brumme, R.: Are partial nutrient balances suitable to evaluate
698 nutrient sustainability of land use systems? Results from a case study in central Sulawesi,
699 Indonesia, *Nutrient Cycling in Agroecosystems*, 72, 201–212, doi:10.1007/s10705-005-
700 1546-2, 2005.
- 701 Drescher, J., Rembold, K., Allen, K., Beckschäfer, P., Buchori, D., Clough, Y., Faust, H.,
702 Fauzi, A. M., Gunawan, D., Hertel, D., Irawan, B., Jaya, I. N. S., Klärner, B., Kleinn, C.,
703 Knohl, A., Kotowska, M. M., Krashevskaya, V., Krishna, V., Leuschner, C., Lorenz, W.,



- 704 Meijide, A., Melati, D., Nomura, M., Pérez-Cruzado, C., Qaim, M., Siregar, I. Z.,
705 Steinebach, S., Tjoa, A., Tschardtke, T., Wick, B., Wiegand, K., Kreft, H., and Scheu, S.:
706 Ecological and socio-economic functions across tropical land use systems after rainforest
707 conversion, *Philosophical Transactions of the Royal Society of London. Series B,*
708 *Biological sciences*, 371, doi:10.1098/rstb.2015.0275, 2016.
- 709 Dubos, B., Snoeck, D., and Flori, A.: Excessive use of fertilizer can increase leaching
710 processes and modify soil reserves in two Ecuadorian oil palm plantations, *Experimental*
711 *Agriculture*, 53, 255–268, doi:10.1017/S0014479716000363, 2017.
- 712 Edy, N., Yelianti, U., Irawan, B., Polle, A., and Pena, R.: Differences in root nitrogen uptake
713 between tropical lowland rainforests and oil palm plantations, *Frontiers in Plant Science*,
714 11, 92, doi:10.3389/fpls.2020.00092, 2020.
- 715 Elsenbeer, H.: Hydrologic flowpaths in tropical rainforest soils - a review, *Hydrological*
716 *Processes*, 15, 1751–1759, doi:10.1002/hyp.237, 2001.
- 717 Fan, Y., Rouspard, O., Bernoux, M., Le Maire, G., Panferov, O., Kotowska, M. M., and
718 Knohl, A.: A sub-canopy structure for simulating oil palm in the Community Land Model
719 (CLM-Palm): phenology, allocation and yield, *Geoscientific Model Development*, 8,
720 3785–3800, doi:10.5194/gmd-8-3785-2015, 2015.
- 721 Feng, X., Porporato, A., and Rodriguez-Iturbe, I.: Changes in rainfall seasonality in the
722 tropics, *Nature Climate Change*, 3, 811–815, doi:10.1038/nclimate1907, 2013.
- 723 Figueiredo, R. O., Markewitz, D., Davidson, E. A., Schuler, A. E., Watrin, O. d. S., and Silva,
724 P. d. S.: Land-use effects on the chemical attributes of low-order streams in the eastern
725 Amazon, *Journal of Geophysical Research*, 115, doi:10.1029/2009JG001200, 2010.
- 726 Food and Agriculture Organization, FAOSTAT: <http://faostat.fao.org/site/339/default.aspx>,
727 last access: 30 March 2020.
- 728 Fukuzawa, K., Shibata, H., Takagi, K., Nomura, M., Kurima, N., Fukazawa, T., Satoh, F., and
729 Sasa, K.: Effects of clear-cutting on nitrogen leaching and fine root dynamics in a cool-



- 730 temperate forested watershed in northern Japan, *Forest Ecology and Management*, 225,
731 257–261, doi:10.1016/j.foreco.2006.01.001, 2006.
- 732 Geist, H. J. and Lambin, E. F.: Proximate causes and underlying driving forces of tropical
733 deforestation, *BioScience*, 52, 143–150, doi:10.1641/0006-
734 3568(2002)052[0143:PCAUDF]2.0.CO;2, 2002.
- 735 Gibbs, H. K., Ruesch, A. S., Achard, F., Clayton, M. K., Holmgren, P., Ramankutty, N., and
736 Foley, J. A.: Tropical forests were the primary sources of new agricultural land in the
737 1980s and 1990s, *Proceedings of the National Academy of Sciences of the United States*
738 of America, 107, 16732–16737, doi:10.1073/pnas.0910275107, 2010.
- 739 Goh, K.-J., Härdter, R., and Fairhurst, T.: Fertilizing for maximum return, *Oil Palm*
740 Management for Large and Sustainable Yields, 279–306, 2003.
- 741 Grass, I., Kubitzka, C., Krishna, V. V., Corre, M. D., Mußhoff, O., Pütz, P., Drescher, J.,
742 Rembold, K., Ariyanti, E. S., Barnes, A. D., Brinkmann, N., Brose, U., Brümmer, B.,
743 Buchori, D., Daniel, R., Darras, K. F. A., Faust, H., Fehrmann, L., Hein, J., Hennings, N.,
744 Hidayat, P., Hölscher, D., Jochum, M., Knohl, A., Kotowska, M. M., Krashevskaya, V.,
745 Kreft, H., Leuschner, C., Lobite, N. J. S., Panjaitan, R., Polle, A., Potapov, A. M.,
746 Purnama, E., Qaim, M., Röhl, A., Scheu, S., Schneider, D., Tjoa, A., Tschardt, T.,
747 Veldkamp, E., and Wollni, M.: Trade-offs between multifunctionality and profit in
748 tropical smallholder landscapes, *Nature Communications*, 11, 1186, doi:10.1038/s41467-
749 020-15013-5, 2020.
- 750 Harmand, J.-M., Ávila, H., Oliver, R., Saint-André, L., and Dambrine, E.: The impact of
751 kaolinite and oxi-hydroxides on nitrate adsorption in deep layers of a Costarican Acrisol
752 under coffee cultivation, *Geoderma*, 158, 216–224, doi:10.1016/j.geoderma.2010.04.032,
753 2010.



- 754 Haron, K., Brookes, P. C., Anderson, J. M., and Zakaria, Z. Z.: Microbial biomass and soil
755 organic matter dynamics in oil palm (*Elaeis Guineensis* Jacq.) plantations, West Malaysia,
756 *Soil Biology and Biochemistry*, 30, 547–552, doi:10.1016/S0038-0717(97)00217-4, 1998.
- 757 Hodge, A.: The plastic plant: root responses to heterogeneous supplies of nutrients, *New*
758 *Phytologist*, 162, 9–24, doi:10.1111/j.1469-8137.2004.01015.x, 2004.
- 759 Hothorn, T., Bretz, F., and Westfall, P.: Simultaneous inference in general parametric models,
760 *Biometrical Journal*, 50, 346–363, 2008.
- 761 Huddell, A. M., Galford, G. L., Tully, K. L., Crowley, C., Palm, C. A., Neill, C., Hickman, J.
762 E., and Menge, D. N. L.: Meta-analysis on the potential for increasing nitrogen losses
763 from intensifying tropical agriculture, *Global Change Biology*, 26, 1–13,
764 doi:10.1111/gcb.14951, 2020.
- 765 Jankowski, K., Neill, C., Davidson, E. A., Macedo, M. N., Costa, C. J., Galford, G. L.,
766 Santos, L. M., Lefebvre, P., Nunes, D., Cerri, C. E. P., McHorney, R., O’Connell, C., and
767 Coe, M. T.: Deep soils modify environmental consequences of increased nitrogen fertilizer
768 use in intensifying Amazon agriculture, *Scientific Reports*, 13478, doi:10.1038/s41598-
769 018-31175-1, 2018.
- 770 Kaspari, M., Yanoviak, S. P., Dudley, R., Yuan, M., and Clay, N. A.: Sodium shortage as a
771 constraint on the carbon cycle in an inland tropical rainforest, *Proceedings of the National*
772 *Academy of Sciences of the United States of America*, 106, 19405–19409,
773 doi:10.1073/pnas.0906448106, 2009.
- 774 Kotowska, M. M., Leuschner, C., Triadiati, T., and Hertel, D.: Conversion of tropical lowland
775 forest reduces nutrient return through litterfall, and alters nutrient use efficiency and
776 seasonality of net primary production, *Oecologia*, 180, 601–618, doi:10.1007/s00442-015-
777 3481-5, 2016.
- 778 Kotowska, M. M., Leuschner, C., Triadiati, T., Meriem, S., and Hertel, D.: Quantifying
779 above- and belowground biomass carbon loss with forest conversion in tropical lowlands



- 780 of Sumatra (Indonesia), *Global Change Biology*, 21, 3620–3634, doi:10.1111/gcb.12979,
781 2015.
- 782 Kurniawan, S., Corre, M. D., Matson, A. L., Schulte-Bisping, H., Utami, S. R., van Straaten,
783 O., and Veldkamp, E.: Conversion of tropical forests to smallholder rubber and oil palm
784 plantations impacts nutrient leaching losses and nutrient retention efficiency in highly
785 weathered soils, *Biogeosciences*, 15, 5131–5154, doi:10.5194/bg-15-5131-2018, 2018.
- 786 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: LmerTest package: tests in linear
787 mixed effects models, *Journal of Statistical Software*, 82, doi:10.18637/jss.v082.i13, 2017.
- 788 Lamade, E., Djegui, N., and Leterme, P.: Estimation of carbon allocation to the roots from
789 soil respiration measurements of oil palm, *Plant and Soil*, 181, 329–339,
790 doi:10.1007/BF00012067, 1996.
- 791 Li, Y. and Ghodrati, M.: Preferential transport of nitrate through soil columns containing root
792 channels, *Soil Science Society of America Journal*, 58, 653–659,
793 doi:10.2136/sssaj1994.03615995005800030003x, 1994.
- 794 Lohse, K. A. and Matson, P.: Consequences of nitrogen additions for soil losses from wet
795 tropical forests, *Ecological Applications*, 15, 1629–1648, doi:10.1890/03-5421, 2005.
- 796 Luke, S. H., Barclay, H., Bidin, K., Chey, V. K., Ewers, R. M., Foster, W. A., Nainar, A.,
797 Pfeifer, M., Reynolds, G., Turner, E. C., Walsh, R. P. D., and Aldridge, D. C.: The effects
798 of catchment and riparian forest quality on stream environmental conditions across a
799 tropical rainforest and oil palm landscape in Malaysian Borneo, *Ecohydrology:*
800 ecosystems, land and water process interactions, *ecohydrogeomorphology*, 10, 1-17,
801 doi:10.1002/eco.1827, 2017.
- 802 Luke, S. H., Purnomo, D., Advento, A. D., Aryawan, A. A. K., Naim, M., Pikstein, R. N., Ps,
803 S., Rambe, T. D. S., Soeprapto, Caliman, J.-P., Snaddon, J. L., Foster, W. A., and Turner,
804 E. C.: Effects of understory vegetation management on plant communities in oil palm



- 805 plantations in Sumatra, Indonesia, *Frontiers in Forests and Global Change*, 2, 1–13,
806 doi:10.3389/ffgc.2019.00033, 2019.
- 807 Mejjide, A., La Rua, C. de, Guillaume, T., Röhl, A., Hassler, E., Stiegler, C., Tjoa, A., June,
808 T., Corre, M. D., Veldkamp, E., and Knohl, A.: Measured greenhouse gas budgets
809 challenge emission savings from palm-oil biodiesel, *Nature Communications*, 11, 1089,
810 doi:10.1038/s41467-020-14852-6, 2020.
- 811 Mejjide, A., Röhl, A., Fan, Y., Herbst, M., Niu, F., Tiedemann, F., June, T., Rauf, A.,
812 Hölscher, D., and Knohl, A.: Controls of water and energy fluxes in oil palm plantations:
813 environmental variables and oil palm age, *Agricultural and Forest Meteorology*, 239, 71–
814 85, doi:10.1016/j.agrformet.2017.02.034, 2017.
- 815 Ministry of Agriculture of Indonesia: Oil palm replanting guideline, Regulation No:
816 18/Permentan/KB.330/5/2016, 2016.
- 817 Mohanty, S., Swain, C. K., Tripathi, R., Sethi, S. K., Bhattacharyya, P., Kumar, A., Raja, R.,
818 Shahid, M., Panda, B. B., Lal, B., Gautam, P., Munda, S., and Nayak, A. K.: Nitrate
819 leaching, nitrous oxide emission and N use efficiency of aerobic rice under different N
820 application strategy, *Archives of Agronomy and Soil Science*, 64, 465–479,
821 doi:10.1080/03650340.2017.1359414, 2018.
- 822 Moradi, A., Teh, C. B. S., Goh, K. J., Husni, A. M. H., and Ishak, C. F.: Effect of four soil
823 and water conservation practices on soil physical processes in a non-terraced oil palm
824 plantation, *Soil and Tillage Research*, 145, 62–71, doi:10.1016/j.still.2014.08.005, 2015.
- 825 Murdiyarso, D., Hergoualc’h, K., and Verchet, L. V.: Opportunities for reducing greenhouse
826 gas emissions in tropical peatlands, *Proceedings of the National Academy of Sciences of
827 the United States of America*, 107, 19655–19660, doi:10.1073/pnas.0911966107, 2010.
- 828 Neill, C., Coe, M. T., Riskin, S. H., Krusche, A. V., Elsenbeer, H., Macedo, M. N.,
829 McHorney, R., Lefebvre, P., Davidson, E. A., Scheffler, R., Figueira, A. M. e. S., Porder,
830 S., and Deegan, L. A.: Watershed responses to Amazon soya bean cropland expansion and



- 831 intensification, *Philosophical Transactions of the Royal Society of London. Series B,*
832 *Biological sciences*, 368, 20120425, doi:10.1098/rstb.2012.0425, 2013.
- 833 Nelson, P. N., Banabas, M., Scotter, D. R., and Webb, M. J.: Using soil water depletion to
834 measure spatial distribution of root activity in oil palm (*Elaeis Guineensis* Jacq.)
835 plantations, *Plant and Soil*, 286, 109–121, doi:10.1007/s11104-006-9030-6, 2006.
- 836 Nyamangara, J., Bergström, L. F., Piha, M. I., and Giller, K. E.: Fertilizer use efficiency and
837 nitrate leaching in a tropical sandy soil, *Journal of environmental quality*, 32, 599–606,
838 doi:10.2134/jeq2003.5990, 2003.
- 839 Olsson, M. O. and Falkengren-Grerup, U.: Partitioning of nitrate uptake between trees and
840 understory in oak forests, *Forest Ecology and Management*, 179, 311–320,
841 doi:10.1016/S0378-1127(02)00544-3, 2003.
- 842 Omoti, U., Ataga, D. O., and Isenmila, A. E.: Leaching losses of nutrients in oil palm
843 plantations determined by tension lysimeters, *Plant and Soil*, 73, 365–376,
844 doi:10.1007/BF02184313, 1983.
- 845 Pahan, I.: Complete guide to oil palm, 8th ed., Penebar Swadaya, Jakarta, Indonesia, 160-164,
846 2010.
- 847 Pardon, L., Bessou, C., Nelson, P. N., Dubos, B., Ollivier, J., Marichal, R., Caliman, J.-P., and
848 Gabrielle, B.: Key unknowns in nitrogen budget for oil palm plantations. A review,
849 *Agronomy for Sustainable Development*, 36, 20, doi:10.1007/s13593-016-0353-2, 2016.
- 850 Pirker, J., Mosnier, A., Kraxner, F., Havlík, P., and Obersteiner, M.: What are the limits to oil
851 palm expansion?, *Global Environmental Change*, 40, 73–81,
852 doi:10.1016/j.gloenvcha.2016.06.007, 2016.
- 853 Priesack, E.: Expert-N model library documentation, Institute of Soil Ecology, 2005.
- 854 R Core Team: R: A language and environment for statistical computing, Foundation for
855 Statistical Computing, Vienna, Austria, 2019.



- 856 Rahman, N., Neergaard, A. de, Magid, J., van de Ven, G. W. J., Giller, K. E., and Bruun, T.
857 B.: Changes in soil organic carbon stocks after conversion from forest to oil palm
858 plantations in Malaysian Borneo, *Environmental Research Letters*, 13, 105001,
859 doi:10.1088/1748-9326/aade0f, 2018.
- 860 Röhl, A., Niu, F., Mejjide, A., Ahongshangbam, J., Ehbrecht, M., Guillaume, T., Gunawan,
861 D., Hardanto, A., Hendrayanto, Hertel, D., Kotowska, M. M., Kreft, H., Kuzyakov, Y.,
862 Leuschner, C., Nomura, M., Polle, A., Rembold, K., Sahner, J., Seidel, D., Zemp, D. C.,
863 Knohl, A., and Hölscher, D.: Transpiration on the rebound in lowland Sumatra,
864 *Agricultural and Forest Meteorology*, 274, 160–171, doi:10.1016/j.agrformet.2019.04.017,
865 2019.
- 866 Roy, E. D., Richards, P. D., Martinelli, L. A., Della Coletta, L., Lins, S. R. M., Vazquez, F.
867 F., Willig, E., Spera, S. A., VanWey, L. K., and Porder, S.: The phosphorus cost of
868 agricultural intensification in the tropics, *Nature plants*, 2, 16043,
869 doi:10.1038/nplants.2016.43, 2016.
- 870 RSPO: Principles and criteria: For the production of sustainable palm oil, 2018.
- 871 Rüegg, J., Quezada, J. C., Santonja, M., Ghazoul, J., Kuzyakov, Y., Buttler, A., and
872 Guillaume, T.: Drivers of soil carbon stabilization in oil palm plantations, *Land
873 Degradation & Development*, 30, 1–12, doi:10.1002/ldr.3380, 2019.
- 874 Schlesinger, W. H. and Bernhardt, E. S.: *Biogeochemistry: An analysis of global change*, 3rd
875 ed., Academic Press, Waltham, Mass, 672 pp., 2013.
- 876 Schroth, G., Rodrigues, M.R.L., and D'Angelo, S. A.: Spatial patterns of nitrogen
877 mineralization, fertilizer distribution and roots explain nitrate leaching from mature
878 Amazonian oil palm plantation, *Soil Use and Management*, 16, 222–229,
879 doi:10.1111/j.1475-2743.2000.tb00197.x, 2000.



- 880 Silver, W. L., Thompson, A. W., Reich, A., Ewel, J. J., and Firestone, M. K.: Nitrogen
881 cycling in tropical plantation forests: potential controls on nitrogen retention, *Ecological*
882 *Applications*, 15, 1604–1614, doi:10.1890/04-1322, 2005.
- 883 Steiner, C., Glaser, B., Geraldtes Teixeira, W., Lehmann, J., Blum, W. E.H., and Zech, W.:
884 Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol
885 amended with compost and charcoal, *Journal of Plant Nutrition and Soil Science*, 171,
886 893–899, doi:10.1002/jpln.200625199, 2008.
- 887 Tarigan, S., Stiegler, C., Wiegand, K., Knohl, A., and Murti Laksono, K.: Relative contribution
888 of evapotranspiration and soil compaction to the fluctuation of catchment discharge: case
889 study from a plantation landscape, *Hydrological Sciences Journal*, 58, 1–10,
890 doi:10.1080/02626667.2020.1739287, 2020.
- 891 Tarigan, S., Wiegand, K., Sunarti, and Slamet, B.: Minimum forest cover required for
892 sustainable water flow regulation of a watershed: a case study in Jambi Province,
893 Indonesia, *Hydrology and Earth System Sciences*, 22, 581–594, doi:10.5194/hess-22-581-
894 2018, 2018.
- 895 Tarigan, S. D., Sunarti, Wiegand, K., Dislich, C., Slamet, B., Heinonen, J., and Meyer, K.:
896 Mitigation options for improving the ecosystem function of water flow regulation in a
897 watershed with rapid expansion of oil palm plantations, *Sustainability of Water Quality*
898 *and Ecology*, 8, 4–13, doi:10.1016/j.swaqe.2016.05.001, 2016.
- 899 Teklu, B. M., Hailu, A., Wiegand, D. A., Scholten, B. S., and van den Brink, P. J.: Impacts of
900 nutrients and pesticides from small- and large-scale agriculture on the water quality of
901 Lake Ziway, Ethiopia, *Environmental Science and Pollution Research*, 25, 13207–13216,
902 doi:10.1007/s11356-016-6714-1, 2018.
- 903 Tokuchi, N., Samejima, H., Hon, J., and Fukushima, K.: Influence of herbicide use in oil palm
904 plantations on stream water chemistry in Sarawak, *Anthropogenic Tropical Forests*, 209–
905 216, doi:10.1007/978-981-13-7513-2_11, 2019.



- 906 Tully, K. L., Lawrence, D., and Scanlon, T. M.: More trees less loss: nitrogen leaching losses
907 decrease with increasing biomass in coffee agroforests, *Agriculture, Ecosystems &*
908 *Environment*, 161, 137–144, doi:10.1016/j.agee.2012.08.002, 2012.
- 909 Tung, P. G. A., Yusoff, M. K., Majid, N. M., Joo, G. K., and Huang, G. H.: Effect of N and K
910 fertilizers on nutrient leaching and groundwater quality under mature oil palm in Sabah
911 during the monsoon period, *American Journal of Applied Sciences*, 6, 1788–1799, 2009.
- 912 van Breemen, N., Mulder, J., and Driscoll, C. T.: Acidification and alkalization of soils, *Plant*
913 *and Soil*, 283–308, 1983.
- 914 van Bruggen, A.H.C., He, M. M., Shin, K., Mai, V., Jeong, K. C., Finckh, M. R., and Morris,
915 J. G.: Environmental and health effects of the herbicide glyphosate, *Science of the Total*
916 *Environment*, 616-617, 255–268, doi:10.1016/j.scitotenv.2017.10.309, 2018.
- 917 van Straaten, O., Corre, M. D., Wolf, K., Tchienkoua, M., Cuellar, E., Matthews, R. B., and
918 Veldkamp, E.: Conversion of lowland tropical forests to tree cash crop plantations loses
919 up to one-half of stored soil organic carbon, *Proceedings of the National Academy of*
920 *Sciences of the United States of America*, 112, 9956–9960, doi:10.1073/pnas.1504628112,
921 2015.
- 922 Vijay, V., Pimm, S. L., Jenkins, C. N., and Smith, S. J.: The impacts of oil palm on recent
923 deforestation and biodiversity loss, *PloS One*, 11, e0159668,
924 doi:10.1371/journal.pone.0159668, 2016.
- 925 Wakelin, S. A., Nelson, P. N., Armour, J. D., Rasiah, V., and Colloff, M. J.: Bacterial
926 community structure and denitrifier (*nir-gene*) abundance in soil water and groundwater
927 beneath agricultural land in tropical North Queensland, Australia, *Soil Research*, 49, 65–
928 76, doi:10.1071/SR10055, 2011.
- 929 WHO: Guidelines for drinking-water quality, 4th ed., World Health Organization, Geneva,
930 2011.



- 931 Wong, M. T.F., Hughes, R., and Rowell, D. L.: Retarded leaching of nitrate in acid soils from
932 the tropics: measurement of the effective anion exchange capacity, *Journal of Soil*
933 *Science*, 655–663, 1990.
- 934 Woodham, C. R., Aryawan, A. A. K., Luke, S. H., Manning, P., Caliman, J.-P., Naim, M.,
935 Turner, E. C., and Slade, E. M.: Effects of replanting and retention of mature oil palm
936 riparian buffers on Ecosystem functioning in oil palm plantations, *Frontiers in Forests and*
937 *Global Change*, 2, 29, doi:10.3389/ffgc.2019.00029, 2019.
- 938



939 **Tables and figures**

940 **Table 1** Soil physical and biochemical characteristics (mean \pm standard errors, $n = 4$ plots) in
 941 the top 50 cm depth for each management zone, averaged across experimental treatments.
 942 Means within a row followed by different letters indicate significant differences among
 943 management zones (one-way ANOVA with Tukey HSD or Kruskal–Wallis H test with multiple
 944 comparisons extension at $P \leq 0.05$). Bulk density measured in the top 10 cm of soil, whereas
 945 all the other parameters are for the 0–50 cm soil depth: element stocks are the sum of the
 946 sampled soil depths (0–5 cm, 5–10 cm, 10–30 cm and 30–50 cm) and the rest are depth-
 947 weighted averages, calculated for each replicate plot. ECEC, effective cation exchange capacity

| Soil properties | | Palm circle | Frond-stacked area | Inter-row |
|-----------------------------------|---------------------------------|-------------------|----------------------|-------------------|
| Bulk density | g cm^{-3} | 1.37 ± 0.01^a | 0.89 ± 0.01^b | 1.36 ± 0.01^b |
| Soil organic C | kg m^{-2} | 6.2 ± 0.6^b | 9.1 ± 0.8^a | 6.4 ± 0.2^b |
| Total N | g m^{-2} | 402 ± 31^b | 571 ± 39^a | 426 ± 15^{ab} |
| soil C:N ratio | | 15.5 ± 0.5^a | 15.7 ± 0.3^a | 15.0 ± 0.5^a |
| ^{15}N natural abundance | ‰ | 5.9 ± 0.1^a | 5.3 ± 0.2^a | 5.7 ± 0.2^a |
| pH | 1:4 (H ₂ O) | 5.05 ± 0.08^a | 5.00 ± 0.08^{ab} | 4.81 ± 0.05^b |
| ECEC | $\text{mmol}_c \text{ kg}^{-1}$ | 35 ± 2^a | 28 ± 2^a | 18 ± 1^b |
| Base saturation | % | 48 ± 3^a | 46 ± 4^a | 20 ± 2^b |
| Aluminum saturation | % | 52 ± 4^b | 50 ± 2^b | 78 ± 2^a |
| Mg | g m^{-2} | 32 ± 3^a | 28 ± 6^a | 9 ± 1^b |
| Ca | g m^{-2} | 169 ± 21^a | 157 ± 15^a | 37 ± 5^b |
| K | g m^{-2} | 39 ± 13^a | 13 ± 1^b | 6 ± 1^b |
| Na | g m^{-2} | 1.5 ± 0.4^a | 0.7 ± 0.2^a | 0.6 ± 0.2^a |
| Al | g m^{-2} | 66 ± 4^b | 71 ± 4^{ab} | 87 ± 3^a |



| | | | | |
|----|-------------------|-----------------|-----------------|-----------------|
| Fe | g m^{-2} | 1.4 ± 0.2^a | 1.8 ± 0.4^a | 1.8 ± 0.5^a |
| Mn | g m^{-2} | 0.7 ± 0.1^b | 1.8 ± 0.3^a | 0.6 ± 0.2^b |
| H | g m^{-2} | 0.2 ± 0.0^a | 0.2 ± 0.0^a | 0.2 ± 0.1^a |

948

949 **Table 2** Annual water balance simulated from March 2017 to February 2018 for each
 950 management zone.

| Water flux (mm yr^{-1}) | Palm circle | Fronnd-stacked area | Inter-row |
|------------------------------------|-------------|---------------------|-----------|
| Precipitation | 2772 | 2772 | 2772 |
| Transpiration | 828 | 448 | 401 |
| Evaporation | 228 | 214 | 434 |
| Interception | 351 | 209 | 209 |
| Runoff | 338 | 0 | 216 |
| Drainage (at 1.5 m depth) | 556 | 1806 | 1179 |

951



952 **Table 3** Annual leaching losses at 1.5 m depth for each experimental treatment from March
 953 2017 to February 2018. Values are area-weighted averages of leaching losses in each
 954 management zone (mean \pm standard error, $n = 4$ plots). Means followed by different letters
 955 indicate differences among experimental treatments (linear-mixed effect models on monthly
 956 values followed by Tukey HSD test for multiple comparisons at $P \leq 0.05$). Treatments: ch =
 957 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
 958 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding. DON =
 959 dissolved organic N; TDN = total dissolved N.

| Element leaching (kg ha ⁻¹ yr ⁻¹) | ch | cw | rh | rw |
|--|------------------------------|-------------------------------|------------------------------|------------------------------|
| NO ₃ ⁻ -N | 71.5 \pm 20.1 ^a | 48.2 \pm 13.0 ^{ab} | 36.3 \pm 20.1 ^b | 30.0 \pm 5.7 ^b |
| NH ₄ ⁺ -N | 1.7 \pm 0.2 ^a | 1.7 \pm 0.1 ^a | 1.8 \pm 0.1 ^a | 1.7 \pm 0.2 ^a |
| DON | 0.5 \pm 0.5 ^a | 0.6 \pm 0.3 ^a | 0.4 \pm 0.1 ^a | 0.3 \pm 0.0 ^a |
| TDN | 73.6 \pm 20.2 ^a | 50.4 \pm 13.1 ^{ab} | 38.4 \pm 8.9 ^b | 32.0 \pm 5.8 ^b |
| Ca | 26.6 \pm 4.3 ^a | 19.4 \pm 4.4 ^b | 18.2 \pm 1.8 ^b | 17.0 \pm 2.1 ^b |
| Mg | 11.6 \pm 2.5 ^a | 7.7 \pm 0.8 ^b | 9.1 \pm 0.7 ^{ab} | 10.8 \pm 3.6 ^{ab} |
| K | 8.1 \pm 1.3 ^a | 6.2 \pm 0.7 ^b | 8.9 \pm 0.6 ^a | 5.7 \pm 1.1 ^b |
| Na | 15.9 \pm 3.5 ^{ab} | 13.6 \pm 2.4 ^b | 18.9 \pm 3.1 ^a | 13.1 \pm 1.2 ^b |
| Mn | 0.3 \pm 0.1 ^a | 0.2 \pm 0.0 ^b | 0.2 \pm 0.0 ^{bc} | 0.1 \pm 0.0 ^c |
| Total Al | 40.8 \pm 11.5 ^a | 20.8 \pm 7.6 ^b | 19.9 \pm 6.8 ^b | 21.8 \pm 3.1 ^b |
| Total S | 2.4 \pm 0.5 ^a | 1.8 \pm 0.4 ^a | 2.1 \pm 0.6 ^a | 4.9 \pm 3.3 ^a |
| Total Fe | 0.2 \pm 0.0 ^a | 0.5 \pm 0.3 ^a | 0.2 \pm 0.0 ^a | 0.5 \pm 0.3 ^a |
| Total P | 0.0 \pm 0.0 ^a | 0.1 \pm 0.0 ^a | 0.0 \pm 0.0 ^a | 0.0 \pm 0.0 ^a |



CI 79.7 ± 15.8^a 36.9 ± 8.3^b 67.7 ± 8.7^a 78.3 ± 7.5^a

960

961

962 **Table 4** N and base cation retention efficiencies in the soil for each management zone and
 963 experimental treatment (means \pm standard error, $n = 4$ plots). Means followed by different
 964 lowercase letters indicate differences among experimental treatments for each management
 965 zone, whereas different uppercase letters indicate differences among management zones for
 966 each experimental treatment (one-way ANOVA with Tukey HSD or Kruskal–Wallis H test
 967 with multiple comparisons extension at $P \leq 0.05$). Weighted-average is based on the areal
 968 coverage of each management zone: 18 % for palm circle, 15 % for frond-stacked area, and 67
 969 % for inter-row. Treatments: ch = conventional fertilization–herbicide; cw = conventional
 970 fertilization–mechanical weeding; rh = reduced fertilization–herbicide; rw = reduced
 971 fertilization–mechanical weeding. See section 2.4 for calculations of N and base cation
 972 retention efficiency.

| | ch | cw | rh | rw |
|--|------------------------|-------------------------|-------------------------|------------------------|
| N retention efficiency ($\text{mg N m}^{-2} \text{d}^{-1} / \text{mg N m}^{-2} \text{d}^{-1}$) | | | | |
| Palm circle | 0.987 ± 0.002^{aA} | 0.982 ± 0.007^{aAB} | 0.986 ± 0.003^{aAB} | 0.997 ± 0.000^{aA} |
| Frond-stacked area | 0.984 ± 0.004^{aA} | 0.989 ± 0.004^{aA} | 0.993 ± 0.001^{aA} | 0.987 ± 0.002^{aA} |
| Inter-row | 0.877 ± 0.025^{aB} | 0.870 ± 0.022^{aB} | 0.900 ± 0.018^{aB} | 0.906 ± 0.039^{aA} |
| Weighted-average | 0.925 ± 0.022^a | 0.934 ± 0.020^a | 0.945 ± 0.012^a | 0.946 ± 0.018^a |
| Base cation retention efficiency ($\text{mol}_c \text{m}^{-2} \text{yr}^{-1} / \text{mol}_c \text{m}^{-2} \text{yr}^{-1}$) | | | | |




| | | | | |
|--------------------|--------------------------|-------------------------|--------------------------|--------------------------|
| Palm circle | $0.967 \pm 0.008^{ab A}$ | $0.982 \pm 0.002^{a A}$ | $0.937 \pm 0.013^{b A}$ | $0.974 \pm 0.010^{ab A}$ |
| FronD-stacked area | $0.884 \pm 0.013^{b A}$ | $0.950 \pm 0.004^{a A}$ | $0.960 \pm 0.002^{a A}$ | $0.928 \pm 0.016^{ab A}$ |
| Inter-row | $0.588 \pm 0.086^{b B}$ | $0.875 \pm 0.022^{a B}$ | $0.704 \pm 0.048^{ab B}$ | $0.822 \pm 0.063^{ab A}$ |
| Weighted-average | 0.876 ± 0.009^b | 0.945 ± 0.007^a | 0.902 ± 0.019^{ab} | 0.934 ± 0.012^{ab} |

973




974 **Figure 1** Pearson correlation test between modeled (red line) and field-measured soil water
975 matric potential (black points) ($n = 50$ field measurements over one year) for each management
976 zone at 30 and 60 cm depths.



977




978 **Figure 2** Monthly water drainage at 1.5 m depth, simulated in each management zone, and
979 daily rainfall from March 2017 to February 2018. The gray shaded area represent the dry season
980 (precipitation $< 140 \text{ mm month}^{-1}$)



981




982 **Figure 3.** Partial cation-anion charge balance of the major solutes (with concentrations > 0.03
 983 mg L⁻¹) in soil water at 1.5 m depth for each experimental treatment in the different
 984 management zones. The concentrations of organic acids (RCOO⁻) and carbonates (HCO₃⁻) are
 985 calculated as the difference between the measured cations and anions. Treatments: ch =
 986 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
 987 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding.



988




989 **Figure 4** Monthly leaching losses at 1.5 m depth (mean \pm standard errors, $n = 4$ plots) for each
990 management zone. Black arrows indicate fertilizer applications and the gray shaded area
991 represents the dry season (precipitation < 140 mm month⁻¹).



992



1993 **Figure 5** Average monthly leaching losses at 1.5 m depth for each experimental treatment from
1994 March 2017 to February 2018. Values are area-weighted averages of leaching losses in each
1995 management zone (means \pm standard errors, $n = 4$ plots). For each parameter, different letters
1996 indicate significant differences among treatments (linear-mixed effect models on monthly
1997 values followed by Tukey HSD test for multiple comparisons at $P \leq 0.05$). Treatments: ch =
1998 conventional fertilization–herbicide; cw = conventional fertilization–mechanical weeding; rh =
1999 reduced fertilization–herbicide; rw = reduced fertilization–mechanical weeding
1000



1001



1002 **Appendices**

1003 **Table A1** Gross N mineralization rates (means \pm SE, $n = 4$ plots) in the top 5 cm soil for each
1004 treatment and management zone in a large-scale plantation in Jambi, Indonesia. Measurements
1005 were done on intact soil cores in February 2018 using the ^{15}N pool dilution technique, as
1006 described in details by Allen et al. (2015). Treatments: ch = conventional fertilization–
1007 herbicide; cw = conventional fertilization–mechanical weeding; rh = reduced fertilization–
1008 herbicide; rw = reduced fertilization–mechanical weeding

Gross N mineralization ($\text{mg N m}^{-2} \text{d}^{-1}$)

| | ch | cw | rh | rw |
|--------------------|----------------|----------------|----------------|----------------|
| palm circle | 2.2 ± 0.6 | 1.9 ± 0.4 | 1.8 ± 0.6 | 3.4 ± 0.2 |
| frond-stacked area | 22.4 ± 3.3 | 32.5 ± 8.0 | 22.4 ± 7.2 | 16.6 ± 5.2 |
| inter-row | 4.8 ± 1.1 | 4.0 ± 0.6 | 3.8 ± 0.8 | 4.4 ± 0.9 |

1009 *Note:* These data are not included in the manuscript to avoid double-publication as these
1010 results were reported in our previous study (Formaglio et al., unpublished data).



1011 **Table A2** Literature comparison of annual N fertilization and total N leaching losses across
 1012 tropical plantations.

| Author | Soil type | rainfall (mm yr ⁻¹) | Type of plantation management | N applied (kg ha ⁻¹ yr ⁻¹) | Total N leaching (kg ha ⁻¹ yr ⁻¹) | Percentage N leached (%) |
|-----------------------|------------|------------------------------------|-------------------------------------|--|---|--------------------------------|
| Present study | loam | 2772 | intensive oil | 260 | 74 | 28 |
| | Acrisol | | palm | | | |
| Present study | loam | 2772 | intensive oil | 130 | 38 | 28 |
| | Acrisol | | palm | | | |
| Omoti et al. 1983 | sandy clay | 2000 | intensive oil | 150 | 9 | 6 |
| | Acrisol | | palm | | | |
| Kurniawan et al. 2018 | loam | 3418 | smallholder | 88 | 11 | 12.5 |
| | Acrisol | | oil palm | | | |
| Tung et al. 2009 | Acrisol | - | intensive oil palm | 128 | 3 (150 days) | 2 |
| Tung et al. 2009 | Acrisol | - | intensive oil palm | 251 | 3 (150 days) | 1 |
| Banabas et al. 2008 | clay loam | 2398 | intensive oil | 100 | 37 | 37 |
| | Andosol | | palm | | | |
| Banabas et al. 2008 | sandy loam | 3657 | intensive oil | 100 | 103 | 103 |
| | Andosol | | palm | | | |
| Cannavo et al. 2013 | clay loam | 2678 | coffee | 250 | 157 | 63 |
| | Andosol | | agroforestry | | | |



| | | | | | | |
|---------------------|--------------|------|--------------|-----|-----|----|
| Tully et al., 2012 | clay loam | 2700 | coffee | 120 | 119 | 99 |
| | Andosol | | agroforestry | | | |
| Armour et al. 2013 | clay Acrisol | 1958 | intensive | 476 | 164 | 34 |
| | | | banana | | | |
| Wakelin et al. 2011 | loam | 2685 | intensive | 305 | 116 | 38 |
| | Acrisol | | banana | | | |

1013


1014

1015 **Figure A1** Lysimeter locations at each treatment plot, with two subplots (blue rectangles) that

1016 each included the three management zones (blue crosses): 1) lysimeters in the palm circle

1017 were at 1 m from the palm trunk, 2) in the frond-stacked area, at about 4 m from the palm

1018 trunk, and 3) in the inter-row, at approximately 4 m from the palm trunk.



1019

1020