

Replies to the comments of Referee #1

The Paper deals with stable Isotope in trees at treeline. It is a valuable contribution to the Journal, there are some Major flaws. In the introduction the authors should mention why Isotope data Derived from tree rings are powerful Tools for climate Reconstructions

5 Following the reviewer's suggestion we added the following sentence in the introduction: "In environments where the trees are rarely moisture stressed, like at the Alps tree-line, the control of $\delta^{13}\text{C}$ is the photosynthetic rate, which depends on predominantly on irradiance and temperature. $\delta^{18}\text{O}$ probably reflects a combination of the direct temperature effect on the isotopic ratio of precipitation and the indirect evaporative enrichment effect. (McCarroll et al. 2003)" We did not mention dD since its paleoclimatic value is still in its infancy and not yet fully understood.

10 There is also evidence that in both Species investigated isotope signatures derived from cellulose are comparable to those from bulk wood.

15 It is known that for carbon, oxygen and hydrogen a correlation exists between cellulose and bulk wood (Gori et al, 2013 and Wieser et al, 2016). However, present work uses samples of subfossil wood, in which the wood composition is variably affected by different states of conservation. For this reason, we chose to analyse the cellulose, although it required additional work.

See Frontiers in plant science 2016,7;799 and agformet 2018,248;251

20 Thank you for the suggestion of the two very interesting papers we missed before. At line 71 we added the following sentence: "This last point is particularly important, because the trees at the Alps tree-line benefit from an enhanced CO_2 fertilization and from a recent temperature increase (Wieser et al. 2016) that may produce a long term trend in cellulose isotopes. For example Wieser et al. (2018) show that the increase of temperature is cancelled by increasing atmospheric CO_2 concentrations in the environment of the Alps treeline under non-limiting water availability. They state that therefore the instantaneous water use efficiency of photosynthesis did not change considerably."

25 In fig 1 and table 1 please give full names of all the sites

We added the site names in table 1. But the addition of the long and bulky names in Fig. 1 would make it difficult to interpret, so we think it is better to use the codes

The results section line 119- 143 should be moved to methods.

30 The indicated section presents the properties of the database we interrogated and an initial analysis to understand which of raw, normalized or scaled data are better suited to detect possible trends in the database. So we are convinced that this section belongs to the Result rather than to the Method section.

In the discussion I miss studies carried out in pine and larch at various treeline Sites in the Swiss and Austrian alps. These data also Show that cembra pine and larch Show similar growth and isotope responses during the last decades. This is in contrast to this manuscript and hence should be discussed.

35 We found only a couple of papers that analysed the age effect in stable isotope in tree rings at mountain treeline, namely Daux et al. (2011) used a larch located at Les Grang, 2050 m.a.s.l. and Esper et al. (2010) used *Pinus uncinata* collected at tree line at about 2300 masl in the Spanish central Pyrenees. This is confirmed by a very recent paper (Büntgen et al, 2020). These papers underlined that the interpretation of the isotopes in the wood of the last decades is particularly complex due to CO_2 fertilization. In this study, we used a population of trees that covers the last 9,000 years, which is better suited to understand the non-climatic trends. In addition, these authors do not mention the cambial age of analysed samples. In case their trees were in the adult phase, the data would not be in contrast with our work.

Replies to the comments of Referee #2

5 The manuscript “Alpine Holocene Tree –Ring Dataset: Age-related trends in the stable isotopes of cellulose show species-specific patterns” by Arosio and co-workers assesses the age-related trends in the stable isotopic compositions of about 200 tree-ring samples collected from the Alps. The total period covered by these samples is approximately the Holocene, though individual samples span up to several hundreds of years.

10 One of the problems in tree ring width analysis is that the ring width often varies with the age of the tree. Thus the non-climatic factors come into play. This problem, to some extent, is encountered in the case of the isotopic records of tree rings as well. Though the effect of the age on tree ring width is removed by employing statistical techniques, a similar procedure, however, is not so common among the isotope dendroclimatologists. In this study, the authors argue that the age-related trend in isotopic data, especially during the early years of tree growth, is quite prevalent. For example, $\delta^{18}\text{O}$ of tree-ring showed inconsistent patterns during the juvenile period of tree growth. So these authors propose a methodology whereby the detrending issue could be avoided. They analyzed a large number of tree ring samples. They suggested that 15 detrending of the tree ring isotopic record is not necessary if the isotopic values are plotted against the cambial age. But various issues need to be fixed before such a method is applied.

We thank the reviewer for the nice summary of our work

For example, which factors are mainly responsible for introducing the so-called age effect in the isotopic values of tree ring cellulose?

20 Thank you for highlighting an interesting point. Unfortunately, the mechanisms that drive age-effects in the isotopic values of cellulose mainly restricted to the juvenile period are not known. In literature there are many hypotheses mostly related to age-dependent changes in physiology. Our work was aimed at assessing the presence or absence of these effects in the alpine treeline environment regarding three isotopes and two tree species. The differences for $\delta^{13}\text{C}$ and δD trends in the juvenile phases of the two species are probably 25 due to different biochemical processes, considering that the trees come from the same environment and that the mean value is the average off all the samples that come from different time epochs and therefore they should not be affected by climate trends. This is presented in the introduction and by fig. 2.

30 Why is the result not similar for both oxygen and hydrogen? Since the pathway for both H and O is essentially the same, that is the soil water, then why the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ trends sometimes show opposite behavior?

35 Oxygen and Hydrogen originate from the source, i.e. soil water, but they are processed by the trees in a completely different way. $\delta^{18}\text{O}$ is generally enriched because of evapotranspiration in the leaves, while δD is generally depleted because of an additional biological fractionation in the photosynthetic processes due to the exchanges with NADH. This is well described by Cormier et al. (2018) (doi: 10.1111/nph.15016) and also in our manuscript (“Larch cellulose is significantly depleted in deuterium isotopes with respect to evergreen conifers” Arosio et al. 2020) that is presently under review in *Frontiers in Earth Science*.

40 Then follows the next question: unless the mechanism is reasonably understood, how can one propose a method to fix the problem? It is known that during environmental stress, such as in a drought season if a particular ring becomes thin, the corresponding $\delta^{13}\text{C}$ would be high. So a long term increasing trend in $\delta^{13}\text{C}$ may be indicative of a prolonged dry condition. But the question may be asked, whether the trend arises as a result of tree ring aging. So the mechanism of causing an age-related trend in an isotopic record must be understood well.

45 The aim of the work was not to propose a method but rather to verify how tree aging in tree line cembra pine and larch affected the tree-ring isotopes, an important issue for paleoclimatic use of the isotopes. It was reported that $\delta^{13}\text{C}$ is a proxy for conditions of drought (McCarroll and Loader, 2003) but in our case we are analysing a large population of trees covering different climatic periods looking for non-climatic signals. In addition, it is known that at the Alps treeline there is no soil water limitation, due to the frequent rain during the growth season. (Wieser et al, 2016). Also, a major canopy effect on $\delta^{13}\text{C}$ in the juvenile phase can reasonably be excluded since the two species lived in the same environments.

In this context, it may be said that the ^{14}C records of tree rings also show age-related trends. But other than the physiological aspect, a physical process, the decay of the radioactive carbon is known to cause such a trend. Hence, by applying a decay correction scheme, the age-related trend on ^{14}C tree ring records can be easily corrected.

5 The physical process of decay does not apply to the stable isotopes we analysed, so the aging effect is related only to physiological/biochemical processes, some of which are known to cause major fractionations. They are complex and involve the relationship between autotrophism (use of the reserves) and heterotrophism (photosynthesis) which vary during the season and during aging and cannot easily be quantified. Furthermore, the interpretation of the ^{14}C may also not be so straight forward since it combines
10 the physical decay but also the non-physical, i.e. physiological/biochemical processes. Depending on the weighting of these processes, a simple interpretation of a physical effect only may be misleading.

In this work, the authors show that isotopic records, especially d^{18}O and d^{13}C , when plotted against the cambial age are not susceptible to age-related trends after a threshold period characterized by juvenile growth. However, the method does not apply hydrogen isotopes. Hence it may not offer a comprehensive
15 solution.

The variation of dD after the juvenile period is small indeed, even smaller than the analytical precision of our measurements of 3‰, and the variation was found in larch and not in cembra pine. Aim of the work was to detect the absence or presence of age trends in the two species for the three isotopes, without the ambition to offer a solution of the paleoclimatic use of isotopes.

20 As mentioned earlier, long term trends in isotopic values are not considered significantly affecting the paleoclimatic interpretation. Hence they are not typically removed. The authors should give evidence that such practice needs reconsideration so that they can justify their work.

We do not agree with the statement “isotopic values are not considered significantly affecting the paleoclimatic interpretation”. In fact, in the introduction we summarized the literature on the data of stable isotopes for paleoclimatic interpretation, stressing the differences among the isotopes, the conflicting results for $\delta^{18}\text{O}$ and the limited data for δD . So there is a lack of agreement about the present or absence of age trends and on how to use the data. For example, recent papers detrended $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ data of tree ring cellulose for a paleo climatic use (Esper et al, 2010 and 2015), while in other papers the values were used undetrended. However, to be clearer, we added a sentence in the introduction quoting examples of papers
25 using undetrended and detrended values of the cellulose stable isotope. “A key question for isotope dendroclimatology is whether isotope ratios of the tree cellulose show age trends or not (McCarroll and Loader, 2004). It is still a controversial issue depending on isotope type and plant species. If tree-age related trends are absent, the analysis and reconstruction of long-term climatic evolutions based on tree-ring isotope series would lose a source of potential bias. Contrary, if they are present, some work suggests to use a detrending procedure (Esper et al. 2010). The same issue concerns the juvenile phase, the values of which can be excluded from paleoclimatic analyses if they show complex age effects; as suggested by (McCarroll and Loader 2004).”
30

The authors have presented all three isotopic records, viz d^{13}C , d^{18}O , and dD , but their inter-relation has not been discussed. d^{18}O and d^{13}C in biological systems (such as corals) often show a strong positive correlation, indicating the presence of kinetic fractionation, and in turn it shows a non-climatic effect. Hence such kind of analysis may be helpful in this context. Similarly, the relation between d^{18}O and dD should also be examined.
35

Thank you for the interesting suggestion. Indeed we have been looking at the correlations between the different isotopes and TRW of various cambial age classes, but the preliminary results show that they are complex and their interpretation needs more work, which is outside of the aim of this work. The correlation between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ in corals is explained by the formation of carbonate, which is very different from the formation of glucose in the photosynthetic and post-photosynthetic processes.
45

Minor issues: Though the manuscript is in general, written well, it contains grammatical errors. (i.e, Line 170-171).

50 Corrected, thank you

Referencing is not done correctly, pls check Line 308-309.

Corrected, thank you

Alpine Holocene Tree-Ring Dataset: Age-related trends in the stable isotopes of cellulose show species-specific patterns.

Tito Arosio^{1,2}, Malin M. Zieher^{1,2,5}, Kurt Nicolussi³, Christian Schlüchter^{2,4}, Markus Leuenberger^{1,2}

¹Climate and Environmental Physics, Physics Institute, University of Bern, 3012 Bern, Switzerland

5 ²Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

³Department of Geography, Universität Innsbruck, 6020 Innsbruck, Austria

⁴Institute of Geological Sciences, University of Bern, 3012 Bern, Switzerland

⁵Swiss Tropical and Public Health Institute, Socinstrasse 57, 4051 Basel, Switzerland

Correspondence to: Tito Arosio (tito.arosio@climate.unibe.ch)

10 **Abstract.** Stable isotopes in tree-ring cellulose are important tools for climatic reconstructions even though their interpretation could be challenging due to non-climate signals, primarily those related to tree ageing. Previous studies on the presence of tree-age related trends during juvenile as well as adult growth phases in δD , $\delta^{18}O$ and $\delta^{13}C$ time series yielded variable results that are not coherent among different plant species. We analysed possible trends in the extracted cellulose of tree-rings of 85 larch trees and 119 cembran pine trees, i.e. in samples of one deciduous and one evergreen conifer species

15 collected at the treeline in the Alps covering nearly the whole Holocene. The age trend analyses of all tree-ring variables were conducted on the basis of mean curves established by averaging the cambial-age aligned tree series. For cambial ages over 100 years, our results prove the absence of any age-related effect in the δD , $\delta^{18}O$ and $\delta^{13}C$ time series for both the evergreen as well the deciduous conifer species, with the only exception of larch δD . However, for lower cambial ages, we found trends that differ for each isotope and species. I.e., mean $\delta^{13}C$ values in larch do not vary with ageing and can be used

20 without detrending, whereas those in cembran pine show a juvenile effect and the data should be detrended. Mean $\delta^{18}O$ values present two distinct ageing phases for both species complicating detrending. Similarly, mean δD values in larch change in the first 50 yr whereas cembran pine between 50-100 yr. Values for these two periods of cambial age for δD and $\delta^{18}O$ should be used with caution for climatic reconstructions, ideally complemented by additional information regarding mechanisms for these trends.

25 1 Introduction

Stable isotopes in tree-ring cellulose are powerful tools for climatic reconstructions (Kress et al. 2010; Nagavciuc et al. 2019). In environments where the trees are rarely moisture stressed, like at the Alps tree-line, the dominant control of $\delta^{13}\text{C}$ is the photosynthetic rate, which depends on predominantly on irradiance and temperature. $\delta^{18}\text{O}$ probably reflects a combination of the direct temperature effect on the isotopic ratio of precipitation isotopic ratio and the indirect evaporative enrichment effect. (McCarroll et al. 2003).

predominantly isotopic ratio of effect An advantage of stable isotope time-series based on tree-rings compared to other isotope time series is the defined dating and temporal, e.g. annual, resolution. A challenge for the climatic interpretation of many tree ring parameters is the presence of non-climate signals, primarily those related to tree ageing. Tree-ring width (TRW) and maximum latewood density show ageing effects, which usually have to be removed with detrending/standardization procedures before using them for climatic reconstruction (Helama 2017). A key question for isotope dendroclimatology is whether isotope ratios of the tree cellulose show age trends or not (McCarroll and Loader, 2004).; It is still a controversial issue depending on isotope type and plant species. If tree-age related trends are absent, the analysis and reconstruction of long-term climatic evolutions based on tree-ring isotope series would lose a source of potential bias. Contrary, if they are present, some work suggests to use a detrending procedure (Esper et al. 2010). The same issue concerns the juvenile phase, the values of which can be excluded from paleoclimatic analyses if they show complex age effects; as suggested by (McCarroll and Loader 2004)~~A key question for isotope dendroclimatology is whether isotope ratios of the tree cellulose also show age trends (McCarroll and Loader 2004), still a controversial issue depending on isotope type and plant species. If tree-age related trends are absent the analysis and reconstruction of long-term climatic evolutions based on tree-ring isotope series would lose a source of potential bias.~~

45 For $\delta^{13}\text{C}$, ageing studies were initiated by Freyer et al. (1979) on many tree species, followed by other investigations on-, e.g., pines, oaks and beeches. All documented an increase for $\delta^{13}\text{C}$ values in the juvenile period (Anderson et al. 2005; Duquesnay et al. 1998; Gagen et al. 2008; Li et al. 2005; McCarroll and Pawellek 2001; Monserud and Marshall 2001; Nagavciuc et al. 2019; Raffalli-Delercé et al. 2004; Treydte et al. 2001). A “long term trend” with continuous increase of $\delta^{13}\text{C}$ values during the entire life of *Pinus sylvestris* was also described (Helama et al. 2015). An early juvenile effect in the first 5 yr was reported for oak (Duffy et al. 2017). Fewer works did not detect any juvenile or long-term trend of $\delta^{13}\text{C}$ values in *Pinus sylvestris* and larch (Gagen et al. 2007; Kilroy et al. 2016; Young et al. 2011). Altogether, most studies found a juvenile trend (increase), but of variable lengths.

An initial work on $\delta^{18}\text{O}$ values found a negative juvenile trend of 300 yr in *Juniperus turkestanica* grown at elevations around 3000 m a.s.l. (Treydte et al. 2006). A similar negative juvenile trend of 300 y was reported also in *Pinus uncinata*,
55 (Esper et al. 2010). A positive trend of 30 yr was found for oak (Labuhn et al. 2014) and positive “long term trends” were

reported for spruce and beeches (Klesse et al. 2018). In contrast, other studies did not find detectable age trends in larch, oak, *Pinus sylvestris*, *Abies alba*, cembra pine (Daux et al. 2011; Duffy et al. 2017; Duffy et al. 2019; Nagavciuc et al. 2019; Saurer et al. 2000; Young et al. 2011). Thus, the studies on $\delta^{18}\text{O}$ trends are rather controversial, showing negative, positive or no age-related trends. The mentioned studies analyzed trees from different geographical origins, including some from sites near the treeline where trees often grow in open space. Such growth situations may have an effect on age trends, i.e., under open space growth there is no neighbourhood competition and therefore no limitation of growth by adjacent trees (Matsushita et al, 2015). The absence of a canopy effect may modify also the isotope composition stored by the trees, as shown for $\delta^{18}\text{O}$ (Daux et al. 2011; Esper et al. 2010; Gagen et al. 2008; Nagavciuc et al. 2019; Young et al. 2011). Furthermore, open space growing limits effects described previously such as tree dominance-suppression or location/exposition effects (wet – dry; sunny – shady) (Leuenberger, 2007) that may lead to age trends. The tree height also imposes a hydraulic limitation and possibly reduces stomatal conductance that may lead to an increase of the cellulose $\delta^{13}\text{C}$ values with increasing age (Brienen et al., 2017).

Changes of δD values in relation to tree ages have been analyzed only in two studies. One found a positive juvenile trend of 20 yr followed by a flat phase in spruce trees (Lipp et al. 1993) and the other identified a positive long-term trend in oak (Mayr et al. 2003).

Helama et al. (2015) suggested that the availability of a suitable database of tree-ring isotopes would allow to detect age trends and would open possibilities to their elimination in order to improve the recognition of long-term climatic evolution. Five issues were indicated as necessary for such a database: i) that it contains a large sample number, ii) that it has no data from “pooled” rings, iii) that the samples are well distributed over calendar years with different climate conditions, iv) that the samples come from timberline or treeline sites, where the distance between the trees is large, i.e., limiting the canopy effect, v) that they do not contain “modern” rings that need to be corrected in $\delta^{13}\text{C}$ for the anthropogenic rise of CO_2 concentration in the atmosphere. This last point is particularly important, because the trees at the Alps tree-line benefit from an enhanced CO_2 fertilization and from a recent temperature increase (Wieser et al. 2016) that may produce a long term trend in cellulose isotopes. For example Wieser et al. (2018-) show that combined effects of the increases of temperature is cancelled by and of increasing the atmospheric CO_2 concentrations in the environment of the Alps treeline under non-limiting water availability. They state that therefore, with the result that the instantaneous water use efficiency of photosynthesis did not change considerably~~This last point is particularly important, because the trees at the Alps tree-line benefit from an enhanced CO_2 fertilization and from a recent temperature increase (Wieser et al. 2016) that may produce a long term trend in cellulose isotopes. For example Wieser et al. (2018-) found that in the environment of the Alps treeline under non-limiting water availability, the temperature increase counteracts the increase of atmospheric CO_2 and leaf conductance for water vapour, the main driver of $\delta^{13}\text{C}$ in tree rings.~~This last point is particularly important, because the trees at Alps tree-line benefit from CO_2 fertilization and from temperature increase (Wieser et al.

~~2016) that may produce a long-term trend in cellulose isotopes. For example Wieser et al. (2018) found that in the environment of Alps treeline under non-limiting water availability, the temperature increase counteracts the increase of atmospheric CO₂ and leaf conductance for water vapour, the main driver of δ¹³C in tree rings.~~

Here we investigated the presence of age trends by utilizing a stable-isotope tree-ring database which was established on the base of the Eastern Alpine Conifer Chronology (Nicolussi et al. 2009). The database consists of i) samples of mainly
95 subfossil wood from 201 trees, ii) the isotope samples are not pooled, iii) the isotope time series with up to multi-centennial length are continuously covering the last ca. 9000 years, iv) it utilizes two different species: deciduous larch and evergreen
cembran pine, v) the wood material was collected at different treeline sites and vi) it contains only 17 trees with rings that grew after the industrial revolution. In the present work we aim to verify the presence of age trends in δD, δ¹⁸O and δ¹³C in comparison with those of cellulose content (CC) and TRW. We considered the whole, multi-centennial cambial age range of
100 the trees to identify and quantify the length and extent of the juvenile and the long-term ageing periods.

2 Material and methods

2.1 Subfossil wood samples and sampling sites in the Alps

Holocene wood sections were available at the Department of Geography of the University of Innsbruck, where the Eastern Alpine Conifer Chronology (EACC) has been established on the base of calendar-dated tree-ring width series (Nicolussi et al.
105 2009). We have utilized a large number of these subfossil wood samples that cover nearly the whole Holocene (Nicolussi et al. 2009). They belong to the deciduous larch (*Larix decidua* Mill.) and the evergreen cembran pine (*Pinus cembra* L.) and have been collected at treeline sites for paleo-climatic studies. The sampling sites are located in different parts of the European Alps covering a SW-NE transect with an elevation range of 1,930 to 2,400 m (Fig. 1). The wood material was collected at 29 different sites, 3 of them have only larch, 15 only cembran pine and 11 sites contain both species. The
110 characteristics of the 201 trees are listed in Table 1. Only 17 of them contain tree rings formed after the industrial revolution, i.e. after ca. 1850 AD. Samples of 5-year spanning wood have been prepared and analyzed for stable isotope ratios, as described before (Ziehmer et al. 2018; Arosio et al., 2020).

2.2 Tree-ring width data and cambial age estimation

The tree-ring width of all the samples were measured with a precision of 0.001 mm as described in Nicolussi et al. (2009).
115 Dated tree samples with relatively wide rings were selected to collect enough material for the isotope measurements. In

subfossil specimen the number of rings available for analyses often do not cover the whole tree lifespan due to effects of decay processes and therefore the cambial age of the first measured ring was estimated from ring curvature for tree samples without preserved pith (Nicolussi et al. 2009)

2.3 Stable isotope analysis

120 The procedure of cellulose extraction, determination of the cellulose content (cellulose dry weight / wood dry weight) (Ziehmer et al. 2018) and the triple-isotope analysis were described before (Loader et al. 2015). Briefly, we used conventional Isotope Ratio Mass Spectrometry (Isoprime 100) coupled to a pyrolysis unit (HEKAtech GmbH, Germany), which is similar to the previously used TC/EA (for technical details see (Leuenberger 2007)). This approach was extended to measurements of non-exchangeable hydrogen of alpha-cellulose using the on-line equilibration method (Filot et al. 2006; 125 Loader et al. 2015). The results are reported in per mil (‰) relative to the Vienna Pee Dee Belemnite (VPDB) for carbon and to Vienna Standard Mean Ocean Water (VSMOW) for hydrogen and oxygen (Coplen 1994). The precision of the measurement is $\pm 3.0\%$ for hydrogen, $\pm 0.3\%$ for oxygen and $\pm 0.15\%$ for carbon (Loader et al. 2015).

2.4 Carbon isotope correction

The burning of fossil fuels and land-use changes of the Industrial Period from about 1850 onwards caused a continuous 130 increase of atmospheric carbon dioxide (CO₂) depleted in $\delta^{13}\text{C}$ (Leuenberger 2007) known as Suess Effect (Suess 1955). This change is reflected in the carbohydrates of the plants, therefore a correction has to be applied to the isotopic series of tree rings. For all the $\delta^{13}\text{C}$ values after the 1000 AD we applied the correction factor described in Leuenberger (2007).

2.5 Age trend analysis

Each tree series was aligned on the cambial age and larch and cembran pine samples were analyzed independently. The 135 values of the three isotope ratios were analyzed as raw, normalized and z-scored data. The normalization consisted in subtracting the mean of the time series of a tree from each raw value of this series. For the analysis of the age-related trends, the isotope series of these three data groups were averaged to mean series for both investigated species under consideration of cambial age of the tree time series. We limited the analysis of these isotope mean series to the cambial age period where the replication number is ≥ 10 (Klesse et al. 2018; Young et al. 2011). We plotted the normalized data of the isotope 140 values versus the cambial age of the trees. Then we applied a linear interpolation in the different parts of the curves to quantify the trend. The division of the curves is different for each isotope, showcasing their different behaviours. The same procedure was applied to TRW and CC series, but for TRW we used the raw in place of the normalized data. To verify the presence of trends we applied a linear fit and compared it with those of the isotopes.

Our aim was to interrogate the stable isotopes of the EACC database for age effects using the well-known age-trends in TRW as comparison. The plot of the cambial age versus calendar age (Fig. 2A) shows that the database includes isotope time series covering age ranges from 15 to 610 years, with only few of them starting from the pith. The cambial ages of the trees are rather uniformly distributed over the entire Holocene, thus avoiding a potential bias in the analysis of age-trend. Figure 150 2A also shows that the time series of the two species, larch (red) and cembran pine (green) are similarly distributed over the Holocene. The sample replication number per cambial age of the trees is shown in figure 2B, with the horizontal line indicating the threshold ≥ 10 , that ranged from 1 to 460yr for cembran pine and -from 10 to 480yr for larch. ~~We considered this replication threshold important in this study.~~

The isotope series of the individual larch trees have a mean length of 273 yr (with a minimum and maximum length of 25 155 and 550 yr, respectively), the mean of their initial measure is 75 yr and the mean final cambial age is 348 yr. The individual cembran pine time series have a mean length of 257 yr (with a minimum and maximum length of 15 and 610 yr, respectively), the mean of their initial data point is 67 yr and the mean final cambial age is 324 yr. Table 1 describes the samples we analyzed and shows that they cover the period 8930 b2k to 2010 A.D., with a maximum cambial age of 725 yr.

The means of raw data of each isotope of larch and cembran pine were plotted versus cambial age (Fig 3B, 4B, 5B). The 160 absolute mean values differ probably because of different geographical origin of the trees or species-specific signature (Arosio et al., 2020), therefore we analyzed also the trend with the normalized data. The geographical effect may influence not only the mean but also the variance of each series, thus altering the age trend. To verify it, we used the z-scored data by dividing the normalized values by the standard deviation for each tree. No consistent difference was found between the normalized and z-score data (supplemental, fig. S1), indicating that the variance of the isotope series was not significantly 165 influenced by geographical factors.

We found that all average isotope series show trend changes only in the first 100 yr of cambial age in agreement with previous reports (Esper, 2015) except δD of larch. Therefore, we analyzed the average series, before and after 100 yr, separately. The trends of the average series with sample replication above or equal to 10 for all subperiods were studied by linear correlations, as in Young et al. (2011).

170 3.1 $\delta^{13}\text{-Carbon}$

Means of the standardized $\delta^{13}\text{C}$ data of all samples from 1 to about 500 cambial yr of both species are shown in figure 3A. Data points with a replication of ≥ 10 are considered, as shown in the replication plot of figure 2A. The plots of the raw 175 and of the normalized data are shown in figure 3B. The mean raw values of cembran pine (cyan) are more depleted than the ones of larch (red) (Fig. 3B). After normalization the mean values of the two species overlap only partially, since in the juvenile phase of the two species show different trends (fig 3B). Mean values for larch are stable, while the cembran pine documents a strong positive trend in the first 100 yr (0.7 ‰/100 yr) followed by a stabilization (Fig. 3C).

3.2 $\delta^{18}\text{-Oxygen}$

The same approach was used to study the variation of $\delta^{18}\text{O}$ of all samples (Fig. 4A) with their mean values in light green. Raw and normalized data are shown in figure 4B. The larch raw data are evidently more $\delta^{18}\text{O}$ enriched than the cembran pine ones, and the normalization strongly reduces the difference between the two species and the two time series are almost overlapping (fig 4B, right). In Figure 4C only the normalized averages series of the two species are plotted. Both of them show a peak in the first 100 yr followed by a phase without major age trend. Linear regression was applied to separate an initial phase of 50 yr with increasing values, followed by a sharp decrease until 100 yr and then a stabilization. The initial increasing phase in the larch was less steep than that in cembran pine, and for the rest the patterns are similar.

185 3.3 $\delta\text{-Deuterium}$

Means of the standardized δD data of all samples from 1 to about 500 cambial yr of the two species are shown in figure 5A. Means of δD raw values clearly indicate that larch is more depleted than cembran pine (fig. 5B, left) as shown before (Arosio et al. 2020). After normalization the two plots partially overlap (fig 5B, right). Figure 5C shows the means of the two species, the pattern of which are rather different. Larch shows a steep initial decrease in the first 50 yr, after a short steep increase within 10 years the values follows a minor increase through all the time, the cembran pine documents an initial slight decrease in the first 50 yr, followed by a steep positive trend of 50 yr and a flat line from 100 yr on (Fig. 5E).

3.4 TRW

The same analysis has also been applied to the non-detrended tree-ring width values, with the difference that in fig 6A the raw data are used. Figure 6A shows all the raw values and the mean value in light green. The plots of the raw and normalized data, expressed in cm, show a similar trend (Fig. 6B). Means with a replication ≥ 10 (Fig 6B and C) show a maximum at around 30 yr, in agreement with previous reports (Bräker 1981), after which the values of both species steadily decrease in two slope sections until 300 yr, thereafter becoming flat (Bräker 1981). For similarity with the isotope analyses, we applied linear regression to the data, instead of the more common exponential regression. For a comparison between the analyses of TRW and For similarity with the isotope values, we applied linear regression to the TRW data, instead of the more common exponential regression.

3.5 Cellulose Content

The same analysis has also been applied to data of the cellulose content. Figure 7A shows all raw values and the mean values in light green. The plots of the raw and normalized data, expressed in percent, show a similar trend (Fig. 7B). Means with a replication ≥ 10 (Fig 7B and C) present a remarkable increase in the first 50 yr in both species, from a cambial age of 51 yr to the end the larch presents a decreasing trend, while the cembran pine shows no trend.

4 Discussion

210 A characteristic of our present work is that the wood samples represent two conifer species, consisting of 201 trees that were collected at 29 different sites at high elevations in the Alps. They were exposed to different environmental conditions such as, e.g., elevation, aspect, slope steepness and water availability. Therefore, we normalized all records by subtracting by the mean of the tree from the raw values (Daux et al. 2011). A different approach was used by Helama et al. (2015) who used the raw data to analyze samples from only three different sites. Here we present analysis of a much more extended database, in time and space, which certainly represents the natural variability realistically. However, there are still issues that requires consideration, in particular the sample replication. Our database does not have a constant sample replication throughout the cambial age of the trees. It is low at the beginning and increases in the first 50 yr, and decreases sharply after 450 yr. This may have some effect on the study of the age trends.

220 As introduced in the results section, we have divided tree ageing in a juvenile period that we deliberately terminated at 100 yr, and the long-term period that lasted until 450 yr. A major conclusion of this study is that the values of δD , $\delta^{18}O$ and $\delta^{13}C$ in long term period from 100 yr to 450 yr did not change significantly, except δD in larch. This is in agreement with previous $\delta^{13}C$ studies on evergreen conifers (Esper et al. 2015; Gagen et al. 2007; Gagen et al. 2008; Klesse et al. 2018; Nagavciuc et al. 2019; Saurer et al. 2004; Young et al. 2011) and with the $\delta^{18}O$ previous studies on larch (Daux et al. 2011; Kilroy et al. 2016; Nagavciuc et al. 2019). This implies that no detrending is necessary of tree isotope data for climate analysis with cambial age in that range, with the exception of δD in larch where a non-significantly trend is present.

225 More complex are the data of the juvenile period, during which the trend behavior differs among the different isotopes and species. In the juvenile phase we found evidence for a positive trend of $\delta^{13}C$ in evergreen cembran pine but not in deciduous larch. These data are in good agreement with studies on evergreen conifers (*Picea abies*, *Pinus sylvestris*, *Pinus uncinata*) that all found an initial positive trend lasting up to 50 yr (Gagen et al. 2007; Gagen et al. 2008), or 100 yr (Klesse et al. 2018) or 200 yr (Esper et al. 2015). Moreover, two previous studies on larch did not report any evident trend for $\delta^{13}C$ in the juvenile period (Daux et al. 2011; Kilroy et al. 2016b). We can conclude that there is a general agreement that deciduous larch and evergreen conifers behave differently in the juvenile period in respect to $\delta^{13}C$ of the cellulose. Further work is needed to understand the reason of this difference.

235 The behavior of $\delta^{18}O$ values in the initial period is rather complex, with a maximum around 50 yr and a decrease up to 100 yr in both species. This is in good agreement with a previous work, that showed, that in *Pinus uncinata* grown at the tree-line have maximal $\delta^{18}O$ values around 20-50 yr followed by a negative trend (Esper et al. 2010) and another study on beech and

240 spruce that found a positive juvenile trend the persisted beyond 50 years of age (Klesse et al. 2018). Altogether, our data show that $\delta^{18}\text{O}$ of cellulose of larch and cembran pine has ageing trends that are similar to those of other tree species, with up and down trends in the first 100 yr and an intermediate maximum at around 50 yr. The significance of these trends remain to be further studied. However, considering the time and space of our database covers, this result seems to be widespread and temporally robust.

245 Our results on δD demonstrate different patterns for larch and cembran pine in the juvenile period, similarly to $\delta^{13}\text{C}$. The evergreen cembran pine displays an initial flat phase of 25 yr, then an increase of 4‰ till 100 yr. This is in partial agreement with a previous work that showed an increase of δD values in the juvenile period, but this lasted only 20 yr (Lipp et al. 1993). Another work measured δD by nitration of the cellulose to remove non-exchangeable hydrogens (Leavitt 2010), but this is certainly not the reason for this difference, as demonstrated by Filot et al. (2006). The difference can be attributed to the different growth environments of the trees, one at an elevation of 330 m near Bad Windsheim (Germany) and the other at a mean altitude of 2100 m, at tree-line sites in the Alps, where tree growth is known to be much slower with prolonged juvenile phase (Körner 2003; Ott 1978). The only other work that analyzed the evolution of δD in cellulose during ageing reported a constant increase of δD values in the first 175 years in oak (Mayr et al. 2003). In our work the deciduous larch showed a different pattern with a strong decrease of δD values in the first 50 yr, followed by a feeble increase smaller than the analytical precision. We have not found other studies that dealt with δD in larch

255

The effect of TRW has been studied for long and is well documented (e.g., Helama et al., 2017). After a very short (< 20 yr) period of increasing TRW values, they consistently decrease, yet with different rates. From 20 to 100 yr the decrease is rather steep thereafter changing to moderate rates. At around 300 yr of cambial age they are flattening out in both species. Understanding the absolute growth rate change is rather complex as discussed for instance in Matsushita et al. (2015). Dependencies of the age-size, growth-size and growth-age relationships are crucial. The fact that our database consists of trees from tree-line sites allows us to state that our derived trends are independent from the so-called crowding effect (influence of neighboring trees). Therefore, it represents mainly the individual variability and the age-size influence on the growth rate. We did not find any dependence of the trends in the different selected time frames within the past 9,000 yr, the behavior we observed should represent the general dependence of the age-size influence.

265

The decay of wood does not influence the carbon and oxygen stable isotope values of the cellulose (Nagavciuc et al. 2018), but that it can impact CC, since the cellulose is decomposed faster than lignin. Yet, it has been shown that CC has the potential to be used as a climate proxy (Ziehmer et al., 2018). In the analysis of tree-age related trends we have to consider that the decay of a trunk is not equal for all parts, and that hardwood in contrast to sapwood presents a decay resistance, which also varies from species to species (Kéérik, 1974). Both investigated species show a similar positive trend in approximately the first 50 yr of cambial age followed by a slight negative trend for larch or no overall trend for cembran pine.

270

This suggests, with reference to our data, that there are probably no (cembran pine) or possibly only minor (larch) influences due to effects of wood decay. This suggests that the CC variation in the first 50 yr is not due to wood decay, but rather a tree-ageing effect.

275

5 Conclusions

The present work confirms the absence of an ageing effect for all three stable isotopes after 100 yr of cambial age in the two conifer species, suggesting that the values older than 100 yr of cambial age can be considered for climate analyses without detrending. The exception is larch that shows a minor increase of δD mean values, smaller than the analytical precision. Before 100 yr the trends differ for each isotope and species, and only the larch $\delta^{13}C$ values can be used without detrending, since they do not vary with ageing. In both species, the $\delta^{18}O$ values present two phases, making the detrending rather challenging. It is similar for δD values in larch that change in the first 50 yr, whereas in cembran pine between 50-100 yr. Again detrending is demanding and should ideally be complemented by additional information regarding an explanation of this behavior. Tree ring cellulose contents show a significant trend for the first 50 years only, in contrast tree ring width curves flattening only after 300 year. Here the application of a regional curve standardization (RCS) is valuable. In summary, for climate reconstructions isotope data older than 100 cambial yr can be use directly, data of the first 100 yr should be used with caution. Therefore, data can be used only after detrending or when compared with data from other age classes covering the same time.

280
285
290

Acknowledgments.

We are grateful to Peter Nyfeler for the precious assistance during measurements of the stable isotopes, to Andrea Thurner and Andreas Österreicher for the preparation of the isotope samples from Alpine sites and the civil service collaborators: Lars Herrmann, Giacomo Ruggia, Jonathan Lamprecht, Yannick Rohrer, Rafael Zuber. The project is funded by the Swiss National Science Foundation (SNF 200021L_144255, SNF 200020_172550) as well as by the Austrian Science Fund (FWF, grant I-1183-N19) and is supported by the Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland (OCCR).\$

Competing interests. The authors declare that they have no conflict of interest

300 Author contribution statement

TA and MZ performed the stable isotope analyses, TA drafted the first version of the manuscript. KN collected the samples and made the crossdating. ML contributed to the evaluation of the results. ML, KN, CS conceived of the presented idea. All authors provided comments to improve the manuscript.

References

- 305 Anderson, W.T., Sternberg L., Pinzon M., Gann T. -Troxler. Childers, D.L and Duever M.: Carbon isotopic composition of cypress trees from South Florida and changing hydrologic conditions. *Dendrochronologia*. 23:1-10. <https://doi.org/10.1016/j.dendro.2005.07.006>, 2005
- Arosio, T., Ziehmer M. M., Nicolussi K., Schlüchter C., Leuenberger M.: Larch cellulose is significantly depleted in deuterium isotopes with respect to evergreen conifers – Submitted, 2020
- 310 Bräker, O.-U. der alterstrend bei jahringdichten und jahringbreiten von nadelhoebyzern und sein ausgleich, 1981
- Brienen, R.J.W., Gloor, E., Clerici, S., Newton, R., Arppe, L., Boom, A., Bottrell, S., Callaghan, M., Heaton, T., Helama, S. and Helle, G.: Tree height strongly affects estimates of water-use efficiency responses to climate and CO₂ using isotopes. *Nature Communications*, 8(1), pp.1-10. <https://doi.org/10.1038/s41467-017-00225-z>, 2017
- Coplen, T.B.. Reporting of stable hydrogen, carbon, and oxygen isotopic abundances (technical report). *Pure and Applied Chemistry*, 66(2), pp.273-276. <https://doi.org/10.1351/pac199466020273>, 1994
- 315 Daux, V., Edouard J., Masson-Delmotte V., Stievenard M., Hoffmann G., Pierre M., Mestre O., Danis P. and Guibal F... Can climate variations be inferred from tree-ring parameters and stable isotopes from *Larix decidua*? Juvenile effects, budmoth outbreaks, and divergence issue. *Earth Planetary Science Letters*. 309:221-233. <https://doi.org/10.1016/j.epsl.2011.07.003>, 2011
- 320 Duffy, J.E., McCarroll D., Barnes A., Ramsey C.B., Davies D., Loader N.J., Miles D. and Young G.H.. Short-lived juvenile effects observed in stable carbon and oxygen isotopes of UK oak trees and historic building timbers. *Chemical Geology*. 472:1-7. <https://doi.org/10.1016/j.chemgeo.2017.09.007>, 2017
- Duffy, J.E., McCarroll D., Loader N.J., Young G.H., Davies D., Miles D. and Bronk Ramsey C. Absence of age-related trends in stable oxygen isotope ratios from oak tree rings. *Global Biogeochemical Cycles*. 33:841-848. <https://doi.org/10.1029/2019GB006195>, 2019
- 325 Duquesnay, A., Breda N., Stievenard M. and Dupouey J.. Changes of tree-ring $\delta^{13}\text{C}$ and water-use efficiency of beech (*Fagus sylvatica* L.) in north-eastern France during the past century. *Plant, Cell Environment*. 21:565-572. <https://doi.org/10.1046/j.1365-3040.1998.00304.x>, 1998
- Esper, J., Frank, D.C., Battipaglia, G., Büntgen, U., Holert, C., Treydte, K., Siegwolf, R. and Saurer, M. Low-frequency noise in $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ tree ring data: A case study of *Pinus uncinata* in the Spanish Pyrenees. *Global Biogeochemical Cycles*, 24(4). <https://doi.org/10.1029/2010GB003772>, 2010
- 330

- Esper, J., Konter, O., Krusic, P.J., Saurer, M., Holzkämper, S. and Büntgen, U., Long-term summer temperature variations in the Pyrenees from detrended stable carbon isotopes. *Geochronometria*, 53:59, 42(1). DOI 10.1515/geochr-2015-0006, 2015
- 335 Esper, J., Klippel, L., Krusic, P.J., Konter, O., Raible, C.C., Xoplaki, E., Luterbacher, J. and Büntgen, U. Eastern Mediterranean summer temperatures since 730 CE from Mt. Smolikas tree-ring densities. *Climate Dynamics*, 54(3), 1367-1382, <https://doi.org/10.1007/s00382-019-05063-x>, 2020
- Filot, M.S., Leuenberger M., Pazdur A. and Boettger T. Rapid online equilibration method to determine the D/H ratios of non-exchangeable hydrogen in cellulose. *Rapid Communications in Mass Spectrometry: An International Journal Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry*. 20:3337-3344. <https://doi.org/10.1002/rcm.2743>, 2006
- 340 ~~*Rapid Communications in Mass Spectrometry: An International Journal Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry*~~. 20:3337-3344. <https://doi.org/10.1002/rcm.2743>, 2006
- Freyer, H.. On the ^{13}C record in tree rings. Part I. ^{13}C variations in northern hemispheric trees during the last 150 years. *J Tellus*. 31:124-137. <https://doi.org/10.3402/tellusa.v31i2.10417>, 1979
- Gagen, M., McCarroll D., Loader N.J., Robertson I., Jalkanen R. and Anchukaitis K. Exorcising the segment length curse: summer temperature reconstruction since AD 1640 using non-detrended stable carbon isotope ratios from pine trees in northern Finland. *The Holocene*. 17:435-446, <https://doi.org/10.1177/0959683607077012>, 2007
- 345 Gagen, M., McCarroll D., Robertson I., Loader N.J. and Jalkanen R.. Do tree ring $\delta^{13}\text{C}$ series from *Pinus sylvestris* in northern Fennoscandia contain long-term non-climatic trends? *Chemical Geology*. 252:42-51, <https://doi.org/10.1016/j.chemgeo.2008.01.013>, 2008
- 350 Helama, S., Melvin, T. M., and Briffa, K. R.: Regional curve standardization: State of the art, *Holocene*, 27, 172–177, <https://doi.org/10.1177/0959683616652709>, 2017
- Helama, S., Arppe L., Timonen M., Mielikäinen K. and Oinonen M.. Age-related trends in subfossil tree-ring $\delta^{13}\text{C}$ data. *Chemical Geology*. 416:28-35, <https://doi.org/10.1016/j.chemgeo.2015.10.019>, 2015
- Kéérik, A.A. Decomposition of wood. *Biology of plant litter decomposition*, 1, p.129, 1974
- 355 Kilroy, E., McCarroll D., Young G.H., Loader N.J. and Bale R.J. Absence of juvenile effects confirmed in stable carbon and oxygen isotopes of European larch trees. *Acta Silvae et Ligni*:27-33, DOI 10.20315/ASetL.111.3, 2016
- Klesse, S., Weigt R., Treydte K., Saurer M., Schmid L., Siegwolf R.T. and Frank D.C. Oxygen isotopes in tree rings are less sensitive to changes in tree size and relative canopy position than carbon isotopes. *Plant, cell environment*. 41:2899-2914, <https://doi.org/10.1111/pce.13424>, 2018
- 360 Körner, C. Alpine plant life: functional plant ecology of high mountain ecosystems; with 47 tables. ed. *Springer Science & Business Media*. Berlin - Heidelberg, DE, 2003
- Kress, A., Saurer, M., Siegwolf, R.T., Frank, D.C., Esper, J. and Bugmann, H. A 350 year drought reconstruction from Alpine tree ring stable isotopes. *Global Biogeochemical Cycles*, 24(2), <https://doi.org/10.1029/2009GB003613>, 2010

- Labuhn, I., Daux V., Pierre M., Stievenard M., Girardclos O., Féron A., Genty D., Masson-Delmotte V. and Mestre O. Tree age, site and climate controls on tree ring cellulose $\delta^{18}\text{O}$: A case study on oak trees from south-western France. *Dendrochronologia*. 32:78-89, <https://doi.org/10.1016/j.dendro.2013.11.001>, 2014
- Leavitt, S.W. Tree-ring C–H–O isotope variability and sampling. *Science of the Total Environment*. 408:5244-5253, 2010
- Leuenberger, M. To what extent can ice core data contribute to the understanding of plant ecological developments of the past? *Terrestrial ecology*. 1:211-233, <https://doi.org/10.1016/j.scitotenv.2010.07.057>, 2007
- 370 Li, Z.-H., Leavitt S.W., Mora C.I. and Liu R.-M. Influence of earlywood–latewood size and isotope differences on long-term tree-ring $\delta^{13}\text{C}$ trends. *Chemical geology*. 216:191-201, <https://doi.org/10.1016/j.chemgeo.2004.11.007>, 2005
- Lipp, J., Trimborn, P., Graff, W. and Becker, B., Climatic significance of D/H ratios in the cellulose of late wood in tree rings from spruce (*Picea abies* L.). In *Isotope techniques in the study of past and current environmental changes in the hydrosphere and the atmosphere*, 1993
- 375 Loader, N., Street-Perrott F., Daley T., Hughes P., Kimak A., Levanič T., Mallon G., Mauquoy D., Robertson I. and Roland T. Simultaneous Determination of Stable Carbon, Oxygen, and Hydrogen Isotopes in Cellulose. *Analytical chemistry*. 87:376-380, <https://doi.org/10.1021/ac502557x>, 2015
- Matsushita, M., Takata, K., Hitsuma, G., Yagihashi, T., Noguchi, M., Shibata, M. and Masaki, T. A novel growth model evaluating age–size effect on long-term trends in tree growth. *Functional Ecology*, 29(10), pp.1250-1259, <https://doi.org/10.1111/1365-2435.12416>, 2015
- 380 Mayr, C., Frenzel B., Friedrich M., Spurk M., Stichler W. and Trimborn P.. Stable carbon-and hydrogen-isotope ratios of subfossil oaks in southern Germany: methodology and application to a composite record for the Holocene. *The Holocene*. 13:393-402, <https://doi.org/10.1191/0959683603hl632rp>, 2003
- McCarroll, D. and Loader N.J.. Stable isotopes in tree rings. *Quaternary Science Reviews*. 23:771-801, <https://doi.org/10.1016/j.quascirev.2003.06.017>, 2004
- 385 McCarroll, D. and Pawellek F.. Stable carbon isotope ratios of *Pinus sylvestris* from northern Finland and the potential for extracting a climate signal from long Fennoscandian chronologies. *The Holocene*. 11:517-526, <https://doi.org/10.1191/095968301680223477>, 2001
- Monserud, R.A. and Marshall J.D.. Time-series analysis of $\delta^{13}\text{C}$ from tree rings. I. Time trends and autocorrelation. *Tree physiology*. 21:1087-1102, <https://doi.org/10.1093/treephys/21.15.1087>, 2001
- 390 Nagavciuc, V., Kern, Z., Perşoiu, A., Keşjár, D. and Popa, I. Aerial decay influence on the stable oxygen and carbon isotope ratios in tree ring cellulose. *Dendrochronologia*, 49, pp.110-117, <https://doi.org/10.1016/j.dendro.2018.03.007>. 2018
- Nagavciuc, V., Ionita M., Perşoiu A., Popa I., Loader N.J. and McCarroll D.. Stable oxygen isotopes in Romanian oak tree rings record summer droughts and associated large-scale circulation patterns over Europe. *Climate Dynamics*. 52:6557-6568, <https://doi.org/10.1007/s00382-018-4530-7>, 2019
- 395


- Nagavciuc, V., Kern, Z., Ionita, M., Hartl, C., Konter, O., Esper, J. and Popa, I., Climate signals in carbon and oxygen isotope ratios of Pinus cembra tree-ring cellulose from the Călimani Mountains, Romania. *International Journal of Climatology*, 40(5), pp.2539-2556, <https://doi.org/10.1002/joc.6349>, 2020
- 400 Nicolussi, K., Kaufmann M., Melvin T.M., Van Der Plicht J., Schiefling P. and Thurner A. A 9111 year long conifer tree-ring chronology for the European Alps: a base for environmental and climatic investigations. *The Holocene*. 19:909-920, <https://doi.org/10.1177/0959683609336565>, 2009
- Ott, E., Über die Abhängigkeit des Radialzuwachses und der Oberhohen bei Fichte und Larche von der Meereshöhe und Exposition im Lotschental. *J For Suisse*. 169:193, 1978
- 405 Raffalli-Delcerce, G., Masson-Delmotte V., Dupouey J., Stievenard M., Breda N. and Moisselin J. Reconstruction of summer droughts using tree-ring cellulose isotopes: a calibration study with living oaks from Brittany (western France). *Tellus B: Chemical Physical Meteorology*. 56:160-174, <https://doi.org/10.3402/tellusb.v56i2.16405>, 2004
- Reinig, F., Nievergelt D., Esper J., Friedrich M., Helle G., Hellmann L., Kromer B., Morganti S., Pauly M. and Sookdeo A... New tree-ring evidence for the Late Glacial period from the northern pre-Alps in eastern Switzerland. *Quaternary Science Reviews*. 186:215-224, <https://doi.org/10.1016/j.quascirev.2018.02.019>, 2018
- 410 Saurer, M., Cherubini P. and Siegwolf R. Oxygen isotopes in tree rings of Abies alba: The climatic significance of interdecadal variations. *Journal of Geophysical Research: Atmospheres*. 105:12461-12470, <https://doi.org/10.1029/2000JD900160>, 2000
- Saurer, M., Siegwolf R.T. and Schweingruber F.H.. Carbon isotope discrimination indicates improving water-use efficiency of trees in northern Eurasia over the last 100 years. *Global Change Biology*. 10:2109-2120, <https://doi.org/10.1111/j.1365-2486.2004.00869.x>, 2004
- 415 Suess, H.E.. Radiocarbon concentration in modern wood. *Science*. 122:415-417, 1955
- Treydte, K., Schleser G.H., Schweingruber F.H. and Winiger M. The climatic significance of $\delta^{13}\text{C}$ in subalpine spruces (Lötschental, Swiss Alps) a case study with respect to altitude, exposure and soil moisture. *Tellus* 53:593-611, <https://doi.org/10.1034/j.1600-0889.2001.530505.x>, 2001
- 420 Treydte, K.S., Schleser G.H., Helle G., Frank D.C., Winiger M., Haug G.H. and Esper J.. The twentieth century was the wettest period in northern Pakistan over the past millennium. *Nature Climate Change*. 440:1179, <https://doi.org/10.1038/nature04743>, 2006
- Young, G.H., Demmler J.C., Gunnarson B.E., Kirchhefer A.J., Loader N.J. and McCarroll D.. Age trends in tree ring growth and isotopic archives: A case study of Pinus sylvestris L. from northwestern Norway. *Global Biogeochemical Cycles*. 25:GB2020, <https://doi.org/10.1029/2010GB003913>, 2011
- 425 Ziehmer M.M., Nicolussi K., Schlüchter C. and Leuenberger M.. Preliminary evaluation of the potential of tree-ring cellulose content as a novel supplementary proxy in dendroclimatology. *Biogeosciences*. 15:1047-1064, <https://doi.org/10.5194/bg-15-1047-2018>, 2018

<u>SITE CODE</u>	<u>SITE NAME</u>	<u>SPECIES</u>	<u>N TREES</u>	<u>MEAN LENGTH (YR)</u>	<u>COORDINATES</u>	<u>ASPECT</u>	<u>ELEVATION (m)</u>
AHMO	Ahrntal, Moaralm	PICE	16	178	47°03'E/12°08'N	SE	1995
AHST	Ahrntal, Starklalm	LADE	2	125	47°05'E/12°11'N	S	2080
		PICE	2	93			
BIH	Paznaun, Bielerhöhe	PICE	1	190	46°91'E/10°10'N	N	2175
EBA	Ötztal, Ebenalm	PICE	11	196	47°01'E/10°95'N	NE	2115
FPCR	Val d'Hérens, Rezentproben Ferpecle	LADE	4	208	46°06'E/7°55'N	WSW	1965
G	Ötztal, Gurgler Zirbenwald	PICE	3	215	46°85'E/11°01'N	NW	2060
GDM	Kaunertal, Daunmoränensee	PICE	15	197	46°88'E/10°71'N	E	2295
GGUA	Ötztal, Gurgler Alm	PICE	3	160	46°85'E/11°N	W	2175
GLI	Kaunertal, Ombrometer	PICE	6	164	46°81'E/10°7'N	NE	2147.5
GP	Kaunertal, Gepatschferner	PICE	5	86	46°86'E/10°73'N	W	2167.5
HIB	Defereggental, Hirschbichl	PICE	1	180	46°9' E/12°25'N	E	2140
KOFL	Ahrntal, Kofler Alm	PICE	1	130	46°95'E/12°1'N	S	2177.5
LFS	Langtaufers, Sandbichl	PICE	2	148	46°81' E/10°7'N	NW	2335
MAZB	Vinschgau, Marzoneralm/B	LADE	6	162	46°58'E/10°95'N	N	2125
		PICE	2	198			
MAZC	Vinschgau, Marzoneralm/C	LADE	4	160	46°58'E/10°95'N	N	2120
MAZE	Vinschgau, Marzoneralm/E	LADE	5	188	46°58'E/10°95'N	N	2125
		PICE	8	124			
MAZF	Vinschgau, Marzoneralm/F	LADE	2	250	46°58'E/10°95'N	N	2105
		PICE	1	155			
MIS	Radurschltal, Miseri	LADE	1	125	46°9'E/10°61'N	N	2252.5
		PICE	1	275			
MM	Val d'Hérens, Glacier du Mont Mine	LADE	6	247	46°03'E/7°916'N	NNE	1995
		PICE	7	260			
MORT	Morteratschgletscher	PICE	3	253	46°41'E/9°933'N	W	2045
RT	Rojental	PICE	2	258	46°8'E/10°46'N	SE	2400
TAH	Passeier, Timmelal	PICE	3	195	46°9'E/11°13'N	S	2117.5
TSC	Val Roseg, Tschiervagletscher	LADE	7	206	46°4'E/9°88'N	NW	2162.5
		PICE	7	256			
UA	Haslital, Unteraargletscher	LADE	17	168	46°56'E/8°21'N	E	1950
		PICE	2	205			
UAZR	Haslital, Unteraargletscher, Rezentproben Nordufer Grimselstausee	PICE	4	173	46°56'E/8°28'N	SSE	1977
ULFI	Ultental, Fiechtsee	LADE	24	243	46°46'E/10°83'N	N	2110
		PICE	4	208			
UWBA	Ultental, Weißbrunnalm	LADE	2	200	46°46'E/10°81'N	NE	2330
VRR	Val Roseg, Rezentproben	LADE	4	219	46°43'E/9°85'N	E	2158.5
		PICE	5	209			

ZER	Mattertal, Zermatt, Findelengletscher	LADE	1	295	46°05'E/7°78'N	N	2315
		PICE	1	110			


435
440
445
450
455

Table 1. Characteristics of the sampling site and of the trees: 11 sites contain both larch and cembran pine samples, 2 sites contain only larch (LADE) specimens and the remaining sites only cembran pine (PICE)




460 **Figure 1.** Map of the location of the 29 sampling sites (© Google Maps 2020): They are situated in the Swiss, the Austrian and the Italian Alps. Information to each site is given in table 1.


A cambial age vs time



B sample number



465 **Figure 2. Cambial age and replication: A, graph of temporal distribution of the all trees and their cambial age. On the X axis the calendar age of the time series is displayed, that goes back to 9,000 yr, on the Y axis the cambial age. Each line represents a tree, in red the larch and in green the cembran pine. B, graph of the replication along the cambial age, the dotted line represents the threshold of 10 that we have considered for the analysis.**



470

Figure 3. Analysis of $\delta^{13}\text{C}$ data: Panel A, normalized $\delta^{13}\text{C}$ values of all larch (LADE, red) and cembra pine (PICE, green) trees, green line corresponds to the mean. Panel B, raw (left) and normalized (right) mean value with corresponding ± 1 standard deviation (grey area), of larch (red) and cembra pine (cyan). Panel C, plots of the mean values with linear approximations for periods 1-100 yr and >100 yr.

475






Figure 4. Analysis of $\delta^{18}\text{O}$ data: Panel A, normalized $\delta^{18}\text{O}$ values of all larch (LADE, red) and cebran pine (PICE, green) trees, green line corresponds to the mean. Panel B, raw (left) and normalized (right) mean value with corresponding ± 1 standard deviation (grey area), of larch (red) and cebran pine (cyan). Panel C, plots of the mean values with linear approximations for periods 1-50 yr, to 51-100 yr and >100 yr.




485


A All trees




B Raw values



Normalized




C Mean values




490 Figure 6. Analysis of Tree Ring Width (TRW) data: Panel A, raw TRW values of all larch (LADE, red) and cembran pine (PICE, green) trees, green line corresponds to the mean. Panel B, raw (left) and normalized (right) mean value with corresponding ± 1 standard deviation (grey area), of larch (red) and cembran pine (cyan). Panel C, plots of the mean values with linear approximations for periods 1-50 yr, to 50-100 yr, to 101-300 yr and >300 yr..


A All trees




B Raw values



Normalized



C Mean values



495 Figure 7. Analysis of Cellulose Content (CC) data: Panel A, raw value of all larch (LADE, red) and cembran pine (PICE, green) trees, green line corresponds to the mean. Panel B, raw (left) and normalized (right) mean value with corresponding ± 1 standard deviation (grey area), of larch (red) and cembran pine (cyan). Panel C, plots of the mean values with linear approximations for periods 1-50 yr and >50 yr.