

Response to Reviewers' Comments

We appreciate the efforts of the reviewers for their insightful and constructive comments. We have addressed concerns in the previous round of review. Below, we provide detailed, point-by-point responses to each of the reviewers' comments. We put the reviewer comments in regular font, author responses in blue, and direct quotes from the revised manuscript *in italic*.

Associate Editor Decision: Reconsider after major revisions (30 Sep 2020) by Yakov Kuzyakov

Comments to the Author:

Dear Authors,

Please incorporate the suggestions of the reviewers in the new version.

Sincerely yours,

Yakov Kuzyakov

Response:

Thanks for handling and reviewing the manuscript. We have incorporated all the suggestions of the reviewers into the revised version.

Associate Editor Decision: Reconsider after major revisions (13 Aug 2020) by Yakov Kuzyakov

Comments to the Author:

Dear Authors,

The reviewers made many useful suggestions and raised some critical points. please incorporate their suggestions and consider the critics

Sincerely yours,

Yakov Kuzyakov

Reviewer #1' comments:

The manuscript ran detailed models to understand the effects of vegetation shift from grassland to woody plants on CaCO₃ dissolution. The manuscript is well written and the conclusions are acceptable. Nevertheless, the model and calculations are far from my expertise and I am unable to judge them.

Response:

We appreciate the reviewer's comments.

Reviewer #2' comments:

The authors study the difference between chemical weathering in grassland and woodland soils through the application of simulations of a model calibrated in the grassland soil region. They aimed to understand the effect of deepening roots on carbonate weathering. In general, the manuscript is well written and structured; the geochemical model and simulation results are well described, using kinetic and thermodynamic equations. It deserves to be published because it deals with some relevant aspects that should be considered in studies dealing with the critical zone, which is the weathering of carbonate minerals and its link with the ecosystem; it also discusses the effects on the global carbon cycle and regional long-term simulations.

Nevertheless, from what the title of the manuscript implies, I was expecting an extensive discussion about the deepening roots and their connection with weathering of carbonates, with a well representation in the geochemical model. The authors did not create a model where it explicitly represent the rhizosphere nor biogeochemical processes besides respired CO₂, despite that the last one was found to be one of the most important variables controlling weathering of carbonates. From my perspective, the deepening roots in the model were represented as pure hydrological feature, their only influence the flow path but not the flow itself, which can vary from plant species (not all plants will have the same water requirements). Although they discussed most of the before mentioned issues under model limitations, this subsection was somehow unclear. I would suggest a revision of the objectives or the model limitations in order to publish the manuscript.

Response:

We appreciate the reviewer's encouraging comments. We agreed with the reviewer that the model does not explicitly include all processes related to deepening roots. To recognize this point and to avoid misleading the readers, we have changed the title to "Deepening roots can enhance carbonate weathering by amplifying CO₂-enriched recharge".

We also cleared the objectives and added more discussions about the model limitations, Line 82-84, 201-205, and 526-534:

"We recognize that rooting characteristics can have multiple influences on water flow paths and water budget, for example, via water uptake and transpiration (Sadras, 2003; Fan et al., 2017; Pierret et al., 2016). This study focuses primarily on their potential influence via the alteration of hydrological flow paths."

"We compared the relative significance of the hydrological (i.e., lateral versus vertical flow partitioning) vs. respiratory (i.e., CO₂ generation) influences of deepening roots. Though other potential differences might be induced by deepening roots (e.g., water uptake, water table, and transpiration), we assumed they remain constant across the grassland and woodland simulations to examine the relative influences of flow path vs. CO₂ generation on carbonate weathering"

"In addition to the alterations of hydrological flow paths, deepening roots may also affect the proportion of plant water uptake or soil water loss through transpiration (Pierret et al., 2016; Zhu et al., 2018), further modifying water fluxes into deep subsurface layers. Representing these dynamics requires a large number of processes that we do not have data to constrain. For example, studies of semiarid woodlands show that woody species predominantly use deep subsurface water while grasses predominantly use soil water from the upper soil layer (Ward et al., 2013). Furthermore, the grassland, with its shallower root distributions, may be more

efficient in using water from small rainfall events than forests with their deeper root distributions (Mazzacavallo and Kulmatiski, 2015). Other studies have also documented significant competition for water uptake in upper soil layers among both woody and grass species (Scholes and Archer, 1997). It remains inconclusive how these water uptake characteristics are best represented in numerical experiments. These processes are therefore not included at this point.”

Specific comments:

Why did the authors not include organic acids in the model or in the discussion? Is it because a small effect on weathering is expected?

Response:

We added the discussion about organic acids, Line 535-545:

“In addition, distributions of microbes and organic acids associated with rooting structures vary with sites and seasons, and root channels and dry conditions with less soil water may trigger calcite re-precipitation. Microbial activities surrounding living and dead roots can also lead to calcite precipitation and infilling of fractures and other macropores, and alter flow pathways (Lambers et al., 2009). Organic acids can decrease soil water pH, increase mineral solubilities through organic-metal complexations, and accelerate chemical weathering (Pittman and Lewan, 2012; Lawrence et al., 2014). Their impacts on carbonate weathering kinetics might be smaller (due to the fast kinetics) compared to their alteration of calcite solubility in natural systems. Such processes further complicate how to represent biogeochemical processes in models. Although we do not explicitly simulate all of these simultaneous and competing processes in our numerical experiments, the model was constrained by the field data in grasslands and woodland that have already integrated these effects in natural systems. Indeed, the weathering rates can be understood as the net weathering rates that are the net difference between dissolution and reprecipitation. This is reflected in lower carbonate weathering rates under low infiltration (low flow) conditions.”

The long-term simulations displays a change in porosity, is this change considered in the short-term model?

Response:

Yes, this is also considered in the short-term model. However, the dissolved calcite volume in the short-term numerical experiments (Scenarios 1-3) is less than 0.5% v/v and has minimal impact on permeability and water-rock contact time in these short-term numerical experiments. We clarified the changes of the solid phase in Line 291-292 and 305-306:

“Note that numerical experiments in Scenario 1-3 focused on the short-term scale with negligible changes in the solid phase.”

“The dissolved mineral volume was negligible (< 0.5% v/v).”

The calibration was done using grassland data, but the water uptake of plants depends also on the species, might it be the case that the roots take more water in woodland soils?

Response:

We agreed that the water uptake may change with plant species and added discussions about the potential impact, Line 526-534:

“In addition to the alterations of hydrological flow paths, deepening roots may also affect the proportion of plant water uptake or soil water loss through transpiration (Pierret et al., 2016;Zhu et al., 2018), further modifying water fluxes into deep subsurface layers. Representing these dynamics require a large number of processes that we do not have data to constrain. For example, studies of semiarid woodlands show that woody species predominantly use deep subsurface water while grasses predominantly use soil water from the upper soil layer (Ward et al., 2013). Furthermore, the grassland, with its shallower root distributions, may be more efficient in using water from small rainfall events than forests with their deeper root distributions (Mazzacavallo and Kulmatiski, 2015).Other studies have also documented significant competition for water uptake in upper soil layers among both woody and grass species (Scholes and Archer, 1997). It remains inconclusive how these water uptake characteristics are best represented in numerical experiments. These processes are therefore not included at this point .”

Under model limitations, the flow partitioned is well explained, however, the points given above are not present, including the difference in water balance (water uptake by plants) among species.

Response:

We added more discussions about the organic acid and other potential differences among species in the section of model limitations, Line 526-534 and 535-545:

“In addition to the alterations of hydrological flow paths, deepening roots may also affect the proportion of plant water uptake or soil water loss through transpiration (Pierret et al., 2016;Zhu et al., 2018), further modifying water fluxes into deep subsurface layers. Representing these dynamics requires a large number of processes that we do not have data to constrain. For example, studies of semiarid woodlands show that woody species predominantly use deep subsurface water while grasses predominantly use soil water from the upper soil layer (Ward et al., 2013). Furthermore, the grassland, with its shallower root distributions, may be more efficient in using water from small rainfall events than forests with their deeper root distributions (Mazzacavallo and Kulmatiski, 2015).Other studies have also documented significant competition for water uptake in upper soil layers among both woody and grass species (Scholes and Archer, 1997). It remains inconclusive how these water uptake characteristics are best represented in numerical experiments. These processes are therefore not included at this point .”

“In addition, distributions of microbes and organic acids associated with rooting structures vary with sites and seasons, and root channels and dry conditions with less soil water may trigger calcite re-precipitation. Microbial activities surrounding living and dead roots can also lead to calcite precipitation and infilling of fractures and other macropores, and alter flow pathways (Lambers et al., 2009). Organic acids can decrease soil water pH, increase mineral solubilities through organic-metal complexations, and accelerate chemical weathering (Pittman and Lewan, 2012;Lawrence et al., 2014). Their impacts on carbonate weathering kinetics might be smaller (due to the fast kinetics) compared to their alteration of calcite solubility in natural systems. Such

processes further complicate how to represent biogeochemical processes in models. Although we do not explicitly simulate all of these simultaneous and competing processes in our numerical experiments, the model was constrained by the field data in grasslands and woodland that have already integrated these effects in natural systems. Indeed, the weathering rates can be understood as the net weathering rates that are the net difference between dissolution and reprecipitation. This is reflected in lower carbonate weathering rates under low infiltration (low flow) conditions.”

Technical comments:

Units are usually separated from the number, including in percentage or permit;

Response:

We modified the units through the whole manuscript.

line 56-57, I did not understand completely this sentence;

Response:

We modified the sentence, Line 60-61.

“In grasslands, the lateral, dense spread of roots in upper soil layers promotes the formation of horizontally-oriented macropores that support near-surface lateral flow (Cheng et al., 2011).”

figure S1 should include coordinates in x and y axis;

Response:

We included coordinates in Figure S1.

line 198, the symbol '>' may be changed into words;

Response:

We modified it into words, Line 210-211:

“Deeper roots in woodlands can increase deep soil permeability by over one order of magnitude (Vergani and Graf, 2016).”

the unit for year is commonly written as 'a' not 'yr'.

Response:

We revised the unit of year into 'a' through the whole manuscript.

Reviewer #3' comments:

This manuscript deals with a topic of paramount importance in the present context of soil C research: the potential consequences of land-use changes (in this case, wood encroachment in grassland areas) in the soil inorganic C dynamics. It offers a comprehensive perspective and succeeds in providing some simulated data not only on the potential consequences of such change on soil CO₂ and water flows, but also on their relative importance as drivers of carbonates weathering in the study area. In this sense, it offers, within the limitations of the modelling approach considered and the available data, a quantitative view of the potential consequences of the mid-term changes in soil pCO₂ and water infiltration when vegetation changes in a soil with calcite accumulation at depth.

In my view, the manuscript is well written, describes well a complicated numerical simulation (the materials in SI are very helpful), and contains interesting information, and a territory-based approach that makes it worth of interest.

The contribution is therefore meaningful and adds to a field where more information is needed. The quality of Figures is very high, both in terms of graphical design and the information they provide.

Response:

We appreciate the encouraging comments and the constructive suggestions below from the reviewer.

There are however some points that were not completely clear to me while reading the manuscript, and therefore I consider that they could be complemented for a clearer understanding. I cannot add much to the thermodynamical considerations in the construction of the model, but, as a soil scientist, there are some questions that came to my mind while revising the manuscript.

- In relation to the soil profile description and the gradient in calcite concentration depicted in Figure 1 (“The gray color gradient reflects the calcite abundance with more calcite in depth”) and Figure 3.b, it is not clear for the reader which was the actual calcite distribution with depth. Only in the supplementary information (Table S1) it is said that the calcite volumetric concentration (m³/m³) was 0 in the upper soil horizon (0-54 cm). In line 324, the calcite-no calcite interface is set at 55 cm. If this is correct, this makes that in the model representing the grassland situation, where 95% of the flow was considered to “exit the soil column to the stream” at 50 cm (line 166-167), there would be almost no contact between soil water and calcite accumulated in the soil profile below 54 cm. Indeed (line 359), in the HF simulations, it is said that water “bypassed the calcite zone”. I understand this can represent the actual situation in the study site, but, if this was the case, I believe this should be more clearly stated in the description of the site and the model: deeper rooting with wood encroachment would actually take soil water enriched in CO₂ to the calcite accumulation areas in the soil profile, where otherwise (grassland), only a very limited amount of it would reach. I also wonder how realistic this can be if the actual distribution of rainfall along the year would be considered (for instance, intense storm episodes or wet vs. dry season rainfall patterns).

Response:

We have Figure 3A showing the depth profile of calcite volume used in all numerical experiments, consistent with the values listed in Table S1. We added clarifications in Line 227:

“b. Calcite distribution (black line) in all cases increases with depth; The values are listed in Table S1 and shown in Figure 3A.”

We also added clarifications about how water flows through the deep calcite-abundant subsurface in the grassland and woodland cases, Line 73-75 and 109-112:

“We hypothesized that deepening roots in woodlands enhance carbonate weathering by promoting deeper recharge and CO₂-carbonate contact in the deep, carbonate-abundant subsurface (Figure 1).”

“The shallow and dense fine roots in the grassland promote lateral macropore development and lateral water flow. In contrast, the woodlands induce vertical macropore development that supports vertical flow (recharge) into the deep, calcite-abundant subsurface compared to the grassland.”

This work aimed to compare the general, averaged behaviors rather than event-scale dynamics. We agreed that the actual distribution of rainfall may affect how and to what extent water reacts with calcite, and added discussions, Line 521-526:

“In natural systems, however, other factors can also influence flow partitioning. Some of these are not explicitly represented in our modeling exercises. For example, contrasts in flow-conducting properties (i.e., porosity and permeability) in shallow and deep zones, physical and chemical heterogeneity in carbonate distribution (Wen and Li, 2018), connectivity between different areas of the catchment in dry and wet times (Wen et al., 2020), and water table and corresponding lateral flow depth associated with the rainfall frequency and intensity (Li et al., 2017; Harman and Cosans, 2019). All these factors may affect the water-calcite contact extent, leading to the changes of carbonate weathering rates (Figure 6).”

- In this sense, there is some confusion about the type of materials water enters in contact with at depth. While the abstract talks about “carbonate rocks” (line. 27), suggesting the parent material, the description of the soil profile (Table S1) denotes the horizon at 146-188 cm as “B” and that at 99-145 cm as “AB”. This would mean that these would be accumulation horizons of materials leached from the upper “A” horizon (0-54 cm). If I understood well this table, their mineral composition according to this table indicates that calcite was not the dominant mineral in any of them.

Response:

We clarified it and added more descriptions about the mineralogy in the methodology, Line 27-29, 89-93 and 165-167:

“These differences in weathering fronts are ultimately caused by the contact time of CO₂-charged water with abundant carbonate minerals in the deep subsurface.”

“The bedrock contains repeating Permian couplets of limestone (1-2 m thick) and mudstone (2-4 m) (Macpherson et al., 2008). The limestone is primarily calcite with traces of dolomite while the mudstone is dominated by illite, chlorite, and mixed-layers of chlorite-illite and chlorite-vermiculite, varying in abundance from major to trace amounts. With an average thickness of 1-2 m in the lowlands, the soil mostly has a carbonate content of less than 25% (Macpherson et al., 2008).”

“In the model, the upper soil layers have more anorthite ($\text{CaAl}_2\text{Si}_2\text{O}_8(\text{s})$) and K-feldspar ($\text{KAlSi}_3\text{O}_8(\text{s})$) and the deeper subsurface contains more calcite (Table S1). The calcite volume increases from 0% in the upper soil layer to 10% in the deep subsurface.”

- In relation to this, a point that is also not clear to me in the manuscript, is to what point re-precipitation of dissolved calcite at depth could determine some of the consequences of increased calcite dissolution with deeper woodland root systems. This would affect both the final fate of Ca and DIC in groundwater, and the assumptions done on the changes in permeability due to increased porosity with calcite depletion (lines 385-394). It is known that in this type of soils, carbonates can re-precipitate for instance around root channels, where water concentration can decrease rapidly compared to the bulk soil matrix.

Response:

We agree that calcite reprecipitation can happen. And they often happen under dry conditions where soils are dried out. This work focuses on average behavior and the first-order principles of hydrological ramification of roots. The model does not explicitly simulate details of root channels and dry conditions that may trigger calcite re-precipitation. However, the model was constrained by the field data in grasslands and woodland that have already integrated these effects in natural systems. In fact, the weathering rates can be understood as the net weathering rates that are the net difference between dissolution and reprecipitation. That is why carbonate weathering rates are lower under low infiltration (low flow) conditions. We discussed the potential impacts of calcite re-precipitation in the section of model limitations, Line 535-545:

“In addition, distributions of microbes and organic acids associated with rooting structures vary with sites and seasons, and root channels and dry conditions with less soil water may trigger calcite re-precipitation. Microbial activities surrounding living and dead roots can also lead to calcite precipitation and infilling of fractures and other macropores, and alter flow pathways (Lambers et al., 2009). Organic acids can decrease soil water pH, increase mineral solubilities through organic-metal complexations, and accelerate chemical weathering (Pittman and Lewan, 2012; Lawrence et al., 2014). Their impacts on carbonate weathering kinetics might be smaller (due to the fast kinetics) compared to their alteration of calcite solubility in natural systems. Such processes further complicate how to represent biogeochemical processes in models. Although we do not explicitly simulate all of these simultaneous and competing processes in our numerical experiments, the model was constrained by the field data in grasslands and woodland that have already integrated these effects in natural systems. Indeed, the weathering rates can be understood as the net weathering rates that are the net difference between dissolution and reprecipitation. This is reflected in lower carbonate weathering rates under low infiltration (low flow) conditions.”

- In relation to the annual average CO₂(g) and CO₂ (aq) concentrations used in the numerical simulations (Table 1), and the idea explained in l.310-315 that some “impositions” on time and depth distribution of soil CO₂ had to be done to the model to capture the variation in alkalinity and Ca data and different horizons, I wonder to what extent this can be considered in the discussion about the use of annual averages to explain processes that can be very dependent of monthly (or even daily) fluctuations (in addition to spatial heterogeneity).

Response:

The numerical experiments in this work focus on the general behaviors using the annual average CO₂(g), which is the dominant driver of Ca and DIC concentrations in soil water. At daily or monthly scales, the variations of rainfall frequency and intensity may change the flow conditions (e.g., water table and lateral flow depth), leading to different extent of water-calcite contact extent and weathering rates. We added discussions about it, Line 521-526:

“In natural systems, however, other factors can also influence flow partitioning. Some of these are not explicitly represented in our modeling exercises. For example, contrasts in flow-conducting properties (i.e., porosity and permeability) in shallow and deep zones, physical and chemical heterogeneity in carbonate distribution (Wen and Li, 2018), connectivity between different areas of the catchment in dry and wet times (Wen et al., 2020), and water table and corresponding lateral flow depth associated with the rainfall frequency and intensity (Li et al., 2017; Harman and Cosans, 2019). All these factors may affect the water-calcite contact extent, leading to the changes of carbonate weathering rates (Figure 6).”

- Finally, a small comment on the term “shallow soil”. It is used repeatedly in the manuscript to refer in fact to the model where grassland roots and subsurface waterflow are considered (in contrast to woodland roots and waterflow). For example, in l. 319 (“soil CO₂ production rate was the highest in the shallow soil”). In my understanding, the soil considered in the model has the same depth in all cases (Figure 1). This makes the term “shallow” (vs. deep?) misleading. I’d suggest to revise this point and either change the term “shallow” and/or to use it to qualify the upper soil layer or the waterflow, not the soil.

Response:

We appreciate this comment. We have modified the terms into “upper soil layer” to avoid confusion.

Reviewer #4! comments:

Compared to soil organic carbon, inorganic carbon in soil (SIC) is usually ignored in global carbon cycle. Increasing evidences indicates that SIC also plays important role in global carbon cycle. Therefore, this manuscript deals with a very interesting topic, i.e., effect of plants on carbonate weathering. However, I’m not an expert on model, and could not review it. After reading it, I feel that this manuscript has presented a very clear concept on the factors that control carbonate weathering, including temperature, hydrological regimes, and soil CO₂ concentration. I just have a comment on it. The deep root systems of plants in some regions (e.g., semi-arid, and semi-humid) may make the deep soil dry, due to the strong transpiration.

Response:

We thank the reviewer for reading the manuscript. We added discussions about water uptake, Line 526-534:

“In addition to the alterations of hydrological flow paths, deepening roots may also affect the proportion of plant water uptake or soil water loss through transpiration (Pierret et al., 2016; Zhu et al., 2018), further modifying water fluxes into deep subsurface layers. Representing these dynamics requires a large number of processes that we do not have data to constrain. For example, studies of semiarid woodlands show that woody species predominantly use deep subsurface water while grasses predominantly use soil water from the upper soil layer (Ward et al., 2013). Furthermore, the grassland, with its shallower root distributions, may be more efficient in using water from small rainfall events than forests with their deeper root distributions (Mazzacavallo and Kulmatiski, 2015). Other studies have also documented significant competition for water uptake in upper soil layers among both woody and grass species (Scholes and Archer, 1997). It remains inconclusive how these water uptake characteristics are best represented in numerical experiments. These processes are therefore not included at this point .”

Reviewer #5' comments:

The manuscript titled “Deepening roots can enhance carbonate weathering” by Wen et al. documents the role that landscape changes (specifically changes from grasslands to woodlands) play in the dissolution of carbonate host rock. The manuscript is well written and addresses the important topic of flow partitioning- vs. soil CO₂-driven weathering in carbonate terrain. The authors ran detailed model simulations to address their research questions:

1. “How and to what degree do rooting characteristics influence carbonate weathering when considering both flow partitioning and soil CO₂ distribution?”
2. “Which factor (flow partitioning or soil CO₂ distribution) predominantly controls weathering?”

From the title and abstract, I was expecting more discussion of rooting depth and its connection to weathering (as another reviewer points out). However, the paper focuses on roots only as they relate to hydrologic flowpaths. Thus, much of the content and modeling is outside my area of expertise so I do not feel I can adequately judge the conclusions of the paper.

Response:

We thank the reviewer for reading the manuscript. Based on comments from all of the reviewers, including Reviewer #5, we have chosen to change the title of the manuscript:

“Deepening roots can enhance carbonate weathering by amplifying CO₂-enriched recharge”

Deepening roots can enhance carbonate weathering by amplifying CO₂-enriched recharge

Hang Wen¹, Pamela L. Sullivan², Gwendolyn L. Macpherson³, Sharon A. Billings⁴, Li Li¹

¹ Department of Civil and Environmental Engineering, Pennsylvania State University, University Park PA 16802, United States

5 ² College of Earth, Ocean, and Atmospheric Science, Oregon State University, Corvallis OR 97331, United States

³ Department of Geology, University of Kansas, Lawrence KS 66045, United States

⁴ Department of Ecology and Evolutionary Biology and Kansas Biological Survey, University of Kansas, Lawrence KS 66045, United States

Correspondence to: Li Li (lili@engr.psu.edu)

10 **Abstract:** Carbonate weathering is essential in regulating atmospheric CO₂ and carbon cycle at the century time scale. Plant roots accelerate weathering by elevating soil CO₂ via respiration. It however remains poorly understood how and how much rooting characteristics (e.g., depth and density distribution) modify flow paths and weathering. We address this knowledge gap using field data from and reactive transport numerical experiments at the Konza Prairie Biological Station (Konza), Kansas (USA), a site where woody encroachment into grasslands is surmised to deepen roots.

Deleted: have been known to

15 Results indicate that deepening roots can enhance weathering in two ways. First, deepening roots can control thermodynamic limits of carbonate dissolution by regulating how much CO₂ transports vertical downward to the deeper carbonate-rich zone. The base-case data and model from Konza reveal that concentrations of Ca and dissolved inorganic Carbon (DIC) are regulated by soil pCO₂ driven by the seasonal soil respiration. This relationship can be encapsulated in equations derived in this work describing the dependence of Ca and DIC on temperature and soil CO₂, which applies in multiple carbonate-dominated catchments. Second, numerical experiments show that roots control weathering rates by regulating recharge (or, vertical water fluxes) into the deeper carbonate zone and export reaction products at dissolution equilibrium. The numerical experiments explored the potential effects of partitioning 40% of infiltrated water to depth in woodlands compared to 5% in grasslands. Soil CO₂ data from wood- and grasslands suggest relatively similar soil CO₂ distribution over depth, and only led to 1% to 12% difference in weathering rates if flow partitioning was kept the same between the two land covers. In contrast, deepening roots can enhance

Deleted: potentially

Deleted: Dissolved Inorganic

Deleted: fluctuation of

Deleted: has been shown to apply

Deleted: the amount of

Deleted: that flush through

Deleted: Numerical

25 weathering by 17% to 207% as infiltration rates increased from 3.7×10^{-2} to 3.7 m/a. These cases are about 1,200 % larger than a case without roots at all, underscoring the essential role of roots in general. Numerical experiments also indicate that weathering fronts in woodlands propagated > 2 times deeper compared to grasslands after 300 years at an infiltration rate of 0.37 m/a. These differences in weathering fronts are ultimately caused by the contact time of CO₂-charged water with abundant carbonate minerals in deep subsurface. We recognize that modeling results are subject to limitations in representing processes and parameters, but these data and numerical experiments prompt the hypothesis that 1) deepening roots in woodlands can enhance carbonate weathering by promoting recharge and CO₂-carbonate contact in the deep, carbonate-abundant subsurface; and 2) the hydrological impacts of rooting characteristics can be more influential than those of soil CO₂ distribution in modulating weathering rates. We call for co-located characterizations of roots, subsurface structure, and soil CO₂ levels, and their linkage to water and water chemistry. These measurements will be essential to illuminate feedback mechanisms of land cover changes, chemical weathering, global carbon cycle, and climate.

Deleted: yr.

Deleted: indicated

Deleted: the

Deleted: yr

Deleted: rocks

Deleted: we propose that the

Deleted: allude to

Deleted: ;

Deleted: ,

Deleted: improve models and

Carbonate weathering has long been considered negligible as a long-term control of atmospheric CO₂ (0.5 to 1 Ma; Berner and Berner, 2012; Winnick and Maher, 2018). Recent studies, however, have underscored its significance in controlling the global carbon cycle at the century time scale that is relevant to modern climate change, owing to its rapid dissolution, fast response to perturbations, and the order of magnitude higher carbon store in carbonate reservoirs compared to the atmosphere (Gaillardet et al., 2019). Carbonate weathering is influenced by many factors, including temperature (Romero-Mujalli et al., 2019b), hydrological regimes (Romero-Mujalli et al., 2019a; Wen and Li, 2018), and soil CO₂ concentrations (Covington et al., 2015) arising from different vegetation types (Calmels et al., 2014). Rapid alteration to any of these factors, either human or climate induced, may change global carbonate weathering fluxes and lead to a departure from the current global atmospheric CO₂ level. This is particularly important given that about 7–12 % of the Earth's continental area is carbonate based and about 25% of the global population completely or partially depend on waters from karst aquifers (Hartmann et al., 2014).

Plant roots have long been recognized as a dominant biotic-driver of chemical weathering and the global carbon cycle (Berner, 1992; Beerling et al., 1998; Brantley et al., 2017a). The growth of forests has been documented to elevate soil pCO₂ and amplify Dissolved Inorganic Carbon (DIC) fluxes (Berner, 1997; Andrews and Schlesinger, 2001). Rooting structure can influence weathering in two ways. First, rooting systems (e.g., grass-, shrub- and woodlands) may affect the distribution of soil carbon (both organic and inorganic), microbe biomass, and soil respiration (Drever, 1994; Jackson et al., 1996; Billings et al., 2018). The relatively deep root distributions of shrublands compared to grasslands may lead to deeper soil carbon profiles (Jackson et al., 1996; Jobbagy and Jackson, 2000), which may help elevate deep CO₂ and acidity that determine carbonate solubility and weathering rates.

Second, plant roots may affect soil structure and hydrological processes. Root trenching and etching can develop porosity (Mottershead et al., 2003; Hasenmueller et al., 2017). Root death and decay can promote the generation of macropores, or more specifically, biopores with connected networks (Angers and Caron, 1998; Zhang et al., 2015). Root channels have been estimated to account for about 70 % of the total described macropores (Noguchi et al., 1997; Beven and Germann, 2013) and for over 70 % of water fluxes through soils (Watson and Luxmoore, 1986). In grasslands, the lateral, dense spread of roots in upper soil layers promotes the formation of horizontally-oriented macropores that support near-surface, lateral flow (Cheng et al., 2011). Highly dense fine roots also increase the abundance of organic matter and promote granular or “sandy” texture soil aggregates that facilitate shallow, near-surface water flow (Oades, 1993; Nippert et al., 2012). In contrast, in shrublands and forests, generally deeper and thicker roots tend to promote a high abundance of macropores and high connectivity to the deep subsurface (Canadell et al., 1996; Nardini et al., 2016), enhancing the drainage of water to the depth (Pawlik et al., 2016).

It is generally known that rooting characteristics vary among plant species and are critical in regulating water budgets, flow paths and storage (Sadras, 2003; Nepstad et al., 1994; Jackson et al., 1996; Cheng et al., 2011; Brunner et al., 2015; Fan et al., 2017). Existing studies however have primarily focused on the role of soil CO₂ and organic acids (Drever, 1994; Lawrence et al., 2014; Gaillardet et al., 2019; Hauser et al., 2020). Systematic studies on coupled effects of hydrological flow paths, flow partitioning and soil CO₂ distribution are missing, owing to the limitation in data that detail rooting effects on flow partitioning and complex hydrological-biogeochemical interactions. Here we ask the questions: how and to what degree do rooting characteristics influence carbonate weathering when considering both flow partitioning and soil CO₂ distribution? Which factor (flow partitioning or soil CO₂ distribution) predominantly controls weathering? We hypothesized that deepening roots in woodlands enhance carbonate weathering by promoting deeper recharge and CO₂-carbonate contact in deep, carbonate-abundant subsurface (Figure 1).

Moved down [1]: about 25% of the global population completely or partially depend on waters from karst aquifers (Hartmann et al., 2014).

Deleted: About 7–12% of the Earth's continental area is carbonate based;

Deleted: Myr

Deleted: type

Moved (insertion) [1]

Deleted: at depth

Deleted: >

Deleted: top soils promote horizontal

Deleted: ,

Deleted: with deep

Deleted: thick

Deleted: have

Deleted: proportion

Deleted: at depth


We tested the hypotheses by a series of numerical experiments integrating reactive transport modelling and water chemistry data from an upland watershed in the Konza Prairie Biological Station, a tallgrass prairie and one of the Long-Term Ecological Research (LTER) sites in the US (Figure S1). We used the calibrated model to carry out numerical experiments for two end-members of vegetation covers, grasslands and woodlands, under flow-partitioning conditions that are characteristic of their rooting structure. These experiments differentiated the impact of biogeochemical and hydrological drivers and bracketed the range of their potential impacts on weathering, thus providing insights on the missing quantitative link between rooting structure, flow paths, and chemical weathering. We recognize that rooting characteristics can have multiple influences on water flow paths and water budget, for example, via water uptake and transpiration (Sadras, 2003; Fan et al., 2017; Pierret et al., 2016). This study focuses primarily on their potential influence via the alteration of hydrological flow paths.

2 Research site and data sources

Site description. Details of the Konza site are in the Supporting Information (SI) and references therein. Here we provide brief information relevant to this work. Konza is a mesic native grassland where experimentally manipulated, long-term burning regimes have led to woody encroachment in up to 70% of the catchment area in some catchments. The mean annual temperature and precipitation are 13°C and ~ 835 mm, respectively (Tsypin and Macpherson, 2012). The bedrock contains repeating Permian couplets of limestone (1-2 m thick) and mudstone (2-4 m) (Macpherson et al., 2008). The limestone is primarily calcite with traces of dolomite while the mudstone is dominated by illite, chlorite, and mixed-layers of chlorite-illite and chlorite-vermiculite, varying in abundance from major to trace amounts. With an average thickness of 1-2 m in the lowlands, soils mostly have carbonate less than 25% (Macpherson et al., 2008). Data suggest that the Konza landscape is undergoing a hydrogeochemical transition that coincides with and may be driven by woody-encroachment. In parallel to these changes is a detectable decline in stream flow and an increase in weathering rates (Macpherson and Sullivan, 2019) and groundwater $p\text{CO}_2$ (Macpherson et al., 2008). Stream water concentration-discharge (C-Q) patterns have exhibited chemodynamic patterns (i.e., solute concentrations are sensitive to changes in discharge) for geogenic species (e.g., Mg and Na) in woody-encroached sites compared to grass sites. Sullivan et al. (2019) hypothesized that concentration-discharge relationships may be affected by woody species with deeper roots, which altered flow paths and mineral-water interactions (Figure 1). We focus on the upland watershed N04d in Konza (Figure S1) that has experienced a 4-year burning interval since 1990 and has seen considerable woody encroachment and changes in hydrologic fluxes (Sullivan et al., 2019).

Data sources. Daily total meteoric precipitation and evapotranspiration were from the Konza data website (ter.konza.ksu.edu/data). Wet chemistry deposition data was from the National Atmospheric Deposition Program NADP (nadp.sws.uiuc.edu). Monthly data of soil gases (16, 84 and 152 cm), soil water (17 and 152 cm), and groundwater (366 cm) from 2009-2010 were used for this work (Tsypin and Macpherson, 2012) (Figure 2A). The sampling points were about 30 m away from the stream. More information on field and laboratory methods were included in the SI.

Deleted: most



145 Figure 1. A conceptual diagram of hydro-biogeochemical interactions in the grassland (A) and woodland (B). The
 150 shallow and dense fine roots in the grassland **promote** lateral macropore development and lateral water flow. **In
 contrast,** the **woodlands induce** vertical macropore development that supports vertical flow (**recharge**) into **the deep,
 calcite-abundant subsurface compared to the grassland**. The gray color gradient reflects the calcite abundance with
 more calcite in depth. M_L and M_G with a unit of mol/a ($= \text{flow rate } Q (\text{m/a}) \times \text{species concentration } C (\text{mol/m}^3) \times \text{unit}$
 cross-section area (m^2)) represent mass fluxes from lateral flow (soil water) and from vertical flow into groundwater,
 respectively. Soil water and groundwater were assumed to eventually flow into stream, **where** Infiltration rate $Q_T = Q_L$
 + Q_G **is the same as discharge that counts for both lateral and vertical water flow.**

3 Reactive transport modeling

3.1 Base case: 1D reactive transport model for the Konza grassland (calibrated with field data)

155 A 1-D reactive transport model was developed using the code CrunchTope (Steeffel et al., 2015). The code solves mass
 conservation equations integrating advective and diffusive / dispersive transport and geochemical reactions. It has been extensively
 used in understanding mineral dissolution, chemical weathering, and biogeochemical reactions (e.g., Lawrence et al. (2014); Wen
 et al. (2016); Deng et al. (2017)). In this study, the base case had a porosity of 0.48 and a depth of 366.0 cm at a resolution 1.0 cm.
 Soil temperature was assumed to decrease linearly from 17 °C at the land surface to 8 °C at 366.0 cm, within typical ranges of
 160 field measurements (Tsy-pin and Macpherson, 2012). Detailed set up of domain, soil mineralogy, initial condition, and precipitation
 chemistry, are in the SI.

Representation of soil CO₂. The model does not explicitly simulate soil respiration (microbial activities and root respiration) that
 produces soil CO₂. Instead, it approximates these processes by having a solid phase “CO₂(g*)” that continuously releases CO₂(g)
 that dissolves into CO₂(aq) (and other relevant aqueous species) at a kinetic rate constant of 10⁻⁹ mol/m²/s (Reaction 0-1 in Table
 165 1) that reproduced the observed soil pCO₂ levels. This value is at the low end of the reported soil respiration rates (10⁻⁹ ~ 10⁻⁵
 mol/m²/s) (Bengtson and Bengtsson, 2007; Ahrens et al., 2015; Carey et al., 2016). The dissolution of CO₂(g*) via CO₂(g) into
 CO₂(aq) follows Henry’s law $C_{\text{CO}_2(\text{aq})} = K_1 p\text{CO}_2$. Here K_1 is the equilibrium constant of Reaction 1, **which equals to Henry’s law**
 constant. The extent of CO₂(g*) dissolution (i.e., the mass change of the solid phase over time) was constrained by $C_{\text{CO}_2(\text{aq})}$, which
 was estimated using temperature-dependent K_1 (following the van’t Hoff equation in Eq. S1) and measured soil CO₂ data at different

- Formatted: Font color: Text 1
- Deleted: promotes
- Deleted: ;
- Formatted: Font color: Text 1
- Deleted: groundwater
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: deeper roots in the woodland induces...
- Formatted: Font color: Text 1
- Formatted: Font color: Text 1
- Deleted: yr
- Deleted: yr
- Deleted: .

Deleted: essentially

horizons (Table 2). These $C_{CO_2(aq)}$ values were then linearly interpolated for individual grid blocks in the model. Finally, these prescribed $C_{CO_2(aq)}$ values were used as the equilibrium constants of the coupled Reaction 0-1 in the form of $CO_2(g^*) \leftrightarrow CO_2(aq)$, such that the soil CO_2 at different depth were represented (Section 4.1). In the base case, the soil profile of $C_{CO_2(aq)}$ values were updated monthly based on the monthly soil CO_2 data (Table 2). More details about the implementation in CrunchTope are included in the SI.

Table 1. Key reactions and kinetic and thermodynamic parameters

Reaction	$\log_{10}K_{eq}$ at 25 °C ^a	Standard enthalpy (ΔH° , kJ/mol)	$\log_{10}k$ (mol/m ² /s) at 25 °C ^d	Specific Surface Area (m ² /g) ^d
Soil CO₂ production through CO₂(g*) and dissolving into CO₂(aq)				
(0) $CO_2(g^*) \leftrightarrow CO_2(g)$	-	-	-9.00	1.0
(1) $CO_2(g) \leftrightarrow CO_2(aq)$	-1.46 ^b	-19.98	-	-
(2) $CO_2(aq) + H_2O \leftrightarrow H^+ + HCO_3^-$	-6.35	9.10	-	-
(3) $HCO_3^- \leftrightarrow H^+ + CO_3^{2-}$	-10.33	14.90	-	-
Chemical weathering				
(4) $CaCO_3(s) + CO_2(aq) + H_2O \leftrightarrow Ca^{2+} + 2HCO_3^-$	-5.12 ^c -4.52 ^c	-15.41	-6.69	0.84
(5) $CaAl_2Si_2O_8(s) + 8H^+ \leftrightarrow Ca^{2+} + 2Al^{3+} + 2H_4SiO_4(aq)$	26.58	-	-11.00	0.045
(6) $KAlSi_3O_8(s) + 4H^+ + 4H_2O \leftrightarrow Al^{3+} + K^+ + 3H_4SiO_4(aq)$	-0.28	-	-12.41	0.20
(7) $Al_2Si_2O_5(OH)_4(s) + 6H^+ \leftrightarrow 2Al^{3+} + 2H_4SiO_4(aq) + H_2O$	6.81	-	-12.97	17.50

- 185 a. Values of K_{eq} were interpolated using the EQ3/6 database (Wolery et al., 1990), except Reaction 1 and 4 (i.e., K_1 and K_4).
- b. By combining the temporal- and spatial-dependent soil pCO_2 data, $CO_2(aq)$ concentrations were calculated through $C_{CO_2(aq)} = K_1 pCO_2$. The prescribed $C_{CO_2(aq)}$ values were used as equilibrium constants in CrunchTope to describe how much soil CO_2 was available for weathering.
- 190 c. Calcite in the **upper** soil (above Horizon B) is mixed with other minerals and has small particle size (Macpherson and Sullivan, 2019), relatively “impure” and therefore has a lower K_{eq} value than those at depth with relatively “pure” calcite. The K_{eq} of the “impure” calcite was calibrated by fitting field data of Ca and alkalinity.
- d. The kinetic rate parameters and specific surface areas were from Palandri and Kharaka (2004), except Reaction 0. The kinetic rate constant of the solid phase $CO_2(g^*)$ dissolution (i.e., soil CO_2 production rate constant) was from (Bengtson and Bengtsson, 2007; Ahrens et al., 2015; Carey et al., 2016); The specific surface area was referred to that of soil organic carbon (Pennell et al., 1995).
- 195 e. These reactions were only used in the base case model as they occurred in **upper** soils in Konza. In the later numerical experiments, these reactions were not included so as to focus on carbonate weathering.

Deleted: shallow

Deleted: shallow

Table 2. Measured $CO_2(g)$ at different depths and corresponding estimated $C_{CO_2(aq)}$

Time	Soil depth				Ways obtained ^{a,b,c}	
I. Grassland (Konza) cases						
	Horizon A (h = 16 cm)	Horizon AB (84 cm)	Horizon B (152 cm)	Groundwater (366 cm)		
Soil T °C	17	15	13	8	Estimated	
K_1	4.1×10^{-2}	4.4×10^{-2}	4.6×10^{-2}	5.4×10^{-2}	Estimated	
1. Base case with monthly updated $CO_2(g)$ (%) and $CO_2(aq)$ (mol/L)						
$CO_2(g)$	July	3.6	6.8	6.6	2.2	Measured
	August	1.4	1.7	7.2	3.9	Measured
	September	0.6	1.2	3.9	4.9	Measured
	October	0.5	1.4	2.5	5.0	Measured
	November	0.6	1.1	2.2	4.0	Measured
	January	0.3	0.8	1.1	3.6	Measured
	March	0.2	0.3	0.5	3.0	Measured
$CO_2(aq)$	July	1.5×10^{-3}	3.0×10^{-3}	3.0×10^{-3}	1.2×10^{-3}	Estimated
	August	5.7×10^{-4}	7.5×10^{-4}	3.3×10^{-3}	2.1×10^{-3}	Estimated
	September	2.5×10^{-4}	6.2×10^{-4}	1.8×10^{-3}	2.6×10^{-3}	Estimated

	October	1.9×10^{-4}	6.2×10^{-4}	1.2×10^{-3}	2.7×10^{-3}	Estimated
	November	2.5×10^{-4}	4.8×10^{-4}	1.0×10^{-3}	2.2×10^{-3}	Estimated
	January	1.2×10^{-4}	3.5×10^{-4}	5.1×10^{-4}	1.9×10^{-3}	Estimated
	March	8.2×10^{-5}	1.3×10^{-4}	2.3×10^{-4}	1.6×10^{-3}	Estimated
2. Numerical experiments with annual average $CO_2(g)$ (%) and $CO_2(aq)$ (mol/L)						
$CO_2(g)$	Annual	1.0 ± 1.2	1.9 ± 2.2	3.4 ± 2.6	3.8 ± 1.0	Measured
$CO_2(aq)$	Annual	$(4.2 \pm 5.0) \times 10^{-4}$	$(8.5 \pm 9.7) \times 10^{-4}$	$(1.6 \pm 1.2) \times 10^{-3}$	$(2.0 \pm 0.5) \times 10^{-3}$	Estimated

II. Woodland cases (Calhoun, South Carolina)

Numerical experiments with annual average $CO_2(g)$ (%) and $CO_2(aq)$ (mol/L)

	(h = 50 cm)	(150 cm)	(300 cm)	(500 cm)	
Soil T °C	16	13	10	8	Estimated
K_1	4.2×10^{-2}	4.6×10^{-2}	4.9×10^{-2}	5.4×10^{-2}	Estimated
$CO_2(g)$	Annual	0.9 ± 0.1	2.7 ± 0.6	3.8 ± 0.6	Measured
$CO_2(aq)$	Annual	$(4.5 \pm 0.7) \times 10^{-4}$	$(1.3 \pm 0.3) \times 10^{-3}$	$(1.9 \pm 0.3) \times 10^{-3}$	Estimated

a. Monthly measured soil CO_2 data for the Konza grassland (base case) were from (Tsy-pin and Macpherson, 2012); The annual-average soil CO_2 used in the grassland experiments was averaged from the corresponding monthly measurements. More information on measurements at the Konza grassland were detailed in the SI. The annual-average soil CO_2 in the woodland experiments was from the Calhoun site in South Carolina where trees (forest) are dominant (Billings et al., 2018).

b. $CO_2(aq)$ were estimated using Henry's law $C_{CO_2(aq)} = K_1 pCO_2$; The temperature-dependent K_1 was calculated following Eq. S1. The $C_{CO_2(aq)}$ values at different soil depths were used to prescribe the available soil CO_2 for chemical weathering.

c. Soil temperature was estimated from the soil water and shallow groundwater temperature (Tsy-pin and Macpherson, 2012; Billings et al., 2018).

Reactions. In the model, the upper soil layers have more anorthite ($CaAl_2Si_2O_8(s)$) and K-feldspar ($KAlSi_3O_8(s)$) and the deeper subsurface contains more calcite (Table S1). The calcite volume increases from 0 % in the upper soil layer to 10 % in the deep subsurface. Table 1 summarizes reactions and thermodynamic and kinetic parameters. Soil CO_2 increases pore water acidity (Reactions 0-2) and accelerates mineral dissolution (Reactions 3-6). Silicate dissolution leads to the precipitation of clay (represented by kaolinite in Reaction 7). These reactions were included in the base case to reproduce field data. The kinetics follows the transition state theory (TST) rate law (Plummer et al., 1978) $r = kA \left(1 - \frac{IAP}{K_{eq}}\right)$, where k is the kinetic rate constant ($mol/m^2/s$), A is the mineral surface area per unit volume (m^2/m^3), IAP is the ion activity product, and K_{eq} is the equilibrium constant. The term IAP/K_{eq} quantifies the extent of disequilibrium: values close to 0 suggest far from equilibrium whereas values close to 1.0 indicate close to equilibrium.

Flow partitioning. Rain water entered soil columns at the annual infiltration rate of $0.37 m/a$, estimated based on the difference between measured precipitation ($0.88 m/a$) and evapotranspiration ($0.51 m/a$). At 50 cm, a lateral flow Q_L (soil water) exited the soil column to the stream at $0.35 m/a$. The rest recharged the deeper domain beyond 50 cm (to the groundwater system) at $0.02 m/a$ (~ 2 % of precipitation) and became groundwater (Figure 1A), a conservative value compared to 2-15 % reported in another study (Steward et al., 2011). The groundwater flow Q_G eventually came out at 366.0 cm and was assumed to enter the stream as part of discharge ($= Q_L + Q_G$). The flow field was implemented in CrunchTope using the "PUMP" option.

Calibration. We used monthly measured alkalinity and Ca concentration data in different horizons (Horizon A, B and groundwater in Figure 2A) from 2009-2010 at the Konza grassland for model calibration. The monthly Nash-Sutcliffe efficiency (NSE), which quantified the residual variance of modeling output compared to measurements, was used for model performance evaluation (Moriassi et al., 2007). NSE values higher than 0.5 are considered acceptable. At Konza, calcite in the upper soil (above Horizon B) is mixed with other minerals, has small particle size, and is considered "impure" (Macpherson and Sullivan, 2019) and therefore

Deleted: The shallow soils

Formatted: Font color: Text 1

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: shallow

240 has a lower K_{eq} value than those at depth with relatively “pure” calcite. The K_{eq} of “impure” calcite (K_I) was calibrated by fitting field data of Ca and alkalinity.

3.2 Numerical experiments

Numerical experiments were set up to understand how and to what extent roots regulate weathering rates and solute transport in grasslands and in woodlands. The base case from Konza was used to represent grasslands, and the Calhoun site (the Calhoun Critical Zone Observatory in South Carolina, USA) was used as a representative woody site (Billings et al., 2018).
245 Grasslands are typically characterized by a high proportion of horizontal macropores induced by dense, lateral-spread of roots mostly at depths less than 0.8 m (Jackson et al., 1996; Frank et al., 2010). These characteristics promote lateral flow (Q_L) at the shallow subsurface (Figure 1A). At Konza, ~~over 90 %~~ of grass roots were at the top 0.5 m, leading to high hydrologic conductivity in top soils (Nippert et al., 2012). In the woodland, a greater proportion of deep roots enhances vertical macropore development (Canadell et al., 1996; Nardini et al., 2016), reduces permeability contrasts at different depths (Vergani and Graf, 2016), and is
250 thought to facilitate more vertical water flow to the depth (Q_G). At the Calhoun site, ~~over 50 %~~ of roots are in the top 0.5 m in woodlands, with the rest penetrating deeper (Jackson et al., 1996; Eberbach, 2003; Billings et al., 2018).

The experiments aimed to compare the general, averaged behaviors rather than event-scale dynamics so the annual-average soil CO_2 data and corresponding prescribed $CO_2(aq)$ concentrations were used (Table 2). The experiments focused on calcite weathering (Reaction 0-4) and excluded silicate weathering reactions (Reaction 5-7). The mineral-dissolution-related
255 parameters from the base case were used for all experiments. ~~We compared the relative significance of the hydrological (i.e., lateral versus vertical flow partitioning) vs. respiratory (i.e., CO_2 generation) influences of deepening roots. Though other potential differences might be induced by deepening roots (e.g., water uptake, water table, and transpiration), we assumed they remain constant across the grassland and woodland simulations to examine the relative influences of flow path vs. CO_2 generation on carbonate weathering.~~

260 Hydrological and biogeochemical differences in grasslands and woodlands. Flow partitioning between lateral shallow flow and vertical recharge flow is challenging to quantify and is subject to large uncertainties under diverse climate, lithology, and land cover conditions. The ratios of lateral flow in ~~upper~~ soils versus the total flow inferred from a tracer study in a grassland vary from ~70 % to ~95 % (Weiler and Naef, 2003). Harman and Cosans (2019) found that the lateral flow rate at ~~upper~~ soils over the overall infiltration can vary between 50 % and 95 %. Deeper roots in woodlands can increase deep soil permeability by ~~over one~~
265 order of magnitude (Vergani and Graf, 2016). Assuming that the vertical, recharged flow water ultimately leaves the watershed as baseflow, the ratio of the lateral versus vertical flow has been reported with a wide range. In forests such as Shale Hills in Pennsylvania and Coal Creek in Colorado, ~7 %-20 % of stream discharge is from groundwater, presumably recharged by vertical flow (Li et al., 2017; Zhi et al., 2019). Values of Q_G/Q_T estimated through base flow separation vary from 20 % to 90 % in forest/wood-dominated watersheds (Price, 2011), often negatively correlating with the proportion of grasslands (Mazvimavi et al.,
270 2004).

Soil respiration rates can vary between $10^{-9} \sim 10^{-5}$ mol/m²/s for both grasslands and woodlands (Bengtson and Bengtsson, 2007; Ahrens et al., 2015; Carey et al., 2016). Soil CO_2 levels may vary by 2-3 orders of magnitude depending on vegetation type and climate conditions (Neff and Hooper, 2002; Breecker et al., 2010). Soil CO_2 levels are further complicated also by their
275 dependence on topographic position, soil depth, and soil moisture, all of which determine the magnitude of microbial and root activities and CO_2 diffusion (Hasenmueller et al., 2015; Billings et al., 2018). There is however no consistent evidence suggesting

Deleted: >

Deleted: >

Deleted: We compared the relative significance of hydrological and biogeochemical effects (soil CO_2 level and distribution) of rooting depths.

Deleted: shallow

Deleted: shallow

Deleted: >

285 which land cover exhibits higher soil respiration rate or soil CO₂ level. Below we describe details of the numerical experiments (Table 3) exploring the influence of hydrological versus biogeochemical impacts of roots.

Table 3. Physical and geochemical characteristics in numerical experiments^a

Cases	Calcite distribution ^b	Schematic	Hydrological and biogeochemical aspects	
			Flow partitioning ^c	Soil CO ₂ distribution ^d
Scenario 1 with flow partitioning (PF)	GrassPF		$Q_L = 95\% Q_T$	Annual average from the Konza grassland (red line)
	WoodPF		$Q_L = 60\% Q_T$ $Q_G = 40\% Q_T$ (blue arrow)	Annual average from the Calhoun forest (blue line)
Scenario 2 with 100% vertical flow (VF)	GrassVF		$Q_L = 0$	Annual average from the Konza grassland (red line)
	WoodVF		$Q_G = Q_T$ (black arrow)	Annual average from the Calhoun forest (blue line)
Scenario 3 with 100% lateral flow (HF)	GrassHF		$Q_L = Q_T$	Annual average from the Konza grassland (red line)
	WoodHF		$Q_G = 0$ (black arrow)	Annual average from the Calhoun forest (blue line)

- 290 a. Diagrams represent the soil profiles of physical and geochemical properties;
 b. Calcite distribution (black line) in all cases increases with depth; **The values are listed in Table S1 and shown in Figure 3A.**
 c. The flow field was implemented in CrunchTope using the “PUMP” option;
 d. Annual-average soil CO₂ used for constraining the grassland (red line) was from the Konza grassland (Tsy-pin and Macpherson, 2012). The annual-average soil CO₂ in the woodland (blue line) was from the Calhoun forest (Billings et al., 2018).

295 **Three scenarios.** Each scenario in Table 3 includes a grassland and woodland case, with their respective profiles of calcite distribution and soil pCO₂ kept the same (column 3 and 4 in Table 3). The only difference in different scenarios is the flow partitioning (column 5 in Table 3). Soil pCO₂ in the grassland (red line in Table 3) were set to reflect the annual average of the Konza site. Soil pCO₂ (blue line) in the woodland is the annual average from the forest-dominant Calhoun site (Billings et al., 2018). In all scenarios, we assumed that grasslands and woodlands had the same total solid phase “CO₂(g*)” producing CO₂ gas and CO₂(aq) but differed in depth distributions. The CO₂(g*) served as the source of soil CO₂ and was constrained by CO₂(g) field data (Table 2). In grasslands, the modeled distribution of CO₂(g*) was steep, with higher density of roots and more abundant CO₂(g*) in the **upper** soils; while the distribution was less steep in woodlands (Jackson et al., 1996). Below the rooting depth, CO₂(g*) was assumed to be smaller (by 10 times) to represent the potential soil CO₂ sources from microbial activities (Billings et al., 2018).

305 Scenario 1 considered flow partitioning (Table 3). With the large permeability contrast of soil and bedrock (over four orders of magnitude) in Konza (Macpherson, 1996), the GrassPF case (with $Q_L = 95\% \times Q_T$ and $Q_G = 5\% \times Q_T$) represents an end-member case for the grassland. The woodland (WoodPF) case was set to have 60% lateral flow and 40% vertical flow into deeper subsurface. This groundwater percentage is at the high end of flow partitioning and serves as an end-member case for woodlands (Vergani and Graf, 2016). Scenarios 2 and 3 had no flow partitioning. Scenario 2 had two cases with 100% vertical flow via the bottom outlet (VF; WoodVF and GrassVF); Scenarios 3 had two cases with 100% horizontal flow (HF; WoodHF and GrassHF) via the shallow outlet at 50 cm. These cases represent the end-member flow cases with 100% lateral flow or 100% vertical flow. In addition, because the two cases have the same flow scheme, they enable the differentiation of effects of soil CO₂ distribution versus hydrology differences. All scenarios were run under infiltration rates from 3.7×10^{-2} to 3.7 m/a ($10^{-4} - 10^2$

Deleted: ¶

Deleted: shallow

Formatted: Not Highlight

Deleted: % ×

Deleted: % ×

Deleted: yr

m/day), the observed daily variation range at Konza. This was to explore the role of flow regimes and identify conditions where most and least significant differences occur. To reproduce the observed soil CO₂ profile, the soil CO₂ production rate (mol/m²/a) at the domain scale (calculated by the mass change of the solid phase CO₂(g*) over time) was assumed to increase with infiltration rates (Figure S2). This is consistent with field observations that soil CO₂ production rate and efflux may increase with rainfall in grassland and forest ecosystems (Harper et al., 2005; Patrick et al., 2007; Wu et al., 2011; Vargas et al., 2012; Jiang et al., 2013). For example, Zhou et al. (2009) documented soil CO₂ production rates increasing from 3.2 to 63.0 mol/m²/a when the annual precipitation increased from 400 to 1,200 mm. In addition, Wu et al. (2011) showed that increasing precipitation from 5 to 2,148 mm enhanced soil respiration by 40% and that a global increase in 2 mm precipitation per decade may lead to an increase of 3.8 mol/m²/a for soil CO₂ production. The simulated soil CO₂ production rates across different infiltration rates here (~0.1-10 mol/m²/a) shown in Figure S2) were close to the reported belowground net primary production (belowground NPP) of typical ecosystems: ~0.8-100 mol/m²/a for grasslands (Gill et al., 2002) and ~0.4 - 40 mol/m²/a for woodlands (Aragao et al., 2009).

Each case was run until steady state, when concentrations at the domain outlet became constant (within ±5%) over time. The time to reach steady state varied from 0.1 to 30 a, depending on infiltration rates. The lateral flux (soil water, $M_L=Q_L \times C_L$) and vertical fluxes (groundwater, $M_G=Q_G \times C_G$) were calculated at 50 and 366 cm, in addition to total fluxes (weathering rates, M_T). These fluxes multiplied with unit cross-section area (m²) convert into rates in units of mol/a.

Carbonate weathering over century time scales: soil property evolution. To compare the propagation of weathering fronts over longer time scales, we carried out two 300-a simulations for Scenario 1 with flow partitioning under the base-case infiltration rate of 0.37 m/a (i.e., Grass_{PF} and Wood_{PF} in Table 3). During this long-term simulation, we updated calcite volume, porosity and permeability. The calcite volume changes were updated in each time step based on corresponding mass changes, which were used to update porosity. Permeability changes were updated based on changes in local porosity following the Kozeny-Carman equation:

$\frac{k_i}{k_{i,0}} = \left(\frac{\phi_i}{\phi_{i,0}}\right)^3 \times \left(\frac{1-\phi_{i,0}}{1-\phi_i}\right)^2$ (Kozeny, 1927; Costa, 2006), where k_i and ϕ_i is permeability and porosity in grid i at time t , and $k_{i,0}$ and $\phi_{i,0}$ are the initial permeability and porosity, respectively.

3.3 Quantification of weathering rates and their dependence on CO₂-carbonate contact

To quantify the overall weathering rates and CO₂-carbonate contact in each scenario, we used the framework from a previously-developed upscaled rate law for dissolution of spatially heterogeneously distributed minerals (Wen and Li, 2018). The rate law says that three characteristics times are important. The equilibrium time τ_{eq} represents the characteristic timescale of mineral dissolution to reach equilibrium in a well-mixed system. The residence time τ_a , i.e., the timescale of advection, quantifies the overall water contact time with the whole domain. It was calculated by the product of domain length (L) and porosity divided by the overall infiltration rate (Q_T): $\tau_a = \frac{L\phi}{Q_T}$. The reactive transport time $\tau_{ad,r}$ quantified the water contact time with calcite as influenced by both advection and diffusion/dispersion. The upscaled rate law is as follows:

$$R_{calcite} = k_{calcite} A_T \left[1 - \exp\left(-\frac{\tau_{eq}}{\tau_a}\right)\right] \left\{1 - \exp\left[-L\left(\frac{\tau_a}{\tau_{ad,r}}\right)\right]\right\}^\alpha \quad (1)$$

Here k is the intrinsic rate constant measured for a mineral in a well-mixed reactor, A_T is the total surface area, L is domain length, α is geostatistical characteristics of spatial heterogeneity, and $\frac{\tau_a}{\tau_{ad,r}}$ is the reactive time ratio quantifying the relative magnitude of the water contact time with the whole domain versus the contact time with the reacting mineral. This rate law consists of two parts: the effective dissolution rates in homogeneous media represented by $k A_T \left[1 - \exp\left(-\frac{\tau_{eq}}{\tau_a}\right)\right]$ and the heterogeneity factor

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yrs

Deleted: yr

Deleted: yr

Deleted: yr

370 $\left\{1 - \exp\left[-L\left(\frac{\tau_a}{\tau_{ad,r}}\right)\right]\right\}^\alpha$ that quantifies effects of preferential flow paths arising from heterogeneous distribution of minerals. When $\frac{\tau_a}{\tau_{ad,r}} > 1$, the water contact time with calcite zone is small, meaning the water is replenished quickly compared to the whole domain, leading to higher CO₂-carbonate interactions. In contrast, small $\frac{\tau_a}{\tau_{ad,r}}$ ratio (<1) reflects water is replenished slowly in the reactive calcite zones, leading to less CO₂-carbonate contact. These different time scales were calculated for Scenarios 1-3 based on the flow characteristics and dissolution thermodynamics and kinetics, as detailed in the SI. Values of τ_{eq} , τ_a , and $\tau_{ad,r}$ for all experiments are listed in Table S2. Note that numerical experiments in Scenario 1-3 focused on the short-term scale, with negligible changes in the solid phase.

4 Results

4.1 The thermodynamics of carbonate dissolution: grassland at Konza as the base case scenario

380 The calibrated model reproduced the temporal dynamics with a Nash-Sutcliffe efficiency (NSE) value > 0.6 and was considered satisfactory (Figure 2). Note that y axis is inverted to display upper soils at the top and deep soils at the bottom to be consistent with their subsurface position shown in Figure 2A. The measured CO₂(g) varied between 0.24 – 7.30% (Figure 2B right axis), one to two orders of magnitude higher than the atmospheric level of 0.04%. The estimated CO₂(aq) (Figure 2B left axis) generally increased with depth except in July and August when horizon B was at peak concentration. The timing of the peaks and valleys varied in different horizons. The CO₂(aq) reached maxima in summer in soil horizons A and B and decreased to Jess than 0.5 mM in winter and spring. The groundwater CO₂(aq) exhibited a delayed peak in September and October, and dampened seasonal variation compared to soil horizons. The temporal trends of alkalinity, DIC, and Ca in groundwater mirrored those of soil CO₂ at their corresponding depths, indicating the predominate control of soil CO₂ on carbonate weathering. The groundwater concentrations of these species were also higher than soil concentrations. The simulated groundwater DIC (approximately summation of CO₂(aq) and alkalinity) was > 6 times higher than that in upper soil (~ 1.0 mM). The dissolved mineral volume was negligible for the simulation period (< 0.5% v/v). Sensitivity analysis revealed that changes in flow velocities influenced concentrations in Horizon A with anorthite as the dominating dissolving mineral (0 – 1.8 m); their effects are negligible in Horizon 390 B and groundwater where fast-dissolving calcite has reached equilibrium.

Deleted: shallow

Deleted: <

Deleted: shallow




Figure 2. (A) Schematic representation of sampling depths in the Konza grassland; Corresponding monthly dynamics of (B) $\text{CO}_2(\text{aq})$ and $\text{CO}_2(\text{g})$, (C) alkalinity, (D) DIC, and (E) Ca concentrations. Lines represent modelling outputs at the corresponding sampling depth of monthly field measurements (dots), including soil water at Horizons A (16 cm) and B (152 cm), and groundwater (366 cm). The lines of $\text{CO}_2(\text{aq})$ and $\text{CO}_2(\text{g})$ in Figure B overlapped. Note that there are no DIC data so no dots in Figure D. The temporal trends of alkalinity, DIC, and Ca mirrored those of soil CO_2 , indicating its predominant control on weathering. Note that concentrations on y-axis from B – E are low to high, to be consistent with the layout in A with deeper water chemistry showing up deeper.

Several measurements/parameters were important in reproducing data, especially soil CO_2 that determined the $\text{CO}_2(\text{aq})$ level and its spatial variation, and equilibrium constant (K_{eq}) of calcite dissolution. Imposition of monthly variations and depth distributions of soil CO_2 were essential to capture the variation of alkalinity and Ca data at different horizons. The imposition of calcite K_{eq} was also critical for reproducing Ca concentrations. Impurities were suggested to affect K_{eq} of natural calcite by a factor of ~ 2.0 (Macpherson and Sullivan, 2019). Calcite K_{eq} in upper soils had to be reduced by a factor of 3.8 in the model to reproduce concentrations in Horizon A. The alkalinity and Ca concentrations were not sensitive to kinetic parameters nor precipitation, because carbonate dissolution was fast and thermodynamically controlled.

4.2 Numerical experiments: the significance of potential hydrology differences

Scenario 1 for hydro-biogeochemical effects with flow partitioning (Wood_{PF} and Grass_{PF}). Figures 3A1-E1 show depth profiles of calcite and soil CO_2 production rates and steady-state concentrations of reaction products. Although the model did not explicitly simulate soil respiration, the soil CO_2 production rate was highest in the upper soil at around $10^{-6.5}$ mol/s and decreased to $\sim 10^{-9.5}$ mol/s at 366 cm (Figure 3B1), consistent with the decline with soil depth observed in natural systems. The $\text{CO}_2(\text{g})$ level (released from the solid phase $\text{CO}_2(\text{g}^*)$) increased with depth due to autotrophic and heterotrophic respiration and downward fluxes of CO_2 -charge water from the upper soil (Figure 3C1). Concentrations of reaction products (Ca, DIC) were lower in the top 40 cm reflecting the lower carbonate-mineral background level and higher at depths over 60 cm. The transition occurred between 35 cm – 60 cm at the vicinity of calcite-no calcite interface at 55 cm, where concentrations of Ca and DIC increased abruptly until reaching equilibrium. This thin transition was driven by fast calcite dissolution and rapid approach to equilibrium, resulting in a short equilibrium distance. The equilibrated DIC and Ca concentrations below 60 cm followed the similar increasing trend of $\text{CO}_2(\text{g})$ with depth in the deep zone (Figure 3C1-E1).

Deleted: shallow


Deleted: shallow

Deleted: shallow


Deleted: >

Formatted: Not Highlight

430 Figures 3F1 show that Ca concentrations in soil water (light color) were lower than groundwater (dark color) and varied with infiltration rates. The difference between soil water and groundwater Ca concentrations was relatively small at low infiltration rates because both reached equilibrium but diverged at high infiltration rates. Higher infiltration rates diluted soil water but not as much for groundwater. As expected, stream concentrations, a mixture of soil water and groundwater (solid line), were in between these values but closely resembled soil water in the grassland. In both cases, stream concentration decreased as infiltration increased, indicating a dilution concentration-discharge relationship.



435 Figure 3. Simulated depth profiles of (A) calcite (vol %), (B) soil CO₂ production rate; (C) CO₂(g) (%), (D) DIC, and (E) Ca at 0.37 m/a, and (F) effluent Ca concentrations at different infiltration rates. From top to bottom rows are Scenario 1 (Grass_{PF} and Wood_{PF}; with flow partitioning, first row), Scenario 2 (Grass_{VF} and Wood_{VF}; 100% vertical flow, second row), and Scenario 3 (Grass_{HF} and Wood_{HF}; 100% lateral flow, last row). Red and blue colors present grassland and woodland, respectively. Lines and empty circles represent modelling outputs while filled circles with error bar in Figure C are the annual average CO₂(g) data (in Table 2). Arrows in Figure A indicate flow conditions. In F, soil water and groundwater refer to concentrations at the lateral (50 cm) and vertical (366 cm) outlets, respectively. Higher fractions of vertical flow in Wood_{PF} led to higher stream Ca concentration compared to Grass_{PF}. In Scenarios 2 and 3 without




445 flow partitioning, stream Ca concentrations were similar. The concentrations of DIC versus discharge are very similar to the Ca concentrations in F.

Scenarios 2-3 for biogeochemical effects (without flow partitioning). Scenarios 2 and 3 were end-member cases that bracket the range of rooting effects. Calcite and soil CO₂ were distributed the same way as their corresponding PF cases (Table 3). The Wood_{VF} and Grass_{VF} cases had 100% flow going downward via the deeper calcite zone maximizing the CO₂-water-calcite contact. 450 In the Wood_{HF} and Grass_{HF} cases (Table 3), all water exited at 50 cm, bypassing the deeper calcite-abundant zone, minimizing the CO₂-water-calcite contact. Figures 3A2-F2 and 3A3-F3 show the VF (Wood_{VF} and Grass_{VF}) and HF (Wood_{HF} and Grass_{HF}) cases, respectively. Similar to the flow partitioning cases, the concentrations of reaction products were low in the shallow zone and increased over 10 times within a short distance ~ 5 cm at the depth of ~ 50 cm. The woodland cases increased slightly more than the grassland cases because of the slightly steeper soil CO₂ distribution (Figure 3C2 and 3C3). Figure 3F2 indicates that the effluent 455 Ca concentrations were slightly higher in Wood_{VF} due to the high soil CO₂ level at the bottom outlet. The Grass_{HF} and Wood_{HF} case almost had the same effluent Ca concentrations at the upper soil (Figure 3F3).

Concentration-discharge relationship and weathering rates in all cases. The VF cases had the highest effluent concentrations and weathering rates, whereas the HF cases had lowest concentrations and weathering rates, and the Grass_{PF} and Wood_{PF} cases fell in between (Figure 4A-B). This is because the VF cases maximized the CO₂-calcite contact with 100% flow through the calcite zone, whereas the HF cases had minimum CO₂-calcite contact with water bypassing the calcite zone (column 3-4 in Table 3). The Grass_{PF} and Wood_{PF} cases allowed different extent of contact prescribed by the amount of flow via the calcite zone. A case run without soil respiration (i.e., no CO₂(g*), not shown) indicated that Ca and DIC concentrations were more than an order of magnitude lower than cases with soil respiration. In addition, the PF and HF cases generally showed dilution patterns with concentration decreasing with infiltration rates, as compared to the VF cases where a chemostatic pattern emerge with almost no changes with infiltration rates. This is because in the VF cases, the concentrations mostly reached equilibrium concentration. In 465 PF and HF cases, a large proportion of water flow through soils with negligible calcite where the water becomes further away from equilibrium as infiltration rates increase.

Deleted: shallow

Formatted: Not Highlight



470 Figure 4. (A-B) Stream water Ca and DIC concentrations, and (C-D) stream fluxes from all scenarios. Stream water was the overall effluent from soil water and groundwater. The differences caused by hydrological differences (VF, HF, and PF) were much larger than the differences within each pair with the same flow partitioning, indicating significant hydrological impacts on weathering. The Ca fluxes in the VF cases (100% vertical flow) were higher than flow partitioning cases because they enabled maximum CO₂-charged water content with unweathered calcite at depth.

475

Within each pair with the same flow pattern, the difference was mainly due to the distribution of soil CO₂. The difference of weathering fluxes was 1-12% and 1-5% between HF cases and VF cases, respectively, much smaller than differences between the PF cases. Comparing the HF and VF cases, differences in weathering fluxes was 73% at 3.7×10^{-2} m/a and 721% at 3.7 m/a, about 1-2 orders of magnitude higher than differences induced by soil CO₂ distribution. Between the Grass_{PF} and Wood_{PF} cases, the differences were in the range of 17%-207% at the flow range of 3.7×10^{-2} - 3.7 m/a. The DIC fluxes showed similar trends (Figure 4D).


480

Although weathering rates generally increased with infiltration rates, the woodland increased more (4.6×10^{-2} to 2.4 mol/a in Wood_{PF}). Estimates of the overall soil CO₂ production rate in Grass_{PF} and Wood_{PF} suggest that the values under the same infiltration rate were all within a difference of 10% (Figure S2), indicating that the difference of weathering rates are mainly driven by the flow partitioning. As a reference, the weathering rate in cases with soil respiration was up to 10 times higher compared to that without soil respiration.

485

Development of reaction fronts at the century scale. To explore the longer-term effects, we ran the PF cases at 0.37 m/a for 300 years. More water flowing vertically through abundant calcite zones in Wood_{PF} resulted in faster weathering and deeper reaction front at a depth of ~210 cm compared to ~95 cm in Grass_{PF} (Figure 5A). The depletion of calcite led to an increase of porosity (Figure 5B), which was over one order of magnitude higher at the domain scale of the woodland than that in the grassland (Figure 5C). Permeability evolution had the similar trend with porosity (not shown here). This indicates that if deeper roots promoted more

490



Deleted:

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

Deleted: yr

water into deeper soils, they would push reaction fronts deeper and controlled the position where chemically unweathered bedrock was transformed into weathered bedrock. At time scales longer than century scale, calcite may become depleted, which ultimately reduces weathering rates (White and Brantley, 2003) and lead to similar weathering fronts in grasslands and woodlands.





Figure 5. Predicted soil profiles of (A) calcite volume change (Δ Calcite = initial calcite volume - current calcite volume) and (B) porosity after 300 years; (C) Predicted temporal evolution of domain-scale porosity in the grassland and woodland. The infiltration rate is 0.37 m/a. Red and blue line represents the case GrassPF and WoodPF, respectively.

4.3 The regulation of weathering rates by CO₂-carbonate contact

Natural systems are characterized by preferential flow paths such that flow distribution is not uniform in space. Weathering in such systems with preferential flow in zones of differing reactivities have been shown to hinge on the contact time between water and the reacting minerals instead of all minerals that are present (Wen and Li, 2018) (Eq. 1). Here we contextualize the weathering rates in different scenarios (symbols) with predictions from an upscaled rate law developed by Wen and Li (2018) that incorporate the effects of heterogeneities in flow paths (Figure 6). The time ratio $\frac{\tau_a}{\tau_{ad,r}}$ compares the domain water contact time (or residence time) with the contact time with dissolving calcite. Note that τ_a is total domain pore volume V_T / total water flow Q_T , and $\tau_{ad,r}$ is total reactive pore volume V_r / total water fluxes passing through reactive zone Q_r . Also note that water passing through the reactive zone stays in the subsurface longer such that it is older water in general. The ratio $\frac{\tau_a}{\tau_{ad,r}}$ is therefore akin to the older water fraction, i.e., the fraction of older water compared to the total water fraction. The older water fraction F_{ow} is the counter part of the young water fraction F_{yw} that has been discussed in the literature (Kirchner, 2016, 2019). In GrassVF and WoodVF (open circles) where all CO₂-charged water flew through the deeper calcite zones, CO₂-calcite interactions reached maximum such that values of $\frac{\tau_a}{\tau_{ad,r}}$ approached 1.0, meaning all water interacted with calcite. Under this condition, weathering rates are the highest among all cases. In contrast, in WoodHF and GrassHF (open diamonds) where all CO₂-charged water bypassed the deeper calcite zone, $\frac{\tau_a}{\tau_{ad,r}}$ can be 1-3 orders of magnitude lower and weathering rates were at their minima. The $\frac{\tau_a}{\tau_{ad,r}}$ value in the water partitioning cases (GrassPF and WoodPF, filled circles) fell in between. At the same τ_a , the WoodPF case with deepening roots promoted CO₂-water-calcite contact (i.e., larger $\frac{\tau_a}{\tau_{ad,r}}$), and dissolved calcite at higher rates.

The magnitude of rate difference also depends on the overall flow rates (or domain contact time τ_a). At fast flow with small τ_a (large symbols and thick lines in Figure 6), flow partitioning has a larger influence. At τ_a of 0.47 years, the weathering



Formatted: Not Highlight



Deleted: yr

Deleted: year

rates in the VF cases were more than an order of magnitude higher than those in the PF cases. The weathering rate in the woodland was over 7 times that of the grassland. In contrast, at τ_a of 47 years, the rate differences between grass- and woody- cases were less than 25%, largely because the dissolution has reached equilibrium. In the VF and HF cases where flow conditions were the same, the soil CO_2 distribution in grassland and woodland differed only slightly, leading to similar values of $\frac{\tau_a}{\tau_{ad,r}}$ and weathering rates and indicating minimal impacts of soil CO_2 distribution. These rates from numerical experiments closely follow the prediction from the upscaled reaction rate law (grey lines, Eq. 1). The rate law predicted that weathering rates increased from HF cases with small $\frac{\tau_a}{\tau_{ad,r}}$ values to VF cases where $\frac{\tau_a}{\tau_{ad,r}}$ approached 1. It also showed that weathering rates reached their maxima in homogeneous domains without flow partitioning, or when the fraction of older, reactive water is essentially 1.0.



535 Figure 6. Calcite weathering rate as a function of the reactive time ratio $\frac{\tau_a}{\tau_{ad,r}}$, a proxy of the older, reactive water fraction compared to the total water fluxes. Symbols are rates from numerical experiments. Grey lines are predictions from the rate law equation at $\alpha = 0.8$ (Wen and Li, 2018). Large to small dots and thick to thin lines are for infiltration rates from $10^{0.6}$ to $10^{-1.4}$ m/a. The red filled stars represent the monthly rates in September (highest soil CO_2) and March (lowest soil CO_2) in the base case at $10^{-0.4}$ (0.37) m/a; the black to grey dash lines represent predictions with increasing soil CO_2 level (i.e., larger τ_{eq}) from Eq. (1). Grey filled star is for the case without soil CO_2 . At any specific infiltration rate or τ_a , the VF and HF cases bracket the two ends whereas the PF cases fall in between. The Wood_{PF} case with deeper roots enhanced the CO_2 -water-calcite contact (i.e., larger $\frac{\tau_a}{\tau_{ad,r}}$ ratios), leading to higher calcite weathering rates compared to grasslands. The differences of weathering rates induced by different soil CO_2 level were relatively small compared to those hydrological changes induced by rooting depth.

5 Discussion

Because carbonate dissolution is thermodynamically controlled and transport limited, the overall weathering rates depend on how much CO_2 -charged water flushes through the carbonate zone. This work indicates that the roots potentially enhance weathering rates in two ways. First, roots can control thermodynamic limits of carbonate dissolution by regulating how much CO_2 is transported downward and enters the carbonate-rich zone. In fact, the base-case grassland data and model reveal that the concentrations of Ca and DIC are regulated by seasonal fluctuation of $p\text{CO}_2$ and soil respiration. Second, roots control how much and how frequently water fluxes through the carbonate zone, exports reaction products at equilibrium such that more dissolution

Formatted: Font: Times New Roman
 Deleted: yr
 Deleted: yr

can occur. The numerical experiments indicate that carbonate weathering at depth hinges on the rate of delivery of CO₂-enriched water. Deepening roots in woodlands that channel more water into the unweathered carbonate can elevate weathering rates by more than an order of magnitude compared to grasslands. Below we elaborate and discuss these two messages.

5.1 The thermodynamics of carbonate weathering: control of temperature and pCO₂

560 The base case data and simulation showed that calcite dissolution reaches equilibrium rapidly and is thermodynamically controlled (Section 4.1), which echoes observations from other weathering studies (Tsypin and Macpherson, 2012; Gaillardet et al., 2019). The extent of dissolution, or solubility indicated in Ca and DIC concentrations, is determined by soil CO₂ levels that are a function of ecosystem functioning and climate. In hot, dry summer, soil respiration reaches its maximum rates in **upper** soil horizons and pCO₂ peaks (Figure 2). In wet and cold winter, soil pCO₂ plummets, leading to much lower Ca and DIC concentrations. Based on Reactions 0-4 (Table 1) and temperature dependence of equilibrium constants, the following equations can be derived (detailed derivation in the SI) for Ca and DIC concentrations in carbonate-dominated landscapes:

$$C_{Ca} = \sqrt[3]{K_{t,25} \exp\left(-\frac{\Delta H_t^\circ}{4R} * \left(\frac{1}{T} - \frac{1}{273.15+25}\right)\right) * pCO_2} \quad (2)$$

$$C_{DIC} = \sqrt[3]{K_{t,25} \exp\left(-\frac{\Delta H_t^\circ}{4R} * \left(\frac{1}{T} - \frac{1}{273.15+25}\right)\right) * pCO_2 + K_{1,25} \exp\left(-\frac{\Delta H_1^\circ}{4R} * \left(\frac{1}{T} - \frac{1}{273.15+25}\right)\right) * pCO_2} \quad (3)$$

570 Here $K_{t,25}$ is the total equilibrium constant $K_t (= \frac{\alpha_{Ca^{2+}} \alpha_{HCO_3^-}^2}{pCO_2} = K_1 K_4)$ of the combined reactions 1 and 4 at 25°C; ΔH_t° is the corresponding standard enthalpy (-35.83 kJ/mol); $K_{1,25}$ and ΔH_1° are the equilibrium constant and standard enthalpy of Reaction 1 (in Table 1); R is the gas constant ($=8.314 \times 10^{-3}$ kJ/K/mol). Eq. 2 and 3 imply that DIC and Ca in **upper** soil water (0.2 m) are lower compared to groundwater (3.6 m) in the base case at Konza, due to lower dissolved CO₂(aq) arising from higher temperature and higher diffusion rates in **upper** soil.

575 Eq. 2 and 3 were tested with **soil pCO₂ and spring water chemistry** data from 8 carbonate-dominated catchments (Dandurand et al., 1982; Lopez-Chicano et al., 2001; Moral et al., 2008; Ozkul et al., 2010; Kanduc et al., 2012; Calmels et al., 2014; Abongwa and Atekwana, 2015; Huang et al., 2015) (also see the SI for details). **As shown in Figure 7**, spring water (as representing groundwater) DIC and Ca concentrations increase with pCO₂, and Eq. 2-3 can describe these data and **DIC and Ca** levels from numerical simulations from this work (empty dots in Figure 7). The lines of Eq. 2-3 describe the relationship at 10 °C, the mean annual temperature, confirming the thermodynamics-control of carbonate dissolution by soil CO₂ levels. The equation lines closely predicted Ca and DIC under high pCO₂ and lower pH conditions (< 8.0), because these conditions ensure the validity of the assumption of negligible CO₃²⁻ in the derivation. The presence of cations and anions other than Ca and DIC can complicate the solution and can bring significant variations of DIC and Ca concentrations under the same pCO₂ conditions.

Deleted: shallow

Deleted: shallow

Deleted: shallow

Deleted: compiled

Deleted: Plotting spring-water chemistry against pCO₂ indicates that

Deleted: (Figure 7),

Deleted: also




Figure 7. (A) Ca and (B) DIC data from the literature (gray dots) and prediction lines of Eq. (2) – (3) at 10 (solid line) and 17 °C (dash line). Measured spring water chemistry are from calcite-dominant catchments in the literature: Abongwa and Atekwana (2015); Lopez-Chicano et al. (2001); Moral et al. (2008); Huang et al. (2015); Ozkul et al. (2010); Dandurand et al. (1982); Calmels et al. (2014); Kanduc et al. (2012); and Tsy-pin and Macpherson (2012).

High temperature ($T = 17^{\circ}\text{C}$) leads to lower DIC and Ca concentrations by about 10% due to the lower calcite and CO_2 solubility at higher T . Higher T also elevates $p\text{CO}_2$ by enhancing soil respiration. Various equations exist for predicting soil $p\text{CO}_2$ based on climate and ecosystem functioning indicators such as Net Primary Production (NPP) (Cerling, 1984; Goddérís et al., 2010; Romero-Mujalli et al., 2019a). These equations can be used together with Eq. 2-3 for the estimation of Ca and DIC concentrations in carbonate-derived waters. Gaillardet et al. (2019) shows that $p\text{CO}_2$ can increase by 2 times with T increase from 9°C to 17°C , which can elevate DIC and Ca concentrations over 50%. Macpherson et al. (2008) observed a 20% increase in groundwater $p\text{CO}_2$ in Konza over a 15-year period and suggested that increased soil respiration due to climate warming may have elevated soil and groundwater $p\text{CO}_2$. Hasenmueller et al. (2015) demonstrated topographic controls on soil CO_2 . Variations of CO_2 and Ca and DIC concentrations therefore are an integrated outcome of climate, soil respiration, subsurface structures, and hydrological conditions.

5.2 Hydrological controls of root-enhanced carbonate weathering

Evidence from field data. Data in Konza show that although soil $p\text{CO}_2$ peaks in summer, these peaks do not occur right away in deeper groundwater until about 2 months later. Macpherson et al. (2008) and Tsy-pin and Macpherson (2012) contributed the delay to the water travel time from soil to groundwater aquifer. The calculation of travel time based on depth difference in soil and groundwater sampling location (214 cm) and average velocity (0.37 m/d) indicate that it will takes 7-8 months in average for water to reach deeper groundwater. The 2 months delay, much shorter than the estimated 7-8 months, suggests that CO_2 -charged water may arrive deeper zones via preferential flow facilitated by roots or other macropores. Macropores can take be large conduits that have been observed in carbonate formations (Hartmann et al. 2014).

The numerical experiments suggest that the root-relevant hydrology can play an essential role in enhancing chemical weathering (Figure 4 and 6). In general, the hydrological enhancement of weathering rates also alludes to the key connection between weathering and the transit (travel) time distribution (McGuire and McDonnell, 2006; Sprenger et al., 2019). In particular, water that routes through lateral paths is younger; water that penetrates deeper is older. Figure 6 says that weathering rates increase with increasing reactive water fraction, until reaching their maxima when all water is in contact with reactive minerals. This aligns with observations at Konza that woody-encroached watersheds exhibit higher Ca fluxes in stream and supports the hypothesis that

Deleted: concentrations

Deleted: field data in

Deleted: .

Deleted: Lines are predictions from the thermodynamic solutions (Eq. 2-3) at 10 (solid line) and 17 °C (dash line).

Formatted: Not Highlight

Deleted: due to higher

Deleted: year

Deleted: via

Deleted: storm events with much faster flow....

Deleted: The enhanced

Formatted: Font color: Auto

deeper roots can enhance mineral-water interaction via deeper flow paths (Sullivan et al., 2019). Deepening roots can also enhance connectivity between shallow and deeper zones, therefore reducing concentration contrasts between soil water and groundwater.

635 This can lead to more chemostatic C-Q relationships as shown in Wood_{PF} ($b = -0.14$ in $C=aQ^b$) compared to Grass_{PF} ($b = -0.33$) (Figure 4). These findings echo conclusions from Zhi et al. (2019), and Zhi and Li (2020) that chemistry differences in shallow versus deeper waters regulate C-Q patterns. The b values from the C-Q relationships coming out of the 1D modeling (Figure 4) suggest that if flow partitioning is the only difference between the grassland and woody watersheds, a C-Q relationship exhibiting dilution with negative b values is expected in the grassland. The Konza stream data in Sullivan et al. (2019) however showed that
640 C-Q slopes in grasslands ($b = -0.003$) and in woody-encroached lands ($b = 0.013$) are both close to zero (Figure 7 and 8 in Sullivan et al. (2019)). The root influence on C-Q relationship therefore remains equivocal.

Deleted: that chemistry differences in shallow water versus groundwater regulate C-Q patterns.

Limitations of the model. This discrepancy may suggest that other catchment features that are not represented in the simple 1D

Formatted: Normal, Space After: 0 pt

645 model can influence C-Q relationships. The model does not explicitly simulate how and to what degree root distribution at depth alters flow pathways. Instead we focus on the first-order principles of hydrological ramification of roots. The numerical experiments took general observations of rooting characteristics in grasslands and woodlands, and assumed that deepening roots in woodlands promote higher flow partitioning into deep subsurface (Canadell et al., 1996; Nardini et al., 2016; Pawlik et al., 2016). In natural systems, however, other factors can also influence flow partitioning. ~~Some of these are not explicitly represented in our modeling exercises.~~ For example, contrasts in flow-conducting properties (i.e., porosity and permeability) in shallow and deep zones, physical and chemical heterogeneity in carbonate distribution (Wen and Li, 2018), connectivity between different
650 areas of the catchment in dry and wet times (Wen et al., 2020), and water table and corresponding lateral flow depth associated with the rainfall frequency and intensity (Li et al., 2017; Harman and Cosans, 2019). All these factors may affect the water-calcite contact, leading to changes of carbonate weathering rates (Figure 6). In addition to the alterations of hydrological flow paths, deepening roots may also affect the proportion of plant water uptake or soil water loss through transpiration (Pierret et al., 2016; Zhu et al., 2018), further modifying water fluxes into deep subsurface layers. For example, studies of semiarid woodlands show
655 that woody species predominantly use deep subsurface water while grasses predominantly use soil water from the upper soil layer (Ward et al., 2013). Furthermore, shallower root distributions in grasslands may be more efficient in using water from small rainfall events than forests with their deeper root distributions (Mazzacavallo and Kulmatiski, 2015). Other studies have also documented significant competition for water uptake in upper soil layers among both woody and grass species (Scholes and Archer, 1997). It remains inconclusive how these water uptake characteristics are best represented in numerical experiments. These processes are
660 therefore not included at this point.

Deleted: This includes, for

Deleted: property

Field Code Changed

Deleted: and

Deleted: . Representing these dynamics require a large number of processes and parameters that we do not have data to constrain at this point.

Field Code Changed

In addition, distributions of microbes and organic acids associated with rooting structures vary with sites and seasons, and root channels and dry conditions with less soil water may trigger calcite re-precipitation. Microbial activities surrounding living and dead roots can also lead to calcite precipitation and infilling of fractures and other macropores, and alter flow pathways (Lambers et al., 2009). Organic acids can decrease soil water pH, increase mineral solubilities through organic-metal complexations, and accelerate chemical weathering (Pittman and Lewan, 2012; Lawrence et al., 2014). Their impacts on carbonate weathering kinetics might be smaller (due to the fast kinetics) compared to their alteration of calcite solubility in natural systems. Such processes further complicate how to represent biogeochemical processes in models. Although we do not explicitly simulate all of these simultaneous and competing processes in our numerical experiments, the model was constrained by the field data in grasslands and woodland that have already integrated these effects in natural systems. Indeed, the weathering rates can be
665 understood as the net weathering rates that are the net difference between dissolution and reprecipitation. This is reflected in lower carbonate weathering rates under low infiltration (low flow) conditions.

The need for root-relevant measurements. The data-model discrepancy highlights the limitation of a simple model but also points to the need for measurements. In fact, because carbonate weathering is transport-limited and depends largely on water flow via carbonate-rich zones, co-located measurements of rooting characteristics, flow, and water chemistry at depth are essential. Root measurements however rarely go deeper than 30 cm (Richter and Billings, 2015). Existing work exploring root influence on weathering focused primarily on effects of soil CO₂ and root exudates (Drever, 1994; Lawrence et al., 2014; Gaillardet et al., 2019). Although it is well known that roots play a paramount role in modulating macropores and subsurface flow (Fan et al., 2017), the interactions between root characteristics, flow partitioning, and chemical weathering has remained poorly understood. Rooting characteristics depend on climate, plant species, topography, soil properties and geology (Canadell et al., 1996; Mazvimavi et al., 2004; Price, 2011; Nardini et al., 2016). Further studies are needed to characterize root distribution beyond 30 cm, how they vary with intrinsic plant species and external conditions, and how and to what extent they alter subsurface flow. The first step could link rooting characteristics, including density and depth, to soil properties and borrow insights from existing relationships between soil properties and subsurface structure. For example, images of roots and pore structures can be used to characterize the spatiotemporal heterogeneity of fluxes (Renard and Allard, 2013). Geostatistical indices such as permeability variance and correlation length can be used to quantify rooting structure, and relate them to flow partitioning and mineral weathering via numerical experiments (Wen and Li, 2017). The combination of numerical reactive transport experiments built on realistic rooting structure can help develop quantitative relationship between roots and flow partitioning, and could support models for estimating the influence of rooting dynamics on water and carbon cycles at the catchment scale.

5.3 Deepening roots enhance carbon fluxes into deep subsurface: a potential carbon sink?

Mounting evidence has shown that the terrestrial system has become a stronger carbon sink in recent decades (Heimann and Reichstein, 2008), potentially counting for the missing carbon sink as large as ~1 Pg C/a in the global carbon budget (Houghton, 2007; Cole et al., 2007). Although still much debated, recent studies have contributed the downward transport of soil-respired CO₂ and DIC into groundwater aquifers in deserts as a possible carbon sink in the global carbon cycle (Ma et al., 2014; Li et al., 2015). Considering the long residence time of DIC in groundwater (10²-10⁴ years) than in the atmosphere (~10¹-10² years; (Archer and Brovkin, 2008)), groundwater may act as a CO₂ storage sink. This work indicated that deepening roots can potentially reroute DIC fluxes to deeper groundwater storage. In particular, vegetation in dry places like deserts often have deep roots (Gupta et al., 2020). In fact, plants are known for growing deeper roots to tap groundwater during droughts (Brunner et al., 2015). Deepening roots can enhance downward water drainage (i.e., high vertical connectivity) to the depth and potentially facilitate the transport of DIC fluxes into deep subsurface. As the pace of climate change accelerates, summer droughts are expected to intensify, which can potentially channel more CO₂ into the deeper subsurface via deepening roots.

With constraints from soil CO₂ data, the simulated CO₂ production rates in Grass_{PF} and Wood_{PF} are similarly at ~0.6 mol C/m²/a under the infiltration rate of 0.37 m/a (Figure S2). This is at the low end of the belowground net production (NPP) estimations at the Konza site (~2.0-30.0 mol C/m²/a) assuming that belowground NPP accounts for 50 % of the total NPP (Lett et al., 2004; Knapp and Ojima, 2014). The simulations showed that in woodlands, the DIC downward fluxes can be > 2.0 times higher than those in grasslands (Figure 4). In other words, more soil-respired CO₂ can transport to groundwater and become stored there for centuries to millennium before entering a stream. At the short time scale, this enhanced downward transport will reduce CO₂ escape back into the atmosphere. This may explain the observations at Konza that woody encroachment increased NPP however soil CO₂ flux was significantly reduced compared to the open grassland (Lett et al., 2004). Changing land cover (e.g., woody encroachment and boreal forest creep) however is not the only mechanism for carbon sink (Stevens et al., 2017; Wang et al., 2020).

Deleted: yr⁻¹

Deleted: yr

Deleted: yr

Deleted: yr

Older aged forests also tend to have deepening roots and may act as the carbon sink (Luysaert et al., 2008), although this is not
725 the case in Konza.

6 Conclusions

This work aims to understand thermodynamic and hydrological control of carbonate weathering driven by deepening
730 roots. Field data and reactive transport simulation for a grassland in the Konza Prairie LTER site suggest that carbonate dissolution
is thermodynamically controlled and seasonal changes in temperature and $p\text{CO}_2$ drives variations of Ca and DIC concentrations in
soil water and groundwater. We derived equations based on reaction thermodynamics (Eq. 2-3) to estimate Ca and DIC as a
function of $p\text{CO}_2$ and temperature, which has been shown to be applicable in other carbonate-dominated systems. The numerical
735 experiments probed the potential effects of deepening roots on weathering by channeling a higher proportion of vertically
downward flow (40% of the total) into deep subsurface with abundant calcite. The results show that deeper penetration of roots
and higher vertical flow (recharge) enhanced CO_2 -carbonate contact. At an infiltration rate of 3.7 m/a , calcite weathering flux in
740 woodlands was 207% higher than that in grasslands. At $3.7 \times 10^{-2} \text{ m/a}$, the weathering flux in woodland was 17% higher. The
hydrological impacts on carbonate weathering were more than 10 times higher than the biogeochemical impacts via elevated soil
 CO_2 alone, underscoring the importance of root presence in weathering. The modeling demonstrates that the weathering rates
depend on flow partitioning (the older water fraction that penetrates deeper) and relative magnitude of water contact time in the
deep carbonate zone. At the century scale, with the higher proportion of vertical flow, the deeper roots pushed the weathering
745 fronts 2-times deeper, and resulted in a 10-times greater increase in porosity and permeability. Broadly, this deeper propagation of
reaction fronts may accelerate rates of channel incision and hillslope erosion (Lebedeva and Brantley, 2013; Brantley et al., 2017b),
and therefore speed up landscape evolution (Phillips, 2005). It alludes to the importance of considering changes in subsurface
hydrological flows associated with shifts in vegetation types and measuring rooting characteristics. This is particularly relevant as
we assess the effects of climate change, land cover and elevated atmosphere CO_2 concentrations on chemical weathering and
carbon cycles.

Data availability. All data used for model parameterization can be acquired from <http://lter.konza.ksu.edu/data>. The input files
necessary to reproduce the results are available from the authors upon request (lili@engr.psu.edu).

Author contributions. H. Wen, P. Sullivan and L. Li initiated the idea and designed the numerical experiments. G.L. Macpherson
750 and S. Billings provided the field data. H. Wen ran the simulations, analyzed simulation results, and wrote the first draft of the
manuscript. All co-authors participated in editing the manuscript.

Competing interests. The authors report no conflicts of interest.

Acknowledgement. We acknowledge funding support from the Konza Prairie LTER program (NSF DEB-1440484), the Signals in
the Soils (SitS, EAR-1331726), and the NSF project (EAR-1911960). We appreciate the assistance of the Konza Prairie Biological
Station. We are grateful for the field data provided by Misha Tsy-pin and Dr. Zachary Brecheisen.

Deleted: flow

Deleted: yr

Deleted: yr

Deleted: .

Deleted: assumed

760 **References**

- Abongwa, P. T., and Atekwana, E. A.: Controls on the chemical and isotopic composition of carbonate springs during evolution to saturation with respect to calcite, *Chem Geol*, 404, 136-149, 10.1016/j.chemgeo.2015.03.024, 2015.
- Ahrens, B., Braakhekke, M. C., Guggenberger, G., Schrupp, M., and Reichstein, M.: Contribution of sorption, DOC transport and microbial interactions to the C-14 age of a soil organic carbon profile: Insights from a calibrated process model, *Soil Biology & Biochemistry*, 88, 390-402, 10.1016/j.soilbio.2015.06.008, 2015.
- Andrews, J. A., and Schlesinger, W. H.: Soil CO₂ dynamics, acidification, and chemical weathering in a temperate forest with experimental CO₂ enrichment, *Global Biogeochemical Cycles*, 15, 149-162, 10.1029/2000gb001278, 2001.
- Angers, D. A., and Caron, J.: Plant-induced changes in soil structure: Processes and feedbacks, *Biogeochemistry*, 42, 55-72, 10.1023/a:1005944025343, 1998.
- 770 Aragao, L., Malhi, Y., Metcalfe, D. B., Silva-Espejo, J. E., Jimenez, E., Navarrete, D., Almeida, S., Costa, A. C. L., Salinas, N., Phillips, O. L., Anderson, L. O., Alvarez, E., Baker, T. R., Goncalvez, P. H., Huaman-Ovalle, J., Mamani-Solorzano, M., Meir, P., Monteagudo, A., Patino, S., Penuela, M. C., Prieto, A., Quesada, C. A., Rozas-Davila, A., Rudas, A., Silva, J. A., and Vasquez, R.: Above- and below-ground net primary productivity across ten Amazonian forests on contrasting soils, *Biogeosciences*, 6, 2759-2778, 10.5194/bg-6-2759-2009, 2009.
- 775 Archer, D., and Brovkin, V.: The millennial atmospheric lifetime of anthropogenic CO₂, *Clim. Change*, 90, 283-297, 10.1007/s10584-008-9413-1, 2008.
- Beerling, D. J., Chaloner, W. G., Woodward, F. I., and Berner, R. A.: The carbon cycle and carbon dioxide over Phanerozoic time: the role of land plants, *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 353, 75-82, doi:10.1098/rstb.1998.0192, 1998.
- 780 Bengtson, P., and Bengtsson, G.: Rapid turnover of DOC in temperate forests accounts for increased CO₂ production at elevated temperatures, *Ecology letters*, 10, 783-790, DOI: 10.1111/j.1461-0248.2007.01072.x, 2007.
- Berner, E. K., and Berner, R. A.: *Global environment: water, air, and geochemical cycles*, Princeton University Press, 2012.
- Berner, R. A.: Weathering, plants, and the long-term carbon cycle, *Geochimica et Cosmochimica Acta*, 56, 3225-3231, doi.org/10.1016/0016-7037(92)90300-8, 1992.
- 785 Berner, R. A.: Paleoclimate - The rise of plants and their effect on weathering and atmospheric CO₂, *Science*, 276, 544-546, 10.1126/science.276.5312.544, 1997.
- Beven, K., and Germann, P.: Macropores and water flow in soils revisited, *Water Resour Res*, 49, 3071-3092, 10.1002/wrcr.20156, 2013.
- Billings, S. A., Hirmas, D., Sullivan, P. L., Lehmeier, C. A., Bagchi, S., Min, K., Brecheisen, Z., Hauser, E., Stair, R., Flournoy, R., and Richter, D. D.: Loss of deep roots limits biogenic agents of soil development that are only partially restored by decades of forest regeneration, *Elementa-Science of the Anthropocene*, 6, 10.1525/elementa.287, 2018.
- Brantley, S. L., Eissenstat, D. M., Marshall, J. A., Godsey, S. E., Balogh-Brunstad, Z., Karwan, D. L., Papuga, S. A., Roering, J., Dawson, T. E., Evaristo, J., Chadwick, O., McDonnell, J. J., and Weathers, K. C.: Reviews and syntheses: on the roles trees play in building and plumbing the critical zone, *Biogeosciences*, 14, 5115-5142, 10.5194/bg-14-5115-2017, 2017a.
- 795 Brantley, S. L., Lebedeva, M. I., Balashov, V. N., Singha, K., Sullivan, P. L., and Stinchcomb, G.: Toward a conceptual model relating chemical reaction fronts to water flow paths in hills, *Geomorphology*, 277, 100-117, 10.1016/j.geomorph.2016.09.027, 2017b.
- Breecker, D. O., Sharp, Z. D., and McFadden, L. D.: Atmospheric CO₂ concentrations during ancient greenhouse climates were similar to those predicted for AD 2100, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 576-580, 10.1073/pnas.0902323106, 2010.
- 800 Brunner, I., Herzog, C., Dawes, M. A., Arend, M., and Sperisen, C.: How tree roots respond to drought, *Frontiers in plant science*, 6, 547, doi.org/10.3389/fpls.2015.00547, 2015.
- Calmels, D., Gaillardet, J., and Francois, L.: Sensitivity of carbonate weathering to soil CO₂ production by biological activity along a temperate climate transect, *Chem Geol*, 390, 74-86, 10.1016/j.chemgeo.2014.10.010, 2014.
- 805 Canadell, J., Jackson, R. B., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E. D.: Maximum rooting depth of vegetation types at the global scale, *Oecologia*, 108, 583-595, 10.1007/bf00329030, 1996.
- Carey, J. C., Tang, J. W., Templer, P. H., Kroeger, K. D., Crowther, T. W., Burton, A. J., Dukes, J. S., Emmett, B., Frey, S. D., Heskell, M. A., Jiang, L., Machmuller, M. B., Mohan, J., Panetta, A. M., Reich, P. B., Reinsch, S., Wang, X., Allison, S. D., Bamminger, C., Bridgman, S., Collins, S. L., De Dato, G., Eddy, W. C., Enquist, B. J., Estiarte, M., Harte, J., Henderson, A., Johnson, B. R., Larsen, K. S., Luo, Y., Marhan, S., Melillo, J. M., Peuelas, J., Pfeifer-Meister, L., Poll, C., Rastetter, E., Reinmann, A. B., Reynolds, L. L., Schmidt, I. K., Shaver, G. R., Strong, A. L., Suseela, V., and Tietema, A.: Temperature response of soil respiration largely unaltered with experimental warming, *Proceedings of the National Academy of Sciences of the United States of America*, 113, 13797-13802, 10.1073/pnas.160536511, 2016.
- 810 Cerling, T. E.: The stable isotopic composition of modern soil carbonate and its relationship to climate, *Earth Planet Sc Lett*, 71, 229-240, 10.1016/0012-821x(84)90089-x, 1984.
- 815 Cheng, J. H., Zhang, H. J., Wang, W., Zhang, Y. Y., and Chen, Y. Z.: Changes in Preferential Flow Path Distribution and Its Affecting Factors in Southwest China, *Soil Science*, 176, 652-660, 10.1097/SS.0b013e31823554ef, 2011.

- Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 10, 171-184, 10.1007/s10021-006-9013-8, 2007.
- Costa, A.: Permeability-porosity relationship: A reexamination of the Kozeny-Carman equation based on a fractal pore-space geometry assumption, *Geophys Res Lett*, 33, 10.1029/2005GL025134, 2006.
- Covington, M. D., Gully, J. D., and Gabrovsek, F.: Natural variations in calcite dissolution rates in streams: Controls, implications, and open questions, *Geophys Res Lett*, 42, 2836-2843, 10.1002/2015gl063044, 2015.
- Dandurand, J. L., Gout, R., Hoefs, J., Menschel, G., Schott, J., and Usdowski, E.: Kinetically controlled variations of major components and carbon and oxygen isotopes in a calcite-precipitating spring, *Chem Geol*, 36, 299-315, 10.1016/0009-2541(82)90053-5, 1982.
- Deng, H., Voltolini, M., Molins, S., Steefel, C., DePaolo, D., Ajo-Franklin, J., and Yang, L.: Alteration and Erosion of Rock Matrix Bordering a Carbonate-Rich Shale Fracture, *Environmental Science & Technology*, 51, 8861-8868, 10.1021/acs.est.7b02063, 2017.
- Drever, J. I.: The effect of land plants on weathering rates of silicate minerals, [Geochimica et Cosmochimica Acta](#), 58, 2325-2332, [doi.org/10.1016/0016-7037\(94\)90013-2](#), 1994.
- Eberbach, P. L.: The eco-hydrology of partly cleared, native ecosystems in southern Australia: a review, *Plant and Soil*, 257, 357-369, 10.1023/A:1027392703312, 2003.
- Fan, Y., Miguez-Macho, G., Jobbagy, E. G., Jackson, R. B., and Otero-Casal, C.: Hydrologic regulation of plant rooting depth, *Proceedings of the National Academy of Sciences of the United States of America*, 114, 10572-10577, 10.1073/pnas.1712381114, 2017.
- Frank, D. A., Pontes, A. W., Maine, E. M., Caruana, J., Raina, R., Raina, S., and Fridley, J. D.: Grassland root communities: species distributions and how they are linked to aboveground abundance, *Ecology*, 91, 3201-3209, 10.1890/09-1831.1, 2010.
- Gaillardet, J., Calmels, D., Romero-Mujalli, G., Zakharova, E., and Hartmann, J.: Global climate control on carbonate weathering intensity, *Chem Geol*, 527, 118762, 10.1016/j.chemgeo.2018.05.009, 2019.
- Gill, R. A., Kelly, R. H., Parton, W. J., Day, K. A., Jackson, R. B., Morgan, J. A., Scurlock, J. M. O., Tieszen, L. L., Castle, J. V., Ojima, D. S., and Zhang, X. S.: Using simple environmental variables to estimate below-ground productivity in grasslands, *Global Ecology and Biogeography*, 11, 79-86, 10.1046/j.1466-822X.2001.00267.x, 2002.
- Godd eris, Y., Williams, J. Z., Schott, J., Pollard, D., and Brantley, S. L.: Time evolution of the mineralogical composition of Mississippi Valley loess over the last 10 kyr: Climate and geochemical modeling, *Geochimica Et Cosmochimica Acta*, 74, 6357-6374, 10.1016/j.gca.2010.08.023, 2010.
- Gupta, A., Rico-Medina, A., and Cano-Delgado, A. I.: The physiology of plant responses to drought, *Science*, 368, 266-269, 10.1126/science.aaz7614, 2020.
- Harman, C. J., and Cosans, C. L.: A low-dimensional model of bedrock weathering and lateral flow coevolution in hillslopes: 2. Controls on weathering and permeability profiles, drainage hydraulics, and solute export pathways, *Hydrological Processes*, 33, 1168-1190, 10.1002/hyp.13385, 2019.
- Harper, C. W., Blair, J. M., Fay, P. A., Knapp, A. K., and Carlisle, J. D.: Increased rainfall variability and reduced rainfall amount decreases soil CO₂ flux in a grassland ecosystem, *Global Change Biology*, 11, 322-334, 10.1111/j.1365-2486.2005.00899.x, 2005.
- Hartmann, A., Goldscheider, N., Wagener, T., Lange, J., and Weiler, M.: Karst water resources in a changing world: Review of hydrological modeling approaches, *Rev. Geophys.*, 52, 218-242, doi:10.1002/2013RG000443, 2014.
- Hasenmueller, E. A., Jin, L. X., Stinchcomb, G. E., Lin, H., Brantley, S. L., and Kaye, J. P.: Topographic controls on the depth distribution of soil CO₂ in a small temperate watershed, *Appl Geochem*, 63, 58-69, 10.1016/j.apgeochem.2015.07.005, 2015.
- Hasenmueller, E. A., Gu, X., Weitzman, J. N., Adams, T. S., Stinchcomb, G. E., Eissenstat, D. M., Drohan, P. J., Brantley, S. L., and Kaye, J. P.: Weathering of rock to regolith: The activity of deep roots in bedrock fractures, *Geoderma*, 300, 11-31, [doi.org/10.1016/j.geoderma.2017.03.020](#), 2017.
- [Hauser, E., Richter, D. D., Markewitz, D., Brecheisen, Z., and Billings, S. A.: Persistent anthropogenic legacies structure depth dependence of regenerating rooting systems and their functions, *Biogeochemistry*, 147, 259-275, 10.1007/s10533-020-00641-2, 2020.](#)
- Heimann, M., and Reichstein, M.: Terrestrial ecosystem carbon dynamics and climate feedbacks, *Nature*, 451, 289-292, 10.1038/nature06591, 2008.
- Houghton, R. A.: Balancing the global carbon budget, *Annual Review of Earth and Planetary Sciences*, 35, 313-347, 10.1146/annurev.earth.35.031306.140057, 2007.
- Huang, F., Zhang, C. L., Xie, Y. C., Li, L., and Cao, J. H.: Inorganic carbon flux and its source in the karst catchment of Maocun, Guilin, China, *Environmental Earth Sciences*, 74, 1079-1089, 10.1007/s12665-015-4478-4, 2015.
- Jackson, R. B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E. D.: A global analysis of root distributions for terrestrial biomes, *Oecologia*, 108, 389-411, 10.1007/bf00333714, 1996.
- Jiang, H., Deng, Q., Zhou, G., Hui, D., Zhang, D., Liu, S., Chu, G., and Li, J.: Responses of soil respiration and its temperature/moisture sensitivity to precipitation in three subtropical forests in southern China, *Biogeosciences*, 10, 3963-3982, 10.5194/bg-10-3963-2013, 2013.

Deleted: Geochim Cosmochim Ac

- Jobbagy, E. G., and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to climate and vegetation, *Ecological Applications*, 10, 423-436, 10.2307/2641104, 2000.
- 880 Kanduc, T., Mori, N., Kocman, D., Stibilj, V., and Grassa, F.: Hydrogeochemistry of Alpine springs from North Slovenia: Insights from stable isotopes, *Chem Geol*, 300, 40-54, 10.1016/j.chemgeo.2012.01.012, 2012.
- [Kirchner, J. W.: Aggregation in environmental systems - Part 1: Seasonal tracer cycles quantify young water fractions, but not mean transit times, in spatially heterogeneous catchments, *Hydrol Earth Syst Sc*, 20, 279-297, 10.5194/hess-20-279-2016, 2016.](#)
- 885 [Kirchner, J. W.: Quantifying new water fractions and transit time distributions using ensemble hydrograph separation: theory and benchmark tests, *Hydrol. Earth Syst. Sci.*, 23, 303-349, 10.5194/hess-23-303-2019, 2019.](#)
- Knapp, A., and Ojima, D.: NPP Grassland: Konza Prairie, USA, 1984-1990, R1, ORNL DAAC, 2014.
- Kozeny, J.: *Über kapillare leitung der wasser in boden*, Royal Academy of Science, Vienna, Proc. Class I, 136, 271-306, 1927.
- [Lambers, H., Mougél, C., Jaillard, B., and Hinsinger, P.: Plant-microbe-soil interactions in the rhizosphere: an evolutionary perspective, *Plant and Soil*, 321, 83-115, 10.1007/s11104-009-0042-x, 2009.](#)
- 890 Lawrence, C., Harden, J., and Maher, K.: Modeling the influence of organic acids on soil weathering, *Geochim Cosmochim Acta*, 139, 487-507, 10.1016/j.gca.2014.05.003, 2014.
- Lebedeva, M. I., and Brantley, S. L.: Exploring geochemical controls on weathering and erosion of convex hillslopes: beyond the empirical regolith production function, *Earth Surface Processes and Landforms*, 38, 1793-1807, 10.1002/esp.3424, 2013.
- Lett, M. S., Knapp, A. K., Briggs, J. M., and Blair, J. M.: Influence of shrub encroachment on aboveground net primary productivity and carbon and nitrogen pools in a mesic grassland, *Canadian Journal of Botany-Revue Canadienne De Botanique*, 82, 1363-1370, 10.1139/b04-088, 2004.
- 895 Li, L., Bao, C., Sullivan, P. L., Brantley, S., Shi, Y. N., and Duffy, C.: Understanding watershed hydrogeochemistry: 2. Synchronized hydrological and geochemical processes drive stream chemostatic behavior, *Water Resour Res*, 53, 2346-2367, 10.1002/2016wr018935, 2017.
- 900 Li, Y., Wang, Y. G., Houghton, R. A., and Tang, L. S.: Hidden carbon sink beneath desert, *Geophys Res Lett*, 42, 5880-5887, 10.1002/2015gl064222, 2015.
- Lopez-Chicano, M., Bouamama, M., Vallejos, A., and Pulido-Bosch, A.: Factors which determine the hydrogeochemical behaviour of karstic springs. A case study from the Betic Cordilleras, Spain, *Appl Geochem*, 16, 1179-1192, 10.1016/s0883-2927(01)00012-9, 2001.
- 905 Luyssaert, S., Schulze, E. D., Börner, A., Knohl, A., Hessenmöller, D., Law, B. E., Ciais, P., and Grace, J.: Old-growth forests as global carbon sinks, *Nature*, 455, 213-215, 10.1038/nature07276, 2008.
- Ma, J., Liu, R., Tang, L. S., Lan, Z. D., and Li, Y.: A downward CO₂ flux seems to have nowhere to go, *Biogeosciences*, 11, 6251-6262, 10.5194/bg-11-6251-2014, 2014.
- Macpherson, G., and Sullivan, P. L. J. C. G.: Dust, impure calcite, and phyloliths: Modeled alternative sources of chemical weathering solutes in shallow groundwater, *Chem Geol*, 527, 118871, 10.1016/j.chemgeo.2018.08.007, 2019.
- 910 Macpherson, G. L.: Hydrogeology of thin limestones: The Konza Prairie Long-Term Ecological Research Site, Northeastern Kansas, *J Hydrol*, 186, 191-228, 10.1016/s0022-1694(96)03029-6, 1996.
- Macpherson, G. L., Roberts, J. A., Blair, J. M., Townsend, M. A., Fowle, D. A., and Beisner, K. R.: Increasing shallow groundwater CO₂ and limestone weathering, Konza Prairie, USA, *Geochimica Et Cosmochimica Acta*, 72, 5581-5599, 10.1016/j.gca.2008.09.004, 2008.
- 915 Mazvimavi, D., Meijerink, A. M. J., and Stein, A.: Prediction of base flows from basin characteristics: a case study from Zimbabwe, *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques*, 49, 703-715, 10.1623/hysj.49.4.703.54428, 2004.
- [Mazzacavallo, M. G., and Kulmatiski, A.: Modelling Water Uptake Provides a New Perspective on Grass and Tree Coexistence, *Plos One*, 10, 10.1371/journal.pone.0144300, 2015.](#)
- 920 [McGuire, K. J., and McDonnell, J. J.: A review and evaluation of catchment transit time modeling, *J Hydrol*, 330, 543-563, 10.1016/j.jhydrol.2006.04.020, 2006.](#)
- Moral, F., Cruz-Sanjulian, J. J., and Olias, M.: Geochemical evolution of groundwater in the carbonate aquifers of Sierra de Segura (Betic Cordillera, southern Spain), *J Hydrol*, 360, 281-296, 10.1016/j.jhydrol.2008.07.012, 2008.
- 925 Moriasi, D. N., Arnold, J. G., Van Liew, M. W., Bingner, R. L., Harmel, R. D., and Veith, T. L.: Model evaluation guidelines for systematic quantification of accuracy in watershed simulations, *Transactions of the Asabe*, 50, 885-900, 10.13031/2013.23153, 2007.
- Mottershead, D. N., Baily, B., Collier, P., and Inkpen, R. J.: Identification and quantification of weathering by plant roots, *Building and Environment*, 38, 1235-1241, 10.1016/s0360-1323(03)00080-5, 2003.
- 930 Nardini, A., Casolo, V., Dal Borgo, A., Savi, T., Stenni, B., Bertoincin, P., Zini, L., and McDowell, N. G.: Rooting depth, water relations and non-structural carbohydrate dynamics in three woody angiosperms differentially affected by an extreme summer drought, *Plant Cell and Environment*, 39, 618-627, 10.1111/pce.12646, 2016.
- Neff, J. C., and Hooper, D. U.: Vegetation and climate controls on potential CO₂, DOC and DON production in northern latitude soils, *Glob. Change Biol.*, 8, 872-884, 10.1046/j.1365-2486.2002.00517.x, 2002.
- 935 Nepstad, D. C., Decarvalho, C. R., Davidson, E. A., Jipp, P. H., Lefebvre, P. A., Negreiros, G. H., Dasilva, E. D., Stone, T. A., Trumbore, S. E., and Vieira, S.: The role of deep roots in the hydrological and carbon cycles of amazonian forests and pastures, *Nature*, 372, 666-669, 10.1038/372666a0, 1994.

- Nippert, J. B., Wieme, R. A., Ocheltree, T. W., and Craine, J. M.: Root characteristics of C-4 grasses limit reliance on deep soil water in tallgrass prairie, *Plant and Soil*, 355, 385-394, 10.1007/s11104-011-1112-4, 2012.
- 940 Noguchi, S., Tsuboyama, Y., Sidle, R. C., and Hosoda, I.: Spatially distributed morphological characteristics of macropores in forest soils of Hitachi Ohta Experimental Watershed, Japan, *Journal of Forest Research*, 2, 207-215, 10.1007/BF02348317, 1997.
- Oades, J. M.: The role of biology in the formation, stabilization and degradation of soil structure, *Geoderma*, 56, 377-400, 10.1016/0016-7061(93)90123-3, 1993.
- 945 Ozkul, M., Gokgoz, A., and Horvatincic, N.: Depositional properties and geochemistry of Holocene perched springline tufa deposits and associated spring waters: a case study from the Denizli Province, Western Turkey, in: *Tufas and Speleothems: Unravelling the Microbial and Physical Controls*, edited by: Pedley, H. M., and Rogerson, M., Geological Society Special Publication, 245-262, 2010.
- Palandri, J. L., and Kharaka, Y. K.: A compilation of rate parameters of water-mineral interaction kinetics for application to geochemical modeling, *Geological Survey Menlo Park CA*, 2004.
- 950 Patrick, L., Cable, J., Potts, D., Ignace, D., Barron-Gafford, G., Griffith, A., Alpert, H., Van Gestel, N., Robertson, T., Huxman, T. E., Zak, J., Loik, M. E., and Tissue, D.: Effects of an increase in summer precipitation on leaf, soil, and ecosystem fluxes of CO₂ and H₂O in a sotol grassland in Big Bend National Park, Texas, *Oecologia*, 151, 704-718, 10.1007/s00442-006-0621-y, 2007.
- Pawlik, L., Phillips, J. D., and Samonil, P.: Roots, rock, and regolith: Biomechanical and biochemical weathering by trees and its impact on hillslopes-A critical literature review, *Earth-Sci Rev*, 159, 142-159, 10.1016/j.earscirev.2016.06.002, 2016.
- Pennell, K. D., Boyd, S. A., and Abriola, L. M.: Surface-area of soil organic-matter reexamined, *Soil Science Society of America Journal*, 59, 1012-1018, 10.2136/sssaj1995.03615995005900040008x, 1995.
- Phillips, J. D.: Weathering instability and landscape evolution, *Geomorphology*, 67, 255-272, 10.1016/j.geomorph.2004.06.012, 2005.
- 960 [Pierret, A., Maeght, J. L., Clement, C., Montoroi, J. P., Hartmann, C., and Gonkhamdee, S.: Understanding deep roots and their functions in ecosystems: an advocacy for more unconventional research, *Annals of Botany*, 118, 621-635, 10.1093/aob/mcw130, 2016.](#)
- [Pittman, E. D., and Lewan, M. D.: Organic acids in geological processes, Springer Science & Business Media, 2012.](#)
- 965 Plummer, L., Wigley, T., and Parkhurst, D.: The kinetics of calcite dissolution in CO₂-water systems at 5 degrees to 60 degrees C and 0.0 to 1.0 atm CO₂, *Am J Sci*, 278, 179-216, 10.2475/ajs.278.2.179, 1978.
- Price, K.: Effects of watershed topography, soils, land use, and climate on baseflow hydrology in humid regions: A review, *Progress in Physical Geography*, 35, 465-492, 10.1177/0309133311402714, 2011.
- Renard, P., and Allard, D.: Connectivity metrics for subsurface flow and transport, *Adv Water Resour*, 51, 168-196, 10.1016/j.advwatres.2011.12.001, 2013.
- 970 Richter, D. D., and Billings, S. A.: 'One physical system': Tansley's ecosystem as Earth's critical zone, *New Phytol.*, 206, 900-912, 10.1111/nph.13338, 2015.
- Romero-Mujalli, G., Hartmann, J., and Börker, J.: Temperature and CO₂ dependency of global carbonate weathering fluxes—Implications for future carbonate weathering research, *Chem Geol*, 527, 118874, 10.1016/j.chemgeo.2018.08.010, 2019a.
- Romero-Mujalli, G., Hartmann, J., Börker, J., Gaillardet, J., and Calmels, D.: Ecosystem controlled soil-rock pCO₂ and carbonate weathering—Constraints by temperature and soil water content, *Chem Geol*, 527, 118634, 10.1016/j.chemgeo.2018.01.030, 2019b.
- 975 [Sadras, V. O.: Influence of size of rainfall events on water-driven processes - I. Water budget of wheat crops in south-eastern Australia, *Aust. J. Agric. Res.*, 54, 341-351, 10.1071/ar02112, 2003.](#)
- [Scholes, R. J., and Archer, S. R.: Tree-grass interactions in savannas, *Annual Review of Ecology and Systematics*, 28, 517-544, 10.1146/annurev.ecolsys.28.1.517, 1997.](#)
- 980 [Sprenger, M., Stumpp, C., Weiler, M., Aeschbach, W., Allen, S. T., Benettin, P., Dubbert, M., Hartmann, A., Hrachowitz, M., Kirchner, J. W., McDonnell, J. J., Orlowski, N., Penna, D., Pfahl, S., Rinderer, M., Rodriguez, N., Schmidt, M., and Werner, C.: The Demographics of Water: A Review of Water Ages in the Critical Zone, *Rev Geophys*, 57, 800-834, 10.1029/2018rg000633, 2019.](#)
- 985 Steefel, C. I., Appelo, C. A. J., Arora, B., Jacques, D., Kalbacher, T., Kolditz, O., Lagneau, V., Lichtner, P. C., Mayer, K. U., Meeussen, J. C. L., Molins, S., Moulton, D., Shao, H., Šimůnek, J., Spycher, N., Yabusaki, S. B., and Yeh, G. T.: Reactive transport codes for subsurface environmental simulation, *Computational Geosciences*, 19, 445-478, 10.1007/s10596-014-9443-x, 2015.
- 990 Stevens, N., Lehmann, C. E. R., Murphy, B. P., and Durigan, G.: Savanna woody encroachment is widespread across three continents, *Glob. Change Biol.*, 23, 235-244, 10.1111/gcb.13409, 2017.
- Steward, D. R., Yang, X., Lauwo, S. Y., Staggenborg, S. A., Macpherson, G. L., and Welch, S. M.: From precipitation to groundwater baseflow in a native prairie ecosystem: a regional study of the Konza LTER in the Flint Hills of Kansas, USA, *Hydrol Earth Syst Sc*, 15, 3181-3194, 10.5194/hess-15-3181-2011, 2011.
- 995 Sullivan, P. L., Stops, M. W., Macpherson, G. L., Li, L., Hirmas, D. R., and Dodds, W. K.: How landscape heterogeneity governs stream water concentration-discharge behavior in carbonate terrains (Konza Prairie, USA), *Chem. Geol.*, 527, 118989, doi.org/10.1016/j.chemgeo.2018.12.002, 2019.

- Tsypin, M., and Macpherson, G. L.: The effect of precipitation events on inorganic carbon in soil and shallow groundwater, Konza Prairie LTER Site, NE Kansas, USA, *Applied Geochemistry*, 27, 2356-2369, 10.1016/j.apgeochem.2012.07.008, 2012.
- 1000 Vargas, R., Collins, S. L., Thomey, M. L., Johnson, J. E., Brown, R. F., Natvig, D. O., and Friggens, M. T.: Precipitation variability and fire influence the temporal dynamics of soil CO₂ efflux in an arid grassland, *Glob. Change Biol.*, 18, 1401-1411, 10.1111/j.1365-2486.2011.02628.x, 2012.
- Vergani, C., and Graf, F.: Soil permeability, aggregate stability and root growth: a pot experiment from a soil bioengineering perspective, *Ecology*, 9, 830-842, 10.1002/eco.1686, 2016.
- 1005 Wang, J. A., Sulla-Menashe, D., Woodcock, C. E., Sonnentag, O., Keeling, R. F., and Friedl, M. A.: Extensive land cover change across Arctic-Boreal Northwestern North America from disturbance and climate forcing, *Glob. Change Biol.*, 26, 807-822, 10.1111/gcb.14804, 2020.
- Ward, D., Wiegand, K., and Getzin, S.: [Walter's two-layer hypothesis revisited: back to the roots!](#), *Oecologia*, 172, 617-630, 10.1007/s00442-012-2538-y, 2013.
- 1010 Watson, K. W., and Luxmoore, R. J.: Estimating macroporosity in a forest watershed by use of a tension infiltrometer, *Soil Science Society of America Journal*, 50, 578-582, 10.2136/sssaj1986.03615995005000030007x, 1986.
- Weiler, M., and Naef, F.: An experimental tracer study of the role of macropores in infiltration in grassland soils, *Hydrological Processes*, 17, 477-493, 10.1002/hyp.1136, 2003.
- 1015 Wen, H., Li, L., Crandall, D., and Hakala, A.: Where lower calcite abundance creates more alteration: enhanced rock matrix diffusivity induced by preferential dissolution, *Energy Fuels*, 30, 4197-4208, 10.1021/acs.energyfuels.5b02932, 2016.
- Wen, H., and Li, L.: An upscaled rate law for magnesite dissolution in heterogeneous porous media, *Geochim Cosmochim Acta*, 210, 289-305, 10.1016/j.gca.2017.04.019, 2017.
- Wen, H., and Li, L.: An upscaled rate law for mineral dissolution in heterogeneous media: The role of time and length scales, *Geochim Cosmochim Acta*, 235, 1-20, 10.1016/j.gca.2018.04.024, 2018.
- 1020 Wen, H., Perdrial, J., Abbott, B. W., Bernal, S., Dupas, R., Godsey, S. E., Harpold, A., Rizzo, D., Underwood, K., Adler, T., Sterle, G., and Li, L.: Temperature controls production but hydrology regulates export of dissolved organic carbon at the catchment scale, *Hydro Earth Syst Sc*, 24, 945-966, 10.5194/hess-24-945-2020, 2020.
- White, A. F., and Brantley, S. L.: The effect of time on the weathering of silicate minerals: why do weathering rates differ in the laboratory and field?, *Chem Geol*, 202, 479-506, 10.1016/j.chemgeo.2003.03.001, 2003.
- 1025 Winnick, M. J., and Maher, K.: Relationships between CO₂, thermodynamic limits on silicate weathering, and the strength of the silicate weathering feedback, *Earth Planet Sc Lett*, 485, 111-120, 10.1016/j.epsl.2018.01.005, 2018.
- Wolery, T. J., Jackson, K. J., Bourcier, W. L., Bruton, C. J., Viani, B. E., Knauss, K. G., and Delany, J. M.: Current Status of the Eq3/6 Software Package for Geochemical Modeling, *Acs Sym Ser*, 416, 104-116, 1990.
- 1030 Wu, Z. T., Dijkstra, P., Koch, G. W., Penuelas, J., and Hungate, B. A.: Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation, *Glob. Change Biol.*, 17, 927-942, 10.1111/j.1365-2486.2010.02302.x, 2011.
- Zhang, Y. H., Niu, J. Z., Yu, X. X., Zhu, W. L., and Du, X. Q.: Effects of fine root length density and root biomass on soil preferential flow in forest ecosystems, *Forest Systems*, 24, 10.5424/fs/2015241-06048, 2015.
- Zhi, W., Li, L., Dong, W. M., Brown, W., Kaye, J., Steefel, C. I., and Williams, K. H.: Distinct Source Water Chemistry Shapes Contrasting Concentration - Discharge Patterns, *Water Resour Res*, 55, 4233-4251, 10.1029/2018wr024257, 2019.
- 1035 [Zhi, W., and Li, L.: The Shallow and Deep Hypothesis: Subsurface Vertical Chemical Contrasts Shape Nitrate Export Patterns from Different Land Uses](#), *Environ Sci Technol*, 10.1021/acs.est.0c01340, 2020.
- Zhou, X. H., Talley, M., and Luo, Y. Q.: Biomass, Litter, and Soil Respiration Along a Precipitation Gradient in Southern Great Plains, USA, *Ecosystems*, 12, 1369-1380, 10.1007/s10021-009-9296-7, 2009.
- 1040 [Zhu, H., Zhang, L. M., and Garg, A.: Investigating plant transpiration-induced soil suction affected by root morphology and root depth](#), *Computers and Geotechnics*, 103, 26-31, 10.1016/j.compgeo.2018.06.019, 2018.