

Patterns in recent and Holocene Pollen Accumulation Rates across Europe; the Pollen Monitoring Programme Database as a tool for vegetation reconstruction

Vojtěch Abraham¹, Sheila Hicks², Helena Svobodová-Svitavská^{1, 3}, Elissaveta Bozilova⁴, Sampson Panajiotidis⁵, Mariana Filipova-Marinova⁶, Christin Eldegard Jensen⁷, Spassimir Tonkov⁴, Irena Agnieszka Pidek⁸, Joanna Świąta-Musznicka⁹, Marcelina Zimny¹⁰, Eliso Kvavadze¹¹, Anna Filbrandt-Czaja¹², Martina Hättestrand¹³, Nurgül Karlıoğlu Kılıç¹⁴, Jana Kosenko¹⁵, Maria Nosova¹⁶, Elena Severova¹⁵, Olga Volkova¹⁵, Margrét Hallsdóttir¹⁷, Laimdota Kalniņa¹⁸, Agnieszka M. Noryśkiewicz¹⁹, Bożena Noryśkiewicz²⁰, Heather Pardoe²¹, Areti Christodoulou²², Tiiu Koff²³, Sonia L. Fontana^{24, 25}, Teija Alenius²⁶, Elisabeth Isaksson²⁷, Heikki Seppä²⁸, Siim Veski²⁹, Anna Pędziszewska⁹, Martin Weiser¹, and Thomas Giesecke³⁰

¹Department of Botany, Faculty of Science, Charles University; Benátská 2; CZ-128 01; Prague; Czech Republic

²P.O. Box 8000, FI-90014 University of Oulu, Finland

³Institute of Botany v.v.i.; Czech Academy of Sciences; Zámek 1; CZ-252 43 Průhonice; Czech Republic

⁴Laboratory of Palynology, Department of Botany, Faculty of Biology, Sofia University, 8 Dragan Tsankov blvd., Sofia 1164, Bulgaria

⁵Lab. of Forest Botany, Faculty of Forestry and Natural Environment, Aristotle University of Thessaloniki, P.O. Box 270, 54124 Thessaloniki, Greece

⁶Museum of Natural History Varna, 41 Maria Louisa Blvd. 9000 Varna; Bulgaria

⁷University of Stavanger, Museum of Archaeology, Peder Klows gate 31A, PB 8600 Forus, NO-4036 Stavanger, Norway

⁸Institute of Earth and Environmental Sciences, Maria Curie-Skłodowska University; al. Krasnicka 2d; 20-718 Lublin; Poland

⁹University of Gdańsk, Faculty of Biology, Department of Plant Ecology, Laboratory of Palaeoecology and Archaeobotany, ul. Wita Stwosza 59, 80-308 Gdańsk, Poland

¹⁰Białowieża Geobotanical Station, Faculty of Biology, University of Warsaw, Sportowa 19, 17-230 Białowieża, Poland

¹¹Georgian National Museum, Putseladze Str.3, Tbilisi 5, Georgia 0105.

¹²Faculty of Biological and Veterinary Sciences, Geobotany and Landscape Planning, Nicolaus Copernicus University in Toruń, 87-100 Toruń, ul. Lwowska 1; Poland

¹³Department of Physical Geography, Stockholm University, SE-106 91 Stockholm, Sweden

¹⁴Department of Forest Botany, Faculty of Forestry, Istanbul University-Cerrahpaşa; Bahçeköy; TR-34473; Istanbul; Turkey

¹⁵Department of Higher Plants, Moscow State University; Leninskie Gory, 1, 12, Moscow, 119234, Russia

¹⁶Main Botanical Garden RAS; Botanicheskaya, 4, Moscow, 127276, Russia

¹⁷Laugarnesvegi 87 íbúð 105, 105 Reykjavík, Iceland

¹⁸Department of Geography and Earth Sciences, University of Latvia; Jelgavas Street 1, LV-1004; Riga, Latvia

¹⁹Institute of Archeology, Faculty of History, Nicolaus Copernicus University in Toruń; Szosa Bydgoska 44/48; 87-100 Toruń; Poland

²⁰Faculty of Earth Sciences and Spatial Management, Nicolaus Copernicus University in Toruń; Lwowska 1, 87-100 Toruń; Poland

²¹Department of Natural Sciences, National Museum Wales, Cathays Park, Cardiff, CF10 3NP, U.K.

²²Department of Forests, Ministry of Agriculture, Rural Development and Environment, P. Box 24136, 1701 Nicosia, Cyprus

²³Tallinn University, School of Natural Sciences and Health, Institute of Ecology, senior researcher. Uus Sadama 5, 10120 Tallinn, Estonia

²⁴Cátedra de Palinología, Facultad de Ciencias Naturales y Museo, UNLP, Calle 64 n°3, 1900 La Plata, Argentina

²⁵Faculty of Resource Management, HAWK University of Applied Sciences and Arts, Büsingenweg 1a, 37077 Göttingen, Germany

²⁶Turku Institute for Advanced Studies (Department of Archaeology), FI-20014 University of Turku

²⁷Norwegian Polar Institute, Fram Centre, N-9296 Tromsø, Norway

²⁸Department of Geosciences and Geography, University of Helsinki, Gustav Hållströmin katu 2, 00014, Helsinki, Finland

²⁹Department of Geology, Tallinn University of Technology, TalTech, Ehitajate tee 5, 19086 Tallinn, Estonia

³⁰Palaeoecology, Department of Physical Geography, Faculty of Geosciences, Utrecht University, P.O. Box 80115, 3508 TC Utrecht, The Netherlands.

Correspondence: Vojtěch Abraham (vojtech.abraham@gmail.com)

Abstract. The collection of modern, spatially extensive pollen data is important for the interpretation of fossil pollen ~~diagrams.~~ ~~Such assemblages and the reconstruction of past vegetation communities in space and time.~~ Modern datasets are readily available for percentage data but lacking for pollen accumulation rates (PAR). Filling this gap has been the motivation of the pollen monitoring network, whose contributors monitored pollen deposition in modified Tauber-traps for several years or decades across ~~European latitudes~~ Europe. Here we present this monitoring dataset consisting of 351 trap locations with a total of 2742 annual samples covering the period from 1981 to 2017. This dataset shows that total ~~PARs~~ PAR are influenced by forest cover and climate parameters, which determine pollen productivity and correlate with latitude. Regional forest cover >80% is indicated by >3200 tree pollen grains cm⁻² y⁻¹. Pollen traps situated beyond 200 km of the distribution of a given tree species ~~are still collecting~~ still collect occasional pollen grains of that species. PAR's of up to 30 grains cm⁻² y⁻¹ in fossil diagram should therefore be interpreted as long distance transport from beyond 200 km from the area of distribution. Comparisons between modern and fossil PAR from the same regions show ~~comparable~~ similar values. Comparisons for temperate taxa often demonstrate that similar ~~high values for temperate taxa in fossils sites~~ trap values are found further south or downhill. While we do not find modern situations comparable to high fossil PAR values of some taxa (e.g., *Corylus*) ~~do not occur~~. CO₂ fertilization and land use may cause high modern PAR 's that are not documented in the fossil record. The modern data is now ~~publically~~ publicly available in the Neotoma Paleoecology Database (<https://apps.neotomadb.org/explorer/>) and serves ~~improving and aids~~ interpretations of fossil PAR data.

Copyright statement. The article is distributed under the Creative Commons Attribution 4.0 License. Unless otherwise stated, associated published material is distributed under the same licence.

1 Introduction

20 1.1 The need for a dataset of modern absolute pollen deposition

Pollen ~~analytical results are usually reported as percentages, which~~ analysis became the widely used method for the reconstruction of the Holocene vegetation. Pollen percentages are a simple ~~and robust representation of the data, while they~~ representation of pollen analytical results, but have a number of well-known ~~biases.~~ The problem of limitations and biases that are often

ignored. One of these is separating locally produced from long distance transported pollen ~~has received attention from the very beginning of modern pollen analysis (Davis, 2000). Particularly at the tree-line the interpretation (Davis, 2000), which is paramount for mapping past changes in plant distributions. Also reconstructing the position of tree-lines~~ from pollen percentage data may be misleading as local treeless vegetation (e.g. ~~tundra) is often low pollen procuring, tundra produces few~~ pollen grains, while distant woodlands (e.g. ~~consisting of boreal trees) may produce much pollen~~. In such situations absolute pollen data are very informative ~~as was already realized by Hesselman (1919) and Malmström (1923)~~. Pollen accumulation rates (PAR) or the number of pollen grains deposited on the sediment surface over a set period of time ~~is better suited, are in theory superior to pollen percentages as they do not suffer the closure effect of percentage data. Thus by using absolute data it is possible~~ to differentiate between ~~low amounts of~~ long distance transported ~~and versus high amounts of~~ locally produced pollen ~~as already realized by (Hesselman, 1919) and (Malmström, 1923)~~. In a seminal publication ~~(Davis and Deevey, 1964) Davis et al. (1964)~~ document the power of using absolute pollen deposition for the interpretation of ~~vegetation change the spread of trees~~ during the postglacial afforestation around Rogers Lake in Connecticut, USA. Another bias in percentage data is the interdependence of values obscuring the quantification in the amount of change of a single taxon. ~~PARs~~ PAR are therefore required when studying the population dynamics of individual trees (Bennett, 1983). ~~While absolute pollen data do not share the artefacts of percentage data, they are often difficult to obtain and subject to a different set of limitations.~~

~~However, obtaining PAR from sediments cores requires One limitation for obtaining reliable PAR from sediment cores is the requirement for~~ accurate chronologies. ~~While sedimentation in lakes occurs generally without abrupt changes, lake Lake~~ internal processes such as re-deposition and sediment focusing and also catchment erosion may bias the resulting values ~~Davis and Brubaker (1973); (1984); Giesecke and Fontana (2008); (Bennett and Buck, 2016) (Davis and Brubaker, 1973; 1984; Giesecke a~~

. These are some reasons why advances in interpreting PAR have been slow. The other reason is that collecting modern PAR values is not as simple as collecting mosses, soil litter or the top sediment of lakes for obtaining modern pollen percentages for a particular vegetation type. Modern rates of pollen accumulation can be obtained from monitoring pollen deposition using pollen traps (Hicks, 1994), as well as from carefully sampling the top ~~sediments~~ sediment of lakes that are either annually laminated or precisely dated (Matthias and Giesecke, 2014). Due to the high inter-annual variability in pollen production (Andersen, 1980; Haselhorst et al., 2020), it is necessary to conduct pollen monitoring over several years to enable comparisons with estimates from sediment cores (Hicks, 1974; Hicks and Hyvärinen, 1999).

For these reasons there are ~~numerable studies investigating only a few investigations of~~ the pollen vegetation relationship using ~~percentage data and comparably few using~~ absolute pollen deposition, ~~while there are numerous studies using percentage data~~. Nevertheless, investigations using pollen traps yielded invaluable insights ~~in~~ into the mode of pollen transport (Tauber, 1967). Also the construction of representation factors for common Europe trees by Andersen (1970), which are still used, ~~were was~~ based on pollen monitoring data from pollen traps. In this way pollen monitoring studies have contributed to the development of models of pollen dispersal and deposition (Gaillard et al., 2008).

~~More recent~~ Several aspects of PAR data have not been exhaustively explored in modern comparison studies and here we will focus on the following three: 1) the influence of climate in combination with forest biomass; 2) the application of PAR to

indicate the local presence of trees versus long distance transport of pollen; using modern PAR values of single taxa to interpret fossil situations.

1.2 Climate and biomass

Recent investigations demonstrate the linear response of ~~the~~ absolute pollen deposition to absolute tree abundance (Seppä et al., 2009, Sugita et al. (2009), Matthias and Giesecke (2014)), which may be used to reconstruct past standing tree biomass of different trees. However, ~~rather than changes in tree biomass, changes in PAR~~ at an annual time resolution ~~correspond to weather conditions of the previous,~~ variability in PAR can be explained by weather conditions during the time of flowering as well as ~~the year of flowering during the previous year~~ (Hicks, 1999, van der Knaap et al. (2010), Nielsen et al. (2010)). Thus the question arises: If weather is determining annual pollen production, could climate ~~or site conditions not determine average~~ rates in addition to determining biomass determine average PAR? Comparing *Pinus* PAR between two pine dominated forest regions in central Sweden and north east Germany shows much higher values in the south, ~~suggest~~ suggesting that PAR may not correspond to tree biomass alone (Matthias and Giesecke, 2014). The relationship between pollen production and weather suggests that more pollen is produced when the primary productivity of the tree is higher. This is also true for fertilization with CO₂ (LaDeau and Clark, 2006). Therefore, climate ~~, soil fertility~~ and even the amount of CO₂ in the atmosphere may determine the pollen productivity of a tree at a given site. ~~Already Welten (1944)~~ Welten (1944) already interpreted the first fossil PAR in this way, suggesting that climate deteriorations may not immediately lead to a decline in forest cover but to the amount of pollen produced. This interpretation of changing PAR was however forgotten. If climate and CO₂ determine pollen productivity than the postglacial increase in PAR at Rodgers Lake could also be due to a change in these parameters. It is ~~therefore impotent~~ , therefore, important to investigate the possible relationship between climate and average PAR in more detail.

20 1.3 Local presence versus long distance transport

Also the initial question on the amount of pollen that may arrive at a site from long distance sources has not been addressed in a systematic way using modern absolute pollen deposition data. Occasional reports of PAR values for the first occurrence of a macrofossil indicating local presence exist (e.g. Giesecke, 2005a), but modern comparisons are lacking. While percentage data are not well suited for detecting distribution limits, a continental scale comparison (Lisitsyna et al., 2011a) provides some guidance on values that can be used for mapping past distribution changes (Giesecke et al., 2017). The PMP dataset presented here provides a continental scale dataset permitting such a comparison with PAR data.

1.4 Modern analogies

Spatially extensive modern pollen percentage datasets provide the possibility of searching for modern analogues for fossil pollen proportions and in this way reconstructing past vegetation and environmental conditions assemblages (Davis et al., 2013; Jackson and Williams, 2004; Overpeck et al., 1985). Modern datasets of absolute pollen deposition are hitherto rarely used ~~as references helping~~ to reconstruct past tree abundances or environmental conditions. By using a network of pollen traps

across the latitudinal tree-line in Finland, Hicks et al. (2001) ~~show~~showed that average modern PAR values can be obtained representing the gradual transition from the boreal forest to tundra. These “modern analogues” were successfully applied to reconstructing Holocene shifts of the latitudinal tree line (Seppä and Hicks, 2006). This idea of building a modern dataset of absolute pollen deposition that can be used as a reference to interpret fossil PAR was the motivation for the establishment of the Pollen Monitoring Programme (PMP, Hicks et al., 1996; 1999; 2001).

1.5 The Pollen Monitoring ~~Program~~Programme (PMP)

The ~~program~~programme was launched in August 1996 at a meeting in Finland, bringing mainly European researchers together. Members of the network changed over the years and monitoring experiments were discontinued or newly started. Although pollen monitoring studies were and are carried out ~~on~~in other continents (e.g. Jantz et al., 2013), the PMP had little success in attracting researchers working outside Europe. The standardisation of the monitoring protocol allowed for easy comparisons between the results in different regions, which were discussed at INQUA in 1999 and led to a special volume published in 2001 (Tinsley and Hicks, 2001) collecting results based on several initial time series (van der Knaap et al., 2001; Koff, 2001; Tinsley, 2001; Tonkov et al., 2001), as well as a first comparative study (Hicks et al., 2001). More individual results were published in the following years (e.g. Gerasimidis et al., 2006; Giesecke and Fontana, 2008; Hätteland et al., 2008; Jensen et al., 2007; Kvavadze, 2001; Pidek, 2007) and comparative studies followed in a second special volume published in 2010 (Giesecke et al., 2010). The data produced by contributors to the PMP were analysed for different questions, including weather parameters determining the amount of pollen production (van der Knaap et al., 2010) and its correlation to masting years in *Fagus* (Pidek et al., 2010). The programme established a database collecting the original data for individual years, as well as general information on the pollen traps installed in the different regions (Fig. 1). The database was developed offline and was thus difficult to access by individual researchers. The paleoecology database Neotoma (Williams et al., 2018) offers a platform to store the PMP data and make it available to researchers worldwide, allowing them to interrogate the data and potentially identify modern analogues to interpret fossil pollen accumulation rates.

The overall purpose of this manuscript is to present an overview of the data in the PMP database and to interrogate this continental scale dataset of modern PAR with the following aims:

- a) Examine the hypothesis that climate as well as regional forest cover explain the variability in PAR.
- b) Study the absolute amount of long distance dispersed pollen encountered in pollen traps beyond the known distribution limits of the parent trees.
- c) Compare modern and fossil pollen accumulation rates by collecting fossil datasets with estimates of PAR from the same regions where the pollen traps were installed. For the most abundant pollen types we explore how the modern situations can provide a reference for the interpretation of the fossil data.




Figure 1. Map of the study area with trap areas (a) and Holocene pollen sites (b). Both datasets are divided into trap regions (colours). – Colours correspond to Fig. 5.

2 Methods

2.1 Study area

Sites in the PMP database were divided into 7 ‘trap regions’ according to ~~the~~their latitude and altitude. These regions were further divided into ‘trap areas’, by grouping 2-10 trap locations according to their spatial proximity. The ~~aretie~~Arctic/alpine Alpine region includes distant trap areas in Spitsbergen and Iceland, northernmost traps in Finland (Utsjoki) and Norway (Lofoten-Vesterålen) and traps in the Scandinavian mountains above the local tree-line (Finnmark, Abisko-Tarfala, Skrubben and Lapps-kardet). The landscape around these traps is often treeless or covered by sparse birch shrubland with *Betula nana* and *B. pubescens* in some locations. The ~~north-boreal~~North Boreal region includes traps in Northern Lapland in Finland, Norway and Sweden with a vegetation dominated by *Betula* and *Pinus sylvestris*. The northern limit of *P. sylvestris* occurs between traps from this region, which are situated at altitudes below 500 m a.s.l. The ~~boreal~~Boreal region includes trap areas situated in Southern Lapland, around the Bay of Bothnia and in Central Sweden. The vegetation is dominated by *Picea abies*, *P. sylvestris* and *Betula* species, with the occurrence of *Alnus incana*. Northernmost populations of *Alnus glutinosa* occur near some sites and the southernmost traps in Central Sweden are situated near northern outpost populations of *Corylus avellana* and *Ulmus glabra*. Traps from the ~~temperate-lowland~~Temperate Lowland region have the widest longitudinal extent including the British Isles, Poland, the Baltic countries and European Russia. Vegetation at trapping locations below 500 m a.s.l. is characterised by *Quercus* and *Fraxinus excelsior* in the ~~West-west~~West-west and *P. sylvestris*, *P. abies* and *Betula* with an admixture of *Quercus*, *Tilia*, *Ulmus* and *C. avellana* in the ~~East~~east. *Fagus sylvatica* and *Carpinus betulus* occur in Poland, *Abies alba* only in south-eastern

Poland. Trapping locations in the ~~temperate region at elevations above 500 m (mid elevation)~~ Mid Altitude Temperate region were separated and include the Krkonoše and Šumava Mountains. Traps in both areas are placed on an elevation gradient from 500 m and 1200 m ~~a.s.l.~~ a.s.l.. The lower slopes of the mountains are dominated by ~~F.~~ F. *Fagus sylvatica*, while the traps are situated on a gradient from ~~P.~~ P. *Picea abies*-dominated forest to the onset of alpine vegetation in Krkonoše. In the Alps and

5 Jura Mountains (Alpine region), traps were placed at even higher elevations, between 1200 m to 3000 m a.s.l., crossing the altitudinal treeline. Trapping locations in ~~the Southeast~~ Temperate/Mediterranean region in the southeastern Europe represent diverse landscapes and vegetation types including grasslands, evergreen and deciduous forests. Some traps are situated in high mountain regions around treeline situations or within the upper mountain forests including Rila (Bulgaria), Pieria (Greece), Timfristos (Greece), Lagodheki (Georgia) and Cyprus. Traps at lower elevations are situated near the Black Sea coast, within

10 the low Strandzha Mountains and European Turkey.

2.2 Data collection

The pollen traps used in the PMP network generally consist of a bucket or bottle large enough to contain the annual surplus in precipitation on a surface of usually 19.6 cm² (5 cm diameter opening) or similar. Many traps had a sloping collar inspired by the design of pollen traps by Tauber (1974), although few collars were truly aerodynamic. The collection of the trap content

15 was generally carried out annually and any special circumstances potentially affecting the annual pollen deposition were noted and stored in the database. For the analyses presented here and data overview, we excluded traps where the pollen record is 2 years or less, as averages may be affected by high inter-annual variability. The only exception is the trap situated in Spitsbergen where there is a two-year record. Pollen accumulation from the two-year record shows little variation and, being the only analogue for a truly arctic and treeless environment, provides important information on long distance pollen transport. We

20 also excluded annual samples with records shorter than 8 months and, in addition, traps or years with spurious values due to particular events or local conditions (Table S1).

Most of the traps in the PMP network were placed in the open vegetation or in forest openings in order to avoid an unrepresentative contribution of individual trees e.g. due to anthers dropping into the traps. Traps were generally installed at ground level mimicking collection conditions relevant for sedimentary archives. ~~In consequence~~ Consequently, tall herbs or grasses

25 might ~~overgrow or cover~~ have overgrown or covered some of the traps potentially leading to higher pollen deposition. Traps not equipped with a mesh occasionally trapped ~~pollen-collecting~~ pollen-collecting insects, leading to enormous counts of insect pollinated taxa e.g. *Calluna*, *Erythranthe guttata*. The presence of insects in the traps is usually noted for the collection year so that careful evaluation of the information in the database can also inform on herbaceous pollen types (Jensen et al., 2007). Including this information in comprehensive database queries is currently not possible and a manual screening of datasets is

30 required when analysing herbaceous pollen types. This problem does not seem to occur in tree pollen taxa. The occurrence of phytophagous insects in the traps were not accompanied by unusual peaks in tree pollen taxa, indicating that the insects inadvertently trapped were primarily collecting pollen from the herbaceous vegetation around the traps. Comprehensive database queries restricted to tree pollen, Poaceae and Cyperaceae should therefore not be affected by the occurrence of insects in the

trap and mainly represent pollen transport via wind, the rainout of pollen from the atmosphere and the gravity component (Tauber, 1967).

Concentrating the content of the traps was carried out either using filter paper or centrifugation and decanting the supernatant. In many cases the trap content was washed onto a paper filter, which was later digested using acetolysis. Pollen quantity was assessed by adding *Lycopodium* spore tablets (Stockmarr, 1971) to each trap before processing. Pollen concentration was obtained from the ratios between pollen grains counted to *Lycopodium* spike counted and *Lycopodium* spike added. Details about *Lycopodium* spike data, as well as details of the pollen trap such as the exact size of the opening are stored in the database. The PMP database was created in the PostgreSQL database system. Names of pollen taxa were unified using the accepted variable names from the European Pollen Database (Giesecke et al., 2019).

2.3 Investigated taxa, climate and ~~environmental parameters~~forest cover

We selected the common tree and shrub taxa of Europe. Pollen taxa generally refer to all the species within the genus. Pollen taxa allowing higher taxonomical resolution, which were consistently separated and excluded from the genus in the whole dataset are marked as “excl.”. Pollen taxa potentially including pollen grains from another genus are indicated by “incl.”: *Abies*, *Alnus* (excl. *A. viridis*), *Betula* (excl. *B. nana*-type), *Carpinus*-type (incl. *C. orientalis*/*Ostrya*-type), *Corylus*, *Fagus*, *Fraxinus* (incl. *F. ornus*), *Juniperus*-type (incl. *Cupressus*, *Tetraclinis*, *Thuja*), *Picea*, *Pinus* (excl. *P. cembra*-type), *Tilia*, *Quercus* (incl. *Q. robur*-type, *Q. cerris*-type and *Q. ilex*-type). Pollen accumulation rates of trees and shrubs were summed as arboreal pollen accumulation (hereafter as “tree PAR”). We also included pollen from the plant families Cyperaceae and Poaceae (excluding cereals). For the purpose of the analysis in this paper we refer to the sum of tree PAR plus Cyperaceae and Poaceae as “total PAR”.

The climate parameters Mean Annual Temperature (MAT) and Annual Precipitation (APrecip) for the trapping locations were obtained from WorldClim 2 (Fick and Hijmans, 2017). For site altitude we used the information supplied by the individual investigator. Comparisons between PAR and forest cover were conducted using the data of the Forest Map of Europe (Kempeneers et al., 2012), which has a grid resolution of 1 km². Forest cover was extracted as a mean of all grid cells within a 10 km radius. We used regression analysis to explore whether individual or combinations of these environmental parameters describing the trapping location can explain the variance in average pollen accumulation of ~~the traps~~total and tree PAR. To balance the contribution of high and low pollen producers in the assessment of the ~~total-PAR~~PAR, we applied correction factors (Table S2, Andersen, 1970). Average modern PAR have a large variance of values between traps of the same region, with the smaller numbers often being the focus of information. For this reason, we often log-transformed PAR values in the different analyses.

2.4 Distribution limits and PAR

Pollen deposition beyond the distribution area of the parent plant was studied by merging the distribution maps of the relevant species included in ~~the pollen type described~~each of the pollen types listed above (Caudullo et al., 2017; San-Miguel-Ayanz et al., 2016). These comparisons were not ~~suitable~~possible for *Alnus*, *Betula*, Cyperaceae, *Juniperus*, *Pinus* and Poaceae as

these taxa are widely distributed in Europe and few traps are located beyond their distribution area. ~~We compared the amount of pollen accumulating from these taxa in pollen traps~~

For each trap location and each pollen taxon we calculated the distance to the nearest area of distribution using GIS (GRASS Development Team, 2018). Initial observations showed that PAR dropped rapidly away from the distribution of the parent tree and did not decline at the same rate at larger distances. We therefore compared distance to the decadic logarithm of PAR, applying linear regression to explore thresholds of long-distance transport (hereafter also as “LDT”) at 200 km from their mapped distribution limits. This distance was chosen as a compromise accounting for uncertainties in the information on distribution limits and available pollen trap data. Pollen traps in the UK are situated beyond the natural distribution limits of several of these trees but were excluded from the comparison as the ~~taxa considered target taxa~~ may be planted in the area.

~~For each trap location and each pollen taxon we calculated the distance to the nearest area of distribution using GIS (GRASS Development Team, 2018). Linear regression between this distance and the decadic logarithm of PAR was used to predict the threshold of long-distance transport (hereafter also as “LDT”). We compare LDT with the characteristic radius for the same taxa. Characteristic radius is a useful measure showing pollen transport predicted by pollen dispersal models. It represents the theoretical proportion of pollen loading at different distances from the source plants (Prentice, 1988) and can thus be easily compared with our empirical values. We used Gaussian Plume model with wind speed 3 m.s⁻¹ (Prentice, 1985).~~

2.5 Comparison between modern and Holocene PAR

To enable the comparison of modern with fossil PAR values the pollen trap data was extracted from the PMP database with above described constraints and all annual samples were averaged within traps. ~~Per~~ For each trap region we selected at least one and a maximum of three Holocene PAR records (Table 1). This distance was chosen as a compromise accounting for uncertainties in the information on distribution limits and available pollen trap data. Holocene PAR estimates often show high variation between samples due to changes in the sedimentary environment. To reduce this effect in this comparison, Holocene data were averaged in 500-year bins. Site and sample compilation resulted in a fossil dataset containing 354 Holocene samples.

We compared trap and fossil PAR datasets in two ways. ~~First, we compared the distribution of logged PAR values and their mean for all taxa in all regions. Differences were measured by~~ For a general comparison of the distribution of fossil and modern values, we plotted the frequency of log-transformed PAR values over all regions and traps per taxon. Per region and taxon we compared the frequency distribution of fossil and modern PAR values using a t-test at 5% level of alpha. ~~Second, we searched for Holocene pollen records containing estimates of PAR in proximity to the locations of pollen traps, so we to identify situations where modern and fossil values are comparable. Secondly, we~~ compared trap and fossil PAR at the level of individual sites or trap areas. ~~Average~~ To facilitate this comparison average trap and fossil PAR values per taxon were submitted to one-dimensional clustering using the R-package Ckmeans.1d.dp (Song and Wang, 2011). ~~The classes produced were used to facilitate the comparison between~~ This method splits the univariate data in the way that the total of within-cluster sums of squares is always minimum. These classes helped us compare trap and fossil data and to ~~match the trap values with analogous situations in the past. The aim of this comparison was to find traps with similarly high values for individual taxa that compared to the highest average fossil PAR~~ link the highest fossil PARs with the trap PARs. We dealt only with the highest

Table 1. Fossil sequences including type and size of the deposit.

| country | region | site | deposit | latitude | longitude | (m) a.s.l. | area (ha) | reference |
|---------|---|----------------|---------|-----------|-----------|------------|-----------|--------------------------------------|
| FIN | aretic <u>Arctic</u> / alpine <u>Alpine</u> | Bruvatnet | lake | 70.17933 | 28.39998 | 119 | 60 | (Hyvärinen, 1975) |
| FIN | aretic <u>Arctic</u> / alpine <u>Alpine</u> | Toskaljavri | lake | 69.19177 | 21.44841 | 704 | 100 | (Seppä et al., 2002) |
| FIN | aretic <u>Arctic</u> / alpine <u>Alpine</u> | Tsuolbmajavri | lake | 68.68915 | 22.05235 | 256 | 14 | (Seppä and Weckström, 2002) |
| FIN | north-boreal <u>North Boreal</u> | Suovalampi | lake | 69.58333 | 28.83333 | 104 | 16 | (Hyvärinen, 1975) |
| FIN | north-boreal <u>North Boreal</u> | Akuvaara | lake | 69.125 | 27.68333 | 170 | 4 | (Hyvärinen, 1975) |
| SWE | boreal <u>Boreal</u> | Abborrtjärnen | lake | 63.88333 | 14.45 | 387 | 3 | (Giesecke, 2005c) |
| SWE | boreal <u>Boreal</u> | Klotjärnen | lake | 61.81667 | 16.53333 | 235 | 1 | (Giesecke and Fontijn-Tekamp, 2005a) |
| SWE | boreal <u>Boreal</u> | Holtjärnen | lake | 60.65 | 14.91667 | 232 | 1 | (Giesecke, 2005a) |
| EST | lowland-temp <u>Lowland Temp.</u> | Rõuge Tõugjärv | lake | 57.73904 | 26.90515 | 114 | 4.2 | (Veski et al., 2012) |
| POL | lowland-temp <u>Lowland Temp.</u> | Suminko | lake | 53.72556 | 17.77278 | 115 | 0.12 | (Pędziszewska et al., 2018) |
| CZE | middle-alt.-temp <u>Middle Alt. Temp.</u> | Prášilské | lake | 49.07551 | 13.40002 | 1079 | 3.7 | (Carter et al., 2018) |
| CZE | middle-alt.-temp <u>Middle Alt. Temp.</u> | Malá niva | peatbog | 48.90789 | 13.81982 | 754 | 65 | (Svobodová et al., 2018) |
| CHE | mountain-temp. <u>Alpine</u> | Sägistalsee | lake | 46.68139 | 7.9775 | 1935 | 7.2 | (van der Knaap et al., 2002) |
| CHE | mountain-temp. <u>Alpine</u> | Bachalpsee | lake | 46.66944 | 8.020833 | 2265 | 8 | (van der Knaap et al., 2002) |
| BGR | temp <u>Temp./medit</u> <u>Medit.</u> | Shabla | lake | 43.58333 | 28.55 | 1 | 1.51 | (Filipova-Marinova et al., 2018) |
| BGR | temp <u>Temp./medit</u> <u>Medit.</u> | Arkutino 2 | lake | 42.3299 | 27.72363 | 0 | 40 | (Bozilova and Beugnot, 2018) |
| BGR | temp <u>Temp./medit</u> <u>Medit.</u> | Ribno | lake | 42.20682 | 23.32346 | 2184 | 3.5 | (Tonkov et al., 2000) |
| GRC | temp <u>Temp./medit</u> <u>Medit.</u> | Voukaria | lake | 38.866667 | 20.833333 | 0 | 10000 | (Jahns, 2004) |

class in each fossil sequence because maximum abundance of several our target taxa was used as a stratigraphic marker of the Holocene period and thus their timing is well known. However, averaging time windows did not smooth out all spuriously high values variation, ~~thus we need to ignore individual.~~ In order to remove this remaining variation of individual time windows, we ignored some high fossil values (Table S3). Thus, we aimed to find modern analogues for fossil situations represented by several bins (more than 500 years). We linked these periods ~~with high fossil PARs~~ containing high fossil PAR to the closest pollen trap, using a matrix of geographical distances between fossil sites and pollen traps. ~~We selected the comparisons for the 8 most abundant taxa to be presented in the main text and show the remaining 7 taxa in the supplementary.~~ All statistical analysis and data visualizations were produced in R (R Core Team, 2019).

3 Results and interpretations

10 3.1 Overview of the PMP database and the environments sampled

The PMP database version 02.02.2020 contains data from 351 trap locations with a total of 2742 annual samples covering the period from 1981 to 2017. Considering the trap ~~reecord~~ records with 3 years and more we obtained 271 mean trap assemblages.

Trapping sites cover a range of altitudes from 0 to 3000 m a.s.l. with annual precipitation ranging from 402 to 1549 mm. Mean annual ~~temperature~~ temperatures (MAT) for the sites fall between -5.7 to 14.1 °C. The forest cover within a 10 km radius of the trapping sites ranges from 0 to 98%. This range of environmental situations has yielded tree pollen accumulation rates from 5 to 86000 grains cm⁻² y⁻¹, with a median value of 5400 grains cm⁻² year⁻¹ (Fig. 2). An overview of the taxonomic composition of the traps (Fig. S1) shows a dominance of pollen from *Pinus* and *Betula* in the traps from boreal and hemiboreal environments, with *Betula* as the taxon with the highest ~~PARs~~ PAR overall. In most northern traps from open environments, Cyperaceae is the dominant NAP pollen type, while Poaceae are dominant in traps from open environments in the south, where they also contribute much higher absolute amounts. The diversity of landscapes and forest types in central and southern Europe is well represented in the pollen composition of traps from this area. ~~Differences in pollen composition and abundance between the high mountain forests of central and southern Europe and the boreal forest in the north are also noticeable~~ (Fig. S1b). Dominance of oak and hornbeam at Temperate/Mediterranean sites in the lowland and pine and birch at Arctic/Alpine and North Boreal sites show similar stability in the Holocene perspective. Vegetation history at the rest of the fossil sites shows a more dynamic development (Fig. S2).

3.2 Dependence of variation in PAR on regional forest cover and climate

Total PAR is generally lower at high latitudes, with the lowest values in the ~~arctic-alpine~~ Arctic/Alpine region (trap area Spitsbergen), where no trees can grow. However, the highest absolute values of tree PAR are not from the southernmost traps but from the ~~lowland-temperate~~ Lowland Temperate region (trap area Tver; Fig. 2). ~~Nevertheless, latitude~~ Latitude alone explains about 11% of the variance in ~~log-transformed~~ log-transformed tree PAR, while MAT and forest cover within 10 km explain 21% and 19% respectively. In combination, these three variables explain 37% of the variation in log-transformed absolute tree pollen deposition. The addition of elevation increased the amount of variance explained to 50% (Table S4a).

Large differences in the pollen productivity between different trees ~~effect~~ affect this relationship. Adjusting the PAR from individual taxa by Andersen factors reduces the bias of differential pollen production between different plants and makes it possible to consider the total amount of pollen deposition including grasses. This adjustment increases the amount of variance explained by the regression model with all 4 explanatory variables to 56% (Table S4b). Due to the inclusion of grasses, the explanatory power of forest cover within 10 km is reduced, while latitude alone explains 37% of the Andersen adjusted ~~log transformed~~ log-transformed total PAR (Fig. 3a, Table S4b).

The regression models consider the full range of the data ~~, while~~ while, due to local factors, there is often a spread of average trap values for different traps in the same region. The traps with the highest regional values do not follow a latitudinal pattern, so the distribution of the minimum average trap values ~~are~~ is more informative (Fig. 3a). These lower values closely follow a latitudinal trend. The average PAR south of 62° latitude and below the altitudinal treeline or close to forests ~~are~~ is generally higher than 1000 grains cm⁻² y⁻¹. An area with low PAR in the south is the coastal grassland in northern Bulgaria. The generally low PAR in this area can be explained by the sparse vegetation cover on thin rendzina soils formed on limestone rock. Adjusting the PAR values by Andersen factors increases the values for this region so that they fit the general latitudinal trend (Fig. 3a). Traps with minimum average PAR values per region also correspond well to the forest cover within 10 km (Fig.




Figure 2. Environmental setting of the trap dataset. (a) trap areas ordered from north (left) to south (right), (b) altitude, (c) mean annual tree PAR, (d) forest cover within 10 km radius, (e) annual precipitation, (f) temperature: MAT - Mean Annual Temperature, (g) temporal coverage of the PMP database.




Figure 3. Latitudinal gradient in pollen accumulation rates of major tree taxa and Poaceae and Cyperaceae (total PAR) adjusted by Andersen values (see Tab S2) of pollen representation (a). Relationship between forest cover within 10 km radius and tree PAR (b). All trap sites (1) and minimum tree PAR per every 3 % of forest cover (2).

3b). ~~Considering~~ Exploring the data showed that a 3% wide bins of the forest cover, traps with the lowest PAR per each bin of the forest cover provide a regression model predicting a tree PAR of 3200 grains cm⁻² year⁻¹ at 80% forest cover within 10 km of the trap.

3.3 Long distance dispersed pollen

- 5 The comparison of PAR with the distribution limit of different tree taxa shows that PAR generally ~~decline~~ declines with distance (Fig. 4). A gradual decline is best documented for *Quercus* where ~~average values at the distribution limit scatter around 100 grains cm⁻² y⁻¹~~ traps cover different distances from the distribution area. This analysis also documents the long-distance transport of many tree pollen, including the heavy pollen of *Picea*. For better comparison of the absolute values between taxa ~~we used regression analysis to estimate~~, we fitted a linear relationship, also to compare the amount of pollen
- 10 at 200 km from the distribution limit (Fig. 4b). This comparison indicates that less than ~~30~~ 80 grains cm⁻² y⁻¹ of *Carpinus*, *Corylus*, *Fagus*, *Fraxinus*, *Quercus* and *Tilia* are deposited beyond 200 km of the distribution of the parent trees. Only *Picea*




Figure 4. Relationship between the distance from the trap site to the nearest area of species distribution and PAR for selected trees. Zero distance represents edge of distribution area. Traps within the distribution area are aggregated in boxplots (a). ~~PARs~~ ~~PAR~~ of the long-distance transport (b) calculated from linear regression at 200 km (Fig. 4a).

shows less than 1 grain $\text{cm}^{-2} \text{y}^{-1}$ at 200 km of the distribution range. In the case of *Fagus* it has to be noted that one trap beyond 200 km of the distribution of the tree recorded more than 30 grains $\text{cm}^{-2} \text{y}^{-1}$. The general threshold of 80 grains $\text{cm}^{-2} \text{y}^{-1}$ suggested here is near the detection limit and the PAR value may be biased by the size of the pollen count in cases where only one grain was encountered.

- 5 PAR at 200 km from the distribution area represent 0.002-36% of the median PAR within the distribution area (Fig. 4), which is 0.2% of pollen loading in average for those seven taxa (Table S5).

3.4 Ranges of modern and fossil PAR values

- The ~~here presented~~ comparison of modern and fossil PAR values shows good agreement in tree PAR. The highest frequency of tree PAR values ranges between 2000 and 10000 grains $\text{cm}^{-2} \text{y}^{-1}$ in both datasets (Fig. 5). Maximum ~~PARs~~ ~~PAR~~ of the trap dataset are higher (often ten times) for all species compared to maximum ~~PARs~~ ~~PAR~~ in the fossil dataset, with the exception of *Corylus*. The frequency distribution of ~~PARs is log-normal~~ ~~PAR is log-normal~~ for *Alnus*, *Tilia* and *Fraxinus*. A strong bimodal distribution of values occurs for some taxa, which is particularly clear for the fossil values can be recognized for fossil samples of *Picea* and *Poaceae* (Fig. 5). Such a bimodality, but less clear, occurs for the modern samples of *Pinus* and for fossil samples of *Abies* and *Fagus*. The highest frequency of trap values for *Carpinus*, *Picea* and *Poaceae*. In the modern samples such a bimodality can be recognized for *Fagus* and *Tilia* is very close to the line of the LDT threshold. The LDT threshold for *Pinus* *Corylus* with a trough at around 1000 grains $\text{cm}^{-2} \text{y}^{-1}$. The *Fraxinus* and *Quercus* separates values of the three boreal regions from the rest of the regions. The trap values of *Picea* from the Arctic/Alpine region are above the LDT threshold. The frequency distribution of modern and fossil ~~PARs~~ ~~PAR~~ corresponds best for *Corylus*, with most values falling between 100 - 300 grains $\text{cm}^{-2} \text{y}^{-1}$ and recent and fossil maxima at around 3000 grains $\text{cm}^{-2} \text{y}^{-1}$. The greatest difference in the
- 10
 - 15

distribution of modern versus fossil PARs occurs for *Juniperus*, where maximum values are around a hundred times larger in the traps. Minimal PARs are about ten times higher in the traps for Poaceae and Cyperaceae, in particular, and the right side of the distribution is shifted upwards. For *Fagus*, *Quercus* and *Carpinus* the fossil PARs show a local maximum in the frequency of low values, which does not occur in the traps. These frequent low values range below the threshold indicating long-distance transported pollen.

Using the 15 selected taxa in seven trap regions and the occurrence of the pollen types, we obtained 92 pairs of trap and fossil data. The frequency distribution of PAR values is similar between modern and fossil samples in 31 pairs based on a t-test and a p-value > 0.05 (Fig. 5, Table S5S6). In this regional comparison, *Betula* shows the best agreement between modern and fossil values. Values are similar in four regions across the gradient, with highest values of 5400 grains cm⁻² y⁻¹ in the lowland temperate region and the lowest values of 34 grains cm⁻² y⁻¹ in the temperate/Mediterranean region, where the parent trees are generally absent. Modern and fossil PARs generally correspond well for the lowland temperate region where, in addition to *Betula*, *Alnus*, *Carpinus*, Cyperaceae, *Fraxinus*, *Picea*, *Pinus* and *Quercus* also have similar values. Although *Corylus* has a good overall agreement, the regions with similar modern to fossil data are shifted, with Holocene values in the boreal region corresponding to modern PARs in the lowland temperate region.

3.5 Taxa specific linkage of the highest average PAR at fossil sites with individual trap values

3.5.1 Introductory notes

To facilitate the comparison of modern and fossil PAR, the combined taxa specific values were submitted to a one-dimensional cluster analysis, which resulted in between 5 and 9 classes of PAR values per taxon (Fig. S1-S2). Comparing the maximum averages of fossil PARs to modern trap data on a site-by-site basis shows that it is possible to find modern comparisons for all fossil situations. We demonstrate the linkage of the highest PAR clusters per each fossil site with individual trap records for the example of selected tree taxa (*Abies*, *Betula*, *Corylus*, *Fagus*, *Picea*, *Pinus*, *Quercus*, *Tilia*; Fig. 6). Detailed description of the remaining taxa main linkage. Detailed descriptions are presented in the supplementary (*Alnus*, *Carpinus*, Cyperaceae, *Fraxinus*, *Juniperus*, Poaceae, arboreal pollen; supplement (Fig. S3).

3.5.1

Modern PAR of 490-3900 grains cm⁻² *Abies* declined in most of the populations, thus Roztoce is the only analogous trap area with PAR for fossil sites in central Europe and similarly the Timfristos trap area is analogous for fossil sites in southern Europe (Fig. 6). *Alnus* retreated from Scandinavia, thus the fossil sites have the closest linkage southwards, on the island Hailuoto or in North Poland. *Corylus*, *Quercus* and *Tilia* (Fig. 7) also declined, thus modern values in North Poland provide analogues for fossil situations in central and northern Europe. Poaceae and Cyperaceae exceed values from fossil examples thus analogous traps are close to the fossil site. Similarly, *Betula* and *Juniperus* have high PAR at only a few trap sites (11900-73900 and 870-8300 grains cm⁻² y⁻¹ are observed in Roztoe, Jura, Rila and Timfristos. These values are produced by the different species: *Abies*




Figure 5. Difference (differ.) between the mean fossil (-) and the mean trap (o) PAR per trap region is shown by length of the vertical segment. Paired histograms of mean annual PAR from [the](#) fossil record (on the left) and from traps (on the right). Colours denote different trap regions and correspond to Fig. 1. Note [logged-log-transformed](#) y-axis. Horizontal lines and numbering on the secondary y-axis denote classes of PAR, for more detail see Fig. 6b) and d) -LDT is threshold for long distance transport.

alba in Roztoče and the Jura mountains and *A. cephalonica* in the Rila mountains and on Timfristos. Generally high fossil values occur in the two Alpine lakes Sägistalsee and Bachalpsee around 7000 years, in Ribno in the Rila mountains around 5000 years and in the two sites in Šumava between 4000 and 1000 years. In all these regions with fossil evidence of high *Abies* populations the modern PAR values in pollen traps are comparably low documenting that the populations have much declined.

5 The *Abies alba* populations in Roztoče provide modern analogues for how dense *Abies* forests may have been in the Alps and Šumava although the forests in Roztoče occur at much lower elevations. Pollen traps on Timfristos mountain provide analogues of the density of Middle Holocene *Abies* forests in the Rila mountains. Pollen traps located far from the distribution limits of *Abies* in Wales, northern Poland, Georgia and Cyprus registered occasional *Abies* pollen grains with PARs of up to 80 grains cm⁻², respectively), whereas the fossil PAR and the rest of the trap dataset are lower, fitting within the two or three

10 first classes. Both taxa find an analogous trap record within the same area, most of the distant links are for the lowest classes (0-5600 and 0-90 grains cm⁻² y⁻¹ respectively). Fossil PAR of *Carpinus* in North Poland and the Black Sea coast fall within the range of trap records from the same area. Trap areas in the Baltics, Šumava and Central Sweden indicate higher fossil than trap PAR of *Carpinus*. All fossil sites within the recent distribution of *Fagus* and *Picea* have the closest analogous trap in the same or adjacent trap area and fossil-trap links are short. Fossil and trap PAR of arboreal pollen and *Pinus* are comparable within the

15 trap areas, except for the Baltics and Rila mountains, respectively, where the fossil is lower than the trap record. On the other hand their record from the Early Holocene at Prášílské Lake is higher than any trap record from the Šumava mountains.

3.5.1

Letting aside traps from the Caucasus and Turkey, trap sites cover the distribution of *Betula pendula* and *B. pubescens*. The highest modern values between 11900-73900 grains cm⁻² y⁻¹ are found in Estonia and Russia as well as in one pollen trap from


20 Hailuoto. These modern PARs exceed values from fossil examples. The highest fossil PAR of around 10000 grains cm⁻² y⁻¹ in this comparison come from northern Poland and Estonia and are thus consistent with the area of high modern values. Whether high abundance of *Betula* is a characteristic of the eastern European forests or a result of frequent disturbance at the forest ecotone or due to recent land-use change is difficult to evaluate based on the available data. Modern and fossil values agree for the sites in central Sweden at PARs between 1900-5600 grains cm⁻² y⁻¹.

25 3.5.1

Modern PARs stay below 2800 grains cm⁻² y⁻¹ except for two traps in north-central Poland. Values above 610 grains cm⁻² y⁻¹ are also found in pollen traps from the Baltic, Wales, Turkey and Georgia. The pollen type mainly comes from *Corylus avellana*, while *C. maxima* occurs in Greece, *C. colchica* in Georgia and *C. colurna* in Turkey and in plantations or as ornamental trees in the rest of the Europe. As discussed in the main text, the selection of fossil sites did not include studies from western and

30 lowland central Europe where values of 10000 grains cm⁻² y⁻¹ are common for the Early Holocene. The highest fossil PARs from the chosen examples were estimated for the Early Holocene from Prášílské situated at 1000 m a.s.l. Thus the high modern values in north-central Poland provide analogues for several fossil situations. Occasional grains and small PARs of *Corylus* pollen are common in traps from boreal regions as well as two traps from the arctic region.

Abies



Mean modern PARs averaged for each trap area (a). b) Range of mean individual trap values classified by one-dimensional clustering. Crossed squares indicate that pollen of the taxon was not found in any trap from the area. c) Map of Europe with the distribution of the species (gray, Caudullo et al., 2017, San-Miguel-Ayán et al. 2016) falling within the pollen taxa, size of symbols shows classes of PAR in recent and the highest PAR per each fossil record. Arrows show the closest trap with the same class of PAR. d) Fossil PAR values with the highest PAR class per each record (see Table 1 for full name) highlighted by the corresponding colour for the class (see b) Note the scale of the x-axis corresponds to the x-axis scale of graph a).

Figure 6. Mean modern PAR averaged for each trap area (a). b) Range of mean individual trap values classified by one-dimensional clustering. Crossed squares indicate that pollen of the taxon was not found in any trap from the area. c) Map of Europe with the distribution of the species (gray, Caudullo et al., 2017, San-Miguel-Ayán et al. 2016) falling within the pollen taxa, size of symbols shows classes of PAR in recent and the highest PAR per each fossil record. Arrows show the closest trap with the same class of PAR. d) Fossil PAR values with the highest PAR class per each record (see Table 1 for full name) highlighted by the corresponding colour for the class (see b) Note the scale of the x-axis corresponds to the x-axis scale of graph a).




Figure 7. See caption Fig. 6




Figure 8. See caption Fig. 6




Figure 9. See caption Fig. 6

3.5.1

PARs of more than 3300 grains cm⁻² y⁻¹ are found in pollen traps in Poland and Georgia and values above 2100 occur in traps from the Strandza, Šumava and Jura mountains, in southern Bulgaria, Czechia and Switzerland respectively. Two species of *Fagus* contribute to the pollen type with *Fagus sylvatica* as the dominant tree across much of Europe and *F. orientalis* occurring in southern Bulgaria, Turkey and Georgia. Fossil PAR increase around 8000 cal. BP at the Bulgarian Black Sea coast, around 7000 in the Šumava Mountains and 1500 years ago in northern Poland. High values in Šumava and northern Poland match the values in adjacent traps. Hardly any *Fagus* pollen occurs in pollen traps outside its modern distribution except sporadic appearances in Tula and the two northernmost trap areas. However, *Fagus* pollen occurs regularly at fossil sites that were assumed to have never been within the distribution of the tree, such as the sites in central Sweden, where *Fagus* grains occurred regularly over the last 3000 years.

3.5.1

Modern *Picea* PARs above 2800 grains cm⁻² y⁻¹ are present in traps from Central Sweden, the Baltic region, Russia, Šumava mountains and the Alps. While some of the modern *Picea* pollen may be released by planted non-native *Picea sitchensis* and *P. pungens*, most of the pollen comes from *Picea abies*, which is also planted in many European regions outside its natural distribution. The highest fossil values in the selection of sites come from the Šumava mountains and are comparable to high trap values from the same region although at lower elevations. Noteworthy are also the generally low fossil *Picea* PARs for




Figure 10. See caption Fig. 6

sites in Central Sweden and Tver region, where the tree is dominating or co-dominating the forest for the last 2000 years. Most pollen traps from beyond the distribution area of *Picea* collect individual *Picea* pollen.

3.5.1

Highest modern PARs exceeding 43600 grains $\text{cm}^{-2} \text{y}^{-1}$ are observed in traps on Cyprus, while the values in traps from the northern boreal forest often stay below 5400 grains $\text{cm}^{-2} \text{y}^{-1}$. *Pinus* PAR values increase from Finnmark (2000 grains $\text{cm}^{-2} \text{y}^{-1}$) and central Sweden (5000 grains $\text{cm}^{-2} \text{y}^{-1}$) to the Baltic and north-central Poland with 35000 grains $\text{cm}^{-2} \text{y}^{-1}$. In northern Europe nearly all *Pinus* pollen comes from *P. sylvestris*, while southern European trapping sites have a higher diversity of trees within the subgenus *Diploxylon* including *P. mugo*, *P. nigra*, *P. brutia* and *P. halapensis*. Pollen of both *Haploxylon* pines *P. cembra* in Alps and *P. peuce* in Rila was separated. High fossil *Pinus* PAR values are estimated for Suminko in northern Poland, matching modern trap values from the same region. In the Šumava mountains *Pinus* was possibly the dominant forest tree during the Early Holocene and declined thereafter, so that modern values from north-central Poland provide the nearest analogue to the Early Holocene situation. The lowest *Pinus* PARs (<125 grains $\text{cm}^{-2} \text{y}^{-1}$) are found in traps from Iceland (2-5 grains $\text{cm}^{-2} \text{y}^{-1}$), Lagodheki and most of the traps from Lofoten-Vesterålen (38-125 grains $\text{cm}^{-2} \text{y}^{-1}$) and single traps from Lapps-kardet, Exmoor England and Zermatt.

3.5.1




Figure 11. See caption Fig. 6

Modern-PARs of *Quercus* within the area of distribution of a parent tree in the genus range between 620-15000 grains cm⁻² y⁻¹. Highest modern values (6000-15000 grains cm⁻² y⁻¹) are found in the traps from the UK, Poland and the southern Balkan. The first two areas host only species belonging to the *Q. robur*-type, whereas the two latter also include species from *Q. cerris*-type and *Q. ilex*-type. The highest fossil values from Suminko seem too high for the region compared to other fossil samples, nevertheless modern values in pollen traps from the region provide comparable high PARs. Also, PARs in recent sediments from some lakes in north-eastern Germany show similar values (Matthias and Giesecke, 2014). PARs to about 1300 grains cm⁻² y⁻¹ are estimated for Røuge-Töugjärv near the distribution limit of *Quercus robur* where the tree only became abundant after 7000 years ago. *Quercus robur* reached its maximum abundance even later in central Sweden where the PAR at Holtjärnen around 3200 years ago suggest its presence. The pollen seems to disperse well and is found in small amounts in most pollen traps beyond its distribution area and fossils samples far to the north of the distribution have also collected *Quercus* pollen during the Holocene, which may be partly due to the abundance of trees of this genus in Europe.

3.5.1

The highest modern PAR 1500-4700 grains cm⁻² y⁻¹ was measured in Poland, European Russia (Tula) and Lagodheki in Georgia, however, this high range has no comparison in the fossil record. Lower values 120-1500 grains cm⁻² y⁻¹ were found in traps from European Turkey and the Alps. Highest fossil PAR 270-1500 was measured in middle altitude and lowland temperate zone during the Middle Holocene. The closest trap analogues to them are in North Poland. Sites on the current edge of *Tilia* distribution and in the mountains (Central Sweden, Alps and Rila) show highest PAR range 120-270 grains cm⁻² y⁻¹,




Figure 12. See caption Fig. 6

also during the Middle Holocene. Trap records corresponding to the lowest class and above the LDT ($5-40 \text{ grains cm}^{-2} \text{ y}^{-1}$) appear in sites within distribution limit of *Tilia* (Baltic, European Russia, Poland), on its edge (Central Sweden, Wales, Black Sea Coast, Greece) or in the mountains (Šumava, Alps).

4 Discussion

5 4.1 Trap PARs and environmental conditions

This overview of European pollen trap data collected by the PMP network demonstrates that modern PARs provide comparable values to fossil records and can thus help in interpreting the fossil signal in spite of the different taphonomic processes that influence PAR values in pollen traps versus lake sediments (Lisitsyna et al., 2011b). The dataset extends across the European latitudinal and altitudinal range and documents general patterns. The latitudinal gradient in PAR is clearly visible in this dataset. Although data on plant biomass and primary productivity are not available for all trapping locations, the regression analysis indicates that mean annual temperature has an influence on the quantity of pollen deposition. The July temperature of the previous year determines the amount of pollen production in *Pinus* near the tree-line (Autio and Hicks, 2004; McCarroll et al., 2003). Evidence from other European regions (van der Knaap et al., 2010; Nielsen et al., 2010) suggests that also the growing season warmth and other climate variables also explain the interannual variability of pollen deposition. On the regional scale, PAR corresponds to plant biomass of the parent tree (Matthias and Giesecke, 2014; Seppä et al., 2009).




Figure 13. See caption Fig. 6

However, differences in forest cover cannot explain the latitudinal gradient in PAR described here, which may, at least in part, result from the latitudinal gradient in primary productivity of trees (Gillman et al., 2015). An increase in primary productivity and pollen production has been shown in a carbon dioxide fertilization experiment (Wayne et al., 2002), which supports the interpretation that average PAR of the same species may vary due to environmental parameters determining its productivity.

5 4.2 Long Distance Transport

Modern ~~PARs~~ PAR from traps near the latitudinal limit of *Pinus* and *Betula* have been used previously to reconstruct past changes in the northern distribution limits of these trees (Seppä and Hicks, 2006). Our LDT result 80 grains cm⁻² y⁻¹ is slightly lower than the range for *Pinus* and *Betula* in arctic-apline zone 100-200 grains cm⁻² y⁻¹ (Seppä and Hicks, 2006). Here we evaluated ~~larger-distances-andtherefore~~ greater distances and, therefore, had to ignore ~~*Pinus* and *Betula*, while-suggesting both~~ species, while indicating some general thresholds for other dominant European trees. The ~~upper-value-of-around-30-suggested~~ threshold of 80 grains cm⁻² y⁻¹ may be used to differentiate local from long distance transport for most wind pollinated trees. However, this value may still lead to false negatives as the absence of a plant cannot be proven by the absence of evidence for presence. The value of 80 grains cm⁻² y⁻¹ for *Picea* agrees well with the fossil PAR value for the tree of 50 *Picea* grains cm⁻² y⁻¹ found in a sample at Klotjärnen just after the occurrence of the first *Picea* bud scale (Giesecke, 2005b). However, these modern thresholds estimated here are likely to depend on the abundance of the parent tree in the larger region rather than properties of the pollen types. A ~~larger-higher~~ threshold would be expected for *Corylus* compared to *Fagus*, based-on-the-fall-speed-of pollen. *Corylus* has a lighter pollen grain than *Fagus*, which can travel more easily over large distances (Table S5). However,

~~Corylus is not very abundant near its northern limit in these more continental areas, while LDT for Picea results too low and Fagus often dominates forests near its presumed limits too high. They do not cover the same distance ranges as the Corylus results. LDT estimate of Fagus is based on a few traps, very distant from the distribution area. Similarly, the occurrences of taxa outside the mapped natural distribution can bias this estimate.~~

- 5 ~~Characteristic radius is a useful measure showing pollen transport predicted by pollen dispersal models. It represents theoretical proportion of pollen loading at different distances from the source plants (Prentice, 1988) and thus can be easily compared with our empirical values. PAR of Considering Carpinus, Corylus, Fraxinus and Fagus, Fraxinus, Picea, Quercus and Tilia, the proportion of pollen loading within the area of distribution and at 200 km from the distribution area represent 5-10 % of the median PAR within the distribution area (Fig. 4). Considering the range of standard terminal velocities of those~~
- 10 ~~taxa and source plants measured empirically by our PAR results (0.002-36%, in average 0.2%) is wider and higher than the theoretical range (0.005-13%, in average 0.05%; Table S6). This mismatch can be caused by too leptokurtic character of the Gaussian Plume model with wind speed windspeed 3 m.s⁻¹ (Abraham et al., 2014), we get 2-13% of pollen loading at the characteristic radius 200 km (Table S6), and fits with previous indication that it underestimates dispersal of pollen with a large grain (Abraham et al., 2014, Theuerkauf et al. (2016)).~~
- 15 ~~Pollen rain beyond the area of distribution is analogous with high elevations with sparse vegetation. However, with the increasing altitude, PAR decreases independently from the actual growth density, because of the worse climatic conditions and low pollen productivity (Markgraf, 1980).~~

4.3 Analogues for vegetation reconstruction

- ~~While the dataset of modern PARs presented here provides analogues for the selection of fossil sites, reliable~~
- 20 ~~The comparisons of modern and fossil PAR values show that pollen traps can characterize the population density of particular taxa in modern situations and thus provide analogues aiding in the interpretation of fossil situations. One caveat of the comparisons conducted here is the lack of fossil sites in the oceanic climate of western Europe. Reliable fossil PAR records were not available from the areas of trapping locations in the UK and pollen trapping locations from northern France or Spain are not available. Moreover, no sites with fossil PAR from western Europe or from low elevations in the Alps are included in this comparison. Consequently,~~
- 25 ~~the This omission has consequences for some of the comparisons. A particular case where PAR could in theory aid in the interpretation of the fossil situation are the large quantities of Corylus pollen deposited in many in early Holocene samples of west European sites during the early Holocene was not considered. There are. However, there are no or limited modern analogues for the highest early Holocene values of for the highest Early Holocene values. Among the fossil sites considered here the highest values are~~
- 30 ~~around 7000 Corylus grains cm⁻² y⁻¹ at Prášliské. Average Early Holocene Corylus PAR at Soppensee in northern Switzerland are 12000 grains cm⁻² y⁻¹ (Lotter, 1999) and at Meerfelder Maar (Kubitz, 2000) in western Germany 18000 grains cm⁻² y⁻¹. Judging from pollen percentages, even higher Early Holocene values should be found in more oceanic situations and the Corylus PAR at Hockham Mere in eastern England may be as high as 40000 grains cm⁻² y⁻¹ for the early Holocene (Bennett, 1983). Modern values in pollen traps from Wales at around 2000 grains cm⁻² y⁻¹ are far below these early Holocene figures and it is likely that modern analogues of sites with high Corylus PARs PAR no longer exist in Europe.~~

Conversely, the high modern PAR values for *Pinus* and *Betula* from Poland and Latvia are not found in the fossil examples. *Pinus* PAR values around 30000 grains cm⁻² y⁻¹ were also obtained from ²¹⁰Pb dated modern lake sediment samples in north eastern Brandenburg (Matthias and Giesecke, 2014). This study evaluated the ~~PARs~~-PAR for the years 1993 and 2009. The increase in *Pinus* PAR values between the first and the second sampling period corresponded with an increase in the amount of standing pine volume in the region. Forestry practices aimed at increasing yield could account for the high *Pinus* values. *Pinus* was extensively planted after the 1950s, even on soils where trees with a lower pollen production would have grown naturally. The fertilization due to increased nitrogen deposition ([Pers-Kamczyc et al., 2020](#)), as well as increased atmospheric carbon dioxide, increase the pollen production not only of *Pinus*. A carbon dioxide enrichment experiment of 19-year old *Pinus taeda* resulted in a twofold probability of reproductive maturity after 3 years (LaDeau and Clark, 2001). The continued experiment also showed that carbon dioxide fertilization increased the number of pollen cones and therefore pollen grains produced per tree (LaDeau and Clark, 2006).

In the case of *Pinus*, the modern dataset includes trap data from Cyprus, where *Pinus brutia* dominates at 1600 m a.s.l., resulting in even higher *Pinus* PAR values compared to those found in the Polish and Baltic regions. The highest *Betula* values come mainly from Russia, where values frequently exceed 30000 grains cm⁻² y⁻¹. We previously discussed such high fossil PAR values for *Corylus*, which is assumed to produce a similar amount of pollen. However, fossil *Betula* ~~PARs~~-PAR in the examples considered here are consistently below 6000 grains cm⁻² y⁻¹ and published early Holocene values rarely exceed 6000 grains cm⁻² y⁻¹ (but see Theuerkauf et al., 2014). Pollen diagrams from the forest steppe ecotone in European Russia are often characterized by high *Betula* ~~percentage~~-percentages (Nosova et al., 2019; Shumilovskikh et al., 2018). However, there are no suitable diagrams with reliable PAR estimates from that region. It is thus difficult to judge whether high modern trap values are associated with recent land-use change or are characteristic of eastern European forests.

The comparison of regional ~~PARs~~-PAR between traps and fossil estimates indicates higher fossil PAR of *Picea*, *Fagus* and *Abies* in middle altitudes of the temperate zone (Fig. 5), which, in the case of *Abies*, represents the Europe-wide decline in *A. alba* (Tinner et al., 2013). *Picea* and *Fagus* dominate central European forests today and *Picea* is planted ~~much~~-well beyond its natural range. However, both trees start flowering rather late in their lives and harvesting the trees at a young age may contribute to lower modern PAR values. Fossil and modern ~~PARs~~-PAR for these trees in Šumava are similar, while only the highest trap values match the Holocene high values. On average there are lower modern PAR values; this may be explained by a lowering of the treeline over the last millennia. This interpretation agrees with REVEALS reconstructions for this region, indicating a decline in the cover abundance of *Picea* and *Fagus* (Abraham et al., 2016; Carter et al., 2018). Within a 60 km radius of the fossil sites, *Picea* decreased in abundance from 70% during the Middle Holocene to 43%, compared to modern abundance. *Fagus* and *Abies* declined from Late Holocene values of 22% and 3% to currently 20% to 1% respectively (Abraham et al., 2016). The abundance of *Abies* in the Roztocze region (SE Poland; Fig. 6) provides a good analogue for the past abundance of the tree in Šumava with maximum PAR of 1000-3000 grains cm⁻² y⁻¹. *Abies* disappeared from the Czech Republic during the Mediaeval Age due to forest management methods (Kozáková et al., 2011), which were not practiced in ~~SE~~-south-east Poland.

Linking the fossil to modern PAR values facilitates the interpretation of the fossil record of individual sites. Unfortunately, the details cannot be discussed here. However, the central Swedish sites Holtjärnen and Klotjärnen provide excellent examples.

These sites are situated north of the modern distribution of *Tilia*, *Corylus*, *Quercus* and near the limit of *Alnus glutinosa*. The fossil PAR values are higher for these taxa than those found in pollen traps at or near these lakes (Giesecke, 2005a; Giesecke and Fontana, 2008). Modern reference values for the ~~PARs~~ PAR of these taxa can be found in northern Poland and Estonia. Moreover, this analogue matching indicates that 3000 years ago the PAR values for *Quercus* at Holtjärnen were sufficiently high to indicate the occurrence of small populations near the lake. Those taxa at both sites also nicely illustrate the potential use of LDT for the interpretation of the fossil record. Fossil PAR values of *Tilia*, *Corylus* and *Quercus* in ~~boreal-region-result~~ Boreal region are above LDT threshold, whereas all trap PAR values ~~are-result~~ below (Fig. 5).

4.4 Limitations and problems

~~There~~ Nevertheless, there are significant differences between the accumulation of pollen in traps and on peatlands and lakes (Lisitsyna et al., 2011b; Pardoe et al., 2010). Differences in pollen trap design and placement in the landscape will influence the values. Trap values are also affected by modern processes that have no impact on the fossil signal, such as pollen from trapped insects. These biases appear minor as indicated by the large consistency of the data collected in the PMP database. Also the comparison of values over this large environmental gradient ~~result-results~~ in the signal being ~~stringer~~ stronger than the noise. Nevertheless, some traps or individual years have unusual values and were removed from the comparison (Table S1). Despite this, the uncertainty of fossil PAR values is much greater than pollen traps, which is primarily due to the added uncertainty coming from sampling a sediment core, combined with the uncertainty of the age model (Maher, 1981). PAR from lake ~~sediments-sediment~~ has additional biases due to differential sedimentation of pollen grains in lakes (Davis and Brubaker, 1973), sediment re-deposition, focussing and catchment erosion (Davis et al., 1984; Pennington, 1979). Although we carefully selected the best available fossil sites, ~~PARs~~ PAR especially from lake Suminko and Rõuge Tõugjärv may be biased by lake internal processes and the addition of stream borne pollen respectively. Nevertheless, their fossil PAR estimates are in the range of values found in pollen traps. Where detailed knowledge ~~on-of~~ the sedimentation process is available, the bias of sediment focussing may be reduced, as in the example of Hockham Mere cited above (see also Bennett and Buck, 2016; Bennett, 1983). Peatlands may thus seem the better choice for obtaining fossil PAR, which may be the case in northern Scandinavia (Barnekow et al., 2007; Finsinger et al., 2013), but frequent changes in the rate of peat growth lead to difficulties assessing the time represented in individual samples at many sites.

The problem of traps collecting ~~high-large~~ amounts of herbaceous pollen brought by insects and small animals was discussed in the method section and for this reason ~~only~~ Poaceae and Cyperaceae are the only herbs selected for our analyses. However, pollen from these two families is also often overrepresented in the pollen traps (Lisitsyna et al., 2011b), as the plants may overhang the trap opening and their pollen may fall directly into the trap. Reduced ~~PARs~~ PAR in the trap may be caused by overgrowth of the vegetation or leaves temporally blocking the opening, while proximity to the forest edge would increase values compared to large open peatlands or lakes. These effects have not been systematically evaluated so far.

Detailed comparisons of vegetation data to ~~PARs~~ PAR hold potential for a better understanding of the spatial representation and processes shaping the pollen signal (Matthias and Giesecke, 2014) and allow estimates of absolute pollen productivity (Sugita et al., 2009) or test pollen dispersal models. However, for this continental scale dataset, available vegetation data have

limited precision. Forest inventory data with the detail essential for this type of study ~~is~~are not available for all traps. The forest cover data presented here has a resolution of 1 km², which is insufficient as the abundance of trees within hundreds of meters of the traps is important. Moreover, without information on standing volume or age structure, the percentage cover used here is a crude measure of the vegetation producing the pollen. Forestry practices like harvesting trees that start flowering at a later age (e.g. *Picea* 30-40 years) reduce the number of trees producing pollen (Matthias et al., 2012) and bias the search for modern analogues. Also, the available mapped distribution limits of trees have large uncertainties precluding more detailed assessments of the quantity of long distance transported pollen using this continental dataset.

5 Conclusions

Comparison of the mean annual PAR from traps and fossil sites showed similar ranges for *Abies*, *Alnus*, *Betula*, *Carpinus*, *Corylus*, *Fagus*, *Fraxinus*, *Picea*, *Pinus*, *Quercus* and *Tilia* at the continental scale. This indicates that there are no major biases hampering the application of the PMP Database data as a modern reference to interpret the fossil record. The dataset clearly shows that climate parameters ~~that~~ correlate with latitude ~~determine~~in determining pollen productivity. The effect of regional forest cover is discernible. Minimum values suggest that an 80% forest cover within 10 km of the trap results in ~~PARs~~PAR above 3200 tree pollen grains cm⁻² year⁻¹.

Assessment of long-distance transport indicates that values below ~~30~~80 grains cm⁻² y⁻¹ for *Carpinus*, *Corylus*, *Fagus*, *Fraxinus*, *Picea*, *Quercus* and *Tilia* may originate from beyond 200 km of a sampling site. This number of ~~30~~80 grains cm⁻² y⁻¹ may therefore be used as a general threshold indicating long distance origin of pollen. The application of these threshold values holds potential to refine and adjust reconstructions of tree distributions.

Code availability. Primary trap and fossil pollen data are available in Neotoma Palaeoecology Database <https://www.neotomadb.org/>. Analysis are based on the WorldClim 2 dataset of Fick and Hijmans (2017), Chorological data for the main European woody species, version 2 by Caudullo et al. (2018), European atlas of forest tree species, 2016th ed. by San-Miguel-Ayán et al. (2016), which are available online: <http://worldclim.org/version2> , <https://data.mendeley.com/datasets/hr5h2hcg4/2> and <http://www.euforgen.org/>, respectively. Forest Map of Europe of Kempeneers et al. (2012) is available on request on authors.

Code for analysis, derived data and code for figures are available in the <https://github.com/vojtechabraham/PMPdatabase>.

Author contributions. Design of the analysis: TG and VA. Data analysis: VA, TG and MW. Manuscript draft: VA and TG. Manuscript comments: HP, SH, CEJ, HSS, HS, SV, SH, MN, MW, SP, ST, TK, AMN and JSM. Data collection: all authors except VA and MW.

Competing interests. The authors declare no competing interests.

Acknowledgements. We thank ~~to all~~ all the collaborators of pollen trapping, especially ~~to~~ those who contributed to the PMP dataset: Pim van der Knaap, Jacqueline van Leeuwen, Karl-Dag Vorren, Heather Tinsley, Lena Barnekow, Małgorzata Latałowa, Enikő Magyari, Larissa Savelieva, Elena Pavlova, Heidi Hyypä, Mari Kuoppamaa and Gunhild Rosqvist. We thank Simon Connor for his detailed review which helped to improve the manuscript, as well as two anonymous reviewers for their comments.

References

- Abraham, V., Oušková, V., and Kuneš, P.: Present-Day Vegetation Helps Quantifying Past Land Cover in Selected Regions of the Czech Republic, *PLoS ONE*, 9, e100117, <https://doi.org/10.1371/journal.pone.0100117>, <http://dx.plos.org/10.1371/journal.pone.0100117>, 2014.
- Abraham, V., Kuneš, P., Petr, L., Svitavská-Svobodová, H., Kozáková, R., Jamrichová, E., Švarcová, M. G., and Pokorný, P.: A pollen-based
5 quantitative reconstruction of the Holocene vegetation updates a perspective on the natural vegetation in the Czech Republic and Slovakia, *Preslia*, 88, 409–434, 2016.
- Andersen, S.: The relative pollen productivity and representation of north European trees, and correction factors for tree pollen spectra., *Danmarks Geologiske Undersøgelse Række II*, 96, 1–99, 1970.
- Andersen, S. T.: Influence of Climatic Variation on Pollen Season Severity in Wind-Pollinated Trees and Herbs, *Grana*, 19, 47–52,
10 <https://doi.org/10.1080/00173138009424986>, 1980.
- Autio, J. and Hicks, S.: Annual variations in pollen deposition and meteorological conditions on the fell Aakenustunturi in northern Finland: Potential for using fossil pollen as a climate proxy, *Grana*, 43, 31–47, <https://doi.org/10.1080/00173130310017409>, 2004.
- Barnekow, L., Loader, N. J., Hicks, S., Froyd, C. A., and Goslar, T.: Strong correlation between summer temperature and pollen accumulation rates for *Pinus sylvestris*, *Picea abies* and *Betula* spp. in a high-resolution record from northern Sweden, *J. Quaternary Sci.*, 22, 653–658,
15 <https://doi.org/10.1002/jqs.1096>, 2007.
- Bennett, K. and Buck, C. E.: Interpretation of lake sediment accumulation rates, *Holocene*, 26, 1092–1102, <https://doi.org/10.1177/0959683616632880>, 2016.
- Bennett, K. D.: Devensian Late-Glacial and Flandrian Vegetational History at Hockham Mere, Norfolk, England, *New Phytol.*, 95, 489–504, <https://doi.org/10.1111/j.1469-8137.1983.tb03513.x>, 1983.
- 20 Bozilova, E. and Beug, H.-J.: On the Holocene history of vegetation in SE Bulgaria (Lake Arkutino, Ropotamo region), *Veg. Hist. Archaeobot.*, 1, 19–32, <https://doi.org/10.1007/BF00190698>, 1992.
- Carter, V. A., Chiverrell, R. C., Clear, J. L., Kuosmanen, N., Moravcová, A., Svoboda, M., Svobodová-Svitavská, H., Leeuwen, V., N, J. F., Knaap, V. D., O, W., and Kuneš, P.: Quantitative Palynology Informing Conservation Ecology in the Bohemian/Bavarian Forests of Central Europe, *Front. Plant Sci.*, 8, <https://doi.org/10.3389/fpls.2017.02268>, 2018.
- 25 Caudullo, G., Welk, E., and San-Miguel-Ayanz, J.: Chorological maps for the main European woody species, *Data in Brief*, 12, 662–666, <https://doi.org/10.1016/j.dib.2017.05.007>, <http://www.sciencedirect.com/science/article/pii/S2352340917301981>, 2017.
- Davis, B. A. S., Zanon, M., Collins, P., Mauri, A., Bakker, J., Barboni, D., Barthelmes, A., Beaudouin, C., Bjune, A. E., Bozilova, E., Bradshaw, R. H. W., Brayshay, B. A., Brewer, S., Brugiapaglia, E., Bunting, J., Connor, S. E., de Beaulieu, J.-L., Edwards, K., Ejarque, A., Fall, P., Florenzano, A., Fyfe, R., Galop, D., Giardini, M., Giesecke, T., Grant, M. J., Guiot, J., Jahns, S., Jankovská, V., Juggins, S.,
30 Kahrmann, M., Karpińska-Kołaczek, M., Kołaczek, P., Kühl, N., Kuneš, P., Lapteva, E. G., Leroy, S. A. G., Leydet, M., Guiot, J., Jahns, S., Jankovská, V., Juggins, S., Kahrmann, M., Karpińska-Kołaczek, M., Kołaczek, P., Kühl, N., Kuneš, P., Lapteva, E. G., Leroy, S. A. G., Leydet, M., López Sáez, J. A., Masi, A., Matthias, I., Mazier, F., Meltsov, V., Mercuri, A. M., Miras, Y., Mitchell, F. J. G., Morris, J. L., Naughton, F., Nielsen, A. B., Novenko, E., Odgaard, B., Ortu, E., Overballe-Petersen, M. V., Pardoe, H. S., Peglar, S. M., Pidek, I. A., Sadori, L., Seppä, H., Severova, E., Shaw, H., Święta Musznicka, J., Theuerkauf, M., Tonkov, S., Veski, S., van der Knaap, W. O., van
35 Leeuwen, J. F. N., Woodbridge, J., Zimny, M., and Kaplan, J. O.: The European Modern Pollen Database (EMPD) project, *Veg. Hist. Archaeobot.*, 22, 521–530, <https://doi.org/10.1007/s00334-012-0388-5>, 2013.

- Davis, M. B.: Palynology after Y2K—Understanding the Source Area of Pollen in Sediments., *Annual Review of Earth & Planetary Sciences*, 28, 1, <https://doi.org/Article>, 2000.
- Davis, M. B. and Brubaker, L. B.: Differential Sedimentation of Pollen Grains in Lakes, *Limnol. Oceanogr.*, 18, 635–646, <https://doi.org/10.4319/lo.1973.18.4.0635>, 1973.
- 5 Davis, M. B. and Deevey, E. S.: Pollen Accumulation Rates: Estimates from Late-Glacial Sediment of Rogers Lake, *Science*, 145, 1293–1295, <https://doi.org/10.1126/science.145.3638.1293>, 1964.
- Davis, M. B., Moeller, R., and Ford, J.: Sediment focusing and pollen influx, in: *Lake sediments and environmental history: studies in palaeolimnology and palaeoecology in honour of Winifred Tutin*, edited by Haworth, E. Y., Lund, J. W. G., and Tutin, W., pp. 261–293, Leicester University Press, 1984.
- 10 Fick, S. E. and Hijmans, R. J.: WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas, *Int. J. Climatol.*, 37, 4302–4315, <https://doi.org/10.1002/joc.5086>, 2017.
- Filipova-Marinova, M.: Palaeoecological investigations of lake Shabla-Ezeretz in NE Bulgaria, *Ecologia Mediterranea*, 11, 147–158, 1985.
- Finsinger, W., Schoning, K., Hicks, S., Lücke, A., Goslar, T., Wagner-Cremer, F., and Hyypä, H.: Climate change during the past 1000 years: a high-temporal-resolution multiproxy record from a mire in northern Finland, *J. Quaternary Sci.*, 28, 152–164, <https://doi.org/10.1002/jqs.2598>, 2013.
- 15 Gaillard, M.-J., Sugita, S., Bunting, M. J., Middleton, R., Broström, A., Caseldine, C., Giesecke, T., Hellman, S. E. V., Hicks, S., Hjelle, K., Langdon, C., Nielsen, A.-B., Poska, A., Stedingk, H., Veski, S., and members, P.: The use of modelling and simulation approach in reconstructing past landscapes from fossil pollen data: a review and results from the POLLANDCAL network, *Veget Hist Archaeobot*, 17, 419–443, <https://doi.org/10.1007/s00334-008-0169-3>, <http://www.springerlink.com/content/1tpg426573513vqg/>, 2008.
- 20 Gerasimidis, A., Panajiotidis, S., Hicks, S., and Athanasiadis, N.: An eight-year record of pollen deposition in the Pieria mountains (N. Greece) and its significance for interpreting fossil pollen assemblages, *Rev. Palaeobot. Palyno.*, 141, 231–243, <https://doi.org/10.1016/j.revpalbo.2006.04.004>, 2006.
- Giesecke, T.: Holocene dynamics of the southern boreal forest in Sweden, *Holocene*, 15, 858–872, <https://doi.org/10.1191/0959683605hl859ra>, 2005a.
- 25 Giesecke, T.: Moving front or population expansion: How did *Picea abies* (L.) Karst. become frequent in central Sweden?, *Quaternary Sci. Rev.*, 24, 2495–2509, <https://doi.org/10.1016/j.quascirev.2005.03.002>, 2005b.
- Giesecke, T.: Holocene forest development in the central Scandes Mountains, Sweden, *Veg. Hist. Archaeobot.*, 14, 133–147, <https://doi.org/10.1007/s00334-005-0070-2>, 2005c.
- Giesecke, T. and Fontana, S. L.: Revisiting pollen accumulation rates from Swedish lake sediments, *Holocene*, 18, 293–305, <https://doi.org/10.1177/0959683607086767>, 2008.
- 30 Giesecke, T., Fontana, S. L., Knaap, W. O., Pardoe, H. S., and Pidek, I. A.: From early pollen trapping experiments to the Pollen Monitoring Programme, *Veg. Hist. Archaeobot.*, 19, 247–258, <https://doi.org/10.1007/s00334-010-0261-3>, 2010.
- Giesecke, T., Brewer, S., Finsinger, W., Leydet, M., and Bradshaw, R. H. W.: Patterns and dynamics of European vegetation change over the last 15,000 years, *Journal of Biogeography*, 44, 1441–1456, <https://doi.org/10.1111/jbi.12974>, <https://onlinelibrary.wiley.com/doi/pdf/10.1111/jbi.12974>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/jbi.12974](https://onlinelibrary.wiley.com/doi/pdf/10.1111/jbi.12974), 2017.
- 35 Giesecke, T., Wolters, S., Leeuwen, J. F. N. v., Knaap, P. W. O. v. d., Leydet, M., and Brewer, S.: Postglacial change of the floristic diversity gradient in Europe, *Nat. Commun.*, 10, 1–7, <https://doi.org/10.1038/s41467-019-13233-y>, 2019.

- Gillman, L. N., Wright, S. D., Cusens, J., McBride, P. D., Malhi, Y., and Whittaker, R. J.: Latitude, productivity and species richness, *Global Ecol. and Biogeogr.*, 24, 107–117, <https://doi.org/10.1111/geb.12245>, 2015.
- GRASS Development Team: Geographic Resources Analysis Support System (GRASS GIS) Software, Open Source Geospatial Foundation, 2018.
- 5 Haselhorst, D. S., Moreno, J. E., and Punyasena, S. W.: Assessing the influence of vegetation structure and phenological variability on pollen-vegetation relationships using a 15-year Neotropical pollen rain record, *Journal of Vegetation Science*, 31, 606–615, <https://doi.org/https://doi.org/10.1111/jvs.12897>, <https://onlinelibrary.wiley.com/doi/abs/10.1111/jvs.12897>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/jvs.12897](https://onlinelibrary.wiley.com/doi/pdf/10.1111/jvs.12897), 2020.
- Hesselman, H.: Om pollenregn på hafvet och fjärrtransport af barrträdspollen., *Geologiska Föreningens i Stockholm Förhandlingar*, 41, 89–99, 1919.
- 10 Hicks, S.: A method of using modern pollen rain values to provide a timescale for pollen diagrams from peat deposits, *Memorana Societas Fauna Flora Fennica*, 49, 21–33, 1974.
- Hicks, S.: Present and past pollen records of Lapland forests, *Rev. Palaeobot. Palyno.*, 82, 17–35, [https://doi.org/10.1016/0034-6667\(94\)90017-5](https://doi.org/10.1016/0034-6667(94)90017-5), 1994.
- 15 Hicks, S.: The relationship between climate and annual pollen deposition at northern tree-lines, *Chemosphere - Global Change Science*, 1, 403–416, [https://doi.org/10.1016/S1465-9972\(99\)00043-4](https://doi.org/10.1016/S1465-9972(99)00043-4), <http://www.sciencedirect.com/science/article/pii/S1465997299000434>, 00034, 1999.
- Hicks, S. and Hyvärinen, H.: Pollen influx values measured in different sedimentary environments and their palaeoecological implications, *Grana*, 38, 228–242, <https://doi.org/10.1080/001731300750044618>, 1999.
- 20 Hicks, S., Ammann, B., Latałowa, M., Pardoe, H. S., and Tinsley, H.: European Pollen Monitoring Programme: Project Description and Guidelines, Oulu University Press, 1996.
- Hicks, S., Tinsley, H., Pardoe, H. S., and Cundill, P. R.: European Pollen Monitoring Programme: Supplement to the Guidelines, Oulu University Press, 1999.
- Hicks, S., Tinsley, H., Huusko, A., Jensen, C., Hättetstrand, M., Gerasimides, A., and Kvavadze, E.: Some comments on spatial variation in arboreal pollen deposition: first records from the Pollen Monitoring Programme (PMP), *Rev. Palaeobot. Palyno.*, 117, 183–194, [https://doi.org/10.1016/S0034-6667\(01\)00086-0](https://doi.org/10.1016/S0034-6667(01)00086-0), 2001.
- 25 Hättetstrand, M., Jensen, C., Hallsdóttir, M., and Vorren, K.-D.: Modern pollen accumulation rates at the north-western fringe of the European boreal forest, *Rev. Palaeobot. Palyno.*, 151, 90–109, <https://doi.org/10.1016/j.revpalbo.2008.03.001>, 2008.
- Hyvärinen, H.: Absolute and relative pollen diagrams from northernmost Fennoscandia, *Fennia - International Journal of Geography*, 142, 1975.
- 30 Jackson, S. T. and Williams, J. W.: Modern analogs in Quaternary paleoecology: Here Today, Gone Yesterday, Gone Tomorrow?, *Annu. Rev. Earth Pl. Sc.*, 32, 495–537, <https://doi.org/10.1146/annurev.earth.32.101802.120435>, 2004.
- Jahns, S.: The Holocene history of vegetation and settlement at the coastal site of Lake Voukaria in Acarnania, western Greece, *Veg. Hist. Archaeobot.*, 14, 55–66, <https://doi.org/10.1007/s00334-004-0053-8>, 2004.
- 35 Jantz, N., Homeier, J., León-Yáñez, S., Moscoso, A., and Behling, H.: Trapping pollen in the tropics — Comparing modern pollen rain spectra of different pollen traps and surface samples across Andean vegetation zones, *Rev. Palaeobot. Palyno.*, 193, 57–69, <https://doi.org/10.1016/j.revpalbo.2013.01.011>, 2013.

- Jensen, C., Vorren, K.-D., and Mørkved, B.: Annual pollen accumulation rate (PAR) at the boreal and alpine forest-line of north-western Norway, with special emphasis on *Pinus sylvestris* and *Betula pubescens*, *Rev. Palaeobot. Palyno.*, 144, 337–361, <https://doi.org/10.1016/j.revpalbo.2006.08.006>, 2007.
- Kempeneers, P., Sedano, F., Seebach, L. M., Strobl, P., and San-Miguel-Ayaz, J.: Data Fusion of Different Spatial Resolution Remote Sensing Images Applied to Forest-Type Mapping, *IEEE T. Geosci. Remote*, 49, 4977 – 4986, <https://doi.org/10.1109/TGRS.2011.2158548>, 2012.
- Koff, T.: Pollen influx into Tauber traps in Estonia in 1997–1998, *Rev. Palaeobot. Palyno.*, 117, 53–62, [https://doi.org/10.1016/S0034-6667\(01\)00076-8](https://doi.org/10.1016/S0034-6667(01)00076-8), 2001.
- Kozáková, R., Šamonil, P., Kuneš, P., Novák, J., Kočár, P., and Kočárová, R.: Contrasting local and regional Holocene histories of *Abies alba* in the Czech Republic in relation to human impact: Evidence from forestry, pollen and anthracological data, *Holocene*, 21, 431–444, <https://doi.org/10.1177/0959683610385721>, 2011.
- Kubitz, B.: Die holozäne Vegetations- und Siedlungsgeschichte in der Westeifel am Beispiel eines hochauflösenden Pollendiagrammes aus dem Meerfelder Maar [History of Holocene vegetation and settlement of the Western Eifel region (Germany) using a high resolution pollenprofile from the Meerfeld Maar], no. 339 in *Dissertationes botanicae*, Borntraeger, 2000.
- Kvavadze, E.: Annual modern pollen deposition in the foothills of the Lagodekhi Reservation (Caucasus, East Georgia), related to vegetation and climate, *Acta Palaeobotanica*, 41, 355–364, 2001.
- LaDeau, S. L. and Clark, J. S.: Rising CO₂ Levels and the Fecundity of Forest Trees, *Science*, 292, 95–98, <https://doi.org/10.1126/science.1057547>, 2001.
- LaDeau, S. L. and Clark, J. S.: Pollen production by *Pinus taeda* growing in elevated atmospheric CO₂, *Funct. Ecol.*, 20, 541–547, <https://doi.org/10.1111/j.1365-2435.2006.01133.x>, 2006.
- Lisitsyna, O. V., Giesecke, T., and Hicks, S.: Exploring pollen percentage threshold values as an indication for the regional presence of major European trees, *Rev. Palaeobot. Palyno.*, 166, 311–324, <https://doi.org/10.1016/j.revpalbo.2011.06.004>, 2011a.
- Lisitsyna, O. V., Hicks, S., and Huusko, A.: Do moss samples, pollen traps and modern lake sediments all collect pollen in the same way? A comparison from the forest limit area of northernmost Europe, *Veg. Hist. Archaeobot.*, 21, 187–199, <https://doi.org/10.1007/s00334-011-0335-x>, 2011b.
- Lotter, A. F.: Late-glacial and Holocene vegetation history and dynamics as shown by pollen and plant macrofossil analyses in annually laminated sediments from Soppensee, central Switzerland, *Veg. Hist. Archaeobot.*, 8, 165–184, <https://doi.org/10.1007/BF02342718>, 1999.
- Maher, L. J.: Statistics for microfossil concentration measurements employing samples spiked with marker grains, *Rev. Palaeobot. Palyno.*, 32, 153–191, [https://doi.org/10.1016/0034-6667\(81\)90002-6](https://doi.org/10.1016/0034-6667(81)90002-6), 1981.
- Malmström, C.: Degerö Stormyr—en botanisk, hydrologisk och utvecklingshistorisk undersökning över ett nordsvenskt myrkomplex., *Meddelanden från Statens Skogsförsöksanstalt*, 20, 1–2, 1923.
- Markgraf, V.: Pollen Dispersal in a Mountain Area, *Grana*, 19, 127–146, <https://doi.org/10.1080/00173138009424995>, <http://www.tandfonline.com/doi/abs/10.1080/00173138009424995>, 1980.
- Matthias, I. and Giesecke, T.: Insights into pollen source area, transport and deposition from modern pollen accumulation rates in lake sediments, *Quaternary Sci. Rev.*, 87, 12–23, <https://doi.org/10.1016/j.quascirev.2013.12.015>, 2014.
- Matthias, I., Nielsen, A. B., and Giesecke, T.: Evaluating the effect of flowering age and forest structure on pollen productivity estimates, *Veg. Hist. Archaeobot.*, 21, 471–484, <https://doi.org/10.1007/s00334-012-0373-z>, 2012.

- McCarroll, D., Jalkanen, R., Hicks, S., Tuovinen, M., Gagen, M., Pawellek, F., Eckstein, D., Schmitt, U., Autio, J., and Heikkinen, O.: Multiproxy dendroclimatology: a pilot study in northern Finland, Holocene, 13, 829–838, <https://doi.org/10.1191/0959683603hl668rp>, 2003.
- Nielsen, A. B., Møller, P. F., Giesecke, T., Stavngaard, B., Fontana, S. L., and Bradshaw, R. H. W.: The effect of climate conditions on inter-annual flowering variability monitored by pollen traps below the canopy in Draved Forest, Denmark, Veg. Hist. Archaeobot., 19, 309–323, <https://doi.org/10.1007/s00334-010-0253-3>, 2010.
- Nosova, M. B., Novenko, E. Y., Severova, E. E., and Volkova, O. A.: Vegetation and climate changes within and around the Polistovo-Lovatskaya mire system (Pskov Oblast, north-western Russia) during the past 10,500 years, Veg. Hist. Archaeobot., 28, 123–140, <https://doi.org/10.1007/s00334-018-0693-8>, 2019.
- 10 Overpeck, J. T., Webb, T., and Prentice, I. C.: Quantitative interpretation of fossil pollen spectra: Dissimilarity coefficients and the method of modern analogs, Quaternary Res., 23, 87–108, [https://doi.org/10.1016/0033-5894\(85\)90074-2](https://doi.org/10.1016/0033-5894(85)90074-2), 1985.
- Pardoe, H. S., Giesecke, T., Knaap, W. O., Svitavská-Svobodová, H., Kvavadze, E. V., Panajiotidis, S., Gerasimidis, A., Pidek, I. A., Zimny, M., Święta Musznicka, J., Latałowa, M., Noryśkiewicz, A. M., Bozilova, E., Tonkov, S., Filipova-Marinova, M. V., Leeuwen, J. F. N., and Kalniņa, L.: Comparing pollen spectra from modified Tauber traps and moss samples: examples from a selection of woodlands across Europe, Veg. Hist. Archaeobot., 19, 271–283, <https://doi.org/10.1007/s00334-010-0258-y>, 2010.
- 15 Pędziszewska, A., Tylmann, W., Witak, M., Piotrowska, N., Maciejewska, E., and Latałowa, M.: Holocene environmental changes reflected by pollen, diatoms, and geochemistry of annually laminated sediments of Lake Suminko in the Kashubian Lake District (N Poland), Rev. Palaeobot. Palyno., 216, 55–75, <https://doi.org/10.1016/j.revpalbo.2015.01.008>, 2015.
- Pennington, W.: The Origin of Pollen in Lake Sediments: An Enclosed Lake Compared with One Receiving Inflow Streams, New Phytol., 83, 189–213, <https://doi.org/10.1111/j.1469-8137.1979.tb00741.x>, 1979.
- 20 Pers-Kamczyc, E., Tyrała-Wierucka, a., Rabska, M., Wrońska-Pilarek, D., and Kamczyc, J.: The higher availability of nutrients increases the production but decreases the quality of pollen grains in *Juniperus communis* L., Journal of Plant Physiology, 248, 153–156, <https://doi.org/10.1016/j.jplph.2020.153156>, <https://www.sciencedirect.com/science/article/pii/S0176161720300456>, 2020.
- Pidek, I. A.: Nine-year record of *Alnus* pollen deposition in the Roztocze region (SE Poland) with relation to vegetation data, Acta Agrobot., 60, 57–64, 2007.
- 25 Pidek, I. A., Svitavská-Svobodová, H., van der Knaap, W. O., Noryśkiewicz, A. M., Filbrandt-Czaja, A., Noryśkiewicz, B., Latałowa, M., Zimny, M., Święta Musznicka, J., Bozilova, E., Tonkov, S., Filipova-Marinova, M., Poska, A., Giesecke, T., and Gikov, A.: Variation in annual pollen accumulation rates of *Fagus* along a N–S transect in Europe based on pollen traps, Veg. Hist. Archaeobot., 19, 259–270, <https://doi.org/10.1007/s00334-010-0248-0>, 2010.
- 30 Prentice, I. C.: Pollen representation, source area, and basin size: Toward a unified theory of pollen analysis, Quaternary Research, 23, 76–86, [https://doi.org/10.1016/0033-5894\(85\)90073-0](https://doi.org/10.1016/0033-5894(85)90073-0), <http://www.sciencedirect.com/ez.statsbiblioteket.dk:2048/science/article/B6WPN-4DV0VT0-73/2/27923d5b1bce5e853b0eb4adde0c0f53>, 1985.
- Prentice, I. C.: Records of vegetation in time and space: the principles of pollen analysis., in: Vegetation history, Handbook of vegetation science., edited by Huntley, B. J. and Webb, T., pp. 17–42, Kluwer Academic Publishers, 00182, 1988.
- 35 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, 2019.
- San-Miguel-Ayanz, J., Rigo, D. d., Caudullo, G., Durrant, T. H., and Mauri, A.: European atlas of forest tree species, Publication Office of the European Union, 2016 edn., 2016.

- Seppä, H. and Hicks, S.: Integration of modern and past pollen accumulation rate (PAR) records across the arctic tree-line: a method for more precise vegetation reconstructions, *Quaternary Sci. Rev.*, 25, 1501–1516, <https://doi.org/10.1016/j.quascirev.2005.12.002>, 2006.
- Seppä, H. and Weckström, J.: Holocene vegetational and limnological changes in the Fennoscandian tree-line area as documented by pollen and diatom records from Lake Tsuolbmajavri, Finland, *Écoscience*, 6, 621–635, <https://doi.org/10.1080/11956860.1999.11682562>, 1999.
- 5 Seppä, H., Nyman, M., Korhola, A., and Weckström, J.: Changes of treelines and alpine vegetation in relation to post-glacial climate dynamics in northern Fennoscandia based on pollen and chironomid records, *J. Quaternary Sci.*, 17, 287–301, <https://doi.org/10.1002/jqs.678>, 2002.
- Seppä, H., Alenius, T., Muukkonen, P., Giesecke, T., Miller, P. A., and Ojala, A. E.: Calibrated pollen accumulation rates as a basis for quantitative tree biomass reconstructions, *Holocene*, 19, 209–220, <https://doi.org/10.1177/0959683608100565>, 2009.
- Shumilovskikh, L. S., Novenko, E., and Giesecke, T.: Long-term dynamics of the East European forest-steppe ecotone, *J. Veg. Sci.*, 29, 416–426, <https://doi.org/10.1111/jvs.12585>, 2018.
- 10 Song, J. and Wang, H.: Ckmeans.1d.dp: Optimal and Fast Univariate Clustering, *Holocene*, 3, 29–33, 2011.
- Stockmarr, J.: Tablets with spores used in absolute pollen analysis, *Pollen et Spores*, 13, 615–621, 1971.
- Sugita, S., Hicks, S., and Sormunen, H.: Absolute pollen productivity and pollen-vegetation relationships in northern Finland, *J. Quaternary Sci.*, 25, 724–736, <https://doi.org/10.1002/jqs.1349>, 2009.
- 15 Svobodová, H., Soukupová, L., and Reille, M.: Diversified development of mountain mires, Bohemian Forest, Central Europe, in the last 13,000 years, *Quatern. Int.*, 91, 123–135, [https://doi.org/10.1016/S1040-6182\(01\)00106-9](https://doi.org/10.1016/S1040-6182(01)00106-9), 2002.
- Tauber, H.: Investigations of the mode of pollen transfer in forested areas, *Review of Palaeobotany and Palynology*, 3, 277–286, [https://doi.org/10.1016/0034-6667\(67\)90060-7](https://doi.org/10.1016/0034-6667(67)90060-7), <http://www.sciencedirect.com/science/article/pii/0034666767900607>, 00198, 1967.
- Tauber, H.: A Static Non-overload Pollen Collector, *New Phytol.*, 73, 359–369, <https://doi.org/10.1111/j.1469-8137.1974.tb04770.x>, 1974.
- 20 Theuerkauf, M., Bos, J. A. A., Jahns, S., Janke, W., Kuparinen, A., Stebich, M., and Joosten, H.: *Corylus* expansion and persistent openness in the early Holocene vegetation of northern central Europe, *Quaternary Sci. Rev.*, 90, 183–198, <https://doi.org/10.1016/j.quascirev.2014.03.002>, 2014.
- Theuerkauf, M., Couwenberg, J., Kuparinen, A., and Liebscher, V.: A matter of dispersal: REVEALSinR introduces state-of-the-art dispersal models to quantitative vegetation reconstruction, *Vegetation History and Archaeobotany*, <https://doi.org/10.1007/s00334-016-0572-0>, <http://link.springer.com/10.1007/s00334-016-0572-0>, 2016.
- 25 Tinner, W., Colombaroli, D., Heiri, O., Henne, P. D., Steinacher, M., Untenecker, J., Vescovi, E., Allen, J. R. M., Carraro, G., Conedera, M., Joos, F., Lotter, A. F., Luterbacher, J., Samartin, S., and Valsecchi, V.: The past ecology of *Abies alba* provides new perspectives on future responses of silver fir forests to global warming, *Ecol. Monogr.*, 83, 419–439, <https://doi.org/10.1890/12-2231.1>, 2013.
- Tinsley, H.: Modern pollen deposition in traps on a transect across an anthropogenic tree-line on Exmoor, southwest England: a note summarising the first three years of data, *Rev. Palaeobot. Palyno.*, 117, 153–158, [https://doi.org/10.1016/S0034-6667\(01\)00083-5](https://doi.org/10.1016/S0034-6667(01)00083-5), 2001.
- 30 Tinsley, H. and Hicks, S.: Preface, *Rev. Palaeobot. Palyno.*, 117, vii–x, [https://doi.org/10.1016/S0034-6667\(01\)00073-2](https://doi.org/10.1016/S0034-6667(01)00073-2), 2001.
- Tonkov, S., Hicks, S., Bozilova, E., and Atanassova, J.: Pollen monitoring in the central Rila Mountains, Southwestern Bulgaria: comparisons between pollen traps and surface samples for the period 1993–1999, *Rev. Palaeobot. Palyno.*, 117, 167–182, [https://doi.org/10.1016/S0034-6667\(01\)00085-9](https://doi.org/10.1016/S0034-6667(01)00085-9), 2001.
- 35 Tonkov, S., Panovska, H., Possnert, G., and Bozilova, E.: The Holocene vegetation history of Northern Pirin Mountain, southwestern Bulgaria: pollen analysis and radiocarbon dating of a core from Lake Ribno Ban derishko, *Holocene*, 12, 201–210, <https://doi.org/10.1191/0959683602hl535rp>, 2002.

- van der Knaap, W. O., van Leeuwen, J. F. N., Fankhauser, A., and Ammann, B.: Palynostratigraphy of the last centuries in Switzerland based on 23 lake and mire deposits: chronostratigraphic pollen markers, regional patterns, and local histories, *Rev. Palaeobot. Palyno.*, 108, 85–142, [https://doi.org/10.1016/S0034-6667\(99\)00035-4](https://doi.org/10.1016/S0034-6667(99)00035-4), 2000.
- van der Knaap, W. O., van Leeuwen, J. F. N., and Ammann, B.: Seven years of annual pollen influx at the forest limit in the Swiss Alps studied by pollen traps: relations to vegetation and climate, *Rev. Palaeobot. Palyno.*, 117, 31–52, [https://doi.org/10.1016/S0034-6667\(01\)00075-6](https://doi.org/10.1016/S0034-6667(01)00075-6), 2001.
- van der Knaap, W. O., van Leeuwen, J. F. N., Svitavská-Svobodová, H., Pidek, I. A., Kvavadze, E., Chichinadze, M., Giesecke, T., Kaszewski, B. M., Oberli, F., Kalniņa, L., Pardoe, H. S., Tinner, W., and Ammann, B.: Annual pollen traps reveal the complexity of climatic control on pollen productivity in Europe and the Caucasus, *Veg. Hist. Archaeobot.*, 19, 285–307, <https://doi.org/10.1007/s00334-010-0250-6>, 2010.
- 10 Veski, S., Amon, L., Heinsalu, A., Reitalu, T., Saarse, L., Stivrins, N., and Vassiljev, J.: Lateglacial vegetation dynamics in the eastern Baltic region between 14,500 and 11,400 cal yr BP: A complete record since the Bølling (GI-1e) to the Holocene, *Quaternary Sci. Rev.*, 40, 39–53, <https://doi.org/10.1016/j.quascirev.2012.02.013>, 2012.
- Wayne, P., Foster, S., Connolly, J., Bazzaz, F., and Epstein, P.: Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO₂-enriched atmospheres, *Ann. Allerg. Asthma Im.*, 88, 279–282, [https://doi.org/10.1016/S1081-1206\(10\)62009-1](https://doi.org/10.1016/S1081-1206(10)62009-1), 2002.
- 15 Welten, M.: Pollenanalytische, stratigraphische und geochronologische Untersuchungen aus dem Faulenseemoos bei Spiez., vol. 21 of *Veröffentlichungen des Geobotanischen Instituts Rübel in Zürich*, Verlag H. Huber, 1944.
- Williams, J. W., Grimm, E. C., Blois, J. L., Charles, D. F., Davis, E. B., Goring, S. J., Graham, R. W., Smith, A. J., Anderson, M., Arroyo-Cabres, J., Ashworth, A. C., Betancourt, J. L., Bills, B. W., Booth, R. K., Buckland, P. I., Curry, B. B., Giesecke, T., Jackson, S. T., Latorre, C., Nichols, J., Purdum, T., Roth, R. E., Stryker, M., and Takahara, H.: The Neotoma Paleoecology Database, a multiproxy, international, community-curated data resource, *Quaternary Res.*, 89, 156–177, <https://doi.org/10.1017/qua.2017.105>, 2018.
- 20