



Enrichment of trace metals from acid sulphate soils in sediments of the Kvarken Archipelago, eastern Gulf of Bothnia, Baltic Sea

Joonas J. Virtasalo¹, Peter Österholm², Aarno T. Kotilainen¹, Mats E. Åström³

¹Marine Geology, Geological Survey of Finland (GTK), Espoo, 02150, Finland

5 ²Department of Geology and Mineralogy, Åbo Akademi University, Turku, 20500, Finland

³Department of Biology and Environmental Science, Linnaeus University, Kalmar, 39182, Sweden

Correspondence to: Joonas J. Virtasalo (joonas.virtasalo@gtk.fi)

Abstract. Rivers draining the acid sulphate soils of western Finland are known to deliver large amounts of trace metals with detrimental environmental consequences to the recipient estuaries in the eastern Gulf of Bothnia, northern Baltic Sea. However, the distribution of these metals in the coastal sea area, and the relevant metal transport mechanisms have been less studied. This study investigates the spatial and temporal distribution of metals in 9 sediment cores, collected from the Kvarken Archipelago, which is the recipient of Laihianjoki and Sulvanjoki Rivers that are impacted by acid sulphate soils. The contents of Cd, Co, Cu, La, Mn, Ni and Zn increase in the cores during the 1960s and 1970s as a consequence of intensive artificial drainage of the acid sulphate soil landscape. The metal deposition has remained at the high level since the 1980s. The metal enrichment in seafloor sediments is currently visible at least 25 km seaward from the river mouths. Comparison to sediment quality guidelines shows that the metal contents are very likely to cause detrimental effects on marine biota more than 12 km out from the river mouths. The dynamic sedimentary environment of the shallow archipelago makes these sediments potential future sources of metals to the ecosystem. Finally, the strong association of metals and nutrients to the same sediment grain size class of 2–6 µm indicates that the transformation of dissolved organic matter and metals to metal-organic aggregates at the river mouths is the key mechanism of seaward trace metal transport, in addition to co-precipitation with Mn-oxyhydroxides identified in previous studies. The large share of terrestrial organic carbon of the total organic C in these sediments (interquartile range = 39–48%) highlights the importance of riverine organic matter supply. These findings are important for the estimation of environmental risks and the management of biologically-sensitive coastal sea ecosystems.

1 Introduction

25 Acid sulphate (AS) soils are regarded as the nastiest soils in the world due to their ability to generate sulphuric acid and extremely low pH to water phase (Dent and Pons, 1995). The global distribution of these soils is estimated at 50 million ha including areas in Australia, Africa, Central and South America, South and Southeast Asia, Scandinavia and Western Europe (Andriessse and van Mensvoort, 2006; Michael et al., 2017). In Europe, the largest occurrences of AS soils are probably found in Finland where the soils are currently being mapped. Current rough estimates point to AS soils occupying an area in the order of 1 million ha in Finland (Anton Boman, personal communication).

35 AS soils in northern Europe are organic-rich sulphide-bearing muds that were originally deposited in the Baltic Sea during its brackish-water phase, which began ca. 7000 years ago in the Gulf of Bothnia (Virtasalo et al., 2007; Häusler et al., 2017). These muds have since emerged above sea level as a result of rapid land uplift (today 4–9 mm/year in western Finland; Mäkinen and Saaranen, 1998; Kakkuri, 2012). Artificial drainage and reclamation of these lands for farming purposes, which was particularly intensive in Finland in the 1960s and 1970s, has caused significant lowering of groundwater level (Saarinen et al., 2010; Yu et al., 2015). The groundwater lowering has enabled rapid oxidation of metal sulphide minerals that are abundant in these muds, producing H₂SO₄ and resulting in AS soils with a pH <4 (Yli-Halla et al., 1999; Sohlenius and Öborn, 2004; Boman et al., 2010). Under these extreme acidic conditions, large quantities of metals are released to the porewater due to the



40 oxidative dissolution of metal sulphides and weathering of silicate minerals. Particularly during high water flow conditions in
spring and autumn, acidic porewater rich in metals (e.g. Al, Cd, Co, La, Mn, Ni and Zn) is flushed to recipient streams, with
detrimental ecological consequences on biodiversity and the community structure of fish, benthic invertebrates and aquatic
plants (Hudd and Kjellman, 2002; Fältmarsch et al., 2008; Sutela and Vehanen, 2017). It is expected that climate change will
increase the acidic runoff and metal loading from AS soils (Saarinen et al., 2010; Veijalainen et al., 2012; Nystrand et al.,
45 2016).

When acidic metal-rich river waters from the boreal AS soil landscape are discharged to estuaries with higher pH and salinity,
the metals are complexed with organic matter or co-precipitated with Al-, Fe- and Mn-oxyhydroxides and consequently
deposited in sediments (Åström and Corin, 2000; Nystrand et al., 2016). Low-density organic aggregates have the capacity to
50 be transported across estuarine gradients, and their role in the seaward trace metal transport has been highlighted in recent
studies, particularly in the Baltic Sea (Gustafsson et al., 2000; Herzog et al., 2020; Jokinen et al., 2020). It has been shown that
trace metals from AS soils are enriched in seafloor sediments near the mouths of rivers draining from the coastal plains of
western Finland compared to background values and to the parent AS soil material (Nordmyr et al., 2008a, 2008b), with
documented deteriorating effects on local benthic invertebrate communities (Wallin et al., 2015). Less is known, however,
55 about the distribution of these metals in sediments further out from the river mouths, although this information is important
for the estimation of ecotoxicological effects and the management of biologically-sensitive coastal sea ecosystems (e.g. de
Souza Machado et al., 2016). The metals distribution pattern is also informative about the seaward metal transport mechanisms.

Permanent sediment deposition in the eastern coastal Gulf of Bothnia is restricted to small patches due to shallow water depths
60 and the openness of the area to dominant southwesterly winds (waves) (Kotilainen et al., 2012). This pattern of sediment
deposition and erosion is in slow but constant change due to the uplift, which makes these recent sediment patches prone to
remobilization and potential secondary sources of metals to the marine ecosystem. In addition, the building and maintenance
of offshore infrastructure such as shipping lanes requires repeated and, on the long term, expanding dredging, which can result
in resuspension and redistribution of metals in the environment (Lehoux et al., 2020). Finally, climate models project stronger
65 westerly winds and shorter ice season in winter over Northern Europe (Ruosteenoja et al., 2019), which have the potential to
increase seafloor erosion and sediment redistribution in the study area. Therefore, understanding the transport and distribution
of metals in seafloor sediments is particularly important in the dynamic eastern coastal Gulf of Bothnia.

This is the first assessment of trace metal distribution in seafloor sediments with distance from rivers running through boreal
70 AS soil landscapes. The studied sea area is in the Kvarken Archipelago, off the town of Vaasa, in the eastern Gulf of Bothnia,
which receives significant amounts of metals from Laihianjoki and Sulvanjoki Rivers that are among the most AS soil impacted
rivers in Finland and Europe (Roos and Åström, 2005). The focus is on sediments deposited before and after the intensive
artificial drainage of the AS soil landscape beginning in the 1960s. The focus is further on metals that are known to be
extensively leached from AS soils (Al, Cd, Co, Cu, La, Mn, Ni and Zn) and much less so from the local industry and other
75 human activities in the area (c.f. Åström and Björklund, 1997; Åström and Corin, 2000; Österholm and Åström, 2002).

2 Study area

The Kvarken Archipelago is located in the Gulf of Bothnia, where it marks the border between the Bothnian Sea in the south
and the Bothnian Bay in the north (Fig. 1). The area was covered by the Fennoscandian Ice Sheet during the latest (Weichselian)
glaciation, which retreated from the area ca. 10 400 years ago (Sauramo, 1929; Saarnisto and Saarinen, 2001; Stroeven et al.,
80 2016). Kvarken is a UNESCO World Heritage Site because it is considered as a prime location demonstrating the effects of



rapid glacioisostatic uplift (today ca. 8 mm/year; Mäkinen & Saaränen, 1998; Kakkuri, 2012) on shoreline displacement and changes in coastal landscape. During and just after deglaciation, the archipelago was submerged to a water depth of 250–280 m, whereas today the area is very shallow (<25 m) and shoaly, with approximately 7000 islands and islets (Breilin et al., 2005; Ojala et al., 2013). The rapid uplift has led to strong seafloor erosion and sediment transport to deep areas further offshore.

85 Till covers ca. 70 % of the modern seafloor in the archipelago (Kotilainen et al., 2012). Glaciolacustrine varved silts and clays, and postglacial lacustrine weakly-layered silty clays cover ca. 18 % of the seafloor, whereas bedrock outcrops are rare (3 % of the seafloor). Patches of recent mud deposition cover only ca. 8 % of the seafloor (Kotilainen et al., 2012).

The Kvarken Archipelago belongs to the continental subarctic climate zone with severe dry winters and almost warm summers.

90 The mean annual air temperature is 4.2 °C, with the mean minimum temperature of 2.1 °C and the mean maximum temperature of 6.6 °C during the period 1981–2010 (Pirinen et al., 2012). The mean annual precipitation is 497 mm. The Bothnian Sea freezes on an annual basis and remains frozen for up to 140–150 days per year. The annual mean sea surface salinity in the archipelago ranges between 3.5 and 4 PSU and the annual mean sea surface temperature between 3.5 and 7 °C. The sea is essentially non-tidal, but irregular water level fluctuations of as much as ±1.5 m take place as a result of variations in wind

95 and atmospheric pressure. Stratification of the shallow waters is governed by thermocline that develops each summer. The area is generally less affected by eutrophication and the associated seafloor oxygen deficiency, which are widespread in the southern and central Baltic Sea (Lundberg et al., 2009).

3 Materials and Methods

3.1 Sediment coring

100 The fieldwork was carried out during the summers 2016–2018 onboard the research vessel *Geomari* of the Geological Survey of Finland. *Geomari* is equipped with a marine geological seismoacoustic survey system, which includes Meridata 28 kHz pinger and Massa TR-61A 3.5–8 kHz CHIRP sub-bottom profilers that were essential for the identification of coring sites that are representative of recent sediment deposition.

105 Altogether 9 sediment cores (Table 1) were collected using a Gemax twin-barrel short gravity corer (core diameter 9 cm), which preserves the soft sediment surface essentially undisturbed. One core of each twin was cut in half lengthwise and cleaned for sedimentological description and photography, whereas the other core was sectioned using a rotary device into 1 cm sample slices. The sample slices were stored in cool and dark until shore-based laboratory analysis.

3.2 Laboratory analyses

110 Sample slices were analysed for ¹³⁷Cs activity content in order to constrain sediment chronology in each core. The ¹³⁷Cs activity of fresh samples was measured for 60 min using a BrightSpec bMCA-USB pulse height analyser coupled to a well-type NaI(Tl) detector at the Geological Survey of Finland (Ojala et al., 2017). Each core was analyzed starting from the uppermost sample slice and progressing downward until near zero (background) activity levels were measured in at least three consecutive samples. No corrections were applied for the results because the aim was only to detect relative ¹³⁷Cs activity peaks. Due to

115 the possible post-depositional downward transport of ¹³⁷Cs through bioturbation and diffusion (Holby and Evans, 1996; Klaminder et al., 2012) the depth of peak ¹³⁷Cs activity (rather than the initial increase) was assumed to represent the fallout from the 1986 Chernobyl nuclear disaster. Sample slices for each core were classified to those deposited in 1986 and later, and those deposited before 1986. The samples deposited before 1986 were further classified to those deposited after and before the year 1960, by calculating the average thickness of sediment deposited annually after 1986 in each core, and estimating the

120 depth of 1960 by assuming a constant sedimentation rate for that core. This approach potentially slightly overestimates the



depth of the year 1960 because of the increasing sediment compaction with core depth. The approach should, therefore, be viewed as conservative to sediments deposited before the year 1960.

125 After the non-destructive ^{137}Cs analysis, fresh sample slices were freeze-dried, homogenized and halved, with one half analysed for multielement composition and the other for grain size distribution at the commercial laboratory Eurofins Labtium Ltd (Kuopio, Finland). The material for multielement analysis was sieved through a 63 μm mesh, and 0.2 g of the passed-through fraction was digested in a four-acid mixture of hydrofluoric acid, perchloric acid, hydrochloric acid and nitric acid (USGS Methods T01 and T20). After evaporation of the acids at 160 $^{\circ}\text{C}$, the resulting gel was dissolved to 1 M HNO_3 , and analysed for element concentrations by inductively coupled plasma-mass spectrometry (ICP-MS), or inductively coupled plasma-optical emission spectrometry (ICP-OES), depending on the element. Because HF dissolves silicate minerals, the digestion is considered as “near-total digestion” (Hall et al., 1996). The commercial sediment reference materials QCGBMS304-6, QCMES-4, QCNIST8704, CO153B and in-house standards were used for assessing measurement accuracy. Element concentrations for all reference materials measured with each sample batch fell well within $\pm 10\%$ of the certified values. Mercury was measured separately by HNO_3 leach of 0.2 g samples through thermal decomposition, amalgamation and atomic absorption spectrometry (US EPA Method 7473). Solid-phase contents of carbon and nitrogen in the samples were analyzed by thermal combustion elemental analysis (TCEA). The pools of inorganic C and N are negligible in this setting (Virtasalo et al., 2005; Jilbert et al., 2018), hence no decalcification was conducted and the total contents are considered equal to organic C and N.

140 To quantify the proportions of terrestrial plant-derived ($\% \text{OC}_{\text{terr}}$) and phytoplankton-derived organic matter in the C pool, a simple binary mixing model was applied for the molar N/C ratio, assuming end-member values of $(\text{N}/\text{C})_{\text{terr}} = 0.04$ and $(\text{N}/\text{C})_{\text{phyt}} = 0.13$ following Goñi et al. (2003) and Jilbert et al. (2018):

$$\% \text{OC}_{\text{terr}} = \frac{(\text{N}/\text{C})_{\text{sample}} - (\text{N}/\text{C})_{\text{phyt}}}{(\text{N}/\text{C})_{\text{terr}} - (\text{N}/\text{C})_{\text{phyt}}} \times 100, \quad (1)$$

145 The model integrates a variety of terrestrial organic-matter sources ranging from fresh vascular plant detritus to more degraded soil organic matter into a single end-member. This is practical since effectively all of the organic matter transported by rivers passes through the soil reservoir before entering the coastal zone, therefore representing a mixture of variably degraded material (Jokinen et al., 2018).

150 Grain size distribution was determined for selected freeze-dried samples by wet-sieving through 20 mm, 6.3 mm, 2 mm, 0.63 mm, 0.2 mm and 0.063 mm ISO 3110/1 test sieves. The samples were pretreated with excess H_2O_2 to remove organic matter prior to the analysis. The $<63 \mu\text{m}$ size fraction was further analyzed down to 0.6 μm using a Micromeritics Sedigraph III 5120 Xray absorption sedimentation analyzer. The sieving results were merged with sedimentation data in Sedigraph software. Median grain size was calculated according to the geometric Folk and Ward (1957) graphical measures implemented in GRADISTAT 4.0 software (Blott and Pye, 2001). Clay is defined as grains finer than 2 μm , whereas mud is clay and silt ($<63 \mu\text{m}$), and sand is 63 μm to 2 mm (Blott and Pye, 2012).

3.3 Statistical analysis

Element contents below detection were rounded to half the detection limits so that approximate values could be used in the analyses.

160 In order to explore relationships between elements in the produced multielement dataset, a robust compositional principal component analysis (PCA) after isometric logratio (ilr) transformation (Filzmoser et al., 2009) was carried out using the



robCompositions 2.0.8 package in the R 3.5.1 software. The resultant loadings and scores were back-transformed to centered logratio (clr) space for meaningful visualization and interpretation in a compositional biplot (Filzmoser et al., 2018).

165 Relationships between elements and grain size classes were explored using partial least squares regression 2 (PLSR2; Tenenhaus, 1998) as implemented in the `plsdepot` 0.1.17 package in the R 3.5.1 software. The PLSR2 results were validated using hierarchical partitioning (Chevan and Sutherland, 1991) as implemented in the `hier.part` 1.0.4 package (Nally and Walsh, 2004) in the R.

4 Results

170 Multielement and grain size data produced in this study are available in PANGAEA (Virtasalo et al., 2020). The studied sediments are poorly sorted with a narrow grain size range: the interquartile range (IQR) of median grain sizes of all sample slices is 1.96–2.54 μm , with a median of 2.18 μm . The sediments are classified as clayey silt according to (Blott and Pye, 2012) and silty clay according to soil taxonomy (Soil Survey Staff, 1999).

4.1 Vertical distribution

175 Peak ^{137}Cs activity is easily distinguishable in all the studied sediment cores, which permits the confident identification of the depth of the Chernobyl fallout year 1986 in each core (Fig. 2). The clearly defined activity peak in each core excludes significant sediment reworking and post-event migration of ^{137}Cs , and supports the estimation of the depth of the year 1960 by assuming a constant sedimentation rate.

180 Contents of metals Cd, Co, Cu, La, Ni and Zn generally begin to increase approximately at the depth of the year 1960 in the studied cores (Fig. 2). An exception is MGGN-2017-19 from the eastern Korshamnsfjärden, close to the rivers, where the metal contents are high and variable with no clear trend below the core depth of 16 cm (early 1980s), but show an upward increasing trend above this level. The metals reach particularly high contents in MGGN-2017-20 from Varisselkä at ca. 1986, after which they begin to decrease at that site. In all cores from Korshamnsfjärden (MGGN-2017-17, MGGN-2017-18, 185 MGGN-2017-19), the increasing trends of Co, Ni, La and Zn continue overall to the core top, whereas the increasing trends of Cu and Cd level out or turn to decrease at ca. 1986. In cores from farther out at sea in Gloppet (MGGN-2018-29, MGGN-2018-31) from ca. 1986 to the core tops, the contents of Co, Ni and La vary at high values, whereas Zn, Cu and Cd show a decreasing trend.

190 Aluminium contents are variable but show no clear trends in the cores, except in MGGN-2017-20 from Varisselkä, where the vertical distribution of Al is similar to other metals in that core (Fig. 2).

Manganese contents are high and variable with an upward increasing trend in cores from Varisselkä, and eastern and middle Korshamnsfjärden (MGGN-2017-18, MGGN-2017-19, MGGN-2017-20; Fig. 2). In cores from farther out at sea in western 195 Korshamnsfjärden and Gloppet (MGGN-2017-17, MGGN-2018-29, MGGN-2018-31), Mn contents are low except a strong increase at the core tops.

Carbon contents increase toward the top in all cores. In cores from Gloppet (MGGN-2018-29, MGGN-2018-31) and from western Korshamnsfjärden (MGGN-2017-17), C contents range between 2 and 3 % prior to ca. 1986, and between 3 and 4 % higher up the cores (Fig. 2). In middle and eastern Korshamnsfjärden (MGGN-2017-18, MGGN-2017-19), C contents vary 200 between 3 and 4 % before ca. 1986, and between 4 and 5 % in the upper core sections. In Varisselkä (MGGN-2017-20), C



205 contents increase strongly between 1960 and 1986, and reach 6 % in the upper section of that core. The terrestrial organic share of the C content before ca. 1986 is 40–50 % in cores from Gloppet and western and middle Korshamnsfjärden, and 50–60 % in cores from eastern Korshamnsfjärden and Varisselkä. After ca. 1986, the share of terrestrial organic carbon decreases in all cores, largely mirroring the upward-increasing C content.

4.2 Statistical relationships

Statistical analyses were carried out on the upper core sections that were deposited after the year 1960 because it is clear in the vertical metal content profiles (Fig. 2) that this interval is the most enriched in metals.

210 The first principal component (PC1) of the robust compositional PCA explains 73.7 % of the total variance. PC2 explains 10.7 %, whereas the rest of the components each explain less than 6 % of the total variance. The metals Co, Ni and Cd cluster along the positive side of PC1 in the biplot (Fig. 3). Also Mn has a strong positive loading on PC1, but it deviates slightly from the other metals. This deviation of Mn is likely explained by its vertical distribution in sediment cores that is similar to the other metals in Varisselkä and in eastern and middle Korshamnsfjärden, but different in cores collected farther out at sea (Fig. 2).

215 A two component PLSR2 model utilizes 82.3 % of the variance of predictor variables (54 elements) to explain 68.3 % of the variance of response variables (13 grain size classes). The metals Cd, Co, Cu, Ni and Zn have strong positive correlations with the grain size classes of 2–4 and 4–6 μm in the PLSR2 (Fig. 4a). Notably, also C and N are positively correlated with these grain size classes.

220 In concordance with the PLSR2, the hierarchical partitioning analysis shows that the 2–4, 4–6 and 1–2 μm classes have the most independent power among grain size classes in predicting Ni contents, and account for 17.8, 13.1 and 11.8 % of the explained variance, respectively (Fig. 4b). The hierarchical partitioning patterns of the other elements identified in the PLSR2 are similar.

225 4.3 Spatial distribution

Metal contents are compared between core sections deposited before the year 1960 and those deposited in 1986 and later in order to explore the magnitude of recent metal enrichment (Fig. 5). Metal contents decrease toward the bottom in all cores; however, in Varisselkä, and Korshamnsfjärden East and Middle, metals do not decrease to low levels comparable to those in western Korshamnsfjärden and Gloppet (Fig. 2). Therefore, mean median metal contents in the pre-1960 sections of the four
230 cores from Gloppet (MGGN-2018-29, MGGN-2018-30, MGGN-2018-31 and MGGN-2018-32) are considered to be representative of the local background values, and are used here as reference values in the depiction of spatial trends.

Median contents of Cd, Co, Cu, La, Mn, Ni and Zn in sediment cores decrease with distance from the Laihianjoki and Sulvanjoki Rivers in the east (Fig. 5; Table 2; Supplement). Comparison between core sections deposited before 1960 and in
235 1986 and later shows that median contents of Cd, Co, Cu, La, Ni and Zn are higher in the upper sections of all cores, with the difference increasing toward east. However, Mn median contents are enriched only in the upper sections of the four cores closest to the rivers (MGGN-2016-8, MGGN-2017-18, MGGN-2017-19, MGGN-2017-20), whereas in the cores further offshore, there is essentially no difference between the upper and lower core sections (Fig. 5). In contrast, median Al contents are essentially uniform in all the cores, and between upper and lower sections.

240 When sediment cores are arranged according to distance from the Laihianjoki and Sulvanjoki Rivers, the pattern of decreasing metal contents with the increasing distance is evident (Fig. 6). The metal median contents also exceed several sediment quality



245 guideline thresholds. The metal contents have not been normalized because the sediment samples contain more than 30 %
clay, and on median 3.7 % C, which means that normalization coefficients according to the Finnish sediment dredging and
dumping guidelines are close to 1.

5 Discussion

Contents of trace metals known to be abundantly leached from AS soils (Al, Cd, Co, Cu, La, Mn, Ni and Zn) have been studied
in sediment cores from the Kvarken Archipelago, which is the recipient sea area of the Laihianjoki and Sulvanjoki Rivers.
These rivers are among the most AS soil impacted rivers in Finland and Europe (Roos and Åström, 2005).

250 5.1 Metal distribution

The median contents of Al are essentially uniform in the studied cores, both laterally with distance from the rivers, and between
the core sections deposited before the year 1960 and in 1986 and later (Fig. 5). The vertical distribution of Al is similar to the
other metals with the highest values in the 1980s in the core from Varisselkä (MGGN-2017-20), but different at the other sites
(Fig. 2). Clearly, the intensive artificial drainage of the AS soil landscape, which began in the 1960s (Saarinen et al., 2010; Yu
255 et al., 2015), has not substantially influenced the delivery of Al to the coring sites other than Varisselkä. IQR of Al contents in
sections deposited in 1986 and later in all cores is 67900–75200 mg/kg (Table 2), and the maximum Al content is 110 000
mg/kg. A higher median Al content (86400 mg/kg; Table 2) has been reported near the mouth of the Vöyrinjoki River
(Nordmyr et al., 2008b), which is situated ca. 37 km northeast from the nearest coring site (Fig. 1b). This is in good agreement
with previous observations, which show that Al to a large extent is deposited very close river mouths together with organic
260 material (Nordmyr et al., 2008a, 2008b; Åström et al., 2012; Nystrand et al., 2016). Wallin et al. (2015) report the Al content
of 59900 mg/kg for a single sample “from the first accumulation basin in the estuary” of the Laihianjoki River (Table 2), but
do not provide coordinates or a map of the sampling location, which makes it difficult to assess the representativeness of the
sample.

265 Manganese median contents are enriched at the four sites closest to the rivers in the east (MGGN-2016-8, MGGN-2017-18,
MGGN-2017-19, MGGN-2017-20) compared to cores collected farther out at sea (Fig. 5). The enrichment is stronger in the
upper core sections deposited in 1986 and later than in the core sections deposited before 1960. The enrichment is in line with
the previously documented increase of metal loading to estuaries in western Finland as a consequence of increased artificial
drainage of the AS soil landscape, beginning in the 1960s (Yu et al., 2015, 2016). The vertical distribution of Mn shows
270 elevated values in upper core sections similar to the other metals at the easternmost sites (Fig. 2; MGGN-2017-18, MGGN-
2017-19, MGGN-2017-20). However, in cores farther out at sea in western Korshamnshjärden and Gløppet, the vertical
distribution of Mn is generally flat except a pronounced increase at the core tops. IQR of Mn in core sections deposited in
1986 and later in the easternmost sites is 3740–7300 mg/kg, whereas it is 548–1293 mg/kg at the sites farther out at sea. The
maximum Mn content in the upper core sections is 14000 mg/kg. Similar Mn contents to the easternmost sites have been
275 reported from the open sea areas of Bothnian Sea and Bothnian Bay: mean 3000 ±1600 mg/kg and 8500 ±5300 mg/kg,
respectively (Leivuori and Niemistö, 1995). Higher Mn contents (median 9013 mg/kg; Table 2) have been reported near the
mouth of the Vöyrinjoki River (Nordmyr et al., 2008b). Manganese enrichment in upper core sections is evident from
Varisselkä (MGGN-2017-18) to eastern and middle Korshamnshjärden (MGGN-2017-19, MGGN-2017-20), which shows that
it is transported longer distances from the rivers than Al. This observation is in line with previous studies, which demonstrate
280 that Mn can travel a long distance before precipitation as Mn-oxyhydroxides and the consequent deposition on the seafloor
(Nordmyr et al., 2008a, 2008b; Nystrand et al., 2016). Even farther out at sea, the strong increase of Mn at the core tops (Fig.
2) is due to the reductive dissolution of buried Mn-oxyhydroxides and associated release of Mn²⁺ into the porewater with



subsequent upward diffusion and oxidative precipitation of Mn as oxyhydroxides in the sediment surface layer (Widerlund and Ingri, 1996; Nordmyr et al., 2008b). The Mn content reported by Wallin et al. (2015) from the Laihianjoki River estuary is comparably low (788 mg/kg; Table 2).
285

The median contents of Cd, Co, Cu, La, Ni and Zn are higher at the four easternmost sites closest to the rivers (MGGN-2016-8, MGGN-2017-18, MGGN-2017-19, MGGN-2017-20) compared to those farther offshore (Fig. 5; Supplement). In contrast to Al and Mn, these metals are enriched in the upper core sections deposited in 1986 and later at all the coring sites compared to the lower sections deposited before 1960. Vertical distributions of these metals show increasing upward trends beginning at ca. 1960 in all cores, except in MGGN-2017-19 (eastern Korshamnsfjärden), where the initial metal contents are high and variable with no clear trend until they begin to increase in the early 1980s (Fig. 2). IQRs of Cd, Ni and Zn, for example, in core sections deposited in 1986 and later are 0.75–1.40 mg/kg, 51–107 mg/kg, and 254–454 mg/kg, respectively (Table 2). Similar Cd contents have been reported from the Laihianjoki River estuary (0.92 mg/kg; Wallin et al., 2015) and the open sea area of Bothnian Bay (mean 0.8 ± 0.3 mg/kg; Leivuori and Niemistö, 1995), whereas lower Cd contents have been reported from the open Bothnian Sea (0.4 ± 0.2 mg/kg; Leivuori and Niemistö, 1995). A higher Ni content has been reported from the Laihianjoki River estuary (130.5 mg/kg; Wallin et al., 2015), and higher Zn contents from the estuaries of Laihianjoki (461 mg/kg; Wallin et al., 2015) and Vöyrinjoki Rivers (maximum 608.5 mg/kg; Nordmyr et al., 2008b). However, the maximum Cd, Ni and Zn contents in the upper core sections are generally 2–3 times higher than previously reported from the area: 3.11 mg/kg, 245 mg/kg, and 835 mg/kg, respectively.
290
295
300

After the strong increase in sedimentary metal contents during the 1960s and 1970s, the metal contents and thus metal loading from the AS soils has stayed overall at the same level since the 1980s (Fig. 2). In Korshamnsfjärden (MGGN-2017-17, MGGN-2017-18, MGGN-2017-19), the contents of Co, Ni, La and Zn generally continue to increase until the core top, whereas the increasing trends of Cu and Cd level out or turn to decrease at ca. 1986. In Gloppet (MGGN-2018-29, MGGN-2018-31), more than 25 km from the river mouths, the contents of Co, Ni and La vary at high values up to the core top, whereas Zn, Cu and Cd begin decrease from ca. 1986 onwards. An exception to this pattern is Varisselkä (MGGN-2017-20), where unexpectedly high contents of Al, Cd, Co, Cu, La, Ni and Zn were deposited in the early to mid-1980s (Fig. 2). In this core, the metal contents decrease toward the core top; however, despite this Varisselkä still has higher metal contents in the sediment surface than the other sites. The decrease in metal contents in Varisselkä parallels the decreasing share of terrestrial organic carbon since the 1980s, which suggests that the decrease in metal contents may be due to the reduced transport of metal-organic aggregates to the site (Section 5.2) as a result of e.g. narrowing of the shallow channel to the southeast (Fig. 5), rather than to a decrease in the metal loading to the archipelago. If the metal loading to the sea area had decreased, it would certainly be visible also in the Korshamnsfjärden cores, which it is not.
305
310

It is worth noting that permanent sediment deposition is today restricted to small patches in the eastern coastal Gulf of Bothnia due to the shallow water depths and openness of the sea area to dominant southwesterly winds (waves) (Kotilainen et al., 2012). As a consequence, finding coring sites that are representative of the recent sediment deposition can be challenging in the area without a guidance from seismoacoustic sub-bottom surveys such as those carried out here. For example, metal contents reported by Wallin et al. (2015) are generally lower than those measured here, although their sampling site supposedly was closer to the source rivers. The four-acid digestion method used in this study generally produces comparable results for metals from AS soils to previous studies that have aimed at analysing “total metal contents” in sediments, although different methods were used (Cook et al., 1997; Nordmyr et al., 2008a, 2008b).
315
320



5.2 Metal transport mechanisms

325 The similar distribution of Cd, Co, Cu, La, Ni and Zn in the studied sea area is supported by the PCA, which shows similar
behaviour of the metals, in particular Cd, Ni, and Co (Fig. 3). Previous studies of water samples, sediment trap material and
seafloor sediments have concluded that Cu and La precipitate readily close to river mouths, whereas Cd, Co, Ni and Zn are
preferentially transported a bit further out where they most likely co-precipitate and are deposited with Mn-oxyhydroxides
(Nordmyr et al., 2008a, 2008b; Nystrand et al., 2016). This study shows that Cd, Co, Cu, La, Ni and Zn all are enriched in
330 sediment cores farther out at sea than Mn, which strongly indicates that other mechanism(s) in addition to the precipitation of
Mn-oxyhydroxides influence their seaward transport and distribution.

Field studies and geochemical modelling show that Cd, Co, Cu, La, Ni and Zn in AS soil impacted rivers are transformed from
dissolved to particulate form as they are discharged to the sea (Nordmyr et al., 2008a, 2008b; Nystrand et al., 2016). Seaward
335 transport of suspended particles is highly dependent on hydrological conditions, with high discharge producing large plumes
of river water by which metals can be transported far from the river mouths (Nystrand et al., 2016). Exceptionally large river
plumes can be caused by extreme events such as that in the late autumn of 2006, when a severely dry summer (maximising
oxidation of sulphides in the AS soils) was succeeded by a severe wet spell (Österholm and Åström, 2008; Saarinen et al.,
2010), causing widespread fish kills in rivers and estuaries in western Finland. Some of the peaks observed in the metal vertical
340 distributions in sediment cores (Fig. 2) may result from such extreme events; however, exceptional events hardly explain the
overall metal enrichment in the cores.

The PLSR2 analysis, supported by hierarchical partitioning, shows that Cd, Co, Cu, Ni and Zn are strongly positively correlated
with sediment grains of the size between 2 and 6 μm (Fig. 4). Also the nutrients C and N have strong positive correlations with
345 the same grain-size range, which indicates that the metals are associated with organic particles. This observation is supported
by recent studies, which demonstrate the importance of metal-organic matter aggregates in land-to-sea transfer of trace metals,
particularly in boreal environments (Jokinen et al., 2020). When riverine dissolved organic matter is transformed to particulate
form as it is discharged to the sea, dissolved metals are also transformed to particulate form, and passively enclosed in the
produced metal-organic matter aggregates (Stolpe and Hassellöv, 2007; Valikhani Samani et al., 2015; Herzog et al., 2020).
350 The low-density organic aggregates can be transported by currents far from the river mouth, as has been demonstrated for the
Kalix River in the Bothnian Bay (Gustafsson et al., 2000), and elsewhere (Regnier and Wollast, 1993; Wang and Wang, 2016;
Pavoni et al., 2020a, 2020b). Organic aggregates in coastal environments are loosely bound and fragile, and have a size range
of tens to thousands of micrometers (Eisma, 1986; Mikkelsen et al., 2006; Lee et al., 2012). The aggregates are easily broken
after deposition by benthic organisms (e.g. Rhoads and Boyer, 1982), by sediment compaction with burial and, ultimately, by
355 the grain size analysis, to their constituent particles, which typically are smaller than 20 μm (Eisma, 1986; Mikkelsen et al.,
2006; Lee et al., 2012). The 2–6 μm size range identified here differs from the local phytoplankton community, which is
dominated by species smaller than 2 μm during summer, and those larger than 10 μm in winter and spring (Andersson et al.,
1996; Paczkowska et al., 2017). The 2–6 μm range is slightly larger than the median sediment grain size, the IQR of median
grain size of the studied samples being 1.96–2.54 μm .

360 The importance of riverborne organic matter in the sea area is demonstrated by the large share of terrestrial organic carbon in
the cores. IQR of the share of terrestrial organic carbon of the total organic C in the cores is 39.2–47.8 %, which is substantially
higher than in, e.g. coastal sea areas of southern Finland, where the terrestrial share usually is less than 30 % (Jilbert et al.,
2018; Jokinen et al., 2018, 2020). The terrestrial share is highest in cores from eastern Korshamnsfjärden (MGGN-2017-19)
365 and Varisselkä (MGGN-2017-20) closest to the river mouths (Fig. 2). The share of terrestrial organic carbon decreases upward
in core sections deposited in 1986 and later, largely mirroring the increase in total C, which indicates that the increase in total



organic C is largely driven by increasing phytoplankton production during the recent ca. 30 years, in line with the observed increasing nutrient levels in the inner Kvarken Archipelago since 1980 (Lundberg et al., 2009).

5.3 Risk assessment

370 Metals associated with particles are eventually settled and buried in sediments, and are therefore less available for the aquatic biota. However, particulate metals in sediments may be toxic to benthic invertebrates via gastrointestinal tract and skin (Eggleton and Thomas, 2004; Wallin et al., 2015). Metals may also be dissolved from sediments to the aqueous phase if seafloor physical-chemical conditions are altered or sediment is bioturbated (Eggleton and Thomas, 2004; de Souza Machado et al., 2016).

375

Finnish sediment dredging and dumping guidelines provide metal content thresholds for the assessment of the suitability of material for offshore dumping (Ympäristöministeriö, 2015). Level 2 thresholds in the guidelines are defined so that metal contents exceeding the levels cause acute toxicity in less than 5 % of marine organisms. Zinc contents in the majority of the samples and Cd contents in half of the samples that were deposited in 1986 and later in Varisselkä (MGGN-2017-20) exceed 380 the level 2 threshold, which means that these sediments are considered unsuitable for offshore disposal (Fig. 6). Nickel contents in almost half of the samples exceed the level 2 threshold as far as 14.2 km from the nearest river mouth (MGGN-2017-17, western Korshamnshjärden).

385 Finnish stakeholders often use North American and Canadian guidelines when assessing the environmental impacts of metals in marine sediments because of similar geological environment (e.g. Vallius, 2015). The North American guidelines determine metal toxicity in sediment relative to two threshold levels: effects range-low (ERL) and effects range-medium (ERM). Metal contents exceeding the ERMs frequently result in adverse effects on biota, whereas metal contents between ERLs and ERMs occasionally result in adverse effects on biota, and metal contents below ERLs rarely result in adverse effects on biota (Long et al., 1995). The Canadian Guidelines for Protection of Aquatic Life consist of the Interim Sediment Quality Guidelines 390 (ISQGs) and the Probable Effect Levels (PELs), which are used to evaluate the biological effects of a contaminant (Canadian Council of Ministers of the Environment, 2001). Contents exceeding the PELs frequently result in adverse effects on biota, whereas levels between the PELs and the ISQGs are associated with infrequently occurring adverse effects. Levels below the ISQG rarely cause adverse effects.

395 More than half of the measured Zn contents exceed the ERL and ISQG levels at each coring site, frequently even in the core sections deposited before 1960 (Fig. 6). Furthermore, the majority of measured Zn contents exceed the ERM and PEL levels as far as 12.6 km from the nearest river mouth (MGGN-2017-18, middle Korshamnshjärden). The majority of measured Ni contents exceed the ERL at each coring site, and the ERM as far as 14.2 km from the nearest river mouth (MGGN-2017-12, western Korshamnshjärden). The majority of Cd contents exceed the ISQL as far as 24.7 km from the nearest river (MGGN-400 2018-30, Glöppet), and the ERL at 12.6 km from the nearest river (MGGN-2017-18, middle Korshamnshjärden). It seems likely that metal loading from AS soils has detrimental effects on biota in the studied sea area. The exotoxicological risk of metal loading from AS soils was previously assessed to be high in the area by Wallin et al. (2015).

6 Conclusions

405 Loading from AS soils has resulted in the strong enrichment of Cd, Co, Cu, La, Mn, Ni and Zn in sediments of the Kvarken Archipelago. The loading intensified in the 1960s and 1970s, when previous studies show that intensive artificial drainage of the coastal AS soil landscape begun. The metal deposition has remained at more or less the same level since the 1980s, however



with fine-scale variability in contents both among the metals and sampling sites. Metal transport from the Laihianjoki and Sulvanjoki Rivers toward open sea largely takes place along Korshamnshjärden. The metal enrichment in seafloor sediments is currently visible at more than 25 km distance from the rivers. Comparison to sediment quality guidelines shows that metal contents in the majority of analysed sub-samples are sufficiently high to very likely have detrimental effects on marine biota more than 12 km from the river mouths. The dynamic nature of the patchy sediment deposition, the rapid uplift of the region, and the predicted increase in storm wave erosion with climate change imply that these sediments are potential future sources of metals to the marine ecosystem. Furthermore, acidic runoff and metal loading from acid sulphate soils have been predicted to increase with the climate change.

Previous studies have identified Mn-oxyhydroxides as a mechanism of metal transport and deposition seaward from the river mouths in the area. This study shows that Cd, Co, Cu, La, Ni and Zn are transported further out at sea than Mn, which requires an additional mechanism of metal transport. The strong association of the metals and nutrients to sediment grains of the same size range (2–6 µm) indicates that the transformation of dissolved organic matter and metals to metal-organic aggregates at the river mouths is the key mechanism of seaward trace metal transport. The large share of terrestrial organic carbon of the total organic C in these sediments (interquartile range 39–48 %) highlights the importance of riverine organic matter supply.

Data availability. Multielement and grain size data are available in PANGAEA (Virtasalo et al., 2020).

Supplement. The supplement related to this article is available online at: [journal web page]

Author contributions. Joonas Virtasalo: Writing – original draft, Conceptualization, Investigation, Formal analysis, Funding acquisition. Mats Åström: Writing – review and editing, Conceptualization. Peter Österholm: Writing – review and editing, Conceptualization. Aarno Kotilainen: Writing – review & editing, Investigation, Funding acquisition, Project administration.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study resulted from the SmartSea project, funded by the Strategic Research Council at the Academy of Finland (grant number 292 985). M.E.Å. additionally acknowledges the Swedish Research Council Formas (grant number 2018-00760). The study has utilized research infrastructure facilities provided by FINMARI (Finnish Marine Research Infrastructure network). Help by the crew of R/V *Geomari* and those who assisted in the field work is gratefully acknowledged. Satu Vuoriainen carried out the ¹³⁷Cs analyses. The reviewers are thanked for comments that helped improve the manuscript.

References

Andersson, A., Hajdu, S., Haecky, P., Kuparinen, J., and Wikner, J.: Succession and growth limitation of phytoplankton in the Gulf of Bothnia (Baltic Sea), *Mar. Biol.*, 126, 791–801, <https://doi.org/10.1007/BF00351346>, 1996.

Andriessse, W., and van Mensvoort, M. E. F.: Acid sulfate soils: distribution and extent, in: *Encyclopedia of Soil Science*, edited by: Lal, R., CRC Press, Boca Raton, U.S.A, 14–19, <https://doi.org/10.1081/E-ESS3-120006641>, 2006.

Åström, M., and Björklund, A.: Geochemistry and acidity of sulphide-bearing postglacial sediments of western Finland, *Environ. Geochem. Hlth.*, 19, 155–164, <https://doi.org/10.1023/A:1018462824486>, 1997.



450 Åström, M., and Corin, N.: Abundance, sources and speciation of trace elements in humus-rich streams affected by acid sulphate soils, *Aquat. Geochem.*, 6, 367–383, <https://doi.org/10.1023/A:1009658231768>, 2000.

Åström, M. E., Österholm, P., Gustafsson, J. P., Nystrand, M., Peltola, P., Nordmyr, L., and Boman, A.: Attenuation of rare earth elements in a boreal estuary, *Geochim. Cosmochim. Ac.*, 96, 105–119, <https://doi.org/10.1016/j.gca.2012.08.004>, 2012.

455 Blott, S. J., and Pye, K.: GRADISTAT: a grain size distribution and statistics package for the analysis of unconsolidated sediments, *Earth Surf. Proc. Land.*, 26, 1237–1248, <https://doi.org/10.1002/esp.261>, 2001.

Blott, S. J., and Pye, K.: Particle size scales and classification of sediment types based on particle size distributions: review and recommended procedures, *Sedimentology*, 59, 2071–2096, <https://doi.org/10.1111/j.1365-3091.2012.01335.x>, 2012.

460

Boman, A., Fröjdö, S., Backlund, K., and Åström, M. E.: Impact of isostatic land uplift and artificial drainage on oxidation of brackish-water sediments rich in metastable iron sulfide, *Geochim. Cosmochim. Ac.*, 74, 1268–1281, <https://doi.org/10.1016/j.gca.2009.11.026>, 2010.

465 Breilin, O., Kotilainen, A., Nenonen, K., and Räsänen, M.: The unique moraine morphology, stratotypes and ongoing geological processes at the Kvarken Archipelago on the land uplift area in the western coast of Finland, in: *Quaternary Studies in the Northern and Arctic Regions of Finland: Proceedings of the Workshop Organized within the Finnish National Committee for Quaternary Research (INQUA), Kilpisjärvi Biological Station, Finland, January 13–14th 2005*, edited by: Ojala, A. E. K., Geological Survey of Finland, Special Paper, 40, 97–111, 2005.

470

Canadian Council of Ministers of the Environment: Canadian sediment quality guidelines for the protection of aquatic life: summary tables. Updated, Canadian Council of Ministers of the Environment, Winnipeg, Canada, 2001.

Chevan, A., and Sutherland, M.: Hierarchical Partitioning, *Am. Stat.*, 45, 90–96, <https://doi.org/10.2307/2684366>, 1991.

475

Cook, F. J., Hicks, W., Gardner, E. A., Carlin, G. D., and Froggatt, D. W.: Export of acidity in drainage water from acid sulphate soils, *Mar. Pollut. Bull.*, 41, 319–326, [https://doi.org/10.1016/S0025-326X\(00\)00138-7](https://doi.org/10.1016/S0025-326X(00)00138-7), 2000.

480 de Souza Machado, A. A., Spencer, K., Kloas, W., Toffolon, M., and Zarfl, C.: Metal fate and effects in estuaries: a review and conceptual model for better understanding of toxicity, *Sci. Total Environ.*, 541, 268–281, <https://doi.org/10.1016/j.scitotenv.2015.09.045>, 2016.

Dent, D. L., and Pons, L. J.: A world perspective on acid sulphate soils, *Geoderma*, 67, 263–276, [https://doi.org/10.1016/0016-7061\(95\)00013-E](https://doi.org/10.1016/0016-7061(95)00013-E), 1995.

485

Eggleton, J., and Thomas, K. V.: A review of factors affecting the release and bioavailability of contaminants during sediment disturbance events, *Environ. Int.*, 30, 973–980, <https://doi.org/10.1016/j.envint.2004.03.001>, 2004.

490 Eisma, D.: Flocculation and de-flocculation of suspended matter in estuaries, *Neth. J. Sea Res.*, 20, 183–199, [https://doi.org/10.1016/0077-7579\(86\)90041-4](https://doi.org/10.1016/0077-7579(86)90041-4), 1986.



- Fältmarsch, R. M., Åström, M. E., and Vuori, K.-M.: Environmental risks of metals mobilised from acid sulphate soils in Finland: a literature review, *Boreal Environ. Res.*, 13, 444–456, 2008.
- 495 Filzmoser, P., Hron, K., and Reimann, C.: Principal component analysis for compositional data with outliers, *Environmetrics*, 20, 621–632, <https://doi.org/10.1002/env.966>, 2009.
- Filzmoser, P., Hron, K., and Templ, M.: *Applied Compositional Data Analysis, with Worked Examples in R*, Springer, Cham, Switzerland, <https://doi.org/10.1007/978-3-319-96422-5>, 2018.
- 500 Folk, R. L., and Ward, W. C.: Brazos river bar: a study in the significance of grain size parameters, *J. Sediment. Petrol.*, 27, 3–26, <https://doi.org/10.1306/74D70646-2B21-11D7-8648000102C1865D>, 1957.
- Goñi, M. A., Teixeira, M. J., and Perkey, D. W.: Sources and distribution of organic matter in a river-dominated estuary (Winyah Bay, SC, USA), *Estuar. Coast. Shelf S.*, 57, 1023–1048, [https://doi.org/10.1016/S0272-7714\(03\)00008-8](https://doi.org/10.1016/S0272-7714(03)00008-8), 2003.
- 505 Gustafsson, Ö., Widerlund, A., Andersson, P. S., Ingri, J., Roos, P., and Ledin, A.: Colloid dynamics and transport of major elements through a boreal river-brackish bay mixing zone, *Mar. Chem.*, 71, 1–21, [https://doi.org/10.1016/S0304-4203\(00\)00035-9](https://doi.org/10.1016/S0304-4203(00)00035-9), 2000.
- 510 Hall, G. E. M.: Analytical perspective on trace element species of interest in exploration, *J. Geochem. Explor.*, 61, 1–19, [https://doi.org/10.1016/S0375-6742\(97\)00046-0](https://doi.org/10.1016/S0375-6742(97)00046-0), 1998.
- Häusler, K., Moros, M., Wacker, L., Hammerschmidt, L., Dellwig, O., Leipe, T., Kotilainen, A., and Arz, H. W.: Mid- to late Holocene environmental separation of the northern and central Baltic Sea basins in response to differential land uplift, *Boreas*, 46, 111–128, [10.1111/bor.12198](https://doi.org/10.1111/bor.12198), 2017.
- 515 Herzog, S. D., Gentile, L., Olsson, U., Persson, P., and Kritzberg, E. S.: Characterization of iron and organic carbon colloids in boreal rivers and their fate at high salinity, *J. Geophys. Res-Bioge.*, 125, e2019JG005517, <https://doi.org/10.1029/2019JG005517>, 2020.
- 520 Holby, O., and Evans, S.: The vertical distribution of Chernobyl-derived radionuclides in a Baltic Sea sediment, *J. Environ. Radioactiv.*, 33, 129–145, [https://doi.org/10.1016/0265-931X\(95\)00089-S](https://doi.org/10.1016/0265-931X(95)00089-S), 1996.
- 525 Hudd, R., and Kjellman, J.: Bad matching between hatching and acidification: a pitfall for the burbot, *Lota lota*, off the river Kyrönjoki, Baltic Sea, *Fish. Res.*, 55, 153–160, [https://doi.org/10.1016/S0165-7836\(01\)00303-4](https://doi.org/10.1016/S0165-7836(01)00303-4), 2002.
- Jilbert, T., Asmala, E., Schröder, C., Tiihonen, R., Myllykangas, J.-P., Virtasalo, J. J., Kotilainen, A., Peltola, P., Ekholm, P., and Hietanen, S.: Impacts of flocculation on the distribution and diagenesis of iron in boreal estuarine sediments, *Biogeosciences*, 15, 1243–1271, <https://doi.org/10.5194/bg-15-1243-2018>, 2018.
- 530



- 535 Jokinen, S. A., Virtasalo, J. J., Jilbert, T., Kaiser, J., Dellwig, O., Arz, H. W., Hänninen, J., Arppe, L., Collander, M., and Saارين, T.: A 1500-year multiproxy record of coastal hypoxia from the northern Baltic Sea indicates unprecedented deoxygenation over the 20th century, *Biogeosciences* 15, 3975–4001, <https://doi.org/10.5194/bg-15-3975-2018>, 2018.
- Jokinen, S. A., Jilbert, T., Tiiponen-Filppula, R., and Koho, K.: Terrestrial organic matter input drives sedimentary trace metal sequestration in a human-impacted boreal estuary, *Sci. Total Environ.*, 717, 137047, <https://doi.org/10.1016/j.scitotenv.2020.137047>, 2020.
- 540 Kakkuri, J.: Fennoscandian land uplift: past, present and future, in: *From the Earth's Core to Outer Space*, edited by: Haapala, I., *Lecture Notes in Earth System Sciences*, 137, 127–136, https://doi.org/10.1007/978-3-642-25550-2_8, 2012.
- Klaminder, J., Appleby, P., Crook, P., and Renberg, I.: Post-deposition diffusion of ^{137}Cs in lake sediment: implications for radiocaesium dating, *Sedimentology*, 59, 2259–2267, <https://doi.org/10.1111/j.1365-3091.2012.01343.x>, 2012.
- 545 Kotilainen, A. T., Kaskela, A. M., Bäck, S., and Leinikki, J.: Submarine De Geer moraines in the Kvarken Archipelago, the Baltic Sea, in: *Seafloor Geomorphology as Benthic Habitat: GeoHab Atlas of Seafloor Geomorphic Features and Benthic Habitats*, edited by: Harris, P. T., and Baker, E. D., Elsevier, Amsterdam, The Netherlands, 289–298, <https://doi.org/10.1016/B978-0-12-385140-6.00017-7>, 2012.
- 550 Lee, B. J., Fettweis, M., Toorman, E., and Molz, F. J.: Multimodality of a particle size distribution of cohesive suspended particulate matters in a coastal zone, *J. Geophys. Res.*, 117, C03014, <https://doi.org/10.1029/2011JC007552>, 2012.
- 555 Lehoux, A. P., Petersen, K., Leppänen, M. T., Snowball, I., and Olsen, M.: Status of contaminated marine sediments in four Nordic countries: assessments, regulations, and remediation approaches, *J. Soil. Sediment.*, in press, <https://doi.org/10.1007/s11368-020-02594-3>, 2020.
- Leivuori, M., and Niemistö, L.: Sedimentation of trace metals in the Gulf of Bothnia, *Chemosphere*, 31, 3839–3856, [https://doi.org/10.1016/0045-6535\(95\)00257-9](https://doi.org/10.1016/0045-6535(95)00257-9), 1995.
- 560 Long, E. R., Macdonald, D. D., Smith, S. L., and Calder, F. D.: Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments, *Environ. Manage.*, 19, 81–97, <https://doi.org/10.1007/BF02472006>, 1995.
- 565 Lundberg, C., Jakobsson, B.-M., and Bonsdorff, E.: The spreading of eutrophication in the eastern coast of the Gulf of Bothnia, northern Baltic Sea - an analysis in time and space, *Estuar. Coast. Shelf S.*, 82, 152–160, <https://doi.org/10.1016/j.ecss.2009.01.005>, 2009.
- 570 Mäkinen, J., and Saaranen, V.: Determination of post-glacial land uplift from the three precise levellings in Finland, *J. Geodesy*, 72, 516–529, <https://doi.org/10.1007/s001900050191>, 1998.
- Michael, P. S., Fitzpatrick, R. W., and Reid, R. J.: Effects of live wetland plant macrophytes on acidification, redox potential and sulphate content in acid sulphate soils, *Soil Use Manage.*, 33, 471–481, <https://doi.org/10.1111/sum.12362>, 2017.



- 575 Mikkelsen, O. A., Hill, P. S., and Milligan, T. G.: Single-grain, microfloc and macrofloc volume variations observed with a LISST-100 and a digital floc camera, *J. Sea Res.*, 55, 87–102, <https://doi.org/10.1016/j.seares.2005.09.003>, 2006.
- Nally, R. M., and Walsh, C. J.: Hierarchical partitioning public-domain software, *Biodivers. Conserv.*, 13, 659–660, <https://doi.org/10.1023/B:BIOC.0000009515.11717.0b>, 2004.
- 580 Nordmyr, L., Åström, M., and Peltola, P.: Metal pollution of estuarine sediments caused by leaching of acid sulphate soils, *Estuar. Coast. Shelf S.*, 76, 141–152, <https://doi.org/10.1016/j.ecss.2007.07.002>, 2008a.
- Nordmyr, L., Österholm, P., and Åström, M.: Estuarine behaviour of metal loads leached from coastal lowland acid sulphate soils, *Mar. Environ. Res.*, 66, 378–393, <https://doi.org/10.1016/j.marenvres.2008.06.001>, 2008b.
- 585 Nystrand, M. I., Österholm, P., Yu, C., and Åström, M.: Distribution and speciation of metals, phosphorus, sulfate and organic material in brackish estuary water affected by acid sulfate soils, *Appl. Geochem.*, 66, 264–274, <https://doi.org/10.1016/j.apgeochem.2016.01.003>, 2016.
- 590 Ojala, A. E. K., Palmu, J.-P., Åberg, A., Åberg, S., and Virkki, H.: Development of an ancient shoreline database to reconstruct the Litorina Sea maximum extension and the highest shoreline of the Baltic Sea basin in Finland, *Bull. Geol. Soc. Finl.*, 85, 127–144, <https://doi.org/10.17741/bgsf/85.2.002>, 2013.
- 595 Ojala, A. E. K., Luoto, T. P., and Virtasalo, J. J.: Establishing a high-resolution surface sediment chronology with multiple dating methods – testing ¹³⁷Cs determination with Nurmijärvi clastic-biogenic varves, *Quat. Geochronol.*, 37, 32–41, <https://doi.org/10.1016/j.quageo.2016.10.005>, 2017.
- Österholm, P., and Åström, M.: Spatial trends and losses of major and trace elements in agricultural acid sulphate soils distributed in the artificially drained Rintala area, W. Finland, *Appl. Geochem.*, 17, 1209–1218, [https://doi.org/10.1016/S0883-2927\(01\)00133-0](https://doi.org/10.1016/S0883-2927(01)00133-0), 2002.
- 600 Österholm, P., and Åström, M.: Meteorological impacts on the water quality in the Pajuluoma acid sulphate area, W. Finland, *Appl. Geochem.*, 23, 1594–1606, <https://doi.org/10.1016/j.apgeochem.2008.01.011>, 2008.
- 605 Paczkowska, J., Rowe, O., Schlüter, L., Legrand, C., Karlson, B., and Andersson, A.: Allochthonous matter: an important factor shaping the phytoplankton community in the Baltic Sea, *J. Plankton Res.*, 39, 23–34, <https://doi.org/10.1093/plankt/fbw081>, 2017.
- 610 Pavoni, E., Crosera, M., Petranich, E., Adami, G., Faganeli, J., and Covelli, S.: Partitioning and mixing behaviour of trace elements at the Isonzo/Soča River mouth (Gulf of Trieste, northern Adriatic Sea), *Mar. Chem.*, 223, 103800, <https://doi.org/10.1016/j.marchem.2020.103800>, 2020a.
- 615 Pavoni, E., Crosera, M., Petranich, E., Oliveri, P., Klun, K., Faganeli, J., Covelli, S., and Adami, G.: Trace elements in the estuarine systems of the Gulf of Trieste (northern Adriatic Sea): a chemometric approach to depict partitioning and behaviour of particulate, colloidal and truly dissolved fractions, *Chemosphere*, 252, 126517, <https://doi.org/10.1016/j.chemosphere.2020.126517>, 2020b.




- 620 Pirinen, P., Simola, H., Aalto, J., Kaukoranta, J.-P., Karlsson, P., and Ruuhela, R.: Climatological statistics of Finland 1981–2010, Finnish Meteorological Institute, Reports 2012:1, 83 pp., 2012.
- Regnier, P., and Wollast, R.: Distribution of trace metals in suspended matter of the Scheldt estuary, *Mar. Chem.*, 43, 3–19, [https://doi.org/10.1016/0304-4203\(93\)90212-7](https://doi.org/10.1016/0304-4203(93)90212-7), 1993.
- 625 Rhoads, D. C., and Boyer, L. F.: The effects of marine benthos on physical properties of sediments: a successional perspective, in: *Animal-Sediment Relations*, edited by: McCall, P. L., and Tevesz, M. J. S., *Topics in Geobiology*, 100, 3–52, https://doi.org/10.1007/978-1-4757-1317-6_1, 1982.
- Roos, M., and Åström, M.: Hydrochemistry of rivers in an acid sulphate soil hotspot area in western Finland, *Agr. Food Sci.*, 630 14, 24–33, <https://doi.org/10.2137/1459606054224075>, 2005.
- Ruosteenoja, K., Vihma, T., and Venäläinen, A.: Projected changes in European and North Atlantic seasonal wind climate derived from CMIP5 simulations, *J. Climate*, 32, 6467–6490, <https://doi.org/10.1175/JCLI-D-19-0023.1>, 2019.
- 635 Saarinen, T., Vuori, K.-M., Alasaarela, E., and Kløve, B.: Long-term trends and variation of acidity, COD_{Mn} and colour in coastal rivers of Western Finland in relation to climate and hydrology, *Sci. Total Environ.*, 408, 5019–5027, <https://doi.org/10.1016/j.scitotenv.2010.07.009>, 2010.
- Saarnisto, M., and Saarinen, T.: Deglaciation chronology of the Scandinavian Ice Sheet from the Lake Onega Basin to the 640 Salpausselkä end moraines, *Global Planet. Change*, 31, 387–405, [https://doi.org/10.1016/S0921-8181\(01\)00131-X](https://doi.org/10.1016/S0921-8181(01)00131-X), 2001.
- Sauramo, M.: The Quaternary geology of Finland, *Bulletin de la Commission Géologique de Finlande*, 86, 1–110, 1929.
- Sohlenius, G., and Öborn, I.: Geochemistry and partitioning of trace metals in acid sulphate soils in Sweden and Finland before 645 and after sulphide oxidation, *Geoderma*, 122, 167–175, <https://doi.org/10.1016/j.geoderma.2004.01.006>, 2004.
- Soil Survey Staff: *Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys*, 2nd edition, Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436, 1999.
- 650 Stolpe, B., and Hassellöv, M.: Changes in size distribution of fresh water nanoscale colloidal matter and associated elements on mixing with seawater, *Geochim. Cosmochim. Ac.*, 71, 3292–3301, <https://doi.org/10.1016/j.gca.2007.04.025>, 2007.
- Stroeven, A. P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B. W., Harbor, J. M., Jansen, J. D., Olsen, L., Caffee, M. W., Fink, D., Lundqvist, J., Rosqvist, G. C., Strömberg, B., and Jansson, K. N.: Deglaciation of 655 Fennoscandia, *Quaternary Sci. Rev.*, 147, 91–121, <https://doi.org/10.1016/j.quascirev.2015.09.016>, 2016.
- Sutela, T., and Vehanen, T.: The effects of acidity and aluminium leached from acid-sulphate soils on riverine fish assemblages, *Boreal Environ. Res.*, 22, 385–391, 2017.
- 660 Tenenhaus, M.: *La régression PLS: theorie et pratique*, Editions Technip, Paris, France, 1998.




- 665 Valikhani Samani, A. R., Karbassi, A. R., Fakhraee, M., Heidari, M., Vaezi, A. R., and Valikhani, Z.: Effect of dissolved organic carbon and salinity on flocculation process of heavy metals during mixing of the Navrud River water with Caspian Seawater, *Desalin. Water Treat.*, 55, 926–934, <https://doi.org/10.1080/19443994.2014.920730>, 2015.
- Vallius, H.: Applying sediment quality guidelines on soft sediments of the Gulf of Finland, Baltic Sea, *Mar. Pollut. Bull.*, 98, 314–319, <https://doi.org/10.1016/j.marpolbul.2015.06.036>, 2015.
- 670 Veijalainen, N., Korhonen, J., Vehviläinen, B., and Koivusalo, H.: Modelling and statistical analysis of catchment water balance and discharge in Finland in 1951–2099 using transient climate scenarios, *J. Water Clim. Change*, 3, 55–78, <https://doi.org/10.2166/wcc.2012.012>, 2012.
- Virtasalo, J. J., Kohonen, T., Vuorinen, I., and Huttula, T.: Sea bottom anoxia in the Archipelago Sea, northern Baltic Sea: implications for phosphorus remineralization at the sediment surface, *Mar. Geol.*, 224, 103–122, <https://doi.org/10.1016/j.margeo.2005.07.010>, 2005.
- 675 Virtasalo, J. J., Kotilainen, A. T., Räsänen, M. E., and Ojala, A. E. K.: Late-glacial and post-glacial deposition in a large, low relief, epicontinental basin: the northern Baltic Sea, *Sedimentology*, 54, 1323–1344, <https://doi.org/10.1111/j.1365-3091.2007.00883.x>, 2007.
- 680 Virtasalo, J. J., Åström, M. E., Österholm, P., and Kotilainen, A. T.: Multielement and grain size data of sediment samples from the Kvarken Archipelago, eastern Gulf of Bothnia, impacted by metal loading from acid sulphate soils, PANGAEA, doi, 2020. Submitted to be made available upon the publication of this paper.
- 685 Wallin, J., Karjalainen, A. K., Schultz, E., Järvistö, J., Leppänen, M., and Vuori, K.-M.: Weight-of-evidence approach in assessment of ecotoxicological risks of acid sulphate soils in the Baltic Sea river estuaries, *Sci. Total Environ.*, 508, 452–461, <https://doi.org/10.1016/j.scitotenv.2014.11.073>, 2015.
- 690 Wang, W., and Wang, W.-X.: Phase partitioning of trace metals in a contaminated estuary influenced by industrial effluent discharge, *Environ. Pollut.*, 214, 35–44, <https://doi.org/10.1016/j.envpol.2016.03.059>, 2016.
- Widerlund, A., and Ingri, J.: Redox cycling of iron and manganese in sediments of the Kalix River estuary, northern Sweden, *Aquat. Geochem.*, 2, 185–201, <https://doi.org/10.1007/BF00121631>, 1996.
- 695 Yli-Halla, M., Puustinen, M., and Koskiahio, J.: Area of cultivated acid sulfate soils in Finland. *Soil Use Manage.*, 15, 62–67, <https://doi.org/10.1111/j.1475-2743.1999.tb00065.x>, 1999.
- Ympäristöministeriö: Sedimenttien ruoppaus- ja läjitysohje, Ympäristöhallinnon ohjeita 1|2015, 72 pp., 2015.
- 700 Yu, C., Virtasalo, J. J., Karlsson, T., Peltola, P., Österholm, P., Burton, E. D., Arppe, L., Hogmalm, J. K., Ojala, A. E. K., and Åström, M. E.: Iron behavior in a northern estuary: large pools of non-sulfidized Fe(II) associated with organic matter, *Chem. Geol.*, 413, 73–85, <https://doi.org/10.1016/j.chemgeo.2015.08.013>, 2015.




705 Yu, C., Virtasalo, J. J., Österholm, P., Burton, E. D., Peltola, P., Ojala, A. E. K., Hogmalm, J. K., and Åström, M.E.: Manganese
accumulation and solid-phase speciation in a 3.5 m thick mud sequence from the estuary of an acidic and Mn-rich creek,
northern Baltic Sea, *Chem. Geol.*, 437, 56–66, <https://doi.org/10.1016/j.chemgeo.2016.05.016>, 2016.



710 **Figure 1. Maps of the Baltic Sea and study area. (a) Map of the Baltic Sea, and probability of acid sulphate soil occurrence in Finland. Black square indicates the location of the study area on the west coast of Finland. (b) Nautical chart of the study area in the Kvarken Archipelago. Red dots indicate the sediment coring sites of this study. Acid sulphate soil probability map: Geological Survey of Finland 2018. Nautical chart: S-57 Finnish Transport Agency 2017.**




715 **Figure 2.** Vertical distributions of ^{137}Cs , carbon, the share of terrestrial organic carbon, and metals typically leached from acid sulphate soils in representative sediment cores in the Kvarken Archipelago. The cores are arranged according to increasing distance from the Laihianjoki and Sulvanjoki Rivers (downward). The dark grey shading indicates the interval deposited between 1960 and 1985, and the light grey shading indicates the interval deposited before the year 1960 in each core. The intervals without shading were deposited in 1986 and later.



720

Figure 3. Biplot for robust compositional PCA of 54 elements in sediments deposited after the year 1960. Thirteen elements with the strongest loadings are shown by arrows.



725 **Figure 4.** (a) Correlation loading plot of the PLSR2 analysis for 54 predictor variables (blue, elements) and 13 response variables (orange, grain size classes in μm). (b) Hierarchical partitioning result plot, showing the independent contribution (%) of each grain size class to predicted Ni contents.





Figure 5. Map of the sea area off the town of Vaasa with median Al, Mn and Ni contents in core sections deposited before 1960 (lower panel) and in 1986 and after (upper panel) indicated. Red dots indicate median contents, whereas the contained dark red dots indicate mean median contents in the pre-1960 sections of four cores from the Gloppet area (MGGN-2018-29, MGGN-2018-30, MGGN-2018-31, MGGN-2018-32) in order to highlight the magnitude of enrichment. Note that core MGGN-2016-8 does not contain sediments deposited before 1960. The Laihianjoki and Sulvanjoki Rivers are outside the map area, in the southeast corner. Nautical chart: S-57 Finnish Transport Agency 2017.

730



735 **Figure 6.** Median Cd, Ni and Zn contents in sediment cores, arranged according to distance from the Laihianjoki and Sulvanjoki
 Rivers, with sediment quality guidelines indicated. Black bars show median contents in core sections deposited in 1986 and later,
 whereas the narrower grey bars show contents in core sections deposited before 1960. Whisker lines above the black bars indicate
 the upper quartile of contents. Note that core MGGN-2016-8 does not contain sediments deposited before 1960. White numbers in
 the Zn panel denote distance to the closest river mouth in km. Blue line indicates level 2 threshold in Finnish sediment dredging and
 740 dumping guidelines (Ympäristöministeriö, 2015). Green solid and dashed lines indicate North American effects range medium
 (ERM) and effects range low (ERL) thresholds, respectively (Long et al., 1995). Purple solid and dashed lines indicate Canadian
 probable effect level (PEL) and interim sediment quality guideline (ISQG) thresholds, respectively (Canadian Council of Ministers
 of the Environment, 2001).



745 **Table 1.** Sediment coring locations, water depths, coring dates, core lengths, and distances to the nearest river mouth.

Sediment core	Latitude	Longitude	Water depth	Coring date	Core length	Distance to
	ETRS-TM35FIN	ETRS-TM35FIN	(m)		(cm)	river (km)
MGGN-2016-8	63°05.040	21°32.182	6	4 Aug 2016	40	8.2
MGGN-2017-17	63°06.828	21°23.579	10	3 July 2017	38	14.2
750 MGGN-2017-18	63°05.781	21°27.339	8	3 July 2017	40	12.6
MGGN-2017-19	63°05.557	21°30.567	7	3 July 2017	40	10.1
MGGN-2017-20	63°06.640	21°32.186	5	3 July 2017	40	10.0
MGGN-2018-29	63°08.983	21°12.564	17	8 Aug 2018	30	26.3
MGGN-2018-30	63°08.023	21°14.327	14	8 Aug 2018	40	24.7
755 MGGN-2018-31	63°06.411	21°13.894	17.5	10 Aug 2018	40	23.8
MGGN-2018-32	63°06.675	21°16.433	15	10 Aug 2018	42	21.4



760 **Table 2.** Median metal and carbon contents in sediment samples, with interquartile ranges in brackets.

Core / Study area	n	Al	Cd	Co	Cu	La	Mn	Ni	Zn	C	Reference
<i>MGGN-2016-8</i>											
1986 and after	40	83650 (72750-87300)	1.17 (1.09-1.3)	70.1 (63.2-80.1)	58.5 (53.7-62.3)	187.6 (165.7-201.9)	1605 (1422-1893)	94.0 (80.0-105.8)	421.5 (372.5-459.3)	5.63 (5.16-5.87)	This study
<i>MGGN-2017-17</i>											
1986 and after	17	66100 (65400-66900)	0.81 (0.71-0.85)	52.8 (45.5-55.8)	31.0 (30.0-31.0)	71.7 (70.7-73.6)	1130 (1080-1640)	56.0 (54.0-57.0)	261.0 (257.0-269.0)	2.74 (2.57-3.05)	This study
1960–1985	15	66400 (65600-66650)	0.91 (0.75-1.08)	40.0 (37.1-43.1)	29.0 (26.0-31.0)	61.2 (55.0-63.5)	954 (921-1030)	50.0 (45.5-53.0)	235.0 (194.0-244.5)	2.18 (2.11-2.22)	
<i>MGGN-2017-18</i>											
1986 and after	22	69700 (67950-71900)	1.41 (1.23-1.52)	97.9 (83.5-100.0)	40.0 (39.0-41.0)	120.0 (111.5-124.8)	5505 (3823-7273)	102.5 (94.3-106.5)	421.0 (396.0-441.5)	4.06 (3.50-4.15)	This study
1960–1985	17	70700 (69200-71900)	1.48 (1.18-1.58)	63.4 (59.6-66.8)	40.0 (33.0-43.0)	98.2 (80.6-103.0)	2350 (1810-2870)	84.0 (77.0-88.0)	369.0 (318.0-384.0)	3.29 (2.89-3.36)	
Before 1960	1	66900	0.74	43.9	29.0	66.1	1700	57.0	253.0	2.63	
<i>MGGN-2017-19</i>											
1986 and after	13	75200 (74100-76100)	1.58 (1.48-1.64)	95.3 (86.9-105.0)	44.0 (43.0-46.0)	151.0 (139.0-156.0)	3510 (3390-5990)	115.0 (102.0-120.0)	470.0 (441.0-489.0)	4.53 (4.10-4.89)	This study
1960–1985	11	75000 (74200-75550)	1.57 (1.34-1.60)	66.0 (64.7-68.2)	42.0 (41.5-44.0)	126.0 (121.5-127.0)	3010 (2050-4405)	86.0 (83.5-88.0)	377.0 (362.5-381.5)	3.76 (3.75-3.92)	
Before 1960	16	74400 (73350-74900)	1.45 (1.41-1.64)	70.7 (63.5-74.6)	42.5 (40.0-44.3)	107.5 (100.8-117.5)	3305 (2333-5240)	91.5 (84.0-95.3)	389.5 (371.5-411.0)	3.64 (3.54-3.77)	
<i>MGGN-2017-20</i>											
1986 and after	14	73350 (72750-74700)	2.51 (2.39-2.71)	164.0 (153.7-180.5)	55.0 (57.0-62.0)	209.0 (192.5-219.5)	5600 (4965-7280)	184.0 (175.3-207.0)	707.0 (687.8-760.0)	5.92 (5.79-6.00)	This study
1960–1985	12	70950 (70500-75425)	2.17 (1.57-2.57)	121.5 (92.93-166.8)	45.5 (32.5-56.8)	128.0 (102.3-173.5)	2660 (1948-5990)	146.0 (111.0-183.3)	532.0 (426.5-662.5)	3.32 (2.88-4.38)	
Before 1960	14	69050 (68400-69875)	0.52 (0.45-0.71)	56.60 (51.23-62.78)	26.0 (25.0-27.0)	74.3 (71.3-76.8)	1630 (1478-1823)	71.5 (69.0-75.8)	282.0 (258.5-289.3)	2.41 (2.35-2.45)	
<i>MGGN-2018-29</i>											
1986 and after	16	70250 (68450-70625)	0.64 (0.58-0.68)	29.1 (26.4-31.7)	36.0 (33.8-37.3)	65.1 (63.6-66.3)	543 (524-592)	45.5 (42.5-47.3)	240.0 (216.3-247.0)	3.27 (3.21-3.40)	This study
1960–1985	13	68000 (65900-69600)	0.86 (0.71-1.05)	24.3 (20.4-26.5)	37.0 (36.0-37.0)	56.4 (54.9-59.3)	584 (564-607)	41.0 (40.0-41.0)	243.0 (214.0-250.0)	3.03 (2.76-3.10)	
Before 1960	1	71200	0.54	19.60	27.0	49.2	542	34.0	159.0	2.37	
<i>MGGN-2018-30</i>											
1986 and after	9	69500 (68500-70600)	0.77 (0.60-0.90)	40.2 (36.1-43.5)	33.0 (31.0-35.0)	68.2 (66.3-68.8)	718 (683-1250)	50.0 (48.0-52.0)	250.0 (228.0-258.0)	3.07 (3.04-3.39)	This study
1960–1985	7	69600 (69400-70500)	1.00 (0.80-1.34)	32.0 (31.2-37.1)	34.0 (30.5-34.5)	58.3 (55.3-62.8)	621 (620-641)	46.0 (43.5-48.0)	225.0 (213.5-266.5)	2.57 (2.54-2.71)	
Before 1960	24	68950 (68150-70000)	0.48 (0.35-0.54)	21.7 (21.1-23.7)	26.0 (25.0-27.0)	49.0 (48.6-49.8)	726 (700-757)	38.0 (37.0-39.0)	150.5 (140.0-163.8)	2.36 (2.27-2.37)	
<i>MGGN-2018-31</i>											
1986 and after	8	69800 (67600-71525)	0.60 (0.52-0.70)	28.9 (27.9-31.7)	32.0 (29.5-34.0)	64.5 (61.6-65.4)	695 (625-1205)	44.5 (42.0-48.5)	216.5 (199.8-231.8)	3.33 (3.15-3.41)	This study
1960–1985	7	70800 (70550-71600)	0.85 (0.65-0.98)	29.7 (28.3-33.3)	34.0 (32.5-37.0)	55.5 (55.2-58.2)	551 (547-573)	45.0 (44.0-45.5)	224.0 (208.5-248.5)	2.63 (2.61-2.76)	
Before 1960	25	70600 (69600-71600)	0.40 (0.27-0.55)	21.6 (19.5-23.1)	27.0 (27.0-29.0)	49.7 (48.9-50.9)	628 (614-642)	38.0 (36.0-39.0)	148.0 (135.0-163.0)	2.36 (2.27-2.40)	
<i>MGGN-2018-32</i>											
1986 and after	6	67350 (66025-68150)	0.64 (0.54-0.72)	32.7 (30.2-35.7)	27.0 (26.3-29.3)	59.5 (59.3-60.2)	909 (605-2085)	42.0 (40.3-43.8)	203.0 (193.5-211.8)	2.70 (2.58-2.99)	This study
1960–1985	6	69400 (68650-70825)	0.56 (0.54-0.75)	29.9 (25.2-31.9)	26.5 (26.0-28.5)	51.3 (49.6-54.2)	557 (535-567)	40.0 (37.0-43.8)	180.5 (159.8-199.8)	2.28 (2.21-2.40)	
Before 1960	30	69850 (69025-70925)	0.25 (0.21-0.32)	18.7 (17.6-19.6)	24.0 (24.0-25.0)	48.7 (47.6-49.4)	633 (607-652)	35.0 (34.0-37.0)	126.0 (120.5-133.3)	2.07 (1.92-2.14)	
<i>Total</i>											
1986 and after	145	71100 (67900-75200)	1.10 (0.75-1.40)	68.5 (42.6-93.0)	41.0 (33.0-55.0)	125.0 (67.8-175.8)	1800 (1120-4090)	88.0 (51.0-107.0)	389.0 (254.0-454.0)	4.10 (3.26-5.39)	This study



825	1960–1985	88	70250 (67925-71950)	1.08 (0.83-1.52)	43.1 (30.8-66.2)	35.5 (30.0-41.3)	64.4 (56.3-107.0)	1030 (595-2355)	54.5 (43.7-86.0)	260.0 (222.5-374.8)	2.79 (2.48-3.36)	
	Before 1960	117	69900 (68600-71400)	0.44 (0.27-0.62)	21.7 (19.5-41.0)	26.0 (24.0-28.0)	49.7 (48.6-68.5)	669 (631-1270)	38.0 (36.0-58.0)	148.0 (131.0-244.0)	2.34 (2.11-2.43)	
	<i>Grand total</i>	350	70300 (68300-72700)	0.86 (0.53-1.32)	45.3 (24.3-72.8)	33.0 (27.0-43.0)	68.4 (51.8-124.8)	1195 (644-2490)	55.0 (40.0-91.0)	258.0 (176.2-393.5)	3.04 (2.36-3.93)	This study
830	<i>Laihianjoki River Estuary</i> (2015) Surface sediment, microwave HNO ₃ digestion	1	59900	0.92	74.3	63.1	n.d.	788	130.5	461	n.d.	Wallin et al.
835	<i>Vöyrinjoki River Estuary</i> 2008b Site B, 1960 and after Four acid digestion	13	86400 (82900-87600)	1.54 (1.32-1.76)	112.1 (106.6-128.6)	59.9 (58.6-64.0)	187 (186-198)	9013 (6854-10923)	83.9 (80.1-98.5)	529.1 (493.6-592.0)	n.d.	Nordmyr et al.
	<i>Bothnian Bay</i> Surface sediment, hydrofluoric acid digestion, mean content	5	n.d.	0.8 ± 0.3	n.d.	46 ± 15	n.d.	8500 ± 5300	n.d.	216 ± 92	4.0 ± 1.2	Leivoori and Niemistö, 1995
840	<i>Bothnian Sea</i> Surface sediment, hydrofluoric acid digestion, mean content	5	n.d.	0.4 ± 0.2	n.d.	30 ± 11	n.d.	3000 ± 1600	n.d.	173 ± 56	2.3 ± 1.0	Leivoori and Niemistö, 1995

Metal contents are in mg/kg dry weight. Carbon content is in % dry weight. N.d. denotes not determined.