

Eukaryotic community composition in the sea surface microlayer across an east-west transect in the Mediterranean Sea

Birthe Zäncker^{1,2}, Michael Cunliffe^{2,3}, Anja Engel¹

¹GEOMAR, Helmholtz Centre for Ocean Research Kiel, 24105 Kiel, Germany

5 ²Marine Biological Association of the UK, Plymouth, PL1 2PB, United Kingdom

³School of Biological and Marine Sciences, University of Plymouth, PL4 8AA, United Kingdom

Correspondence to: Anja Engel (aengel@geomar.de)

Abstract. The sea surface microlayer (SML) represents the boundary layer at the air-sea interface. Microbial eukaryotes in the SML potentially influence air-sea gas exchange directly by taking up and producing gases, and indirectly by excreting and degrading organic matter, which may modify the viscoelastic properties of the SML. However, little is known about the distribution of microbial eukaryotes in the SML. We studied the composition of the microbial community, transparent exopolymer particles and polysaccharides in the SML during the PEACETIME cruise along a west-east transect in the Mediterranean Sea, covering the western basin, Tyrrhenian Sea and Ionian Sea. At the stations located in the Ionian Sea, fungi – likely of continental origin and delivered by atmospheric deposition - were found in high relative abundances making up a significant proportion of the sequences recovered. Concomitantly, bacterial and picophytoplankton counts decreased from west to east, while transparent exopolymer particle (TEP) abundance and total carbohydrate (TCHO) concentrations remained constant in all basins. Our results suggest that the presence of substrates for fungi, such as *Cladosporium* known to take up phytoplankton-derived polysaccharides, in combination with decreased substrate competition by bacteria might favour fungal dominance in the neuston of the Ionian Sea and other low nutrient low chlorophyll (LNLC) regions.

20 **1 Introduction**

The sea surface microlayer (SML) constitutes a 1 to 100 μm thick boundary layer between the ocean and the atmosphere (Cunliffe and Murrell, 2010; Liss and Duce, 2005; Zhang et al., 2003) with distinct physical and chemical properties compared to the underlying water (Cunliffe et al., 2013; Zhang et al., 2003). The SML potentially has a substantial influence on air-sea exchange processes, such as gas transfer and sea spray aerosol formation (Cunliffe et al., 2013; Engel et al., 2017; Freney et al., 2020; Sellegri et al., in prep.).

The microbial food web plays a crucial role in ocean biogeochemistry and has been vastly studied. Despite the fact that microbes in the SML can directly and indirectly influence air-sea gas exchange, few studies have looked at the microbial community composition in the SML, and have mainly focussed on bacteria (Agogu e et al., 2005; Joux et al., 2006; Obernosterer et al., 2008) with little attention to microbial eukaryotes (Taylor and Cunliffe, 2014). While phytoplankton throughout the water column play an important role in the ocean as primary producers, phytonuston in the SML (Apts, 1989;

Hardy and Apts, 1984; Naumann, 1917), might have an additional crucial role by impacting air-sea gas exchange (Ploug, 2008; Upstill-Goddard et al., 2003). Early microscopic observations of the SML reported mostly diatoms, dinoflagellates and cyanobacteria (Hardy et al., 1988). More recent studies using 18S rRNA gene sequencing found a decreased protist diversity in the SML compared to underlying water with chrysophytes and diatoms enriched in the SML (Cunliffe and Murrell, 2010; 35 Taylor and Cunliffe, 2014).

Not only phytoneuston, but also zooneuston, bacterioneuston and myconeuston might influence air-sea gas exchange processes by either parasitizing phytoneuston and thus impacting the primary productivity, or by degrading organic matter available in the SML and releasing CO₂. While some studies have explored bacterioneuston diversity in the Mediterranean Sea (Agogue et al., 2005; Joux et al., 2006), fungi have not yet been characterized in the SML in this region. Fungi are however abundant 40 in marine environments (Gladfelter et al., 2019; Grossart et al., 2019; Hassett et al., 2019), living a saprotrophic or parasitic lifestyle and have been found in the Mediterranean Sea before (Garzoli et al., 2015; Gnani et al., 2017) and in the myconeuston studied at other locations (Taylor and Cunliffe, 2014).

Phytoplankton and phytoneuston can release precursors such as carbohydrates which can aggregate and form gelatinous particles such as transparent exopolymer particles (TEP) (Chin et al., 1998; Engel et al., 2004; Verdugo et al., 2004). TEP 45 contain mainly polysaccharides (Mopper et al., 1995; Passow, 2002), occur ubiquitously in the ocean (Alldredge et al., 1993; Passow, 2002), and are an important structural component of the SML (Wurl and Holmes, 2008). Due to their stickiness TEP can aggregate with other denser particles (Azetsu-Scott and Passow, 2004; Engel, 2000; Passow and Alldredge, 1995), eventually sink out of the euphotic layer into the deep ocean leading to carbon export (Engel et al., 2004). However, the rate of TEP-related carbon export does not only depend on TEP production by phytoplankton, but also on their microbial 50 degradation.

Few studies have looked at spatial distribution of the microbial eukaryote communities in the SML and possible environmental drivers of community composition in the open Mediterranean Sea, a characteristic low nutrient low chlorophyll (LNLC) region (Durrieu de Madron et al., 2011). The anti-estuarine circulations at the Strait of Gibraltar and the Straits of Sicily, transport low-nutrient surface waters into the basins, and deeper waters out of the basins, resulting in oligotrophic conditions and ultra- 55 oligotrophic conditions in the western and eastern Mediterranean basin, respectively (Krom et al., 2004; Mermex Group et al., 2011; Pujo-Pay et al., 2011; Tanhua et al., 2013). The present study focuses on TEP as important structural components of the SML and their precursors, carbohydrates, as well as microbial eukaryotes distribution, focusing on the myconeuston community composition in the SML using samples collected during the PEACETIME cruise in the Mediterranean Sea during May and June 2017.

60 **2 Material and methods**

2.1 Sampling

Samples were collected onboard the RV *Pourquoi pas?* from the 10th May to the 11th June 2017. Water from the SML and the underlying water (ULW; 20 cm below the SML) was collected at 12 stations from 2.9°E to 19.8°E and 35.5°N to 42.0°N (Fig. 1). SML samples were collected from a zodiac using a glass plate sampler (Cunliffe and Wurl, 2014; Harvey, 1966). The dimensions of the silicate glass plate (50 x 26 cm) resulted in an effective sampling surface area of 2,600 cm² considering both sides. To avoid contamination during sampling, the zodiac was positioned upwind and in front of the research vessel. The glass plate was immersed and withdrawn perpendicular to the sea surface. With a Teflon wiper, SML samples were collected in acid cleaned and rinsed bottles (Cunliffe and Wurl, 2014). Approximately 1.5 L of SML sample was collected in the course of 1 h. Sampling times are listed in table 1. All sampling equipment was acid-cleaned (10 % HCl), rinsed with Milli-Q and copiously rinsed with seawater from the respective depth once the sampling site was reached. The ULW samples were collected simultaneously with two acid-cleaned and MilliQ rinsed glass bottles by immersing the closed bottles and opening them at app. 20 cm.

2.2 Gel particle determination

The abundance and area of TEP was measured microscopically (Engel, 2009). The sample volume (10-30 ml) was chosen according to the prevailing TEP concentrations. Samples were filtered onto 0.4 µm Nucleopore membranes (Whatman) and stained with 1 ml Alcian Blue solution (0.2 g l⁻¹ w/v) for 3 seconds. Filters were mounted on Cytoclear[®] slides and stored at -20°C until analysis. Two filters per sample with 30 images each were analyzed using a Zeiss Axio Scope.A1 (Zeiss) equipped with a Zeiss AxioCam MRC. The pictures with a resolution of 1388 x 1040 pixels were saved using AxioVision LE64 Rel. 4.8 (Zeiss). All particles larger than 0.2 µm² were analyzed. ImageJ was subsequently used for image analysis (Schneider et al., 2012). A filter prepared with 10 ml MilliQ water served as a blank.

2.3 Bacterioplankton and bacterioneuston abundance

Bacterial cell numbers were determined from 2 ml samples fixed with 100 µl glutaraldehyde (GDA, 1 % final concentration). Samples were stored at -20°C and stained with SYBR Green I (Molecular Probes) to determine abundance using a Becton & Dickinson FACScalibur flow cytometer equipped with a 488 nm laser. Bacterial cells were detected by the unique signature in a plot of side scatter (SSC) vs. green fluorescence (FL1). Yellow-green latex beads (Polysciences, 0.5 µm) were used as internal standards.

2.4 Picophytoplankton and picophytoneuston abundance

Picophytoplankton and picophytoneuston cell numbers were determined from 2 ml samples fixed and stored as for bacterial abundances. Samples were filtered through a 50 µm filter and analyzed with a flow cytometer (similar as in section 2.3).

90 Enumeration of cells was conducted using a high flow rate (app. 39-41 $\mu\text{l min}^{-1}$). The forward or right-angle light scatter (FALS, RALS) as well as the phycoerythrin and chl *a* related fluorescent signal was used to distinguish the cells. Cell counts were analyzed using the CellQuest Pro-Software (BD Biosciences). The method used here (fixative addition + slow freezing) follows recommendations by Lepesteur et al. (1993).

2.5 Total combined carbohydrates

95 Samples (20 ml) for total hydrolysable carbohydrates (TCHO) > 1 kDa were filled into precombusted glass vials (8h, 500°C) and stored at -20°C. In the home lab, TCHO analysis was carried out using high performance anion exchange chromatography with pulsed amperometric detection (HPAEC-PAD) on a Dionex ICS 3000 ion chromatography system (Engel and Händel 2011). Prior to analysis, samples were desalinated by membrane dialysis (1 kDa MWCO, Spectra Por) at 1°C for 5h and hydrolyzed for 20 h at 100°C in HCl (0.8 M final concentration) with subsequent neutralization using acid evaporation (N₂,
100 for 5 h at 50°C). Two replicates were analysed for each sample.

2.6 DNA extraction and eukaryote 18S rRNA gene sequencing

Water samples for sequencing (400 ml each) were passed through a 100 μm pore size mesh in order to remove meta-zooplankton that could dominate the 18S sequences and subsequently filtered onto a Durapore membrane (Millipore, 47 mm, 0.2 μm) and immediately stored at -80°C. In order to improve cell accessibility for the DNA extraction, filters in cryogenic
105 tubes were immersed in liquid nitrogen and the filter was crushed with a pestle. DNA was extracted according to a modified protocol from Zhou et al. (1996) by Wietz et al. (2015). The protocol included bead-beating, phenol-chloroform-isoamyl alcohol purification, isopropanol precipitation and ethanol washing. An additional protein-removal step by salting was used to avoid protein contamination.

Library preparation and sequencing was conducted at the Integrated Microbiome Resource at Dalhousie University, Halifax,
110 Canada and is described in detail elsewhere (Comeau et al., 2017). Samples were PCR-amplified in two dilutions (1:1 and 1:10) using the 18S rRNA gene primers E572F and E1009R (Comeau et al., 2011). Prior to pooling, samples were cleaned up and normalized using the Invitrogen SequalPrep 96-well Plate kit (Thermo Fisher Scientific). Sequencing was conducted according to Comeau et al. (2017) on an Illumina MiSeq using 300+300 bp paired-end V3 chemistry.

Sequences were processed using the DADA2 pipeline (Callahan et al., 2016) and sequences shorter than 400 bp, longer than
115 444 bp, with more than 8 homopolymers or any ambiguous bases were discarded. Sequences were aligned with the 18S rRNA gene sequences of the SILVA 132 alignment database (Quast et al., 2013). Subsequently, sequences that aligned outside of most of the dataset and chimeras were removed. Sequences were classified using the SILVA 132 database (Quast et al., 2013) and deposited at the European Nucleotide Archive (ENA accession number PRJEB23731). Sequences were not subsampled and sequence numbers per sample ranged from 1063 (S8 SML) to 43,027 (S5 SML). However, for principal component
120 analysis (PCA), all samples were subsampled down to 1063 sequences.

2.7 Statistical analyses

Statistical analyses and maps were produced using R (R Core Team, 2014) and bathymetry from NOAA (National Oceanic and Atmospheric Administration). The enrichment factor (EF) was used to compare the concentration of parameter A in the SML ($[A]_{SML}$) to the concentration in the ULW ($[A]_{ULW}$) and was calculated as follows (Eq. 1; World Health Organization, 125 1995):

$$EF = \frac{[A]_{SML}}{[A]_{ULW}} \quad (1)$$

An $EF > 1$ indicates enrichment, $EF < 1$ indicates depletion and $EF = 1$ indicates no difference between the SML and the ULW. The significance of differences between the SML and ULW and between the basins of eukaryote sequences and biogeochemical parameters were tested using the Kruskal-Wallis test and PERMANOVA. Correlations were calculated using 130 Spearman's rank correlation.

2.8 Data obtained from the ship

Wind speed, surface water salinity and temperature were obtained at 5 m depth from the RV *Pourquoi pas?* system. Radiation measurements were obtained with a Li-Cor Radiation Sensor (Li-200SZ) at wavelengths of 400 to 1100 nm. All parameters were measured every 5 min during sampling on the zodiac and averaged over the sampling period for statistical analyses (Table 135 1).

3 Results

3.1 Microbial eukaryote community composition in the SML and ULW

The eukaryotic communities in the SML and the ULW were similar (ANOSIM, $p=0.039$, $R=0.1002$). However, differences were detected in the eukaryotic community composition (Fig. 2) of the basins sampled (western Mediterranean, Tyrrhenian 140 Sea and Ionian Sea). ANOSIM showed that the differences in the eukaryotic community composition were slightly larger across basins than between SML and ULW ($p=0.0025$, $R=0.2263$). However, the overall diversity and evenness (based on shannon and pielou indices) were not significantly different between basins (Fig. S1).

16 orders were found in relative abundances over 5 % of the total eukaryotic community in one or more of all 12 stations (Fig. 3). The communities in the SML and ULW at most stations were similar, with Dinophyceae and Syndiniales (Dinoflagellata), 145 and an unidentified Eukaryote class dominating the eukaryotic community. Zooneuston were found in most of the SML samples, but rarely ($n=2$) in the ULW samples. Zooneuston were comprised of Ploimida (Rotifer), Maxillopoda (Cyclopoida and Calanoida) and Scyphozoa (Semaestomeae).

Myconeuston and mycoplankton were found in high relative abundances in three ULW samples and in the corresponding SML samples (Stations 7, 8 and ION_2) in the Ionian Sea. In the ULW of station 7, fungi made up more than half (54 %) of the total 150 number of retrieved sequences. The vast majority of fungal amplicon sequence variants (ASVs) (64 out of 69) belonged to

Ascomycota and Mucoromycota with the remaining five belonging to the Chytridiomycota (n=3), Basidiomycota and Neocallimastigomycota. All fungal ASVs that were recovered throughout the cruise and their relative abundance are shown in Figure 4. While fungal ASVs made up a significant amount of sequences in the Ionian Sea (stations to the right of fig. 4), they were barely detectable at the other stations (p=0.014 for differences in fungal ASVs between basins tested with PERMANOVA).

3.2 Concentrations and SML enrichments of microorganisms and organic matter

Bacterial numbers did not show any significant differences between layers (Fig. 5A). In the SML, bacterial abundances ranged from 2.0×10^5 to 1.0×10^6 cells ml⁻¹ with an average of $5.2 \times 10^5 \pm 2.3 \times 10^5$ cells ml⁻¹. In the ULW, bacterial numbers were on average $4.6 \times 10^5 \pm 1.5 \times 10^5$ cells ml⁻¹ (range of 2.2×10^5 to 6.9×10^5 cells ml⁻¹) (Fig. 5).

160 Picophytoneuston (0.2 – 20 μm size range) abundance was on average $3.3 \times 10^3 \pm 1.9 \times 10^3$ cells ml⁻¹ in the SML and picophytoplankton abundance was on average $2.3 \times 10^3 \pm 1.7 \times 10^3$ cells ml⁻¹ in the ULW (range of 1.4×10^3 to 8.5×10^3 cells ml⁻¹ in the SML, 9.5×10^2 to 7.1×10^3 cells ml⁻¹ in the ULW). Overall, cell counts determined by flow cytometry were significantly higher in the SML than in the ULW (p=0.002, n=12; Fig. 5B).

TEP concentration averaged $1.4 \times 10^7 \pm 9.7 \times 10^6$ particles l⁻¹ (ranging between 3.6×10^6 and 3.7×10^7 TEP l⁻¹) in the SML. In 165 the ULW, the average TEP concentration was $3.6 \times 10^6 \pm 2.1 \times 10^6$ particles l⁻¹ (ranging between 6.8×10^5 and 7.5×10^6 TEP l⁻¹) in the ULW. TEP area in the SML was on average $9.7 \times 10^7 \pm 1.2 \times 10^8$ mm² l⁻¹ (1.5×10^7 to 4.5×10^8 mm² l⁻¹). TEP area was lower in the ULW with an average of $2.3 \times 10^7 \pm 1.1 \times 10^7$ (2.9×10^6 to 3.9×10^7 mm² l⁻¹). Both TEP abundance and area were significantly enriched in the SML (Fig. 5; p=0.01 and p=0.007, respectively). While irradiation, water temperature and salinity did not correlate with TEP abundance or area, wind speed did have a significant negative correlation with TEP 170 abundance in the SML ($R^2 = -0.73$) and TEP area in the SML ($R^2 = -0.75$) and the enrichment factor for TEP area ($R^2 = -0.63$). TCHO concentrations were similar between the SML and ULW (Fig. 5E), with no significant differences between depths (778 ± 294 nM (range 562 to 1684 nM) in the SML and 605 ± 97 nM (range 525 to 885 nM) in the ULW).

4 Discussion

4.1 Eukaryotic diversity in the surface of the Mediterranean Sea

175 The eukaryotic community composition between the SML and the ULW differed only slightly, with higher horizontal heterogeneity and significant differences between the communities of the Western, Tyrrhenian and Ionian basins. The shannon diversity did not differ significantly between depths or basins, however, a slight decrease of species richness from west to east could be observed (Fig. S1), possibly related to the transition from oligotrophic to ultra oligotrophic conditions given the more pronounced water exchange with the Atlantic in the western basin (Reddaway and Bigg, 1996).

180 No diatoms were present at high relative abundances in our samples. In seasonal studies in the Mediterranean Sea, diatom contribution can be significant during blooms in March and April, but later in the year, as the water column stratifies, , their

contribution decreases (Marty et al., 2002). Even though diatoms most likely were not dominant in the samples, the extremely low abundance (< 1%) of diatoms in the samples might also indicate a bias of the primers used or removal of larger cells and aggregates during the pre-filtration step. Another potential bias is the dominance of dinoflagellate genera (Fig. 3).
185 Dinoflagellates have a large number of 18S rRNA gene copies in comparison to other phytoplankton groups, and therefore their abundance in 18S rRNA gene sequencing is often overestimated (Godhe et al., 2008; Guo et al., 2016).
Previous studies suggested various factors that potentially drive the phytoplankton community composition. In addition to buoyancy of cells, high levels of UV-radiation could potentially cause damage by photoinhibition. While dinoflagellates, one of the dominating phytoplankton groups, can produce photoprotective compounds, including mycosporine-like amino acids (MAAs) (Carreto et al., 1990; Häder et al., 2007), they can still be inhibited by high UV radiation (Ekelund, 1991).
190 In the present study, inhibition by UV radiation is not indicated in the data since phytoplankton was enriched in the SML despite high radiation values (e.g. stations 4 and 7; Table 1). At the same time, TEP were significantly enriched during this sampling campaign while the phytoplankton community did not show significant differences. Previous studies suggested that TEP can protect phytoplankton and bacteria from UV radiation (Elasri and Miller, 1999; Ortega-Retuerta et al., 2009). Process oriented
195 studies would be needed to determine whether TEP production was higher in the SML due to UV protection of phytoplankton or whether TEP formation rates were higher in the SML due to wind and wave shear at the surface (Carlson, 1993; Cunliffe et al., 2013).

4.2 Fungi in the Ionian Sea

Fungi were prevalent in the Ionian Sea (more than half of the sequences retrieved at one station), while scarce in the western
200 basin and the Tyrrhenian Sea (Fig. 3 and 4). Most of the fungal ASVs present in the Ionian Sea belonged to Ascomycota and Mucoromycota. The question arises as to what drives the higher fungal relative abundances in this region of the Mediterranean Sea. While fungi, like dinoflagellates and other eukaryotic groups, can have varying amounts of 18S rDNA gene copy numbers, the patchy distribution of fungi found in this study makes a consistent bias unlikely. Marine fungi can live a saprotrophic lifestyle, degrading and recycling high molecular weight organic matter (Christmas and Cunliffe, 2020; Cunliffe et al., 2017)
205 and potentially competing with functionally similar bacteria. Some marine fungi are also phytoplankton parasites, potentially altering phytoplankton community composition through selective parasitism (Amend et al., 2019; Grossart et al., 2019).. At present, we have a very limited understanding of diversity and functional role of fungi in the SML (myconeuston). One previous study of the coastal myconeuston in the Western English Channel off Plymouth (UK) showed that the SML was dominated by both Ascomycota and Basidiomycota (Taylor and Cunliffe, 2014), while our results support the dominance of
210 Ascomycota in the SML.

So far, not many studies have looked at fungi in LNLC regions. A global comparison of fungal distribution (Hassett et al., 2020) has found that fungal diversity determined by amplicon sequencing varies between different oceanic regions with *Exophiala*, belonging to Ascomycota, dominating the Ligurian Sea and an unclassified Ascomycota being the most abundant taxon in other regions.

215 ASVs identified as Solanales (Nicotiana) had quite high relative abundances in the easternmost stations. Since Solanales are land plants, presence of their DNA could suggest a terrestrial influence on the Ionian Sea, linked to wet or dry deposition that occurred before and/or during our sampling period in this basin. This is also corroborated by air mass trajectory backtracking using the HYSPLIT model (Fig. S3) which showed that aerosols were most likely of continental origin (Fu et al., in prep). This is also confirmed by atmospheric measurements indicating that chemical composition of dry and wet depositions were
220 influenced by Eastern European air masses (Desboeufs et al., in prep.). Station FAST_2 in the western basin was highly influenced by dust input in the area (Guieu et al., 2020; Tovar-Sánchez et al., 2020). This coincided not only with a strong increase in TEP abundance in the SML, but also with a distinct increase in the relative abundance of unidentified dinoflagellates in the SML (Fig. 3). The details of the impact of dust input on the organic matter and microbial community composition in the SML and ULW are discussed elsewhere (Engel et al., in prep). However, no fungi were found at station FAST_2 neither in
225 the SML nor in the ULW (Fig. 4), showing that dust input does not necessarily deposit fungi to the surface ocean, this potentially also holds true for the Ionian Sea.. In addition to atmospheric inputs, riverine inputs can also influence the Mediterranean Sea (Martin et al., 1989). However, the Ionian Sea itself does not experience vast riverine input and riverine influence is less pronounced in the open sea, making riverine sources of mycophyta unlikely. Ascomycota and Mucoromycota have been recovered from a variety of marine environments (Bovio, 2019; Grossart et al., 2019; Hassett et al., 2019), implying
230 that they also might be thriving in the SML of the Mediterranean Sea rather than originating solely from terrestrial sources. Overall, the most abundant fungal ASV in the Ionian Sea (ASV 8) was identified as belonging to genus *Cladosporium* which has been found in marine environments before (Cunliffe et al., 2017). Another explanation for the high relative abundance of fungi in the Ionian Sea might be that they are more adapted to dealing with the low nutrient conditions found in the more eastern basin of the Mediterranean Sea.

235 Bacterial and microalgal numbers determined by flow cytometry decreased significantly from west to east, with bacteria showing the largest decline (Tovar-Sánchez et al., 2020). While total microalgal abundances determined by flow cytometry were low in the SML and in the ULW, they were comparable to other studies on phytoplankton abundance in the SML of the Mediterranean Sea (Joux et al., 2006). The microalgal numbers between 5 and 200 m water depth (data not shown) were higher than at the air-sea interface. Even though overall bacterial numbers decrease, further molecular analyses would be needed to
240 determine if the bacterial community is changing from west to east and if certain bacterial taxa can benefit from the ultra oligotrophic conditions. In contrast to bacteria and phytoplankton, spatial trends in TCHO and TEP, as well as DOC in the SML and DOC and POC in the ULW did not show significant differences between the Ionian Sea and the other basins (Freney et al., 2020; Trueblood et al., 2020). TEP are often enriched in the SML (Engel and Galgani, 2016; Jennings et al., 2017; Wurl et al., 2009; Wurl and Holmes, 2008), in particular over oligotrophic regions (Jennings et al., 2017; Zäncker et al., 2017). This
245 is in good accordance with trends observed in the present study (Durrieu de Madron et al., 2011; Fogg, 1995; Wikner and Hagstrom, 1988). Wind speed correlated negatively with TEP abundance and area in the SML, showing that wind can negatively affect TEP concentrations at the air-sea interface as has been previously suggested (Sun et al., 2018).

Since exchange of water with the Atlantic is most pronounced in the western basin and an anti-estuarine circulation prevails in the Mediterranean Sea, nutrient limitation increases eastwards. TEP production has been shown to be independent of stoichiometric ratios in the surrounding water in a previous study (Corzo et al., 2000). Since in the SML, light limitation rarely occurs and TEP can potentially protect phytoplankton from light damage (Elasri and Miller, 1999; Ortega-Retuerta et al., 2009), phytoplankton might still photosynthesize and excrete carbohydrates that assemble to TEP. This would not only explain the lack of difference in TEP abundances between basins, but also TCHO concentrations. TCHO can also be produced by cell lysis (due to nutrient depletion).

TCHO and TEP could also provide available substrate and microhabitats for marine fungi with reduced competition by bacteria in the Ionian Sea. *Malassezia* and *Cladosporium* have been shown to assimilate carbon derived from TEP-associated algal polysaccharides in the English Channel (Cunliffe et al., 2017), highlighting that *Cladosporium* and other fungi might be able to profit from the decreased bacterial competition in the Ionian Sea. Further, previous studies have shown that the Eastern Mediterranean Sea contains higher concentrations of organic pollutants (Berrojalbiz et al., 2011a, 2011b) while a *Cladosporium* strain has the capacity to degrade polycyclic aromatic hydrocarbons (Birolli et al., 2018), highlighting another potential substrate for the fungi detected in the Ionian Sea.

5 Conclusions

The present study shows that bacteria and picophytoplankton numbers decrease from west to east in the Mediterranean Sea, in contrast organic matter such as microgels and TCHO are still prevalent in surface waters. Our findings from the Ionian Sea suggest that accumulation of organic substrates in the surface under oligotrophic conditions may favour certain taxa such as fungi which can benefit from decreased competition by bacteria. LNLC regions, where phytoplankton and bacterial counts are typically low, but TEP enrichment is high in the SML, might be a specific ecosystem where fungi are able to thrive and to control organic matter degradation.

Acknowledgements

We would like to thank the chief scientist, Cécile Guieu and Karine Desboeufs, of the PEACETIME cruise on the RV *Pourquoi pas?*. We would also like to thank the captain and crew of the *Pourquoi pas?* for technical assistance in the field. This work is a contribution of the PEACETIME project (<http://peacetime-project.org>), a joint initiative of the MERMEX and ChArMEX components supported by CNRS-INSU, IFREMER, CEA, and Météo-France as part of the programme MISTRALS coordinated by INSU (doi: 10.17600/17000300).

We thank Jon Roa for his help in analyzing the total combined carbohydrates and Tania Klüver for analyzing the flow cytometry cell counts. We would also like to thank ISOS (Kiel, Germany), for funding part of this work with a PhD-Miniproposal Grant.

Data availability

All biogeochemical data will be made available at the French INSU/CNRS LEFE CYBER database (data manager, webmaster: Catherine Schmechtig). All sequence data is available at the European Nucleotide Archive (ENA accession number PRJEB23731).

Author contributions

BZ, MC and AE wrote the paper and contributed to the data analysis. BZ participated in the sample treatment.

Competing interests

The authors declare that they have no conflict of interest

References

- Agogu , H., Casamayor, E. O., Bourrain, M., Obernosterer, I., Joux, F., Herndl, G. J. and Lebaron, P.: A survey on bacteria inhabiting the sea surface microlayer of coastal ecosystems, *FEMS Microbiol. Ecol.*, 54(2), 269–280, doi:10.1016/j.femsec.2005.04.002, 2005.
- Allredge, A. L., Passow, U. and Logan, B. E.: The abundance and significance of a class of large, transparent organic particles in the ocean, *Deep. Res. I*, 40(6), 1131–1140, 1993.
- Amend, A., Burgaud, G., Cunliffe, M., Edgcomb, V. P., Ettinger, C. L., Gutierrez, M. H., Heitman, J., Hom, E. F. Y., Ianiri, G., Jones, A. C., Kagami, M., Picard, K. T., Quandt, C. A., Raghukumar, S., Riquelme, M., Stajich, J., Vargas-mu iz, J., Walker, A. K., Yarden, O. and Gladfelter, A. S.: Fungi in the Marine Environment : Open Questions and Unsolved Problems, *MBio*, 10(2), 1–15, 2019.
- Apts, J. T. H. C. W.: Photosynthetic carbon reduction: high rates in the sea-surface microlayer, *Mar. Biol.*, 101(3), 411–417, doi:10.1007/bf00428138, 1989.
- Azetsu-Scott, K. and Passow, U.: Ascending marine particles: Significance of transparent exopolymer particles (TEP) in the upper ocean, *Limnol. Oceanogr.*, 49(3), 741–748, doi:10.4319/lo.2004.49.3.0741, 2004.
- Berrojalbiz, N., Dachs, J., Ojeda, M. J., Valle, M. C., Jim nez, J. C., Wollgast, J., Ghiani, M., Hanke, G. and Zaldivar, J. M.: Biogeochemical and physical controls on concentrations of polycyclic aromatic hydrocarbons in water and plankton of the Mediterranean and Black Seas, *Global Biogeochem. Cycles*, 25, 1–14, doi:10.1029/2010GB003775, 2011a.
- Berrojalbiz, N., Dachs, J., Vento, S., Del, Jos, M., Valle, M. C., Castro-jim, J., Mariani, G., Wollgast, J. and Hanke, G.: Persistent Organic Pollutants in Mediterranean Seawater and Processes Affecting Their Accumulation in Plankton, *Environ. Sci. Technol.*, 45, 4315–4322, doi:10.1021/es103742w, 2011b.

- Birolli, W. G., Santos, D. D. A., Alvarenga, N. and Garcia, A. C. F. S.: Biodegradation of anthracene and several PAHs by the marine-derived fungus *Cladosporium* sp. CBMAI 1237 Biodegradation of anthracene and several PAHs by the marine-derived fungus *Cladosporium* sp. CBMAI 1237, *Mar. Pollut. Bull.*, 192(2), 525–533, doi:10.1016/j.marpolbul.2017.10.023, 2018.
- Bovio, E.: Marine fungi from sponges: biodiversity, chemodiversity and biotechnological applications., 2019.
- 310 Callahan, B. J., Mcmurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. and Holmes, S. P.: DADA2: High resolution sample inference from Illumina amplicon data, *Nat. Methods*, 13(7), 581–583, doi:10.1038/nmeth.3869.DADA2, 2016.
- Carlson, D.: The Early Diagenesis of Organic Matter: Reaction at the Air-Sea Interface, in *Organic Geochemistry*1, pp. 225–268., 1993.
- Carreto, J. I., Carignan, M. O., Daleo, G. and DeMarco, S. G.: Occurrence of mycosporine-like amino acids in the red-tide
315 dinoflagellate *Alexandrium excavatum*:uv-photoprotective compounds?., *J. Plankton Res.*, 12(5), 909–921 [online] Available from: <http://193.144.41.80:8086/2431.pdf>, 1990.
- Chin, W.-C., Orellana, M. V. and Verdugo, P.: Spontaneous assembly of marine dissolved organic matter into polymer gels, *Nature*, 394(July), 1998.
- Christmas, N. and Cunliffe, M.: Depth-dependent mycoplankton glycoside hydrolase gene activity in the open ocean —
320 evidence from the Tara Oceans eukaryote metatranscriptomes, *ISME J.*, 14, 2361–2365, doi:10.1038/s41396-020-0687-2, 2020.
- Comeau, A. M., Li, W. K. W., Tremblay, J. É., Carmack, E. C. and Lovejoy, C.: Arctic ocean microbial community structure before and after the 2007 record sea ice minimum, *PLoS One*, 6(11), doi:10.1371/journal.pone.0027492, 2011.
- Comeau, A. M., Douglas, G. M. and Langille, M. G. I.: Microbiome Helper: a Custom and Streamlined Workflow for
325 Microbiome Research, *mSystems*, 2(1), e00127-16, doi:10.1128/mSystems.00127-16, 2017.
- Corzo, A., Morillo, J. A. and Rodriguez, S.: Production of transparent exopolymer particles (TEP) in cultures of *Chaetoceros calcitrans* under nitrogen limitation, *Aquat. Mar. Ecol.*, 23, 63–72, 2000.
- Cunliffe, M. and Murrell, J. C.: The sea-surface microlayer is a gelatinous biofilm., *ISME J.*, 3(9), 1001–1003, doi:10.1038/ismej.2009.69, 2009.
- 330 Cunliffe, M. and Murrell, J. C.: Eukarya 18S rRNA gene diversity in the sea surface microlayer: implications for the structure of the neustonic microbial loop., *ISME J.*, 4(3), 455–458, doi:10.1038/ismej.2009.133, 2010.
- Cunliffe, M. and Wurl, O.: Guide to best practices to study the ocean’s surface, Plymouth. [online] Available from: https://repository.oceanbestpractices.org/bitstream/handle/11329/261/SCOR_GuideSeaSurface_2014.pdf?sequence=1&isAll
owed=y, 2014.
- 335 Cunliffe, M., Engel, A., Frka, S., Gašparović, B., Guitart, C., Murrell, J. C., Salter, M., Stolle, C., Upstill-Goddard, R. and Wurl, O.: Sea surface microlayers: A unified physicochemical and biological perspective of the air-ocean interface, *Prog. Oceanogr.*, 109, 104–116, doi:10.1016/j.pocean.2012.08.004, 2013.
- Cunliffe, M., Hollingsworth, A., Bain, C., Sharma, V. and Taylor, J. D.: Algal polysaccharide utilisation by saprotrophic planktonic marine fungi, *Fungal Ecol.*, 30, 135–138, doi:10.1016/j.funeco.2017.08.009, 2017.

- 340 Desboeufs, K., Doussin, J. F., Giorio, C., Triquet, S., Fu, Y., Dulac, F., Garcia-Nieto, D., Chazette, P., Feron, A., Formenti, P., Gaimoz, C., Maisonneuve, F., Riffault, V., Saiz-Lopez, A., Siour, G., Zapf, P. and Guieu, C.: ProcEss studies at the Air/sEa Interface after dust deposition in the MEditerranean sea (PEAcEtIME) cruise: Atmospheric overview, *Biogeosciences*, n.d.
- Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel, C., Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O., Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B., Bouloubassi, I., Brunet, C., Cadiou, J. F., Carlotti, F., Chami, M., Charmasson, S., Charrière, B., Dachs, J., Doxaran, D., Dutay, J. C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F., Fernandez, C., Fowler, S., Francour, P., Gaertner, J. C., Galzin, R., Gasparini, S., Ghiglione, J. F., Gonzalez, J. L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L. E., Jacquet, S. H. M., Jeffrey, W. H., Joux, F., Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C., Lemé, R., Loÿe-Pilot, M. D., Mallet, M., Méjanelle, L., Mélin, F., Mellon, C., Mérigot, B., Merle, P. L., Migon, C., Miller, W. L., Mortier, L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T., Petrenko, A., Poggiale, J. C., Prieur, L., Pujo-Pay, M., Pulido-Villena, Raimbault, P., Rees, A. P., Ridame, C., Rontani, J. F., Ruiz Pino, D., Sicre, M. A., Taillandier, V., Tamburini, C., Tanaka, T., Taupier-Letage, I., Tedetti, M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F., Tronczynski, J., Ulses, C., Van Wambeke, F., Vantrepotte, V., Vaz, S. and Verney, R.: Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, *Prog. Oceanogr.*, 91(2), 97–166, doi:10.1016/j.pocean.2011.02.003, 2011.
- 345 Ekelund, N. G. A.: The Effects of UV-B Radiation on Dinoflagellates, *J. Plant Physiol.*, 138(3), 274–278, 1991.
- Elasri, M. O. and Miller, R. V.: Study of the Response of a Biofilm Bacterial Community to UV Radiation Study of the Response of a Biofilm Bacterial Community to UV Radiation, *Appl. Environ. Microbiol.*, 65(5), 2025–2031, doi:0099-2240, 1999.
- Engel, A.: The role of transparent exopolymer particles (TEP) in the increase in apparent particle stickiness (alpha) during the decline of a diatom bloom, *J. Plankton Res.*, 22(3), 485–497, doi:10.1093/plankt/22.3.485, 2000.
- Engel, A.: Determination of marine gel particles, in *Practical Guidelines for the Analysis of Seawater*, edited by O. Wurl, pp. 125–142, CRC Press Taylor & Francis Group, Boca Raton, FL., 2009.
- Engel, A. and Galgani, L.: The organic sea-surface microlayer in the upwelling region off the coast of Peru and potential implications for air-sea exchange processes, *Biogeosciences*, 13(4), 989–1007, doi:10.5194/bg-13-989-2016, 2016.
- 365 Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E. and Zondervan, I.: Polysaccharide aggregation as a potential sink of marine dissolved organic carbon, *Nature*, 428(April), 929–932, doi:10.1038/nature02506.1., 2004.
- Engel, A., Bange, H. W., Cunliffe, M., Burrows, S. M., Friedrichs, G., Galgani, L., Herrmann, H., Hertkorn, N., Johnson, M., Liss, P. S., Quinn, P. K., Schartau, M., Soloviev, A., Stolle, C., Upstill-Goddard, R. C., van Pinxteren, M. and Zäncker, B.: The Ocean's Vital Skin: Toward an Integrated Understanding of the Sea Surface Microlayer, *Front. Mar. Sci.*, 4(May), 165, doi:10.3389/fmars.2017.00165, 2017.
- 370 Fogg, G. E.: Some comments on picoplankton and its importance in the pelagic ecosystem, *Aquat. Microb. Ecol.*, 9(1), 33–39, doi:10.3354/ame009033, 1995.
- Frenay, E., Sellegrì, K., Nicosia, A., Trueblood, J. T. and Rinaldi, M.: Mediterranean nascent sea spray organic aerosol and

- relationships with seawater biogeochemistry, *Atmos. Chem. Phys.*, (June), 1–23, 2020.
- 375 Garzoli, L., Gnani, G., Tamma, F., Tosi, S., Varese, G. C. and Picco, A. M.: Sink or swim: Updated knowledge on marine fungi associated with wood substrates in the Mediterranean Sea and hints about their potential to remediate hydrocarbons, *Prog. Oceanogr.*, 137, 140–148, doi:10.1016/j.pocean.2015.05.028, 2015.
- Gladfelter, A. S., James, T. Y. and Amend, A. S.: Marine fungi, *Curr. Biol.*, 29, 2019.
- Gnani, G., Garzoli, L., Poli, A., Prigione, V., Burgaud, G. and Varese, G. C.: The culturable mycobiota of *Flabellia petiolata*:
380 First survey of marine fungi associated to a Mediterranean green alga, *PLoS One*, 12(4), 1–20, doi:10.1371/journal.pone.0175941, 2017.
- Godhe, A., Asplund, M. E., Härnström, K., Saravanan, V., Tyagi, A. and Karunasagar, I.: Quantification of diatom and dinoflagellate biomasses in coastal marine seawater samples by real-time PCR, *Appl. Environ. Microbiol.*, 74(23), 7174–7182, doi:10.1128/AEM.01298-08, 2008.
- 385 Grossart, H.-P., Van den Wyngaert, S., Kagami, M., Wurzbacher, C., Cunliffe, M. and Rojas-Jimenez, K.: Fungi in aquatic ecosystems, *Nat. Rev. Microbiol.*, 17(6), 339–354, 2019.
- Guieu, C., D’Ortenzio, F., Dulac, F., Taillandier, V., Doglioli, A., Petrenko, A., Barrillon, S., Mallet, M., Pierre, N. and Desboeufs, K.: Process studies at the air-sea interface after atmospheric deposition in the Mediterranean Sea : objectives and strategy of the PEACETIME oceanographic campaign (May – June 2017), 2020.
- 390 Guo, L., Sui, Z. and Liu, Y.: Quantitative analysis of dinoflagellates and diatoms community via Miseq sequencing of actin gene and v9 region of 18S rDNA, *Sci. Rep.*, 6(May), 1–9, doi:10.1038/srep34709, 2016.
- Häder, D.-P., Kumar, H. D., Smith, R. C. and Worrest, R. C.: Effects of solar UV radiation on aquatic ecosystems and interactions with climate change, *Photochem. Photobiol. Sci.*, 6(3), 267–285, doi:10.1039/B700020K, 2007.
- Hardy, J. T. and Apts, C. W.: The sea-surface microlayer: phytoneuston productivity and effects of atmospheric particulate
395 matter, *Mar. Biol.*, 82, 293–300, 1984.
- Hardy, J. T., Coley, J. A., Antrim, L. D. and Kiessler, S. L.: A hydrophobic large-volume sampler for collecting aquatic surface microlayers: characterization and comparison with the glass plate method, *Can. J. Fish. Aquat. Sci.*, 45(5), 822–826, 1988.
- Harvey, G.: Microlayer collection from the sea surface: a new method and initial results, *Limnol Ocean.*, 11(4), 608–613, 1966.
- Hassett, B. T., Borrego, E. T., Vonnahme, T. R., Rämä, T., Kolomiets, M. V. and Gradinger, R.: Arctic marine fungi: Biomass,
400 functional genes, and putative ecological roles, *ISME Journal*, 13(6), 1484–1496, 2019.
- Hassett, B. T., Vonnahme, T. R., Peng, X. and Jones, E. B. G.: Global diversity and geography of planktonic marine fungi, *Bot. Mar.*, 63(2), 121–139, 2020.
- Jennings, M. K., Passow, U., Wozniak, A. S. and Hansell, D. A.: Distribution of transparent exopolymer particles (TEP) across an organic carbon gradient in the western North Atlantic Ocean, *Mar. Chem.*, 190, 1–12, doi:10.1016/j.marchem.2017.01.002,
405 2017.
- Joux, F., Agogue, H., Obernosterer, I., Dupuy, C., Reinthaler, T., Herndl, G. J. and Lebaron, P.: Microbial community structure in the sea surface microlayer at two contrasting sites in the northwestern Mediterranean Sea., *Aquat. Microb. Ecol.*, 42, 91–

104, doi:10.3354/ame042091, 2006.

- Krom, M. D., Herut, B. and Mantoura, R. F. C.: Nutrient budget for the Eastern Mediterranean: Implications for phosphorus
410 limitation, *Limnol. Oceanogr.*, 49(5), 1582–1592, doi:10.4319/lo.2004.49.5.1582, 2004.
- Lepesteur, M., Martin, J. M. and Fleury, A.: A comparative study of d methods for phytoplank flow cytometry, *Mar. Ecol. Prog. Ser.*, 93(1/2), 55–63, 1993.
- Liss, P. S. and Duce, R. A.: *The sea surface and global change*, Cambridge University Press, Cambridge., 2005.
- Martin, J. M., Elbaz-Poulichet, F., Guieu, C., Loÿe-Pilot, M. D. and Han, G.: River versus atmospheric input of material to the
415 mediterranean sea: an overview, *Mar. Chem.*, 28(1–3), 159–182, doi:10.1016/0304-4203(89)90193-X, 1989.
- Marty, J. C., Chiavérini, J., Pizay, M. D. and Avril, B.: Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991-1999), *Deep. Res. Part II Top. Stud. Oceanogr.*, 49(11), 1965–1985, doi:10.1016/S0967-0645(02)00022-X, 2002.
- Mermex Group, T., Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D’Ortenzio, F., Estournel, C.,
420 Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O., Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B., Bouloubassi, I., Brunet, C., Cadiou, J. F., Carlotti, F., Chami, M., Charmasson, S., Charrière, B., Dachs, J., Doxaran, D., Dutay, J. C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F., Fernandez, C., Fowler, S., Francour, P., Gaertner, J. C., Galzin, R., Gasparini, S., Ghiglione, J. F., Gonzalez, J. L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L. E., Jacquet, S. H. M., Jeffrey, W. H., Joux, F., Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C.,
425 Lemé, R., Loÿe-Pilot, M. D., Mallet, M., Méjanelle, L., Mélin, F., Mellon, C., Mériçot, B., Merle, P. L., Migon, C., Miller, W. L., Mortier, L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T., Petrenko, A., Poggiale, J. C., Prieur, L., Pujo-Pay, M., Pulido-Villena, Raimbault, P., Rees, A. P., Ridame, C., Rontani, J. F., Ruiz Pino, D., Sicre, M. A., Taillandier, V., Tamburini, C., Tanaka, T., Taupier-Letage, I., Tedetti, M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F., Tronczynski, J., Ulses, C., Van Wambeke, F., Vantrepotte, V., Vaz, S. and Verney, R.: Marine ecosystems’ responses to climatic and
430 anthropogenic forcings in the Mediterranean, *Prog. Oceanogr.*, 91(2), 97–166, doi:10.1016/j.pocean.2011.02.003, 2011.
- Mopper, K., Zhou, J., Ramana, K. S., Passow, U., Dam, H. G. and Drapeau, D. T.: The role of surface active carbohydrates in the flocculation of a diatom bloom in a mesocosm, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 42(1), 43–73, 1995.
- Naumann, E.: Über das Neuston des Süßwassers, *Biol. Cent.*, 37(2), 98–106, 1917.
- Obernosterer, I., Catala, P., Lami, R., Caparros, J., Ras, J., Bricaud, a, Dupuy, C., Van Wambeke, F. and Lebaron, P.:
435 Biochemical characteristics and bacterial community structure of the sea surface microlayer in the South Pacific Ocean, *Biogeosciences Discuss.*, 4(4), 693–705, doi:10.5194/bgd-4-2809-2007, 2008.
- Ortega-Retuerta, E., Passow, U., Duarte, C. and Reche, I.: Effects of ultraviolet B radiation on (not so) transparent exopolymer particles, *Biogeosciences*, 6, 3071–3080, doi:10.5194/bgd-6-7599-2009, 2009.
- Passow, U.: Transparent Exopolymer Particles in Aquatic Environments, *Prog. Oceanogr.*, 55(3–4), 287–333,
440 doi:10.1016/S0079-6611(02)00138-6, 2002.
- Passow, U. and Alldredge, A. L.: Aggregation of a diatom bloom in a mesocosm: The role of transparent exopolymer particles

- (TEP), *Deep. Res. Part II*, 42(1), 99–109, doi:10.1016/0967-0645(95)00006-C, 1995.
- Ploug, H.: Cyanobacterial surface blooms formed by *Aphanizomenon* sp. and *Nodularia spumigena* in the Baltic Sea: Small-scale fluxes, pH, and oxygen microenvironments, *Limnol. Oceanogr.*, 53(3), 914–921, doi:10.4319/lo.2008.53.3.0914, 2008.
- 445 Pujo-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J. F., Goyet, C., Moutin, T. and Prieur, L.: Integrated survey of elemental stoichiometry (C , N , P) from the Western to Eastern Mediterranean Sea, *Biogeosciences*, 8(4), 883–899, doi:10.5194/bgd-7-7315-2010, 2011.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, *Nucleic Acids Res.*, 41(D1), 590–596, 450 doi:10.1093/nar/gks1219, 2013.
- R Core Team: R: A language and environment for statistical computing. Vienne, Austria: R Foundation for Statistical Computing, Vienna., 2014.
- Reddaway, J. M. and Bigg, G. R.: Climatic change over the Mediterranean Sea and links to the more general atmospheric circulation, *Int. J. Clim. Chang.*, 16, 651–661, 1996.
- 455 Schneider, C. a, Rasband, W. S. and Eliceiri, K. W.: NIH Image to ImageJ: 25 years of image analysis, *Nat. Methods*, 9(7), 671–675, doi:10.1038/nmeth.2089, 2012.
- Sellegrì, K., Nicosia, A., Freney, E., Uitz, J., Thyssen, M., Grégori, G., Engel, A., Zäncker, B., Haëntjens, N., Mas, S., Picard, D., Saint-Macary, A., Peltola, M., Rose, C., Trueblood, J., Lefevre, D., D’Anna, B., Desboeuf, K., Meskhidze, N., Guieu, C. and Law, C. S.: Surface ocean microbiota determine cloud precursors, *Sci. Rep.*, n.d.
- 460 Sun, C., Sperling, M. and Engel, A.: Effect of wind speed on the size distribution of biogenic gel particles in the sea surface microlayer : Insights from a wind wave channel experiment , (October), 1–39, doi:10.5194/bg-15-3577-2018, 2018.
- Tanhua, T., Hainbucher, D., Schroeder, K., Cardin, V., Álvarez, M. and Civitarese, G.: The Mediterranean Sea system: A review and an introduction to the special issue, *Ocean Sci.*, 9(5), 789–803, doi:10.5194/os-9-789-2013, 2013.
- Taylor, J. D. and Cunliffe, M.: High-throughput sequencing reveals neustonic and planktonic microbial eukaryote diversity in 465 coastal waters, *J. Phycol.*, 50(5), 960–965, doi:10.1111/jpy.12228, 2014.
- Tovar-Sánchez, A., Rodríguez-Romero, A., Engel, A., Zäncker, B., Fu, F., Marañón, E., Pérez-Lorenzo, M., Bressac, M., Wagener, T., Triquet, S., Siour, G., Desboeufs, K. and Guieu, C.: Characterizing the surface microlayer in the Mediterranean Sea: Trace metal concentrations and microbial plankton abundance, *Biogeosciences*, 17(8), 2349–2364, doi:10.5194/bg-17-2349-2020, 2020.
- 470 Trueblood, J. V, Nicosia, A., Engel, A., Zäncker, B., Rinaldi, M., Thyssen, M., Obernosterer, I., Dinasquet, J., Belosi, F., Tovar-, A., Rodriguez-romero, A., Santachiara, G., Guieu, C. and Sellegrì, K.: A Two-Component Parameterization of Marine Ice Nucleating Particles Based on Seawater Biology and Sea Spray Aerosol Measurements in the Mediterranean Sea, *Atmos. Chem. Phys.*, (June), 1–17, 2020.
- Upstill-Goddard, R. C., Frost, T., Henry, G. R., Franklin, M., Murrell, J. C. and Owens, N. J. P.: Bacterioneuston control of 475 air-water methane exchange determined with a laboratory gas exchange tank, *Global Biogeochem. Cycles*, 17(4), 1108,

doi:10.1029/2003GB002043, 2003.

Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U. and Santschi, P. H.: The oceanic gel phase: A bridge in the DOM-POM continuum, *Mar. Chem.*, 92(1-4 SPEC. ISS.), 67–85, doi:10.1029/2002GL016046, 2004.

480 Wietz, M., Wemheuer, B., Simon, H., Giebel, H. A., Seibt, M. A., Daniel, R., Brinkhoff, T. and Simon, M.: Bacterial community dynamics during polysaccharide degradation at contrasting sites in the Southern and Atlantic Oceans, *Environ. Microbiol.*, 17(10), 3822–3831, doi:10.1111/1462-2920.12842, 2015.

Wikner, J. and Hagstrom, A.: Evidence for a tightly coupled nanoplanktonic predator-prey link regulating the bacterivores in the marine environment, *Mar. Ecol. Prog. Ser.*, 50, 137–145, doi:10.3354/meps050137, 1988.

World Health Organization: The sea-surface microlayer and its role in global change, Geneva., 1995.


485 Wurl, O. and Holmes, M.: The gelatinous nature of the sea-surface microlayer, *Mar. Chem.*, 110(1–2), 89–97, doi:10.1016/j.marchem.2008.02.009, 2008.

Wurl, O., Miller, L., Röttgers, R. and Vagle, S.: The distribution and fate of surface-active substances in the sea-surface microlayer and water column, *Mar. Chem.*, 115(1–2), 1–9, doi:10.1016/j.marchem.2009.04.007, 2009.

490 Zäncker, B., Bracher, A., Röttgers, R. and Engel, A.: Variations of the Organic Matter Composition in the Sea Surface Microlayer: A Comparison between Open Ocean, Coastal, and Upwelling Sites Off the Peruvian Coast, *Front. Microbiol.*, 8(December), 1–17, doi:10.3389/fmicb.2017.02369, 2017.


Zhang, Z., Liu, L., Liu, C. and Cai, W.: Studies on the sea surface microlayer: II. The layer of sudden change of physical and chemical properties, *J. Colloid Interface Sci.*, 264(1), 148–159, doi:10.1016/S0021-9797(03)00390-4, 2003.

495 Zhou, J., Bruns, M. A. and Tiedje, J. M.: DNA recovery from soils of diverse composition, *Appl. Environ. Microbiol.*, 62(2), 316–322, doi:D - NLM: PMC167800 EDAT- 1996/02/01 MHDA- 1996/02/01 00:01 CRDT- 1996/02/01 00:00 PST - ppublish, 1996.




500


Figure 1: Map of the stations sampled during the PEACETIME cruise in the Mediterranean Sea in May/June 2017. Stations FAST and TYRR were sampled twice. Colours represent sampled basins (blue: western basin, green: Tyrrhenian Sea, red: Ionian Sea).




510 **Figure 2: Principal Component Analyses (PCA) using the eukaryotic community composition at the ASV level (see text for a detailed description) with environmental factors plotted. Colours distinguish the three different basins sampled (blue: western basin, green: Tyrrhenian Sea, red: Ionian Sea).**



515 **Figure 3: Eukaryotic community composition at the order level (taxa over 5 percent in at least one of the samples are displayed). Stations ordered from west to east with brackets indicating the western Mediterranean (blue), the Tyrrhenian Sea (green) and the Ionian Sea (red).**



520 **Figure 4: Heatmap of fungal relative ASV abundances in all sequences samples. Colour brackets indicate the different basins as in Fig. 3 and Fig. 2. Grey corresponds to the absence of the ASV in the respective sample.**



525 **Figure 5: Boxplots of bacteria (A) and phytoplankton (B) abundance as well as area (C) and concentrations (D) of Transparent Exopolymer Particles (TEP) and total carbohydrates (TCHO) (E) for each basin in the Mediterranean Sea: western basin (west), Tyrrhenian Sea (tyr) and Ionian Sea (ion). Blue stars mark significant SML enrichment/depletion, green stars mark significant differences between the three basins (Kruskal-Wallis tests used for significance levels). Significance levels: *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$. Black dots correspond to outliers, grey dots to the measured values and concentrations.**

530

Table 1: Name, position and environmental conditions at the stations sampled throughout the cruise. Temperature and salinity were collected at 5 m water depth.

station	Latitude	Longitude	Local time	wind speed [m s ⁻¹]	water temperature [°C]	salinity [PSU]	irradiation [W m ⁻²]
S1	41.8918	6.3333	15:45	9.7	16.4	38.2	1297.8
S3	39.1333	7.6835	10:00	2.9	18.7	37.2	2343.2
S4	37.9832	7.9768	10:30	3.5	19.8	37.1	2270.2
TYRR_1	39.34	12.5928	11:00	3.4	20.3	37.8	2253.1
TYRR_2	39.3398	12.5928	12:30	2.5	21.1	37.7	2311.1
S6	38.8077	14.4997	9:00	5.2	20.4	37.4	2215.5
SAV	37.8401	18.1658	12:00	1.5	20.1	38.5	
S7	36.6035	18.1658	7:00	2.5	20.8	38.5	16.8
ION_2	35.4892	19.7765	9:45	6.4	21.1	38.8	1235.3
S8	36.2103	16.631	7:45	1.9	21.2	37.9	2144.0
FAST_2	37.946	2.9102	8:30	3.1	21.7	36.7	627.4
FAST_6	37.0466	2.9168	8:30	5.1	21.9	36.6	1787.1