

# Estimating maximum fine fraction organic carbon in UK grasslands

Kirsty C. Paterson<sup>1,2</sup>, Joanna M. Cloy<sup>1</sup>, Robert. M. Rees<sup>1</sup>, Elizabeth M. Baggs<sup>2</sup>, Hugh Martineau<sup>3</sup>, Dario Fornara<sup>4</sup>, Andrew J. Macdonald<sup>5</sup>, and Sarah Buckingham<sup>1</sup>

5 <sup>1</sup> Scotland's Rural College, West Mains Road, Edinburgh, EH9 3JG, United Kingdom.

<sup>2</sup> Global Academy of Agriculture and Food Security, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, United Kingdom.

<sup>3</sup> Treberfydd Farm, Wales, United Kingdom.

<sup>4</sup> Agri-Food & Biosciences Institute (AFBI), Newforge Lane, BT9 5PX, Belfast, United Kingdom.

10 <sup>5</sup> Sustainable Agriculture Sciences Department, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, United Kingdom.

*Correspondence to:* Sarah Buckingham ([sarah.buckingham@sruc.ac.uk](mailto:sarah.buckingham@sruc.ac.uk))

**Abstract.** Soil organic carbon (SOC) sequestration across agroecosystems worldwide can contribute to mitigate the effects of  
15 climate change by reducing levels of atmospheric CO<sub>2</sub>. Stabilisation of organic carbon (OC) in the fine soil fraction (< 20 µm)  
is considered an important long-term store of SOC and the saturation deficit (difference between measured OC and estimated  
maximum OC in the fine fraction) is frequently used to assess SOC sequestration potential following the linear regression  
equation developed by Hassink, (1997). However, this approach is often taken without any assessment of the fit of the equation  
to the soils being studied. The statistical limitations of linear regression have previously been noted, giving rise to the proposed  
20 use of boundary line analysis and quantile regression to provide more robust estimates of maximum SOC stabilisation. The  
objectives of this work were to assess the suitability of the Hassink, (1997) equation to estimate maximum fine fraction OC in  
UK grassland soils of varying sward ages and to evaluate the linear regression, boundary line and quantile regression methods  
to estimate maximum fine fraction OC. A chronosequence of 10 grasslands was sampled, in order to assess the relationship  
between sward age (time since last reseeding event) and the measured and predicted maximum fine fraction OC. Significantly  
25 different regression equations show that the Hassink, (1997) equation does not accurately reflect maximum fine fraction OC  
in UK grasslands when determined using the proportion of fine soil fraction (< 20 µm, %) and measured fine fraction OC (g  
C kg<sup>-1</sup> soil). The QR estimate of maximum SOC stabilisation was almost double that of linear regression and BL analysis (0.89  
± 0.074, 0.43 ± 0.017 and 0.57 ± 0.052 g C kg<sup>-1</sup> soil, respectively). Sward age had an inconsistent effect on the measured  
variables and potential maximum fine fraction OC. Fine fraction OC across the grasslands made up 4.5 to 55.9% of total SOC,  
30 implying that there may be either high potential for additional C sequestration in the fine fraction of these soils, or stabilisation  
in aggregates is predominant in these grassland soils. This work highlights the need to ensure that methods used to predict  
maximum fine fraction OC reflect the soil *in situ*, resulting in more accurate assessments of carbon sequestration potential.

## 1. Introduction

35 Carbon (C) sequestration in soils offers a significant opportunity to remove CO<sub>2</sub> from the atmosphere and store it into  
long lived C pools (Lal, 2004; Powlson et al., 2011), with co-benefits for soil structure and functioning (Lorenz and Lal, 2018;  
Smith, 2012; Soussana et al., 2004). However, to utilise soils as a CO<sub>2</sub> drawdown mechanism, accurate estimates of their  
storage capability are required. With respect to soil organic carbon (SOC) sequestration, organic carbon (OC) stabilised via  
40 adsorption to mineral surfaces in the fine soil fraction (< 20 µm) is often regarded as the most important due to its longer  
residence time (Baldock and Skjemstad, 2000; Six et al., 2002). There is empirical evidence that there is an upper protective  
capacity limit, or saturation point of the mineral stabilised OC pool (Six et al., 2002; Stewart et al., 2007). Potential SOC  
sequestration (or saturation deficit) can be estimated by subtracting the current fine fraction OC from the estimated maximum  
fine fraction OC (Angers et al., 2011).

Hassink, (1997) compared pairs of Dutch arable and grassland soils and found that while soil bulk SOC contents  
45 significantly differed among soils, fine fraction OC did not. These findings led to the idea that the saturation point of the fine  
soil fraction could be estimated by linear regression using the mass proportion of fine fraction in a soil sample (%) and the  
current fine fraction OC (g C kg<sup>-1</sup> soil). Several iterations of the concept have been proposed to overcome the limitations of  
linear regression. For example, boundary line analysis uses a defined upper or lower subset of a data set to estimate the  
boundary line, when a limiting response to an independent variable(s) along a boundary is supported (Lark and Milne, 2016;  
50 Schmidt et al., 2000). Using the upper 90<sup>th</sup> percentile of a data set, boundary line analysis overcomes the limitation of linear  
regression depicting the mean response to the independent variable (Feng et al., 2013; Shatar and Mcbratney, 2004), which is  
thought to cause an underestimation of sequestration potential. Quantile regression estimates the response of a specific quartile  
using the entire data set. It also makes no assumptions regarding homogeneity of variance, thus increasing the robustness of  
the estimated maximum fine fraction OC, as sample size is not reduced as in BL analysis (Beare et al., 2014; Cade and Noon,  
55 2003). Using a forced zero intercept overcomes the contradiction of a positive intercept indicating the presence of fine fraction  
OC without any fine soil fraction (Beare et al., 2014; Feng et al., 2013; Liang et al., 2009). These suggestions have been  
proposed to improve estimates of maximum fine fraction OC. However, several studies use the original equation presented by  
Hassink, (1997) to estimate sequestration potential at different scales (e.g. Angers et al., 2011; Chen et al., 2019; Lilly and  
Baggaley, 2013; Wiesmeier et al., 2014). There is a need for validity checks to determine the suitability of the Hassink, (1997)  
60 linear regression equation to predict maximum fine fraction OC of the soils in the respective studies. Without this sequestration  
potentials may be both over and underestimated.

Within the UK, human-managed grasslands are the dominant land use, covering 36% of the land area (Ward et al., 2016). Managed grasslands are planted and maintained to increase agricultural productivity through fertiliser and liming applications, and the re-seeding of swards. The high levels of disturbance associated with re-seeding events by mould board ploughing and harrowing in particular, result in changes in soil structure, notably the breaking up of aggregates, nutrient cycling and SOC mineralisation (Carolan and Fornara, 2016; Drewer et al., 2017; Soussana et al., 2004). Organo-mineral associations form the basis of microaggregates (Balcock and Skjemstad, 2000), and thus the destruction of aggregates makes the organo-mineral stabilised OC in the fine fraction, more accessible for mineralisation by the soil microbial community. Additionally, the release of other organic carbon pools may induce a priming effect, potentially enhancing the losses from the typically stable mineral associated OC in the fine fraction. The long-term effect of such re-seeding event on SOC dynamics is understudied, it is therefore important to understand how disturbance might affect OC in the fine fraction, and thus the SOC sequestration ability of managed grasslands.

The objectives of this study were (i) to assess the suitability of the Hassink, (1997) equation to estimate maximum fine fraction OC in UK grassland soils, (ii) to evaluate the linear regression, boundary line and quantile regression methods to estimate maximum fine fraction OC, and (iii) to explore the relationship between sward age (time since last reseeded event), and current and predicted maximum fine fraction OC. We hypothesised that i) the linear regression equation developed using UK grassland soils would be significantly different to that of Hassink, (1997), and that ii) grasslands with an older sward age, would have a greater proportion of total SOC stabilised in fine fraction ( $< 20 \mu\text{m}$ ) and a lower sequestration potential.

## 2. Materials and Methods

### 2.1 Site Description and Sampling

Ten grassland chronosequences covering a wide range of soil types, land use and climatic conditions were identified across the UK in 2016. The sites included the range of agricultural activity associated with UK grasslands (upland grazing, dairy, and mixed grazing), variations in soil type (organo-mineral, mineral and chalk) and the majority of UK climatic zones (Table 1). At each location, five to eight individual fields of different sward age (represented by years since a ploughing and reseeded event), ranging from 1 to 179 years, were identified for sampling. In each field, areas were avoided which had different applications of manure, soil types or topography, headlands, areas near gates, where lime or manure had previously been dumped, or where livestock congregate. Two replicate soil cores were collected to a depth of 30 cm using a soil auger with a 2.5 cm diameter steel core and bulked to give a single composite sample. This was repeated 10 times in each field at regular intervals in a 'W' shape across the field totalling 10 replicate samples per field per site. Intact soil cores for determining bulk densities were collected at three locations in each field at two depths (10 to 15 cm and 20 to 25 cm) using intact rings (7.5 cm

diameter, 5 cm height). Replicate samples were sieved to 2 mm and fresh subsamples were used to determine soil pH in water. Remaining sieved soils were dried at 40°C and ball milled prior to determination of total C and N contents (% by mass) using a Flash 2000 elemental analyser. Intact soils were dried at 107°C and weighed to calculate dry bulk densities, any stones were removed.

## 2.2 Soil fractionation

The fine fraction (< 20 µm) of the soil was separated using a combined ultrasonic dispersion and sedimentation method adapted from Hassink, (1997). Briefly, 20 g of dried sieved soil was soaked in 100 mL of deionised water for 24 hours. The suspension was then sonicated with a Microson XL2000 Ultrasonicator for 20 minutes at 20 W in 50 mL centrifuge tubes, surrounded by ice to prevent overheating. The separated samples were recombined in 150 mL tubes, and shaken end over end to disperse the soil water suspension. Sedimentation times were determined using a table applying Stokes Law, for 20 µm particles, a particle density of 2.65 g cm<sup>-3</sup> and sedimentation depth of 5 cm at temperatures between 20°C and 35°C (Jackson, 2005). After the appropriate sedimentation time, the fine fraction was siphoned off the soil suspension. The fine fraction was dried for 24 hours at 107°C and ball milled prior to total C and N analysis (% by mass) using a Flash 2000 Elemental Analyser, to determine the current OC content of the fine fraction. At each site, a minimum of 3 fields varying in age (young, intermediate, and old at that location) were selected, and 3 of the 10 replicate field samples were selected at random for fractionation.

Hydrochloric acid (HCl) fumigation was used to remove carbonates from the Plumpton samples. Ball-milled samples, in silver capsules, were moistened with deionised water (1:4 sample:water ratio) to aid the efficiency of carbonate removal by HCl fumes (Dhillon et al., 2015). The samples were placed in a vacuum desiccator with a beaker of 100 mL of 12 M HCl, for 24 hours and subsequently dried in a ventilated oven at 60°C for 16 hours, to remove excess moisture and HCl (Dhillon et al., 2015). Total C and N contents were determined as outlined above.

## 2.3 Statistical analyses

All statistical analyses were carried out using R software version 3.5.3 (Team, 2019). Significant differences were determined by ANOVA's and by post-hoc Tukey tests ( $\alpha = 0.05$ ). Where assumptions of normality and variance were not satisfied by testing (Shapiro Wilkens and Levenes Test) significant differences were identified using Kruskal test and post hoc Dunn test. A Kendal tau ( $\tau$ ) correlation matrix was produced using the 'corrplot' package (Wei and Simko, 2017).

### 120 2.3.1 Regression analyses

Linear regression was used to predict maximum fine fraction OC, with the mass proportion of fine fraction (< 20  $\mu\text{m}$ , %) in a sample and the measured OC of the fine fraction ( $\text{g C kg}^{-1}$  soil) as the independent and dependent variables, respectively. Regression equations were developed for the combined UK data set, and the individual sites. Linear regression with a forced zero intercept was used with data from this study and the data published in Hassink (1997).

125           Boundary line analyses were performed as an alternative to linear regression, both with and without a forced zero intercept to predict maximum fine fraction OC for all UK sites. The data was organised by mass proportion of the fine fraction (%) and divided into subgroups at 5, 10 and 15% intervals. The 10% interval reflects the method of Feng et al. (2013), whilst the 5 and 15% intervals were used to assess the effect of interval on estimation of maximum fine fraction OC. The groups were then ordered by measured fine fraction OC ( $\text{g C kg}^{-1}$  soil), and the values in the 90th percentile were used to plot the boundary  
130 line. Boundary line analysis was not used for individual sites as it resulted in too few data points. Quantile regression analysis was performed in Rstudio using the ‘quantreg’ package (Koenker, 2019), for the 90th and median percentiles ( $\tau = 0.90$  and  $\tau = 0.50$ ). Forcing the intercept to zero overcomes the paradox of having C stabilised as MAOC without any fine fraction in the soil. Significant differences between slopes were identified using the ‘lsmeans’ package (Lenth, 2016), followed by post-hoc Tukey tests ( $\alpha = 0.05$ ).

135

### 2.3.2 Carbon saturation ratio

The carbon saturation ratio was determined in order to identify the degree of saturation across the sites, when estimating maximum fine fraction OC, using the Hassink (1997), UK, and site-specific linear regression equations both with and without a forced zero intercept, and the equations generated by boundary line and quantile regression analyses. The carbon saturation  
140 ratio was calculated by dividing the current fine fraction OC by the estimated maximum fine fraction OC content. Values < 1 were deemed under saturated, = 1 as at saturation and > 1 as oversaturated.

## 3. Results

### 3.1 Current C concentrations

145 The measured total SOC and fine fraction OC concentrations exhibited variation within the grassland sites (Fig. 1). Total SOC varied from 8.2 to 85.8  $\text{g C kg}^{-1}$  soil, with a median of 32.7  $\text{g C kg}^{-1}$  soil. Hillsborough, Overton and Plumpton had significantly higher total SOC, whilst Harpenden and Llangorse had the lowest total SOC ( $P < 0.05$ ) (Fig. 1). The measured fine fraction OC ranged from 1.4 to 20.9  $\text{g C kg}^{-1}$  soil, with a median of 6.2  $\text{g C kg}^{-1}$  soil. Overton had the highest total fine fraction OC

( $P < 0.05$ ) and was the only organically managed site (Fig. 1). The proportion of OC stabilised in the fine fraction ( $< 20 \mu\text{m}$ ) had high variability across the UK sites accounting for 4.5 to 50.1% of total SOC with a median of 17.5%. The proportion of total SOC stabilised in the fine fraction ( $< 20 \mu\text{m}$ ), and proportion of fine fraction in a sample did not significantly differ in Harpenden and Overton, however they have significantly different measured fine fraction OC contents ( $\text{g C kg}^{-1}$  soil) ( $P < 0.05$ ), indicating different saturation potentials (Fig. 1). Soil C:N ratio was positively correlated with fine fraction C:N (0.30,  $P < 0.0001$ ), Table 2, however there was no relationship between bulk soil C:N ratio and proportion of fine fraction (data not shown). The fine fraction C:N ratio was significantly different between the sites, Figure 1, however the mean of all the data showed little deviation,  $9.84 \pm 1.00$  (mean  $\pm$  standard deviation). Full details of all the measured properties of bulk and fine fraction, per field are presented in Table A1.

The significance of correlations between the measured soil properties, time since reseeded and known environmental factors were analysed. The matrix of Kendall tau ( $\tau$ ) correlation coefficients in Table 2, revealed that measured fine fraction OC was positively correlated with median annual temperature ( $\tau = 0.13$ ,  $P < 0.05$ ), %N ( $\tau = 0.26$ ,  $P < 0.0001$ ) and %C ( $\tau = 0.27$ ,  $P < 0.0001$ ) in the bulk soil, and negatively correlated with mean annual rainfall ( $\tau = -0.36$ ,  $P < 0.0001$ ), and %N ( $\tau = -0.15$ ,  $P < 0.05$ ) in the fine fraction. Mass proportion of fine fraction and measured fine fraction OC ( $\text{g C kg}^{-1}$  soil) were positively correlated in cambisols ( $R^2 = 0.61$ ,  $P < 0.05$ ), gleysols ( $R^2 = 0.76$ ,  $P < 0.05$ ), podzols ( $R^2 = 0.93$ ,  $P < 0.05$ ), and stagnosols ( $R^2 = 0.88$ ,  $P < 0.05$ ) (Fig. 2). However, the proportion of total SOC stabilised in the fine fraction ( $< 20 \mu\text{m}$ ), was greatest in luvisols ( $P < 0.05$ ) (Fig. 3).

### 3.2 Estimated maximum fine fraction organic carbon

The slope generated from the UK data used to estimate MAOC (Table 3) was significantly different ( $P < 0.05$ ) to the slope reported in Hassink (1997). There was no significant difference between the slopes generated from the UK data, the data from Hassink (1997) when estimated by linear regression with a forced zero intercept. Significantly different ( $P < 0.05$ ) slopes were found between the individual UK sites, owing to the range in the proportion of the fine fraction within each sample, from 1.85 to 51.8%, (Tables A3 and A4).

Coefficients from boundary line analysis are presented in (Table 3). There was no significant difference in slopes between the 5, 10, and 15% fine fraction intervals used. The median percentile quantile regression analysis had a similar slope to the boundary line and linear regression with forced zero intercept. Quantile regression using the 90<sup>th</sup> percentile resulted in the steepest slope of all estimation methods (Table 3). The C saturation ratios revealed the difference in number of samples with potential to sequester more C (Table 4). The Hassink (1997) linear regression equation, without a forced zero intercept, predicted the greatest number of unsaturated sites, followed by the 90<sup>th</sup> percentile quantile regression, with a forced zero

intercept. There was no clear relationship between oversaturated sites and proportion of silt and clay contents as oversaturation  
180 occurred across all proportions, indicated by points above the lines in Fig. 4.

### 3.3 Effect of sward age on current C concentrations and estimated maximum MAOC

Sward age (years since last reseeding event) had a weak positive correlation with the mass proportion of the fine fraction (%)  
(Table 2). When grouped in five year intervals, significant differences were found between age group and the mass proportion  
185 of the fine fraction (%), measured fine fraction OC ( $< 20 \mu\text{m}$ ), ( $\text{g C kg}^{-1}$  soil), and the C:N ratio of the fine fraction (Table 5),  
however there was no consistent increase or decrease with sward age. At the individual sites, significant differences were  
observed between fields, with some properties, but again there was no consistent effect of sward age (Tables A3 and A4).

## 4. Discussion

### 4.1 Estimation of maximum fine fraction organic carbon

190 Determining the potential C sequestration capacity of soils is essential to predict the influence of land management for climate  
change mitigation. The determination of saturation deficit using the mass proportion of the fine fraction and current fine  
fraction OC content is an established method with a strong grounding in correlation between the variables. However, the  
Hassink (1997) linear regression equation has been used to estimate sequestration potentials, without prior testing to determine  
its applicability to the soils in question (e.g. Angers et al., 2011; Chen et al., 2019; Lilly and Baggaley, 2013; Wiesmeier et al.,  
195 2014). This may have potentially over or underestimated sequestration potential, which may have repercussions for decisions  
made regarding land management. The significantly different slopes for the linear regression equations (Table 3) shows that  
the Hassink (1997) regression equation is not suitable for estimating maximum fine fraction OC in UK grasslands. Previous  
concerns have focused on the potential for the equation developed by Hassink (1997) to underestimate maximum fine fraction  
OC, as linear regression represents the mean response of the independent variable, rather than the maximum. For the UK  
200 grasslands in this study estimating maximum fine fraction OC using the Hassink (1997) regression approach resulted in a  
significant overestimation of fine fraction OC sequestration potential. Future work using maximum fine fraction OC  
predictions equations reported in the literature (e.g. Beare et al., 2014; Feng et al., 2013; Hassink, 1997; Six et al., 2002) should  
first conduct a validity test, and determine if the regression equations match the soils in question or a subset of the data, to  
ensure results are not significantly over or underestimated.

205 To overcome the contradiction of an intercept greater than zero, indicating that C is stabilised in the fine fraction  
without any fine fraction, a forced zero intercept was used. The linear regression slopes with a forced zero intercept were not  
significantly different, and were similar to that of Feng et al. (2013),  $0.42 \pm 0.002$ . Liang et al. (2009) reported a lower slope

of 0.36 in Chinese black soils, whilst Beare et al. (2014) reported a slope of  $0.70 \pm 0.03$  in long-term New Zealand pastures. The range of reported values, and differences across the UK sites (Tables A3 and A4), suggest that the effect of the proportion of fine fraction of a sample on fine fraction OC is not consistent and likely reflects differences in pedogenic and environmental conditions, land management and possibly the fine fraction OC isolation method. It may be that the use of the mass proportion of fine fraction to predict maximum fine fraction OC is only suited on larger scales, rather than smaller, site specific scales, as indicated by the variability in this study.

Boundary line analysis and quantile regression have been suggested as alternatives to overcome the limitations of linear regression. The estimation of maximum fine fraction OC was greatest when using quantile regression ( $\tau = 0.90$ ), whereas boundary line estimates at 5 and 10% intervals were similar to quantile regression ( $\tau = 0.50$ ), and those estimated from linear regression (Table 3). The use of the median percentile quantile regression highlights the closeness of linear regression predictions being more indicative of mean values, thus underestimating SOC sequestration potential. The boundary line estimate of Feng et al. (2013),  $0.89 \pm 0.05$ , was nearly double their linear regression; this was not the case in our study. Boundary line analysis uses a subset of data to estimate, in this case, an upper limit, the data set used by Feng et al. (2013) had a wider spread of measured fine fraction OC of 0.9 to 71.7 g C kg<sup>-1</sup> soil, compared to 1.72 to 18.29 g C kg<sup>-1</sup> soil in our UK soils. Therefore, the upper subset of data was composed of higher values giving a steeper slope and demonstrates that the C sequestration estimate generated by boundary line analysis is biased by the range of data.

The strength of using quantile regression analysis is that it makes no assumptions of homogeneity of variance and uses the entire data set to estimate the upper limit of a response. The measured fine fraction OC in the UK sites lacks homogeneity of variance (Fig. 4), where the variation in the measured fine fraction OC increases with the proportion of fine fraction. Therefore, of the methods explored in this study for our grassland soils, we consider the quantile regression at the 90<sup>th</sup> percentile estimate of maximum fine fraction OC to be the most robust. This method results in the greatest number of unsaturated samples (Table 4) suggesting great potential for additional sequestration. When examining the estimated OC input versus existing fine fraction OC using estimates generated by quantile regression at the 90<sup>th</sup> percentile a positive correlation between current fine fraction OC and estimated C input (Kendall tau ( $\tau$ ); 0.323,  $P < 0.001$ ), was observed for the entire data set. However, this was not the case at the site level (Fig. A1). Where in some instances increasing fine fraction OC (g C kg<sup>-1</sup> soil) was associated with increased estimated C input until saturation, such as Aberyswyth, Myerscough and Plumpton. Therefore, despite a higher fine fraction OC contents these samples are furthest from saturation. In contrast the opposite was true for Crichton and Hillsborough (and Harpenden, Kirkton and Overton, although not statistically significant) implying that for these sites samples with a higher fine fraction OC are closer to saturation. It is unclear why this is the case particularly as in all sites, bar Harpenden, there is a positive regression between mass proportion of the fine fraction and fine fraction OC (Table A3). Meaning that higher fine fraction OC is also associated with higher mass proportion of the fine soil fraction. It is likely that the OM input to the soils with the higher mass proportion of fine fraction is insufficient to bridge the gap between



240 current and estimated maximum fine fraction OC, as it is not possible to identify any other effect due to pedogenic or environmental conditions measured in this work.

Estimating maximum fine fraction OC on the basis of mass proportion of fine fraction is likely to be an oversimplification of the dynamics of fine fraction OC accrual. Other parameters such as mineralogy, soil microbial community, environmental conditions (e.g precipitation, Table 2) and land management, can significantly influence fine fraction OC stabilisation (Cotrufo et al., 2015; Kallenbach et al., 2016). This work has identified some soil and environmental properties that may play a role in fine fraction OC stabilisation such as median annual temperature, %N and %C in the bulk soil, mean annual rainfall and %N in the fine fraction (Table 2). Warmer median annual temperatures may enhance plant productivity and microbial processing, the by-products of which are important precursors to fine fraction OC (Cotrufo et al., 2013). It would be interesting to know at which point higher temperatures have a deleterious effect on fine fraction OC accumulation. Mean annual rainfall and %N in the fine fraction were negatively correlated to fine fraction OC. It was anticipated that fine fraction OC would be positively correlated with fine fraction N, as nitrogen rich microbial by-products have been found to form new organo-mineral associations onto which OC preferentially binds (Kopittke et al., 2018). These bonds may have been disturbed during the fractionation process, resulting in an N rich fine fraction with less OC content.

The influence of soil type of fine fraction OC was also evident in our results as all soil types had statistically significant positive correlations between the mass proportion of fine fraction and measured fine fraction OC, except for leptosols and luvisols (Fig. 2). However, these soil types exhibited the greatest proportion of total SOC stabilised in the fine fraction (Fig.3). Luvisols have a high base saturation facilitating more fine fraction OC stabilisation via complexation of organic ligands by free  $\text{Ca}^{2+}$  (Chen et al., 2020). Identifying soils where a greater proportion of total SOC is stored in the fine fraction is important to identify where fine fraction OC needs to be protected, but also where it can be enhanced.

260 Whilst we consider the quantile regression at the 90th percentile method to provide the most robust estimate of maximum fine fraction OC in the sites studied, further experimental work to test the saturation level of these soils, would help to validate this. Incubation studies which force an unsaturated soil to its 'saturation' level and the effect of influencing variables, mentioned above will help to elucidate the factors controlling fine fraction OC saturation. In particular further empirical evidence of how to manipulate fine fraction OC stabilisation processes in a way that is practical for grassland management to promote the formation of new organo-mineral associations, and understanding their stability will be important for establishing the true potential of additional carbon sequestration across managed grasslands.

### 4.3 Effect of sward age on fine fraction OC

It was anticipated that for fields of an older sward age, a greater proportion of total SOC would be stabilised as fine fraction OC, as tillage breaks up macroaggregates making OC in the fine fraction available for mineralisation. Alternatively, fine

fraction OC is less sensitive to disturbance than particulate organic matter (POM), resulting in the accumulation of POM as the fine fraction OC pool remains stable, if sufficiently saturated. The results seem to support neither hypothesis. The proportion of total SOC stabilised in the fine fraction was not consistently higher in the oldest field, and in some instances was significantly less, such as Aberystwyth (Table A2). When grouped in five year intervals, significant differences in C:N ratio of the fine fraction, the proportion of fine fraction in a sample (%) by mass, measured fine fraction OC ( $\text{g C kg}^{-1}$  soil) and the relative proportion of measured fine fraction OC of the total SOC content of the bulk soil were found between age groups (Table 5), however, there was no consistent trend in the results. This data does not support the hypothesis that older swards will have a greater proportion of SOC stabilised in the fine soil fraction, and a reduced potential for additional C sequestration. Equally there was no negative correlation between sward age and the proportion of total SOC stabilised which would be supportive of the alternate hypothesis. From the data it appears that fine fraction OC makes up a greater proportion of SOC with increasing sward age when comparing the less than 5 years, 6 to 10 and 11 to 15 years age groups. However there is a significant decrease in the amount of SOC that is stabilised in the fine fraction in the 16 to 20 years group, this is likely due to fields in this age range originating from Crichton, Hillsborough and Plumpton, which have some of the lowest mass proportion of the fine fraction (Figure 1, C). The sward age analysis may also be confounded by the variation of the proportion of fine fraction, particularly on soil properties influenced by mass proportion of fine fraction such as %C and %N and current fine fraction OC ( $\text{g C kg}^{-1}$  soil). However, it was not possible to conduct robust ANCOVA's with a grouping variable with more than two levels. It may be possible to elucidate the relationship better from a wider study with more samples per age group as our 16 to 20 years group only has 9 values compared to 48 in the less than 5 years group.

Fine fraction OC only accounted for 4.5 to 50.12% indicating high OC storage in other soil pools such as POM, or different aggregate fractions. The fine roots of grassland flora species promote aggregate formation (O'Brien and Jastrow, 2013; Rasse et al., 2005), which may be a dominant stabilisation process in grasslands. However previous work has found no effect of sward age or the frequency of grassland reseeded on the % C in differing aggregate fractions ( $> 2000 \mu\text{m}$ ,  $250\text{--}2000 \mu\text{m}$ ,  $53\text{--}250 \mu\text{m}$  and  $< 53 \mu\text{m}$ ) (Carolan and Fornara, 2016; Fornara et al., 2020). The impact of reseeded disturbance may be offset due to the high density of roots in grasslands by facilitating aggregate reformation. Additionally, dissolved organic carbon from below ground inputs is more efficiently stabilised in organo-mineral associations than above ground dissolved organic carbon (litter leachate) (Sokol and Bradford, 2019). The narrow rhizosphere to bulk soil ratio in grasslands, may make the fine fraction OC in grasslands more resilient to disturbance events.

## 5. Conclusions

Estimating the long-term sequestration of soil C in the fine fraction is difficult due to the lack of reliable methodologies that can be widely applied to all soils. Our study has demonstrated that the Hassink (1997) linear regression equation is not suitable

to estimate maximum fine fraction OC in a range of UK grassland soils. Therefore, caution should be applied to estimates of maximum fine fraction OC obtained using the Hassink (1997) equation, in instances where it may not accurately reflect fine fraction OC of the soil *in situ*. After exploring various univariate estimation methods, we recommend the use of quantile regression at the 90<sup>th</sup> percentile to overcome the shortfalls of linear regression. However, such a simple estimate is unlikely to accurately reflect the dynamics of fine fraction OC stabilisation. This work has helped to identify some key parameters which play a role in fine fraction OC stabilisation, such median annual temperature, mean annual precipitation, bulk soil %C and %N and fine fraction %N. Further work to understand how these parameters influence fine fraction OC dynamics, will help to accurately assess the feasibility of achieving soil carbon sequestration targets. Our results showed little evidence of the impact of time since last reseeded event on the OC in the fine soil fraction. However, improving our understanding of SOC stabilisation processes, and their resilience to grassland management is essential to ensure that current SOC is not only enhanced but also protected.

### **Author contribution**

KCP, SB, JMC, RMR and EMB formulated the research question and study design. KCP conducted the experimental work, data analysis, and prepared the manuscript draft. All authors contributed to editing and reviewing of the manuscript.

### **315 Data Availability**

All data resulting from this study are available from the authors upon request to Sarah Buckingham ([sarah.buckingham@sruc.ac.uk](mailto:sarah.buckingham@sruc.ac.uk))

### **Competing Interest**

The authors declare that they have no conflict of interest.

### **320 Acknowledgements**

We acknowledge funding from Business Environment, Industry and Strategy, Ricardo AEA, Scottish Government's Strategic Research Programme and the Global Academy of Agriculture and Food Security, University of Edinburgh. We are grateful to John Parker and Lydia Guo for their assistance in both the field and laboratory. We also thank Mr Steve Freeman for technical assistance, and Margaret Glendinning and Sarah Perryman for access to information and data from the Electronic Rothamsted

325 Archive (e-RA). The Rothamsted Long-term Experiments National Capability (LTE-NCG) is supported by the UK  
Biotechnology and Biological Sciences Research Council (BBS/E/C/000J0300) and the Lawes Agricultural Trust. The authors  
are also grateful to collaborators at the local field sites for supplying soil samples and management information.

### 330 **References**

- Angers, D. A., Arrouays, D., Saby, N. P. A. and Walter, C.: Estimating and mapping the carbon saturation deficit of French agricultural topsoils, *Soil Use Manag.*, 27(4), 448–452, doi:10.1111/j.1475-2743.2011.00366.x, 2011.
- Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Organic Geochemistry*, 31, 697–710., *Org. Geochem.*, 31, 697–710, doi:10.1016/S0146-6380(00)00049-8,  
335 2000.
- Beare, M. H., McNeill, S. J., Curtin, D., Parfitt, R. L., Jones, H. S., Dodd, M. B. and Sharp, J.: Estimating the organic carbon stabilisation capacity and saturation deficit of soils: A New Zealand case study, *Biogeochemistry*, 120(1–3), 71–87, doi:10.1007/s10533-014-9982-1, 2014.
- Cade, B. S. and Noon, B. R.: A gentle introduction to quantile regression for ecologists, *Front. Ecol. Environ.*, 1(8), 412–420,  
340 doi:10.1890/1540-9295(2003)001[0412:AGITQR]2.0.CO;2, 2003.
- Carolan, R. and Fornara, D. A.: Soil carbon cycling and storage along a chronosequence of re-seeded grasslands: Do soil carbon stocks increase with grassland age?, *Agric. Ecosyst. Environ.*, 218, 126–132, doi:10.1016/j.agee.2015.11.021, 2016.
- Chen, J., Xiao, W., Zheng, C. and Zhu, B.: Nitrogen addition has contrasting effects on particulate and mineral-associated soil organic carbon in a subtropical forest, *Soil Biol. Biochem.*, 142, 107708, doi:10.1016/j.soilbio.2020.107708, 2020.
- 345 Chen, S., Arrouays, D., Angers, D. A., Martin, M. P. and Walter, C.: Soil carbon stocks under different land uses and the applicability of the soil carbon saturation concept, *Soil Tillage Res.*, 188(September 2018), 53–58, doi:10.1016/j.still.2018.11.001, 2019.
- Cotrufo, F. M., Ranalli, M. G., Haddix, M. L., Six, J. and Lugato, E.: Soil carbon storage informed by particulate and mineral-associated organic matter, *Nat. Geosci.*, 12, 989–994, doi:10.1038/s41561-019-0484-6, 2019.
- 350 Cotrufo, M. F., Soong, J. L., Horton, A. J., Campbell, E. E., Haddix, M. L., Wall, D. H. and Parton, W. J.: Formation of soil organic matter via biochemical and physical pathways of litter mass loss, *Nat. Geosci.*, 8(10), 776–779, doi:10.1038/ngeo2520, 2015.
- Dhillon, G. S., Amichev, B. Y., de Freitas, R. and Van Rees, K.: Accurate and Precise Measurement of Organic Carbon Content

- in Carbonate-Rich Soils, *Commun. Soil Sci. Plant Anal.*, 46(21), 2707–2720, doi:10.1080/00103624.2015.1089271, 2015.
- 355 Drewer, J., Anderson, M., Levy, P. E., Scholtes, B., Helfter, C., Parker, J., Rees, R. M. and Skiba, U. M.: The impact of ploughing intensively managed temperate grasslands on N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> fluxes, *Plant Soil*, 411(1–2), 193–208, doi:10.1007/s11104-016-3023-x, 2017.
- Feng, W., Plante, A. F. and Six, J.: Improving estimates of maximal organic carbon stabilization by fine soil particles, *Biogeochemistry*, 112(1–3), 81–93, doi:10.1007/s10533-011-9679-7, 2013.
- 360 Fornara, D., Olave, R. and Higgins, A.: Evidence of low response of soil carbon stocks to grassland intensification, *Agric. Ecosyst. Environ.*, doi:10.1016/j.agee.2019.106705, 2020.
- Gulde, S., Chung, H., Amelung, W., Chang, C. and Six, J.: Soil Carbon Saturation Controls Labile and Stable Carbon Pool Dynamics, *Soil Sci. Soc. Am. J.*, 72(3), 605, doi:10.2136/sssaj2007.0251, 2008.
- Hassink, J.: The capacity of soils to preserve organic C and N by their association with clay and silt particles, *Plant Soil*, 365 191(1), 77–87, doi:10.1023/A:1004213929699, 1997.
- Jackson, M. L.: *Soil chemical analysis: advanced course*, University of Wisconsin, Wisconsin, USA., 2005.
- Kallenbach, C. M., Frey, S. D. and Grandy, A. S.: Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls, *Nat. Commun.*, 7, 1–10, doi:10.1038/ncomms13630, 2016.
- Kleber, M., Sollins, P. and Sutton, R.: A conceptual model of organo-mineral interactions in soils: Self-assembly of organic 370 molecular fragments into zonal structures on mineral surfaces, *Biogeochemistry*, 85(1), 9–24, doi:10.1007/s10533-007-9103-5, 2007.
- Koenker, R.: *quantreg: Quantile Regression*, [online] Available from: <https://cran.r-project.org/package=quantreg>, 2019.
- Lal, R.: Agricultural activities and the global carbon cycle, *Nutr. Cycl. Agroecosystems*, 70, 103–116, 2004.
- Lark, R. M. and Milne, A. E.: Boundary line analysis of the effect of water-filled pore space on nitrous oxide emission from 375 cores of arable soil, *Eur. J. Soil Sci.*, 67(2), 148–159, doi:10.1111/ejss.12318, 2016.
- Lenth, R. V.: Least-Squares Means: The R Package {lsmeans}, *J. Stat. Softw.*, 69(1), 1–33, doi:10.18637/jss.v069.i01, 2016.
- Liang, A., Yang, X., Zhang, X., McLaughlin, N., Shen, Y. and Li, W.: Soil organic carbon changes in particle-size fractions following cultivation of Black soils in China, *Soil Tillage Res.*, 105(1), 21–26, doi:10.1016/j.still.2009.05.002, 2009.
- Lilly, A. and Bagdaley, N. J.: The potential for Scottish cultivated topsoils to lose or gain soil organic carbon, *Soil Use Manag.*, 380 29(1), 39–47, doi:10.1111/sum.12009, 2013.
- Lorenz, K. and Lal, R.: *Carbon Sequestration in Agricultural Soils.*, 2018.
- O’Brien, S. L. and Jastrow, J. D.: Physical and chemical protection in hierarchical soil aggregates regulates soil carbon and nitrogen recovery in restored perennial grasslands, *Soil Biol. Biochem.*, 61, 1–13, doi:10.1016/j.soilbio.2013.01.031, 2013.
- Powlson, D. S., Whitmore, A. P. and Goulding, K. W. T.: Soil carbon sequestration to mitigate climate change: A critical re- 385 examination to identify the true and the false, *Eur. J. Soil Sci.*, 62(1), 42–55, doi:10.1111/j.1365-2389.2010.01342.x, 2011.

- Rasse, D. P., Rumpel, C. and Dignac, M. F.: Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation, *Plant Soil*, 269(1–2), 341–356, doi:10.1007/s11104-004-0907-y, 2005.
- Schmidt, U., Thoni, H. and Kaupenjohann, M.: Using a boundary line approach to analyze N<sub>2</sub>O flux data from agricultural soils, *Nutr. Cycl. Agroecosystems*, 57, 119–129, doi:10.1201/9780203739310, 2000.
- 390 Shatar, T. M. and Mcbratney, A. B.: Boundary-line analysis of field-scale yield response to soil properties, *J. Agric. Sci.*, 142(5), 553–560, doi:10.1017/S0021859604004642, 2004.
- Six, J., Conant, R. T., Paul, E. A. and Paustian, K.: Stabilization mechanisms of SOM implications for C saturation of soils, *Plant Soil*, 241(2), 155–176, doi:10.1023/A:1016125726789, 2002.
- Smith, P.: Soils and climate change, *Curr. Opin. Environ. Sustain.*, 4, 539–544, doi:10.1007/978-94-017-7453-6\_10, 2012.
- 395 Smith, P.: Do grasslands act as a perpetual sink for carbon?, *Glob. Chang. Biol.*, 20(9), 2708–2711, doi:10.1111/gcb.12561, 2014.
- Sokol, N. W. and Bradford, M. A.: Microbial formation of stable soil carbon is more efficient from belowground than aboveground input, *Nat. Geosci.*, 12(1), 46–53, doi:10.1038/s41561-018-0258-6, 2019.
- Soussana, J.-F., Soussana, J.-F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T. and Arrouays, D.: Carbon  
400 cycling and sequestration opportunities in temperate grasslands, *Soil Use Manag.*, 20(2), 219–230, doi:10.1079/SUM2003234, 2004.
- Stewart, C. E., Paustian, K., Conant, R. T., Plante, A. F. and Six, J.: Soil carbon saturation: Concept, evidence and evaluation, *Biogeochemistry*, 86(1), 19–31, doi:10.1007/s10533-007-9140-0, 2007.
- Stewart, C. E., Paustian, K., Conant, R. T., Plante, A. F. and Six, J.: Soil carbon saturation: Implications for measurable carbon  
405 pool dynamics in long-term incubations, *Soil Biol. Biochem.*, 41(2), 357–366, doi:10.1016/j.soilbio.2008.11.011, 2009.
- Team, R. C.: R: A Language and Environment for Statistical Computing, [online] Available from: <https://www.r-project.org/>, 2019.
- Ward, S. E., Smart, S. M., Quirk, H., Tallwin, J. R. B., Mortimer, S. R., Shiel, R. S., Wilby, A. and Bardgett, R. D.: Legacy effects of grassland management on soil carbon to depth, *Glob. Chang. Biol.*, 22(8), 2929–2938, doi:10.1111/gcb.13246, 2016.
- 410 Wei, T. and Simko, V.: R package “corrplot”: Visualization of a Correlation Matrix, [online] Available from: <https://github.com/taiyun/corrplot>, 2017.
- Wiesmeier, M., Hübner, R., Spörlein, P., Geuß, U., Hangen, E., Reischl, A., Schilling, B., von Lützow, M. and Kögel-Knabner, I.: Carbon sequestration potential of soils in southeast Germany derived from stable soil organic carbon saturation, *Glob. Chang. Biol.*, 20(2), 653–665, doi:10.1111/gcb.12384, 2014.

415

**Table 1 Summary of UK grassland site characteristics.**

Site	Age range (years)	Land Use <sup>a</sup>	Mean Annual Temperature (°C) <sup>b</sup>	Mean Annual Rainfall (mm) <sup>a</sup>	Elevation (m.a.s.l)	WRB Soil Type <sup>c</sup>	Soil Texture <sup>c</sup>
<b>Aberystwyth</b> (52°25'N 04°02'W)	2 to 33	UpG	9.5 to 11	1000	20 to 65	ST, CM	Clay to sandy loam
<b>Crichton</b> (55°02'N 03°35'W)	1 to 20	DP	9.5 to 9.9	1100	5 to 50	CM	Clay loam to sandy loam
<b>Easter Bush</b> (55°51'N 03°52'W)	3 to 6	MG	6 to 9	< 700	215 to 265	GL	Clay loam to sandy loam
<b>Harpenden</b> (51°48'N 00°22'W)	22 to 179	UnG	9.5 to 10.5	700	120 to 130	LV	Silty clay loam
<b>Hillsborough</b> (54°27' N 6°04' W)	1 to 37	DP	8.5 to 10	900	120	CM	Clay loam
<b>Kirkton</b> (56°25'N 04°39'W)	1 to 35	UpG	8 to 9.4	2528	163 to 170	PZ	Clay loam to sandy loam
<b>Llangorse</b> (51°55'N 03°16'W)	2.5 to 25	MG	8 to 10	1000		CM	Loam/ Clay to Silty loam
<b>Myerscough</b> (53°51'N 02°46'W)	2 to 48.4	MG	9 to 10.5	1000	8 to 15	GL	Clay to sandy loam
<b>Overton</b> (51°48'N 02°08'W)	3 to 50	MGO	9 to 11	800	240 to 276	LP	Clay loam to silty loam
<b>Plumpton</b> (50°54'N 00°04'W)	1 to 20	MG	9.5 to 11	800	49 to 85 & 160 to 215	ST	Clay to clay loam, Chalky clay to chalky loam

<sup>a</sup> Land Use; DP; Dairy pasture, MG; Mixed grazing; MGO; Mixed grazing organic, UpG; Upland grazing, UnG; Ungrazed.

420 <sup>b</sup> Mean annual temperature and rainfall estimated from Met Office climatic region summaries, averaged over 1981 to 2010.

<sup>c</sup> World Reference Base (WRB) Soil Type: ST; Stagnosols, CM; Cambisols, GL;Gleysol, LV; Luvisols; PZ; Podzol; LP; Leptosol .Soil type and texture determined from GPS locations and UK Soil Observatory Map viewer.

425 **Table 2. Correlation matrix of Kendal tau ( $\tau$ ) coefficients for bulk and fine fraction (<20  $\mu$ m) soil properties, sward age and known environmental parameters.**

		Bulk Soil									Fine Fraction		
		Temp.	Prec.	Age	%N	%C	C:N	pH	%SC	Fine fraction OC	%N	%C	C:N
	Temp.	1											
	Prec.	-0.05	1										
Bulk soil	Age	0.15	-0.11	1									
	%N	0.23***	-0.07	0	1								
	%C	0.16*	0.06	0.04	0.73***	1							
	C:N	-0.25***	-0.05	0.02	- <sup>a</sup>	- <sup>a</sup>	1						
	pH	0.07	-0.30***	-0.07	-0.04	0.03	0.02	1					
	%SC	0.26***	-0.43***	0.14*	0.12	0.12	-0.01	0.21***	1				
	Fine fraction OC	0.13*	-0.36***	0.1	0.26***	0.27***	0.01	0.12	- <sup>a</sup>	1			
Fine Fraction	%N	-0.32***	0.28***	-0.07	0.17**	0.14*	-0.02	-0.27***	-0.47***	-0.15**	1		
	%C	-0.33***	0.25***	-0.09	0.18**	0.17**	0.06	-0.25***	-0.47***	- <sup>a</sup>	0.87***	1	
	C:N	-0.21***	-0.08	-0.16	0.11**	0.21***	0.30***	-0.05	-0.15*	-0.02	- <sup>a</sup>	- <sup>a</sup>	1

<sup>a</sup> No correlation calculated as one variable used to calculate the other.

Age; years since last reseeding event, Temp; median value from the mean annual temperature range ( $^{\circ}$ C), Prec.; mean annual rainfall (mm), %SC; mass proportion of fine fraction in a sample (%), Fine fraction OC; measured fine fraction OC (g C kg<sup>-1</sup> bulk soil).

430

Level of significance: \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\* $P < 0.0001$

435



**Table 3. Analyses coefficients for the estimation of maximum fine fraction organic carbon by linear regression (LR), linear regression with forced zero intercept (LR\_0), boundary line (BL) and quantile regression (QR). Lettering indicates slopes which were significantly different within a method ( $P < 0.05$ ).**

Method		Slope ( $\pm 1$ SEM)	<i>P</i> slope	Intercept ( $\pm 1$ SEM)	<i>P</i> intercept	RMSE	n	R <sup>2</sup>
LR	Hassink, (1997)	0.37 <sup>a</sup>		4.07			40	
	All UK	0.32 $\pm$ 0.023 <sup>b</sup>	***	2.86 $\pm$ 0.368	***	2.58	129	0.61
LR_0	Hassink, (1997) <sup>a</sup>	0.45 $\pm$ 0.02	***			4.97	40	0.94
	All UK	0.47 $\pm$ 0.017	***			3.13	129	0.85
BL	5% intervals	0.48 $\pm$ 0.058	***			5.89	19	0.79
	10% intervals	0.48 $\pm$ 0.070	***			6.36	15	0.77
	15% intervals	0.56 $\pm$ 0.056	***			4.77	14	0.89
QR	QR ( $\tau = 0.90$ )	0.92 $\pm$ 0.071	***			7.90	129	0.90
	QR ( $\tau = 0.50$ )	0.49 $\pm$ 0.032	***			3.15	129	0.66

RMSE, root mean square error.

Level of significance: \*\*\*  $P < 0.001$

<sup>a</sup>Data extracted from Hassink, (1997) used to generate slope value with forced zero intercept.

450 **Table 4. Carbon saturation ratios calculated from the estimated maximum fine fraction organic carbon by linear regression (LR), linear regression with forced zero intercept (LR\_0), boundary line (BL) and quantile regression (QR). Values < 1 indicate unsaturated, = 1 at saturation and > 1 are oversaturated samples.**

<b>Method</b>		<b>No. of unsaturated samples (n = 129)</b>	<b>Mean ratio</b>	<b>Median</b>
<b>LR</b>	Hassink, (1997)	105	0.77	0.73
	UK	75	0.98	0.94
	UK site specific	71	1	0.99
<b>Forced 0 intercept</b>				
<b>LR_0</b>	Hassink (1997)	30	1.52	1.44
	UK	34	1.47	1.39
	UK site specific	57	1.09	1.04
<b>BL</b>	5%	38	1.42	1.34
	10%	36	1.43	1.35
	15%	50	1.22	1.15
<b>QR</b>	50 <sup>th</sup>	38	1.4	1.32
	90 <sup>th</sup>	99	0.74	0.7

455


460

465

**Table 5. Effect of sward age grouped at five year intervals on selected soil properties. Values are means  $\pm$  standard error of the mean, and different letters indicate age groups which are significantly different ( $P < 0.05$ ), by columns.**

<b>Age</b>	<b>n</b>	<b>C:N</b>	<b>% SC</b>	<b>Fine fraction organic carbon (g C kg<sup>-1</sup> soil)</b>	<b>Fine fraction organic carbon (% of SOC<sub>total</sub>)</b>
0 to 5	48	10.18 $\pm$ 0.15 <sup>a</sup>	10.00 $\pm$ 1.41 <sup>a</sup>	5.68 $\pm$ 0.49 <sup>a</sup>	18.32 $\pm$ 1.5 <sup>ab</sup>
6 to 10	18	9.79 $\pm$ 0.26 <sup>ab</sup>	14.47 $\pm$ 1.69 <sup>b</sup>	8.58 $\pm$ 0.59 <sup>b</sup>	24.94 $\pm$ 1.35 <sup>c</sup>
11 to 15	15	9.33 $\pm$ 0.11 <sup>b</sup>	15.27 $\pm$ 2.98 <sup>ab</sup>	9.17 $\pm$ 1.66 <sup>ab</sup>	20.66 $\pm$ 2.55 <sup>abc</sup>
16 to 20	9	10.41 $\pm$ 0.31 <sup>a</sup>	6.10 $\pm$ 0.77 <sup>a</sup>	4.68 $\pm$ 0.44 <sup>a</sup>	11.54 $\pm$ 1.12 <sup>a</sup>
21+	39	9.50 $\pm$ 0.11 <sup>b</sup>	15.19 $\pm$ 1.69 <sup>b</sup>	7.44 $\pm$ 0.69 <sup>ab</sup>	23.21 $\pm$ 1.94 <sup>bc</sup>


Age; years since last reseeding event, C:N ratio of the fine fraction, %SC; proportion of fine fraction in a sample (%) by mass, 470 Fine fraction organic carbon (% of SOC<sub>total</sub>); relative proportion of measured fine fraction OC of the total SOC content of the bulk soil.



475


480

**Figure 1.** Measured total SOC (g C kg<sup>-1</sup>soil) (A) total fine fraction organic carbon (g C kg<sup>-1</sup>soil) (B), mass proportion of fine fraction (< 20 μm, %) (C), relative proportion of measured fine fraction organic carbon of the total SOC content of the bulk soil (D) and fine fraction C:N ratio (E), for each of the grassland sites; Aberyswyth (A), Crichton (C), Easter Bush (E), Hillsborough (H), Harpenden (Ha), Kirkton (K), Llangorse (L), Myerscough (M), Overton (O) and Plumpton (P). Boxes represent the 25<sup>th</sup> and 75<sup>th</sup> percentile, with lines showing the median value. Whiskers show the lowest and highest values with outliers indicates as crosses (> 1.5 times the interquartile range). Lettering indicates significant differences between soils ( $P < 0.05$ ).



485 **Figure 2. Relationships between mass proportion of the fine fraction (%) and fine fraction organic carbon (g C kg<sup>-1</sup> soil) in the soil types used in this study.**


490



495

Figure 3. Relative proportion of measured measured fine fraction organic carbon of the total SOC content of the bulk soil for the different soil types used in this study. Lettering indicates significant differences at  $P < 0.05$ .

500



505

Figure 4. Measured fine fraction organic carbon ( $\text{g C kg}^{-1}$  soil) in relation to mass proportion of fine fraction of a soil sample (%). Line of best fit represent (A) linear regression method of Hassink, (1997) and data from this study, and (B) boundary line (BL) using 15% intervals, and quantile regression analysis (QR) at 90<sup>th</sup> and 50<sup>th</sup> percentiles.

510

515

**Table A1. Bulk soil properties for each UK site. Values are means of ten replicates in each field,  $\pm$  one standard error of the mean. Except Harpenden where values are means of five replicates per field. Lettering indicates values which are significantly different, within a site ( $P < 0.05$ ).**

Site	Age		pH	C:N	C (g C kg <sup>-1</sup> soil)	C stock (t C ha <sup>-1</sup> )	N stock (t N ha <sup>-1</sup> )
	(year s)	BD <sup>a</sup>					
Aberyswyth	2	1 $\pm$ 0.01 <sup>a</sup>	5.20 $\pm$ 0.05 <sup>a</sup>	9.70 $\pm$ 0.05 <sup>b</sup>	26.95 $\pm$ 0.63 <sup>b</sup>	73.61 $\pm$ 1.73 <sup>b</sup>	7.59 $\pm$ 0.16 <sup>b</sup>
	6	0.98 $\pm$ 0.04 <sup>a</sup>	4.70 $\pm$ 0.04 <sup>bc</sup>	9.68 $\pm$ 0.08 <sup>b</sup>	26.7 $\pm$ 0.82 <sup>b</sup>	73.86 $\pm$ 2.28 <sup>b</sup>	7.62 $\pm$ 0.18 <sup>b</sup>
	11	0.82 $\pm$ 0.05 <sup>b</sup>	5.12 $\pm$ 0.06 <sup>a</sup>	10.46 $\pm$ 0.09 <sup>a</sup>	29.72 $\pm$ 0.83 <sup>b</sup>	76.97 $\pm$ 2.15 <sup>b</sup>	7.36 $\pm$ 0.20 <sup>b</sup>
	31	0.74 $\pm$ 0.05 <sup>b</sup>	4.99 $\pm$ 0.09 <sup>ab</sup>	10.54 $\pm$ 0.22 <sup>a</sup>	29.4 $\pm$ 1.63 <sup>b</sup>	74.23 $\pm$ 4.12 <sup>b</sup>	7.01 $\pm$ 0.31 <sup>b</sup>
	33	0.69 $\pm$ 0.03 <sup>b</sup>	4.18 $\pm$ 0.02 <sup>c</sup>	10.59 $\pm$ 0.10 <sup>a</sup>	38.19 $\pm$ 1.97 <sup>b</sup>	95.67 $\pm$ 4.92 <sup>a</sup>	9.01 $\pm$ 0.40 <sup>a</sup>
Crichton	1	0.92 $\pm$ 0.03	5.14 $\pm$ 0.03 <sup>ab</sup>	12.19 $\pm$ 0.08 <sup>ab</sup>	34.66 $\pm$ 0.66 <sup>a</sup>	82.40 $\pm$ 1.56	6.76 $\pm$ 0.11 <sup>b</sup>
	3	0.99 $\pm$ 0.07	5.65 $\pm$ 0.06 <sup>b</sup>	11.73 $\pm$ 0.11 <sup>bc</sup>	29.94 $\pm$ 1.37 <sup>ab</sup>	74.63 $\pm$ 3.41	6.36 $\pm$ 0.26 <sup>b</sup>
	15	0.93 $\pm$ 0.05	4.77 $\pm$ 0.04 <sup>ac</sup>	9.90 $\pm$ 0.88 <sup>c</sup>	30.85 $\pm$ 3.06 <sup>b</sup>	79.73 $\pm$ 7.91	7.98 $\pm$ 0.23 <sup>a</sup>
	20	0.93 $\pm$ 0.04	4.54 $\pm$ 0.03 <sup>c</sup>	13.21 $\pm$ 0.14 <sup>a</sup>	27.26 $\pm$ 0.87 <sup>b</sup>	66.62 $\pm$ 2.11	5.04 $\pm$ 0.15 <sup>c</sup>
Easter Bush	3	1.02 $\pm$ 0.04 <sup>abc</sup>	5.45 $\pm$ 0.06 <sup>ab</sup>	13.03 $\pm$ 0.11 <sup>bc</sup>	32.46 $\pm$ 1.29 <sup>a</sup>	93.52 $\pm$ 3.72 <sup>a</sup>	7.17 $\pm$ 0.26 <sup>a</sup>
	5	1.19 $\pm$ 0.03 <sup>a</sup>	5.44 $\pm$ 0.06 <sup>ab</sup>	12.84 $\pm$ 0.21 <sup>bc</sup>	26.41 $\pm$ 0.54 <sup>b</sup>	74.45 $\pm$ 1.52 <sup>b</sup>	5.80 $\pm$ 0.10 <sup>bc</sup>
	5	0.84 $\pm$ 0.06 <sup>c</sup>	5.67 $\pm$ 0.04 <sup>a</sup>	11.74 $\pm$ 0.17 <sup>d</sup>	27.50 $\pm$ 1.0 <sup>b</sup>	58.15 $\pm$ 2.12 <sup>c</sup>	4.94 $\pm$ 0.14 <sup>d</sup>
	6	0.96 $\pm$ 0.05 <sup>bc</sup>	5.32 $\pm$ 0.06 <sup>b</sup>	12.45 $\pm$ 0.13 <sup>c</sup>	30.46 $\pm$ 1.93 <sup>ab</sup>	71.16 $\pm$ 4.50 <sup>bc</sup>	5.72 $\pm$ 0.36 <sup>cd</sup>
	6	1.12 $\pm$ 0.05 <sup>ab</sup>	5.81 $\pm$ 0.20 <sup>a</sup>	14.15 $\pm$ 0.13 <sup>a</sup>	28.95 $\pm$ 0.99 <sup>ab</sup>	75.69 $\pm$ 2.59 <sup>b</sup>	5.35 $\pm$ 0.17 <sup>cd</sup>
Harpenden	8	1.12 $\pm$ 0.03 <sup>ab</sup>	4.99 $\pm$ 0.04 <sup>c</sup>	13.43 $\pm$ 0.11 <sup>b</sup>	33.03 $\pm$ 0.50 <sup>a</sup>	89.43 $\pm$ 1.34 <sup>a</sup>	6.66 $\pm$ 0.10 <sup>ab</sup>
	22	1.37 $\pm$ 0.07	7.37 $\pm$ 0.04 <sup>a</sup>	12.09 $\pm$ 0.2	16.06 $\pm$ 0.59 <sup>c</sup>	25.37 $\pm$ 0.93 <sup>c</sup>	3.3 $\pm$ 0.12 <sup>c</sup>
	68	1.12 $\pm$ 0.08	5.85 $\pm$ 0.12 <sup>ab</sup>	12.34 $\pm$ 0.08	19.8 $\pm$ 0.63 <sup>b</sup>	50.49 $\pm$ 1.59 <sup>b</sup>	4.06 $\pm$ 0.13 <sup>b</sup>
Hillsborough	179	1.09 $\pm$ 0.14	5.63 $\pm$ 0.06 <sup>b</sup>	12.8 $\pm$ 0.26	28.7 $\pm$ 1.47 <sup>a</sup>	72.98 $\pm$ 3.74 <sup>a</sup>	5.89 $\pm$ 0.30 <sup>a</sup>
	1	1.79 $\pm$ 0.10	6.31 $\pm$ 0.07 <sup>a</sup>	11.25 $\pm$ 0.12 <sup>ab</sup>	46.68 $\pm$ 2.04	120.16 $\pm$ 5.26 <sup>ab</sup>	10.69 $\pm$ 0.51 <sup>ab</sup>
	7	1.88 $\pm$ 0.08	5.10 $\pm$ 0.04 <sup>b</sup>	11.46 $\pm$ 0.11 <sup>b</sup>	42.85 $\pm$ 1.52	108.86 $\pm$ 3.87 <sup>b</sup>	9.51 $\pm$ 0.34 <sup>bc</sup>
	16	1.79 $\pm$ 0.05	5.33 $\pm$ 0.08 <sup>b</sup>	10.87 $\pm$ 0.06 <sup>c</sup>	42.36 $\pm$ 1.98	111.63 $\pm$ 5.21 <sup>ab</sup>	10.27 $\pm$ 0.47 <sup>ab</sup>
	23	1.75 $\pm$ 0.05	4.76 $\pm$ 0.03 <sup>c</sup>	11.33 $\pm$ 0.09 <sup>ab</sup>	46.44 $\pm$ 1.78	125.43 $\pm$ 4.82 <sup>a</sup>	11.08 $\pm$ 0.45 <sup>a</sup>
Kirkton	37	1.69 $\pm$ 0.06	5.13 $\pm$ 0.06 <sup>b</sup>	10.34 $\pm$ 0.77 <sup>ac</sup>	40.90 $\pm$ 3.10	86.04 $\pm$ 6.52 <sup>c</sup>	8.38 $\pm$ 0.24 <sup>c</sup>
	1	0.9 $\pm$ 0.04	4.78 $\pm$ 0.04 <sup>c</sup>	12.13 $\pm$ 0.11 <sup>c</sup>	27.90 $\pm$ 0.81 <sup>c</sup>	82.03 $\pm$ 2.39 <sup>b</sup>	6.77 $\pm$ 0.22
	3	0.95 $\pm$ 0.04	5.49 $\pm$ 0.06 <sup>a</sup>	12.61 $\pm$ 0.15 <sup>b</sup>	36.67 $\pm$ 1.56 <sup>a</sup>	98.19 $\pm$ 4.17 <sup>ab</sup>	7.79 $\pm$ 0.31



	5	0.83 ± 0.06	5.15 ± 0.03 <sup>b</sup>	13.56 ± 0.08 <sup>a</sup>	34.83 ± 1.84 <sup>ab</sup>	103.03 ± 5.45 <sup>a</sup>	7.59 ± 0.38
	35	0.97 ± 0.06	4.72 ± 0.07 <sup>c</sup>	11.67 ± 0.13 <sup>d</sup>	30.51 ± 1.48 <sup>bc</sup>	90.50 ± 4.38 <sup>ab</sup>	7.72 ± 0.32
Llangorse	2.5	1.01 ± 0.04	5.14 ± 0.08 <sup>c</sup>	9.21 ± 0.09	17.83 ± 0.42	49.75 ± 1.18	5.40 ± 0.11 <sup>ab</sup>
	5	0.93 ± 0.04	5.44 ± 0.03 <sup>b</sup>	9.40 ± 0.07	18.60 ± 0.45	50.80 ± 1.22	5.40 ± 0.10 <sup>b</sup>
	15	0.94 ± 0.06	5.68 ± 0.03 <sup>a</sup>	9.36 ± 0.17	19.42 ± 0.38	53.70 ± 1.06	5.74 ± 0.12 <sup>ab</sup>
	25	1.06 ± 0.03	5.54 ± 0.07 <sup>ab</sup>	9.16 ± 0.87	19.73 ± 2.52	55.10 ± 7.05	6.18 ± 0.34 <sup>a</sup>
Myerscough	2	1.22 ± 0.02 <sup>ab</sup>	4.97 ± 0.05 <sup>b</sup>	13.58 ± 0.24 <sup>bc</sup>	27.47 ± 0.65 <sup>c</sup>	82.25 ± 1.96 <sup>c</sup>	6.07 ± 0.15 <sup>bc</sup>
	6	1.10 ± 0.04 <sup>b</sup>	5.59 ± 0.05 <sup>a</sup>	11.79 ± 0.76 <sup>c</sup>	41.44 ± 2.73 <sup>a</sup>	124.05 ± 8.17 <sup>a</sup>	10.56 ± 0.28 <sup>a</sup>
	13	0.93 ± 0.05 <sup>b</sup>	5.00 ± 0.20 <sup>b</sup>	13.12 ± 0.43 <sup>c</sup>	44.82 ± 2.34 <sup>a</sup>	134.45 ± 7.01 <sup>a</sup>	10.30 ± 0.71 <sup>a</sup>
	34	1.29 ± 0.02 <sup>a</sup>	5.99 ± 0.13 <sup>a</sup>	17.20 ± 1.12 <sup>ab</sup>	37.58 ± 1.45 <sup>ab</sup>	112.46 ± 4.36 <sup>ab</sup>	6.71 ± 0.30 <sup>b</sup>
	48.4	1.44 ± 0.06 <sup>a</sup>	5.77 ± 0.02 <sup>a</sup>	22.10 ± 1.46 <sup>a</sup>	29.86 ± 1.96 <sup>bc</sup>	88.97 ± 5.85 <sup>bc</sup>	4.03 ± 0.08 <sup>c</sup>
Overton	3	0.98 ± 0.09 <sup>a</sup>	6.58 ± 0.12 <sup>b</sup>	9.76 ± 0.05 <sup>b</sup>	32.77 ± 0.84 <sup>c</sup>	83.02 ± 2.13 <sup>b</sup>	8.51 ± 0.23 <sup>b</sup>
	12	0.38 ± 0.03 <sup>b</sup>	6.83 ± 0.03 <sup>b</sup>	10.18 ± 0.12 <sup>ab</sup>	70.18 ± 1.92 <sup>a</sup>	81.20 ± 2.23 <sup>b</sup>	7.99 ± 0.23 <sup>b</sup>
	22	0.71 ± 0.07 <sup>ab</sup>	7.36 ± 0.04 <sup>a</sup>	10.68 ± 0.39 <sup>a</sup>	59.88 ± 3.86 <sup>b</sup>	132.75 ± 8.56 <sup>a</sup>	12.33 ± 0.39 <sup>a</sup>
	50	1.74 ± 0.9 <sup>a</sup>	4.63 ± 0.08 <sup>c</sup>	10.14 ± 0.14 <sup>ab</sup>	51.18 ± 2.84 <sup>b</sup>	153.08 ± 8.50 <sup>a</sup>	15.08 ± 0.80 <sup>a</sup>
Plumpton	1	0.99 ± 0.02 <sup>a</sup>	6.34 ± 0.08 <sup>b</sup>	10.85 ± 0.08 <sup>ab</sup>	40.92 ± 1.21 <sup>b</sup>	122.21 ± 3.61 <sup>b</sup>	11.26 ± 0.28 <sup>b</sup>
	5	1.08 ± 0.03 <sup>a</sup>	7.15 ± 0.06 <sup>a</sup>	11.27 ± 0.41 <sup>a</sup>	45.55 ± 0.61 <sup>b</sup>	132.09 ± 1.77 <sup>b</sup>	11.87 ± 0.48 <sup>b</sup>
	20	0.72 ± 0.04 <sup>b</sup>	5.38 ± 0.21 <sup>c</sup>	10.54 ± 0.17 <sup>b</sup>	58.08 ± 2.36 <sup>a</sup>	163.23 ± 6.62 <sup>a</sup>	15.47 ± 0.56 <sup>a</sup>

<sup>a</sup>Bulk density (BD), means and SEM of six samples, except Harpenden with two samples per field, corrected for stones.

525

530

535

**Table A2. Fine fraction (<20 µm) soil properties for each UK site. Values are means of three replicates in each field, ± one standard error of the mean. Lettering indicates values which are significantly different, within a site ( $P < 0.05$ ).**

Location	Age (years)	%N	%C	C:N	%Fine Fraction	Organic Carbon (g C kg <sup>-1</sup> bulk soil) <sup>a</sup>	Organic Carbon (% of SOC <sub>total</sub> )
Aberyswyth	2	0.48 ± 0.01 <sup>b</sup>	4.16 ± 0.08	8.62 ± 0.08 <sup>ab</sup>	19.08 ± 1.04 <sup>a</sup>	7.93 ± 0.45 <sup>a</sup>	0.30 ± 0.01 <sup>a</sup>
	6	0.52 ± 0.04 <sup>b</sup>	4.14 ± 0.30	8.05 ± 0.17 <sup>b</sup>	14.47 ± 1.18 <sup>ab</sup>	5.92 ± 0.22 <sup>ab</sup>	0.21 ± 0.02 <sup>ab</sup>
	11	0.55 ± 0.03 <sup>b</sup>	4.89 ± 0.25	8.86 ± 0.06 <sup>ab</sup>	18.02 ± 1.51 <sup>ab</sup>	8.77 ± 0.56 <sup>a</sup>	0.29 ± 0.02 <sup>ab</sup>
	31	0.61 ± 0.05 <sup>ab</sup>	5.78 ± 0.62	9.51 ± 0.30 <sup>ab</sup>	13.78 ± 0.49 <sup>b</sup>	8.02 ± 1.15 <sup>a</sup>	0.27 ± 0.02 <sup>ab</sup>
	33	0.76 ± 0.03 <sup>a</sup>	7.57 ± 0.37	9.96 ± 0.23 <sup>a</sup>	5.02 ± 0.22 <sup>c</sup>	3.81 ± 0.36 <sup>b</sup>	0.10 ± 0.01 <sup>b</sup>
Crichton	1	1.01 ± 0.06	10.53 ± 0.83	10.40 ± 0.23 <sup>ab</sup>	4.00 ± 0.45	4.24 ± 0.69	0.12 ± 0.02
	3	1.15 ± 0.27	11.17 ± 2.28	9.84 ± 0.35 <sup>b</sup>	3.28 ± 0.23	3.75 ± 1.00	0.13 ± 0.04
	15	1.02 ± 0.12	9.76 ± 1.20	9.52 ± 0.12 <sup>b</sup>	3.52 ± 0.26	3.38 ± 0.31	0.10 ± 0.02
	20	0.82 ± 0.05	9.07 ± 0.72	11.03 ± 0.24 <sup>a</sup>	3.37 ± 0.3	3.01 ± 0.09	0.11 ± 0.01
Easter Bush	3	0.65 ± 0.04	7.15 ± 0.50	11.00 ± 0.13 <sup>ab</sup>	14.38 ± 1.56 <sup>ab</sup>	10.27 ± 1.19 <sup>a</sup>	0.30 ± 0.01
	5	0.65 ± 0.04	6.91 ± 0.50	10.57 ± 0.06 <sup>bc</sup>	12.17 ± 0.9 <sup>ab</sup>	8.34 ± 0.43 <sup>ab</sup>	0.32 ± 0.02
	5	0.67 ± 0.02	6.62 ± 0.23	9.83 ± 0.07 <sup>c</sup>	9.55 ± 0.73 <sup>b</sup>	6.32 ± 0.51 <sup>b</sup>	0.23 ± 0.03
	6	0.68 ± 0.03	7.81 ± 0.43	9.85 ± 0.24 <sup>c</sup>	9.75 ± 0.23 <sup>b</sup>	6.88 ± 1.13 <sup>ab</sup>	0.23 ± 0.01
	6	0.72 ± 0.12	7.11 ± 1.31	11.43 ± 0.16 <sup>a</sup>	10.58 ± 1.04 <sup>b</sup>	8.22 ± 0.70 <sup>ab</sup>	0.27 ± 0.02
	8	0.59 ± 0.04	6.07 ± 0.30	10.35 ± 0.28 <sup>bc</sup>	16.47 ± 1.3 <sup>a</sup>	9.91 ± 0.26 <sup>ab</sup>	0.30 ± 0.01
Harpenden	22	0.23 ± 0.01 <sup>b</sup>	1.90 ± 0.04 <sup>c</sup>	8.20 ± 0.26 <sup>b</sup>	36.15 ± 4.77 <sup>a</sup>	6.82 ± 0.75 <sup>ab</sup>	0.42 ± 0.04
	68	0.32 ± 0.01 <sup>b</sup>	3.08 ± 0.06 <sup>b</sup>	9.54 ± 0.21 <sup>a</sup>	22.27 ± 0.92 <sup>b</sup>	6.86 ± 0.28 <sup>ab</sup>	0.36 ± 0.02
	179	0.46 ± 0.03 <sup>a</sup>	4.35 ± 0.36 <sup>a</sup>	9.54 ± 0.12 <sup>a</sup>	20.83 ± 1.64 <sup>b</sup>	9.02 ± 0.91 <sup>a</sup>	0.32 ± 0.01
Hillsborough	1	0.90 ± 0.08	8.97 ± 1.14	9.91 ± 0.34	7.37 ± 0.12	6.86 ± 1.93	0.14 ± 0.03
	7	1.04 ± 0.06	10.23 ± 0.91	9.80 ± 0.31	8.05 ± 0.08	8.15 ± 0.96	0.17 ± 0.02
	16	0.99 ± 0.04	9.36 ± 0.32	9.46 ± 0.03	6.33 ± 0.19	5.92 ± 0.13	0.15 ± 0.01
	23	1.15 ± 0.01	11.11 ± 0.13	9.70 ± 0.18	4.58 ± 0.27	5.10 ± 0.36	0.12 ± 0.01
	37	1.04 ± 0.04	10.12 ± 0.35	9.76 ± 0.04	7.15 ± 0.33	7.22 ± 0.10	0.27 ± 0.11
Kirkton	1	0.91 ± 0.03	9.27 ± 0.12 <sup>b</sup>	10.15 ± 0.24 <sup>b</sup>	3.90 ± 0.1	3.62 ± 0.13	0.14 ± 0.01
	3	1.01 ± 0.04	10.63 ± 0.33 <sup>a</sup>	10.56 ± 0.27 <sup>ab</sup>	3.02 ± 0.03	3.20 ± 0.07	0.08 ± 0.00
	5	0.88 ± 0.03	10.23 ± 0.16 <sup>ab</sup>	11.66 ± 0.31 <sup>a</sup>	4.62 ± 0.95	4.75 ± 1.03	0.13 ± 0.02

	35	0.96 ± 0.03	9.22 ± 0.40 <sup>b</sup>	9.64 ± 0.39 <sup>b</sup>	4.23 ± 0.42	3.93 ± 0.51	0.14 ± 0.00
Llangorse	2.5	0.51 ± 0.03 <sup>b</sup>	4.76 ± 0.29 <sup>b</sup>	9.36 ± 0.07	6.00 ± 0.32 <sup>a</sup>	2.83 ± 0.10 <sup>a</sup>	0.16 ± 0.01 <sup>a</sup>
	5	0.88 ± 0.08 <sup>a</sup>	8.29 ± 0.80 <sup>a</sup>	9.43 ± 0.07	2.65 ± 0.43 <sup>b</sup>	2.13 ± 0.11 <sup>ab</sup>	0.11 ± 0.01 <sup>ab</sup>
	15	0.67 ± 0.10 <sup>ab</sup>	6.06 ± 0.78 <sup>ab</sup>	9.11 ± 0.24	3.23 ± 1.03 <sup>b</sup>	1.81 ± 0.34 <sup>b</sup>	0.09 ± 0.02 <sup>b</sup>
	25	0.62 ± 0.06 <sup>ab</sup>	5.32 ± 0.54 <sup>b</sup>	8.60 ± 0.05	3.27 ± 0.22 <sup>b</sup>	1.72 ± 0.14 <sup>b</sup>	0.07 ± 0.01 <sup>b</sup>
Myerscough	2	0.63 ± 0.08	6.60 ± 0.76	10.43 ± 0.12	5.23 ± 0.66 <sup>c</sup>	3.35 ± 0.09 <sup>c</sup>	0.12 ± 0.00 <sup>b</sup>
	6	0.49 ± 0.03	4.57 ± 0.29	9.31 ± 0.23	27.50 ± 3.85 <sup>a</sup>	12.39 ± 1.24 <sup>a</sup>	0.31 ± 0.04 <sup>a</sup>
	13	0.50 ± 0.05	4.83 ± 0.60	9.52 ± 0.32	30.88 ± 4.39 <sup>a</sup>	14.45 ± 0.92 <sup>a</sup>	0.30 ± 0.02 <sup>a</sup>
	34	0.47 ± 0.01	4.28 ± 0.13	9.12 ± 0.07	18.72 ± 0.04 <sup>ab</sup>	8.02 ± 0.24 <sup>b</sup>	0.21 ± 0.02 <sup>ab</sup>
	48.4	0.47 ± 0.02	4.47 ± 0.36	9.58 ± 0.36	12.08 ± 0.74 <sup>bc</sup>	5.35 ± 0.14 <sup>bc</sup>	0.17 ± 0.02 <sup>b</sup>
Overton	3	0.42 ± 0.03 <sup>c</sup>	3.57 ± 0.31 <sup>c</sup>	8.45 ± 0.18 <sup>b</sup>	38.65 ± 6.58	13.57 ± 1.77	0.41 ± 0.05
	12	0.88 ± 0.05 <sup>a</sup>	8.52 ± 0.59 <sup>a</sup>	9.64 ± 0.11 <sup>a</sup>	20.70 ± 2.41	17.45 ± 1.75	0.25 ± 0.02
	22	0.61 ± 0.02 <sup>b</sup>	6.36 ± 0.18 <sup>b</sup>	10.36 ± 0.15 <sup>a</sup>	19.85 ± 4.39	12.52 ± 2.50	0.23 ± 0.07
	50	0.63 ± 0.05 <sup>b</sup>	6.23 ± 0.29 <sup>b</sup>	10.04 ± 0.45 <sup>a</sup>	29.50 ± 2.23	18.29 ± 0.86	0.34 ± 0.04
Plumpton	1	0.35 ± 0.02 <sup>b</sup>	3.81 ± 0.18 <sup>b</sup>	10.87 ± 0.09	19.50 ± 5.61	7.23 ± 1.74	0.18 ± 0.05
	5	0.36 ± 0.04 <sup>b</sup>	4.19 ± 0.49 <sup>b</sup>	11.76 ± 0.91	6.60 ± 2.08	2.56 ± 0.49	0.06 ± 0.01
	20	0.56 ± 0.02 <sup>a</sup>	5.96 ± 0.23 <sup>a</sup>	10.75 ± 0.62	8.60 ± 0.3	5.11 ± 0.21	0.08 ± 0.01

%Fine Fraction; mass proportion of fine fraction in a sample (%).

540 <sup>a</sup> MAOC (g C kg<sup>-1</sup> bulk soil) accounts for the proportion of fine fraction per kilogram of bulk soil.

545

550 **Table A3. Linear regression coefficients for the estimation of maximum fine fraction organic carbon (g C kg<sup>-1</sup> soil). Lettering indicates slopes which are significantly different ( $P < 0.05$ ).**

Site	Slope ( $\pm 1$ SEM)	<i>P</i> slope	Intercept ( $\pm 1$ SEM)	<i>P</i> intercept	RMSE	n	R <sup>2</sup>
<b>Aberyswyth</b>	0.33 $\pm$ 0.059 <sup>bc</sup>	***	2.28 $\pm$ 0.892	*	1.11	15	0.70
<b>Crichton</b>	1.14 $\pm$ 0.470 <sup>abcd</sup>	*	-0.44 $\pm$ 1.684	Ns	0.79	12	0.37
<b>Easter Bush</b>	0.49 $\pm$ 0.094 <sup>d</sup>	***	2.33 $\pm$ 1.172	Ns	1.10	18	0.63
<b>Harpenden</b>	-0.02 $\pm$ 0.07 <sup>a</sup>	Ns	8.01 $\pm$ 1.837	**	1.42	9	0.01
<b>Hillsborough</b>	0.97 $\pm$ 0.148 <sup>d</sup>	***	0.16 $\pm$ 1.02	Ns	0.84	15	0.77
<b>Kirkton</b>	1.01 $\pm$ 0.088 <sup>abcd</sup>	***	-0.11 $\pm$ 0.357	Ns	0.26	12	0.93
<b>Llangorse</b>	0.29 $\pm$ 0.055 <sup>abc</sup>	***	1.03 $\pm$ 0.225	***	0.27	12	0.73
<b>Mysercough</b>	0.40 $\pm$ 0.031 <sup>bcd</sup>	***	1.07 $\pm$ 0.669	Ns	1.14	15	0.93
<b>Overton</b>	0.12 $\pm$ 0.109 <sup>cd</sup>	Ns	12.16 $\pm$ 3.142	**	3.35	12	0.11
<b>Plumpton</b>	0.30 $\pm$ 0.042 <sup>ab</sup>	***	1.45 $\pm$ 0.573	*	0.82	9	0.88

RMSE: Root mean square error.

Level of significance: \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\* $P < 0.001$

555


560

**Table A4. Linear regression coefficients for the estimation of maximum fine fraction organic carbon (g C kg<sup>-1</sup> soil) with a forced zero intercept. Lettering indicates slopes, that are significantly different ( $P < 0.05$ ).**

<b>Site</b>	<b>Slope (<math>\pm 1</math> SEM)</b>	<b><i>P</i></b>	<b>RMSE</b>	<b>n</b>	<b>R<sup>2</sup></b>
<b>Aberyswyth</b>	0.47 $\pm$ 0.024 <sup>bc</sup>	***	1.357	15	0.96
<b>Crichton</b>	1.02 $\pm$ 0.067 <sup>cdef</sup>	***	0.796	12	0.95
<b>Easter Bush</b>	0.67 $\pm$ 0.024 <sup>e</sup>	***	1.231	18	0.98
<b>Harpenden</b>	0.26 $\pm$ 0.035 <sup>a</sup>	***	2.739	9	0.87
<b>Hillsborough</b>	0.99 $\pm$ 0.033 <sup>f</sup>	***	0.842	15	0.99
<b>Kirkton</b>	0.98 $\pm$ 0.0197 <sup>def</sup>	***	0.265	12	0.99
<b>Llangorse</b>	0.52 $\pm$ 0.035 <sup>abcdef</sup>	***	0.474	12	0.95
<b>Mysercough</b>	0.45 $\pm$ 0.016 <sup>b</sup>	***	1.255	15	0.98
<b>Overton</b>	0.52 $\pm$ 0.055 <sup>bcd</sup>	***	5.297	12	0.89
<b>Plumpton</b>	0.39 $\pm$ 0.030 <sup>ab</sup>	***	1.141	9	0.96

RMSE: Root mean square error.

Level of significance: \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\* $P < 0.001$



570 **Figure A1.** Estimated fine fraction OC input (g C kg<sup>-1</sup>soil) compared to measured fine fraction OC (g C kg<sup>-1</sup>soil) in each of the sites studied. The estimated OC input was predicted using quantile regression at the 90<sup>th</sup> percentile.