
1 SUPPLEMENTAL INFORMATION

2 **Reviews and syntheses**

3 **The biogeochemical cycle of Si in the modern ocean**

4 Paul J. Tréguer, Jill N. Sutton, Mark Brzezinski, Matthew A. Charette, Timothy Devries,
5 Stephanie Dutkiewicz, Claudia Ehlert, Jon Hawkings, Aude Leynaert, Su Mei Liu, Natalia
6 Llopis Monferrer, María López-Acosta, Manuel Maldonado, Shaily Rahman, Lihua Ran,
7 Olivier Rouxel

8
9
10
11 This document complements the review article. It provides detailed legends for Figures 1, 2 and
12 4, and a few additional comments to the main text. Annex 1 shows data for the determination
13 of biogenic silica (bSi) production measured by isotopic techniques.

14 15 **Section 1. Introduction**

16 **Detailed legend of Fig. 1 and flux abbreviations**

17 Schematic view of the Si cycle in the modern world ocean (input, output, and biological Si
18 fluxes), and possible balance (total Si inputs = total Si outputs = 14.8 Tmol-Si yr⁻¹) that is in
19 reasonable agreement with the individual range of each flux (F), see Tables 1 and 2. All fluxes
20 are in Tmol-Si yr⁻¹.

21 *-Inputs:* Rivers: $F_{R(dSi+aSi)} = 8.1$; Aeolian inputs: $F_A = 0.5$; Glacial meltwater: $F_{ISMW} = 0.3$;
22 Submarine groundwater: $F_{GW} = 2.3$; Dissolution of minerals: $F_W = 1.9$; Hydrothermal: $F_H =$
23 1.7

24 *-Outputs:* Burial: $F_{B(\text{net deposit})} = 7.4$; Sponges: $F_{SP} = 1.7$; Reverse weathering: $F_{RW} = 5.7$.

25 *-Biological and other fluxes:* Uptake (pelagic production) = $F_{P_{gross}} = 255$; Recycling (surface)
26 $F_{D(surface)} = 143$ (D: P = 0.56 according to Tréguer and De La Rocha 2013); Export: $F_E = 112$;
27 Recycling (deep) $F_{D(deep)} = 28.8$ ($F_{D(deep)}/F_E=0.25$ according to Tréguer and De La Rocha 2013);
28 Rain = $F_{S(rain)} = 84.0$; Recycling (sediment-water interface) $F_{D(benthic)} = 76.6$; upwelling,
29 diffusion: $F_{upw/ed} = 104.6$.

30 *Flux abbreviations:* *dSi input fluxes:* F_R total river net discharge, F_A dissolution of aeolian -
31 transported siliceous dusts, F_{GW} submarine groundwater discharge, F_W dissolution of siliceous
32 material transported from land on the continental margins, and of basalt, F_H hydrothermal activity
33 of the oceanic ridges (axis + off axis), F_{ISMW} ice shelf melt water flux (subglacial melt water +
34 basal melting of ice shelves + melting of icebergs);

35 *Si output fluxes:* $F_{B(\text{net deposit})}$ long-term burial of biogenic silica, F_{SP} siliceous sponges, F_{RW}
36 reverse weathering by formation of authigenic silicate minerals;

37 *Biological fluxes:* $F_{P\text{gross}}$ production of biogenic silica due to diatoms and other pelagic
38 silicifiers, $F_{E(\text{export})}$ export flux of biogenic silica to the deep reservoir, $F_{S(\text{rain})}$ part of the export
39 flux that reaches the sediment - water interface, $F_{B(\text{net deposit})}$ long-term accumulation of biogenic
40 opal in coastal and abyssal, sediments.

41 Note that the dSi uptake due to benthic organisms is not represented on Fig. 1. Indeed, the total
42 bSi production of benthic diatoms is presently unknown but it should be $< 5 \text{ Tmol-Si yr}^{-1}$
43 (Leynaert et al., work in progress), and that the total bSi production of sponges ($6.1 (\pm 5.9)$
44 Tmol-Si yr^{-1}) is still preliminary (Maldonado et al., work in progress).

45 *Other fluxes:* $F_{D(\text{surface})}$ recycling of Si by dissolution of the biogenic silica in the surface
46 reservoir, $F_{D(\text{deep})}$ part of the export flux that dissolves in the deep reservoir, $F_{\text{upw/ed}}$ transfer of
47 dSi from the deep to the surface reservoir by upwelling or eddy diffusion. $F_{D(\text{benthic})}$, flux at the
48 sediment - water interface, is according to Tréguer and De La Rocha 2013).

49

50 **Section 2 -The input fluxes**

51 **2.1 Detailed legend of Figure 2.**

52 Schematic view of the low temperature processes that control the dissolution of (either
53 amorphous or crystallized) siliceous minerals in seawater in the coastal zone and in the deep
54 ocean, feeding F_{GW} and F_w . These processes (white arrows) correspond to low or medium
55 energy flux dissipated per volume of a given siliceous particle in the coastal zone, in the
56 continental margins, and in the abysses, and to high kinetic energy flux dissipated in the surf
57 zone. Inputs of siliceous biogenic and lithogenic silica into the ocean are mainly due to
58 suspended matter transferred from the continent. Rivers support the main transfer of Si to the
59 coastal zone, either as dSi or as aSi. Abundant transfer of siliceous suspended matter into the
60 ocean is also expected from river mouths and deltas through dissemination of suspended matter
61 in the coastal zone, in the continental margin, and beyond to the abysses. In sandy and
62 permeable soil zones dSi is also transferred from the continent to the coastal zone through
63 submarine groundwater discharge processes either as net fresh water inputs or as
64 brackish/seawater recycling due to tidal pumping. In the surf zone, low temperature dissolution
65 of grains of lithogenic silica (quartz, feldspar, etc.) could be intense under the pressure of the
66 intensive and continuous shaking due to waves. For the sake of clarity, we distinguish the
67 processes at work in a quiet zone that receive river inputs from those occurring in a sand beach
68 zone subject to strong dissipation of energy due to wave motion. Note that neither the low-

69 temperature dissolution of wind-borne siliceous material (F_A) nor that occurring in high-
70 temperature hydrothermal systems (F_H) are represented on this figure.

71 **2.2 Dissolution of minerals**

72 Regarding the marine component of F_W dissolution of minerals, Fabre et al. (2019) (main text
73 reference list), focused on wave and tidal action prevailing in the intertidal surf zone (Fig. 2).
74 From laboratory experiment with pure quartz, they showed that quartz grains submitted to
75 violent agitation are capable of substantial dissolution of silica at time scale of days. According
76 to these authors, the flux of dissolution of siliceous material from sandy beaches is $F_{\text{dissolution}} =$
77 $k(T^\circ\text{C}) \times S_{\text{reactive}} \times (C^* - C_{\text{sw}})$. It corresponds to a net input of dSi to the ocean. In this equation,
78 k ($f(T^\circ\text{C})$) in m s^{-1} is the mass transfer coefficient, S_{reactive} in m is the reactive surface of sand
79 grains, C^* in mol-Si L^{-1} is the temperature-dependent solubility limit of sand at thermodynamic
80 equilibrium, and C_{sw} , in mol-Si L^{-1} is the coastal seawater silicic acid concentration. Assuming
81 that all sandy beaches is composed of quartz, Fabre et al. (2019) calculated a global flux of 3.2
82 (± 1.0) Tmol Si yr^{-1} . However, this estimate is not well constrained.

83 Firstly, it is clear that the mineral composition of the world ocean beaches is not entirely
84 composed of quartz. Indeed, the composition of sandy beaches is variable and represent 31%
85 of the coastline of the continents at world scale (Luijendijk et al, 2018, main text reference list).
86 Sandy beaches are composed both of non-siliceous materials (mostly calcareous), and diverse
87 siliceous types of materials (amorphous silica, quartz, feldspars, clays, etc.). These siliceous
88 materials have different solubility and dissolution rates in seawater (S1: Lerman et al., 1975;
89 S2: Hurd et al., 1979), and they can be more or less coated by organic and metals, which affect
90 silica dissolution (e.g. S3: Loucaides et al. 2010; S4: Wiley, 1975). These differing sand
91 compositions directly affects the value of S_{reactive} as well as that of C^* .

92 Secondly, in surf zones the mixing conditions of sand and seawater can be very variable, both
93 over time and at local and regional scale, thus affecting the solid to liquid phase ratio, and the
94 values of the C^* .

95 Thirdly, Fabre et al (2019)'s value for C_{sw} (i.e. 85.4 μM), is far too high for coastal waters,
96 irrespective of its regional context in the world ocean.

97 Finally, C_{sw} can be seasonally variable, particularly in temperate zones.

98 Also note that, by definition, this flux is already included in the marine SGD estimate according
99 to Cho et al. (2018) (main text reference list)

100 *REFERENCES:*

101 S1. Lerman, A., Mackenzie F.T., Bricker, O.P. Rates of dissolution of aluminosilicates in seawater, *EPSL*, 25: 82-
102 88.

103 S2. Hurd, D.C., Fraley, C., Fugate J.K. Silica apparent solubilities and rates of dissolution and precipitation (1979)
104 DOI: 10.1021/bk-1979-0093.ch021

105 S3: Loucaides et al. Seawater-mediated interactions between diatomaceous silica and terrigenous sediments:
106 Results from long-term incubation experiments, *Mar. Chem.*, 3, 241-251.

107 S4: Wiley, J.D. Rates of dissolution of aluminosilicates in seawater. *Mar. Chem.*, 3, 241-251.

108

109 **Section 3 -The output fluxes**

110 **Deposition of sponge silica in marine sediments**

111 The annual rate of sponge silica deposition to the sediments, unlike in diatoms, cannot be easily
112 calculated from annual production rates. The longevity of sponges, which ranges from months
113 or years to even centuries or millennia (S5: Elwood et al., 2007; S6 : McMurray et al., 2008;
114 Jochum et al. 2017, main text reference list), decouples the process of skeleton production —
115 which slowly accumulates bSi over the sponge lifespan — from the process of releasing the
116 accumulated bSi into the sediments — which occurs within weeks to months after sponge death.
117 The deposition of sponge bSi is also decoupled from the rain of planktonic, and therefore, from
118 the rate of sediment deposition. The reason of this is in the benthic nature of sponges. Sponges
119 live already attached to the bottom, so their bSi does no transit through the water column once
120 sponges die. When the flesh of the body gets progressively rotten, the spicules (i.e. skeletal
121 pieces of sponges) fall directly on the seafloor at the site where the sponge was growing forming
122 a spicule patch (S7: Laguionie-Marchais et al., 2015), which can be slowly disaggregated over
123 time by the action of scavengers and other benthic macrofauna, also by the action of bottom
124 currents, turbidity currents included.

125 **REFERENCES:**

126 S5. Ellwood, M. J., M. Kelly, and B. Richer De Forges. Silica banding in the deep-sea lithistid sponge
127 *Corallistes undulatus*: Investigating the potential influence of diet and environment on growth. *Limnol.*
128 *Oceanogr.* 52: 1865-1873 (2007).

129 S6. McMurray, S. E., J. E. Blum, and J. R. Pawlik. Redwood of the reef: growth and age of the giant barrel
130 sponge *Xestospongia muta* in the Florida Keys. *Mar. Biol.* 155: 159-171 (2008).

131 S7. Laguionie-Marchais C., Kuhn L. A., Huffard C. L., Ruhl H. A., Smith K. L., Jr., Spatial and temporal
132 variation in sponge spicule patches at Station M, northeast Pacific. *Mar. Biol.* 162, 617-624 (2015).

133

134 **Section 4 -The biological fluxes**

135 **4.1 Biogenic silica production as measured by isotopic techniques**

136 **Annex 1** shows all bSi production data with corresponding references.

137 **4.2 bSi pelagic production uncertainties**

138 The bSi pelagic production estimates from satellite NPP products, ocean biogeochemical
139 models, and empirical studies each have their own uncertainty. When NPP is extrapolated to
140 silica production, values are multiplied by estimates of the fraction of primary productivity done
141 by diatoms and then a Si:C ratio. Of these, the choice of a Si:C ratio for the HNLC regions is
142 the most uncertain. Our chosen value of 0.52 is 4 fold higher than for nutrient-replete temperate
143 diatoms (Brzezinski, 1985, see text reference list), but field observations suggest anywhere
144 from no effect to an 8-fold increase (S8: Marchetti & Harrison, 2007, S9 : Timmermans et al.,
145 2004). Biases in satellite NPP models also contribute to uncertainties in estimates of Si
146 production. Particularly relevant are potential biases in Southern Ocean chlorophyll
147 concentrations (and consequently, NPP), which may be underestimated in the Southern Ocean
148 by as much as a factor of 3-4 (S10: Johnson et al., 2013).

149 For the biogeochemical models the two main sources of uncertainty are the extrapolation from
150 silica export to gross silica production using D:P ratios (Table 1 of Tréguer & De La Rocha,
151 2013, main text reference list), as well as uncertainties in the parametrization of Southern Ocean
152 physics and biology. For the seven GOBMs that report separate estimates of net silica
153 production for the Southern Ocean, when Southern Ocean silica production is regressed against
154 total global silica production the fitted line has an R^2 value of 0.96 with a slope equal to 1.05
155 (± 0.11). This points to model parameterizations of Southern Ocean physics and/or biology as
156 the major determinant of differences in global silica production estimates among GOBMs.

157 For the global bSi pelagic production estimates derived from field data, extrapolation over both
158 time and space is required. Few empirical studies of bSi production over an entire year are
159 available. The vast majority of annual estimates determined for Longhurst provinces are
160 extrapolated from field programs lasting a few weeks or less. Data sparsity remains a problem,
161 and in the analysis presented here some Longhurst zones contain only a single measurement
162 and nearly half of zones have no data at all (main text Fig. 3).

163 REFERENCES

- 164 S8. Marchetti, A., Harrison, P.J. Coupled changes in the cell morphology and elemental (C, N, and Si)
165 composition of the pennate diatom *Pseudo-nitzschia* due to iron deficiency. *Limnol. Oceanogr.* **52**, 2270–
166 2284 (2007).
- 167 S9. Timmermans, K.R., van der Wagt, B., de Baar, H.J.W. Growth rates, half-saturation constants, and
168 silicate, nitrate, and phosphate depletion in relation to iron availability of four large, open-ocean diatoms
169 from the Southern Ocean. *Limnol. Oceanogr.* **49**, 2141–2151 (2004).
- 170 S10. Johnson, R., Strutton, P.G., Wright, S.W., McMinn, A., Meiners, K.M. Three improved satellite
171 chlorophyll algorithms for the Southern Ocean. *J. Geophys. Res. Ocean.* **118**, 3694–3703 (2013).

172 4.3 bSi production: contribution of benthic diatoms

173 Determination of bSi production by microphytobenthos is still in its infancy. However, we can
174 already foresee its potential impact in two ways. Through intensive studies in a temperate
175 subtidal ecosystem (S11: Ni Longphuir et al. 2009, S12 : Leynaert et al. 2009, and S13 :
176 Chatterjee et al., 2013) we conservatively estimate that benthic diatoms can produce 1 mol-Si
177 m⁻² yr⁻¹. Extrapolation to the photic area of the world coastal ocean gives 6.8 Tmol-Si yr⁻¹.
178 Cahoon (1999) (S14) proposed a global estimate of annual benthic microalgal primary
179 production (based on the analysis of 85 worldwide studies) of 514 × 10¹² gC yr⁻¹. Applying the
180 Brzezinski (1985) (main text reference list) mean Si/C molar ratio for marine diatom of 0.13 to
181 convert this carbon production in silica production, gives 5.4 Tmol-Si yr⁻¹. Both estimates are
182 in good agreement. Therefore, the biogenic silica production by the microphytobenthos might
183 represent about 2.5% of the global bSi production. It remains to be determined what proportion
184 of this flux will finally contributes to the net sink of bSi. Preliminary studies have shown that
185 bSi dissolution rates of benthic diatoms are 10 times slower than pelagic diatoms measured in
186 the same conditions.

187 REFERENCES

- 188 S11. Ni Longphuir, S. *et al.* Diurnal heterogeneity in silicic acid fluxes in shallow coastal sites: causes and
189 implications. *Estuar. Coast. Shelf Sci.* **82**, 495-502 (2009).
- 190 S12. Leynaert, A., Longphuir, S. N., Claquin, P., Chauvaud, L. & Ragueneau, O. No limit? The multiphasic
191 uptake of silicic acid by benthic diatoms. *Limnol. Oceanogr.* **54**, 571–576 (2009).
- 192 S13. Chatterjee, A. *et al.* Comparative dynamics of pelagic and benthic micro-algae in a coastal ecosystem.
193 *Estuarine, Coastal and Shelf Science* **133**, 67–77 (2013).
- 194 S14. Cahoon, L. B. The role of benthic microalgae in neritic ecosystems. in *Oceanography and Marine*
195 *Biology. An Annual Review.* vol. 37, CRC Press (1999).

196

197 Section 5 – Discussion

198 **5.1 Preliminary note to Fig. 4:** Depiction of a schematic Si cycle in the coastal and continental
199 margin zone (CCMZ), excluding coastal upwelling, linked to the rest of the world ocean (« open
200 ocean » zone including upwelling and polar zones). In principle, the CCMZ comprises proximal
201 and distal coastal zones, as defined by Laruelle et al. (2009) (main text reference list), which
202 includes coastal upwelling in the distal coastal zone. However, in coastal upwelling zones both
203 physical and biogeochemical dynamics are markedly different from those in the CCMZ *as*
204 represented by Jeandel et al. 2016 (see main text reference list). Indeed, if in the CCMZ the
205 transfer of material from land to ocean plays a major role in physical and biogeochemical
206 processes, in coastal upwelling zones the bSi production is mainly fueled by dSi flux from
207 below (i.e. from deeper in the water column), as it does for the global ocean *sensu lato* (Fig. 1).

208 Therefore, in our synthesis quantifying the Si cycle in the « boundary exchange » zone, coastal
209 upwelling is discarded from the CCMZ and conceptually incorporated to the « open ocean »
210 zone, as shown in Fig. 4. This figure represents a possible Si cycle *assuming steady state* in the
211 CCMZ and in the rest of the world ocean (so called « open ocean »), that is with total inputs =
212 total outputs = 14.8 Tmol-Si yr⁻¹.

213 **5.2 Detailed legend of Fig. 4**

214 Input and dSi fluxes (grey arrows), outputs and biological fluxes (black arrows). In this steady-
215 state scenario total inputs = total outputs = 14.8 Tmol-Si yr⁻¹ (consistent with main text Figure
216 1). Note that (1) in the CCMZ the burial flux of bSi (3.7 Tmol-Si yr⁻¹) and the reverse
217 weathering flux (5.7 Tmol-Si yr⁻¹) (authigenic siliceous material) are fed by the export flux of
218 biogenic silica, and (2) the « open ocean » deficiency in dSi (2.9 Tmol-Si yr⁻¹) is made up by a
219 transfer from the CCMZ. For details about those inputs and outputs refer to the « inputs » and
220 « outputs » sections of the main text. For abbreviations, see detailed legend of Fig. 1 in
221 Supplement (section 1).

222 *Additional comments of Figure 4:*

223 -Estimates of burial rate (planktonic bSi): in the CCMZ (excluding coastal upwelling),
224 according to Rahman et al. 2017 (main text reference), $F_{BCCMZ} = 3.7$ Tmol-Si yr⁻¹; in the « open
225 ocean » zone, consistent with Fig. 1, $F_{Bopenocean} = 7.4 - 3.7 = 3.7$ Tmol-Si yr⁻¹.

226 -Estimates of reverse weathering flux: in Fig. 4 CCMZ F_{RWCCMZ} equals 5.7 Tmol-Si yr⁻¹,
227 compatible with $F_R = 4.7 (\pm 2.1)$ Tmol-Si yr⁻¹ (Rahman et al. 2017, see main text reference list).
228 Reverse weathering in the « open ocean » remains unquantified.

229 -Following Annex 1, bSi production in the CCMZ* is 13% of the total production (255 Tmol-
230 Si yr⁻¹). Consequently, F_{Pgross} is 33 and 222 Tmol-Si yr⁻¹, for CCMZ and “open ocean”,
231 respectively.

232 -According to Tréguer and De La Rocha (2013) (main text reference list), the pelagic
233 production to dissolution ratio (D:P) being 0.51 and 0.57, for CCMZ and “open ocean”,
234 respectively. $F_{D(surface)}$ for those compartments is 17 and 124 Tmol-Si yr⁻¹, and F_E is 16 and 98
235 Tmol-Si yr⁻¹, for CCMZ and “open ocean”, respectively.

236 Note than every component of the inputs, outputs, and biological Si fluxes, although interacting
237 between each other, is determined by independent methods, that is there is no overlap in the
238 counting of these fluxes. Therefore, the export production (16.0 Tmol-Si yr⁻¹) in particular feeds
239 both the Si burial rate (3.7 Tmol-Si yr⁻¹) and the reverse weathering flux (5.7 Tmol-Si yr⁻¹).

240 Figure 4 also shows that the “open ocean” bSi production is mostly fueled by dSi inputs from
241 below (91.7 Tmol-Si yr⁻¹), the CCMZ only providing 2.9 Tmol-Si yr⁻¹ to the “open ocean”.

242 *Although specific estuarine sites have been studied (e.g. S15 DeMaster 1983, S16, S17
243 Raimonet al. 2013a & b), no global estimate is presently available for the estuarine bSi
244 production. Given that most of estuarine waters are turbid we anticipate that the contribution of
245 estuaries to the total bSi of the coastal zone should be small.

246 *REFERENCES*

- 247 S15. DeMaster D.J., Knapp, G.B., Nittrouer, C.A. Biological uptake and accumulation of silica on the Amazon
248 continental shelf. *Geochim. Cosmochim. Acta.* **47**, 1713-23 (1983).
- 249 S16. Raimonet M. et al. Spatio-temporal variability in benthic silica cycling in two macrotidalestuaries: Causes
250 and consequences for local to global studies. *Est. Coast. Shelf Sci.*, **119**, 31-43 (2013a).
- 251 S17. Raimonet, M. et al. Strong gradient of benthic biogeochemical processes along a macrotidal temperate
252 estuary: focus on P and Si cycles. *Biogeochemistry* **117**, 399-417 (2013b).

253

254

255

256

Annex 1. Data of biogenic silica production as measured from isotopic techniques

All data													
Domain	Ocean	Province	System	sampling period	range pBSi	moyenne	longitude	latitude	prod. duration	annual PBSi	annual PBSi	References	References
					mmol-Si m ⁻² day ⁻¹	mmol-Si m ⁻² day ⁻¹			d	mmol m ⁻² an ⁻¹	10 ⁹ mol an ⁻¹		
Coastal	Pacific	ALSK	Bering Sea coastal	May/July 1979 - 1980	7 - 51	20,9	-163	57	159	3317	1957	Banahan and Goering, 1986	18
Polar	Southern Ocean	CCSZ	Ross Sea	jan/feb 1990		34,0	-175	-76,5	79	2698	5396	Nelson et al., 1991	19
Polar	Southern Ocean	CCSZ	Pacific, Ross Sea		7,1 - 93	38,0	166	-75	79	3015	6030	Nelson and Smith, 1986	20
Polar	Southern Ocean	CCSZ	Indian sector	spring 95/summer 95		5,3	62	-66	79	421	841	Caubert 1998	21
Polar	Southern Ocean	CCSZ	Indian sector			57,6	140,42	-66,42	79	4570	9141	Beucher et al., 2004	22
Polar	Southern Ocean	PFZ	Weddell Scotia	1992	19,6-60,7	27,6	-6	-49,5	130	3582	32237	Quéguiner et al., 2002	23
Polar	Southern Ocean	PFZ	Weddell sea (EPOS 2)	1988/1989	11,2-20,6	15,9	-49	-57	130	2063	18571	Quéguiner et al., 1991	24
Polar	Southern Ocean	PFZ	Indian sector	spring 95/summer 94	1,7-2,2	2,0	62	-50	130	260	2336	Caubert 1998	21
Polar	Southern Ocean	POOZ	Pacific	oct/nov 97		12,2	-170	-58	114	1385	16625	Brzezinski et al., 2001	25
Polar	Southern Ocean	POOZ	Pacific	dec 97		27,1	-170	-60	114	3077	36928	Brzezinski et al., 2001	25
Polar	Southern Ocean	POOZ	East of New Zealand			3,6	-175	-61	114	404	4851	Nelson and Gordon, 1982	26
Polar	Southern Ocean	POOZ	Weddell Scotia	1992		3,5	-6	-52	114	397	4769	Quéguiner et al., 2002	23
Polar	Southern Ocean	POOZ	Pacific	summer 1998	0,6 - 2,58	2,6	142	-52	114	293	3516	Quéguiner et al., 2001	27
Polar	Southern Ocean	POOZ	Kerguelen KEOPS 2	2011	3,09 - 47,9	23,2	72	-48,5	114	2634	31614	Closset et al 2014	28
Polar	Southern Ocean	POOZ	Indian sector	spring 95/summer 94	2,3-3,9	3,1	62	-54	114	352	4224	Caubert 1998	21
Polar	Southern Ocean	POOZ	Indian sector			10,2	142	-55	114	1158	13899	Beucher et al., 2004	22
Polar	Southern Ocean	SIZ	Weddell Sea	1990	1,97 - 3,18	2,6	-40	-70	114	294	6176	Leynaert et al., 1993	29
Polar	Southern Ocean	SIZ	Pacific	janv-98		22,6	-170	-61	114	2566	53893	Brzezinski et al., 2001	25
Polar	Southern Ocean	SIZ	Pacific	feb march 98		5,5	-170	-62	114	625	13116	Brzezinski et al., 2001	25
Polar	Southern Ocean	SIZ	East of New Zealand			6,8	-175	-64	114	768	16120	Nelson and Gordon, 1982	26
Polar	Southern Ocean	SIZ	Weddell Sea (EPOS 2)	1988/1989	2,3 - 22,9	10,1	-49	-60	114	1147	24085	Tréguer et al., 1991	30
Polar	Southern Ocean	SIZ	Weddell Scotia	1992		4,4	-6	-57	114	500	10493	Quéguiner et al., 2002	23
Polar	Southern Ocean	SIZ	Weddell sea (EPOS 2)	1988/1989	6,0-20,0	10,8	-49	-60,5	114	1226	25754	Quéguiner et al., 1991	24
Polar	Southern Ocean	SIZ	Indian sector	spring 95/sum. 94	5,7-8,9	7,0	62	-58,5	114	795	16693	Caubert 1998	21
Polar	Southern Ocean	SIZ	Indian sector			7,9	138,68	-65	114	897	18839	Beucher et al., 2004	22
Polar	Southern Ocean	SUBANT	Indian sector	Jan/feb 1999	0,25 - 0,93	0,5	63	-45	95	47	1353	Leblanc et al., 2002	31
Polar	Southern Ocean	SUBANT	Pacific	Summer 1998	0,6 - 2,58	1,1	142	-47	95	102	2955	Quéguiner et al., 2001	27
Polar	Southern Ocean	SUBANT	East of New Zealand			2,0	-175	-54	95	190	5523	Nelson and Gordon, 1982	26
Polar	Southern Ocean	SUBANT	Indian sector			1,7	146	-48	95	162	4694	Beucher et al., 2004	22
Polar	Pacific	BERS	Bering Sea outer	May/July 1979 - 1981	1,8 - 25	10,1	-167	55,5	95	962	3741	Banahan and Goering, 1988	18
Coastal	Pacific	CAMR	Baja California		2,3 - 1140	89,0	-114	27	365	32485	40931	Nelson and Goering, 1978	32
Coastal	Pacific	CCAL	Coast. upw Monterey	April 1995	13 - 1140	202,0	-122,2	36,8	175	35262	33852	Brzezinski et al., 1997	33
Coastal	Pacific	CCAL	Coast. upw Monterey	April 1992	22 - 40	32,0	-122,2	36,2	175	5586	5363	Brzezinski et al., 1997	33
Coastal	Pacific	CCAL	Costa Rica Dome	July 2011	0,11 - 5,57	1,4	-122	33	175	251	241	Krause et al., 2015	34
Coastal	Pacific	CCAL	Santa Barbara	1996-1997	2,4-57,3	34,2	-120	34	175	5970	5731	Shipe and Brzezinski, 2001	35
Coastal	Pacific	CHIL	Peru Chile upw - BIOSOPE	Oct/Nov 2004	42 - 52	47,0	-72	-35	190	8950	23361	Leblanc et al 2018	36
Coastal	Pacific	CHIL		March/May 1977	0,1 - 4,04	1,2	-78	-13	190	226	590	Nelson et al., 1981	37
Coastal	Atlantic	CNRY	NW Africa upw	May 1974	14 - 40	23,0	-17,5	21,5	222	5106	4085	Nelson and Goering, 1977	38
Westerlies	Mediterranean	MEDI	Mediterranean (SOFI)	Dec -Jan 1997/98	0,80	0,8	-1	36	127	105	325	Leblanc et al., 2004	39
Westerlies	Mediterranean	MEDI	Mediterranean (SOFI)	1999/2000	0,14-1,4	0,4	5	43	127	48	149	Leblanc et al 2003	40
Westerlies	Atlantic	NADR	N Atlantic (PAP)		0,5-1,3	0,9	-16	48,8	127	114	400	Ragueneau et al., 1997	41
Westerlies	Atlantic	NADR	NE Atlantic	May 2001	6 - 166	35,0	-15	60	127	4445	15558	Brown et al., 2003	42
Westerlies	Atlantic	NAST (E)	N Atlantic -POMME	March/April/ sept 2001	0,04-11,2	0,8	-18	42	127	96	36	Leblanc et al 2005	43
Westerlies	Atlantic	NAST (W)	Warm-Core Ring	April/June 1982	4,4 - 11,7	7,1	-63	37	127	901	5228	Brzezinski and Nelson, 1989	44
Westerlies	Atlantic	NAST (W)	Sargasso Sea (BATS)	1991-1994	0,2 - 1,6	0,7	-64	31,1	365	240	1392	Nelson and Brzezinski, 1997	45
Westerlies	Atlantic	NAST (W)	Mode-water eddy	April 2007	0,13 - 1,12	0,6	-64	32,5	127	81	471	Krause et al., 2010	46
Westerlies	Atlantic	NAST (W)	Sargasso Sea	May/March 1989	0,2 - 1,48	0,6	-65	32	127	81	471	Brzezinski and Kosman, 1996	47
Westerlies	Atlantic	NADR	N. Atlantic (BENGAL)			0,9	30	46	127	114	400	Ragueneau et al., 2000	48
Trades	Pacific	NPTG	Central North Pacific	August 1995	0,5 - 2,9	1,2	-155	28	365	453	9545	Brzezinski et al 1998	49
Trades	Pacific	NPTG	HOT-ALOHA		0,09 - 0,49	0,2	-158	22,5	365	63	1329	Brzezinski et al., 2011	50
Trades	Pacific	NPTG	Subtropical front	summer bloom	0,10 - 1,74	0,6	-150	27	365	219	4619	Krause et al., 2013	51
Trades	Pacific	PEQD	Eq. upwelling	nov-96	2,50	2,5	-180	0	365	913	9435	Leynaert et al., 2001	52
Trades	Pacific	PEQD	Eq. upwelling	nov-96	0,1 - 0,65	0,4	-180	5	365	146	1510	Leynaert et al., 2001	52
Trades	Pacific	PEQD	Eq. upwelling	nov-96	0,1 - 0,65	0,4	-180	-5	365	146	1510	Leynaert et al., 2001	52
Trades	Pacific	PEQD	Eastern equa. Upw.	2004	0,6 - 2,59	1,6	-110	-1	365	584	6039	Krause et al., 2011	53
Trades	Pacific	PEQD	Eastern equa. Upw.	2005	0,3 - 2,49	1,3	-125	0	365	475	4906	Krause et al., 2011	53
Westerlies	Pacific	PSAG	N Pacific (OSP)			5,1	-145	50	111	566	1812	Wong and Matar, 1999	54
Polar	Atlantic	SARC	Svalbard	May 2016	0,27 - 1,46	0,8	25	77	127	102	237	Krause et al 2018	55
Westerlies	Pacific	SPSG	SE Gyre BIOSOPE	Oct/Nov 2004	0,04 - 0,2	0,1	-90	-31	127	15	568	Leblanc et al 2018	36
Westerlies	Pacific	SPSG	HNLC - BIOSOPE	Oct/Nov 2004	0,8 - 5,6	3,6	-138	-8	127	457	17043	Leblanc et al 2018	36
Trades	Pacific	WARM	W equat. oligotrophic	oct-94	0,8 - 2,1	1,4	-168	0	365	511	8575	Blain et al., 1997	56
Trades	Pacific	WARM	W equat. oligotrophic			3,9	-160	0	365	1424	23886	Blain et al., 1997	56
Coastal	Atlantic	NECS	Bay of Brest	2001		2,5	-4,6	48,3	365	900	1224	Del Amo et al	57
Westerlies	Atlantic	NADR	AMT	April- May 2004		0,26	-16,78	47,45	127	32	113	Poulton et al, 2006	58
Westerlies	Atlantic	NASE	AMT	April- May 2004		0,43	-20,16	38,5	127	54	241	Poulton et al, 2006	58
Westerlies	Atlantic	NASW	AMT	April- May 2004		0,15	-36,4	29,2	127	19	110	Poulton et al, 2006	58
Trades	Atlantic	NATR	AMT	April- May 2004		0,14	-33,4	22,2	365	51	423	Poulton et al, 2006	58
Trades	Atlantic	SATL	AMT	April- May 2004		0,31	-30,37	-27,48	190	60	1058	Poulton et al, 2006	58
Trades	Atlantic	WTRA	AMT	April- May 2004		0,58	-27,06	6,1	365	212	1135	Poulton et al, 2006	58
Westerlies	Atlantic	NADR	Celtic Sea	nov-14		1,00	-8,83	48,75	127	127	445	Poulton et al, 2019	59

258

259

260

261

262

Note that « d », duration of the productive period in days, was extrapolated from the bloom phenology for each province (calculated as the number of day where the chloro concentration is greater than the average concentration between the maximum and the minimum values).

Average per province

Domain	Province	Surface 10 ⁶ km ²	Average mmol-Si m ⁻² day ⁻¹	Prod. duration day	Annual PBSi mmol-Si m ⁻² an ⁻¹	Annual PBSi 10 ⁹ mol-Si an ⁻¹
Coastal	CNRY	0,8	23,0	190	4 380	3 504
Coastal	NECS	1,4	2,5	159	391	532
Polar	SARC	2,3	0,8	127	102	237
Westerlies	NADR	3,5	9,3	127	1 176	4 116
Westerlies	NAST (E)	0,4	0,6	127	75	28
Westerlies	NAST (W)	5,8	1,8	127	233	1 353
Westerlies	MEDI	3,1	0,6	127	77	237
Coastal	ALSK	0,6	20,9	159	3 317	1 957
Coastal	CAMR	1,3	89,0	365	32 485	40 931
Coastal	CCAL	1,0	67,4	175	11 767	11 297
Coastal	CHIL	2,6	24,1	190	4 588	11 975
Polar	BERS	3,9	10,1	95	962	3 741
Trades	NPTG	21,1	0,7	365	245	5 164
Trades	PEQD	10,3	1,2	365	453	4 680
Trades	WARM	16,8	2,7	365	967	16 230
Westerlies	PSAE	3,2	5,1	111	567	1 813
Westerlies	SPSG	37,3	1,9	127	236	8 806
Westerlies	SUBANT	29,0	1,3	95	125	3 631
Polar	PFZ	9,0	15,2	130	1 968	17 715
Polar	POOZ	12,0	10,7	114	1 213	14 553
Polar	SIZ	21,0	8,6	114	980	20 574
Westerlies	NASW	5,8	0,2	127	19	110
Trades	NATR	8,3	0,1	365	51	423
Trades	SATL	17,8	0,3	190	60	1 058
Trades	WTRA	5,4	0,6	365	212	1 135
Polar	CCSZ	2,0	33,7	79	2676,0	5352,0

Average per Ocean Basin

Ocean	Surface 10 ⁶ km ²	portion in AO	Surface considered*	Annual PBSi mmol m ⁻² an ⁻¹	Annual PBSi 10 ⁹ mol an ⁻¹	% PBSi Total
Atlantic	106	20	86	250	21 501	8,6
Pacific	166	32	134	1 088	145 736	58,6
Southern Ocean*			73	586	42 760	17,2
Arctic	14		14	102	1 422	0,6
Indien	75	21			37 188	15,0
Sub Total	361		307		211 419	
				Annual World PBSi	248 607	

Average per domain (Longhurst & Tréguer and Jacques for the OA)

Domain	Surface 10 ⁶ km ²			Annual PBSi mmol m ⁻² an ⁻¹	Annual PBSi 10 ⁹ mol an ⁻¹	% PBSi Total
Coastal	29			1 276	37 017	13,0
upw	8			12 026	101 019	35,5
Polar*	54			1 238	66 827	23,5
Westerlies*	130			228	29 647	10,4
Trades	140			360	50 416	17,7
				Annual World PBSi	284 926	

*Surface considered: this is the ocean surface minus the Antarctic ocean

*Polar: surface includes the Antarctic Ocean (from the polar front poleward, i.e. 44 10⁶ km²) + arctic zones as defined by Longhurst

*Westerlies: includes the subantarctic zone

*Southern Ocean: includes the Antarctic Ocean and the subantarctic zone (i.e. respectively 44 10⁶ km² + 29 10⁶ km²)

263
264

265 REFERENCES

266 S18. Banahan, S. & Goering, J. J. The production of biogenic silica and its accumulation on the southeastern
267 Bering Sea shelf. *Cont. Shelf Res.* **5**, 199–213 (1986).

-
- 268 S19. Nelson, D. M., Ahern, J. A. & Herlihy, L. J. Cycling of biogenic silica within the upper water column of
269 the Ross Sea. *Mar. Chem.* **35**, 461–476 (1991).
- 270 S20. Nelson, D. M. & Smith Jr, W. O. Phytoplankton bloom dynamics of the western Ross Sea ice edge. II.
271 Mesoscale cycling of nitrogen and silicon. *Deep Sea Res.* **33**, 1389–1412 (1986).
- 272 S21. Caubert, T. Production de silice biogène dans l’océan Austral: détermination par spectrométrie de masse.
273 PhD, Université de Bretagne Occidentale, Brest, 190pp. (1998).
- 274 S22. Beucher, C., Tréguer, P., Corvaisier, R., Hapette, A. & Elskens, M. Production and dissolution of
275 biosilica, and changing microphytoplankton dominance in the Bay of Brest (France). *Mar. Ecol. Prog.
276 Ser.* **267**, 57–69 (2004).
- 277 S23. Quéguiner, B. & Brzezinski, M. A. Biogenic silica production rates and particulate organic matter
278 distribution in the Atlantic sector of the Southern Ocean during austral spring 1992. *Deep Sea Res. II:
279 Topical Studies in Oceanogr.* **49**, 1765–1786 (2002).
- 280 S24. Quéguiner, B., Tréguer, P. & Nelson, D. M. The production of biogenic silica in the Weddell and Scotia
281 Seas. *Mar. Chem.* **35**, 449–459 (1991).
- 282 S25. Brzezinski, M. A., Nelson, D. M., Franck, V. M. & Sigmon, D. E. Silicon dynamics within an intense
283 open-ocean diatom bloom in the Pacific sector of the Southern Ocean. *Deep Sea Res. Part II: Topical
284 Studies in Oceanography* **48**, 3997–4018 (2001).
- 285 S26. Nelson, D. M. & Gordon, L. I. Production and pelagic dissolution of biogenic silica in the Southern
286 Ocean. *Geochim. Cosmochim. Acta* **46**, 491–501 (1982).
- 287 S27. Quéguiner, B. Biogenic silica production in the Australian sector of the Subantarctic Zone of the Southern
288 Ocean in late summer 1998. *J. Geophys. Res.* **106**, 31627–31636 (2001).
- 289 S28. Closset, I. *et al.* Seasonal evolution of net and regenerated silica production around a natural Fe-fertilized
290 area in the Southern Ocean estimated with Si isotopic approaches. *Biogeosciences* **11**, 5827–5846 (2014).
- 291 S29. Leynaert, A., Nelson, D., Quéguiner, B. & Tréguer, P. The silica cycle in the Antarctic Ocean: is the
292 Weddell Sea atypical? *Mar. Ecol. Prog. Ser.* **96**, 1–15 (1993).
- 293 S30. Tréguer, P. *et al.* Production of biogenic silica in the Weddell-Scotia Seas measured with ³²Si. *Limnol.
294 Oceanogr.* **36**, 1217–1227 (1991).
- 295 S31. Leblanc, K. *et al.* Particulate biogenic silica and carbon production rates and particulate matter
296 distribution in the Indian sector of the Subantarctic Ocean. *Deep Sea Res. Part II: Topical Studies in
297 Oceanogr.* **49**, 3189–3206 (2002).
- 298 S32. Nelson, D. M. & Goering, J. J. Assimilation of silicic acid by phytoplankton in the Baja California and
299 northwest Africa upwelling systems 1: Silicic acid uptake. *Limnol. Oceanogr.* **23**, 508–517 (1978).
- 300 S33. Brzezinski, M. A. & Phillips, D. R. Evaluation of ³²Si as a tracer for measuring silica production rates in
301 marine waters. *Limnol. Oceanogr.* **42**, 856–865 (1997).
- 302 S34. Krause, J. W. *et al.* Variability in diatom contributions to biomass, organic matter production and export
303 across a frontal gradient in the California Current Ecosystem. *J. Geophys. Res. Oceans* **120**, 1032–1047
304 (2015).
- 305 S35. Shipe, R. F. & Brzezinski, M. A. A time series study of silica production and flux in an eastern boundary
306 region: Santa Barbara Basin, California. *Glob. Biogeochem. Cycles* **15**, 517–531 (2001).
- 307 S36. Leblanc, K. *et al.* Silicon cycle in the Tropical South Pacific: evidence for an active pico-sized siliceous

-
- 308 plankton. *Biogeosciences* (2018).doi:10.5194/bg-2018-149.
- 309 S37. Nelson, D. M., Goering, J. J. & Boisseau, D. W. Consumption and regeneration of silicic acid in three
310 coastal upwelling systems. *Coast. Est. Sci.* **1**, 242–256 (1981).
- 311 S38. Nelson, D. M. & Goering, J. J. Near-surface silica dissolution in the upwelling region off northwest
312 Africa. *Deep Sea Research* **24**, 65–73 (1977).
- 313 S39. Leblanc, K. Siliceous phytoplankton production and export related to trans-frontal dynamics of the
314 Almeria-Oran frontal system (western Mediterranean Sea) during winter. *J. Geophys. Res.* **109**, C07010
315 (2004).
- 316 S40. Leblanc, K., Quéguiner, B., Garcia, N., Rimmelin, P. & Raimbault, P. Silicon cycle in the NW
317 Mediterranean Sea: seasonal study of a coastal oligotrophic site. *Oceanol. Acta* **26**, 339–355 (2003).
- 318 S41. Ragueneau, O. *et al.* OPALEO (OPAL as PALEO-productivity proxy), Synthesis and perspectives, in
319 Final Report From the First 2 International Workshop (Brest 96, Corvallis 97), edited by O. Ragueneau,
320 A. Leynaert, and P. Tréguer, Brest, France (1997).
- 321 S42. Brown, L., Sanders, R., Savidge, G. & Lucas, C. H. The uptake of silica during the spring bloom in the
322 Northeast Atlantic Ocean. *Limnol. Oceanogr.* **48**, 1831–1845 (2003).
- 323 S43. Leblanc, K. *et al.* A seasonal study of diatom dynamics in the North Atlantic during the POMME
324 experiment (2001): Evidence for Si limitation of the spring bloom. *J. Geophys. Res.* **110**, C07S14 (2005).
- 325 S44. Brzezinski, M. A. & Nelson, D. M. Seasonal changes in the silicon cycle within a Gulf Stream warm-
326 core ring. *Deep Sea Res. Part A. Oceanogr. Res. Papers* **36**, 1009–1030 (1989).
- 327 S45. Nelson, D. M. & Brzezinski, M. A. Diatom growth and productivity in an oligo-trophic midocean gyre:
328 A 3-yr record from the Sargasso Sea near Bermuda. *Limnol. Oceanogr.* **42**, 473–486 (1997).
- 329 S46. Krause, J. W., Nelson, D. M. & Lomas, M. W. Production, dissolution, accumulation, and potential export
330 of biogenic silica in a Sargasso Sea mode-water eddy. *Limnol. Oceanogr.* **55**, 11 (2010).
- 331 S47. Brzezinski, M. & Kosman, C. Silica production in the Sargasso Sea during spring 1989. *Mar. Ecol. Prog.*
332 *Ser.* **142**, 39–45 (1996).
- 333 S48. Ragueneau, O. *et al.* A review of the Si cycle in the modern ocean: recent progress and missing gaps in
334 the application of biogenic opal as a paleoproductivity proxy. *Glob. Planet. Change* **26**, 317–365 (2000).
- 335 S49. Brzezinski, M., Villareal, T. & Lipschultz, F. Silica production and the contribution of diatoms to new
336 and primary production in the central North Pacific. *Mar. Ecol. Prog. Ser.* **167**, 89–104 (1998).
- 337 S50. Brzezinski, M. A. *et al.* The annual silica cycle of the North Pacific subtropical gyre. *Deep Sea Res. Part*
338 *I: Oceanogr. Res. Papers* **58**, 988–1001 (2011).
- 339 S51. Krause, J. W., Brzezinski, M. A., Villareal, T. A. & Wilson, C. Biogenic silica cycling during summer
340 phytoplankton blooms in the North Pacific subtropical gyre. *Deep Sea Res. Part I: Oceanogr. Research*
341 *Papers* **71**, 49–60 (2013).
- 342 S52. Leynaert, A., Tréguer, P., Lancelot, C. & Rodier, M. Silicon limitation of biogenic silica production in
343 the Equatorial Pacific. *Deep Sea Res. Part I* **48**, 639–660 (2001).
- 344 S53. Krause, J. W., Nelson, D. M. & Brzezinski, M. A. Biogenic silica production and the diatom contribution
345 to primary production and nitrate uptake in the eastern equatorial Pacific Ocean. *Deep Sea Res. Part II:*
346 *Topical Studies in Oceanogr.* **58**, 434–448 (2011).
- 347 S54. Wong, C. S. & Matear, R. J. Sporadic silicate limitation of phytoplankton productivity in the subarctic

348 NE Pacific. *Deep Sea Res. Part II: Topical Studies in Oceanogr.* **46**, 2539–2555 (1999).

349 S55. Krause, J.W. *et al.* Biogenic silica production and diatom dynamics in the Svalbard region during spring.

350 *Biogeosciences* **15**, 6503–6517 (2018).

351 S56. Blain, S. *et al.* Biomass, growth rates and limitation of Equatorial Pacific diatoms. *Deep Sea Res. Part I:*

352 *Oceanogr. Res. Papers* **44**, 1255–1275 (1997).

353 S57. Del Amo, Y., Queguiner, B., Treguer, P., Breton, H. & Lampert, L. Impacts of high-nitrate freshwater

354 inputs on macrotidal ecosystems. 11. Specific role of the silicic acid pump in the year-round dominance

355 of diatoms in the Bay of Brest (France). *Mar. Ecol. Progr. Ser.* **161**, 225–237 (1997).

356 S58. Poulton A.J. *et al.* (2006) Phytoplankton mineralization in the tropical and subtropical Atlantic Ocean.

357 *Glob. Biogeochem. Cycles* **20**, GB4002 (2006).

358 S59. Poulton A.J. *et al.* (2019) Dissolution dominates silica cycling in a shelf sea autumn bloom. *Geophys.*

359 *Res. Lett.* **46**, 6765 – 6774 (2019).

360

361