

Comparing modified substrate induced respiration with selective inhibition (SIRIN) and N₂O isotope approaches to estimate fungal contribution to denitrification in three arable soils under anoxic conditions

5 Lena Rohe^{1, 2, 3}, Traute-Heidi Anderson², Heinz Flessa², Anette ~~Giesemann~~²Goeske², Dominika Lewicka-Szczebak^{2, 4}, Nicole Wrage-Mönnig⁵, Reinhard Well²

¹Helmholtz Centre for Environmental Research – UFZ, Department Soil System Sciences, Theodor-Lieser Str. 4, Halle, Germany

²Thünen Institute of Climate Smart Agriculture, Bundesallee 65, Braunschweig, Germany

10 ³University of Göttingen, Department of Crop Sciences, Institute of Grassland Science, von-Siebold-Str. 8, 37075 Göttingen, Germany

⁴~~Institute of Geological Sciences, University of Wrocław, pl. M. Borna 9, 50-204 Wrocław, Poland University of Göttingen, Centre for Stable Isotope Research and Analysis, Büsgenweg 2, 37077 Göttingen, Germany~~

15 ⁵University of Rostock, Agricultural and Environmental Faculty, Grassland and Fodder Sciences, Justus-Liebig-Weg 6, Rostock, Germany

Correspondence to: Lena Rohe (lena.rohe@ufz.de)

Keywords: selective growth inhibition, ¹⁵N site preference, fungal denitrification, C₂H₂, isotope endmember mixing approach, *SP_{N2O}* mixing balance, *SP/δ¹⁸O* mapping approach

20 Abstract

~~The co-existence of many N₂O production pathways in soil hampers differentiation of microbial pathways. It is still not proven. The question whether fungi have a large contribution are major significant contributors to the soil emissions of the greenhouse gas nitrous oxide (N₂O) from denitrification has not yet been resolved. Here, as far as we know~~
25 ~~The co-existing of many N₂O production pathways in soil hampers differentiation of microbial pathways. To our knowledge, for the first time~~ three approaches to independently investigate the fungal fraction contributing to N₂O from denitrification were used simultaneously ~~for the first time~~ (modified substrate induced respiration with selective inhibition (SIRIN) approach, and two isotopic approaches, i.e. endmember mixing approach (IEM) using the ¹⁵N site preference of N₂O produced (*SP_{N2O}*), and the *SP/δ¹⁸O* mapping approach (*SP/δ¹⁸O* Map)). This enabled a comparison of methods and a quantification of the importance of fungal
30 ~~denitrification in soil. Pure culture studies provide evidence of the ability of soil fungi to produce nitrous oxide (N₂O) during denitrification. Soil studies with selective inhibition indicated a possible dominance of fungal compared to bacterial N₂O production in soil, which drew more attention to fungal denitrification. Analyzing the isotopic composition of N₂O, especially the ¹⁵N site preference of N₂O produced (*SP_{N2O}*), showed that N₂O of pure bacterial or fungal cultures differed in *SP_{N2O}* values, which might enable the quantification of fungal N₂O based on the isotopic endmember signatures of N₂O produced by fungi and bacteria.~~
35 ~~This study aimed to identify the fungal contribution to N₂O emissions and determine under anaerobic conditions in incubated repacked soil samples by using different approaches to disentangle sources of N₂O. Three approaches were established (modified substrate induced respiration with selective inhibition (SIRIN) approach, endmember mixing approach (IEM) and the *SP/δ¹⁸O* mapping approach (*SP/δ¹⁸O* Map) to independently investigate the fungal fraction contributing to N₂O from denitrification.~~ Three soils were incubated ~~with~~ ~~four treatments of the SIRIN approach~~ under anaerobic conditions to promote denitrification ~~with four treatments of the a modified substrate~~

~~induced respiration with selective inhibition (SIRIN) approach.~~ While one treatment without microbial inhibition served as a control, the other three treatments were amended with inhibitors to selectively inhibit bacterial, fungal or bacterial and fungal growth. These treatments were performed in three varieties. In one variety, the ^{15}N tracer technique was used to estimate the effect of N_2O reduction on N_2O produced, while two other varieties were performed under natural isotopic conditions ~~but~~ with and without acetylene.

~~Three approaches were established to estimate the N_2O production by a fungal community in soil: i) A modification of the SIRIN approach was used to calculate N_2O evolved from selected organism groups, and ii) $SP_{\text{N}_2\text{O}}$ values from the acetylated treatment were used in the isotope endmember mixing approach (IEM), and iii) the $SP/\delta^{18}\text{O}$ mapping approach ($SP/\delta^{18}\text{O}$ Map) was used to estimate the fungal contribution to N_2O production and N_2O reduction under anaerobic conditions from the non-acetylated treatment.~~

~~All three approaches revealed a small fungal contribution to N_2O fluxes (f_{FD}) under anaerobic conditions in the soils tested. Quantifying the fungal fraction with modified SIRIN was not successful –due to large amounts of uninhibited N_2O production. In only one soil, f_{FD} using modified SIRIN could be estimated using modified SIRIN and resulted in $28\pm 9\%$, which was possibly overestimated as results obtained by IEM and $SP/\delta^{18}\text{O}$ Map for this soil resulted in f_{FD} of below 15 and 20 %, respectively. As a consequence of the unsuccessful SIRIN approach, estimation of fungal $SP_{\text{N}_2\text{O}}$ values was impossible. For this soil, $SP_{\text{N}_2\text{O}}$ values of the fungal fraction determined with modified SIRIN could be compared with fungal $SP_{\text{N}_2\text{O}}$ endmember values previously reported in the literature and indicated...~~

~~The three approaches tested revealed a small fungal contribution to N_2O fluxes under anaerobic conditions in the soils tested. Quantifying the fungal fraction with modified SIRIN was only possible in one soil and totaled 0.28 ± 0.09 . This was higher than the results obtained by IEM and $SP/\delta^{18}\text{O}$ Map, which accounted zero to 0.20 of N_2O produced to the fungal community.~~

~~To our knowledge, this study was the first attempt to quantify the fungal contribution to anaerobic N_2O production by simultaneous application of three approaches, i.e. modified SIRIN, IEM and $SP/\delta^{18}\text{O}$ Map. While all successful methods coincided by suggesting a small or missing fungal contribution, further studies under conditions ensuring with stimulated larger fungal N_2O fluxes by adding fungal C substrates sources preferred by fungi and an improved modified SIRIN approach, including alternative inhibitors, are needed to better cross-validate the methods.~~

1. Introduction

The greenhouse gas nitrous oxide (N_2O) contributes to global warming and to the depletion of the ozone layer in the stratosphere (Crutzen, 1970; IPCC, 2013). The largest anthropogenic N_2O emissions originate from agricultural soils and are mainly produced during microbial nitrification, nitrifier denitrification and denitrification (Firestone and Davidson, 1989; Bremner, 1997; IPCC, 2013; Wrage-Mönnig et al., 2018). In order to find mitigation strategies for N_2O emissions from arable soils, it is important to understand N_2O sources and sinks and thus improve knowledge about the production pathways and the microorganisms involved.

~~Denitrification describes describes the stepwise reduction of nitrate (NO_3^-) to dinitrogen (N_2), with the intermediates nitrite (NO_2^-), nitric oxide (NO) and N_2O (Knowles, 1982). While this entire reaction chain including the ability to reduce N_2O to N_2 is found among bacterial denitrifiers, most fungi lack N_2O reductase (Nos).~~ For a long time, it was believed that solely bacteria are involved in N_2O formation during denitrification (Firestone and Davidson, 1989); however, also several fungi are capable of denitrification (Bollag and Tung, 1972; Shoun et al.,

1992). Denitrification describes the reduction of nitrate (NO_3^-) to dinitrogen (N_2), with the intermediates nitrite (NO_2^-), nitric oxide (NO) and N_2O (Knowles, 1982). While this entire reaction chain including the ability to reduce N_2O to N_2 is found among bacterial denitrifiers, most fungi lack N_2O reductase (Nos). Recently, pure culture studies showed that N_2O from fungal denitrification was often accompanied with N_2O from abiotic production (Phillips et al., 2016a; Phillips et al., 2016b), which may lead to overestimate the importance of fungal N_2O production. Other studies indicated that although only some fungal species (e.g. *Fusarium* strains) are performing respiratory denitrification, these may produce substantial amounts of N_2O performing respiratory denitrification with substantial amounts of N_2O production (Higgins et al., 2018; Keuschnig et al., 2020). Even though only a few fungal species were identified to be capable of respiratory denitrification, N_2O produced by fungi may thus contribute largely to N_2O from denitrification in soil, since, firstly, fungi dominate the biomass in soil (up to 96 %) compared to bacteria in general and thus fungi could potentially play a dominant role in N_2O production (Ruzicka et al., 2000; Braker and Conrad, 2011). Thus, a respiratory fungal-to-bacterial (F:B) ratio of 4 is typical for arable soils (Anderson and Domsch, 1975; Blagodatskaya and Anderson, 1998). Secondly, the fact that due to a lacking N_2O reductase (Nos) (Shoun et al., 1992; Shoun et al., 2012; Higgins et al., 2018), N_2O is the major end product of fungal denitrification. This led to the assumption that the potential activity of fungal N_2O production in soil may exceed that of bacteria, provided that both microbial groups have the same specific denitrification activity (Shoun et al., 1992; Sutka et al., 2008). However, co-occurring processes may also contribute to N_2O production, such as co-denitrification, i.e. a hybrid N_2O is formed using one N atom from NO_2^- and one N atom from compounds like azide or ammonium (NH_4^+) (Shoun and Tanimoto, 1991; Shoun et al., 1992; Tanimoto et al., 1992; Spott et al., 2011), and also abiotic N_2O formation (Phillips et al., 2016a; Phillips et al., 2016b). This could potentially lead to overestimation of the importance of fungal N_2O production. Although there are methodological approaches to disentangle sources of N_2O , it is still challenging to clearly attribute N_2O emitted from soil to bacterial or fungal denitrification.

One approach to differentiate between N_2O produced by fungi and bacteria during denitrification comprises the addition of two antibiotics to soil incubation experiments, i.e. streptomycin and cycloheximide to inhibit bacterial or fungal protein biosynthesis, i.e. growth, respectively. Co-denitrification was found to often co-occur with fungal denitrification. During this fungal pathway, a hybrid N_2O is formed using one N atom from NO_2^- and one N atom from compounds like azide or ammonium (NH_4^+) for N_2O production. A ^{15}N tracing approach was used to identify and quantify co-denitrification, which contributed about 92% to N_2O produced in an incubation experiment with a grassland soil under anaerobic conditions (Laughlin and Stevens, 2002). This again stresses the large potential N_2O production by fungi. However, in pure culture studies, not only co-denitrification, but also abiotic N_2O formation may co-occur with fungal denitrification (Phillips et al., 2016a; Phillips et al., 2016b; Rohe et al., 2017) contribute to N_2O production but potentially ion of pathway differentiation is still challenging.

Soil incubation experiments could serve to differentiate between N_2O produced by fungi and bacteria during denitrification by the application of two antibiotics: streptomycin and cycloheximide, which inhibit bacterial or fungal growth, respectively, by inhibition of the protein biosynthesis. This method is known as substrate induced respiration with selective inhibition (SIRIN) and was originally developed to determine the bacterial or fungal contribution to CO_2 respiration (Anderson and Domsch, 1975). A few studies used a modification of this method for N_2O analysis (Laughlin and Stevens, 2002; Crenshaw et al., 2008; Blagodatskaya et al., 2010; Long et al., 2013) and found a greater decrease of N_2O production with fungal than with bacterial growth inhibition (i.e.g. 89 vs. 23 % decrease, respectively (Laughlin and Stevens, 2002)). This indicated, indicating that fungi might

dominate N₂O production (Laughlin and Stevens, 2002; McLain and Martens, 2006; Crenshaw et al., 2008; Blagodatskaya et al., 2010; Long et al., 2013; Chen et al., 2014; Chen et al., 2015). However, difficulties of this method may be to achieve complete inhibition of selective groups (Ladan and Jacinthe, 2016) and to avoid shifts in the structure of microbial communities as response of pre-incubation or duration of experiments.

Another opportunity to distinguish between N₂O from bacterial and fungal denitrification and other pathways analysing is the analysis of the isotopic composition of N₂O might be a promising tool to distinguish between N₂O from bacterial and fungal denitrification and other pathways.

Especially, the isotopomer ratios of N₂O (i.e. N₂O molecules with the same bulk ¹⁵N isotopic enrichment but showing different positions of ¹⁵N in the linear N₂O molecule (Ostrom and Ostrom, 2017)) in pure culture studies showed differences in N₂O of bacterial and fungal denitrification (Sutka et al., 2006; Sutka et al., 2008; Frame and Casciotti, 2010; Rohe et al., 2014a; Rohe et al., 2017). and might be suitable for distinguishing between N₂O produced by bacteria or fungi under denitrifying conditions.

Isotopomer ratios of N₂O can be expressed as ¹⁵N site preference (*SP*_{N₂O}), i.e. the difference between δ¹⁵N of the central and terminal N-position of the asymmetric N₂O molecule (Toyoda and Yoshida, 1999). The *SP*_{N₂O} values of N₂O of six pure fungal cultures was between 16 and 37 ‰ (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017), whereas several bacterial cultures produced N₂O with *SP*_{N₂O} values between -7.5 and +3.5 ‰ during denitrification (Toyoda et al., 2005; Sutka et al., 2006; Rohe et al., 2017). While it is

generally assumed that *SP*_{N₂O} values of N₂O produced by fungal pure cultures during denitrification are transferable to N₂O produced by fungal soil communities, this has not yet been proven. Until now, studies reporting possible ranges of fungal contributions to N₂O fluxes from soil were based on *SP*_{N₂O} values of pure cultures (Köster et al., 2013b; Zou et al., 2014; Lewicka-Szczepak et al., 2017; Senbayram et al., 2018; Senbayram et al., 2020; Lewicka-Szczepak et al., 2014), but uncertainty of this approach arose from the large ranges of fungal *SP*_{N₂O} values (Sutka et al., 2008; Maeda et al., 2015; Rohe et al., 2017). It would thus be useful to constrain fungal *SP*_{N₂O} values for a specific soil or soil type.

However, the *SP*_{N₂O} value of N₂O produced by pure bacterial cultures during nitrification is approximately 33 ‰ and thus interferes with that *SP*_{N₂O} values of fungal denitrification (Sutka et al., 2006; Sutka et al., 2008; Rohe et al., 2014a). This demonstrates the difficulty to use solely *SP*_{N₂O} values as an indicator for different organism groups contributing to N₂O production from soil, where different pathways may co-occur.

~~While it is generally assumed that *SP*_{prodN₂O} values of N₂O produced by fungal pure cultures during denitrification is transferable to N₂O produced by fungal soil communities, this has not yet been proven. Until now, studies reporting possible ranges of fungal contributions to N₂O fluxes from soil were based on *SP*_{prodN₂O} values of pure cultures, but uncertainty of this approach arose from the fact that the full range of *SP*_{prodN₂O} values is between 16 and 37 ‰ have been reported. It would thus be useful to constrain fungal *SP*_{prodN₂O} values for a specific soil or soil type.~~

Based on the above cited ranges for the isotopomer endmembers of fungal and bacterial denitrification, and assuming that only fungal and bacterial denitrification are responsible for N₂O production, the fraction of fungal N₂O can be calculated using the isotope endmember mixing approach (IEM) with *SP*_{N₂O} values of N₂O produced in soil (*SP*_{prod}), provided N₂O reduction does not occur (Ostrom et al., 2010; Ostrom and Ostrom, 2011).

If there is a N₂O reduction, *SP*_{N₂O} and also δ¹⁵N and δ¹⁸O values of produced N₂O (δ¹⁵N^{bulk}_{N₂O} and δ¹⁸O_{N₂O}, respectively) are affected by isotopic fractionation (Ostrom et al., 2007; Ostrom and Ostrom, 2011). This means that the ¹⁴N¹⁶O bond of N₂O is preferentially broken compared to ¹⁴N¹⁸O or ¹⁵N¹⁶O, resulting in residual N₂O that is relatively isotopically enriched in ¹⁵N and ¹⁸O and shows larger *SP*_{N₂O} values compared to *SP*_{N₂O} values of N₂O from denitrification without the reduction step (Popp et al., 2002; Ostrom et al., 2007).

165 In controlled laboratory experiments, the N_2O reduction to N_2 can be inhibited using acetylene (C_2H_2) during anaerobic incubation experiments (Yoshinari and Knowles, 1976; Groffman et al., 2006; Well and Flessa, 2009; Nadeem et al., 2013). Hence, C_2H_2 inhibition might be suitable to quantify SP_{prod} values in soils exhibiting significant N_2O reduction and would thus allow quantification of fungal N_2O fluxes based on SP_{prod} values. However, problems due to incomplete inhibition of N_2O reduction and unwanted inhibition of other pathways may occur (Wrage et al., 2004b; Wrage et al., 2004a). Another possibility to quantify N_2O reduction to N_2 during 170 denitrification is also possible with ^{15}N tracing experiments using ^{15}N enriched substrates and analysing $^{15}N_2$ fluxes (Well et al., 2006; Lewicka-Szczebak et al., 2014). The ^{15}N tracer approaches also enables to distinguish between N_2O from fungal denitrification and co-denitrification, i.e. a hybrid N_2O is formed using one N atom from NO_2^- and one N atom from compounds like azide or ammonium (NH_4^+) for N_2O production (Tanimoto et al., 1992; Laughlin and Stevens, 2002; Rohe et al., 2017).

175 The N_2O reduction can be quantified using N_2O natural abundance isotopic signatures (i.e. N_2O with differing number or positions of N or O isotopes (Ostrom and Ostrom, 2017)), which also enables simultaneous differentiation of selected pathways producing N_2O (i.e. N_2O with differing number or positions of N or O isotopes). Here, the isotope mapping approach uses isotope fractionation factors together with $\delta^{15}N$ values of precursors ($\delta^{15}N_{NOx}$) as well as $\delta^{15}N^{bulk}_{N_2O}$ and SP_{N_2O} values of N_2O produced were used (Toyoda et al., 2011). 180 Recently, this isotope mapping approach was further developed (SP/ $\delta^{18}O$ Map) using $\delta^{18}O_{N_2O}$ and SP_{N_2O} values and $\delta^{18}O$ values of precursors (Lewicka-Szczebak et al., 2017) by using and different slopes of N_2O reduction and mixing lines in the $\delta^{18}O$ – SP isotope plot. While SP_{N_2O} values are independent of isotopic signatures of the precursors, $\delta^{15}N^{bulk}_{N_2O}$ and $\delta^{18}O_{N_2O}$ result from the isotopic signature of the precursor and isotopic fractionation during N_2O production (Toyoda et al., 2005; Frame and Casciotti, 2010). Regarding $\delta^{18}O_{N_2O}$, a complete exchange 185 of oxygen (O) between NO_3^- and soil water can be assumed and consequently, one can use the $\delta^{18}O$ values of soil water for interpretation of $\delta^{18}O_{N_2O}$ values (Kool et al., 2009; Snider et al., 2009; Lewicka-Szczebak et al., 2016). However, interpretation of $\delta^{18}O_{N_2O}$ values from different microbial groups may be more complex due to incomplete O exchange, because variations in the extent of O exchange between water and N oxides affect the final $\delta^{18}O_{N_2O}$ value (Garber and Hollocher, 1982; Aerssens et al., 1986; Kool et al., 2007; Rohe et al., 2014b; Rohe et al., 2017). 190 Importantly, fungal and bacterial N_2O showed different ranges for $\delta^{18}O_{N_2O}$ values, hence this isotopic signature may also be helpful in differentiation of these pathways (Lewicka-Szczebak et al., 2016). This SP/ $\delta^{18}O$ Map approach thus allows for differentiation and estimation of the contributions of isotope effects due to N_2O reduction and admixture of fungal N_2O . Hence, N_2O reduction can be estimated together with the N_2O mixing due to application of two isotopic signatures of N_2O . Thus, the N_2O reduction to N_2 does not affect the outcome of the 195 SP/ $\delta^{18}O$ Map.

200 In controlled laboratory experiments the N_2O reduction to N_2 can be inhibited using acetylene (C_2H_2) during anaerobic incubation experiments (Yoshinari and Knowles, 1976; Groffman et al., 2006; Well and Flessa, 2009; Nadeem et al., 2013). Hence, C_2H_2 inhibition might be suitable to quantify SP_{prod} values in soils exhibiting significant N_2O reduction and would thus allow quantification of fungal N_2O fluxes based on SP_{prod} values. Another possibility to quantify the N_2O reduction to N_2 during denitrification is also possible with ^{15}N tracing experiments using ^{15}N enriched substrates and analysing $^{15}N_2$ fluxes (Well et al., 2006; Lewicka-Szczebak et al., 2014). The ^{15}N tracer approaches also enables to distinguish between N_2O from fungal denitrification and co-denitrification (Laughlin and Stevens, 2002; Rohe et al., 2017). In a laboratory experiment using a grassland soil under anaerobic conditions, co-denitrification was found to contribute about 92 % to N_2O produced, while only 8

~~205 % resulted from denitrification (Laughlin and Stevens, 2002). This again stresses the large potential N₂O production by fungi and also the need of pathway differentiation.~~

~~210 Although SP_{N_2O} values are independent of isotopic signatures of the precursors, $\delta^{15}N$ and $\delta^{18}O$ values of produced N₂O ($\delta^{15}N^{bulk}_{N_2O}$ and $\delta^{18}O_{N_2O}$, respectively) result from the isotopic signature of the precursor and isotopic fractionation during N₂O production (Toyoda et al., 2005; Frame and Caseiotti, 2010). (Lewicka-Szezebak et al., 2014; Kool et al., 2009; Snider et al., 2009) Interpretation of $\delta^{18}O_{N_2O}$ values is even more complex due to incomplete O exchange, because O exchange during denitrification between water and denitrification intermediates alters affect the final $\delta^{18}O_{N_2O}$ value (Garber and Hollocher, 1982; Aerssens et al., 1986; Kool et al., 2007; Rohe et al., 2014b; Rohe et al., 2017). However, recently, fungal and bacterial N₂O showed different ranges for $\delta^{18}O_{N_2O}$ values and this isotopic signature may also be helpful in differentiation of these pathways (Lewicka-Szezebak et al., 2016).~~

~~215 Moreover, $\delta^{15}N^{bulk}_{N_2O}$, $\delta^{18}O_{N_2O}$ and SP_{N_2O} values are in the course of denitrification affected by isotopic fractionation due to N₂O reduction. During N₂O reduction, the ¹⁴N¹⁶O bond is preferentially broken compared to ¹⁴N¹⁸O or ¹⁵N¹⁶O, resulting in residual N₂O, that is relatively isotopically enriched in ¹⁵N and ¹⁸O and shows larger SP_{N_2O} values compared to SP_{N_2O} values of N₂O from denitrification without the reduction step (Popp et al., 2002; Ostrom et al., 2007). Quantification of N₂O reduction to N₂ during denitrification is possible by and analyzing ¹⁵N₂ fluxes in ¹⁵N tracing experiments using ¹⁵N enriched substrates (Well et al., 2006; Lewicka-Szezebak et al., 2014). To quantify N₂O reduction and the pathways producing N₂O based on N₂O isotopocules (i.e. N₂O with differing number or positions of N or O isotopes (Ostrom and Ostrom, 2017)), in latter i.e. the isotope mapping approach was developed using isotope fractionation factors together with $\delta^{15}N^{bulk}$ values of N₂O precursors ($\delta^{15}N_{NO_3}$) as well as $\delta^{15}N^{bulk}_{N_2O}$ and SP_{N_2O} values of N₂O produced (Toyoda et al., 2011). Recently, this isotope mapping approach was further developed (SP/ $\delta^{18}O$ Map) using $\delta^{18}O_{N_2O}$ and SP_{N_2O} values of N₂O and $\delta^{18}O$ values of precursors (Lewicka-Szezebak et al., 2014; Lewicka-Szezebak et al., 2017). This approach uses different slopes of N₂O reduction and mixing lines in the $\delta^{18}O$ —SP isotope plot and allows for differentiation of isotope effects due to N₂O reduction and admixture of fungal N₂O. Hence, N₂O reduction can be estimated together with the N₂O mixing due to application of two isotopic signatures of N₂O. For the SP/ $\delta^{18}O$ Map, the inhibition of N₂O reduction is not needed.~~

~~230 Based on the above cited ranges for the isotopomer endmembers of fungal and bacterial denitrification, and assuming that only fungi and bacteria are responsible for N₂O production, the fraction of fungal N₂O can be calculated using the isotope endmember mixing approach (IEM) with SP_{N_2O} values of N₂O produced in soil (SP_{prod}), provided N₂O reduction, which is altering SP_{N_2O} values of emitted N₂O, does not occur (Ostrom et al., 2010; Ostrom and Ostrom, 2011). This can be ensured in laboratory experiments by inhibiting N₂O reduction to N₂ using acetylene (C₂H₂) during anaerobic incubation experiments (Yoshinari and Knowles, 1976; Groffman et al., 2006; Well and Flessa, 2009; Nadeem et al., 2013). Hence, C₂H₂ inhibition might be suitable to quantify SP_{prod} values in soils exhibiting significant N₂O reduction and would thus allow quantification of fungal N₂O fluxes based on SP_{prod} values. For the SP/ $\delta^{18}O$ Map, the inhibition of N₂O reduction is not needed. Hence, N₂O reduction can be estimated together with the N₂O mixing due to application of two isotopic signatures of N₂O. While it is generally assumed that SP_{prod} values of N₂O produced by fungal pure cultures is transferable to N₂O produced by fungal soil communities, this has not yet been proven. Until now, studies reporting possible ranges of fungal contributions to N₂O fluxes from soil were based on SP_{prod} values of pure arose from the fact that the full range of~~

245 SP_{prod} values is between 16 and 37%. It would thus be useful to constrain fungal SP_{prod} values for a specific soil or soil type.

So far, the described methods for distinguishing between fungal and bacterial N_2O emission have not been evaluated and compared in the same soil and their accuracy and possible bias remains unknown. A better knowledge of the comparability of the methods would enable comparison of results of studies using different methods and thus further improve our understanding of processes of N_2O production. It would also reveal weaknesses of approaches and might lead to the development of better methods.

250 Therefore, this study aims at (i) determining the fungal contribution to N_2O production by denitrification under anoxic conditions and glucose addition using three arable soils and three approaches (modified SIRIN, IEM and the $SP/\delta^{18}O$ Map), and to assess their usefulness in soil studies and thus assess factors of potential bias of the methods and (ii) to estimate the SP_{N_2O} values from fungal soil communities and thus to evaluate the transferability of the pure culture range of the fungal SP_{N_2O} endmember values. We hypothesized that the fungal fraction contributing to N_2O from denitrification in different soils using a modified SIRIN approach and isotopic methods will be correlated but not exactly matched due to limited inhabitability of microbial communities and variability in SP_{N_2O} endmember values. Furthermore, successful application of the modified SIRIN approach with determined fungal fraction contributing to N_2O from denitrification will yield fungal SP_{N_2O} endmember values within the range of values previously reported in the literature.

255 Therefore, this study aims at (i) determining the fungal contribution on to N_2O production by denitrification under anoxic conditions and glucose addition using three arable soils and three approaches (modified SIRIN, IEM and the $SP/\delta^{18}O$ Map), and to assess their usefulness in soil studies and thus assess factors of potential bias of the methods and (ii) to estimate the SP_{N_2O} values from a fungal soil communities and thus to evaluate the transferability of the pure culture range of the fungal SP_{N_2O} endmember values. - modified SIRIN, IEM and the $SP/\delta^{18}O$ Map, (ii) to compare the fungal contribution on N_2O production determined by these approaches and thus assess factors of potential bias of the methods, and (iii) to estimate the SP_{N_2O} values from a fungal soil community and thus to evaluate the transferability of the pure culture range of the fungal SP_{N_2O} endmember values.

270 2. Materials and Methods

2.1 Soil samples

275 All experiments were conducted with three arable soils differing in texture to provide different conditions for denitrification, soils differing in texture, C_{org} content, C/N ratio and pH. Thus, it was assumed that the soils harbour different denitrifying communities, i.e. different fractions of bacteria and fungi contributing to denitrification. One of the soils was sampled during a second season to evaluate if the fungal fraction contributing to N_2O production is soil-specific or can be subject to seasonal change of microbial communities. As one this soil was sampled at two different time points, we conducted four experiments and named the different experiments "Soil 1.1", "Soil 1.2", "Soil 2", and "Soil 3": - Experiment Soil 1.1 and Soil 1.2 with loamy sand (Soil 1) sampled in December 2012 and in June 2011, respectively, Experiment Soil 2 with sand sampled in January 2013, Experiment and Soil 3 with silt loam sampled in December 2012, and Experiment 4 with loamy sand sampled in June 2011 (Table 1).


280 Soil samples of the upper 30 cm were collected in plastic bags aerated via cotton wool stoppers and stored at 6 °C for maximally two months. To get information about the initial soil status, the mineral nitrogen content (N_{min}) of

soil samples was determined before and after fertilization by extracting NO_3^- and NH_4^+ with 0.01 M calcium chloride dihydrate ($\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$) according to ISO 14255 and analysing NO_3^- and NH_4^+ concentrations in the extracts with a Continuous-Flow-Analyser (SKALAR, Germany) directly after sample collection. ~~To get information about the initial soil status,~~ Other soil characteristics (C and N content, soil pH value, isotopic values of soil NO_3^- and NO_2^-) were analysed with samples of *Soil 1.1*, *Soil 2* and *Soil 3*. Total contents of C and N in soil samples were ~~analyzed~~ analysed by dry combustion of ~~grinded~~ ground samples (LECO TruSpec, Germany). The soil pH was measured in 0.01 M CaCl_2 . ~~The mineral nitrogen content (N_{min}) of soil samples was determined before and after fertilization by extracting NO_3^- and NH_4^+ with 0.01 M calcium chloride dihydrate ($\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$) according to ISO 14255 and analyzing NO_3^- and NH_4^+ concentrations in the extracts with a Continuous Flow-Analyzer (SKALAR, Germany).~~ The $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values of NO_3^- and NO_2^- ($\delta^{15}\text{N}_{\text{NO}_x}$ and $\delta^{18}\text{O}_{\text{NO}_x}$, respectively) in soil extracts (with 0.01 M calcium chloride dihydrate ($\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$)) were ~~analyzed~~ analysed by the bacterial denitrifier method (Casciotti et al., 2002) (Table 1).

The three soils were also sampled in summer 2010 for pre-experiments to gain information on the ~~R~~ respiratory biomass ~~of the three soils was analyzed with~~ by analyzing the substrate induced respiration (SIR) according to Anderson and Domsch (1978) and the respiratory F:B ratio was ~~analyzed~~ analysed with substrate induced respiration with selective inhibition (SIRIN) ~~in summer 2010~~ by a computer-generated selectivity analysis: "SIR-SBA 4.00" (Heinemeyer, copyright MasCo Analytik, Hildesheim, Germany) (Anderson and Domsch, 1975) (Table 1). The scheme of glucose and growth inhibitor combinations is listed below in section "Methodological Methodical approach". ~~For further~~ The characteristics of the soils, ~~see are listed in~~ Table 1.

2.2 Methodological Methodical approach

The experimental setup with various measures is presented in the following sections and illustrated in Figure 1. Important terms used and its descriptions are listed in Supplementary Material, Table S1.






Figure 1: The methodical approach comprised a pre-experiment with substrate induced respiration (SIR) to estimate the optimal glucose concentration ($c_{opt}(\text{glucose})$) and the fungal-to-bacterial ration in the soil (F:B ratio), and the substrate induced respiration with selective inhibition approach (SIRIN) to determine the optimal inhibitor concentration ($c_{opt}(\text{streptomycin})$ and $c_{opt}(\text{cycloheximide})$). The initial soil status, i.e. ammonium and nitrate concentration of the soil ($c(\text{NH}_4^+)$ and $c(\text{NO}_3^-)$, respectively), was measured in N_{min} extracts and the isotopic signature of soil NO_3^- was analysed by the denitrifier method. The incubation experiment comprised the SIRIN approach with three experimental varieties: without acetylene (-C₂H₂), with C₂H₂ (+C₂H₂), and without C₂H₂ but with ¹⁵N labelled NO_3^- (traced), while NO_3^- with natural isotopic composition was added to the other two varieties. Produced gas was analysed for its concentration ($c(\text{CO}_2)$ and $c(\text{N}_2\text{O})$) using gas chromatography (GC) and N_2O was further analysed by isotope ratio mass spectrometry (IRMS) for its isotopic composition. Please refer to the Material & Methods section for more information.

2.2.1 SIRIN pre-experiment

As in most studies applying the SIRIN method on N_2O emissions (e. g. Laughlin and Stevens, 2002; Chen et al., 2014; Ladan and Jacinthe, 2016), a pre-experiment was conducted with samples collected in 2010, in order to get information about optimal substrate and inhibitor concentrations for substrate induced respiration with growth inhibition. The pre-experiments of the present study were conducted in two steps- as described in the original methods, i.e. -CO₂ production under oxic conditions was analysed to estimate the substrate induced respiration by the SIR method (Anderson and Domsch, 1978) and the substrate induced respiration with selective inhibition by the SIRIN method (Anderson and Domsch, 1975) as follows.

In a first pre-experiment (Figure 1), the SIR method (Anderson and Domsch, 1978) was used to get information about the amount of respiratory biomass in soil- under oxic conditions. In this pre-experiment glucose served as substrate to initiate microbial growth (Anderson and Domsch, 1975). To this end, we added different concentrations of glucose (0.50, 0.75, 1.0, 1.5, 2.0, 3.0, 4.0, 5.0, 6.0 mg g⁻¹ dry weight (dw) soil) to find the optimal glucose concentration ($c_{opt}(\text{glucose})$), which is the glucose concentration that causes maximum initial respiration rates by analysing CO₂ production (Anderson and Domsch, 1978). $c_{opt}(\text{glucose})$ was 1.0 mg g⁻¹ for Experiment Soil 2 (sand) and 1.5 mg g⁻¹ for Experiment Soils 1, 3 and 4 (loamy sand and silt loam). Glucose served as substrate to initiate microbial growth (Anderson and Domsch, 1975).

We conducted In a second pre-experiment (Figure 1), the SIRIN method was used according to Anderson and Domsch (1975) for determining the respiratory F:B ratio- according to Anderson and Domsch (1975). The $c_{opt}(\text{glucose})$ determined in the first pre-experiment was used, while s Selectivity of the inhibitor combinations of streptomycin (bacterial respiratory inhibitor) and cycloheximide (fungal respiratory inhibitor) were tested with the following three concentrations, (0.75, 1.0, 1.5 mg g⁻¹ dw, respectively). The optimal concentration for inhibition of fungal respiration was 0.75 mg g⁻¹ dw soil cycloheximide ($c_{opt}(\text{cycloheximide})$) and for bacterial respiratory

340 inhibition 1.0 mg g^{-1} dw soil streptomycin ($c_{opt}(\text{streptomycin})$). ~~According to As~~ in the first pre-experiment, CO_2 production under oxic conditions was analysed. The determined optimal concentrations of glucose, streptomycin and cycloheximide were used in the modified SIRIN approach, on the assumption that concentrations optimal for CO_2 respiration also allow denitrification. Examples of respiration curves derived from SIR and SIRIN pre-experiments are represented in Figure S1 and S2, respectively.

Table 1: Soil characteristics of three arable soils from Germany used for incubation experiments (~~Exp-Soil~~) (standard deviation in brackets). Except for NH₄⁺ and NO₃⁻, soil characteristics (C, N, pH, δ¹⁵N_{NO_x} and δ¹⁸O_{NO_x}) of loamy sand were only analysed once for samples collected in 2012.

<u>Soil (Year)</u>	<u>Soil texture</u>	<u>Soil type (WRB)</u>	<u>Location</u>	<u>C content [%]</u>	<u>N content [%]</u>	<u>NH₄⁺ [mg N kg⁻¹]</u>	<u>NO₃⁻ [mg N kg⁻¹]</u>	<u>pH (CaCl₂)</u>	<u>δ¹⁵N_{NO_x} [‰]^c</u>	<u>δ¹⁸O_{NO_x} [‰]^c</u>	<u>F:B^f</u>	<u>Biomass^g [µg C gdw⁻¹ soil]</u>
1.1 (2012)	Loamy sand	Haplic Luvisol	Braunschweig ^a	1.43 (<0.01)	0.10 (<0.01)	0.4 (<0.1)	14.1 (2.1)	5.67	3.98	-4.82	2.6	234
4-1.2 (2011)	sand	Luvisol		/	/	1.0 (0.4)	11.0 (0.3)	/	/	/	/	/
2 (2013)	Sand	Gleyic Podzol	Wennebostel ^b	2.31 (0.04)	0.14 (<0.01)	1.9 (0.2)	6.6 (0.2)	5.54	0.73	-2.68	2.6	161
3 (2013)	Silt loam	Haplic Luvisol	Göttingen ^c	1.62 (0.02)	0.13 (<0.01)	n.d. ^d	22.7 (<0.1)	7.38	4.18	2.32	4.9	389

^aExperimental Station of the Friedrich-Löffler Institute, Braunschweig, Germany

^bprivate agricultural field North of Hannover, water protection area Fuhrberger Feld, Germany

^cReinshof Experimental Farm, Georg-August-University, Göttingen, Germany

^dnot detectable (i.e. below detection limit of 0.06 mg kg⁻¹ NH₄⁺-N)

^eIsotopic values of natural soil NO₃⁻ using the denitrifier method (Casciotti et al., 2002).

^fRespiratory fungal-to-bacterial (F:B) ratio ~~analyzed~~ analysed by SIRIN method (Anderson and Domsch, 1973, 1975) in a pre-experiment in 2010.

^gRespiratory biomass ~~analyzed~~ analysed by CO₂ production from SIR method (Anderson and Domsch, 1978) in a pre-experiment in 2010..

2.2.2 Soil incubation with selective inhibition to determine N₂O forming processes

The experimental design included two ~~factors~~ approaches, (i.) microbial inhibition by fungal and/or bacterial inhibitors and (ii.) activity of N₂O reductase ~~analyzed-analysed~~ either by inhibition with C₂H₂ or quantification by ¹⁵N tracing (Figure 1). To address the microbial inhibition factor approach (i.), the SIRIN method for determination of the respiratory F:B ratio based on CO₂ emission was modified to determine N₂O production by microbial groups. However, in contrast to previous studies by Laughlin and Stevens (2002), McLain and Martens (2006), Blagodatskaya et al. (2010) and Long et al. (2013), we did not pre-incubate the soil with the growth inhibitors, as this could result in changes of the microbial community (e.g. preferential growth of selected organisms). We intended to disturb microbial communities as little as possible.

The soil was sieved (2 mm) and pre-incubated at 22 °C for five to seven days in the dark with cotton wool stoppers to allow respiration and aerobic conditions in soil bags. Four microbial inhibitor treatments (each in triplicate) with *c_{opt}*(glucose) for each soil were established:

- A Control, without growth inhibitors
- B With streptomycin sulfate (C₄₂H₈₄N₁₄O₃₆S₃) to inhibit bacterial growth
- C With cycloheximide (C₁₅H₂₃NO₄) to inhibit fungal growth
- D With streptomycin and cycloheximide, to inhibit bacterial and fungal growth

To address ~~factor~~ the other approach (ii.), all microbial inhibitor treatments were conducted in three N₂O reductase varieties, i.e.: with ¹⁵N-NO₃⁻ fertilizer (variety “*traced*”) to quantify N₂O reduction to N₂, with natural abundance NO₃⁻ and 10 kPa C₂H₂ in the headspace (variety “+C₂H₂”) to block N₂O reductase, and with natural abundance NO₃⁻ but without blocking N₂O reductase, i.e. no C₂H₂ added (variety “-C₂H₂”) (Figure 1). In total, there were 48 experimental treatments and 144 vessels (four Soils Experiments with four inhibitor treatments (A, B, C, D) and three varieties (*traced*, +C₂H₂ and -C₂H₂), each in triplicates).

The soil was adjusted to 80% water filled pore space (WFPS) with distilled water ~~and S~~ simultaneously to that, ~~the soil was~~ fertilized with ~~NO₃⁻~~ (varieties -C₂H₂, +C₂H₂, and *traced*). ~~The soil sample used with Soil 1.2 was incubated prior to the other soils and was amended with 60 mg N kg⁻¹ NaNO₃, while in agreement with other experiments conducted in our laboratory, 50 mg N kg⁻¹ KNO₃ were used with Soil 1.1, 2 and 3 (varieties -C₂H₂ and +C₂H₂ with 50 mg N kg⁻¹ KNO₃, in Experiment 1, 2 and 3 and with 60 mg N kg⁻¹ NaNO₃, in Experiment 4 and *traced* variety with 50 mg N kg⁻¹ ¹⁵N KNO₃ in Experiment 1, 2 and 3 and 60 mg N kg⁻¹ ¹⁵N KNO₃ in Experiment 4. In variety *traced*, NO₃⁻ with a ¹⁵N-enrichment labeling of 50 atom% (at%) was used.~~ For each treatment, we incubated 100 g dw soil in 850 mL preserving jars (J. WECK GmbH u. Co KG, Wehr, Germany) with gas inlet and outlet equipped with ~~three-three~~ port luer lock plastic stopcocks (Braun, Melsungen, Germany). According to the original SIRIN method (Anderson and Domsch, 1973, 1978) ~~and~~ a mixture of *c_{opt}*(glucose) and carrier material talcum (5 mg talcum g dw⁻¹) was added to soil of treatment A and together with the growth inhibitors to the soil of treatments B, C and D. The soil and additives of each treatment were mixed for 90 seconds with a handheld electric mixer. During packing, the soil density was adjusted to an expected target soil density of 1.6 g cm⁻³ in Experiment 1, Soil 1.1, 1.2, and 2 ~~and 4~~ and of 1.3 g cm⁻³ in Experiment Soil 3 to imitate field conditions. To ultimately achieve denitrifying conditions in all treatments and to avoid catalytic NO decomposition in the +C₂H₂ variety (Nadeem et al., 2013), the headspace of the closed jars was flushed with N₂ to exchange the headspace 10 times. Directly following, 85 mL of the gas in the headspace in variety +C₂H₂ were exchanged by pure C₂H₂ resulting in 10 kPa C₂H₂ in the headspace. The manual sample collection of 14 mL gas in duplicates with a plastic syringe was

performed after six, eight and ten hours (*Experiment 1 Soil 1.1*, 2 and 3) or two, four and eight hours (*Experiment 4 Soil 1.2*) of incubation time, respectively. The removed gas was replaced by the same amount of N₂.

400 2.3 Gas analysis

Gas samples were ~~analyzed~~analysed for N₂O and CO₂ concentrations ($c(N_2O)$ and $c(CO_2)$) with gas chromatography (GC, Agilent 7890A, Agilent, Böblingen, Germany) (*Figure 1*). The analytical precision of measurements was derived from analysing laboratory standards of different concentrations (0.5-1,000 ppm N₂O and 340-10,000 ppm CO₂) and resulted in a measurement precision of 1 % for N₂O and 0.5 % for CO₂. The instrumental detection limit of N₂O was $4 \mu\text{g N kg}^{-1} \text{h}^{-1}$ with a measurement precision of 1% and for of CO₂ it was $137 \mu\text{g C kg}^{-1} \text{h}^{-1}$ the detection limit was C h^{-1} with a measurement precision of 0.5%. As a control, N₂ and O₂ concentrations in the samples were ~~analyzed~~analysed with GC to ensure anaerobic conditions during the incubation for N₂O production from denitrification. CO₂ and N₂O production rates were calculated by averaging the measured N₂O production, i.e. between the time point of flushing with N₂ (t=0) and six, eight or ten hours (or two, four and eight hours with *Soil 1.2*).

The N₂O isotopic analysis of the gas samples of varieties $-C_2H_2$ and $+C_2H_2$ (*Figure 1*) were performed on a pre-concentrator (PreCon, Thermo-Finnigan, Bremen, Germany) interfaced with a GC (Trace Gas Ultra, Thermo Scientific, Bremen, Germany) and ~~analyzed~~analysed by isotope ratio mass spectrometry (IRMS, Delta V, Thermo Fisher Scientific, Bremen, Germany) (Brand, 1995; Toyoda and Yoshida, 1999; Köster et al., 2013b). A laboratory standard N₂O gas was used for calibration, having $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$, $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ and $SP_{\text{N}_2\text{O}}$ values of -1.06 ‰, 40.22 ‰, and -2.13 ‰, respectively, in three concentrations (5, 10 and 20 ppm). The analytical precision was 0.1 ‰, 0.2 ‰ and 1.5 ‰ for $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$, $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ and $SP_{\text{N}_2\text{O}}$ values, respectively. H₂O and CO₂ were trapped with magnesium perchlorate and ascarite, respectively, to prevent any interference with N₂O analysis.

The gas samples of variety *traced* from *Experiment 1 Soil 1.1*, 2, and 3 were ~~analyzed~~analysed for the 29/28 and 30/28 ratios of N₂ according to Lewicka-Szczebak et al. (2013) using a modified GasBench II preparation system coupled to ~~IRMS~~ (MAT 253, Thermo Scientific, Bremen, Germany). The gas samples of variety *traced* from *Experiment 4 Soil 1.2* were ~~analyzed~~analysed at the Centre for Stable Isotope Research and Analysis (University of Göttingen, Germany). The N₂ produced was ~~analyzed~~analysed using an elemental ~~analyzer~~analyser (Carlo Erba ANA 1500) that was coupled to dual inlet IRMS (Finnigan MAT 251) (Well et al., 1998; Well et al., 2006). Isotopic values of N₂O of *Experiment 4 Soil 1.2* (variety *traced*) were ~~analyzed~~analysed in the same lab using a pre-concentration unit coupled to IRMS (Precon-DeltaXP, Thermo Scientific, Bremen, Germany) (Well et al., 2006). Isotope ratios were used applying the non-random distribution approach to calculate the fraction of N₂ and N₂O originating from the ¹⁵N-labelled N pool as well as the ¹⁵N enrichment of that N pool (a_p) (Bergsma et al., 2001; Spott et al., 2006).

430 2.4 Inhibitor effects

For interpretation of N₂O or CO₂ production, the validity of the experimental results with respect to fungal and bacterial N₂O fluxes was checked using a flux balance comparing the sum of bacterial and fungal inhibition effects (treatments B and C) to the dual inhibition effect (treatment D):

$$D = A - [(A - B) + (A - C)] \quad (\text{Eq. 1})$$

435 With A , B , C and D representing the N_2O production rates of the last sampling time of treatment A , B , C and D , respectively. Assuming that in the other three treatments (A , B and C) non-inhibitable N_2O production was equal to treatment D , N_2O produced by bacteria ~~or~~ and fungi should show the following relation between the four treatments:

$$(A - D) = (B - D) + (C - D) \quad (\text{Eq. 2})$$

440 The fungal contribution to N_2O production during denitrification with microbial inhibition (f_{FDmi}) can be calculated, when N_2O production of treatment D is significantly smaller than N_2O production of treatments A , B and C by:

$$f_{FDmi} = \frac{(A-C)}{(A-D)} \quad (\text{Eq. 3})$$

445 A detailed discussion of inhibitor effects and difficulties with organisms that were not inhibited or abiotic sources (treatment D) is presented in section 4.1.

2.5 Isotope methods

2.5.1 Isotope endmember mixing approach (IEM)

The fungal fraction (f_{FD}) contributing to N_2O production from denitrification in soil samples was calculated according to the isotope mixing model (IEM) proposed by Ostrom et al. (2010), which was established for calculating the bacterial fraction (f_{BD}) of N_2O production. Assuming that bacteria (BD) and fungi (FD) are the only ~~sources of N_2O microorganisms responsible for denitrification~~ in soil, the ^{15}N site preference values of produced N_2O (SP_{prod}) results from the SP_{N_2O} mixing balance:

$$SP_{prod} = f_{FD} * SP_{FD} + f_{BD} * SP_{BD} \quad (\text{Eq. 4})$$

455 where f_{FD} and f_{BD} represent the fraction of N_2O produced by fungi and other N_2O sources than fungal denitrification, respectively, and SP_{FD} and SP_{BD} are the respective SP_{N_2O} endmember values (Ostrom et al., 2010; Ostrom and Ostrom, 2011). This calculation was based on the assumption that the sum of f_{BD} and f_{FD} equals 1 and that N_2O reduction to N_2 is negligible. The mean SP_{FD} value was assumed to be 33.6 ‰ (Sutka et al., 2008; Maeda et al., 2015; Rohe et al., 2014a; Rohe et al., 2017) and the SP_{BD} value from heterotrophic denitrification was assumed with minimum and maximum values from -7.5 to +3.7 ‰ (Yu et al., 2020). For this IEM approach, only results from variety + C_2H_2 could be used to calculate the fungal fraction contributing to N_2O production ($f_{FD,SP}$), as microorganisms of this variety produce N_2O that is not affected by reduction to N_2 . The $f_{FD,SP}$ contributing to N_2O production during denitrification was calculated ~~from using~~ the measured SP_{N_2O} value ~~from treatment A~~ of variety + C_2H_2 as SP_{prod} value ~~(Eq. 4) in~~ Eq. 4 that was solved for f_{FD} ($f_{FD} = 1 - ((SP_{prod} - SP_{FD}) / (SP_{BD} - SP_{FD}))$). ~~By applying this equation, a range for $f_{FD,SP}$ is received when using minimum and maximum SP_{BD} values.~~

465 Based on SP_{N_2O} values from - C_2H_2 variety, it was possible to solve Eq. 4 also to estimate the maximum potential fungal contribution to denitrification ($f_{FD,SPpot}$) assuming that we did not have any estimations for N_2O reduction. While bacterial denitrification and nitrifier denitrification would result in low SP_{N_2O} values ($SP_{BD/ND} = -10.7$ to +3.7 ‰ (Frame and Casciotti, 2010; Yu et al., 2020)), large SP_{N_2O} values would be expected from fungal denitrification and nitrification ($SP_{FB/N} = 16$ to 37 ‰ (Sutka et al., 2008; Decock and Six, 2013; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017)). N_2O reduction could have further increased the SP_{prod} values. If the contribution of this process on SP_{prod} values cannot be precisely estimated, by neglecting these effects we can determine the maximal potential fungal contribution. f_{FD} calculated from Eq. 4 (variety - C_2H_2) would thus be lower if N_2O reduction had

occurred. However, assuming the impact of N_2O reduction on SP_{N_2O} was negligible, this IEM enabled to calculate the maximum potential f_{FD} as $f_{FD, SP_{pot}} = 1 - ((SP_{N_2O} - SP_{FD/N}) / (SP_{BD/ND} - SP_{FD/N}))$.

In case successful inhibition (modified SIRIN approach), Eq. 4 was solved for the SP_{FD} value using F_{FD} , F_{BD} , and SP_{prod} values of the respective variety.

2.5.2 Product ratio [$N_2O/(N_2+N_2O)$] of denitrification

The variety *traced* served to assess N_2O reduction during denitrification in each experiment. The product ratio of denitrification [$N_2O/(N_2+N_2O)$] as given by the variety *traced* (r_{15N}) was calculated as:

$$r_{15N} = \frac{{}^{15}N_{N_2O}}{{}^{15}N_{N_2} + {}^{15}N_{N_2O}} \quad (\text{Eq. 5})$$

with ${}^{15}N_{N_2O}$ and ${}^{15}N_{N_2}$ representing N_2O and N_2 produced in the ${}^{15}N$ -labeled fertilizer pool. To check the effectiveness of C_2H_2 in blocking the N_2O reduction, r_{15N} was compared with $r_{C_2H_2}$, where the latter can be calculated from N_2O production rates of varieties $-C_2H_2$ and $+C_2H_2$:

$$r_{C_2H_2} = \frac{N_2O_{-C_2H_2}}{N_2O_{+C_2H_2}} \quad (\text{Eq. 6})$$

with $N_2O_{-C_2H_2}$ and $N_2O_{+C_2H_2}$ representing the N_2O produced in varieties $-C_2H_2$ and $+C_2H_2$, respectively.

It was possible to assess the completeness of blockage of N_2O reduction by C_2H_2 with the experimental setup as follows. If r_{15N} and $r_{C_2H_2}$ were in agreement, a complete blockage of N_2O reduction could be assumed. This enabled us to estimate reduction effects on the isotopic signatures of N_2O by comparing the $\delta\theta$ values, i.e. isotopic values of N_2O produced without N_2O reduction effects of variety $+C_2H_2$, with isotopic values of N_2O of variety $-C_2H_2$.

The information on the product ratio was used as an additional possibility to calculate the f_{FD} also for variety $-C_2H_2$. The Rayleigh-type model presented by Lewicka-Szczebak et al. (2017) and Senbayram et al. (2018) for similar closed-system incubations was used to calculate the ${}^{15}N$ site preference values of the originally produced N_2O of variety $-C_2H_2$ (SP_{prod}). SP values of emitted N_2O , i.e. after partial reduction of produced N_2O (SP_{N_2O-r}), were corrected with the net isotope effect of N_2O reduction (ηr) and the r_{15N} as follows:

$$SP_{prod} = SP_{N_2O-r} + \eta r \ln(r_{15N}) \quad (\text{Eq. 7})$$

According to Yu et al. (2020) the ηr was assumed to be -6 %. Secondly, Eq. 4 was used to calculate the f_{FD} by using SP_{prod} values of variety $-C_2H_2$ ($f_{FD, SP_{calc}}$) obtained from Eq. 7.

2.5.2.3 $SP/\delta^{18}O$ isotope mapping approach ($SP/\delta^{18}O$ Map)

The f_{FD} contributing to N_2O production from denitrification in soil samples was also estimated with the $SP/\delta^{18}O$ Map ($f_{FD, MAP}$) (Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2020). This method allows for estimating both the f_{FD} and N_2O product ratio [$N_2O/(N_2+N_2O)$] (r_{Map}). For precise estimations, the $\delta^{18}O$ values of soil water ($\delta^{18}O_{H_2O}$) applied in the experiments are needed and these values were not determined.

However, since we have independent information on the N_2O product ratio from the *traced* variety (r_{15N}), we can calculate the possible $\delta^{18}O_{H_2O}$ values of soil to get the nearest N_2O product ratios in natural and ${}^{15}N$ treatments.

The fitting of $\delta^{18}O_{H_2O}$ values ($f_{FD, MAP}$) was performed for mean, minimal and maximal values of SP_{BD} (-1.9, -7.5 and 3.7 ‰, respectively) and aimed at obtaining the minimal difference between r_{Map} and that measured in the *traced* variety, i.e.; the minimal value of $(r_{15N} - r_{Map})^2$ (according to least squares method) and that measured with for $-C_2H_2$ and $+C_2H_2$, i.e. $r_{C_2H_2}$ variety (for explanation of the product ratio see next section 2.5.2). This further allows calculation of obtaining the possible ranges for f_{FD} for particular $\delta^{18}O_{H_2O}$ fitted values (Table 4) based on the $SP/\delta^{18}O$ mapping approach (Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2020). Namely, the fitted

$\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values are applied to properly correct the $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values of the mixing endmembers (*BD* and *FD*), which depend on the ambient water. Afterwards, the corrected values of mixing endmembers are applied to calculate the f_{FD} values. The calculations with this approach may be performed assuming two different scenarios of the interplay between N_2O mixing and reduction (Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2020), but for this study both scenarios yielded almost identical results (maximal differences of 0.02 in N_2O product ratio and 2 % for f_{FD} was/were found), due to f_{BD} near 100 %. Hence, we only provide the results assuming the reduction of bacterial N_2O followed by mixing with fungal N_2O . In the following, all calculated fractions are presented in percent (%).

2.5.3 Product ratio [$\text{N}_2\text{O}/(\text{N}_2+\text{N}_2\text{O})$] of denitrification

The variety *traced* served to assess N_2O reduction during denitrification in each experiment. The product ratio of denitrification [$\text{N}_2\text{O}/(\text{N}_2+\text{N}_2\text{O})$] as given by the variety *traced* (^{15}N) was calculated as:

$$\overline{^{15}\text{N}} = \frac{^{15}\text{N}_{\text{N}_2\text{O}}}{^{15}\text{N}_{\text{N}_2} + ^{15}\text{N}_{\text{N}_2\text{O}}} \quad (\text{Eq. 5})$$

with $^{15}\text{N}_{\text{N}_2\text{O}}$ and $^{15}\text{N}_{\text{N}_2}$ representing N_2O and N_2 produced in the ^{15}N labeled fertilizer pool. To check the effectiveness of C_2H_2 to block the N_2O reduction, ^{15}N was compared with C_2H_2 , where the latter can be calculated from N_2O production rates of varieties $-\text{C}_2\text{H}_2$ and $+\text{C}_2\text{H}_2$:

$$\overline{\text{C}_2\text{H}_2} = \frac{\text{N}_2\text{O}_{-\text{C}_2\text{H}_2}}{\text{N}_2\text{O}_{+\text{C}_2\text{H}_2}} \quad (\text{Eq. 6})$$

with $\text{N}_2\text{O}_{-\text{C}_2\text{H}_2}$ and $\text{N}_2\text{O}_{+\text{C}_2\text{H}_2}$ representing the N_2O produced in varieties $-\text{C}_2\text{H}_2$ and $+\text{C}_2\text{H}_2$, respectively.

If product ratio ^{15}N and product ratio C_2H_2 were in agreement, a complete blockage of N_2O reduction could be assumed. This enabled us to estimate reduction effects on the isotopic signatures of N_2O by comparing the isotopic values of N_2O produced without N_2O reduction effects of variety $+\text{C}_2\text{H}_2$ ($\delta 0$ values) with isotopic values of N_2O of variety $-\text{C}_2\text{H}_2$.

The information on the product ratio was used as an additional possibility to calculate the also for variety $-\text{C}_2\text{H}_2$. First, the Rayleigh type model presented by Lewicka-Szczebak et al. (2017) and Senbayram et al. (2018) for similar closed system incubations, the ^{15}N site preference values of the originally produced N_2O , i.e. without its reduction to N_2O (SP_{prod}), of variety $-\text{C}_2\text{H}_2$ (SP_{prod}) was calculated by correcting SP values of emitted N_2O , i.e. after partial reduction of produced N_2O ($SP_{\text{N}_2\text{O}}$) from variety $-\text{C}_2\text{H}_2$ with the net isotope effect of N_2O reduction (ηr) and the ^{15}N as follows:

$$SP_{\text{prod}} = SP_{\text{N}_2\text{O}} + \eta r \ln(\overline{^{15}\text{N}}) \quad (\text{Eq. 7})$$

According to (Yu et al., 2020) the ηr was assumed to be 6%. Secondly, Eq. 4 was used to calculate the by using SP_{prod} values of variety $-\text{C}_2\text{H}_2$ ($_{-}\text{SP}_{\text{eate}}$) obtained from Eq. 7

2.6 Sources-Other sources of N_2O produced

Assuming that denitrification is/was the only process-producing/source of N_2O in the incubation experiment, the expected ^{15}N enrichment in N_2O produced ($^{15}\text{N}_{\text{N}_2\text{O}_{\text{exp}}}$) was given by

$$^{15}\text{N}_{\text{N}_2\text{O}_{\text{exp}}} [\text{at}\%] = \frac{(N_{\text{soil}} \times ^{15}\text{N}_{\text{nat}}) + (N_{\text{fert}} \times ^{15}\text{N}_{\text{fert}})}{N^{\text{bulk}}} \quad (\text{Eq. 8})$$

with N_{soil} , N_{fert} and N^{bulk} describing the amount of N [mg] in unfertilized soil samples (Table 1), fertilizer and fertilized soil samples, respectively, and $^{15}\text{N}_{\text{nat}}$ and $^{15}\text{N}_{\text{fert}}$ is standing for the ^{15}N enrichment under natural conditions

(0.3663 at%) and in fertilizer (50 at%), respectively. Comparison of measured ^{15}N enrichment in N_2O and $^{15}\text{N}_{\text{N}_2\text{O}_{\text{exp}}}$ gave information about the contribution of processes other than denitrification to N_2O production.

550 2.7 Statistical Analysis

We conducted several three-way analyses of variance (ANOVA) to test significant effects of soil, experimental variety and treatment on N_2O production, CO_2 production, and $SP_{\text{N}_2\text{O}}$, $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$ and $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values. The pairwise comparison with Tukey's HSD test ~~was made allowed~~ to find differences between soils, varieties and treatments influencing N_2O production, CO_2 production, and isotopic values. Significant effects of soils and treatments on $r_{\text{C}_2\text{H}_2}$ and $r_{^{15}\text{N}}$ were tested by two-way ANOVA, while differences between soils and treatments influencing the product ratios were tested with pairwise comparison with Tukey's HSD test. Effects of varieties $-\text{C}_2\text{H}_2$ and *traced* on N_2O and CO_2 production were tested by ANOVA. For this ANOVA, the N_2O production rate had to be \log_{10} -transformed to achieve homogeneity of variance and normality. The significance level α was 0.054 for every ANOVA. For some ANOVAs treatments were excluded, when replicates were $n < 3$. This was the case when only one or two samples out of three replicates could be analysed. This is denoted in the captions of tables (Table 2 and 3). The N_2O or CO_2 production rates of variety $+\text{C}_2\text{H}_2$ were followed over three sampling times by regression. For statistical analysis, we used the program R (R Core Team, 2013). Excel Solver tool was used to determine the $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values in the application of $SP/\delta^{18}\text{O}$ Map calculations.

560 3. Results

565 3.1 N_2O production rates

N_2O and CO_2 production rates of all treatments were similar in magnitude in almost all cases and mostly indistinguishable (Table 2, Figure 2). CO_2 production rates were determined to get additionally information about the denitrifying process. N_2O production rates exhibited increasing trends with ongoing incubation time for every soil with large variations within the treatments (Figure 2). Contrary to that, CO_2 production rates showed decreasing trends (Figure 42, exemplarily shown for data of variety $+\text{C}_2\text{H}_2$). Calculations of inhibitor effects were based on average N_2O and CO_2 production rates of the entire incubation period, i.e. ~~10-ten~~ hours of incubation time for ~~Experiment 1~~ Soil 1.1, 2 and 3 and ~~8-eight~~ hours for ~~Experiment 4~~ Soil 1.2.


N_2O and CO_2 production rates of all $+\text{C}_2\text{H}_2$ varieties differed significantly among soils ($P < 0.001$ ~~$P < 0.001$~~) and N_2O production rates differed also significantly among treatments ($P < 0.001$ ~~$P < 0.001$~~). Largest N_2O production rates of about ~~5.5555~~ to ~~6.1613~~ $\mu\text{g N kg}^{-1}\text{h}^{-1}$ ~~was~~ ere obtained in ~~Experiment 1~~ Soil 1.1 and 3, respectively, while in ~~Experiment~~ Soil 2 and ~~4-1.2~~ N_2O production rates were ~~lower~~ smaller (~~2.6271~~ and ~~2.7264~~ $\mu\text{g N kg}^{-1}\text{h}^{-1}$, respectively). N_2O and CO_2 production rates were significantly larger in variety $+\text{C}_2\text{H}_2$ than in variety $-\text{C}_2\text{H}_2$ of ~~Experiment 1~~ Soil 1.1, ~~1.2 3~~ and ~~3 4~~ ($P < 0.001$, $P = 0.002$, and $P < 0.001$ ~~$P < 0.010$ and $P < 0.010$~~ for N_2O production rate and $P = < 0.02701$, $P = < 0.010027$, and $P = 0.008$ for CO_2 production rate, respectively) (Table 2), while $-\text{C}_2\text{H}_2$ and $+\text{C}_2\text{H}_2$ varieties of ~~Experiment~~ Soil 2 did not differ in N_2O and CO_2 production rates ($P = 0.6402$ and $P = 0.288342$, respectively).


585

Figure 12: Time series of average N₂O and CO₂ production rates during incubation of variety +C₂H₂ at the three sample collection times of each soil (Experiment 1 Soil 1 to 34) for treatment A without growth inhibitors, B with bacterial growth inhibition, C with fungal growth inhibition, and D with bacterial and fungal growth inhibition; P-values for linear regressions (significance level $\alpha \leq 0.05$). For all significant regressions, R²-values were ≥ 0.46 and in the case of non-significance, R²-values were ≤ 0.40 .


590

n.d.: There was no detectable CO₂ production in Experiment 4 Soil 1.2 at the first sampling time after 2 hours. (Figure is continued on next page)





-Figure 4-2 continued.



595

Without blockage of N₂O reductase (variety -C₂H₂), N₂O production rates of treatment A varied significantly among experiments-Soils with mean values between 1.6175 and 3.6355 μg N kg⁻¹ h⁻¹ ($P \leq \leq 0.001$) (Table 2). In Experiment 1Soil 1.1, N₂O production rate was significantly larger (2.7272 μg N kg⁻¹ h⁻¹) than in Experiment 4Soil 1.2 (1.6175 μg N kg⁻¹ h⁻¹) ($P = 0.028$) in variety -C₂H₂. In most cases of the three varieties (-C₂H₂, +C₂H₂, and traced) The inhibitor application of each variety revealed in most cases that treatment A (without growth inhibitors) produced most N₂O, followed by either treatment B (bacterial growth inhibitor; more N₂O compared to treatment C in ExperimentSoils 1.2, 2, and 3-and 4) or treatments C (fungal growth inhibitor; more N₂O compared to

600

605 treatment B in ~~Experiment 1~~Soil 1.1). ~~Smallest N₂O production rates were in most cases found in treatment D (In~~
~~varieties -C₂H₂, +C₂H₂ and traced varieties, non-inhibitable organisms-N₂O production(treatment D) showed~~
610 ~~smallest N₂O production rates in most cases (i. e. except of for~~ variety -C₂H₂ of ~~Experiment 1~~Soil 1.1, varieties -
C₂H₂ and traced of Experiment Soil 3 and variety traced of ~~Experiment 4~~Soil 1.2). Microbial inhibitor treatments
differed significantly in N₂O fluxes of variety +C₂H₂ of each ~~experiment~~Soil (always $P \leq 0.040042$), while this
was not the case for inhibitor treatments of varieties -C₂H₂ and traced of ~~Experiment 4~~Soil 1.2 ($P = 0.154$ and
 $P = 0.154$, respectively). Significant deviations of treatments without (A) or with full inhibition (D) were found in
615 the following cases (Table 2): N₂O production rate of treatment A was significantly larger compared to the other
three treatments of ~~Experiment 1~~Soil 1.1 (+C₂H₂ and -C₂H₂), ~~Experiment Soil 2~~ (+(-C₂H₂) and Experiment Soil 3
(+C₂H₂); treatment D was significantly smaller compared to the other three treatments in ~~Experiment Soil 2~~ (+
(+C₂H₂) only and compared to treatments A and C ~~of in Experiment 1~~Soil 1.1 (+C₂H₂). A detailed discussion of
620 inhibitor effects and difficulties with organisms that were not inhibited or abiotic sources is presented in section
4.1. Comparing varieties -C₂H₂ and traced, N₂O and CO₂ rates did not differ ($P = 0.991$ for N₂O production rate
and $P = 0.490$ for CO₂ production rate, respectively), confirming that ¹⁵N-labeling did not affect N₂O and CO₂
processes.

Table 2: Average CO₂ and N₂O production rates and N₂O isotopic values of N₂O of the last sample collection with and without C₂H₂ application in the headspace (varieties -C₂H₂ and +C₂H₂) of each soil (~~Experiment 1~~ *Soil 1* ~~to 4~~) for treatments A without, B with bacterial, C with fungal, and D with bacterial and fungal growth inhibition, respectively (standard deviation in brackets, *n* = 3).

Treatment/ variety	mean N ₂ O [µg N kg ⁻¹ h ⁻¹]	mean CO ₂ [µg C kg ⁻¹ h ⁻¹]	δ ¹⁸ O _{N₂O} [‰]	δ ¹⁵ N ^{bulk} _{N₂O} [‰]	SP _{N₂O} [‰]
Experiment 1 <i>Soil 1.1</i> (Loamy sand, winter 2012)					
A / -C ₂ H ₂	272.0 (38.4)a	1233.8 (170.5)a	13.1 (0.2)a	-21.9 (1.7)a	1.6 (0.8)a
B / -C ₂ H ₂	180.9 (16.8)b	1284.8 (168.0)a	13.0 (<0.1)*	-24.2 (0.7)*	-1.3 (0.2)*
C / -C ₂ H ₂	203.1 (14.4)ab	1124.8 (54.8)a	14.6 (0.4)a	-20.0 (0.8)a	-1.6 (0.5)a
D / -C ₂ H ₂	207.8 (32.6)ab	1371.7 (35.3)a	15.2 (0.5)*	-20.2 (1.8)*	-0.3 (0.5)*
A / +C ₂ H ₂	554.9 (46.5)a	1700.9 (98.1)a	8.5 (0.1)a	-22.1 (0.3)a	-0.4 (0.3)a
B / +C ₂ H ₂	353.5 (14.0)b	1610.7 (47.2)a	7.5 (0.1)a	-26.1 (0.2)a	-1.2 (1.0)ba
C / +C ₂ H ₂	441.8 (18.5)c	1604.1 (60.3)a	9.3 (0.2)a	-22.4 (0.4)a	-0.9 (0.4)ba
D / +C ₂ H ₂	331.0 (20.5)b	1438.0 (141.9)a	7.8 (0.3)*	-24.2 (0.1)*	-2.3 (0.7)*
<i>Soil 1.2</i> (Loamy sand, summer 2011)					
A / -C ₂ H ₂	175.3 (6.6)a	2448.5 (135.8)a	25.7 (0.3)a	-30.6 (0.2)a	12.1 (1.6)a
B / -C ₂ H ₂	121.3 (74.0)a	2091.3 (19.5)b	28.0 (5.0)a	-32.3 (0.7)a	7.7 (1.4)b
C / -C ₂ H ₂	104.5 (5.3)a	1844.7 (192.1)b	29.3 (0.1)a	-30.0 (0.5)a	4.3 (1.0)c
D / -C ₂ H ₂	73.8 (63.0)a	1632.2 (115.3)b	28.9 (1.2)a	-31.8 (2.2)a	3.4 (2.0)c
A / +C ₂ H ₂	263.5 (31.7)a	2076.6 (305.3)a	13.5 (0.5)*	-34.7 (0.1)*	-1.0**
B / +C ₂ H ₂	233.0 (15.6)ab	1794.9 (238.9)a	14.3 (1.7)a	-33.8 (0.9)a	-4.9 (0.9)a
C / +C ₂ H ₂	119.5 (102.7)b	1736.8 (424.7)a	19.0 (7.0)a	-33.1 (2.8)a	-1.7 (2.7)a
D / +C ₂ H ₂	161.6 (7.6)ab	1497.0 (138.7)a	14.8 (0.5)a	-35.7 (0.2)a	-4.9 (0.7)a
Experiment 2 <i>Soil 2</i> (Sand, winter 2012)					
A / -C ₂ H ₂	315.0 (35.0)a	1316.7 (97.7)a	15.5 (1.8)a	-18.9 (2.6)a	-0.9 (2.5)a
B / -C ₂ H ₂	241.7 (3.0)b	1209.2 (24.6)a	15.0 (1.3)a	-23.4 (2.5)ab	-0.8 (<0.1)a
C / -C ₂ H ₂	247.6 (22.8)b	1201.9 (48.2)a	14.3 (0.1)a	-21.8 (0.2)ab	-1.8 (0.2)a
D / -C ₂ H ₂	198.4 (26.8)b	1102.4 (101.7)a	13.4 (0.3)a	-24.5 (0.1)ab	-1.2 (0.3)a
A / +C ₂ H ₂	270.9 (36.3)a	1271.6 (203.5)a	12.6 (0.3)a	-18.9 (4.6)a	-1.4 (0.3)a
B / +C ₂ H ₂	263.1 (19.1)a	1338.7 (71.9)a	12.3 (0.1)a	-24.6 (0.2)b	-2.0 (0.2)a
C / +C ₂ H ₂	247.3 (15.9)a	1220.2 (50.0)a	12.7 (0.1)*	-23.3 (0.2)*	-1.7 (0.4)*
D / +C ₂ H ₂	187.3 (21.8)b	1173.1 (55.1)a	12.2 (0.3)a	-26.0 (0.1)b	-1.5 (0.9)a
Experiment 3 <i>Soil 3</i> (Silt loam, winter 2013)					
A / -C ₂ H ₂	355.0 (18.4)a	1227.6 (95.2)a	26.0 (0.5)a	-20.8 (0.5)a	-0.5 (0.4)a
B / -C ₂ H ₂	325.4 (36.3)ab	1159.3 (178.2)a	24.1 (0.2)ba	-22.0 (0.2)ba	-0.1 (0.4)a
C / -C ₂ H ₂	278.9 (9.8)ab	1056.0 (59.6)a	27.3 (0.1)ba	-20.6 (0.3)a	0.6 (0.2)a
D / -C ₂ H ₂	291.1 (38.5)ab	1118.5 (70.3)a	26.3 (0.3)a	-21.0 (0.1)a	-0.04 (0.182)a
A / +C ₂ H ₂	612.8 (25.2)a	1332.5 (116.9)a	15.2 (0.1)a	-25.6 (0.8)a	-2.8 (0.2)a
B / +C ₂ H ₂	546.9 (27.5)b	1235.7 (83.4)a	14.9 (0.2)a	-26.3 (<0.1)a	-3.5 (0.4)a
C / +C ₂ H ₂	519.8 (19.2)b	1173.5 (25.7)a	16.2 (<0.1)*	-25.2 (0.1)*	-4.0 (0.4)*
D / +C ₂ H ₂	511.7 (3.5)b	1295.6 (63.3)a	16.0 (0.1)ba	-25.1 (0.1)a	-4.3 (0.5)a
Experiment 4 (Loamy sand, summer 2011)					
A / -C ₂ H ₂	175.3 (6.6)a	2448.5 (135.8)a	25.7 (0.3)a	-30.6 (0.2)a	12.1 (1.6)a
B / -C ₂ H ₂	121.3 (74.0)a	2091.3 (19.5)b	28.0 (5.0)a	-32.3 (0.7)a	7.7 (1.4)b

C/-C₂H₂	104.5 (5.3)a	1844.7 (192.1)b	29.3 (0.1)a	-30.0 (0.5)a	4.3 (1.0)e
D/-C₂H₂	73.8 (63.0)a	1632.2 (115.3)b	28.9 (1.2)a	-31.8 (2.2)a	3.4 (2.0)e
A/+C₂H₂	263.5 (31.7)a	2076.6 (305.3)a	13.5 (0.5)*	-34.7 (0.1)*	-1.0**
B/+C₂H₂	233.0 (15.6)a	1794.9 (238.9)a	14.3 (1.7)a	-33.8 (0.9)a	-4.9 (0.9)a
C/+C₂H₂	119.5 (102.7)a	1736.8 (424.7)a	19.0 (7.0)a	-33.1 (2.8)a	-1.7 (2.7)b
D/+C₂H₂	161.6 (7.6)a	1497.0 (138.7)a	14.8 (0.5)a	-35.7 (0.2)a	-4.9 (0.7)e

Letters denote significant differences ($P < 0.105$) among treatments and varieties within a soil.

Asterisks indicate that only two samples (*) or one sample (***) of triplicates were ~~analyzable~~ analysable due to logistical difficulties.

3.2 Isotopologues of N₂O produced in different varieties and treatments

3.2.1 Variety -C₂H₂

SP_{N₂O} values of all Soils and inhibitor treatments of variety -C₂H₂ were within a range of -1.8 to 12.1 ‰ (Table 2) and differed among inhibitor treatments ($P = 0.037$). SP_{N₂O} values in variety -C₂H₂ of Soil 1.2 was particularly large (3.4 to 12.1 ‰) compared to the other Soils (1.6 to -1.6 ‰). SP_{N₂O} values of variety -C₂H₂ were significantly larger than SP_{N₂O} values of variety +C₂H₂ ($P < 0.001$) (up to 2.4, 1.5, 4.6 and 4.1 ‰ in Soil 1.1, 2, 3 and 4, respectively). Generally, most SP_{prod} values of variety -C₂H₂ (Eq. 7) were smaller than SP_{N₂O} values of variety -C₂H₂ but still larger than SP_{N₂O} values of variety +C₂H₂ and are presented in Table S2 (Supplementary Material).

3.2.1.2 Variety +C₂H₂

SP_{N₂O} values of all ~~experiment~~ Soils, and all treatments of variety +C₂H₂ were within a narrow range between -4.9 and -0.4 ‰ (Table 2), ~~and differed only significantly among treatments of Experiment 4 ($P = 0.002$).~~ In general, there were only small differences among treatments: SP_{N₂O} values of treatments A in variety +C₂H₂ differed significantly among soils ($P < 0.001$), with largest SP_{N₂O} values in ~~Experiment 1~~ Soil 1.1 (-0.4 ‰) and smallest SP_{N₂O} values in ~~Experiment~~ Soil 3 (-2.8 ‰). SP_{N₂O} values of treatment D in variety +C₂H₂ of all soils varied between -1.5 and -4.9 ‰, but only SP_{N₂O} values of ~~Experiment~~ Soil 2 differed significantly from SP_{N₂O} values of the other ~~Experiment~~ Soils ($P = 0.006$). For treatments B of variety +C₂H₂, SP_{N₂O} values differed only significantly between ~~Experiment 1~~ Soil 1.1 and 4.1.2, 2 and 4.1.2, and 1.1 and 3 (each $P = 0.002$). SP_{N₂O} values from treatment C in variety +C₂H₂ did not differ significantly ($P = 0.600$). For every soil, we found significantly larger $\delta^{18}\text{O}_{\text{N}_2\text{O}}$, $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$ and SP_{N₂O} values in variety -C₂H₂ than in variety +C₂H₂ ($P < 0.001$), except for ~~Experiment~~ Soil 2, where $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$ values of variety -C₂H₂ were indistinguishable from those of variety +C₂H₂ ($P = 0.400$). However, only in a few varieties there were significant differences in $\delta^{18}\text{O}_{\text{N}_2\text{O}}$, $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$ or SP_{N₂O} values between treatments with fungal and bacterial inhibition (B and C, respectively) (Table 2). As explained in section 3.3, N₂O reduction blockage in varieties +C₂H₂ was successful in most cases (~~Experiment~~ Soil 2, 3 and 4). SP_{N₂O} values of this variety are thus assumed to be valid estimates of $\delta\theta$, i.e. SP_{prod} values of N₂O production, and can thus be used for applying the IEM.

~~3.2.2 Variety -C₂H₂~~

655 SP_{N_2O} values of all experiments and inhibitor treatments of variety $-C_2H_2$ were within a range of 1.8 to 12.1‰ (Table 2) and did not differ among inhibitor treatments ($P=0.037$). SP_{N_2O} values in variety $-C_2H_2$ of Experiment 4 was particularly large (3.4–12.1‰) compared to the other experiments (1.6 to 1.6‰). As already stated above, SP_{N_2O} values of variety $-C_2H_2$ were significantly larger than SP_{N_2O} values of variety $+C_2H_2$ (up to 2.4, 1.5, 4.6 and 4.1‰ in Experiment 1, 2, 3 and 4, respectively). Generally, most SP_{prod} values of variety $-C_2H_2$ (Eq. 7) were smaller than SP_{N_2O} values of variety $-C_2H_2$ but still larger than SP_{N_2O} values of variety $+C_2H_2$ and are presented in Table S1 (supplementary Material).

3.2.3 Variety traced

665 The ^{15}N -labeling of N_2O ($^{15}N_{N_2O}$) or N_2 produced ($^{15}N_{N_2}$) gave information about the incorporated N from ^{15}N -labeled NO_3^- into N_2O or N_2 as well as about the N_2O reduction to N_2 . Microorganisms in each treatment used the ^{15}N -labeled NO_3^- in variety traced (Table 3) and expected $^{15}N_{N_2O}$ depended on the initial N abundance in NO_3^- of unfertilized soil (Eq. 7). Experiment 4 Soil 1.2 is the only one showing a large discrepancy between measured (about 30 at%) and calculated $^{15}N_{N_2O_exp}$ (49 at%) in N_2O , whereas the other experiment Soils showed close agreement (Table 3).

3.3 Product ratios of denitrification and efficiency of N_2O reductase blockage by C_2H_2

670 $r_{C_2H_2}$ as well as product ratio $r_{^{15}N}$ of determined with Experiment Soil 2 were significantly larger than of with the other experiment Soils ($P \leq 0.001$) (Table 3). $r_{^{15}N}$ of treatment B was significantly larger than of treatment C and D of Experiment 4 Soil 1.2 ($P = 0.032$), while all other treatments of other Soils soils did not differ. $r_{C_2H_2}$ did not differ significantly among treatments ($P = 0.400$). In order to test the efficiency of blockage of N_2O reduction by C_2H_2 application, $r_{C_2H_2}$ (Eq. 5) was compared with $r_{^{15}N}$ (Eq. 6). In Experiment 1 Soil 1.1, $r_{C_2H_2}$ was by far smaller than $r_{^{15}N}$, while both calculated product ratios were in similar ranges in the other three experiment Soils and thus a successful blockage of N_2O reduction was assumed for those experiment Soils.

680 Table 3: Average CO_2 and N_2O production rates of the last sample collection after 10 or 8 hours of variety traced, respectively, with ^{15}N labeling in N_2O ($^{15}N_{N_2O}$) and the calculated $r_{^{15}N}$ of variety traced and $r_{C_2H_2}$ calculated from N_2O production rates of variety $-C_2H_2$ and $+C_2H_2$ of each soil (Experiment 1 Soil 1 to 43) for treatments A without, B with bacterial, C with fungal, and D with bacterial and fungal growth inhibition, respectively (standard deviation in brackets, $n = 3$).

Treatment	mean N_2O [$\mu g N kg^{-1} h^{-1}$]	mean CO_2 [$\mu g N kg^{-1} h^{-1}$]*	$^{15}N_{N_2O}$ [at %]	$^{15}N_{N_2O_exp}$ [at %] ^a	Calc. total $r_{^{15}N}$ ^{b*}	Calc. total $r_{C_2H_2}$ ^{c*}
Experiment 1 Soil 1.1 (Loamy Sand, 2012)						
A	255.6 (43.5)	1310.0 (167.3)	36.8 (0.1)	39	0.80 (0.02)	0.48 (0.07)
B	154.5 (29.6)	1153.5 (238.4)	36.4 (0.2)		0.76 (0.02)	0.48 (0.05)
C	191.6 (30.7)	1219.6 (109.1)	36.9 (<0.1)		0.72 (0.05)	0.45 (0.04)
D	148.1 (1.9)	1253.8 (54.5)	36.8 (0.1)		0.69 (0.02)	0.54 (0.05)
Soil 1.2 (Loamy Sand, 2011)						
A	156.9 (62.7)	3111.4 (1252.5)	31.1**	49	0.54 (0.05)	0.63 (0.10)
B	169.2 (6.1)	2314.6 (307.1)	26.5**		0.59 (0.03)	0.63 (0.17)
C	117.2 (3.1)	1785.6 (79.3)	30.1 (1.1)*		0.50 (0.01)	0.62 (0.02)
D	115.2 (3.1)	1706.7 (38.1)	33.5 (0.5)*		0.50 (0.01)	0.53 (0.12)
Experiment Soil 2 (Sand, 2012)						
A	240.7 (0.95)	1286.2 (5.6)	43.2 (<0.1)	44	0.94 (0.01)	1.04 (0.10)

B	<u>185.1 (3.9)</u>	<u>1157.4 (17.3)</u>	43.0 (0.1)		0.94 (0.01)	0.81 (0.04)
C	<u>241.1 (13.4)</u>	<u>1282.1 (63.4)</u>	43.2 (0.1)		0.95 (0.01)	0.99 (0.09)
D	<u>167.3 (34.9)</u>	<u>1199.0 (34.6)</u>	42.7 (0.1)		0.93 (0.01)	0.98 (0.04)
Experiment Soil 3 (Silt loam, 2013)						
A	<u>285.9 (20.4)</u>	<u>1044.0 (46.6)</u>	35.8 (<0.1)		0.62 (<0.01)	0.52 (0.04)
B	<u>320.5 (14.7)</u>	<u>1204.2 (86.5)</u>	35.5 (<0.1)	34	0.62 (0.01)	0.59 (0.02)
C	<u>216.4 (34.9)</u>	<u>980.5 (202.5)</u>	35.5 (<0.1)		0.59 (0.02)	0.48 (0.04)
D	<u>231.4 (11.4)</u>	<u>988.5 (74.4)</u>	35.3 (<0.1)		0.62 (0.01)	0.51 (0.04)
Experiment 4 (Loamy Sand, 2011)						
A	⊖	⊖	<u>31.1**</u>		<u>0.54 (0.05)</u>	<u>0.63 (0.10)</u>
B	⊖	⊖	<u>26.5**</u>	49	<u>0.59 (0.03)</u>	<u>0.63 (0.17)</u>
C	⊖	⊖	<u>30.1*</u>		<u>0.50 (0.01)</u>	<u>0.62 (0.02)</u>
D	⊖	⊖	<u>33.5*</u>		<u>0.50 (0.01)</u>	<u>0.53 (0.12)</u>

Asterisks indicate that only two samples (*) or one sample (**) were ~~analyzed~~ analysed, due to logistical difficulties.

^a $^{15}N_{N_2O_{exp}}$ [at %] was calculated from Eq. 78.

^b $r_{L_{15N}}$ = [N₂O/(N₂+N₂O)] with N₂O or N₂ production rates from variety traced; see Eq. 5

^c $r_{C_2H_2}$ = [N₂O-C₂H₂/N₂O+C₂H₂] with N₂O production rate from varieties -C₂H₂ and +C₂H₂; see Eq. 6, cf. Table 2

685

3.4 Fungal contribution to N₂O production from denitrification by microbial inhibitor approach (modified SIRIN)

When calculating f_{FDmi} , N₂O production rates of treatment D must be significantly smaller compared to the other three treatments and the flux balance according to Eq. 1 and 2 must be consistent. Taking the large ranges of N₂O production rates of each treatment (minimum and maximum values) into account, for each Soil (A-D) was indistinguishable from ((B-D)+(C-D)) (Eq. 2), showing good agreement between Eqs. 1 and 2. However, N₂O production in treatment D was large within all varieties. Only with -Soil 2 of the variety +C₂H₂, the N₂O production rates of treatment D were significantly smaller than those of the other three treatments. Thus, for Soil 2, f_{FDmi} could be calculated (Eq. 3) and amounted to 28 ± 9 % (Table 5) with a corresponding This was only the case in Experiment 2 of variety +C₂H₂. The calculated F_{FDmi} (Eq. 3) was 0.28 ± 0.90 (Table 5). The respective flux fungal N₂O production rate of fungal N₂O was $0.2423.7 \pm 0.081.8$ $\mu\text{g N kg}^{-1} \text{ h}^{-1}$. Although the N₂O production rate of Treatment D was smaller than that of treatment A (Soil 2), it must be pointed out, that due to the large amount of non-inhibitable production (treatment D), even the result for Soil 2 is actually very unsure. For all other experimentSoils, calculation of f_{FDmi} was not possible, i.e. SIRIN was not successful.

690

695

3.5 Fungal contribution to N₂O production from denitrification by the SP endmember mixing approach (IEM) and SP/ $\delta^{18}\text{O}$ isotope mapping approach (SP/ $\delta^{18}\text{O}$ Map)

The IEM revealed that $f_{FD_{SP}}$ was small in all Soils (≤ 15 %, ≤ 14 %, ≤ 9 %, and ≤ 11 % with Soil 1 to 3, respectively) (Table 5). Regardless of influence of N₂O reduction on SP_{N_2O} values, only in Soil 1.2 $f_{FD_{SP_{pot}}}$ could have reached 66%, while fungal denitrification could not have dominated with the other three soils (Table 5).

705

When applying SP/ $\delta^{18}\text{O}$ Map, we can assess the plausibility of the determined f_{FD} values based on the $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values obtained from the fitting ($\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value in Table 4) and the fitting outcome, i.e. the difference between $r_{L_{15N}}$ and r_{MAP} (Diff-in, see Table 4). The most probable $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value for our experimentSoils can be assumed based on the fact that Braunschweig tap water was added to soil used and the original soil water also represents the isotope characteristics typical for this region, which is about -7.4 ‰ (long-term mean Braunschweig precipitation water

(Stumpp et al., 2014)). Thus, in the presented application of SP/ $\delta^{18}\text{O}$ Map, $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values were fitted and it has to be pointed out that the precision of such calculations can be improved by measuring $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ instead. Depending on the season and evaporative losses, $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ this value may slightly vary and the most possible range of soil water in our ~~experiment~~Soils may vary from about -11 to -4‰ as observed in other experiments ~~conducted~~ used in our laboratory ~~experiments with~~ similar conditions (Lewicka-Szczebak et al., 2014; Rohe et al., 2014a; Lewicka-Szczebak et al., 2017; Rohe et al., 2017). Taking this into account, we can say that for ~~Experiment 1~~Soil 1.1, the fungal contribution must be below ~~0.022~~‰, because to obtain any larger f_{FD} values, unrealistically small $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -14.9‰) must be fitted (see Table 4). For ~~Experiment~~Soil 2, both the smaller $f_{\text{FD_MAP}}$ values of ~~0.01~~‰ and the larger ones up to ~~0.15~~‰ are possible, since they are associated with very realistic $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -6.3 and -10.1, respectively) and identical *Diff* of 0.04 (Table 4). For ~~Experiment~~Soil 3, the only plausible fitting can be obtained for the smallest SP_{BD} values, which are associated with a $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value of -5.6‰ (Table 4). Although the *Diff* for this fitting is slightly higher, the other fittings must be rejected due to unrealistic $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -1.7 and +3.7‰), hence $f_{\text{FD_MAP}}$ values must be ~~between 0.04- and 0.09~~‰. Similarly, for ~~Experiment 4~~Soil 1.2, the only plausible fitting can be obtained for the smallest SP_{BD} values, which are associated with a $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value of -6.8‰ (Table 4) and indicate $f_{\text{FD_MAP}}$ values from ~~0.11 to 0.20~~‰. Here this fitting also shows clearly the smallest *Diff* of only 0.01 (Table 4). However, except for ~~Experiment 4~~Soil 1.2, where the *Diff* is smallest for the last fitting, the *Diff* values for other ~~experiment~~Soils are very similar for different fittings with the largest values in ~~Experiment~~Soil 3. A better fit (showing smaller *Diff* values) was not possible with any other combination of SP_{BD} and $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values. Since the precision of $r_{15\text{N}}$ (expressed in standard deviation in Table 3) was always ≤ 0.05 , this uncertainty of $r_{15\text{N}}$ did not reduce the precision of the fitting (compare large ranges of $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ and r_{MAP} values, respectively, in Table 4). The $f_{\text{FD_SP}}$ ranged between 0 and approximately ~~0.15~~‰ (Table 5). The results obtained from SP/ $\delta^{18}\text{O}$ Map show $f_{\text{FD_MAP}}$ reaching up to ~~0.14, 0.15, and 0.09 and 0.20~~‰ for ~~Experiment~~Soils 1.1, 1.2, 2, and 3, and 4, respectively (Figure 3, Table 4, Table 5). **Importantly, due to the fitting procedure applied the estimations of $f_{\text{FD_MAP}}$ values are based not only on $SP_{\text{N}_2\text{O}}$ and $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values but also on the results obtained in the ^{15}N treatment ($r_{15\text{N}}$ values).**

735

740

Table 4: Summary of the results provided by SP/ $\delta^{18}\text{O}$ Map for fraction of fungal denitrification ($f_{\text{FD_MAP}}$) and N_2O product ratio (r_{MAP}) in the acetylated (+ C_2H_2) and non-acetylated (- C_2H_2) ~~treatments varieties~~ for ~~3~~three possible $SP_{\text{N}_2\text{O}}$ values from bacterial denitrification (SP_{BD}): mean (-1.9‰), maximal (3.7‰), and minimal (-7.5‰). The $\delta^{18}\text{O}$ values of soil water ($\delta^{18}\text{O}_{\text{H}_2\text{O}}$) were fitted to get the lowest difference (*Diff*) between product ratio determined with ^{15}N treatment ($r_{15\text{N}}$) and SP/ $\delta^{18}\text{O}$ Map (~~$r_{15\text{N}}$ and~~ r_{MAP}). The most plausible fittings are ~~shown in bolded~~ (see discussion for reasons of this choice).

745

Experiment Soil	Variety	$r_{15\text{N}}$	SP_{BD} [‰]	$\delta^{18}\text{O}_{\text{H}_2\text{O}}$ [‰]	r_{MAP}	<i>Diff</i>	$f_{\text{FD_MAP}}$ [%]*
<u>1.1</u>	- C_2H_2	0.66	-1.9	-11.2	0.66	0.00	<u>-0.01-1</u>
	+ C_2H_2	1	-1.9	-11.2	1.00	0.00	<u>0.02</u>
	- C_2H_2	0.66	3.7	-6.1	0.65	0.01	<u>-0.14-14</u>
	+ C_2H_2	1	3.7	-6.1	1.00	0.00	<u>-0.16-16</u>

	$-C_2H_2$	0.66	-7.5	-14.9	0.66	0.00	0.08
	$+C_2H_2$	1	-7.5	-14.9	1.00	0.00	0.14
<u>1.2</u>	<u>$-C_2H_2$</u>	<u>0.60</u>	<u>-1.9</u>	<u>-3.3</u>	<u>0.66</u>	<u>0.06</u>	<u>15</u>
	<u>$+C_2H_2$</u>	<u>1</u>	<u>-1.9</u>	<u>-3.3</u>	<u>0.96</u>	<u>0.04</u>	<u>-30</u>
	$-C_2H_2$	0.60	3.7	1.5	0.72	0.12	8
	$+C_2H_2$	1	3.7	1.5	0.91	0.09	-21
	$-C_2H_2$	0.60	-7.5	-6.8	0.61	0.01	20
	$+C_2H_2$	1	-7.5	-6.8	0.99	0.01	11
2	$-C_2H_2$	0.94	-1.9	-6.3	0.90	0.04	0.01
	$+C_2H_2$	1	-1.9	-6.3	1.04	0.04	0.01
	$-C_2H_2$	0.94	3.7	-1.2	0.90	0.04	-0.16-16
	$+C_2H_2$	1	3.7	-1.2	1.04	0.04	-0.18-18
	$-C_2H_2$	0.94	-7.5	-10.1	0.90	0.04	0.13
	$+C_2H_2$	1	-7.5	-10.1	1.04	0.04	0.15
3	$-C_2H_2$	0.61	-1.9	-1.7	0.54	0.07	-0.03-3
	$+C_2H_2$	1	-1.9	-1.7	1.04	0.04	-0.05-5
	$-C_2H_2$	0.61	3.7	3.7	0.54	0.07	-0.14-14
	$+C_2H_2$	1	3.7	3.7	1.03	0.03	-0.24-24
	$-C_2H_2$	0.61	-7.5	-5.6	0.53	0.08	0.04
	$+C_2H_2$	1	-7.5	-5.6	1.04	0.04	0.09
<u>4</u>	<u>$-C_2H_2$</u>	<u>0.60</u>	<u>-1.9</u>	<u>-3.3</u>	<u>0.66</u>	<u>0.06</u>	<u>0.15</u>
	<u>$+C_2H_2$</u>	<u>1</u>	<u>-1.9</u>	<u>-3.3</u>	<u>0.96</u>	<u>0.04</u>	<u>-0.03</u>
	$-C_2H_2$	0.60	3.7	1.5	0.72	0.12	0.08
	$+C_2H_2$	1	3.7	1.5	0.91	0.09	-0.21
	$-C_2H_2$	0.60	-7.5	-6.8	0.61	0.01	0.20
	$+C_2H_2$	1	-7.5	-6.8	0.99	0.01	0.11

*Negative values for $f_{FD\ MAP}$ are non-realistic and therefore discarded for further interpretation.

750

755

760

765

770

775

780

Table 5: Ranges of the fraction of N₂O produced by fungi (f_{FD}) from four soil experiments/Soils using four different approaches: Fungal fraction was calculated using a) the microbial inhibitor approach (modified SIRIN) (f_{FDmi}), b) the isotopomer endmember mixing approach (IEM) by SP isotope mixing balance using b) variety +C₂H₂ ($f_{FD SP}$), c) $f_{FD SPpot}$ the IEM for results from variety -C₂H₂ assuming the SP effect of N₂O reduction was negligible, -d), the IEM by SPN₂O isotope mixing balance (IEM) for results from variety -C₂H₂ with reduction correction to calculate the SPN₂O values ($f_{FD SPcalc}$), and e) the $\delta^{18}O/SP$ Map ($f_{FD MAP}$) with $\delta^{18}O_{N_2O}$ and SPN₂O values from variety -C₂H₂ and variety +C₂H₂. Negative values by IEM and $\delta^{18}O/SP$ Map are assumed to be zero.

Experiment/Soil	f_{FDmi} [%] ^a	$f_{FD SP}$ [%] ^{b*}	$f_{FD SPpot}$ [%] ^{c*}	$f_{FD SPcalc}$ [%] ^{d*}	$f_{FD MAP}$ [%] ^{e*}
1.1	n.d.	0-14 to 0-15	-12 to 39	-60-0 to 19	0-0.02
1.2	n.d.	-23 to 11	10 to 66	1 to 21	11 to 20
2	0-19 to 0-37	0-18 to 0-14	-14 to 36	-120 to 0-15	0-01 to 0-15
3	n.d.	0-25 to 0-09	-11 to 40	-90 to 0-18	0-04 to 0-09
4	n.d.	0-0-11		0-0-21	0-11-0-20

^aFungal fraction of N₂O production calculated by Eq. 3 taking variations of three replicates into account.

^bFungal fraction of N₂O production calculated by Eq. 4 for variety +C₂H₂ with assuming SPN₂O values of N₂O produced by bacteria were 3.7 ‰ (resulting in negative fraction and therefore set to zero) or -7.5 ‰ (Yu et al., 2020) and by fungi on average 33.6 ‰ (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017). Using the minimum and maximum SPN₂O values known for bacteria resulted in a $f_{FD SP}$ range.

^cMaximum potential fungal fraction of N₂O production calculated by Eq. 4 as an average range for all treatments of variety -C₂H₂ assuming SPN₂O values of N₂O produced by bacterial denitrification or nitrifier denitrification were between 3.7 and -10.7 ‰ (Frame and Casciotti, 2010; Yu et al., 2020) or produced by fungal denitrification or nitrification were between 16 and 37 ‰ (Sutka et al., 2008; Decock and Six, 2013; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017). Using the minimum and maximum SPN₂O values known from pure cultures resulted in the given $f_{FD SPpot}$ range. Here, the effect of partial reduction of N₂O could not be included on SPN₂O values was assumed to be negligible. -measured of the four replicates the given

^dEq. 4 to solve for fungal fraction in variety -C₂H₂ with assuming SPN₂O values of N₂O produced by bacteria was 3.7 (resulting in negative fraction and therefore set to zero) or -7.5 ‰ and using reduction correction with $\eta_r = -6$ ‰ to calculate SP_{prod} values (Senbayram et al., 2018; Yu et al., 2020). Using the minimum and maximum SPN₂O values known for bacteria resulted in a $f_{FD SPcalc}$ range.

^eFungal fraction of N₂O production calculated by SP/ $\delta^{18}O$ Map with assuming most probable SPN₂O values from bacterial denitrification (according to Table 4). -Using the minimum and maximum SPN₂O values known for bacteria and ranges of fitted $\delta^{18}O_{N_2O}$ values (the fitting is based also on results obtained in ¹⁵N treatment) values resulted in a $f_{FD MAP}$ range.

*Negative values for $f_{FD SP}$, $f_{FD SPpot}$, $f_{FD SPcalc}$, $f_{FD MAP}$ are non-realistic and therefore discarded for further interpretation.

n.d.-not determined because of insufficient inhibition.

785

790

795

800

805




Figure 23: SP/ $\delta^{18}\text{O}$ isotope mapping approach (SP/ $\delta^{18}\text{O}$ Map) to estimate the contribution of bacteria or fungi to N_2O produced according to Lewicka-Szczebak et al. (2017) and Lewicka-Szczebak et al. (2020). The isotopic values for natural abundance treatments with acetylene addition (+ C_2H_2 , empty symbols) and without acetylene addition (- C_2H_2 , corresponding filled symbols) are shown for four experiment Soils (1 to 3-4). The grey rectangles indicate expected ranges of isotopic signatures for heterotrophic bacterial denitrification (BD) and fungal denitrification (FD) (Yu et al., 2020). The black solid line is the mixing line connecting the average expected values for BD and FD, while the red solid line is the mean reduction (for the mean SP values for BD) line and the red dashed line is the minimum reduction line (for the minimal $\text{SP}_{\text{N}_2\text{O}}$ values for BD).

3.6 $\text{SP}_{\text{N}_2\text{O}}$ values of N_2O produced by the fungal soil community

Solving Eq. 4 for SP_{FD} enables to calculate $\text{SP}_{\text{N}_2\text{O}}$ values from the fungal soil community for Experiment 2 (Table 6). Estimates for the ranges of F_{FD} and F_{BD} from the results (+ C_2H_2) of the modified SIRIN were obtained ($F_{\text{FDmi}}=0.19-0.37$ and $F_{\text{BD}}=1-F_{\text{FDmi}}$ resulted in a range between 0.63 and 0.81, respectively, see section “3.4 Fungal contribution to N_2O production from denitrification by microbial inhibitor approach (modified SIRIN)”). The SP_{prod} values of N_2O ($\text{SP}_{\text{prod}}=-1.4$ ‰) of the respective treatment A (Table 2, variety + C_2H_2) served to calculate $\text{SP}_{\text{N}_2\text{O}}$ values for fungal denitrification for Experiment 2. Assuming -7.5 or -3.7 ‰ for the bacterial $\text{SP}_{\text{N}_2\text{O}}$ endmember values of N_2O (Toyoda et al., 2005; Sutka et al., 2006; Yu et al., 2020) resulted in SP_{FD} values between -10 ‰ ($\text{SP}_{\text{BD}}=3.7$ ‰) and 25 ‰ ($\text{SP}_{\text{BD}}=-7.5$ ‰) (Table 6). The respective SP_{FD} value for variety - C_2H_2 was in a very similar range between -17 ‰ and 27 ‰ (Table 6) using SP_{prod} values ($\text{SP}_{\text{prod}}=-1.0$ ‰) of the respective treatment A (Table S1). (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015)

Table 6: SP_{FD} values (i.e. $\text{SP}_{\text{N}_2\text{O}}$ values of N_2O produced by fungi) by solving Eq. 4 using F_{FDmi} and F_{BD} from results of modified SIRIN approach and using SP_{prod} values of varieties + C_2H_2 and - C_2H_2 of Experiment 2.

Treatment	SP_{prod} [‰]	SP_{BD} [‰] ^a	F_{FDmi} ^b	F_{BD} ^b	SP_{FD} [‰]
+ C_2H_2	-1.4	-7.5	0.19	0.81	25
		3.7	0.19	0.81	-23
		-7.5	0.37	0.63	9
		3.7	0.37	0.63	-10

		-7.5	0.19	0.81	27
		3.7	0.19	0.81	-17
$-C_2H_2$	-1.0	-7.5	0.37	0.63	10
		3.7	0.37	0.63	-9

SP_{N_2O} endmember values of bacterial denitrification were taken for calculation (Eq. 4) according to studies with pure cultures (Toyoda et al., 2005; Sutka et al., 2006; Yu et al., 2020).

^bRanges of F_{FDmi} and F_{BD} were calculated using the modified SIRIN approach.

4. Discussion

To our knowledge, this was the first attempt to determine SP_{N_2O} values by fungi or bacteria from soil communities using microbial growth inhibitors with a modification of SIRIN and comparing microbial inhibitor and isotopic approaches (IEM and SP/ $\delta^{18}O$ Map) to estimate fungal contribution to N_2O production from denitrification in anoxic incubation. Using The IEM isotopic approaches revealed that the fungal contribution to N_2O production was small ($f_{FD_SP} \leq 0.15\%$ or $f_{FD_MAP} \leq 0.20\%$) in the three soils tested (Table 5). A dominant contribution of fungi over bacteria was also excluded by the potential maximum fungal denitrification for Soil 1, 2, and 3 (f_{FD_SPpot} between 37 and 40 %, Table 5), even though effects of N_2O reduction are not included. The modified SIRIN approach was not successful, because large amounts of non-inhibitable N_2O production were observed with all four Soils (Table 2, Table 3). The fungal fraction producing N_2O during denitrification (f_{FDmi}) was only one experiment estimated for Soil 2, with where significantly smaller N_2O production in treatment D was observed compared to that of treatment A and resulted modified SIRIN allowed the calculation of the fungal fraction producing N_2O during denitrification (F_{FDmi} between in a range of -0.19 and to 0.37 % in Experiment 2), which was larger was probably overestimated due to uncertainties resulting from the large N_2O production of non-inhibitable sources, than the by two isotope approaches (≤ 0.20). While the three approaches coincided in showing dominance of bacterial denitrification, the isotopic approaches yielded similar small estimates of for f_{FD} ($\leq 20\%$) and thus did not confirm largest f_{FDmi} of Experiment Soil 2. The strict application of the SIRIN method prescribes proof of selectivity of the inhibitors (i.e., streptomycin should not inhibit fungi and cycloheximide should not inhibit bacteria). The All SIRIN results obtained with respect to N_2O production by the fungal or bacterial fraction were rather unsatisfactory, thus fungal SP_{N_2O} values could not be assessed, and the overall results led to unsolved questions, which are discussed in the following sections.

4.1 Experimental setup and inhibitor effects

Inhibitor effects, expressed by smaller N_2O production with selective inhibitors (treatments B, C and D) compared to treatments without inhibitors (A), were only minor in the present study. In accordance with other studies, N_2O production was analysed after the addition of glucose as substrate (Laughlin and Stevens, 2002; McLain and Martens, 2006; Blagodatskaya et al., 2010; Long et al., 2013). Glucose initiates the growth of active heterotrophic organisms. Since pure cultures were shown to synthesize enzymes capable of denitrification within two to three hours (USEPA, 1993), a pre-incubation of soil under anaerobic conditions is not needed. Thus, when gas sample collection started organisms should have produced denitrifying enzymes and microbial growth of initially active organisms should have started too. However, in accordance to Anderson and Domsch (1975) experimental duration should be as short as possible to ensure the CO_2 production by initially active organisms only. Thus, short-time incubation is recommended when conducting a modified SIRIN approach, as the incubation period should cause

changes in conditions for microorganisms and initiate growth on the one hand, while it should avoid the consumption of inhibitors as C sources on the other.

With incubation time, production rates of CO₂ decreased, probably because experimental incubation conditions provoked unfavourable conditions and physiological changes, e. g. due to anaerobic conditions or local substrate depletion (e. g. C supplied as glucose). Decreasing CO₂ fluxes might also be explained by CO₂ accumulation in pore space as this effect is shown by modelled diffusive fluxes from soil in closed systems (Well et al., 2019).

Previous studies found much larger inhibitor effects by pre-incubating the soil with selective inhibitors (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013; Chen et al., 2014). It is therefore important to discuss considerable differences among the experimental design of the present study compared to that of other studies (e. g., Laughlin and Stevens, 2002; Blagodatskaya et al., 2010).

The conventional practice of SIRIN implies determination of $c_{opt}(\text{glucose})$, $c_{opt}(\text{streptomycin})$ or $c_{opt}(\text{cycloheximide})$ with an "Ultragas 3" CO₂ analyser (WösthoffCo., Bochum) (Anderson and Domsch, 1973) with continuous gas flow. We used this method to determine optimal concentrations for SIRIN in the pre-experiment and used these concentrations for the modified SIRIN approach as well. This optimization procedure was not used in other studies (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013). We supposed that optimal concentrations for CO₂ respiration should work as well for denitrification, if both inhibitors inhibit the denitrification process as well. However, although SIRIN has so far been tested with isolated cultures and soils for microbial growth for CO₂ production only (Anderson and Domsch, 1973, 1975), information on N₂O producing processes, especially denitrification, is still lacking and should be investigated in further studies. Additionally, as presented by Ladan and Jacinthe (2016) the bactericide bronopol and the fungicide captan were more effective inhibitors than streptomycin or cycloheximide and should be included when evaluating inhibition approaches and isotopic endmember approaches.

Previous studies that found much larger inhibitor effects were conducted after pre-incubating the soil with selective inhibitors (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013; Chen et al., 2014).-The In contrast to that, the experimental design of our incubation setup was without soil pre-incubation with selective inhibitors to minimize disturbance of the soil microbial community and our approach was thus -however, but this was- in agreement with the original SIRIN method for respiration (Anderson and Domsch, 1973, 1975, 1978) without soil pre incubation with selective inhibitors to minimize disturbance of the soil microbial community.

Another study performing similar experiments without pre-incubation with inhibitors did not find effectiveness of application of both antibiotics during long-term application (up to 48 h) (Ladan and Jacinthe, 2016)-, although streptomycin and cycloheximide are commonly used to inhibit denitrification of selective groups. Nevertheless, as we expected that pre-incubation with selective inhibitors would induce changes in the F:B ratio of soil, we decided to conduct the modified SIRIN approach without a pre-incubation step. This assumption was supported by findings of Blagodatskaya et al. (2010), where pre-incubation of about one to twenty hours with cycloheximide resulted in increasing inhibitor efficiency with time, while this was not the case when pre-incubating with streptomycin. Consequently This suggests that -microbial communities- might change after exposition to cycloheximide.

In the present study, even with both growth inhibitors (treatment D), N₂O production was large in all experiments, i.e. in most cases not significantly smaller than in the other three treatments A, B or C. Thus, we suppose similar contributions of non-inhibitable organisms and processes in all treatments. Non-inhibitable organisms could be, for example, bacteria or fungi that are not in growth stage or may be not affected by inhibitors. Recently, Pan et

915 al. (2019) summarized findings of other studies and pointed out that some microorganisms can use inhibitors as growth substrates, that dead organisms may serve as energy sources for others, and that interactions of microbial species may change due to non-inhabitable organisms occurring in soil communities. Non-inhabitable organisms could be archaea as well, which are also known to be capable of denitrification (Philippot et al., 2007; Hayatsu et al., 2008). It is known that archaea are not affected by streptomycin or cycloheximide (Seo and DeLaune, 2010). However, effects of archaeal occurrence in soil or secondary effects on fungi or bacteria were not tested in this study. Additionally, abiotic N₂O production cannot be quantified with the experimental setup, but might be contributing to each inhibitor treatment.

920 In summary, the present experimental setup without pre-incubating soil samples with selective inhibitors was not successful in complete inhibition of bacterial or fungal denitrifiers. Although pre-incubation with selective inhibitors may lead to more successful inhibition, we do not recommend this due to induced changes in soil communities. For further studies focusing on application of modified SIRIN to determine the fraction of bacterial or fungal N₂O derived from denitrification a method validation using also different inhibitors is recommended.

925 Inhibitor application without pre incubating with inhibitors was contrary to previous studies focusing on N₂O production (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013) and we suppose expected that pre incubation with selective inhibitors would induce that pre incubation with selective inhibitors changes the F:B ratio compared to the undisturbed soil considerably more than soil incubation without this pre incubation step. Additionally, although Blagodatskaya et al. (2010), did not find more inhibitor efficiency after a period of 1 to 20 hours of pre incubation with streptomycin, they found greater inhibitor effects of cycloheximide with pre incubation phases. This could indicate that the microbial distribution changed after exposition to this inhibitor. Anderson and Domsch (1975) stated already that CO₂ production of initially active organisms can only be ensured up to six or eight hours of experimental duration and biomass activity is changed by both inhibitors. conditions

930 It has to be noticed that pre incubation in previous studies was without glucose, while N₂O production was analyzed after the addition of glucose as substrate in the present as well as previous studies (Laughlin and Stevens, 2002; McLain and Martens, 2006; Blagodatskaya et al., 2010; Long et al., 2013). Glucose initiates the growth of active heterotrophic organisms. Pre incubation under denitrifying conditions is not needed for microorganisms to produce denitrifying enzymes as pure cultures synthesized enzymes capable of denitrification within two to three hours (USEPA, 1993). We started gas sample collection after two or four hours, when organisms should have produced denitrifying enzymes and microbial growth of initially active organisms should have started. With incubation time, production rates of CO₂ decreased, probably because experimental incubation conditions provoked unfavorable conditions and physiological changes, e.g. due to increasing partial pressure within the closed jars.

935 The conventional practice of SIRIN implies determination of $c_{opt}(\text{glucose})$, $c_{opt}(\text{streptomycin})$ or $c_{opt}(\text{cycloheximide})$ with an "Ultragas 3" CO₂ analyzer (WösthoffCo., Bochum) (Anderson and Domsch, 1973) with continuous gas flow and we used this method to determine optimal concentrations for SIRIN and used these concentrations for the modified SIRIN approach as well. This optimization procedure was not used in other studies (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013). We supposed that optimal concentrations for CO₂ respiration should work as well for denitrification, if both inhibitors are apt to inhibit the

940

950 denitrification process as well. However, although SIRIN has so far been tested with isolated cultures and soils for microbial growth on agar and for CO₂ production (Anderson and Domsch, 1975, 1973), but information on N₂O

producing processes, especially denitrification, is still lacking and should be investigated in further studies. ~~and processes can~~

~~completely, using also different inhibitors~~

4.2 Inhibitor effects

~~Even with both growth inhibitors (treatment D) N₂O production was large in all experiments, i.e., often not significantly smaller than in the other three treatments. Thus, we suppose similar contributions of non-inhibitable organisms in all treatments. Non-inhibitable organisms could be, for example, bacteria or fungi that are not in growth stage or may be not affected by inhibitors. Pan et al. (2019) These organisms could be archaea as well, which are also known to be capable of denitrification (Philippot et al., 2007; Hayatsu et al., 2008). It is known, that archaea are not affected by streptomycin or cycloheximide (Seo and DeLaune, 2010). However, effects of archaeal occurrence in soil or secondary effects on fungi or bacteria were not tested in this study. As stated before, Ladan and Jacinthe (2016) did not find effective inhibition of denitrification by either inhibitor for denitrification although streptomycin and cycloheximide are commonly used to inhibit denitrification of selective groups. Thus, similar experiments with different inhibitors, such as the bactericide bronopol and the fungicide captan presented by Ladan and Jacinthe (2016), should be conducted to evaluate inhibition approaches and isotopic endmember approaches.~~

4.3-2 ~~Is SIRIN without~~ ~~Is~~ ~~C₂H₂ suitable~~ ~~application a suitable and necessary treatment for examining to~~ ~~examine the fungal contribution to N₂O production in soil?~~

In order to determine SP_{N_2O} values without alteration by partial reduction of N₂O to N₂, C₂H₂ was used to quantitatively block N₂O reduction during denitrification. We found the expected effect of C₂H₂ application, i.e. larger N₂O production rates in variety +C₂H₂ compared to variety -C₂H₂. Calculated product ratios varied between 0.5 and 0.95 (r_{15N}) in all ~~Soils~~soils, showing that N₂O reduction can have significant effects on measured N₂O production and isotopic values. ~~The product ratio is controlled by the reaction rate or by the activity of enzymes capable of N₂O reduction (Nos) in the system.~~

The calculated $r_{C_2H_2}$ was within the same range as r_{15N} in ~~Experiment~~Soil 1.2, 2, and 3 and 4 (maximal 9% difference), ~~providing the~~indicating effective blockage of N₂O reductase in variety +C₂H₂ in these ~~Soils~~soils. Only in ~~Experiment 1~~Soil 1.1, r_{15N} and $r_{C_2H_2}$ differed by about 34% with larger calculated reduction in the ~~tracer-traced~~ variety, which might ~~be explained by potential~~point to incomplete inhibition by the C₂H₂ method. ~~Nadeem et al. (2013) found some~~Artifacts with C₂H₂ ~~were found in previous studies, which~~resultinged in smaller N₂O production rates due to NO oxidation accelerated by C₂H₂ application in the presence of very small ~~oxygen (O)~~ amounts ($\geq 0.19 \text{ mL L}^{-1}$) (Bollmann and Conrad, 1997b, a; Nadeem et al., 2013). Moreover, incomplete C₂H₂ diffusion into denitrifying aggregates might also lead to incomplete N₂O reductase blockage (Groffman et al., 2006). Both potential methodological errors cannot be excluded for ~~Experiment 1~~Soil 1.1.

For the other three ~~experiment~~Soils (1.2, 2, and 3 and 4), it can be supposed that the isotopic signature of N₂O of variety +C₂H₂ showed isotopic signatures of produced N₂O without influences of N₂O reduction (SP_{prod}). By comparing varieties -C₂H₂ and +C₂H₂, isotopologue values of all ~~these~~Soils (except $\delta^{15N}_{bulk_{N_2O}}$ values of ~~Experiment~~Soil 2) of variety -C₂H₂ were significantly larger than those ~~that~~of variety +C₂H₂. The enrichment of residual N₂O in heavy isotopes results from the isotope effect associated with N₂O reduction (Jinuntuya-Nortman

et al., 2008; Well and Flessa, 2009; Lewicka-Szczebak et al., 2014). This explains why C₂H₂ application is essential for ~~analyzing analysing~~ N₂O produced by different microbial ~~organism~~ groups from soil, ~~using solely the modified SIRIN approach without additional isotopic approaches.~~ This has particular relevance for experiments with ~~modified SIRIN approaches.~~

995 ~~Moreover, when applying~~ Although the modified SIRIN approach presented here was not successful, it should be ~~noted that comparable soil incubation experiments without quantifying N₂O reduction, potentially overestimate fungal denitrification is potentially overestimated~~ due to the impact of SIRIN inhibitors on N₂O reduction.

1000 ~~It is evident that~~ Of course, N₂O fluxes represent net N₂O production, i.e. the difference between gross N₂O production by the microbial community and N₂O reduction, mainly by heterotrophic bacterial denitrifiers (Müller and Clough, 2014). ~~The goal of SIRIN application has been to determine the contribution of fungi and bacteria, respectively, to net N₂O production.~~ It has been shown that N₂O released by microorganisms to air-filled pore space can be partially consumed by denitrifiers before being emitted (Clough et al., 1998). This means that fungal N₂O can also be subject to reduction by bacterial denitrifiers. Consequently, ~~inhibiting successful inhibition of~~

1005 ~~bacterial denitrification by SIRIN would enhance lead the measured flux of fungal to an overestimation of fungal contribution to N₂O production.~~ Until now, this effect has not been considered in ~~previous~~ SIRIN papers on fungal N₂O (e. g. Laughlin and Stevens, 2002; Ladan and Jacinthe, 2016; Chen et al., 2014). This effect can only be evaluated by measuring N₂O reduction in all inhibitor treatments ~~as in our study~~. If true, the N₂O reduction with bacterial inhibition should be smaller than that of the treatments without inhibition or with fungal inhibition.

1010 ~~Though~~ However, with fungal inhibition, N₂O reduction is also assumed to be smaller ~~than~~ that without inhibition, because N₂O produced by fungi is missed for bacterial reduction.

1015 ~~As the product ratio in soil denitrification exhibited the full range from 0 to 1, this effect can be quite relevant and must thus be considered in future studies. Therefore, we recommend to estimate the effectiveness of C₂H₂ in blocking the N₂O reductase by performing parallel ¹⁵N approaches with and without C₂H₂ in studies using the modified SIRIN to determine the fraction of bacterial or fungal N₂O production.~~

1020 ~~The product ratio is a measure for the N₂O reduction to N₂. However, regarding the ¹⁵N, there was no evidence of different N₂O reduction effects between the SIRIN treatments. The ¹³C₂H₂ also revealed indistinguishable values between SIRIN treatments in Experiment 1 and 4, but it was slightly larger in Experiment 3 with bacterial inhibition compared to the other treatments. However, this effect was very small, which would only cause small overestimation of fungal contribution. The smallest N₂O reduction was found in Experiment 2 (¹³C₂H₂ values near 1), with smallest ¹³C₂H₂ with bacterial inhibition (0.81). This could result in an overestimation of bacterial contribution, since with blockage of N₂O reduction, gross N₂O production of bacteria is measured.~~

1025 ~~The ¹⁵N and ¹³C₂H₂ were between 0.5 and 1 and N₂O reduction was thus never consuming most of the produced N₂O. Hence, both the C₂H₂ and Streptomycin effects on SIRIN results were probably low. But However, as the product ratio in soil denitrification exhibited the full range from 0 to 1, meaning that this effect can be quite relevant and must thus be considered in future studies. Therefore, we recommend~~

~~As the product ratio in soil denitrification exhibited the full range from 0 to 1, this effect can be quite relevant and must thus be considered in future studies. Therefore, we recommend to estimate the effectiveness of C₂H₂ in blocking the N₂O reductase by performing parallel ¹⁵N approaches with and without C₂H₂ in studies using the modified SIRIN to determine the fraction of bacterial or fungal N₂O production.~~

~~The product ratio is a measure for the N₂O reduction to N₂. However, regarding the ¹⁵N, there was no evidence of different N₂O reduction effects between the SIRIN treatments. The ¹³C₂H₂ also revealed indistinguishable values between SIRIN treatments in Experiment 1 and 4, but it was slightly larger in Experiment 3 with bacterial inhibition compared to the other treatments. However, this effect was very small, which would only cause small overestimation of fungal contribution. The smallest N₂O reduction was found in Experiment 2 (¹³C₂H₂ values near 1), with smallest ¹³C₂H₂ with bacterial inhibition (0.81). This could result in an overestimation of bacterial contribution, since with blockage of N₂O reduction, gross N₂O production of bacteria is measured.~~

~~The ¹⁵N and ¹³C₂H₂ were between 0.5 and 1 and N₂O reduction was thus never consuming most of the produced N₂O. Hence, both the C₂H₂ and Streptomycin effects on SIRIN results were probably low. But However, as the product ratio in soil denitrification exhibited the full range from 0 to 1, meaning that this effect can be quite relevant and must thus be considered in future studies. Therefore, we recommend~~

4.4.3 *SP*_{N₂O} values of N₂O produced by microbial communities

~~As discussed above, all N₂O fluxes of modified SIRIN treatments of Soil 1.1, 1.2, and 3 were largely affected dominated by N₂O from non-inhibitable organisms or processes, which of course have an impact on *SP*_{N₂O} values of all SIRIN treatments. This made it impossible to calculate *SP*_{N₂O} values for active bacteria or fungi~~

1030 (modified SIRIN B and C), also with *Soil 2*, where a relatively large N₂O production was observed with treatment
D (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015). ~~This is discussed in more detail in (see section 3.4).~~
1035 ~~Despite this,~~ the SP_{N_2O} values from +C₂H₂ variety as well as SP_{prod} values (i.e. reduction corrected SP_{N_2O} values
of -C₂H₂ variety) values of each *Soil*, represented by treatment A of (modified SIRIN), indicated predominantly
bacteria to be responsible for N₂O production during denitrification, assuming that results of SP_{N_2O} values of
denitrification by pure bacterial cultures ~~is~~ ~~are~~ transferable to bacteria of soil communities contributing to
denitrification. ~~Also in many soil incubation studies, small SP_{N_2O} values (without reduction effects) within the~~
~~range of bacterial pure cultures have been found. The latter assumption has been confirmed repeatedly in soil~~
~~incubation studies, where in absence of N₂O reduction smallest SP_{N_2O} values have been found that were within the~~
~~range of bacterial pure cultures~~ (Lewicka-Szczebak et al., 2015; Lewicka-Szczebak et al., 2017; Senbayram et al.,
1040 2018). Therefore, there was ~~so far~~ no unequivocal evidence of fungi contributing to N₂O production during
denitrification ~~in soils~~, although ~~here~~, the isotopic approaches ~~revealed were consistent with a fungal~~ contribution
~~to N₂O production during denitrification~~ of up to 0.20 % ~~on of~~ N₂O production during denitrification.
The SP_{N_2O} values of ~~treatment A within~~ variety +C₂H₂ ~~within treatment A showed that are~~ ~~not affected by reduction~~
~~effects the signature of produced N₂O was not affected by reduction effects~~ and ~~therefore~~ might give evidence of
1045 the microbial community contributing to N₂O production ~~regarding differences in SP_{N_2O} values of pure bacterial~~
~~or fungal culture studies~~ (Sutka et al., 2006; Sutka et al., 2008; Frame and Casciotti, 2010; Rohe et al., 2014a).
However, variations in SP_{N_2O} values of treatments A of variety +C₂H₂ ~~are~~ ~~were~~ very small and do not give a clear
evidence of any differences in microbial soil community producing N₂O. ~~Lewicka-Szczebak et al. (2014) analyzed~~
~~analysed~~ SP_{N_2O} values of denitrification with blockage of N₂O reduction by C₂H₂ for the same soils as used in the
1050 present study ~~for Experiment (Soil 1.1 and 1.2 as well as Experiment Soil 3)~~ and revealed SP_{N_2O} values between
-3.6 and -2.1 ‰, which is similar to the respective SP_{N_2O} values of the present study from -4.9 to -0.4 ‰. This
reinforces the conclusion that bacteria dominated gross N₂O production under anoxic conditions in both ~~these~~
~~soils studies. Obviously,~~
 ~~SP_{prod} values (variety -C₂H₂) differed from SP_{N_2O} values (variety +C₂H₂), which may result from deviations~~
1055 ~~between the actual fractionation factor that was not estimated in the present study and the used fractionation factor~~
~~of -6 ‰ adapted from the literature (Yu et al., 2020). If so, we could assume smaller fractionation effects in the~~
~~present study as decreasing this average fractionation factor would lead to increasing SP_{prod} values, which in turn~~
~~would result in values more similar to SP_{N_2O} values of variety -C₂H₂.~~
~~However, other studies found larger SP_{N_2O} values of produced N₂O (up to +621 ‰) unaffected by the reduction~~
1060 ~~effect of up to +6 ‰ (Köster et al., 2013a) (Senbayram et al., 2018, 2020), most probably as a result of larger~~
~~contributions of fungi to N₂O production. However, those results were obtained in an experimental setup with~~
~~ambient oxygen concentration, without glucose amendment and without C₂H₂ inhibition of N₂O reduction since~~
~~N₂ gas fluxes were directly measured. It was also discussed before that short time incubations under static~~
~~conditions as presented here, may promote bacterial over fungal growth, which may also be transferable to~~
1065 ~~denitrification activity by both organism groups (Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2014).~~
~~Additionally to this, the selection use of glucose as substrate in the selected concentration may further promote~~
~~bacteria compared to fungi even more (Koranda et al., 2014; Reischke et al., 2014).~~

4.5 $\delta^{18}O_{N_2O}$ values

1070 The analysis of $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values can give information about O exchange between water and denitrification intermediates by various microorganisms. The range of $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values in our study for variety +C₂H₂ (7.5 to 19.0 %) was quite similar to the range found by for the same soils (4.8 to 16.3 %), where almost complete O exchange with soil water was documented. Hence, for this study the O exchange was probably also very high. However, there were no remarkable differences in $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values among treatments within one variety and soil and therefore we assume no differences in O exchange among the treatments.

1075 The information on $\delta^{18}\text{O}_{\text{N}_2\text{O}}$ values combined with known $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values is also precious information for differentiation between N₂O mixing and reduction processes (Lewicka-Szezebak et al., 2017). However, for this study, $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values were not analyzed. However, due to parallel *traced* variety experiments, we could determine possible $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values for the particular SP_{N₂O} values of bacterial denitrification mixing endmembers (Table 4). Since the $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value for the particular geographic region can be assessed based on the known isotopic signatures of meteoric waters, the most plausible ranges of $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values can be used to indicate the plausible ranges of α_{MAP} values. In case of precisely determined $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values, the calculated α_{MAP} values could be more precise, however, here we show that in case of missing $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values but known product ratio, the SP/ $\delta^{18}\text{O}$ Map can also provide information on N₂O production pathway contributions.

1085 4.6.4 Potential influence of Hybrid N₂O denitrification

1090 When one N atom in N₂O originates from labeled NO₃⁻ and the other one from an unlabeled N source, this results in α_p values and ¹⁵N enrichment of produced N₂O smaller than the respective enrichment of the NO₃⁻ pool. The ¹⁵N enrichment of N₂O in Soil 1.2 was about 60 % smaller than the ¹⁵N enrichment in soil NO₃⁻, leading to the assumption that N₂O was produced not only by denitrification. We also calculated α_p values of the other three Soils (data not shown) which coincided with the ¹⁵N enrichment of N₂O (Table 3), showing no indication of hybrid N₂O. Since α_p would not be affected by contributions of unlabeled N₂O we can thus exclude the possibility that this smaller enrichment could be caused by dilution of enriched N₂O from denitrification by N₂O production from an unknown N source and thus verified that this was due to formation of hybrid N₂O, potentially via co-denitrification (Spott et al., 2011). The influence of co-denitrification, which is predominantly associated to fungi (Spott et al., 2011), may have a large impact on N₂O production, since Laughlin and Stevens (2002) found N₂O production in their experiment derived to 92% from co-denitrification and only 8% from denitrification. So far, there is no study on SP_{N₂O} values of N₂O produced by co-denitrification. Co-denitrification could have been a contributing process in Experiment 1. When N in N₂O originates only from ¹⁵N labeled soil NO₃⁻, measured $\delta^{15}\text{N}^{\text{bulk}}_{\text{N}_2\text{O}}$ values as well as the ¹⁵N enrichment of the labelled N pool producing N₂O (α_p) should show identical ¹⁵N enrichment to the labeled soil NO₃⁻. During co-denitrification, when one N atom in N₂O originates from labeled NO₃⁻ and the other one from another unlabeled and unknown N source, this results in α_p values and ¹⁵N enrichment of produced N₂O smaller than the respective enrichment of the NO₃⁻ pool. The ¹⁵N enrichment of soil NO₃⁻ was about 60% larger than the analyzed ¹⁵N enrichment in N₂O, leading to the assumption that N₂O was produced not only by denitrification. We also calculated α_p values of the other three experiments (data not shown) which coincided with the ¹⁵N enrichment of N₂O (Table 3). Since α_p would not be affected by contributions of unlabelled N₂O we can thus exclude the possibility that this smaller enrichment could be caused by dilution of enriched N₂O from denitrification by N₂O production from an unknown N source and thus verified that this was due to formation of hybrid N₂O, probably via co-denitrification (Spott et al., 2011). In the other experiments there was no indication

of co-denitrification being relevant for N₂O production since ¹⁵N enrichments of NO₃⁻ and N₂O coincided. The question arises, why hybrid N₂O formation was only found when the loamy sand was sampled in summer (June, Experiment 1) but not when it was sampled during winter (December, Experiment 1). Information on substrates for co-denitrification, i.e. NO₂⁻ and NH₄⁺ or certain organic N compounds could have been different due to seasonal effects. Moreover, seasonal impacts on microbial community could have been relevant. Since these possible factors were not assessed in our study and their impact on co-denitrification is still poorly understood, it is currently not possible to give an answer here. But since SP_{N₂O} values of the acetylated treatments of Soil 1.2 coincided with the SP_{N₂O} value range of bacterial denitrification and also with SP_{N₂O} values of the other Soils, our data give no indication that the SP_{N₂O} values of hybrid N₂O, potentially produced during co-denitrification, differed from that of bacterial denitrification. It was however, remarkable that the maximum potential contribution of fungal denitrification to N₂O ($f_{FD, SP_{pot}}$) was higher for Soil 1.2 compared to that of Soil 1.1 from the winter period. Soil 1.2 was the only soil where $f_{FD, SP_{pot}}$ exceeded 50%, thus fungi may potentially dominate N₂O emissions only in this Soil. Thus, only the SP_{N₂O} values in Experiment 4 might be influenced by co-denitrification. But since SP_{N₂O} values of the acetylated treatments of Experiment 4 coincided with the SP_{N₂O} value range of bacterial denitrification and also with SP_{N₂O} values of the other experiments, our data give no indication that co-denitrification produces N₂O with SP_{N₂O} values differing from bacterial denitrification.

4.7-5 Calculating Steps towards quantifying the fungal fraction contributing to N₂O production and SP_{FD} values

Due to the inefficiency of the inhibition of microbial inhibition regarding N₂O production in most cases, calculation of f_{FDmi} contributing to N₂O production was only possible for Experiment Soil 2 only, although even this calculated value included inaccuracies. The isotopic approaches, however, which are independent of modified SIRIN results, yielded similar estimates of f_{FD} for all Soils, while it has to be emphasised that estimations based on stable isotope approaches do not rely on N₂O production of modified SIRIN results. As recently published (Wu et al., 2019), uncertainty analysis is a complex issue and large uncertainties of the results from the SP/ $\delta^{18}O$ Map approach can be assumed when all the possible sources of errors are taken into account. Regarding the presented application of SP/ $\delta^{18}O$ Map, calculation would be more precise when measuring $\delta^{18}O_{H_2O}$ rather than using the fitted $\delta^{18}O_{H_2O}$ values, as discussed above. Still, the analysis of $\delta^{18}O_{N_2O}$ values can give information about O exchange between water and denitrification intermediates by various microorganisms (Aerssens et al., 1986; Kool et al., 2007; Rohe et al., 2014b; Rohe et al., 2017). The range of $\delta^{18}O_{N_2O}$ values in our study for variety +C₂H₂ (7.5 to 19.0 ‰) was quite similar to the range found by Lewicka-Szczebak et al. (2014) for the same soils (4.8 to 16.3 ‰), where almost complete O exchange with soil water was documented. Hence, for this study the O exchange was probably also very high. There were also no remarkable differences in $\delta^{18}O_{N_2O}$ values among treatments within one variety and soil and therefore we assume no differences in O exchange among the treatments. The information on $\delta^{18}O_{N_2O}$ values combined with known $\delta^{18}O_{H_2O}$ values is also precious information for differentiation between N₂O mixing and reduction processes (Lewicka-Szczebak et al., 2017). Due to parallel traced variety experiments, possible $\delta^{18}O_{H_2O}$ values for the particular SP_{N₂O} values of bacterial denitrification mixing endmembers could be determined (Table 4). Since the $\delta^{18}O_{H_2O}$ value for the particular geographic region can be assessed based on the known isotopic signatures of meteoric waters (Lewicka-Szczebak et al., 2014; Stumpp et al., 2014; Lewicka-Szczebak et al., 2017; Buchen et al., 2018), the most plausible ranges of $\delta^{18}O_{H_2O}$ values can be used to indicate the plausible ranges of $f_{FD, MAP}$ values. Here we showed that in case of missing $\delta^{18}O_{H_2O}$ values but known product ratio, the SP/ $\delta^{18}O$ Map

1150 can also provide information on N₂O production pathway contributions. Comparing the modified SIRIN with the isotopic approaches revealed that the fungal fraction contribution to N₂O production was consistently estimated to be smaller (about 0.28 % in modified SIRIN, ≤0.15 % with IEM, ≤0.20 % with SP/δ¹⁸O Map) than the bacterial fraction. Although we did not obtain a very clear picture of various microorganisms contributing to N₂O production due to the large uncertainties of the calculated fractions, all approaches coincided by showing dominance of bacterial N₂O. In contrast to SIRIN, the isotopic approaches yielded similar estimates of F_{FD} for all experiments. 1155 This was supported by estimates for maximum potential contribution of fungal denitrification to N₂O in variety - C₂H₂ (f_{FD} SP_{prod}) for Soil 1.1, 2 and 3. In some soil studies using helium incubations, the SP_{prod} values obtained by correction for the reduction effect on SP_{N_2O} values showed significantly larger values than SP_{N_2O} of bacterial denitrification (Köster et al., 2013a; Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2014; Senbayram et al., 2018; Senbayram et al., 2020). However, those results were obtained in an experimental setup with ambient oxygen concentration. Short incubations under static conditions as presented here may, however, promote bacterial over fungal growth, which may also be transferable to denitrification activity by both organism groups (Lewicka-Szczebak et al., 2014; Lewicka-Szczebak et al., 2017). Therefore, it can be supposed Obviously, that based on the estimations from isotopic approaches, various soils may largely differ in the microbial community that contributes to N₂O from denitrification. 1160

1165 The However, all our three tested soils seemed to contain a microbial community where fungi have minor contributions to N₂O emissions from denitrification compared to bacteria. However, this may also have been due to the applied experimental setup favoring bacterial denitrification by static and strictly anoxic conditions. Additionally, the use of glucose as substrate in the selected concentration may further promote bacteria compared to fungi (Koranda et al., 2014; Reischke et al., 2014). and due to the choice of glucose as substrate. Senbayram et al. (2018) could show in an incubation experiment with sufficient NO₃⁻ supply, that fungal contribution to denitrification was larger with straw compared to a control without straw addition. Thus, experimental conditions need to be carefully set and more information is needed here in order to get a good representation of soil conditions in incubation experiments. 1170

1175 The fungal SP_{FD} values (section 3.6 “ SP of N₂O produced by the fungal soil community”) by SIRIN were highly variable with values between -23 and +25 ‰, which is smaller than the SP_{N_2O} range of N₂O known from pure cultures (16–37 ‰) (Sutka et al., 2008; Rohe et al., 2014a). Unfortunately, both ranges exhibit a large overlap but also some discrepancy, which precludes a clear conclusion whether or not Experiment 2 yielded valid estimates of fungal SP_{N_2O} values. There may be different reasons why estimating the SP_{N_2O} values using SIRIN of the fungal community was imprecise: the fungal fraction contributing to denitrification of the tested soils was only small compared to that of bacteria, SP_{N_2O} values were estimated using a large endmember range known from pure culture studies only, and possible SIRIN artefacts may have occurred as discussed above. The isotopic approaches should thus be further investigated with soils, where presumable fungi are presumed to contribute largely to N₂O production during (e. g. acid forest soils, or litter-amended arable soils) (Senbayram et al., 2018) and using SIRIN with more suitable inhibitors (Ladan and Jacinthe, 2016). The critical question whether the isotopic signatures of 1185 fungal N₂O determined in pure culture studies are transferable to natural soil conditions can not be fully answered with this study due to large uncertainties associated with the results of the SIRIN method. The latter precluded determination of making the SP_{N_2O} values of N₂O from fungal denitrification. Further experiments would be needed with improved selective inhibition to assure that SP_{N_2O} values known from a few pure cultures or soil isolates (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015) are true for fungal soil communities as well.

1190 This could be accompanied by studies mixing various fungal species known to occur in soil or by isolating fungal communities from soil and conduct similar experiments under anoxic conditions with supply of electron acceptors and C sources to investigate denitrification. In such incubations, parallel ^{15}N tracing experiments should be conducted to assure denitrification as the dominating process for N_2O production and quantify the possible contribution of co-denitrification.

1195 5. Conclusions

Based on the presented results we conclude that the modified SIRIN approach in the form presented here is not appropriate to estimate the contribution of selected communities (bacteria or fungi) on denitrification from soil. Here, the quantification of the fungal fraction with modified SIRIN could be done with one soil only and was possibly overestimated when compared with the results of isotopic approaches. Both isotope approaches (IEM and $\text{SP}/\delta^{18}\text{O}$ Map) revealed similar results of the fungal fraction contributing to denitrification and thus could be recommended as equally suitable for future studies. The present study could show that consideration of N_2O reduction is indispensable. It has to be pointed out, however, that the fungal fraction estimated applies only for the soil under presented experimental conditions, i.e. anaerobic conditions and with glucose amendment, but not for the investigated soil in general.

1200 Further studies are needed to cross-validate methods, e. g. with improved inhibitor approaches or molecular-based methods. Due to the mentioned difficulties selective inhibitor and isotopic approaches coincided in showing dominance of bacterial denitrification. Neither the modified SIRIN approach, nor IEM or $\text{SP}/\delta^{18}\text{O}$ Map approaches yielded larger contributions of the fungal N_2O fraction in any experiment. Both selective growth inhibitors of modified SIRIN confirmed the expected effect on N_2O production only in one out of four experiments. However, it has to be pointed out, that quantifying the fungal fraction with modified SIRIN was done with one soil only and was possibly overestimated when compared the results of isotopic approaches. According to this, the $\text{SP}_{\text{N}_2\text{O}}$ values of fungal N_2O could not be calculated from this modified SIRIN treatment did not appear to be a valid estimate of this value and need further evaluation. There might be several potential artefacts in the modified SIRIN approach should be, where further studies should focus on investigated, e.g. including the effectiveness of inhibitors, changes in microbial community during pre-incubation with inhibitors and effects of bacterial consumption of N_2O produced by fungi in the presence of bacterial growth inhibitors. The present study could show that consideration of N_2O reduction in further studies is indispensable inevitably necessary. Further studies should also determine the range of $\text{SP}_{\text{N}_2\text{O}}$ values known from fungal denitrification in soils as well as the effect of specific inhibitors on microbial groups producing N_2O and reducing N_2O during denitrification.

1220 *Data availability.* Gas emission and isotopic data are available from the authors on request.

Author contribution. HF, NWM, RW and THA designed the experiment. LR carried out the experiment at Thünen Institute for Climate-Smart Agriculture in Braunschweig. AG, DLS and RW helped with isotopic analysis and DLS performed the $\delta^{18}\text{O}/\text{SP}$ Map. LR, RW and DLS prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

1225 *Acknowledgements.* Many thanks are due to Jens Dyckmans for ¹⁵N analysis and to Martina Heuer for N₂O isotopic analyses. This joint research project was financially supported by the State of Lower-Saxony and the Volkswagen Foundation, Hanover, Germany. Further financial support was provided by the German Research Foundation (grant LE 3367/1-1 to DLS, grant WE 1904/8-1 to LR and RW, and the research unit 2337: “Denitrification in Agricultural Soils: Integrated Control and Modeling at Various Scales (DASIM)”, grant WE 1904/10-1 to RW and
1230 WR 211/1-2 to NWM).

References

- Aerssens, E., Tiedje, J. M., and Averill, B. A.: Isotope labeling studies on the mechanisms of N-bond formation in denitrification, *J. Biol. Chem.*, 261, 9652-9656, 1986.
- 1235 Anderson, J. P. E., and Domsch, K. H.: Quantification of bacterial and fungal contributions to soil respiration, *Archiv Fur Mikrobiologie*, 93, 113-127, doi:10.1007/BF00424942, 1973.
- Anderson, J. P. E., and Domsch, K. H.: Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soil, *Canadian Journal of Microbiology*, 21, 314-322, doi: 10.1139/m75-045, 1975.
- 1240 Anderson, J. P. E., and Domsch, K. H.: Physiological method for quantitative measurement of microbial biomass in soils, *Soil Biol. Biochem.*, 10, 215-221, doi:10.1016/0038-0717(78)90099-8, 1978.
- 1245 Bergsma, T. T., Ostrom, N. E., Emmons, M., and Robertson, G. P.: Measuring simultaneous fluxes from soil of N₂O and N₂ in the field using the ¹⁵N-Gas "nonequilibrium" technique, *Environmental Science & Technology*, 35, 4307-4312, doi: 10.1021/es010885u, 2001.
- 1250 Blagodatskaya, E., Dannenmann, M., Gasche, R., and Butterbach-Bahl, K.: Microclimate and forest management alter fungal-to-bacterial ratio and N₂O-emission during rewetting in the forest floor and mineral soil of mountainous beech forests, *Biogeochemistry*, 97, 55-70, doi:10.1007/s10533-009-9310-3, 2010.
- Blagodatskaya, E. V., and Anderson, T.-H.: Interactive effects of pH and substrate quality on the fungal-to-bacterial ratio and qCO₂ of microbial communities in forest soils, *Soil Biol. Biochem.*, 30, 1269-1274, doi: 10.1016/S0038-0717(98)00050-9, 1998.
- 1255 Bollag, J. M., and Tung, G.: Nitrous oxide release by soil fungi, *Soil Biol. Biochem.*, 4, 271-276, doi: 10.1016/0038-0717(72)90021-1, 1972.
- 1260 Bollmann, A., and Conrad, R.: Acetylene blockage technique leads to underestimation of denitrification rates in oxic soils due to scavenging of intermediate nitric oxide, *Soil Biol. Biochem.*, 29, 1067-1077, 1997a.
- Bollmann, A., and Conrad, R.: Enhancement by acetylene of the decomposition of nitric oxide in soil, *Soil Biol. Biochem.*, 29, 1057-1066, 1997b.
- 1265 Braker, G., and Conrad, R.: Diversity, structure, and size of N₂O-producing microbial communities in soils-What matters for their functioning?, in: *Advances in Applied Microbiology*, Vol 75, edited by: Laskin, A. I., Sariaslani, S., and Gadd, G. M., *Advances in Applied Microbiology*, 33-70, 2011.
- 1270 Brand, W. A.: Precon: A fully automated interface for the pre-GC concentration of trace gases in air for isotopic analysis, *Isot. Environ. Health S.*, 31, 277-284, doi: 10.1080/10256019508036271, 1995.
- Bremner, J.: Sources of nitrous oxide in soils, *Nutrient Cycling in Agroecosystems*, 49, 7-16, doi: 10.1023/A:1009798022569, 1997.
- 1275 Buchen, C., Lewicka-Szczepak, D., Flessa, H., and Well, R.: Estimating N₂O processes during grassland renewal and grassland conversion to maize cropping using N₂O isotopocules, *Rapid Commun. Mass Spec.*, 32, 1053-1067, doi: 10.1002/rcm.8132, 2018.

- 1280 Casciotti, K. L., Sigman, D. M., Hastings, M. G., Böhlke, J. K., and Hilkert, A.: Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, 74, 4905-4912, doi: 10.1021/ac020113w, 2002.
- 1285 Chen, H., Mothapo, N. V., and Shi, W.: The significant contribution of fungi to soil N₂O production across diverse ecosystems, *Appl. Soil Ecol.*, 73, 70-77, doi: 10.1016/j.apsoil.2013.08.011, 2014.
- Chen, H. H., Mothapo, N. V., and Shi, W.: Soil Moisture and pH Control Relative Contributions of Fungi and Bacteria to N₂O Production, *Microb. Ecol.*, 69, 180-191, doi: 10.1007/s00248-014-0488-0, 2015.
- 1290 Clough, T. J., Jarvis, S. C., Dixon, E. R., Stevens, R. J., Laughlin, R. J., and Hatch, D. J.: Carbon induced subsoil denitrification of ¹⁵N-labelled nitrate in 1 m deep soil columns, *Soil Biol. Biochem.*, 31, 31-41, doi: 10.1016/S0038-0717(98)00097-2, 1998.
- 1295 Crenshaw, C. L., Lauber, C., Sinsabaugh, R. L., and Staveland, L. K.: Fungal control of nitrous oxide production in semiarid grassland, *Biogeochemistry*, 87, 17-27, doi: 10.1007/s10533-007-9165-4, 2008.
- Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal of the Royal Meteorological Society*, 96, 320-325, doi: 10.1002/qj.49709640815, 1970.
- 1300 Decock, C., and Six, J.: How reliable is the intramolecular distribution of ¹⁵N in N₂O to source partition N₂O emitted from soil?, *Soil Biol. Biochem.*, 65, 114-127, <http://dx.doi.org/10.1016/j.soilbio.2013.05.012>, 2013.
- Frame, C. H., and Casciotti, K. L.: Biogeochemical controls and isotopic signatures of nitrous oxide production by a marine ammonia-oxidizing bacterium, *Biogeosciences*, 7, 2695-2709, doi: 10.5194/bg-7-2695-2010, 2010.
- 1305 Garber, E. A., and Hollocher, T. C.: ¹⁵N, ¹⁸O tracer studies on the activation of nitrite by denitrifying bacteria. Nitrite/water-oxygen exchange and nitrosation reactions as indicators of electrophilic catalysis, *J. Biol. Chem.*, 257, 8091-8097, doi: 10.1007/BF00399539, 1982.
- 1310 Groffman, P. M., Altabet, M. A., Böhlke, J. K., Butterbach-Bahl, K., David, M. B., Firestone, M. K., Giblin, A. E., Kana, T. M., Nielsen, L. P., and Voytek, M. A.: Methods for measuring denitrification: Diverse approaches to a difficult problem, *Ecological Applications*, 16, 2091-2122, 2006.
- 1315 Hayatsu, M., Tago, K., and Saito, M.: Various players in the nitrogen cycle: Diversity and functions of the microorganisms involved in nitrification and denitrification, *Soil Science & Plant Nutrition*, 54, 33-45, <https://doi.org/10.1111/j.1747-0765.2007.00195.x>, 2008.
- 1320 Higgins, S. A., Schadt, C. W., Matheny, P. B., and Löffler, F. E.: Phylogenomics reveal the dynamic evolution of fungal nitric oxide reductases and their relationship to secondary metabolism, *Genome Biology and Evolution*, 10, 2474-2489, doi: 10.1093/gbe/evy187, 2018.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.
- 1325 Jinuntuya-Nortman, M., Sutka, R. L., Ostrom, P. H., Gandhi, H., and Ostrom, N. E.: Isotopologue fractionation during microbial reduction of N₂O within soil mesocosms as a function of water-filled pore space, *Soil Biol. Biochem.*, 40, 2273-2280, doi: 10.1016/j.soilbio.2008.05.016, 2008.
- 1330 Keuschnig, C., Gorfer, M., Li, G., Mania, D., Frostegård, Å., Bakken, L., and Larose, C.: NO and N₂O transformations of diverse fungi in hypoxia: evidence for anaerobic respiration only in *Fusarium* strains, *Environmental Microbiology*, 22, 2182-2195, doi: 10.1111/1462-2920.14980, 2020.
- Knowles, R.: Denitrification, *Microbiol. Rev.*, 46, 43-70, 1982.
- 1335 Kool, D. M., Wrage, N., Oenema, O., Dolfing, J., and Van Groenigen, J. W.: Oxygen exchange between (de)nitrification intermediates and H₂O and its implications for source determination of NO₃⁻ and N₂O: a review, *Rapid Commun. Mass Spec.*, 21, 3569-3578, doi: 10.1002/rcm.3249, 2007.

- 1340 Kool, D. M., Wrage, N., Oenema, O., Harris, D., and Van Groenigen, J. W.: The ^{18}O signature of biogenic nitrous oxide is determined by O exchange with water, *Rapid Commun. Mass Spec.*, 23, 104-108, doi: 10.1002/rcm.3859, 2009.
- 1345 Koranda, M., Kaiser, C., Fuchslueger, L., Kitzler, B., Sessitsch, A., Zechmeister-Boltenstern, S., and Richter, A.: Fungal and bacterial utilization of organic substrates depends on substrate complexity and N availability, *FEMS Microbiology Ecology*, 87, 142-152, doi: 10.1111/1574-6941.12214, 2014.
- 1350 Köster, J. R., Well, R., Dittert, K., Giesemann, A., Lewicka-Szczebak, D., Muehling, K.-H., Herrmann, A., Lammel, J., and Senbayram, M.: Soil denitrification potential and its influence on N_2O reduction and N_2O isotopomer ratios, *Rapid Commun. Mass Spec.*, 27, 2363-2373, doi: 10.1002/rcm.6699, 2013a.
- 1355 Köster, J. R., Well, R., Tuzson, B., Bol, R., Dittert, K., Giesemann, A., Emmenegger, L., Manninen, A., Cárdenas, L., and Mohn, J.: Novel laser spectroscopic technique for continuous analysis of N_2O isotopomers – application and intercomparison with isotope ratio mass spectrometry, *Rapid Commun. Mass Spec.*, 27, 216-222, doi: 10.1002/rcm.6434, 2013b.
- 1360 Ladan, S., and Jacinthe, P.-A.: Evaluation of antibacterial and antifungal compounds for selective inhibition of denitrification in soils, *Environmental Science: Processes & Impacts*, 18, 1519-1529, doi: 10.1039/C6EM00456C, 2016.
- 1365 Laughlin, R. J., and Stevens, R. J.: Evidence for fungal dominance of denitrification and codenitrification in a grassland soil, *Soil Sc. Soc. Am. J.*, 66, 1540-1548, doi: 10.2136/sssaj2002.1540, 2002.
- Lewicka-Szczebak, D., Well, R., Giesemann, A., Rohe, L., and Wolf, U.: An enhanced technique for automated determination of ^{15}N signatures of N_2 , ($\text{N}_2+\text{N}_2\text{O}$) and N_2O in gas samples, *Rapid Commun. Mass Spec.*, 27, 1548-1558, <https://doi.org/10.1002/rcm.6605>, 2013.
- 1370 Lewicka-Szczebak, D., Well, R., Köster, J. R., Fuß, R., Senbayram, M., Dittert, K., and Flessa, H.: Experimental determinations of isotopic fractionation factors associated with N_2O production and reduction during denitrification in soils, *Geochim. Cosmochim. Ac.*, 134, 55-73, doi: 10.1016/j.gca.2014.03.010, 2014.
- Lewicka-Szczebak, D., Well, R., Bol, R., Gregory, A. S., Matthews, G. P., Misselbrook, T., Whalley, W. R., and Cardenas, L. M.: Isotope fractionation factors controlling isotopocule signatures of soil-emitted N_2O produced by denitrification processes of various rates, *Rapid Commun. Mass Spec.*, 29, 269-282, doi: 10.1002/rcm.7102, 2015.
- 1375 Lewicka-Szczebak, D., Dyckmans, J., Kaiser, J., Marca, A., Augustin, J., and Well, R.: Oxygen isotope fractionation during N_2O production by soil denitrification, *Biogeosciences*, 13, 1129-1144, doi: 10.5194/bg-13-1129-2016, 2016.
- 1380 Lewicka-Szczebak, D., Augustin, J., Giesemann, A., and Well, R.: Quantifying N_2O reduction to N_2 based on N_2O isotopocules – validation with independent methods (helium incubation and ^{15}N gas flux method), *Biogeosciences*, 14, 711-732, doi: 10.5194/bg-14-711-2017, 2017.
- 1385 Lewicka-Szczebak, D., Lewicki, M. P., and Well, R.: N_2O isotope approaches for source partitioning of N_2O production and estimation of N_2O reduction – validation with the ^{15}N gas-flux method in laboratory and field studies, *Biogeosciences*, 17, 5513-5537, doi: 10.5194/bg-17-5513-2020, 2020.
- 1390 Long, A., Heitman, J., Tobias, C., Philips, R., and Song, B.: Co-occurring anammox, denitrification, and codenitrification in agricultural soils, *Applied and Environmental Microbiology*, 79, 168-176, doi: 10.1128/aem.02520-12, 2013.
- Maeda, K., Spor, A., Edel-Hermann, V., Heraud, C., Breuil, M.-C., Bizouard, F., Toyoda, S., Yoshida, N., Steinberg, C., and Philippot, L.: N_2O production, a widespread trait in fungi, *Scientific Reports*, 5, 9697, doi: 10.1038/srep09697, 2015.
- 1395 McLain, J. E. T., and Martens, D. A.: N_2O production by heterotrophic N transformations in a semiarid soil, *Appl. Soil. Ecol.*, 32, 253-263, 2006.
- Müller, C., and Clough, T. J.: Advances in understanding nitrogen flows and transformations: gaps and research pathways, *J. Agric. Sci.*, 152, S34-S44, <https://doi.org/10.1017/s0021859613000610>, 2014.

- 1400 Nadeem, S., Dorsch, P., and Bakken, L. R.: Autoxidation and acetylene-accelerated oxidation of NO in a 2-phase system: Implications for the expression of denitrification in ex situ experiments, *Soil Biol. Biochem.*, 57, 606-614, doi: 10.1016/j.soilbio.2012.10.007, 2013.
- 1405 Ostrom, N., and Ostrom, P.: The Isotopomers of Nitrous Oxide: Analytical Considerations and Application to Resolution of Microbial Production Pathways, in: *Handbook of Environmental Isotope Geochemistry*, edited by: Baskaran, M., *Advances in Isotope Geochemistry*, Springer Berlin Heidelberg, 453-476, 2011.
- 1410 Ostrom, N. E., Pitt, A., Sutka, R., Ostrom, P. H., Grandy, A. S., Huizinga, K. M., and Robertson, G. P.: Isotopologue effects during N₂O reduction in soils and in pure cultures of denitrifiers, *Journal of Geophysical Research-Biogeosciences*, 112, doi: 10.1029/2006jg000287, 2007.
- 1415 Ostrom, N. E., Sutka, R., Ostrom, P. H., Grandy, A. S., Huizinga, K. M., Gandhi, H., von Fischer, J. C., and Robertson, G. P.: Isotopologue data reveal bacterial denitrification as the primary source of N₂O during a high flux event following cultivation of a native temperate grassland, *Soil Biol. Biochem.*, 42, 499-506, doi: 10.1016/j.soilbio.2009.12.003, 2010.
- 1420 Ostrom, N. E., and Ostrom, P. H.: Mining the isotopic complexity of nitrous oxide: a review of challenges and opportunities, *Biogeochemistry*, 132, 359-372, doi: 10.1007/s10533-017-0301-5, 2017.
- Pan, Y. S., Wu, Y. C., Li, X. Z., Zeng, J., and Lin, X. G.: Continuing Impacts of Selective Inhibition on Bacterial and Fungal Communities in an Agricultural Soil, *Microb. Ecol.*, 78, 927-935, doi: 10.1007/s00248-019-01364-0, 2019.
- 1425 Philippot, L., Hallin, S., and Schloter, M.: Ecology of denitrifying prokaryotes in agricultural soil, in: *Advances in Agronomy*, edited by: Donald, L. S., Academic Press, 249-305, 2007.
- 1430 Phillips, R., Grelet, G., McMillan, A., Song, B., Weir, B., Palmada, T., and Tobias, C.: Fungal denitrification: *Bipolaris sorokiniana* exclusively denitrifies inorganic nitrogen in the presence and absence of oxygen, *FEMS Microbiol. Lett.*, 363, 5, doi: 10.1093/femsle/fnw007, 2016a.
- Phillips, R. L., Song, B., McMillan, A. M. S., Grelet, G., Weir, B. S., Palmada, T., and Tobias, C.: Chemical formation of hybrid di-nitrogen calls fungal codenitrification into question, *Scientific Reports*, 6, 39077, doi: 10.1038/srep39077, 2016b.
- 1435 Popp, B. N., Westley, M. B., Toyoda, S., Miwa, T., Dore, J. E., Yoshida, N., Rust, T. M., Sansone, F. J., Russ, M. E., Ostrom, N. E., and Ostrom, P. H.: Nitrogen and oxygen isotopomeric constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochem. Cy.*, 16, doi: 10.1029/2001gb001806, 2002.
- 1440 Reischke, S., Rousk, J., and Bååth, E.: The effects of glucose loading rates on bacterial and fungal growth in soil, *Soil Biol. Biochem.*, 70, 88-95, doi: 10.1016/j.soilbio.2013.12.011, 2014.
- 1445 Rohe, L., Anderson, T.-H., Braker, G., Flessa, H., Giesemann, A., Lewicka-Szczebak, D., Wrage-Mönnig, N., and Well, R.: Dual isotope and isotopomer signatures of nitrous oxide from fungal denitrification – a pure culture study, *Rapid Communications in Mass Spectrometry* 28, 1893-1903, doi: 10.1002/rcm.6975, 2014a.
- Rohe, L., Anderson, T.-H., Braker, G., Flessa, H., Giesemann, A., Wrage-Mönnig, N., and Well, R.: Fungal oxygen exchange between denitrification intermediates and water, *Rapid Commun. Mass Spec.*, 28, 377-384, doi: 10.1002/rcm.6790, 2014b.
- 1450 Rohe, L., Well, R., and Lewicka-Szczebak, D.: Use of oxygen isotopes to differentiate between nitrous oxide produced by fungi or bacteria during denitrification, *Rapid Commun. Mass Spec.*, 31, 1297-1312, doi: 10.1002/rcm.7909, 2017.
- 1455 Ruzicka, S., Edgerton, D., Norman, M., and Hill, T.: The utility of ergosterol as a bioindicator of fungi in temperate soils, *Soil Biol. Biochem.*, 32, 989-1005, doi: 10.1016/S0038-0717(00)00009-2, 2000.
- 1460 Senbayram, M., Well, R., Bol, R., Chadwick, D. R., Jones, D. L., and Wu, D.: Interaction of straw amendment and soil NO₃⁻ content controls fungal denitrification and denitrification product stoichiometry in a sandy soil, *Soil Biol. Biochem.*, 126, 204-212, doi: 10.1016/j.soilbio.2018.09.005, 2018.

- Senbayram, M., Well, R., Shan, J., Bol, R., Burkart, S., Jones, D. L., and Wu, D.: Rhizosphere processes in nitrate-rich barley soil tripled both N₂O and N₂ losses due to enhanced bacterial and fungal denitrification, *Plant and Soil*, 448, 509-522, doi: 10.1007/s11104-020-04457-9, 2020.
- 1465 Seo, D. C., and DeLaune, R. D.: Fungal and bacterial mediated denitrification in wetlands: Influence of sediment redox condition, *Water Research*, 44, 2441-2450, doi: 10.1016/j.watres.2010.01.006, 2010.
- 1470 Shoun, H., and Tanimoto, T.: Denitrification by the fungus *Fusarium oxysporum* and involvement of cytochrome P-450 in the respiratory nitrite reduction, *J. Biol. Chem.*, 266, 11078-11082, 1991.
- Shoun, H., Kim, D.-H., Uchiyama, H., and Sugiyama, J.: Denitrification by fungi, *FEMS Microbiol. Lett.*, 94, 277-281, 1992.
- 1475 Shoun, H., Fushinobu, S., Jiang, L., Kim, S. W., and Wakagi, T.: Fungal denitrification and nitric oxide reductase cytochrome P450nor, *Philos. T. Roy. Soc.*, 367, 1186-1194, 10.1098/rstb.2011.0335, 2012.
- 1480 Snider, D. M., Schiff, S. L., and Spoelstra, J.: ¹⁵N/¹⁴N and ¹⁸O/¹⁶O stable isotope ratios of nitrous oxide produced during denitrification in temperate forest soils, *Geochim. Cosmochim. Ac.*, 73, 877-888, 10.1016/j.gca.2008.11.004, 2009.
- 1485 Spott, O., Russow, R., Apelt, B., and Stange, C. F.: A ¹⁵N-aided artificial atmosphere gas flow technique for online determination of soil N₂ release using the zeolite K strolith SX6 , *Rapid Commun. Mass Spec.*, 20, 3267-3274, <https://doi.org/10.1002/rcm.2722>, 2006.
- Spott, O., Russow, R., and Stange, C. F.: Formation of hybrid N₂O and hybrid N₂ due to codenitrification: First review of a barely considered process of microbially mediated N-nitrosation, *Soil Biol. Biochem.*, 43, 1995-2011, doi: 10.1016/j.soilbio.2011.06.014, 2011.
- 1490 Stump, C., Klaus, J., and Stichler, W.: Analysis of long-term stable isotopic composition in German precipitation, *Journal of Hydrology*, 517, 351-361, doi: 10.1016/j.jhydrol.2014.05.034, 2014.
- 1495 Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Breznak, J. A., Gandhi, H., Pitt, A. J., and Li, F.: Distinguishing nitrous oxide production from nitrification and denitrification on the basis of isotopomer abundances, *Appl. Environ. Microbiol.*, 72, 638-644, doi: 10.1128/aem.72.1.638-644.2006, 2006.
- Sutka, R. L., Adams, G. C., Ostrom, N. E., and Ostrom, P. H.: Isotopologue fractionation during N₂O production by fungal denitrification, *Rapid Commun. Mass Spec.*, 22, 3989-3996, doi: 10.1002/rcm.3820, 2008.
- 1500 Tanimoto, T., Hatano, K., Kim, D. H., Uchiyama, H., and Shoun, H.: Co-denitrification by the denitrifying system of fungus *Fusarium oxysporum*, *FEMS Microbiol. Lett.*, 93, 177-180, 1992.
- 1505 Toyoda, S., and Yoshida, N.: Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer, *Anal. Chem.*, 71, 4711-4718, 10.1021/ac9904563, 1999.
- Toyoda, S., Mutobe, H., Yamagishi, H., Yoshida, N., and Tanji, Y.: Fractionation of N₂O isotopomers during production by denitrifier, *Soil Biol. Biochem.*, 37, 1535-1545, doi: 10.1016/j.soilbio.2005.01.009, 2005.
- 1510 Toyoda, S., Yano, M., Nishimura, S.-i., Akiyama, H., Hayakawa, A., Koba, K., Sudo, S., Yagi, K., Makabe, A., Tobar, Y., Ogawa, N. O., Ohkouchi, N., Yamada, K., and Yoshida, N.: Characterization and production and consumption processes of N₂O emitted from temperate agricultural soils determined via isotopomer ratio analysis, *Global Biogeochem. Cy.*, 25, doi: 10.1029/2009gb003769, 2011.
- 1515 USEPA: Manual: Nitrogen Control, Office of Water, Washington, D.C. , EPA/625/R-93 /010, 1993.
- Well, R., Becker, K.-W., Meyer, B., Langel, R., and Reineking, A.: Continuous flow equilibration for mass spectrometric analysis of dinitrogen emissions, *Soil Sc. Soc. Am. J.*, 62, 906-910, doi: 10.2136/sssaj1998.03615995006200040008x, 1998.

- 1520 Well, R., Kurganova, I., de Gerenyu, V. L., and Flessa, H.: Isotopomer signatures of soil-emitted N₂O under different moisture conditions - A microcosm study with arable loess soil, *Soil Biol. Biochem.*, 38, 2923-2933, 10.1016/j.soilbio.2006.05.003, 2006.
- 1525 Well, R., and Flessa, H.: Isotopologue signatures of N₂O produced by denitrification in soils, *Journal of Geophysical Research: Biogeosciences*, 114, G02020, doi: 10.1029/2008jg000804, 2009.
- 1530 Well, R., Burkart, S., Giesemann, A., Grosz, B., Köster, J. R., and Lewicka-Szczebak, D.: Improvement of the ¹⁵N gas flux method for in situ measurement of soil denitrification and its product stoichiometry, *Rapid Commun. Mass Spec.*, 33, 437-448, <https://doi.org/10.1002/rcm.8363>, 2019.
- 1535 Wrage-Mönnig, N., Horn, M. A., Well, R., Müller, C., Velthof, G., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited, *Soil Biol. Biochem.*, 123, A3-A16, doi: 10.1016/j.soilbio.2018.03.020, 2018.
- 1540 Wrage, N., Velthof, G. L., Laanbroek, H. J., and Oenema, O.: Nitrous oxide production in grassland soils: assessing the contribution of nitrifier denitrification, *Soil Biol. Biochem.*, 36, 229-236, 10.1016/j.soilbio.2003.09.009, 2004a.
- 1545 Wrage, N., Velthof, G. L., Oenema, O., and Laanbroek, H. J.: Acetylene and oxygen as inhibitors of nitrous oxide production in *Nitrosomonas europaea* and *Nitrospira briensis*: a cautionary tale, *Fems Microbiology Ecology*, 47, 13-18, 10.1016/s0168-6496(03)00220-4, 2004b.
- 1550 Wu, D., Well, R., Cárdenas, L. M., Fuß, R., Lewicka-Szczebak, D., Köster, J. R., Brüggemann, N., and Bol, R.: Quantifying N₂O reduction to N₂ during denitrification in soils via isotopic mapping approach: Model evaluation and uncertainty analysis, *Environmental Research*, 179, 108806, <https://doi.org/10.1016/j.envres.2019.108806>, 2019.
- 1555 Yoshinari, T., and Knowles, R.: Acetylene inhibition of nitrous-oxide reduction by denitrifying bacteria, *Biochemical and Biophysical Research Communications*, 69, 705-710, doi: 10.1016/0006-291x(76)90932-3, 1976.
- 1560 Yu, L., Harris, E., Lewicka-Szczebak, D., Barthel, M., Blomberg, M. R. A., Harris, S. J., Johnson, M. S., Lehmann, M. F., Liisberg, J., Müller, C., Ostrom, N. E., Six, J., Toyoda, S., Yoshida, N., and Mohn, J.: What can we learn from N₂O isotope data? - Analytics, processes and modelling, *Rapid Commun Mass Spectrom*, doi: 10.1002/rcm.8858, 2020.
- Zou, Y., Hirono, Y., Yanai, Y., Hattori, S., Toyoda, S., and Yoshida, N.: Isotopomer analysis of nitrous oxide accumulated in soil cultivated with tea (*Camellia sinensis*) in Shizuoka, central Japan, *Soil Biol. Biochem.*, 77, 276-291, doi: 10.1016/j.soilbio.2014.06.016, 2014.

Supplementary Material for

Comparing modified substrate induced respiration with selective inhibition (SIRIN) and N₂O isotope approaches to estimate fungal contribution to denitrification in three arable soils under anoxic conditions

5 Lena Rohe^{1, 2, 3}, Traute-Heidi Anderson², Heinz Flessa², Anette **Giesemann²Goeske²**, Dominika Lewicka-Szczebak^{2, 4}, Nicole Wrage-Mönnig⁵, Reinhard Well²

¹Helmholtz Centre for Environmental Research – UFZ, Department Soil System Sciences, Theodor-Lieser Str. 4, Halle, Germany

²Thünen Institute of Climate Smart Agriculture, Bundesallee 65, Braunschweig, Germany

10 ³University of Göttingen, Department of Crop Sciences, Institute of Grassland Science, von-Siebold-Str. 8, 37075 Göttingen, Germany

⁴~~Institute of Geological Sciences, University of Wrocław, pl. M. Borna 9, 50-204 Wrocław, Poland
University of Göttingen, Centre for Stable Isotope Research and Analysis, Büsgenweg 2, 37077 Göttingen, Germany~~

15 ⁵University of Rostock, Agricultural and Environmental Faculty, Grassland and Fodder Sciences, Justus-Liebig-Weg 6, Rostock, Germany

Correspondence to: Lena Rohe (lena.rohe@ufz.de)

Table S1: Important terms used in the present study and descriptions of terms with presenting the associated sections.

<u>Term</u>	<u>Description</u>	<u>Eq.</u>	<u>Section</u>
<u>NO₃⁻</u>	<u>Nitrate: electron acceptor for denitrification</u>	/	<u>1, 2</u>
<u>NO₂⁻</u>	<u>Nitrite: electron acceptor for denitrification</u>		
<u>NO</u>	<u>Nitrogen monoxide: intermediate of denitrification</u>		
<u>N₂O</u>	<u>Nitrous oxide: intermediate or product of denitrification</u>		
<u>N₂</u>	<u>Dinitrogen: end product of denitrification</u>		
<u>KNO₃</u>	<u>Potassium nitrate: electron acceptor for denitrification</u>		
<u>NH₄⁺</u>	<u>Ammonia</u>		
<u>CO₂</u>	<u>Carbon dioxide: product of respiration</u>		
<u>C₂H₂</u>	<u>Acetylene used to block the N₂O reductase</u>		
<u>O</u>	<u>oxygen</u>		
<u>Nos</u>	<u>N₂O reductase</u>	/	<u>1</u>
<u>δ¹⁵N^{bulk}_{N₂O}</u>	<u>δ¹⁵N values of produced N₂O</u>	/	<u>1</u>
<u>δ¹⁵N_{NO_x}</u>	<u>δ¹⁵N^{bulk} values of N₂O precursors NO₃⁻ or NO₂⁻</u>	/	<u>2.1</u>
<u>SP_{N₂O}</u>	<u>¹⁵N site preference of N₂O; i.e. difference between δ¹⁵N of the central and terminal N-position of the asymmetric N₂O molecule (Toyoda and Yoshida, 1999).</u>	/	<u>1, 2.3, 2.5</u>
<u>δ¹⁸O_{N₂O}</u>	<u>δ¹⁸O values of produced N₂O</u>	/	<u>1</u>
<u>δ¹⁸O_{NO_x}</u>	<u>δ¹⁸O values of N₂O precursors NO₃⁻ or NO₂⁻</u>	/	<u>1</u>
<u>δ¹⁸O_{H₂O}</u>	<u>δ¹⁸O values of water (H₂O)</u>	/	<u>1, 2.5.2</u>
<u>Soil 1.1</u>	<u>loamy sand sampled in December 2012</u>	/	<u>2.1;</u>
<u>Soil 1.2</u>	<u>loamy sand sampled in June 2011</u>		<u>Table 1</u>
<u>Soil 2</u>	<u>sand sampled in January 2013</u>		
<u>Soil 3</u>	<u>silt loam sampled in December 2012</u>		
<u>F:B</u>	<u>Respiratory fungal-to-bacterial ratio analysed by SIRIN method (Anderson and Domsch, 1973, 1975)</u>	/	<u>1, 2.2;</u> <u>Table 1</u>
<u>SIR</u>	<u>Substrate-induced respiration (Anderson and Domsch, 1973, 1975, 1978)</u>	/	<u>2.2.1;</u> <u>Table 1</u>
<u>c_{opt}(cycloheximide), c_{opt}(streptomycin)</u>	<u>optimal concentration for inhibition of fungal respiration</u>		<u>2.1</u>
<u>SIRIN</u>	<u>Substrate-induced respiration with selective inhibition (Anderson and Domsch, 1973, 1975)</u>	<u>1, 2,</u> <u>3</u>	<u>1, 2.2.1,</u> <u>2.2.2, 2.4</u>

<u>treatment A</u>	<u>without addition of inhibitor, but amended with glucose</u>		
<u>treatment B</u>	<u>with addition of inhibitor for bacterial growth (streptomycin) and glucose</u>		
<u>treatment C</u>	<u>with addition of inhibitor for fungal growth (cycloheximide) and glucose</u>		
<u>treatment B</u>	<u>with addition of bot inhibitors (streptomycin, cycloheximide) and glucose</u>		
<u>f_{Dmi}</u>	<u>fungal contribution to N_2O production during denitrification with microbial inhibition</u>	<u>3</u>	<u>Table 5</u>
<u>Variety traced</u>	<u>^{15}N tracer technique was used to estimate the effect of N_2O reduction on N_2O produced</u>	<u>/</u>	<u>1; 2.2.2; Figure 1</u>
<u>Variety +C_2H_2</u>	<u>Natural isotopic conditions and C_2H_2 addition to the headspace (10 kPa) to block N_2O reduction</u>		
<u>Variety -C_2H_2</u>	<u>Natural isotopic conditions and no C_2H_2 addition to the headspace</u>		
<u>WFPS</u>	<u>Water filled pore space</u>	<u>/</u>	<u>2.2</u>
<u>GC</u>	<u>Gas chromatography</u>	<u>/</u>	<u>2.3</u>
<u>$c(N_2O)$, $c(CO_2)$</u>	<u>N_2O and CO_2 concentrations analysed by GC</u>	<u>/</u>	<u>2.3, Figure 1</u>
<u>IRMS</u>	<u>Isotope ratio mass spectrometry</u>	<u>/</u>	<u>2.5</u>
<u>IEM</u>	<u>the isotope endmember mixing approach proposed by Ostrom et al. (2010)</u>	<u>/</u>	<u>1, 2.5.1</u>
<u>SP_{prod}</u>	<u>SP_{N_2O} values of N_2O produced in soil</u>	<u>4</u>	<u>1, 2.5.1</u>
<u>f_{FD}</u>	<u>Fraction of fungi contributing to N_2O production during denitrification</u>	<u>4</u>	<u>2.5.1</u>
<u>f_{BD}</u>	<u>Fraction of bacteria contributing to N_2O production during denitrification</u>	<u>4</u>	<u>2.5.1</u>
<u>SP_{FD}</u>	<u>SP_{N_2O} values produced by fungi contributing to N_2O production during denitrification</u>	<u>4</u>	<u>2.5.1</u>
<u>SP_{BD}</u>	<u>SP_{N_2O} values produced by bacteria contributing to N_2O production during denitrification</u>	<u>4</u>	<u>2.5.1</u>
<u>f_{FD_SP}</u>	<u>SP_{N_2O} values produced by fungi calculated with IEM using results of variety +C_2H_2, assuming SP_{N_2O} values of N_2O produced by bacteria were 3.7 ‰ (resulting in negative fraction and therefore set to zero) or -7.5 ‰. Using the minimum and maximum SP_{N_2O} values known for bacteria resulted in a f_{FD_SP} range.</u>	<u>4</u>	<u>2.5.1, Table 5</u>
<u>f_{FD_SPpot}</u>	<u>Maximum potential fungal fraction of N_2O production calculated by with IEM for all treatments of variety -C_2H_2 assuming SP_{N_2O} values of N_2O produced by bacterial denitrification or nitrifier denitrification were between 3.7 and -10.7 ‰ (Frame and Casciotti, 2010; Yu et al., 2020) or produced by fungal denitrification or nitrification were between 16 and 37 ‰ (Sutka et al., 2008; Decock and Six, 2013; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017). Here, the effect of potential partial reduction of N_2O could not be included.</u>	<u>4</u>	<u>2.5.1, Table 5</u>
<u>SP/$\delta^{18}O$ Map</u>	<u>isotope mapping approach was further developed (SP/$\delta^{18}O$ Map) using $\delta^{18}O_{N_2O}$ and SP_{N_2O} values of N_2O and $\delta^{18}O$ values of precursors (Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2020)</u>	<u>/</u>	<u>1, 2.5.2</u>
<u>$F_{FD}f_{FD_MAP}$</u>	<u>f_{FD} contributing to N_2O production from denitrification in soil samples estimated with the SP/$\delta^{18}O$ Map</u>	<u>/</u>	<u>2.5.2, Table 4, Table 5</u>
<u>r_{MAP}</u>	<u>N_2O product ratio [$N_2O/(N_2+N_2O)$] estimated with the SP/$\delta^{18}O$ Map</u>	<u>/</u>	<u>2.5.2</u>
<u>r_{15N}</u>	<u>N_2O product ratio [$N_2O/(N_2+N_2O)$] derived from variety traced</u>	<u>5</u>	<u>2.5.3</u>
<u>$^{15}N_{N_2O}$, $^{15}N_{N_2}$</u>	<u>^{15}N-labeling of N_2O or N_2 produced</u>	<u>5</u>	<u>2.5.3</u>
<u>$r_{C_2H_2}$</u>	<u>N_2O product ratio [$N_2O/(N_2+N_2O)$] calculated from N_2O production rates of varieties -C_2H_2 and +C_2H_2</u>	<u>6</u>	<u>2.5.3</u>

$\frac{N_2O_{-C_2H_2}}{N_2O_{+C_2H_2}}$	N_2O produced in varieties $-C_2H_2$ and $+C_2H_2$, respectively	6	2.5.3
SP_{N_2O-r}	^{15}N site preference values of produced N_2O , i.e. without its reduction to N_2O (SP_{prod}), of variety $-C_2H_2$	7	2.5.3
η_r	Net isotope effect of N_2O reduction	7	2.5.3
$\delta\theta$	isotopic values of N_2O produced without N_2O reduction effects of variety $+C_2H_2$	7	2.5.3
f_{FD_SPcalc}	From variety $-C_2H_2$, SP_{N_2O} values of N_2O produced by bacteria was 3.7 (resulting in negative fraction and therefore set to zero) or -7.5 ‰ and using reduction correction with $\eta_r = -6$ ‰ to calculate SP_{prod} values (Senbayram et al., 2018; Yu et al., 2020). Using the minimum and maximum SP_{N_2O} values known for bacteria resulted in a f_{FD_SP} range.	7	2.5.3, Table 5
a_p	calculate the fraction of N_2 and N_2O originating from the ^{15}N -labelled N pool as well as the ^{15}N enrichment of that N pool	7	4.4
$^{15}N_{N_2O_exp}$	expected ^{15}N enrichment in N_2O produced assuming that denitrification is the only process producing N_2O in the incubation experiment	8	2.6
$N_{soil}, N_{fert}, N^{bulk}$	amount of N [mg] in unfertilized soil samples	8	2.6
$^{15}N_{nat}, ^{15}N_{fert}$	^{15}N enrichment under natural conditions (0.3663 at%) and in fertilizer (50 at%), respectively	8	2.6

20 **Determining optimal concentrations for SIR and SIRIN**

As described in the Material and Methods section, optimal concentrations of glucose or inhibitors streptomycin and cycloheximide were determined by SIR or SIRIN method using an automated incubation system (using an "Ultragas 3" CO_2 analyser (WösthoffCo., Bochum) with continuous gas flow) and analysed with the software "SIR-SBA 4.00" (Heinemeyer, copyright MasCo Analytik, Hildesheim, Germany) (Anderson and Domsch, 1973, 1975, 1978). This program enabled to analyse respiration curves for biomass and F:B ratio in soil. However, as data were generated by this software of the incubation system raw data could not be exported and it is thus not possible to represent all tested concentrations and replicates for one soil in one figure. Therefore, results for one representative replicate with glucose concentrations between 0.75 and 2 $mg\ g^{-1}$ soil as an example is presented for Soils 1 to 3 (Figure S1). Additionally, one representative respiration curve of pre-experiments using the SIRIN approach is represented as an example for each with optimum concentrations of streptomycin and cycloheximide (Figure S2).





Figure 1: Respiration curves of pre-experiments derived from data analysis using the computer program “SIR-SBA 4.00” (Heinemeyer, copyright MasCo Analytik, Hildesheim, Germany) for Soils 1 -3. Here results for experiments with glucose concentrations between 0.5 and 2 mg g⁻¹ are presented as examples for one replicate each.



40

Figure S2: Respiration curves of the pre-experiment for SIRIN approach derived from data analysis using the computer program "SIR-SBA 4.00" (Heinemeyer, copyright MasCo Analytik, Hildesheim, Germany) with optimum inhibitor concentrations. The examples represent treatment A without growth inhibition, treatment B with 1.0 mg g⁻¹ dw soil

streptomycin, treatment C with 0.75 mg g⁻¹ dw soil cycloheximide and D with both inhibitors for experiments with *Soil I-3*. Results show curves as an example for one replicate each.

45

Table S4S2: SP values of produced N₂O, i.e. without its reduction to N₂, of variety -C₂H₂ (SP_{prod}) calculated by the Rayleigh-type model according to Lewicka-Szczebak et al. (2017) and Senbayram et al. (2018) (Eq. 7) using the isotope effect of N₂O reduction from the literature (-6‰) (Yu et al., 2020) and the *product-ratio*_{15N}.

Experiment	Treatment/variety	SP _{prod}
<i>Experiment-Soil I.1</i> (Loamy sand, winter 2012)	A / -C ₂ H ₂	0.91
	B / -C ₂ H ₂	0.37
	C / -C ₂ H ₂	1.06
	D / -C ₂ H ₂	-0.03
<i>Soil I.2</i> (Loamy sand, summer 2011)	A / -C ₂ H ₂	2.71
	B / -C ₂ H ₂	-1.80
	C / -C ₂ H ₂	2.40
	D / -C ₂ H ₂	-0.71
<i>Soil Experiment 2</i> (Sand, winter 2012)	A / -C ₂ H ₂	-1.00
	B / -C ₂ H ₂	-1.64
	C / -C ₂ H ₂	-1.40
	D / -C ₂ H ₂	-1.03
<i>Soil Experiment 3</i> (Silt loam, winter 2013)	A / -C ₂ H ₂	0.02
	B / -C ₂ H ₂	-0.62
	C / -C ₂ H ₂	-0.89
	D / -C ₂ H ₂	-1.43
<i>Experiment 4</i> (Loamy sand, summer 2011)	A / -C ₂ H ₂	2.71
	B / -C ₂ H ₂	-1.80
	C / -C ₂ H ₂	2.40
	D / -C ₂ H ₂	-0.71

50 **References:**

Anderson, J. P. E., and Domsch, K. H.: Quantification of bacterial and fungal contributions to soil respiration, *Archiv Fur Mikrobiologie*, 93, 113-127, doi:10.1007/BF00424942, 1973.

55 Anderson, J. P. E., and Domsch, K. H.: Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soil, *Canadian Journal of Microbiology*, 21, 314-322, doi: 10.1139/m75-045, 1975.

Anderson, J. P. E., and Domsch, K. H.: Physiological method for quantitative measurement of microbial biomass in soils, *Soil Biol. Biochem.*, 10, 215-221, doi:10.1016/0038-0717(78)90099-8, 1978.

60 Lewicka-Szczebak, D., Augustin, J., Gieseemann, A., and Well, R.: Quantifying N₂O reduction to N₂ based on N₂O isotopocules – validation with independent methods (helium incubation and ¹⁵N gas flux method), *Biogeosciences*, 14, 711-732, doi: 10.5194/bg-14-711-2017, 2017.

65 Lewicka-Szczebak, D., Lewicki, M. P., and Well, R.: N₂O isotope approaches for source partitioning of N₂O production and estimation of N₂O reduction – validation with the ¹⁵N gas-flux method in laboratory and field studies, *Biogeosciences*, 17, 5513-5537, 10.5194/bg-17-5513-2020, 2020.

70 Ostrom, N. E., Sutka, R., Ostrom, P. H., Grandy, A. S., Huizinga, K. M., Gandhi, H., von Fischer, J. C., and Robertson, G. P.: Isotopologue data reveal bacterial denitrification as the primary source of N₂O during a high flux event following cultivation of a native temperate grassland, *Soil Biol. Biochem.*, 42, 499-506, doi: 10.1016/j.soilbio.2009.12.003, 2010.

75 Senbayram, M., Well, R., Bol, R., Chadwick, D. R., Jones, D. L., and Wu, D.: Interaction of straw amendment and soil NO₃⁻ content controls fungal denitrification and denitrification product stoichiometry in a sandy soil, *Soil Biol. Biochem.*, 126, 204-212, doi: 10.1016/j.soilbio.2018.09.005, 2018.

Toyoda, S., and Yoshida, N.: Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer, *Anal. Chem.*, 71, 4711-4718, 10.1021/ac9904563, 1999.

80 Yu, L., Harris, E., Lewicka-Szczebak, D., Barthel, M., Blomberg, M. R. A., Harris, S. J., Johnson, M. S., Lehmann, M. F., Liisberg, J., Müller, C., Ostrom, N. E., Six, J., Toyoda, S., Yoshida, N., and Mohn, J.: What can we learn from N₂O isotope data? - Analytics, processes and modelling, *Rapid Commun Mass Spectrom*, doi: 10.1002/rcm.8858, 2020.

85