

Comparing modified substrate induced respiration with selective inhibition (SIRIN) and N₂O isotope approaches to estimate fungal contribution to denitrification in three arable soils under anoxic conditions

5 Lena Rohe^{1,2,3}, Traute-Heidi Anderson², Heinz Flessa², Anette Goeske², Dominika Lewicka-Szczebak^{2,4}, Nicole Wrage-Mönnig⁵, Reinhard Well²

¹Helmholtz Centre for Environmental Research – UFZ, Department Soil System Sciences, Theodor-Lieser Str. 4, Halle, Germany

²Thünen Institute of Climate Smart Agriculture, Bundesallee 65, Braunschweig, Germany

10 ³University of Göttingen, Department of Crop Sciences, Institute of Grassland Science, von-Siebold-Str. 8, 37075 Göttingen, Germany

⁴Institute of Geological Sciences, University of Wrocław, pl. M. Borna 9, 50-204 Wrocław, Poland

⁵University of Rostock, Agricultural and Environmental Faculty, Grassland and Fodder Sciences, Justus-Liebig-Weg 6, Rostock, Germany

15 *Correspondence to:* Lena Rohe (lena.rohe@ufz.de)

Keywords: selective growth inhibition, ¹⁵N site preference, fungal denitrification, C₂H₂, isotope endmember mixing approach, *SP_{N2O}* mixing balance, SP/ $\delta^{18}\text{O}$ mapping approach

Abstract

20 The coexistence of many N₂O production pathways in soil hampers differentiation of microbial pathways. The question whether fungi are significant contributors to soil emissions of the greenhouse gas nitrous oxide (N₂O) from denitrification has not yet been resolved. Here, as far as we know, three approaches to independently investigate the fungal fraction contributing to N₂O from denitrification were used simultaneously for the first time (modified substrate induced respiration with selective inhibition (SIRIN) approach, and two isotopic approaches, 25 i.e. endmember mixing approach (IEM) using the ¹⁵N site preference of N₂O produced (*SP_{N2O}*), and the SP/ $\delta^{18}\text{O}$ mapping approach (SP/ $\delta^{18}\text{O}$ Map)). This enabled a comparison of methods and a quantification of the importance of fungal denitrification in soil.

30 Three soils were incubated in four treatments of the SIRIN approach under anaerobic conditions to promote denitrification. While one treatment without microbial inhibition served as a control, the other three treatments were amended with inhibitors to selectively inhibit bacterial, fungal or bacterial and fungal growth. These treatments were performed in three varieties. In one variety, the ¹⁵N tracer technique was used to estimate the effect of N₂O reduction on N₂O produced, while two other varieties were performed under natural isotopic conditions with and without acetylene.

35 All three approaches revealed a small fungal contribution to N₂O fluxes (f_{FD}) under anaerobic conditions in the soils tested. Quantifying the fungal fraction with modified SIRIN was not successful due to large amounts of uninhibited N₂O production. In only one soil, f_{FD} could be estimated using modified SIRIN and resulted in 28±9 %, which was possibly overestimated as results obtained by IEM and SP/ $\delta^{18}\text{O}$ Map for this soil resulted in f_{FD} of below 15 and 20 %, respectively. As a consequence of the unsuccessful SIRIN approach, estimation of fungal *SP_{N2O}* values was impossible.

40 While all successful methods coincided by suggesting a small or missing fungal contribution, further studies with
stimulated fungal N₂O fluxes by added fungal C substrates and an improved modified SIRIN approach, including
alternative inhibitors, are needed to better cross-validate the methods.

1. Introduction

45 The greenhouse gas nitrous oxide (N₂O) contributes to global warming and to the depletion of the ozone layer in
the stratosphere (Crutzen, 1970; IPCC, 2013). The largest anthropogenic N₂O emissions originate from agricultural
soils and are mainly produced during microbial nitrification, nitrifier denitrification and denitrification (Firestone
and Davidson, 1989; Bremner, 1997; IPCC, 2013; Wrage-Mönnig et al., 2018). In order to find mitigation
strategies for N₂O emissions from arable soils, it is important to understand N₂O sources and sinks and thus
improve knowledge about the production pathways and the microorganisms involved.

50 Denitrification describes the stepwise reduction of nitrate (NO₃⁻) to dinitrogen (N₂), with the intermediates nitrite
(NO₂⁻), nitric oxide (NO) and N₂O (Knowles, 1982). For a long time, it was believed that solely bacteria are
involved in N₂O formation during denitrification (Firestone and Davidson, 1989); however, also several fungi are
capable of denitrification (Bollag and Tung, 1972; Shoun et al., 1992). Pure culture studies indicated that although
only some fungal species (e.g. *Fusarium* strains) are performing respiratory denitrification, these may produce
55 substantial amounts of N₂O (Higgins et al., 2018; Keuschnig et al., 2020). N₂O produced by fungi may thus
contribute largely to N₂O from denitrification in soil, since, firstly, fungi dominate the biomass in soil (up to 96
%) compared to bacteria in general (Ruzicka et al., 2000; Braker and Conrad, 2011). A respiratory fungal-to-
bacterial (F:B) ratio of 4 is typical for arable soils (Anderson and Domsch, 1975; Blagodatskaya and Anderson,
1998). Secondly, due to a lacking N₂O reductase (Nos) (Shoun et al., 1992; Shoun et al., 2012; Higgins et al.,
60 2018), N₂O is the major end product of fungal denitrification. However, although there are methodological
approaches to disentangle sources of N₂O, it is still challenging to clearly attribute N₂O emitted from soil to
bacterial or fungal denitrification.

One approach to differentiate between N₂O produced by fungi and bacteria during denitrification comprises the
addition of two antibiotics to soil incubation experiments, i.e. streptomycin and cycloheximide to inhibit bacterial
65 or fungal protein biosynthesis, i.e. growth, respectively. This method is known as substrate induced respiration
with selective inhibition (SIRIN) and was originally developed to determine the bacterial or fungal contribution to
CO₂ respiration (Anderson and Domsch, 1975). A few studies used a modification of this method for N₂O analysis
(Laughlin and Stevens, 2002; Crenshaw et al., 2008; Blagodatskaya et al., 2010; Long et al., 2013) and found a
greater decrease of N₂O production with fungal than with bacterial growth inhibition (i.e. 89 vs. 23 % decrease,
70 respectively (Laughlin and Stevens, 2002)). This indicated that fungi might dominate N₂O production (Laughlin
and Stevens, 2002; McLain and Martens, 2006; Crenshaw et al., 2008; Blagodatskaya et al., 2010; Long et al.,
2013; Chen et al., 2014; Chen et al., 2015). However, difficulties of this method may be to achieve complete
inhibition of selective groups (Ladan and Jacinthe, 2016) and to avoid shifts in the structure of microbial
communities as response of pre-incubation or duration of experiments. Another opportunity to distinguish between
75 N₂O from bacterial and fungal denitrification and other pathways is the analysis of the isotopic composition of
N₂O. Especially the isotopomer ratios of N₂O (i.e. N₂O molecules with the same bulk ¹⁵N isotopic enrichment but
different positions of ¹⁵N in the linear N₂O molecule (Ostrom and Ostrom, 2017)) in pure culture studies showed
differences in N₂O of bacterial and fungal denitrification (Sutka et al., 2006; Sutka et al., 2008; Frame and
Casciotti, 2010; Rohe et al., 2014a; Rohe et al., 2017). Isotopomer ratios of N₂O can be expressed as ¹⁵N site

80 preference (SP_{N_2O}), i.e. the difference between $\delta^{15}N$ of the central and terminal N-position of the asymmetric N_2O molecule (Toyoda and Yoshida, 1999). The SP_{N_2O} values of N_2O of six pure fungal cultures was between 16 and 37 ‰ (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017), whereas several bacterial cultures produced N_2O with SP_{N_2O} values between -7.5 and +3.5 ‰ during denitrification (Toyoda et al., 2005; Sutka et al., 2006; Rohe et al., 2017). While it is generally assumed that SP_{N_2O} values of N_2O produced by fungal
85 pure cultures during denitrification are transferable to N_2O produced by fungal soil communities, this has not yet been proven. Until now, studies reporting possible ranges of fungal contributions to N_2O fluxes from soil were based on SP_{N_2O} values of pure cultures (Köster et al., 2013b; Zou et al., 2014; Lewicka-Szczebak et al., 2017; Senbayram et al., 2018; Senbayram et al., 2020; Lewicka-Szczebak et al., 2014), but uncertainty of this approach arose from the large ranges of fungal SP_{N_2O} values (Sutka et al., 2008; Maeda et al., 2015; Rohe et al., 2017). It
90 would thus be useful to constrain fungal SP_{N_2O} values for a specific soil or soil type.

The SP_{N_2O} value of N_2O produced by pure bacterial cultures during nitrification is approximately 33 ‰ and thus interferes with that of fungal denitrification (Sutka et al., 2006; Sutka et al., 2008; Rohe et al., 2014a). This demonstrates the difficulty to use solely SP_{N_2O} values as an indicator for different organism groups contributing to
95 N_2O production from soil, where different pathways may co-occur. Based on the above cited ranges for the isotopomer endmembers of fungal and bacterial denitrification, and assuming that only fungal and bacterial denitrification are responsible for N_2O production, the fraction of fungal N_2O can be calculated using the isotope endmember mixing approach (IEM) with SP_{N_2O} values of N_2O produced in soil (SP_{prod}), provided N_2O reduction does not occur (Ostrom et al., 2010; Ostrom and Ostrom, 2011). If there is a N_2O reduction, SP_{N_2O} and also $\delta^{15}N$ and $\delta^{18}O$ values of produced N_2O ($\delta^{15}N^{bulk}_{N_2O}$ and $\delta^{18}O_{N_2O}$, respectively) are affected by isotopic fractionation
100 (Ostrom et al., 2007; Ostrom and Ostrom, 2011). This means that the $^{14}N^{16}O$ bond of N_2O is preferentially broken compared to $^{14}N^{18}O$ or $^{15}N^{16}O$, resulting in N_2O that is isotopically enriched in ^{15}N and ^{18}O and shows larger SP_{N_2O} values compared to N_2O from denitrification without the reduction step (Popp et al., 2002; Ostrom et al., 2007).

In controlled laboratory experiments, the N_2O reduction to N_2 can be inhibited using acetylene (C_2H_2) during anaerobic incubation experiments (Yoshinari and Knowles, 1976; Groffman et al., 2006; Well and Flessa, 2009; Nadeem et al., 2013). Hence, C_2H_2 inhibition might be suitable to quantify SP_{prod} values in soils exhibiting
105 significant N_2O reduction and would thus allow quantification of fungal N_2O fluxes based on SP_{prod} values. However, problems due to incomplete inhibition of N_2O reduction and unwanted inhibition of other pathways may occur (Wrage et al., 2004b; Wrage et al., 2004a). Another possibility to quantify N_2O reduction to N_2 during denitrification is also possible with ^{15}N tracing experiments using ^{15}N enriched substrates and analysing $^{15}N_2$ fluxes
110 (Well et al., 2006; Lewicka-Szczebak et al., 2014). The ^{15}N tracer approach also enables to distinguish between N_2O from fungal denitrification and co-denitrification, i.e. a hybrid N_2O is formed using one N atom from NO_2^- and one N atom from compounds like azide or ammonium (NH_4^+) for N_2O production (Tanimoto et al., 1992; Laughlin and Stevens, 2002; Rohe et al., 2017).

N_2O reduction can be quantified using N_2O natural abundance isotopic signatures, which also enables simultaneous
115 differentiation of selected pathways producing N_2O . Here, the isotope mapping approach uses isotope fractionation factors together with $\delta^{15}N$ values of precursors ($\delta^{15}N_{NO_x}$) as well as $\delta^{15}N^{bulk}_{N_2O}$ and SP_{N_2O} values of N_2O produced (Toyoda et al., 2011). Recently, this isotope mapping approach was further developed (SP/ $\delta^{18}O$ Map) using $\delta^{18}O_{N_2O}$ and SP_{N_2O} values and $\delta^{18}O$ values of precursors (Lewicka-Szczebak et al., 2017) and different slopes of N_2O reduction and mixing lines in the $\delta^{18}O$ – SP isotope plot. While SP_{N_2O} values are independent of isotopic signatures
120 of the precursors, $\delta^{15}N^{bulk}_{N_2O}$ and $\delta^{18}O_{N_2O}$ result from the isotopic signature of the precursor and isotopic

fractionation during N₂O production (Toyoda et al., 2005; Frame and Casciotti, 2010). Regarding δ¹⁸O_{N₂O}, a complete exchange of oxygen (O) between NO₃⁻ and soil water can be assumed and consequently, one can use the δ¹⁸O values of soil water for interpretation of δ¹⁸O_{N₂O} values (Kool et al., 2009; Snider et al., 2009; Lewicka-Szczebak et al., 2016). However, interpretation of δ¹⁸O_{N₂O} values from different microbial groups may be more complex due to incomplete O exchange, because variations in the extent of O exchange between water and N oxides affect the final δ¹⁸O_{N₂O} value (Garber and Hollocher, 1982; Aerssens et al., 1986; Kool et al., 2007; Rohe et al., 2014b; Rohe et al., 2017). Importantly, fungal and bacterial N₂O showed different ranges for δ¹⁸O_{N₂O} values, hence this isotopic signature may also be helpful in differentiation of these pathways (Lewicka-Szczebak et al., 2016). This SP/δ¹⁸O Map approach thus allows for an estimation of the contributions of N₂O reduction and admixture of fungal N₂O.

So far, the described methods for distinguishing between fungal and bacterial N₂O emission have not been compared in the same soil and their accuracy and possible bias remains unknown. A better knowledge of the comparability of the methods would enable comparison of results of studies using different methods and thus further improve our understanding of processes of N₂O production. It would also reveal weaknesses of approaches and might lead to the development of better methods.

Therefore, this study aims at (i) determining the fungal contribution to N₂O production by denitrification under anoxic conditions and glucose addition using three arable soils and three approaches (modified SIRIN, IEM and the SP/δ¹⁸O Map), and to assess their usefulness in soil studies and thus assess factors of potential bias of the methods and (ii) to estimate the SP_{N₂O} values from fungal soil communities and thus to evaluate the transferability of the pure culture range of the fungal SP_{N₂O} endmember values. We hypothesized that the fungal fraction contributing to N₂O from denitrification in different soils using a modified SIRIN approach and isotopic methods will be correlated but not exactly matched due to limited inhabitability of microbial communities and variability in SP_{N₂O} endmember values. Furthermore, successful application of the modified SIRIN approach with determined fungal fraction contributing to N₂O from denitrification will yield fungal SP_{N₂O} endmember values within the range of values previously reported in the literature.

2. Materials and Methods

2.1 Soil samples


All experiments were conducted with three arable soils differing in texture, C_{org} content, C/N ratio and pH. Thus it was assumed that the soils harbour different denitrifying communities, i.e. different fractions of bacteria and fungi contributing to denitrification. One of the soils was sampled during a second season to evaluate if the fungal fraction contributing to N₂O production is soil-specific or can be subject to seasonal change of microbial communities. As this soil was sampled at two different time points, we conducted four experiments and named the different experiments “Soil 1.1”, “Soil 1.2”, “Soil 2”, and “Soil 3”: *Soil 1.1 and Soil 1.2* with loamy sand (*Soil 1*) sampled in December 2012 and in June 2011, respectively, *Soil 2* with sand sampled in January 2013, and *Soil 3* with silt loam sampled in December 2012 (Table 1).

Soil samples of the upper 30 cm were collected in plastic bags aerated via cotton wool stoppers and stored at 6 °C for maximally two months. To get information about the initial soil status, the mineral nitrogen content (N_{min}) of soil samples was determined before and after fertilization by extracting NO₃⁻ and NH₄⁺ with 0.01 M calcium chloride dihydrate (CaCl₂ · 2 H₂O) according to ISO 14255 and analysing NO₃⁻ and NH₄⁺ concentrations in the

160 extracts with a Continuous-Flow-Analyser (SKALAR, Germany) directly after sample collection. Other soil characteristics (C and N content, soil pH value, isotopic values of soil NO_3^- and NO_2^-) were analysed with samples of *Soil 1.1*, *Soil 2* and *Soil 3*. Total contents of C and N in soil samples were analysed by dry combustion of ground samples (LECO TruSpec, Germany). The soil pH was measured in 0.01 M CaCl_2 . The $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values of NO_3^- and NO_2^- ($\delta^{15}\text{N}_{\text{NO}_x}$ and $\delta^{18}\text{O}_{\text{NO}_x}$, respectively) in soil extracts (with 0.01 M calcium chloride dihydrate ($\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$)) were analysed by the bacterial denitrifier method (Casciotti et al., 2002) (Table 1).
 165 The three soils were also sampled in summer 2010 for pre-experiments to gain information on the respiratory biomass by analyzing the substrate induced respiration (SIR) according to Anderson and Domsch (1978) and the respiratory F:B ratio was analysed with substrate induced respiration with selective inhibition (SIRIN) by a computer-generated selectivity analysis: “SIR-SBA 4.00” (Heinemeyer, copyright MasCo Analytik, Hildesheim, Germany) (Anderson and Domsch, 1975) (Table 1). The scheme of glucose and growth inhibitor combinations is listed below in section “Methodical approach”. The characteristics of the soils are listed in Table 1.

2.2 Methodical approach

The experimental setup with various measures is presented in the following sections and illustrated in Figure 1. Important terms used and its descriptions are listed in Supplementary Material, Table S1.



175
 180
 185
 Figure 1: The methodical approach comprised a pre-experiment with substrate induced respiration (SIR) to estimate the optimal glucose concentration ($c_{\text{opt}}(\text{glucose})$) and the fungal-to-bacterial ration in the soil (F:B ratio), and the substrate induced respiration with selective inhibition approach (SIRIN) to determine the optimal inhibitor concentration ($c_{\text{opt}}(\text{streptomycin})$ and $c_{\text{opt}}(\text{cycloheximide})$). The initial soil status, i.e. ammonium and nitrate concentration of the soil ($c(\text{NH}_4^+)$ and $c(\text{NO}_3^-)$, respectively), was measured in N_{min} extracts and the isotopic signature of soil NO_3^- was analysed by the denitrifier method. The incubation experiment comprised the SIRIN approach with three experimental varieties: without acetylene ($-\text{C}_2\text{H}_2$), with C_2H_2 ($+\text{C}_2\text{H}_2$), and without C_2H_2 but with ^{15}N labelled NO_3^- (traced), while NO_3^- with natural isotopic composition was added to the other two varieties. Produced gas was analysed for its concentration ($c(\text{CO}_2)$ and $c(\text{N}_2\text{O})$) using gas chromatography (GC) and N_2O was further analysed by isotope ratio mass spectrometry (IRMS) for its isotopic composition. Please refer to the Material & Methods section for more information.

2.2.1 SIRIN pre-experiment

As in most studies applying the SIRIN method on N_2O emissions (e. g. Laughlin and Stevens, 2002; Chen et al., 2014; Ladan and Jacinthe, 2016), a pre-experiment was conducted with samples collected in 2010, in order to get
 190 information about optimal substrate and inhibitor concentrations for substrate induced respiration with growth

inhibition. The pre-experiments of the present study were conducted in two steps as described in the original methods, i.e. CO₂ production under oxic conditions was analysed to estimate the substrate induced respiration by the SIR method (Anderson and Domsch, 1978) and the substrate induced respiration with selective inhibition by the SIRIN method (Anderson and Domsch, 1975) as follows.

195 In a first pre-experiment (Figure 1), the SIR method (Anderson and Domsch, 1978) was used to get information about the amount of respiratory biomass in soil under oxic conditions. In this pre-experiment glucose served as substrate to initiate microbial growth (Anderson and Domsch, 1975). To this end, we added different concentrations of glucose (0.50, 0.75, 1.0, 1.5, 2.0, 3.0, 4.0, 5.0, 6.0 mg g⁻¹ dry weight (dw) soil) to find the optimal glucose concentration ($c_{opt}(\text{glucose})$), which is the glucose concentration that causes maximum initial respiration rates by analysing CO₂ production (Anderson and Domsch, 1978). $c_{opt}(\text{glucose})$ was 1.0 mg g⁻¹ for *Soil 2* (sand) and 1.5 mg g⁻¹ for *Soils 1, 3* and *4* (loamy sand and silt loam).

200 In a second pre-experiment (Figure 1), the SIRIN method was used according to Anderson and Domsch (1975) for determining the respiratory F:B ratio. The $c_{opt}(\text{glucose})$ determined in the first pre-experiment was used, while selectivity of the inhibitor combinations of streptomycin (bacterial respiratory inhibitor) and cycloheximide (fungal respiratory inhibitor) were tested with three concentrations (0.75, 1.0, 1.5 mg g⁻¹ dw, respectively). The optimal concentration for inhibition of fungal respiration was 0.75 mg g⁻¹ dw soil cycloheximide ($c_{opt}(\text{cycloheximide})$) and for bacterial respiratory inhibition 1.0 mg g⁻¹ dw soil streptomycin ($c_{opt}(\text{streptomycin})$). As in the first pre-experiment, CO₂ production under oxic conditions was analysed. The determined optimal concentrations of glucose, streptomycin and cycloheximide were used in the modified SIRIN approach, on the assumption that concentrations optimal for CO₂ respiration also allow denitrification. Examples of respiration curves derived from SIR and SIRIN pre-experiments are represented in Figure S1 and S2, respectively.

205

210

Table 1: Soil characteristics of three arable soils from Germany used for incubation experiments (*Soil*) (standard deviation in brackets). Except for NH_4^+ and NO_3^- , soil characteristics (C, N, pH, $\delta^{15}\text{N}_{\text{NO}_x}$ and $\delta^{18}\text{O}_{\text{NO}_x}$) of loamy sand were only analysed once for samples collected in 2012.

<i>Soil</i> (Year)	<i>Soil</i> texture	<i>Soil</i> type (WRB)	Location	C content [%]	N content [%]	NH_4^+ [mg N kg ⁻¹]	NO_3^- [mg N kg ⁻¹]	pH (CaCl ₂)	$\delta^{15}\text{N}_{\text{NO}_x}$ [‰] ^e	$\delta^{18}\text{O}_{\text{NO}_x}$ [‰] ^e	F:B ^f	Biomass ^g [µg C gdw ⁻¹ soil]
1.1 (2012)	Loamy sand	Haplic Luvisol	Braun- schweig ^a	1.43 (<0.01)	0.10 (<0.01)	0.4 (<0.1)	14.1 (2.1)	5.67	3.98	-4.82	2.6	234
1.2 (2011)				/	/	1.0 (0.4)	11.0 (0.3)	/	/	/	/	/
2 (2013)	Sand	Gleyic Podzol	Wenne- bostel ^b	2.31 (0.04)	0.14 (<0.01)	1.9 (0.2)	6.6 (0.2)	5.54	0.73	-2.68	2.6	161
3 (2013)	Silt loam	Haplic Luvisol	Götting- en ^c	1.62 (0.02)	0.13 (<0.01)	n.d. ^d	22.7 (<0.1)	7.38	4.18	2.32	4.9	389

215 ^aExperimental Station of the Friedrich-Löffler Institute, Braunschweig, Germany

^bprivate agricultural field North of Hannover, water protection area Fuhrberger Feld, Germany

^cReinshof Experimental Farm, Georg-August-University, Göttingen, Germany

^dnot detectable (i.e. below detection limit of 0.06 mg kg⁻¹ NH₄⁺-N)

^eIsotopic values of natural soil NO₃⁻ using the denitrifier method (Casciotti et al., 2002).

220 ^fRespiratory fungal-to-bacterial (F:B) ratio analysed by SIRIN method (Anderson and Domsch, 1973, 1975) in a pre-experiment in 2010.

^gRespiratory biomass analysed by CO₂ production from SIR method (Anderson and Domsch, 1978) in a pre-experiment in 2010..

2.2.2 Soil incubation with selective inhibition to determine N₂O forming processes

225 The experimental design included two approaches, (i.) microbial inhibition by fungal and/or bacterial inhibitors and (ii.) activity of N₂O reductase analysed either by inhibition with C₂H₂ or quantification by ¹⁵N tracing (Figure 1). To address the microbial inhibition approach (i.), the SIRIN method for determination of the respiratory F:B ratio based on CO₂ emission was modified to determine N₂O production by microbial groups. However, in contrast to previous studies by Laughlin and Stevens (2002), McLain and Martens (2006), Blagodatskaya et al. (2010) and
230 Long et al. (2013), we did not pre-incubate the soil with the growth inhibitors, as this could result in changes of the microbial community (e.g. preferential growth of selected organisms). We intended to disturb microbial communities as little as possible.

The soil was sieved (2 mm) and pre-incubated at 22 °C for five to seven days in the dark with cotton wool stoppers to allow respiration and aerobic conditions in soil bags. Four microbial inhibitor treatments (each in triplicate)
235 with *c_{opt}*(glucose) for each soil were established:

- A Control, without growth inhibitors
- B With streptomycin sulfate (C₄₂H₈₄N₁₄O₃₆S₃) to inhibit bacterial growth
- C With cycloheximide (C₁₅H₂₃NO₄) to inhibit fungal growth
- D With streptomycin and cycloheximide, to inhibit bacterial and fungal growth

240 To address the other approach (ii.), all microbial inhibitor treatments were conducted in three varieties, i.e.: with ¹⁵N-NO₃⁻ fertilizer (variety “*traced*”) to quantify N₂O reduction to N₂, with natural abundance NO₃⁻ and 10 kPa C₂H₂ in the headspace (variety “+C₂H₂”) to block N₂O reductase, and with natural abundance NO₃⁻ but without blocking N₂O reductase, i.e. no C₂H₂ added (variety “-C₂H₂”) (Figure 1). In total, there were 48 experimental treatments and 144 vessels (four *Soils* with four inhibitor treatments (A, B, C, D) and three varieties (*traced*, +C₂H₂
245 and -C₂H₂), each in triplicate).

The soil was adjusted to 80 % water filled pore space (WFPS) with distilled water. Simultaneously, the soil was fertilized with NO₃⁻ (varieties -C₂H₂, +C₂H₂, and *traced*). The soil sample used with *Soil 1.2* was incubated prior to the other soils and was amended with 60 mg N kg⁻¹ NaNO₃, while in agreement with other experiments conducted in our laboratory, 50 mg N kg⁻¹ KNO₃ were used with *Soil 1.1*, 2 and 3. In variety *traced*, NO₃⁻ with a
250 ¹⁵N enrichment of 50 atom% (at%) was used. For each treatment, we incubated 100 g dw soil in 850 mL preserving jars (J. WECK GmbH u. Co KG, Wehr, Germany) with gas inlet and outlet equipped with three-port luer lock plastic stopcocks (Braun, Melsungen, Germany). According to the original SIRIN method (Anderson and Domsch, 1973, 1978) a mixture of *c_{opt}*(glucose) and carrier material talcum (5 mg talcum g dw⁻¹) was added to soil of treatment A and together with the growth inhibitors to the soil of treatments B, C and D. The soil and additives of
255 each treatment were mixed for 90 seconds with a handheld electric mixer. During packing, the soil density was adjusted to an expected target soil density of 1.6 g cm⁻³ in *Soil 1.1*, 1.2, and 2 and of 1.3 g cm⁻³ in *Soil 3* to imitate field conditions. To ultimately achieve denitrifying conditions in all treatments and to avoid catalytic NO decomposition in the +C₂H₂ variety (Nadeem et al., 2013), the headspace of the closed jars was flushed with N₂ to exchange the headspace 10 times. Directly following, 85 mL of the gas in the headspace in variety +C₂H₂ were
260 exchanged by pure C₂H₂ resulting in 10 kPa C₂H₂ in the headspace. The manual sample collection of 14 mL gas in duplicates with a plastic syringe was performed after six, eight and ten hours (*Soil 1.1*, 2 and 3) or two, four and eight hours (*Soil 1.2*) of incubation time, respectively. The removed gas was replaced by the same amount of N₂.

2.3 Gas analysis

Gas samples were analysed for N₂O and CO₂ concentrations ($c(N_2O)$ and $c(CO_2)$) with gas chromatography (GC, Agilent 7890A, Agilent, Böblingen, Germany) (Figure 1). The analytical precision of measurements was derived from analysing laboratory standards of different concentrations (0.5-1,000 ppm N₂O and 340-10,000 ppm CO₂) and resulted in a measurement precision of 1 % for N₂O and 0.5 % for CO₂. The instrumental detection limit of N₂O was 4 µg N kg⁻¹ h⁻¹ and of CO₂ it was 137 µg C kg⁻¹ h⁻¹. As a control, N₂ and O₂ concentrations in the samples were analysed with GC to ensure anaerobic conditions during the incubation for N₂O production from denitrification. CO₂ and N₂O production rates were calculated by averaging the measured N₂O production, i.e. between the time point of flushing with N₂ (t=0) and six, eight or ten hours (or two, four and eight hours with *Soil I.2*).

The N₂O isotopic analysis of the gas samples of varieties -C₂H₂ and +C₂H₂ (Figure 1) were performed on a pre-concentrator (PreCon, Thermo-Finnigan, Bremen, Germany) interfaced with a GC (Trace Gas Ultra, Thermo Scientific, Bremen, Germany) and analysed by isotope ratio mass spectrometry (IRMS, Delta V, Thermo Fisher Scientific, Bremen, Germany) (Brand, 1995; Toyoda and Yoshida, 1999; Köster et al., 2013b). A laboratory standard N₂O gas was used for calibration, having $\delta^{15}N^{bulk}_{N_2O}$, $\delta^{18}O_{N_2O}$ and SP_{N_2O} values of -1.06 ‰, 40.22 ‰, and -2.13 ‰, respectively, in three concentrations (5, 10 and 20 ppm). The analytical precision was 0.1 ‰, 0.2 ‰ and 1.5 ‰ for $\delta^{15}N^{bulk}_{N_2O}$, $\delta^{18}O_{N_2O}$ and SP_{N_2O} values, respectively. H₂O and CO₂ were trapped with magnesium perchlorate and ascarite, respectively, to prevent any interference with N₂O analysis.

The gas samples of variety *traced* from *Soil I.1*, 2, and 3 were analysed for the 29/28 and 30/28 ratios of N₂ according to Lewicka-Szczepak et al. (2013) using a modified GasBench II preparation system coupled to IRMS (MAT 253, Thermo Scientific, Bremen, Germany). The gas samples of variety *traced* from *Soil I.2* were analysed at the Centre for Stable Isotope Research and Analysis (University of Göttingen, Germany). The N₂ produced was analysed using an elemental analyser (Carlo Erba ANA 1500) that was coupled to dual inlet IRMS (Finnigan MAT 251) (Well et al., 1998; Well et al., 2006). Isotopic values of N₂O of *Soil I.2* (variety *traced*) were analysed in the same lab using a pre-concentration unit coupled to IRMS (Precon-DeltaXP, Thermo Scientific, Bremen, Germany) (Well et al., 2006). Isotope ratios were used applying the non-random distribution approach to calculate the fraction of N₂ and N₂O originating from the ¹⁵N-labelled N pool as well as the ¹⁵N enrichment of that N pool (a_p) (Bergsma et al., 2001; Spott et al., 2006).

2.4 Inhibitor effects

For interpretation of N₂O or CO₂ production, the validity of the experimental results with respect to fungal and bacterial N₂O fluxes was checked using a flux balance comparing the sum of bacterial and fungal inhibition effects (treatments B and C) to the dual inhibition effect (treatment D):

$$D = A - [(A - B) + (A - C)] \quad (\text{Eq. 1})$$

With A , B , C and D representing the N₂O production rates of the last sampling time of treatment A , B , C and D , respectively. Assuming that in the other three treatments (A , B and C) non-inhibitable N₂O production was equal to treatment D , N₂O produced by bacteria and fungi should show the following relation between the four treatments:

$$(A - D) = (B - D) + (C - D) \quad (\text{Eq. 2})$$

The fungal contribution to N₂O production during denitrification with microbial inhibition (f_{FDmi}) can be calculated, when N₂O production of treatment D is significantly smaller than N₂O production of treatments A, B and C by:

$$f_{FDmi} = \frac{(A-C)}{(A-D)} \quad (\text{Eq. 3})$$

305 A detailed discussion of inhibitor effects and difficulties with organisms that were not inhibited or abiotic sources (treatment D) is presented in section 4.1.

2.5 Isotope methods

2.5.1 Isotope endmember mixing approach (IEM)

310 The fungal fraction (f_{FD}) contributing to N₂O production from denitrification in soil samples was calculated according to the isotope mixing model (IEM) proposed by Ostrom et al. (2010), which was established for calculating the bacterial fraction (f_{BD}) of N₂O production. Assuming that bacteria (BD) and fungi (FD) are the only sources of N₂O in soil, the ¹⁵N site preference values of produced N₂O (SP_{prod}) results from the SP_{N_2O} mixing balance:

$$SP_{prod} = f_{FD} * SP_{FD} + f_{BD} * SP_{BD} \quad (\text{Eq. 4})$$

315 where f_{FD} and f_{BD} represent the fraction of N₂O produced by fungi and other N₂O sources than fungal denitrification, respectively, and SP_{FD} and SP_{BD} are the respective SP_{N_2O} endmember values (Ostrom et al., 2010; Ostrom and Ostrom, 2011). This calculation was based on the assumption that the sum of f_{BD} and f_{FD} equals 1 and that N₂O reduction to N₂ is negligible. The mean SP_{FD} value was assumed to be 33.6 ‰ (Sutka et al., 2008; Maeda et al., 2015; Rohe et al., 2014a; Rohe et al., 2017) and the SP_{BD} value from heterotrophic denitrification was assumed with minimum and maximum values from -7.5 to +3.7 ‰ (Yu et al., 2020). For this IEM approach, only results from variety +C₂H₂ could be used to calculate the fungal fraction contributing to N₂O production (f_{FD_SP}), as microorganisms of this variety produce N₂O that is not affected by reduction to N₂. The f_{FD_SP} contributing to N₂O production during denitrification was calculated using the measured SP_{N_2O} value of variety +C₂H₂ as SP_{prod} value in Eq. 4 that was solved for f_{FD} ($f_{FD} = 1 - ((SP_{prod} - SP_{FD}) / (SP_{BD} - SP_{FD}))$). By applying this equation, a range for f_{FD_SP} is received when using minimum and maximum SP_{BD} values.

320 Based on SP_{N_2O} values from -C₂H₂ variety, it was possible to solve Eq. 4 also to estimate the maximum potential fungal contribution to denitrification (f_{FD_SPpot}) assuming that we did not have any estimations for N₂O reduction. While bacterial denitrification and nitrifier denitrification would result in low SP_{N_2O} values ($SP_{BD/ND} = -10.7$ to +3.7 ‰ (Frame and Casciotti, 2010; Yu et al., 2020)), large SP_{N_2O} values would be expected from fungal denitrification and nitrification ($SP_{FB/N} = 16$ to 37 ‰ (Sutka et al., 2008; Decock and Six, 2013; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017)). N₂O reduction could have further increased the SP_{prod} values. If the contribution of this process on SP_{prod} values cannot be precisely estimated, by neglecting these effects we can determine the maximal potential fungal contribution. f_{FD} calculated from Eq. 4 (variety -C₂H₂) would thus be lower if N₂O reduction had occurred. However, assuming the impact of N₂O reduction on SP_{N_2O} was negligible, this IEM enabled to calculate 335 the maximum potential f_{FD} as $f_{FD_SPpot} = 1 - ((SP_{N_2O} - SP_{FD/N}) / (SP_{BD/ND} - SP_{FD/N}))$.

2.5.2 Product ratio [N₂O/(N₂+N₂O)] of denitrification

340 The variety *traced* served to assess N₂O reduction during denitrification in each experiment. The product ratio of denitrification [N₂O/(N₂+N₂O)] as given by the variety *traced* (r_{15N}) was calculated as:

$$r_{15N} = \frac{{}^{15}N_{N_2O}}{{}^{15}N_{N_2} + {}^{15}N_{N_2O}} \quad (\text{Eq. 5})$$

with ${}^{15}N_{N_2O}$ and ${}^{15}N_{N_2}$ representing N₂O and N₂ produced in the ¹⁵N-labeled fertilizer pool. To check the effectiveness of C₂H₂ in blocking the N₂O reduction, r_{15N} was compared with $r_{C_2H_2}$, where the latter can be calculated from N₂O production rates of varieties -C₂H₂ and +C₂H₂:

$$345 \quad r_{C_2H_2} = \frac{N_2O_{-C_2H_2}}{N_2O_{+C_2H_2}} \quad (\text{Eq. 6})$$

with $N_2O_{-C_2H_2}$ and $N_2O_{+C_2H_2}$ representing the N₂O produced in varieties -C₂H₂ and +C₂H₂, respectively.

It was possible to assess the completeness of blockage of N₂O reduction by C₂H₂ with the experimental setup as follows. If r_{15N} and $r_{C_2H_2}$ were in agreement, a complete blockage of N₂O reduction could be assumed. This enabled us to estimate reduction effects on the isotopic signatures of N₂O by comparing the $\delta\theta$ values, i.e. isotopic values of N₂O produced without N₂O reduction effects of variety +C₂H₂, with isotopic values of N₂O of variety -C₂H₂.

350 The information on the product ratio was used as an additional possibility to calculate the f_{FD} also for variety -C₂H₂. The Rayleigh-type model presented by Lewicka-Szczepak et al. (2017) and Senbayram et al. (2018) for similar closed-system incubations was used to calculate the ¹⁵N site preference values of the originally produced N₂O of variety -C₂H₂ (SP_{prod}). SP values of emitted N₂O, i.e. after partial reduction of produced N₂O (SP_{N_2O-r}), were corrected with the net isotope effect of N₂O reduction (ηr) and the r_{15N} as follows:

$$355 \quad SP_{prod} = SP_{N_2O-r} + \eta r \ln(r_{15N}) \quad (\text{Eq. 7})$$

According to Yu et al. (2020) the ηr was assumed to be -6 ‰. Subsequently, Eq. 4 was used to calculate the f_{FD} by using SP_{prod} values of variety -C₂H₂ (f_{FD_SPcalc}) obtained from Eq. 7.

2.5.3 SP/ $\delta^{18}O$ isotope mapping approach (SP/ $\delta^{18}O$ Map)

360 The f_{FD} contributing to N₂O production from denitrification in soil samples was also estimated with the SP/ $\delta^{18}O$ Map (f_{FD_MAP}) (Lewicka-Szczepak et al., 2017; Lewicka-Szczepak et al., 2020). This method allows for estimating both the f_{FD} and N₂O product ratio [N₂O/(N₂+N₂O)] (r_{Map}). For precise estimations, the $\delta^{18}O$ values of soil water ($\delta^{18}O_{H_2O}$) applied in the experiments are needed and these values were not determined. However, since we have independent information on the N₂O product ratio from the *traced* variety (r_{15N}), we can calculate the possible $\delta^{18}O_{H_2O}$ values of soil to get the nearest N₂O product ratios in natural and ¹⁵N treatments. The fitting of $\delta^{18}O_{H_2O}$ values was performed for mean, minimal und maximal values of SP_{BD} (-1.9, -7.5 and 3.7 ‰, respectively) and aimed at obtaining the minimal difference between r_{Map} and that measured in the *traced* variety, i.e. the minimal value of $(r_{15N} - r_{Map})^2$ (according to least squares method) (for explanation of the product ratio see section 2.5.2). This further allows calculation of the possible ranges for f_{FD} for particular $\delta^{18}O_{H_2O}$ fitted values (Table 4) based on $\delta^{18}O_{H_2O}$ values are applied to properly correct the $\delta^{18}O_{N_2O}$ values of the mixing endmembers (*BD* and *FD*), which depend on the ambient water. Afterwards, the corrected values of mixing endmembers are applied to calculate the f_{FD} values. The calculations with this approach may be performed assuming two different scenarios of the interplay between N₂O mixing and reduction (Lewicka-Szczepak et al., 2017; Lewicka-Szczepak et al., 2020), but for this
375 study both scenarios yielded almost identical results (maximal differences of 0.02 in N₂O product ratio and 2 %

for f_{FD} were found), due to f_{BD} near 100 %. Hence, we only provide the results assuming the reduction of bacterial N_2O followed by mixing with fungal N_2O . In the following, all calculated fractions are presented in percent (%).

2.6 Other sources of N_2O

380 Assuming that denitrification was the only source of N_2O in the incubation experiment, the expected ^{15}N enrichment in N_2O produced ($^{15}N_{N_2O_exp}$) was given by

$$^{15}N_{N_2O_exp} [at\%] = \frac{(N_{soil} \times ^{15}N_{nat}) + (N_{fert} \times ^{15}N_{fert})}{N^{bulk}} \quad (\text{Eq. 8})$$

with N_{soil} , N_{fert} and N^{bulk} describing the amount of N [mg] in unfertilized soil samples (Table 1), fertilizer and fertilized soil samples, respectively, and $^{15}N_{nat}$ and $^{15}N_{fert}$ the ^{15}N enrichment under natural conditions (0.3663 at%) and in fertilizer (50 at%), respectively. Comparison of measured ^{15}N enrichment in N_2O and $^{15}N_{N_2O_exp}$ gave
385 information about the contribution of processes other than denitrification to N_2O production.

2.7 Statistical Analysis

We conducted several three-way analyses of variance (ANOVA) to test significant effects of soil, experimental variety and treatment on N_2O production, CO_2 production, and SP_{N_2O} , $\delta^{15}N^{bulk}_{N_2O}$ and $\delta^{18}O_{N_2O}$ values. The pairwise
390 comparison with Tukey's HSD test allowed to find differences between soils, varieties and treatments influencing N_2O production, CO_2 production, and isotopic values. Significant effects of soils and treatments on $r_{C_2H_2}$ and r_{15N} were tested by two-way ANOVA, while differences between soils and treatments influencing the product ratios were tested with pairwise comparison with Tukey's HSD test. Effects of varieties $-C_2H_2$ and *traced* on N_2O and CO_2 production were tested by ANOVA. For this ANOVA, the N_2O production rate had to be \log_{10} -transformed to
395 achieve homogeneity of variance and normality. The significance level α was 0.05 for every ANOVA. For some ANOVAs treatments were excluded, when replicates were $n < 3$. This was the case when only one or two samples out of three replicates could be analysed. This is denoted in the captions of tables (Table 2 and 3). The N_2O or CO_2 production rates of variety $+C_2H_2$ were followed over three sampling times by regression. For statistical analysis, we used the program R (R Core Team, 2013). Excel Solver tool was used to determine the $\delta^{18}O_{H_2O}$ values in the
400 application of SP/ $\delta^{18}O$ Map calculations.

3. Results


3.1 N_2O production rates

N_2O and CO_2 production rates of all treatments were similar in magnitude in almost all cases and mostly indistinguishable (Table 2, Figure 2). CO_2 production rates were determined to get additional information about
405 the denitrifying process. N_2O production rates exhibited increasing trends with ongoing incubation time for every soil with large variations within the treatments (Figure 2). Contrary to that, CO_2 production rates showed decreasing trends (Figure 2, exemplarily shown for data of variety $+C_2H_2$). Calculations of inhibitor effects were based on average N_2O and CO_2 production rates of the entire incubation period, i.e. ten hours of incubation time for *Soil 1.1*, 2 and 3 and eight hours for *Soil 1.2*.

410 N_2O and CO_2 production rates of all $+C_2H_2$ varieties differed significantly among soils ($P < 0.001$) and N_2O production rates differed also significantly among treatments ($P < 0.001$). Largest N_2O production rates of about 555 to 613 $\mu\text{g N kg}^{-1} \text{h}^{-1}$ were obtained in *Soil 1.1* and 3, respectively, while in *Soil 2* and *1.2* N_2O production rates

415 were smaller (271 and 264 $\mu\text{g N kg}^{-1}\text{h}^{-1}$, respectively). N_2O and CO_2 production rates were significantly larger in variety $+C_2H_2$ than in variety $-C_2H_2$ of *Soil 1.1*, *1.2* and *3* ($P < 0.001$, $P = 0.002$, and $P < 0.001$ for N_2O production rate and $P < 0.001$, $P = 0.027$, and $P = 0.008$ for CO_2 production rate, respectively) (Table 2), while $-C_2H_2$ and $+C_2H_2$ varieties of *Soil 2* did not differ in N_2O and CO_2 production rates ($P = 0.640$ and $P = 0.342$, respectively).

420 **Figure 2: Time series of average N_2O and CO_2 production rates during incubation of variety $+C_2H_2$ at the three sample collection times of each soil (*Soil 1* to *3*) for treatment A without growth inhibitors, B with bacterial growth inhibition, C with fungal growth inhibition, and D with bacterial and fungal growth inhibition; P -values for linear regressions (significance level $\alpha \leq 0.05$). For all significant regressions, R^2 -values were ≥ 0.46 and in the case of non-significance, R^2 -values were ≤ 0.40 . n.d.: There was no detectable CO_2 production in *Soil 1.2* at the first sampling time after 2 hours. (Figure is continued on next page)**



425




Figure 2 continued.

Without blockage of N₂O reductase (variety -C₂H₂), N₂O production rates of treatment A varied significantly
 430 among Soils with mean values between 175 and 355 µg N kg⁻¹ h⁻¹ ($P < 0.001$) (Table 2). In Soil 1.1, N₂O
 production rate was significantly larger (272 µg N kg⁻¹ h⁻¹) than in Soil 1.2 (175 µg N kg⁻¹ h⁻¹) ($P = 0.028$) in
 variety -C₂H₂. In most cases of the three varieties (-C₂H₂, +C₂H₂, and traced) treatment A (without growth
 inhibitors) produced most N₂O, followed by either treatment B (bacterial growth inhibitor; more N₂O compared
 to treatment C in Soils 1.2, 2, and 3) or treatment C (fungal growth inhibitor; more N₂O compared to treatment B
 435 in Soil 1.1). Smallest N₂O production rates were in most cases found in treatment D (non-inhibitable N₂O
 production) (except for variety -C₂H₂ of Soil 1.1, varieties -C₂H₂ and traced of Soil 3 and variety traced of Soil
 1.2). Microbial inhibitor treatments differed significantly in N₂O fluxes of variety +C₂H₂ of each Soil (always
 $P \leq 0.042$), while this was not the case for inhibitor treatments of varieties -C₂H₂ and traced of Soil 1.2 ($P = 0.154$
 and $P = 0.154$, respectively). Significant deviations of treatments without (A) or with full inhibition (D) were
 440 found in the following cases (Table 2): N₂O production rate of treatment A was significantly larger compared to
 the other three treatments of Soil 1.1 (+C₂H₂ and -C₂H₂), Soil 2 (-C₂H₂) and Soil 3 (+C₂H₂); treatment D was

445 significantly smaller compared to the other three treatments in *Soil 2* (+ C_2H_2) only and compared to treatments A and C in *Soil 1.1* (+ C_2H_2). A detailed discussion of inhibitor effects and difficulties with organisms that were not inhibited or abiotic sources is presented in section 4.1. Comparing varieties - C_2H_2 and *traced*, N_2O and CO_2 rates did not differ ($P = 0.991$ for N_2O production rate and $P = 0.490$ for CO_2 production rate, respectively), confirming that ^{15}N -labeling did not affect N_2O and CO_2 processes.

Table 2: Average CO₂ and N₂O production rates and N₂O isotopic values of N₂O of the last sample collection with and without C₂H₂ application in the headspace (varieties -C₂H₂ and +C₂H₂) of each soil (Soil 1 to 3) for treatments A without, B with bacterial, C with fungal, and D with bacterial and fungal growth inhibition, respectively (standard deviation in brackets, n = 3).

Treatment/ variety	mean N ₂ O [μg N kg ⁻¹ h ⁻¹]	mean CO ₂ [μg C kg ⁻¹ h ⁻¹]	δ ¹⁸ O _{N₂O} [‰]	δ ¹⁵ N ^{bulk} _{N₂O} [‰]	SP _{N₂O} [‰]
<i>Soil 1.1</i> (Loamy sand, winter 2012)					
A / -C ₂ H ₂	272.0 (38.4)a	1233.8 (170.5)a	13.1 (0.2)a	-21.9 (1.7)a	1.6 (0.8)a
B / -C ₂ H ₂	180.9 (16.8)b	1284.8 (168.0)a	13.0 (<0.1)*	-24.2 (0.7)*	-1.3 (0.2)*
C / -C ₂ H ₂	203.1 (14.4)ab	1124.8 (54.8)a	14.6 (0.4)a	-20.0 (0.8)a	-1.6 (0.5)a
D / -C ₂ H ₂	207.8 (32.6)ab	1371.7 (35.3)a	15.2 (0.5)*	-20.2 (1.8)*	-0.3 (0.5)*
A / +C ₂ H ₂	554.9 (46.5)a	1700.9 (98.1)a	8.5 (0.1)a	-22.1 (0.3)a	-0.4 (0.3)a
B / +C ₂ H ₂	353.5 (14.0)b	1610.7 (47.2)a	7.5 (0.1)a	-26.1 (0.2)a	-1.2 (1.0)a
C / +C ₂ H ₂	441.8 (18.5)c	1604.1 (60.3)a	9.3 (0.2)a	-22.4 (0.4)a	-0.9 (0.4)a
D / +C ₂ H ₂	331.0 (20.5)b	1438.0 (141.9)a	7.8 (0.3)*	-24.2 (0.1)*	-2.3 (0.7)*
<i>Soil 1.2</i> (Loamy sand, summer 2011)					
A / -C ₂ H ₂	175.3 (6.6)a	2448.5 (135.8)a	25.7 (0.3)a	-30.6 (0.2)a	12.1 (1.6)a
B / -C ₂ H ₂	121.3 (74.0)a	2091.3 (19.5)b	28.0 (5.0)a	-32.3 (0.7)a	7.7 (1.4)b
C / -C ₂ H ₂	104.5 (5.3)a	1844.7 (192.1)b	29.3 (0.1)a	-30.0 (0.5)a	4.3 (1.0)c
D / -C ₂ H ₂	73.8 (63.0)a	1632.2 (115.3)b	28.9 (1.2)a	-31.8 (2.2)a	3.4 (2.0)c
A / +C ₂ H ₂	263.5 (31.7)a	2076.6 (305.3)a	13.5 (0.5)*	-34.7 (0.1)*	-1.0**
B / +C ₂ H ₂	233.0 (15.6)ab	1794.9 (238.9)a	14.3 (1.7)a	-33.8 (0.9)a	-4.9 (0.9)a
C / +C ₂ H ₂	119.5 (102.7)b	1736.8 (424.7)a	19.0 (7.0)a	-33.1 (2.8)a	-1.7 (2.7)a
D / +C ₂ H ₂	161.6 (7.6)ab	1497.0 (138.7)a	14.8 (0.5)a	-35.7 (0.2)a	-4.9 (0.7)a
<i>Soil 2</i> (Sand, winter 2012)					
A / -C ₂ H ₂	315.0 (35.0)a	1316.7 (97.7)a	15.5 (1.8)a	-18.9 (2.6)a	-0.9 (2.5)a
B / -C ₂ H ₂	241.7 (3.0)b	1209.2 (24.6)a	15.0 (1.3)a	-23.4 (2.5)ab	-0.8 (<0.1)a
C / -C ₂ H ₂	247.6 (22.8)b	1201.9 (48.2)a	14.3 (0.1)a	-21.8 (0.2)ab	-1.8 (0.2)a
D / -C ₂ H ₂	198.4 (26.8)b	1102.4 (101.7)a	13.4 (0.3)a	-24.5 (0.1)b	-1.2 (0.3)a
A / +C ₂ H ₂	270.9 (36.3)a	1271.6 (203.5)a	12.6 (0.3)a	-18.9 (4.6)a	-1.4 (0.3)a
B / +C ₂ H ₂	263.1 (19.1)a	1338.7 (71.9)a	12.3 (0.1)a	-24.6 (0.2)b	-2.0 (0.2)a
C / +C ₂ H ₂	247.3 (15.9)a	1220.2 (50.0)a	12.7 (0.1)*	-23.3 (0.2)*	-1.7 (0.4)*
D / +C ₂ H ₂	187.3 (21.8)b	1173.1 (55.1)a	12.2 (0.3)a	-26.0 (0.1)b	-1.5 (0.9)a
<i>Soil 3</i> (Silt loam, winter 2013)					
A / -C ₂ H ₂	355.0 (18.4)a	1227.6 (95.2)a	26.0 (0.5)a	-20.8 (0.5)a	-0.5 (0.4)a
B / -C ₂ H ₂	325.4 (36.3)ab	1159.3 (178.2)a	24.1 (0.2)a	-22.0 (0.2)a	-0.1 (0.4)a
C / -C ₂ H ₂	278.9 (9.8)b	1056.0 (59.6)a	27.3 (0.1)a	-20.6 (0.3)a	0.6 (0.2)a
D / -C ₂ H ₂	291.1 (38.5)ab	1118.5 (70.3)a	26.3 (0.3)a	-21.0 (0.1)a	0.0 (0.2)a
A / +C ₂ H ₂	612.8 (25.2)a	1332.5 (116.9)a	15.2 (0.1)a	-25.6 (0.8)a	-2.8 (0.2)a
B / +C ₂ H ₂	546.9 (27.5)b	1235.7 (83.4)a	14.9 (0.2)a	-26.3 (<0.1)a	-3.5 (0.4)a
C / +C ₂ H ₂	519.8 (19.2)b	1173.5 (25.7)a	16.2 (<0.1)*	-25.2 (0.1)*	-4.0 (0.4)*
D / +C ₂ H ₂	511.7 (3.5)b	1295.6 (63.3)a	16.0 (0.1)a	-25.1 (0.1)a	-4.3 (0.5)a

Letters denote significant differences (P < 0.05) among treatments and varieties within a soil.

Asterisks indicate that only two samples (*) or one sample (**) of triplicates were analysable due to logistical difficulties.

3.2 Isotopologues of N₂O produced in different varieties and treatments

3.2.1 Variety -C₂H₂

460 SP_{N_2O} values of all *Soils* and inhibitor treatments of variety -C₂H₂ were within a range of -1.8 to 12.1 ‰ (Table 2) and differed among inhibitor treatments ($P = 0.037$). SP_{N_2O} values in variety -C₂H₂ of *Soil 1.2* was particularly large (3.4 to 12.1 ‰) compared to the other *Soils* (1.6 to -1.6 ‰). SP_{N_2O} values of variety -C₂H₂ were significantly larger than SP_{N_2O} values of variety +C₂H₂ ($P < 0.001$) (up to 2.4, 1.5, 4.6 and 4.1 ‰ in *Soil 1.1*, 2, 3 and 4, respectively). Generally, most SP_{prod} values of variety -C₂H₂ (Eq. 7) were smaller than SP_{N_2O} values of variety -C₂H₂ but still larger than SP_{N_2O} values of variety +C₂H₂ and are presented in Table S2 (Supplementary Material).

3.2.2 Variety +C₂H₂

465 SP_{N_2O} values of all *Soils* and all treatments of variety +C₂H₂ were within a narrow range between -4.9 and -0.4 ‰ (Table 2). In general, there were only small differences among treatments: SP_{N_2O} values of treatments A in variety +C₂H₂ differed significantly among soils ($P < 0.001$), with largest SP_{N_2O} values in *Soil 1.1* (-0.4 ‰) and smallest SP_{N_2O} values in *Soil 3* (-2.8 ‰). SP_{N_2O} values of treatment D in variety +C₂H₂ of all soils varied between -1.5 and -4.9 ‰, but only SP_{N_2O} values of *Soil 2* differed significantly from SP_{N_2O} values of the other *Soils* ($P = 0.006$). For 470 treatments B of variety +C₂H₂, SP_{N_2O} values differed only significantly between *Soil 1.1* and 1.2, 2 and 1.2, and 1.1 and 3 (each $P = 0.002$). SP_{N_2O} values from treatment C in variety +C₂H₂ did not differ significantly ($P = 0.600$). For every soil, we found significantly larger $\delta^{18}O_{N_2O}$, $\delta^{15}N^{bulk}_{N_2O}$ and SP_{N_2O} values in variety -C₂H₂ than in variety +C₂H₂ ($P < 0.001$), except for *Soil 2*, where $\delta^{15}N^{bulk}_{N_2O}$ values of variety -C₂H₂ were indistinguishable from those of variety +C₂H₂ ($P = 0.400$). However, only in a few varieties there were significant differences in $\delta^{18}O_{N_2O}$, 475 $\delta^{15}N^{bulk}_{N_2O}$ or SP_{N_2O} values between treatments with fungal and bacterial inhibition (B and C, respectively) (Table 2). As explained in section 3.3, N₂O reduction blockage in varieties +C₂H₂ was successful in most cases (*Soil 2*, 3 and 4). SP_{N_2O} values of this variety are thus assumed to be valid estimates of $\delta\theta$, i.e. SP_{prod} values of N₂O production, and can thus be used for applying the IEM.

3.2.3 Variety traced

480 The ¹⁵N-labeling of N₂O (¹⁵N_{N₂O}) or N₂ produced (¹⁵N_{N₂}) gave information about the incorporated N from ¹⁵N-labeled NO₃⁻ into N₂O or N₂ as well as about the N₂O reduction to N₂. Microorganisms in each treatment used the ¹⁵N-labeled NO₃⁻ in variety traced (Table 3) and expected ¹⁵N_{N₂O} depended on the initial N abundance in NO₃⁻ of unfertilized soil (Eq. 7). *Soil 1.2* is the only one showing a large discrepancy between measured (about 30 at%) and calculated ¹⁵N_{N₂O_{exp} (49 at%) in N₂O, whereas the other *Soils* showed close agreement (Table 3).}

485 3.3 Product ratios of denitrification and efficiency of N₂O reductase blockage by C₂H₂

$r_{C_2H_2}$ as well as r_{15N} determined with *Soil 2* were significantly larger than with the other *Soils* ($P \leq 0.001$) (Table 3). r_{15N} of treatment B was significantly larger than of treatment C and D of *Soil 1.2* ($P = 0.032$), while all other treatments of other *Soils* did not differ. $r_{C_2H_2}$ did not differ significantly among treatments ($P = 0.400$). In order to test the efficiency of blockage of N₂O reduction by C₂H₂ application, $r_{C_2H_2}$ (Eq. 5) was compared with r_{15N} (Eq. 490 6). In *Soil 1.1*, $r_{C_2H_2}$ was by far smaller than r_{15N} , while both calculated product ratios were in similar ranges in the other three *Soils* and thus a successful blockage of N₂O reduction was assumed for those *Soils*.

Table 3: Average CO₂ and N₂O production rates of the last sample collection after 10 or 8 hours of variety *traced*, respectively, with ¹⁵N labeling in N₂O (¹⁵N_{N2O}) and the calculated r_{15N} of variety *traced* and r_{C2H2} calculated from N₂O production rates of variety -C₂H₂ and +C₂H₂ of each soil (*Soil 1* to 3) for treatments A without, B with bacterial, C with fungal, and D with bacterial and fungal growth inhibition, respectively (standard deviation in brackets, $n = 3$).

Treatment	mean N ₂ O [μg N kg ⁻¹ h ⁻¹]	mean CO ₂ [μg N kg ⁻¹ h ⁻¹]	¹⁵ N _{N2O} [at%] ¹⁵ N _{N2O_exp} [at%] ^a	Calc. total r_{15N} ^b	Calc. total r_{C2H2} ^c
<i>Soil 1.1</i> (Loamy Sand, 2012)					
A	255.6 (43.5)	1310.0 (167.3)	36.8 (0.1)	0.80 (0.02)	0.48 (0.07)
B	154.5 (29.6)	1153.5 (238.4)	36.4 (0.2)	0.76 (0.02)	0.48 (0.05)
C	191.6 (30.7)	1219.6 (109.1)	36.9 (<0.1)	0.72 (0.05)	0.45 (0.04)
D	148.1 (1.9)	1253.8 (54.5)	36.8 (0.1)	0.69 (0.02)	0.54 (0.05)
<i>Soil 1.2</i> (Loamy Sand, 2011)					
A	156.9 (62.7)	3111.4 (1252.5)	31.1**	0.54 (0.05)	0.63 (0.10)
B	169.2 (6.1)	2314.6 (307.1)	26.5**	0.59 (0.03)	0.63 (0.17)
C	117.2 (3.1)	1785.6 (79.3)	30.1 (1.1)*	0.50 (0.01)	0.62 (0.02)
D	115.2 (3.1)	1706.7 (38.1)	33.5 (0.5)*	0.50 (0.01)	0.53 (0.12)
<i>Soil 2</i> (Sand, 2012)					
A	240.7 (0.95)	1286.2 (5.6)	43.2 (<0.1)	0.94 (0.01)	1.04 (0.10)
B	185.1 (3.9)	1157.4 (17.3)	43.0 (0.1)	0.94 (0.01)	0.81 (0.04)
C	241.1 (13.4)	1282.1 (63.4)	43.2 (0.1)	0.95 (0.01)	0.99 (0.09)
D	167.3 (34.9)	1199.0 (34.6)	42.7 (0.1)	0.93 (0.01)	0.98 (0.04)
<i>Soil 3</i> (Silt loam, 2013)					
A	285.9 (20.4)	1044.0 (46.6)	35.8 (<0.1)	0.62 (<0.01)	0.52 (0.04)
B	320.5 (14.7)	1204.2 (86.5)	35.5 (<0.1)	0.62 (0.01)	0.59 (0.02)
C	216.4 (34.9)	980.5 (202.5)	35.5 (<0.1)	0.59 (0.02)	0.48 (0.04)
D	231.4 (11.4)	988.5 (74.4)	35.3 (<0.1)	0.62 (0.01)	0.51 (0.04)

Asterisks indicate that only two samples (*) or one sample (**) were analysed due to logistical difficulties.

^a $r_{15N_{N2O_{exp}}}$ [at%] was calculated from Eq. 8.

^b $r_{15N} = [N_2O/(N_2+N_2O)]$ with N₂O or N₂ production rates from variety *traced*; see Eq. 5

^c $r_{C2H2} = [N_2O-C_2H_2/N_2O+C_2H_2]$ with N₂O production rate from varieties -C₂H₂ and +C₂H₂; see Eq. 6, cf. Table 2

500 3.4 Fungal contribution to N₂O production from denitrification by microbial inhibitor approach (modified SIRIN)

When calculating f_{FDmi} , N₂O production rates of treatment D must be significantly smaller compared to the other three treatments and the flux balance according to Eq. 1 and 2 must be consistent. Taking the large ranges of N₂O production rates of each treatment (minimum and maximum values) into account, for each *Soil* (A-D) was indistinguishable from ((B-D)+(C-D)) (Eq. 2), showing good agreement between Eqs. 1 and 2. However, N₂O production in treatment D was large within all varieties. Only with *Soil 2* of the variety +C₂H₂, the N₂O production rates of treatment D were significantly smaller than those of the other three treatments. Thus, for *Soil 2*, f_{FDmi} could be calculated (Eq. 3) and amounted to $28 \pm 9\%$ (Table 5) with a corresponding fungal N₂O production rate of $23.7 \pm 1.8 \mu\text{g N kg}^{-1} \text{ h}^{-1}$. Although the N₂O production rate of Treatment D was smaller than that of treatment A (*Soil 2*), it must be pointed out that due to the large amount of non-inhibitable production (treatment D), even the result for *Soil 2* is actually very unsure. For all other *Soils*, calculation of f_{FDmi} was not possible, i.e. SIRIN was not successful.

3.5 Fungal contribution to N₂O production from denitrification by the SP endmember mixing approach (IEM) and SP/ $\delta^{18}\text{O}$ isotope mapping approach (SP/ $\delta^{18}\text{O}$ Map)

515 The IEM revealed that f_{FD_SP} was small in all *Soils* ($\leq 15\%$, $\leq 14\%$, $\leq 9\%$, and $\leq 11\%$ with *Soil 1* to *3*, respectively) (Table 5). Regardless of influence of N₂O reduction on SP_{N_2O} values, only in *Soil 1.2* f_{FD_SPpot} could have reached 66%, while fungal denitrification could not have dominated with the other three soils (Table 5).

When applying SP/ $\delta^{18}\text{O}$ Map, we can assess the plausibility of the determined f_{FD} values based on the $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values obtained from the fitting ($\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value in Table 4) and the fitting outcome, i.e. the difference between $r_{15\text{N}}$ and r_{MAP} (*Diff*, see Table 4). The most probable $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value for our *Soils* can be assumed based on the fact that Braunschweig tap water was used and the original soil water also represents the isotope characteristics typical for this region, which is about -7.4% (long-term mean Braunschweig precipitation water (Stumpp et al., 2014)). Thus, in the presented application of SP/ $\delta^{18}\text{O}$ Map, $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values were fitted and it has to be pointed out that the precision of such calculations can be improved by measuring $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ instead. Depending on the season and evaporative losses, $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ may slightly vary and the most possible range of soil water in our *Soils* may vary from about -11 to -4% as observed in other experiments used in our laboratory experiments with similar conditions (Lewicka-Szczebak et al., 2014; Rohe et al., 2014a; Lewicka-Szczebak et al., 2017; Rohe et al., 2017). Taking this into account, we can say that for *Soil 1.1*, the fungal contribution must be below 2% , because to obtain any larger f_{FD} values, unrealistically small $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -14.9%) must be fitted (see Table 4). For *Soil 2*, both the smaller f_{FD_MAP} values of 1% and the larger ones up to 15% are possible, since they are associated with very realistic $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -6.3 and -10.1 , respectively) and identical *Diff* of 0.04 (Table 4). For *Soil 3*, the only plausible fitting can be obtained for the smallest SP_{BD} values, which are associated with a $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value of -5.6% (Table 4). Although the *Diff* for this fitting is slightly higher, the other fittings must be rejected due to unrealistic $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values (of -1.7 and $+3.7\%$), hence f_{FD_MAP} values must be between 4 and 9% . Similarly, for *Soil 1.2*, the only plausible fitting can be obtained for the smallest SP_{BD} values, which are associated with a $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value of -6.8% (Table 4) and indicate f_{FD_MAP} values from 11 to 20% . Here this fitting also shows clearly the smallest *Diff* of only 0.01 (Table 4). However, except for *Soil 1.2*, where the *Diff* is smallest for the last fitting, the *Diff* values for other *Soils* are very similar for different fittings with the largest values in *Soil 3*. A better fit (showing smaller *Diff* values) was not possible with any other combination of SP_{BD} and $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values. Since the precision of $r_{15\text{N}}$ (expressed in standard deviation in Table 3) was always ≤ 0.05 , this uncertainty of $r_{15\text{N}}$ did not reduce the precision of the fitting (compare large ranges of $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ and r_{MAP} values, respectively, in Table 4). The f_{FD_SP} ranged between 0 and approximately 15% (Table 5). The results obtained from SP/ $\delta^{18}\text{O}$ Map show f_{FD_MAP} reaching up to 14 , 20 , 15 , and 9% for *Soils 1.1*, *1.2*, *2*, and *3*, respectively (Figure 3, Table 4, Table 5). Importantly, due to the fitting procedure applied the estimations of f_{FD_MAP} values are based not only on SP_{N_2O} and $\delta^{18}\text{O}_{N_2O}$ values but also on the results obtained in the ^{15}N treatment ($r_{15\text{N}}$ values).

550

555

Table 4: Summary of the results provided by SP/ $\delta^{18}\text{O}$ Map for fraction of fungal denitrification (f_{FD_MAP}) and N_2O product ratio (r_{MAP}) in the acetylated ($+C_2H_2$) and non-acetylated ($-C_2H_2$) varieties for three possible SP_{N_2O} values from bacterial denitrification (SP_{BD}): mean (-1.9 ‰), maximal (3.7 ‰), and minimal (-7.5 ‰). The $\delta^{18}\text{O}$ values of soil water ($\delta^{18}\text{O}_{H_2O}$) were fitted to get the lowest difference (*Diff*) between product ratio determined with ^{15}N treatment (r_{15N}) and SP/ $\delta^{18}\text{O}$ Map (r_{MAP}). The most plausible fittings are shown in bold (see discussion for reasons of this choice).

<i>Soil</i>	<i>Variety</i>	r_{15N}	SP_{BD} [‰]	$\delta^{18}\text{O}_{H_2O}$ [‰]	r_{MAP}	<i>Diff</i>	f_{FD_MAP} [%]*
1.1	$-C_2H_2$	0.66	-1.9	-11.2	0.66	0.00	-1
	$+C_2H_2$	1	-1.9	-11.2	1.00	0.00	2
	$-C_2H_2$	0.66	3.7	-6.1	0.65	0.01	-14
	$+C_2H_2$	1	3.7	-6.1	1.00	0.00	-16
	$-C_2H_2$	0.66	-7.5	-14.9	0.66	0.00	8
	$+C_2H_2$	1	-7.5	-14.9	1.00	0.00	14
1.2	$-C_2H_2$	0.60	-1.9	-3.3	0.66	0.06	15
	$+C_2H_2$	1	-1.9	-3.3	0.96	0.04	-30
	$-C_2H_2$	0.60	3.7	1.5	0.72	0.12	8
	$+C_2H_2$	1	3.7	1.5	0.91	0.09	-21
	$-C_2H_2$	0.60	-7.5	-6.8	0.61	0.01	20
	$+C_2H_2$	1	-7.5	-6.8	0.99	0.01	11
2	$-C_2H_2$	0.94	-1.9	-6.3	0.90	0.04	1
	$+C_2H_2$	1	-1.9	-6.3	1.04	0.04	1
	$-C_2H_2$	0.94	3.7	-1.2	0.90	0.04	-16
	$+C_2H_2$	1	3.7	-1.2	1.04	0.04	-18
	$-C_2H_2$	0.94	-7.5	-10.1	0.90	0.04	13
	$+C_2H_2$	1	-7.5	-10.1	1.04	0.04	15
3	$-C_2H_2$	0.61	-1.9	-1.7	0.54	0.07	-3
	$+C_2H_2$	1	-1.9	-1.7	1.04	0.04	-5
	$-C_2H_2$	0.61	3.7	3.7	0.54	0.07	-14
	$+C_2H_2$	1	3.7	3.7	1.03	0.03	-24
	$-C_2H_2$	0.61	-7.5	-5.6	0.53	0.08	4
	$+C_2H_2$	1	-7.5	-5.6	1.04	0.04	9

*Negative values for f_{FD_MAP} are non-realistic and therefore discarded for further interpretation.

560

565

570

575

580

Table 5: Ranges of the fraction of N₂O produced by fungi (f_{FD}) from four Soils using different approaches: Fungal fraction was calculated using a) the microbial inhibitor approach (modified SIRIN) (f_{FDmi}), the isotopomer endmember mixing approach (IEM) by SP isotope mixing balance using b) variety +C₂H₂ (f_{FD_SP}), c) f_{FD_SPpot} for results from variety -C₂H₂ assuming the SP effect of N₂O reduction was negligible, d) for results from variety -C₂H₂ with reduction correction to calculate the SP_{N2O} values (f_{FD_SPcalc}), and e) the $\delta^{18}O/SP$ Map (f_{FD_MAP}) with $\delta^{18}O_{N2O}$ and SP_{N2O} values from variety -C₂H₂ and variety +C₂H₂.

<i>Soil</i>	f_{FDmi} [%] ^a	f_{FD_SP} [%] ^{b*}	f_{FD_SPpot} [%] ^{c*}	f_{FD_SPcalc} [%] ^{d*}	f_{FD_MAP} [%] ^{e*}
1.1	n.d.	-14 to 15	-12 to 39	-6 to 19	<2
1.2	n.d.	-23 to 11	10 to 66	1 to 21	11 to 20
2	19 to 37	-18 to 14	-14 to 36	-12 to 15	1 to 15
3	n.d.	-25 to 9	-11 to 40	-9 to 18	4 to 9

585

^aFungal fraction of N₂O production calculated by Eq. 3 taking variations of three replicates into account

^bFungal fraction of N₂O production calculated by Eq. 4 for variety +C₂H₂ with assuming SP_{N2O} values of N₂O produced by bacteria were 3.7 ‰ or -7.5 ‰ (Yu et al., 2020) and by fungi on average 33.6 ‰ (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017). Using the minimum and maximum SP_{N2O} values known for bacteria resulted in a f_{FD_SP} range.

590

^cMaximum potential fungal fraction of N₂O production calculated by Eq. 4 as an average range for all treatments of variety -C₂H₂ assuming SP_{N2O} values of N₂O produced by bacterial denitrification or nitrifier denitrification were between 3.7 and -10.7 ‰ (Frame and Casciotti, 2010; Yu et al., 2020) or produced by fungal denitrification or nitrification were between 16 and 37 ‰ (Sutka et al., 2008; Decock and Six, 2013; Rohe et al., 2014a; Maeda et al., 2015; Rohe et al., 2017). Using the minimum and maximum SP_{N2O} values known from pure cultures resulted in the given f_{FD_SPpot} range. Here, the effect of partial reduction of N₂O on SP_{N2O} values was assumed to be negligible.

595

^dEq. 4 to solve for fungal fraction in variety -C₂H₂ with assuming SP_{N2O} values of N₂O produced by bacteria was 3.7 (resulting in negative fraction and therefore set to zero) or -7.5 ‰ and using reduction correction with $\eta_r = -6$ ‰ to calculate SP_{prod} values (Senbayram et al., 2018; Yu et al., 2020). Using the minimum and maximum SP_{N2O} values known for bacteria resulted in a f_{FD_SPcalc} range.


600

^eFungal fraction of N₂O production calculated by SP/ $\delta^{18}O$ Map with assuming most probable SP_{N2O} values from bacterial denitrification (according to Table 4). Using the minimum and maximum SP_{N2O} values known for bacteria and ranges of fitted $\delta^{18}O_{H2O}$ values (the fitting is based also on results obtained in ¹⁵N treatment) resulted in a f_{FD_MAP} range.

605

*Negative values for f_{FD_SP} , f_{FD_SPpot} , f_{FD_SPcalc} , f_{FD_MAP} are non-realistic and therefore discarded for further interpretation.

n.d.-not determined because of insufficient inhibition.



610 Figure 3: SP/ $\delta^{18}\text{O}$ isotope mapping approach (SP/ $\delta^{18}\text{O}$ Map) to estimate the contribution of bacteria or fungi to N_2O
 produced according to Lewicka-Szczebak et al. (2017) and Lewicka-Szczebak et al. (2020). The isotopic values for
 natural abundance treatments with acetylene addition (+ C_2H_2 , empty symbols) and without acetylene addition (- C_2H_2 ,
 615 corresponding filled symbols) are shown for four Soils (1 to 3). The grey rectangles indicate expected ranges of isotopic
 signatures for heterotrophic bacterial denitrification (BD) and fungal denitrification (FD) (Yu et al., 2020). The black
 solid line is the mixing line connecting the average expected values for BD and FD, while the red solid line is the mean
 reduction (for the mean SP values for BD) line and the red dashed line is the minimum reduction line (for the minimal
 SP $_{\text{N}_2\text{O}}$ values for BD).

4. Discussion

620 To our knowledge, this was the first attempt to determine SP $_{\text{N}_2\text{O}}$ values by fungi or bacteria from soil communities
 using microbial growth inhibitors with a modification of SIRIN and comparing microbial inhibitor and isotopic
 approaches (IEM and SP/ $\delta^{18}\text{O}$ Map) to estimate fungal contribution to N_2O production from denitrification in
 anoxic incubation. The isotopic approaches revealed that the fungal contribution to N_2O production was small
 ($f_{\text{FD}_{\text{SP}}} \leq 15\%$ or $f_{\text{FD}_{\text{MAP}}} \leq 20\%$) in the soils tested (Table 5). A dominant contribution of fungi over bacteria was
 625 also excluded by the potential maximum fungal denitrification for Soil 1.1, 2, and 3 ($f_{\text{FD}_{\text{SPpot}}}$ between 37 and 40
 %, Table 5), even though effects of N_2O reduction are not included. The modified SIRIN approach was not
 successful, because large amounts of non-inhibitable N_2O production were observed with all four Soils (Table 2,
 Table 3). The fungal fraction producing N_2O during denitrification ($f_{\text{FD}_{\text{mi}}}$) was only estimated for Soil 2, where
 significantly smaller N_2O production in treatment D was observed compared to that of treatment A and resulted in
 630 a range of 19 to 37 %, which was probably overestimated due to uncertainties resulting from the large N_2O
 production of non-inhibitable sources. While the three approaches coincided in showing dominance of bacterial
 denitrification, the isotopic approaches yielded small estimates for $f_{\text{FD}} (\leq 20\%)$ and thus did not confirm largest
 $f_{\text{FD}_{\text{mi}}}$ of Soil 2. The strict application of the SIRIN method prescribes proof of selectivity of the inhibitors (i.e.
 streptomycin should not inhibit fungi and cycloheximide should not inhibit bacteria). All SIRIN results obtained
 635 with respect to N_2O production by the fungal or bacterial fraction were unsatisfactory, thus fungal SP $_{\text{N}_2\text{O}}$ values
 could not be assessed, and the overall results led to unsolved questions, which are discussed in the following
 sections.

4.1 Experimental setup and inhibitor effects

In accordance with other studies, N₂O production was analysed after the addition of glucose as substrate (Laughlin and Stevens, 2002; McLain and Martens, 2006; Blagodatskaya et al., 2010; Long et al., 2013). Glucose initiates the growth of active heterotrophic organisms. Since pure cultures were shown to synthesize enzymes capable of denitrification within two to three hours (USEPA, 1993), pre-incubation of soil under anaerobic conditions is not needed. Thus, when gas sample collection started organisms should have produced denitrifying enzymes and microbial growth of initially active organisms should have started too. However, in accordance to Anderson and Domsch (1975) experimental duration should be as short as possible to ensure the CO₂ production by initially active organisms only. Thus, short-time incubation is recommended when conducting a modified SIRIN approach, as the incubation period should cause changes in conditions for microorganisms and initiate growth on the one hand, while it should avoid the consumption of inhibitors as C sources on the other.

With incubation time, production rates of CO₂ decreased, probably because experimental incubation conditions provoked unfavourable conditions and physiological changes, e. g. due to anaerobic conditions or local substrate depletion (e. g. C supplied as glucose). Decreasing CO₂ fluxes might also be explained by CO₂ accumulation in pore space as this effect is shown by modelled diffusive fluxes from soil in closed systems (Well et al., 2019).

Previous studies found much larger inhibitor effects (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013; Chen et al., 2014). It is therefore important to discuss considerable differences among the experimental design of the present study compared to that of other studies (e. g., Laughlin and Stevens, 2002; Blagodatskaya et al., 2010).

The conventional practice of SIRIN implies determination of $c_{opt}(\text{glucose})$, $c_{opt}(\text{streptomycin})$ or $c_{opt}(\text{cycloheximide})$ with an "Ultragas 3" CO₂ analyser (WösthoffCo., Bochum) (Anderson and Domsch, 1973) with continuous gas flow. We used this method to determine optimal concentrations for SIRIN in the pre-experiment and used these concentrations for the modified SIRIN approach as well. This optimization procedure was not used in other studies (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013). We supposed that optimal concentrations for CO₂ respiration should work as well for denitrification, if both inhibitors inhibit the denitrification process as well. However, although SIRIN has so far been tested with isolated cultures and soils for microbial growth for CO₂ production only (Anderson and Domsch, 1973, 1975), information on N₂O producing processes, especially denitrification, is still lacking and should be investigated in further studies. Additionally, as presented by Ladan and Jacinthe (2016) the bactericide bronopol and the fungicide captan were more effective inhibitors than streptomycin or cycloheximide and should be included when evaluating inhibition approaches and isotopic endmember approaches.

Previous studies that found much larger inhibitor effects were conducted after pre-incubating the soil with selective inhibitors (Laughlin and Stevens, 2002; Blagodatskaya et al., 2010; Long et al., 2013; Chen et al., 2014). In contrast to that, the experimental design of our incubation setup was without soil pre-incubation with selective inhibitors to minimize disturbance of the soil microbial community and our approach was thus in agreement with the original SIRIN method for respiration (Anderson and Domsch, 1973, 1975, 1978). Another study performing similar experiments without pre-incubation with inhibitors did not find effectiveness of application of both antibiotics during long-term application (up to 48 h) (Ladan and Jacinthe, 2016), although streptomycin and cycloheximide are commonly used to inhibit denitrification of selective groups. Nevertheless, as we expected that pre-incubation with selective inhibitors would induce changes in the F:B ratio of soil, we decided to conduct the modified SIRIN approach without a pre-incubation step. This assumption was supported by findings of Blagodatskaya et al. (2010),

680 where pre-incubation of about one to twenty hours with cycloheximide resulted in increasing inhibitor efficiency with time, while this was not the case when pre-incubating with streptomycin. This suggests that microbial communities might change after exposition to cycloheximide.

In the present study, even with both growth inhibitors (treatment D), N₂O production was large in all experiments, i.e. in most cases not significantly smaller than in treatments A, B or C. Thus, we suppose similar contributions of non-inhibitable organisms and processes in all treatments. Non-inhibitable organisms could be, for example, 685 bacteria or fungi that are not in growth stage or may be not affected by inhibitors. Recently, Pan et al. (2019) summarized findings of other studies and pointed out that some microorganisms can use inhibitors as growth substrates, that dead organisms may serve as energy sources for others, and that interactions of microbial species may change due to non-inhibitable organisms occurring in soil communities. Non-inhibitable organisms could be archaea as well, which are also known to be capable of denitrification (Philippot et al., 2007; Hayatsu et al., 2008). 690 It is known that archaea are not affected by streptomycin or cycloheximide (Seo and DeLaune, 2010). However, effects of archaeal occurrence in soil or secondary effects on fungi or bacteria were not tested in this study. Additionally, abiotic N₂O production cannot be quantified with the experimental setup, but might be contributing to each inhibitor treatment.

In summary, the present experimental setup without pre-incubating soil samples with selective inhibitors was not 695 successful in complete inhibition of bacterial or fungal denitrifiers. Although pre-incubation with selective inhibitors may lead to more successful inhibition, we do not recommend this due to induced changes in soil communities. For further studies focusing on application of modified SIRIN to determine the fraction of bacterial or fungal N₂O derived from denitrification a method validation using also different inhibitors is recommended.

700 **4.2 Is C₂H₂ application a suitable and necessary treatment for examining the fungal contribution to N₂O production in soil?**

In order to determine SP_{N_2O} values without alteration by partial reduction of N₂O to N₂, C₂H₂ was used to quantitatively block N₂O reduction during denitrification. We found the expected effect of C₂H₂ application, i.e. larger N₂O production rates in variety +C₂H₂ compared to variety -C₂H₂. Calculated product ratios varied between 0.5 and 0.95 (r_{15N}) in all *Soils*, showing that N₂O reduction can have significant effects on measured N₂O 705 production and isotopic values.

The calculated $r_{C_2H_2}$ was within the same range as r_{15N} in *Soil 1.2*, *2*, and *3* (maximal 9 % difference), indicating effective blockage of N₂O reductase in variety +C₂H₂ in these *Soils*. Only in *Soil 1.1*, r_{15N} and $r_{C_2H_2}$ differed by about 34 % with larger calculated reduction in the *traced* variety, which might point to incomplete inhibition by the C₂H₂ method. Artifacts with C₂H₂ were found in previous studies, resulting in smaller N₂O production rates 710 due to NO oxidation accelerated by C₂H₂ application in the presence of very small O amounts (Bollmann and Conrad, 1997a, b; Nadeem et al., 2013). Moreover, incomplete C₂H₂ diffusion into denitrifying aggregates might also lead to incomplete N₂O reductase blockage (Groffman et al., 2006). Both potential methodological errors cannot be excluded for *Soil 1.1*.

For the other three *Soils* (*1.2*, *2*, and *3*), it can be supposed that the isotopic signature of N₂O of variety +C₂H₂ 715 showed isotopic signatures of produced N₂O without influences of N₂O reduction (SP_{prod}). By comparing varieties -C₂H₂ and +C₂H₂, isotopologue values of all these *Soils* (except $\delta^{15N}_{bulk_{N_2O}}$ values of *Soil 2*) of variety -C₂H₂ were significantly larger than those of variety +C₂H₂. The enrichment of residual N₂O in heavy isotopes results from the isotope effect associated with N₂O reduction (Jinuntuya-Nortman et al., 2008; Well and Flessa, 2009; Lewicka-

720 Szczebak et al., 2014). This explains why C_2H_2 application is essential for analysing N_2O produced by different microbial groups from soil. This has particular relevance for experiments with modified SIRIN approaches. Although the modified SIRIN approach presented here was not successful, it should be noted that comparable soil incubation experiments without quantifying N_2O reduction potentially overestimate fungal denitrification due to the impact of SIRIN inhibitors on N_2O reduction.

725 Of course, N_2O fluxes represent net N_2O production, i.e. the difference between gross N_2O production by the microbial community and N_2O reduction, mainly by heterotrophic bacterial denitrifiers (Müller and Clough, 2014). It has been shown that N_2O released by microorganisms to air-filled pore space can be partially consumed by denitrifiers before being emitted (Clough et al., 1998). This means that fungal N_2O can also be subject to reduction by bacterial denitrifiers. Consequently, successful inhibition of bacterial denitrification by SIRIN would enhance the measured flux of fungal N_2O . Until now, this effect has not been considered in SIRIN papers on fungal N_2O 730 (e. g. Laughlin and Stevens, 2002; Ladan and Jacinthe, 2016; Chen et al., 2014). This effect can only be evaluated by measuring N_2O reduction in all inhibitor treatments. If true, the N_2O reduction with bacterial inhibition should be smaller than that of the treatments without inhibition or with fungal inhibition. However, with fungal inhibition, N_2O reduction is also assumed to be smaller than without inhibition, because N_2O produced by fungi is missed for bacterial reduction.

735 As the product ratio in soil denitrification exhibited the full range from 0 to 1, this effect can be quite relevant and must thus be considered in future studies. Therefore, we recommend to estimate the effectiveness of C_2H_2 in blocking the N_2O reductase by performing parallel ^{15}N approaches with and without C_2H_2 in studies using the modified SIRIN to determine the fraction of bacterial or fungal N_2O production.

4.3 SP_{N_2O} values of N_2O produced by microbial communities

740 As discussed above, all N_2O fluxes of modified SIRIN treatments of *Soil 1.1*, *1.2*, and *3* were dominated by N_2O from non-inhibitable organisms or processes. This made it impossible to calculate SP_{N_2O} values for active bacteria or fungi (modified SIRIN B and C), also with *Soil 2*, where a relatively large N_2O production was observed with treatment D (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015) (see section 3.4).

745 Despite this, the SP_{N_2O} values from $+C_2H_2$ variety as well as SP_{prod} values (i.e. reduction corrected SP_{N_2O} values of $-C_2H_2$ variety) of each *Soil*, represented by treatment A of modified SIRIN, indicated predominantly bacteria to be responsible for N_2O production during denitrification, assuming that results of SP_{N_2O} values of denitrification by pure bacterial cultures are transferable to bacteria of soil communities contributing to denitrification. Also in many soil incubation studies, SP_{N_2O} values (without reduction effects) within the range of bacterial pure cultures have been found (Lewicka-Szczebak et al., 2015; Lewicka-Szczebak et al., 2017; Senbayram et al., 2018). 750 Therefore, there was so far no unequivocal evidence of fungi contributing to N_2O production during denitrification in soils, although here, the isotopic approaches were consistent with a fungal contribution of up to 20 % of N_2O production during denitrification.

The SP_{N_2O} values of variety $+C_2H_2$ within treatment A are not affected by reduction effects and therefore might give evidence of the microbial community contributing to N_2O production (Sutka et al., 2006; Sutka et al., 2008; 755 Frame and Casciotti, 2010; Rohe et al., 2014a). However, variations in SP_{N_2O} values of treatments A of variety $+C_2H_2$ were very small and do not give a clear evidence of any differences in microbial soil community producing N_2O . Lewicka-Szczebak et al. (2014) analysed SP_{N_2O} values of denitrification with blockage of N_2O reduction by C_2H_2 for the same soils as used in the present study (*Soil 1.1* and *1.2* as well as *Soil 3*) and revealed SP_{N_2O} values

between -3.6 and -2.1 ‰, which is similar to the respective SP_{N_2O} values of the present study from -4.9 to -0.4 ‰. This reinforces the conclusion that bacteria dominated gross N_2O production under anoxic conditions in both studies.

SP_{prod} values (variety $-C_2H_2$) differed from SP_{N_2O} values (variety $+C_2H_2$), which may result from deviations between the actual fractionation factor that was not estimated in the present study and the used fractionation factor of -6 ‰ adapted from the literature (Yu et al., 2020). If so, we could assume smaller fractionation effects in the present study as decreasing this average fractionation factor would lead to increasing SP_{prod} values, which in turn would result in values more similar to SP_{N_2O} values of variety $-C_2H_2$.

4.4 Potential influence of hybrid N_2O

When one N atom in N_2O originates from labeled NO_3^- and the other one from an unlabeled N source, this results in a_p values and ^{15}N enrichment of produced N_2O smaller than the respective enrichment of the NO_3^- pool. The ^{15}N enrichment of N_2O in *Soil 1.2* was about 60 % smaller than the ^{15}N enrichment in soil NO_3^- , leading to the assumption that N_2O was produced not only by denitrification. We also calculated a_p values of the other three *Soils* (data not shown) which coincided with the ^{15}N enrichment of N_2O (Table 3), showing no indication of hybrid N_2O . Since a_p would not be affected by contributions of unlabeled N_2O we can thus exclude the possibility that this smaller enrichment could be caused by dilution of enriched N_2O from denitrification by N_2O production from an unknown N source and thus verified that this was due to formation of hybrid N_2O , potentially via co-denitrification (Spott et al., 2011). So far, there is no study on SP_{N_2O} values of N_2O produced by co-denitrification. But since SP_{N_2O} values of the acetylated treatments of *Soil 1.2* coincided with the SP_{N_2O} value range of bacterial denitrification and also with SP_{N_2O} values of the other *Soils*, our data give no indication that the SP_{N_2O} values of hybrid N_2O , potentially produced during co-denitrification, differed from that of bacterial denitrification. It was however, remarkable that the maximum potential contribution of fungal denitrification to N_2O (f_{FD_SPpot}) was higher for *Soil 1.2* compared to that of *Soil 1.1* from the winter period. *Soil 1.2* was the only soil where f_{FD_SPpot} exceeded 50%, thus fungi may potentially dominate N_2O emissions only in this *Soil*.

4.5 Steps towards quantifying the fungal fraction contributing to N_2O production

Due to the inefficiency of the inhibition of microbial N_2O production in most cases, calculation of f_{FDmi} contributing to N_2O production was possible for *Soil 2* only, although even this calculated value included inaccuracies. The isotopic approaches, however, which are independent of modified SIRIN results, yielded similar estimates of f_{FD} for all *Soils*. As recently published (Wu et al., 2019), uncertainty analysis is a complex issue and large uncertainties of the results from the SP/ $\delta^{18}O$ Map approach can be assumed when all the possible sources of errors are taken into account. Regarding the presented application of SP/ $\delta^{18}O$ Map, calculation would be more precise when measuring $\delta^{18}O_{H_2O}$ rather than using the fitted $\delta^{18}O_{H_2O}$ values. Still, the analysis of $\delta^{18}O_{N_2O}$ values can give information about O exchange between water and denitrification intermediates by various microorganisms (Aeressens et al., 1986; Kool et al., 2007; Rohe et al., 2014b; Rohe et al., 2017). The range of $\delta^{18}O_{N_2O}$ values in our study for variety $+C_2H_2$ (7.5 to 19.0 ‰) was quite similar to the range found by Lewicka-Szczebak et al. (2014) for the same soils (4.8 to 16.3 ‰), where almost complete O exchange with soil water was documented. Hence, for this study the O exchange was probably also very high. There were also no remarkable differences in $\delta^{18}O_{N_2O}$ values among treatments within one variety and soil and therefore we assume no differences in O exchange among the treatments. The information on $\delta^{18}O_{N_2O}$ values combined with known $\delta^{18}O_{H_2O}$ values is also precious

information for differentiation between N₂O mixing and reduction processes (Lewicka-Szczebak et al., 2017). Due to parallel *traced* variety experiments, possible $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values for the particular SP_{N₂O} values of bacterial denitrification mixing endmembers could be determined (Table 4). Since the $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ value for the particular geographic region can be assessed based on the known isotopic signatures of meteoric waters (Lewicka-Szczebak et al., 2014; Stumpp et al., 2014; Lewicka-Szczebak et al., 2017; Buchen et al., 2018), the most plausible ranges of $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values can be used to indicate the plausible ranges of $f_{\text{FD_MAP}}$ values. Here we showed that in case of missing $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ values but known product ratio, the SP/ $\delta^{18}\text{O}$ Map can also provide information on N₂O production pathway contributions. Comparing the modified SIRIN with the isotopic approaches revealed that the fungal contribution to N₂O production was consistently estimated to be smaller (about 28 % in modified SIRIN, ≤ 15 % with IEM, ≤ 20 % with SP/ $\delta^{18}\text{O}$ Map) than the bacterial fraction. This was supported by estimates for maximum potential contribution of fungal denitrification to N₂O in variety -C₂H₂ ($f_{\text{FD_SPpot}}$) for *Soil 1.1*, 2 and 3. In some soil studies using helium incubations, the SP_{Prod} values obtained by correction for the reduction effect on $SP_{\text{N}_2\text{O}}$ values showed significantly larger values than SP_{N₂O} of bacterial denitrification (Köster et al., 2013a; Lewicka-Szczebak et al., 2017; Lewicka-Szczebak et al., 2014; Senbayram et al., 2018; Senbayram et al., 2020). However, those results were obtained in an experimental setup with ambient oxygen concentration. Short incubations under static conditions as presented here may, however, promote bacterial over fungal growth, which may also be transferable to denitrification activity by both organism groups (Lewicka-Szczebak et al., 2014; Lewicka-Szczebak et al., 2017). Obviously, based on the estimations from isotopic approaches, soils may largely differ in the microbial community that contributes to N₂O from denitrification.

However, all our tested soils seemed to contain a microbial community where fungi have minor contributions to N₂O emissions from denitrification compared to bacteria. This may also have been due to the applied experimental setup favoring bacterial denitrification by static and strictly anoxic conditions. Additionally, the use of glucose as substrate in the selected concentration may further promote bacteria compared to fungi (Koranda et al., 2014; Reischke et al., 2014). Senbayram et al. (2018) could show in an incubation experiment with sufficient NO₃⁻ supply, that fungal contribution to denitrification was larger with straw compared to a control without straw addition. Thus, experimental conditions need to be carefully set and more information is needed here in order to get a good representation of soil conditions in incubation experiments.

The isotopic approaches should be further investigated with soils where fungi are presumed to contribute largely to N₂O production (e. g. acid forest soils, or litter-amended arable soils) (Senbayram et al., 2018) and using SIRIN with more suitable inhibitors (Ladan and Jacinthe, 2016). The critical question whether the isotopic signatures of fungal N₂O determined in pure culture studies are transferable to natural soil conditions could not be answered with this study due to large uncertainties associated with the results of the SIRIN method. The latter precluded determination of $SP_{\text{N}_2\text{O}}$ values of N₂O from fungal denitrification. Further experiments would be needed with improved selective inhibition to assure that $SP_{\text{N}_2\text{O}}$ values known from a few pure cultures or soil isolates (Sutka et al., 2008; Rohe et al., 2014a; Maeda et al., 2015) are true for fungal soil communities as well. This could be accompanied by studies mixing various fungal species known to occur in soil or by isolating fungal communities from soil and conduct similar experiments under anoxic conditions with supply of electron acceptors and C sources to investigate denitrification. In such incubations, parallel ¹⁵N tracing experiments should be conducted to assure denitrification as the dominating process for N₂O production and quantify the possible contribution of co-denitrification.

5. Conclusions

840 Based on the presented results we conclude that the modified SIRIN approach in the form presented here is not appropriate to estimate the contribution of selected communities (bacteria or fungi) on denitrification from soil. The quantification of the fungal fraction with modified SIRIN could be done with one soil only and was possibly overestimated when compared with the results of isotopic approaches. Both isotope approaches (IEM and SP/ $\delta^{18}\text{O}$ Map) revealed similar results of the fungal fraction contributing to denitrification and thus could be recommended as equally suitable for future studies. The present study could show that consideration of N_2O reduction is
845 indispensable. It has to be pointed out, however, that the fungal fraction estimated applies only for the soil under presented experimental conditions, i.e. anaerobic conditions and with glucose amendment, but not for the investigated soil in general.

Further studies are needed to cross-validate methods, e. g. with improved inhibitor approaches or molecular-based methods. Due to the mentioned difficulties, the $SP_{\text{N}_2\text{O}}$ values of fungal N_2O could not be calculated from the
850 modified SIRIN approach. Several potential artefacts in the modified SIRIN approach should be further investigated, e.g. the effectiveness of inhibitors, changes in microbial community during pre-incubation with inhibitors and effects of bacterial consumption of N_2O produced by fungi.

Data availability. Gas emission and isotopic data are available from the authors on request.

Author contribution. HF, NWM, RW and THA designed the experiment. LR carried out the experiment at Thünen
855 Institute for Climate-Smart Agriculture in Braunschweig. AG, DLS and RW helped with isotopic analysis and DLS performed the $\delta^{18}\text{O}$ /SP Map. LR, RW and DLS prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Many thanks are due to Jens Dyckmans for ^{15}N analysis and to Martina Heuer for N_2O isotopic analyses. This joint research project was financially supported by the State of Lower-Saxony and the Volkswagen
860 Foundation, Hanover, Germany. Further financial support was provided by the German Research Foundation (grant LE 3367/1-1 to DLS, grant WE 1904/8-1 to LR and RW, and the research unit 2337: “Denitrification in Agricultural Soils: Integrated Control and Modeling at Various Scales (DASIM)”, grant WE 1904/10-1 to RW and WR 211/1-2 to NWM).

References

865 Aerssens, E., Tiedje, J. M., and Averill, B. A.: Isotope labeling studies on the mechanisms of N-bond formation in denitrification, *J. Biol. Chem.*, 261, 9652-9656, 1986.

Anderson, J. P. E., and Domsch, K. H.: Quantification of bacterial and fungal contributions to soil respiration, *Archiv Fur Mikrobiologie*, 93, 113-127, doi:10.1007/BF00424942, 1973.
870

Anderson, J. P. E., and Domsch, K. H.: Measurement of bacterial and fungal contributions to respiration of selected agricultural and forest soil, *Canadian Journal of Microbiology*, 21, 314-322, doi: 10.1139/m75-045, 1975.

- 875 Anderson, J. P. E., and Domsch, K. H.: Physiological method for quantitative measurement of microbial biomass in soils, *Soil Biol. Biochem.*, 10, 215-221, doi:10.1016/0038-0717(78)90099-8, 1978.
- Bergsma, T. T., Ostrom, N. E., Emmons, M., and Robertson, G. P.: Measuring simultaneous fluxes from soil of N₂O and N₂ in the field using the ¹⁵N-Gas "nonequilibrium" technique, *Environmental Science & Technology*, 35, 4307-4312, doi: 10.1021/es010885u, 2001.
- 880 Blagodatskaya, E., Dannenmann, M., Gasche, R., and Butterbach-Bahl, K.: Microclimate and forest management alter fungal-to-bacterial ratio and N₂O-emission during rewetting in the forest floor and mineral soil of mountainous beech forests, *Biogeochemistry*, 97, 55-70, doi:10.1007/s10533-009-9310-3, 2010.
- 885 Blagodatskaya, E. V., and Anderson, T.-H.: Interactive effects of pH and substrate quality on the fungal-to-bacterial ratio and qCO₂ of microbial communities in forest soils, *Soil Biol. Biochem.*, 30, 1269-1274, doi: 10.1016/S0038-0717(98)00050-9, 1998.
- 890 Bollag, J. M., and Tung, G.: Nitrous oxide release by soil fungi, *Soil Biol. Biochem.*, 4, 271-276, doi: 10.1016/0038-0717(72)90021-1, 1972.
- Bollmann, A., and Conrad, R.: Enhancement by acetylene of the decomposition of nitric oxide in soil, *Soil Biol. Biochem.*, 29, 1057-1066, 1997a.
- 895 Bollmann, A., and Conrad, R.: Acetylene blockage technique leads to underestimation of denitrification rates in oxic soils due to scavenging of intermediate nitric oxide, *Soil Biol. Biochem.*, 29, 1067-1077, 1997b.
- 900 Braker, G., and Conrad, R.: Diversity, structure, and size of N₂O-producing microbial communities in soils-What matters for their functioning?, in: *Advances in Applied Microbiology*, Vol 75, edited by: Laskin, A. I., Sariaslani, S., and Gadd, G. M., *Advances in Applied Microbiology*, 33-70, 2011.
- Brand, W. A.: Precon: A fully automated interface for the pre-GC concentration of trace gases in air for isotopic analysis, *Isot. Environ. Health S.*, 31, 277-284, doi: 10.1080/10256019508036271, 1995.
- 905 Bremner, J.: Sources of nitrous oxide in soils, *Nutrient Cycling in Agroecosystems*, 49, 7-16, doi: 10.1023/A:1009798022569, 1997.
- 910 Buchen, C., Lewicka-Szczebak, D., Flessa, H., and Well, R.: Estimating N₂O processes during grassland renewal and grassland conversion to maize cropping using N₂O isotopocules, *Rapid Commun. Mass Spec.*, 32, 1053-1067, doi: 10.1002/rcm.8132, 2018.
- 915 Casciotti, K. L., Sigman, D. M., Hastings, M. G., Böhlke, J. K., and Hilkert, A.: Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, 74, 4905-4912, doi: 10.1021/ac020113w, 2002.
- 920 Chen, H., Mothapo, N. V., and Shi, W.: The significant contribution of fungi to soil N₂O production across diverse ecosystems, *Appl. Soil Ecol.*, 73, 70-77, doi: 10.1016/j.apsoil.2013.08.011, 2014.
- Chen, H. H., Mothapo, N. V., and Shi, W.: Soil Moisture and pH Control Relative Contributions of Fungi and Bacteria to N₂O Production, *Microb. Ecol.*, 69, 180-191, doi: 10.1007/s00248-014-0488-0, 2015.
- 925 Clough, T. J., Jarvis, S. C., Dixon, E. R., Stevens, R. J., Laughlin, R. J., and Hatch, D. J.: Carbon induced subsoil denitrification of ¹⁵N-labelled nitrate in 1 m deep soil columns, *Soil Biol. Biochem.*, 31, 31-41, doi: 10.1016/S0038-0717(98)00097-2, 1998.
- Crenshaw, C. L., Lauber, C., Sinsabaugh, R. L., and Stavely, L. K.: Fungal control of nitrous oxide production in semiarid grassland, *Biogeochemistry*, 87, 17-27, doi: 10.1007/s10533-007-9165-4, 2008.
- 930 Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, *Quarterly Journal of the Royal Meteorological Society*, 96, 320-325, doi: 10.1002/qj.49709640815, 1970.
- Decock, C., and Six, J.: How reliable is the intramolecular distribution of ¹⁵N in N₂O to source partition N₂O emitted from soil?, *Soil Biol. Biochem.*, 65, 114-127, <http://dx.doi.org/10.1016/j.soilbio.2013.05.012>, 2013.

- 935 Frame, C. H., and Casciotti, K. L.: Biogeochemical controls and isotopic signatures of nitrous oxide production by a marine ammonia-oxidizing bacterium, *Biogeochemistry*, 7, 2695-2709, doi: 10.5194/bg-7-2695-2010, 2010.
- Garber, E. A., and Hollocher, T. C.: ^{15}N , ^{18}O tracer studies on the activation of nitrite by denitrifying bacteria. Nitrite/water-oxygen exchange and nitrosation reactions as indicators of electrophilic catalysis, *J. Biol. Chem.*, 257, 8091-8097, doi: 10.1007/BF00399539, 1982.
- 940 Groffman, P. M., Altabet, M. A., Bohlke, J. K., Butterbach-Bahl, K., David, M. B., Firestone, M. K., Giblin, A. E., Kana, T. M., Nielsen, L. P., and Voytek, M. A.: Methods for measuring denitrification: Diverse approaches to a difficult problem, *Ecological Applications*, 16, 2091-2122, 2006.
- 945 Hayatsu, M., Tago, K., and Saito, M.: Various players in the nitrogen cycle: Diversity and functions of the microorganisms involved in nitrification and denitrification, *Soil Science & Plant Nutrition*, 54, 33-45, <https://doi.org/10.1111/j.1747-0765.2007.00195.x>, 2008.
- 950 Higgins, S. A., Schadt, C. W., Matheny, P. B., and Löffler, F. E.: Phylogenomics reveal the dynamic evolution of fungal nitric oxide reductases and their relationship to secondary metabolism, *Genome Biology and Evolution*, 10, 2474-2489, doi: 10.1093/gbe/evy187, 2018.
- 955 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.
- Jinuntuya-Nortman, M., Sutka, R. L., Ostrom, P. H., Gandhi, H., and Ostrom, N. E.: Isotopologue fractionation during microbial reduction of N_2O within soil mesocosms as a function of water-filled pore space, *Soil Biol. Biochem.*, 40, 2273-2280, doi: 10.1016/j.soilbio.2008.05.016, 2008.
- 960 Keuschnig, C., Gorfer, M., Li, G., Mania, D., Frostegård, Å., Bakken, L., and Larose, C.: NO and N_2O transformations of diverse fungi in hypoxia: evidence for anaerobic respiration only in *Fusarium* strains, *Environmental Microbiology*, 22, 2182-2195, doi: 10.1111/1462-2920.14980, 2020.
- 965 Knowles, R.: Denitrification, *Microbiol. Rev.*, 46, 43-70, 1982.
- Kool, D. M., Wrage, N., Oenema, O., Dolfing, J., and Van Groenigen, J. W.: Oxygen exchange between (de)nitrification intermediates and H_2O and its implications for source determination of NO_3^- and N_2O : a review, *Rapid Commun. Mass Spec.*, 21, 3569-3578, doi: 10.1002/rcm.3249, 2007.
- 970 Kool, D. M., Wrage, N., Oenema, O., Harris, D., and Van Groenigen, J. W.: The ^{18}O signature of biogenic nitrous oxide is determined by O exchange with water, *Rapid Commun. Mass Spec.*, 23, 104-108, 10.1002/rcm.3859, 2009.
- 975 Koranda, M., Kaiser, C., Fuchslueger, L., Kitzler, B., Sessitsch, A., Zechmeister-Boltenstern, S., and Richter, A.: Fungal and bacterial utilization of organic substrates depends on substrate complexity and N availability, *FEMS Microbiology Ecology*, 87, 142-152, doi: 10.1111/1574-6941.12214, 2014.
- 980 Köster, J. R., Well, R., Dittert, K., Giesemann, A., Lewicka-Szczebak, D., Muehling, K.-H., Herrmann, A., Lammel, J., and Senbayram, M.: Soil denitrification potential and its influence on N_2O reduction and N_2O isotopomer ratios, *Rapid Commun. Mass Spec.*, 27, 2363-2373, doi: 10.1002/rcm.6699, 2013a.
- 985 Köster, J. R., Well, R., Tuzson, B., Bol, R., Dittert, K., Giesemann, A., Emmenegger, L., Manninen, A., Cárdenas, L., and Mohn, J.: Novel laser spectroscopic technique for continuous analysis of N_2O isotopomers – application and intercomparison with isotope ratio mass spectrometry, *Rapid Commun. Mass Spec.*, 27, 216-222, doi: 10.1002/rcm.6434, 2013b.
- 990 Ladan, S., and Jacinthe, P.-A.: Evaluation of antibacterial and antifungal compounds for selective inhibition of denitrification in soils, *Environmental Science: Processes & Impacts*, 18, 1519-1529, doi: 10.1039/C6EM00456C, 2016.
- Laughlin, R. J., and Stevens, R. J.: Evidence for fungal dominance of denitrification and codenitrification in a grassland soil, *Soil Sc. Soc. Am. J.*, 66, 1540-1548, doi: 10.2136/sssaj2002.1540, 2002.

Lewicka-Szczebak, D., Well, R., Giesemann, A., Rohe, L., and Wolf, U.: An enhanced technique for automated determination of ^{15}N signatures of N_2 , $(\text{N}_2+\text{N}_2\text{O})$ and N_2O in gas samples, *Rapid Commun. Mass Spec.*, 27, 1548-1558, <https://doi.org/10.1002/rcm.6605>, 2013.

1000 Lewicka-Szczebak, D., Well, R., Köster, J. R., Fuß, R., Senbayram, M., Dittert, K., and Flessa, H.: Experimental determinations of isotopic fractionation factors associated with N_2O production and reduction during denitrification in soils, *Geochim. Cosmochim. Ac.*, 134, 55-73, doi: 10.1016/j.gca.2014.03.010, 2014.

1005 Lewicka-Szczebak, D., Well, R., Bol, R., Gregory, A. S., Matthews, G. P., Misselbrook, T., Whalley, W. R., and Cardenas, L. M.: Isotope fractionation factors controlling isotopocule signatures of soil-emitted N_2O produced by denitrification processes of various rates, *Rapid Commun. Mass Spec.*, 29, 269-282, doi: 10.1002/rcm.7102, 2015.

1010 Lewicka-Szczebak, D., Dyckmans, J., Kaiser, J., Marca, A., Augustin, J., and Well, R.: Oxygen isotope fractionation during N_2O production by soil denitrification, *Biogeosciences*, 13, 1129-1144, doi: 10.5194/bg-13-1129-2016, 2016.

1015 Lewicka-Szczebak, D., Augustin, J., Giesemann, A., and Well, R.: Quantifying N_2O reduction to N_2 based on N_2O isotopocules – validation with independent methods (helium incubation and ^{15}N gas flux method), *Biogeosciences*, 14, 711-732, doi: 10.5194/bg-14-711-2017, 2017.

Lewicka-Szczebak, D., Lewicki, M. P., and Well, R.: N_2O isotope approaches for source partitioning of N_2O production and estimation of N_2O reduction – validation with the ^{15}N gas-flux method in laboratory and field studies, *Biogeosciences*, 17, 5513-5537, 10.5194/bg-17-5513-2020, 2020.

1020 Long, A., Heitman, J., Tobias, C., Philips, R., and Song, B.: Co-occurring anammox, denitrification, and codenitrification in agricultural soils, *Applied and Environmental Microbiology*, 79, 168-176, 10.1128/aem.02520-12, 2013.

1025 Maeda, K., Spor, A., Edel-Hermann, V., Heraud, C., Breuil, M.-C., Bizouard, F., Toyoda, S., Yoshida, N., Steinberg, C., and Philippot, L.: N_2O production, a widespread trait in fungi, *Scientific Reports*, 5, 9697, doi: 10.1038/srep09697, 2015.

1030 McLain, J. E. T., and Martens, D. A.: N_2O production by heterotrophic N transformations in a semiarid soil, *Appl. Soil. Ecol.*, 32, 253-263, 2006.

Müller, C., and Clough, T. J.: Advances in understanding nitrogen flows and transformations: gaps and research pathways, *J. Agric. Sci.*, 152, S34-S44, <https://doi.org/10.1017/s0021859613000610>, 2014.

1035 Nadeem, S., Dorsch, P., and Bakken, L. R.: Autoxidation and acetylene-accelerated oxidation of NO in a 2-phase system: Implications for the expression of denitrification in ex situ experiments, *Soil Biol. Biochem.*, 57, 606-614, doi: 10.1016/j.soilbio.2012.10.007, 2013.

1040 Ostrom, N., and Ostrom, P.: The Isotopomers of Nitrous Oxide: Analytical Considerations and Application to Resolution of Microbial Production Pathways, in: *Handbook of Environmental Isotope Geochemistry*, edited by: Baskaran, M., *Advances in Isotope Geochemistry*, Springer Berlin Heidelberg, 453-476, 2011.

1045 Ostrom, N. E., Pitt, A., Sutka, R., Ostrom, P. H., Grandy, A. S., Huizinga, K. M., and Robertson, G. P.: Isotopologue effects during N_2O reduction in soils and in pure cultures of denitrifiers, *Journal of Geophysical Research-Biogeosciences*, 112, doi: 10.1029/2006jg000287, 2007.

Ostrom, N. E., Sutka, R., Ostrom, P. H., Grandy, A. S., Huizinga, K. M., Gandhi, H., von Fischer, J. C., and Robertson, G. P.: Isotopologue data reveal bacterial denitrification as the primary source of N_2O during a high flux event following cultivation of a native temperate grassland, *Soil Biol. Biochem.*, 42, 499-506, doi: 10.1016/j.soilbio.2009.12.003, 2010.

1050 Ostrom, N. E., and Ostrom, P. H.: Mining the isotopic complexity of nitrous oxide: a review of challenges and opportunities, *Biogeochemistry*, 132, 359-372, doi: 10.1007/s10533-017-0301-5, 2017.

1055 Pan, Y. S., Wu, Y. C., Li, X. Z., Zeng, J., and Lin, X. G.: Continuing Impacts of Selective Inhibition on Bacterial and Fungal Communities in an Agricultural Soil, *Microb. Ecol.*, 78, 927-935, 10.1007/s00248-019-01364-0, 2019.

- Philippot, L., Hallin, S., and Schloter, M.: Ecology of denitrifying prokaryotes in agricultural soil, in: *Advances in Agronomy*, edited by: Donald, L. S., Academic Press, 249-305, 2007.
- 1060 Popp, B. N., Westley, M. B., Toyoda, S., Miwa, T., Dore, J. E., Yoshida, N., Rust, T. M., Sansone, F. J., Russ, M. E., Ostrom, N. E., and Ostrom, P. H.: Nitrogen and oxygen isotopomeric constraints on the origins and sea-to-air flux of N₂O in the oligotrophic subtropical North Pacific gyre, *Global Biogeochem. Cy.*, 16, doi: 10.1029/2001gb001806, 2002.
- 1065 Reischke, S., Rousk, J., and Bååth, E.: The effects of glucose loading rates on bacterial and fungal growth in soil, *Soil Biol. Biochem.*, 70, 88-95, doi: 10.1016/j.soilbio.2013.12.011, 2014.
- 1070 Rohe, L., Anderson, T.-H., Braker, G., Flessa, H., Giesemann, A., Lewicka-Szczebak, D., Wrage-Mönnig, N., and Well, R.: Dual isotope and isotopomer signatures of nitrous oxide from fungal denitrification – a pure culture study, *Rapid Communications in Mass Spectrometry* 28, 1893-1903, doi: 10.1002/rcm.6975, 2014a.
- 1075 Rohe, L., Anderson, T.-H., Braker, G., Flessa, H., Giesemann, A., Wrage-Mönnig, N., and Well, R.: Fungal oxygen exchange between denitrification intermediates and water, *Rapid Commun. Mass Spec.*, 28, 377-384, doi: 10.1002/rcm.6790, 2014b.
- 1080 Rohe, L., Well, R., and Lewicka-Szczebak, D.: Use of oxygen isotopes to differentiate between nitrous oxide produced by fungi or bacteria during denitrification, *Rapid Commun. Mass Spec.*, 31, 1297-1312, doi: 10.1002/rcm.7909, 2017.
- 1085 Ruzicka, S., Edgerton, D., Norman, M., and Hill, T.: The utility of ergosterol as a bioindicator of fungi in temperate soils, *Soil Biol. Biochem.*, 32, 989-1005, doi: 10.1016/S0038-0717(00)00009-2, 2000.
- 1090 Senbayram, M., Well, R., Bol, R., Chadwick, D. R., Jones, D. L., and Wu, D.: Interaction of straw amendment and soil NO₃⁻ content controls fungal denitrification and denitrification product stoichiometry in a sandy soil, *Soil Biol. Biochem.*, 126, 204-212, doi: 10.1016/j.soilbio.2018.09.005, 2018.
- 1095 Senbayram, M., Well, R., Shan, J., Bol, R., Burkart, S., Jones, D. L., and Wu, D.: Rhizosphere processes in nitrate-rich barley soil tripled both N₂O and N₂ losses due to enhanced bacterial and fungal denitrification, *Plant and Soil*, 448, 509-522, doi: 10.1007/s11104-020-04457-9, 2020.
- 1100 Seo, D. C., and DeLaune, R. D.: Fungal and bacterial mediated denitrification in wetlands: Influence of sediment redox condition, *Water Research*, 44, 2441-2450, doi: 10.1016/j.watres.2010.01.006, 2010.
- 1105 Shoun, H., Kim, D.-H., Uchiyama, H., and Sugiyama, J.: Denitrification by fungi, *FEMS Microbiol. Lett.*, 94, 277-281, 1992.
- 1110 Shoun, H., Fushinobu, S., Jiang, L., Kim, S. W., and Wakagi, T.: Fungal denitrification and nitric oxide reductase cytochrome P450nor, *Philos. T. Roy. Soc.*, 367, 1186-1194, doi: 10.1098/rstb.2011.0335, 2012.
- 1115 Snider, D. M., Schiff, S. L., and Spoelstra, J.: ¹⁵N/¹⁴N and ¹⁸O/¹⁶O stable isotope ratios of nitrous oxide produced during denitrification in temperate forest soils, *Geochim. Cosmochim. Ac.*, 73, 877-888, doi: 10.1016/j.gca.2008.11.004, 2009.
- 1120 Spott, O., Russow, R., Apelt, B., and Stange, C. F.: A ¹⁵N-aided artificial atmosphere gas flow technique for online determination of soil N₂ release using the zeolite Kőstrolith SX6®, *Rapid Commun. Mass Spec.*, 20, 3267-3274, <https://doi.org/10.1002/rcm.2722>, 2006.
- 1125 Spott, O., Russow, R., and Stange, C. F.: Formation of hybrid N₂O and hybrid N₂ due to codenitrification: First review of a barely considered process of microbially mediated N-nitrosation, *Soil Biol. Biochem.*, 43, 1995-2011, doi: 10.1016/j.soilbio.2011.06.014, 2011.
- 1130 Stumpp, C., Klaus, J., and Stichler, W.: Analysis of long-term stable isotopic composition in German precipitation, *Journal of Hydrology*, 517, 351-361, doi: 10.1016/j.jhydrol.2014.05.034, 2014.

- 1115 Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Breznak, J. A., Gandhi, H., Pitt, A. J., and Li, F.: Distinguishing nitrous oxide production from nitrification and denitrification on the basis of isotopomer abundances, *Appl. Environ. Microbiol.*, 72, 638-644, doi: 10.1128/aem.72.1.638-644.2006, 2006.
- 1120 Sutka, R. L., Adams, G. C., Ostrom, N. E., and Ostrom, P. H.: Isotopologue fractionation during N₂O production by fungal denitrification, *Rapid Commun. Mass Spec.*, 22, 3989-3996, doi: 10.1002/rcm.3820, 2008.
- Tanimoto, T., Hatano, K., Kim, D. H., Uchiyama, H., and Shoun, H.: Co-denitrification by the denitrifying system of fungus *Fusarium oxysporum*, *FEMS Microbiol. Lett.*, 93, 177-180, 1992.
- 1125 Toyoda, S., and Yoshida, N.: Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer, *Anal. Chem.*, 71, 4711-4718, 10.1021/ac9904563, 1999.
- 1130 Toyoda, S., Mutobe, H., Yamagishi, H., Yoshida, N., and Tanji, Y.: Fractionation of N₂O isotopomers during production by denitrifier, *Soil Biol. Biochem.*, 37, 1535-1545, doi: 10.1016/j.soilbio.2005.01.009, 2005.
- 1135 Toyoda, S., Yano, M., Nishimura, S.-i., Akiyama, H., Hayakawa, A., Koba, K., Sudo, S., Yagi, K., Makabe, A., Tobari, Y., Ogawa, N. O., Ohkouchi, N., Yamada, K., and Yoshida, N.: Characterization and production and consumption processes of N₂O emitted from temperate agricultural soils determined via isotopomer ratio analysis, *Global Biogeochem. Cy.*, 25, doi: 10.1029/2009gb003769, 2011.
- USEPA: Manuel: Nitrogen Control, Office of Water, Washington, D.C. , EPA/625/R-93 /010, 1993.
- 1140 Well, R., Becker, K.-W., Meyer, B., Langel, R., and Reineking, A.: Continuous flow equilibration for mass spectrometric analysis of dinitrogen emissions, *Soil Sc. Soc. Am. J.*, 62, 906-910, doi: 10.2136/sssaj1998.03615995006200040008x, 1998.
- 1145 Well, R., Kurganova, I., de Gerenyu, V. L., and Flessa, H.: Isotopomer signatures of soil-emitted N₂O under different moisture conditions - A microcosm study with arable loess soil, *Soil Biol. Biochem.*, 38, 2923-2933, 10.1016/j.soilbio.2006.05.003, 2006.
- Well, R., and Flessa, H.: Isotopologue signatures of N₂O produced by denitrification in soils, *Journal of Geophysical Research: Biogeosciences*, 114, G02020, doi: 10.1029/2008jg000804, 2009.
- 1150 Well, R., Burkart, S., Giesemann, A., Grosz, B., Köster, J. R., and Lewicka-Szczebak, D.: Improvement of the ¹⁵N gas flux method for in situ measurement of soil denitrification and its product stoichiometry, *Rapid Commun. Mass Spec.*, 33, 437-448, <https://doi.org/10.1002/rcm.8363>, 2019.
- 1155 Wrage-Mönnig, N., Horn, M. A., Well, R., Müller, C., Velthof, G., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited, *Soil Biol. Biochem.*, 123, A3-A16, doi: 10.1016/j.soilbio.2018.03.020, 2018.
- 1160 Wrage, N., Velthof, G. L., Laanbroek, H. J., and Oenema, O.: Nitrous oxide production in grassland soils: assessing the contribution of nitrifier denitrification, *Soil Biol. Biochem.*, 36, 229-236, 10.1016/j.soilbio.2003.09.009, 2004a.
- Wrage, N., Velthof, G. L., Oenema, O., and Laanbroek, H. J.: Acetylene and oxygen as inhibitors of nitrous oxide production in *Nitrosomonas europaea* and *Nitrosospira briensis*: a cautionary tale, *Fems Microbiology Ecology*, 47, 13-18, 10.1016/s0168-6496(03)00220-4, 2004b.
- 1165 Wu, D., Well, R., Cárdenas, L. M., Fuß, R., Lewicka-Szczebak, D., Köster, J. R., Brüggemann, N., and Bol, R.: Quantifying N₂O reduction to N₂ during denitrification in soils via isotopic mapping approach: Model evaluation and uncertainty analysis, *Environmental Research*, 179, 108806, <https://doi.org/10.1016/j.envres.2019.108806>, 2019.
- 1170 Yoshinari, T., and Knowles, R.: Acetylene inhibition of nitrous-oxide reduction by denitrifying bacteria, *Biochemical and Biophysical Research Communications*, 69, 705-710, doi: 10.1016/0006-291x(76)90932-3, 1976.
- 1175 Yu, L., Harris, E., Lewicka-Szczebak, D., Barthel, M., Blomberg, M. R. A., Harris, S. J., Johnson, M. S., Lehmann, M. F., Liisberg, J., Müller, C., Ostrom, N. E., Six, J., Toyoda, S., Yoshida, N., and Mohn, J.: What can we learn

from N₂O isotope data? - Analytics, processes and modelling, Rapid Commun Mass Spectrom, doi: 10.1002/rcm.8858, 2020.

1180 Zou, Y., Hirono, Y., Yanai, Y., Hattori, S., Toyoda, S., and Yoshida, N.: Isotopomer analysis of nitrous oxide accumulated in soil cultivated with tea (*Camellia sinensis*) in Shizuoka, central Japan, Soil Biol. Biochem., 77, 276-291, doi: 10.1016/j.soilbio.2014.06.016, 2014.