

Denitrification by benthic foraminifera and their contribution to N-loss from a fjord environment

Constance Choquel^{1*}, Emmanuelle Geslin¹, Edouard Metzger¹, Helena L. Filipsson², Nils Risgaard-Petersen³, Patrick Launeau¹, Manuel Giraud¹, Thierry Jauffrais^{4,1}, Bruno Jesus^{5,6} and Aurélia Mouret¹

1: UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France

2: Department of Geology, Lund University, Sweden

3: Department of Biology, Aquatic Biology, Aarhus University, Denmark

4: Ifremer, IRD, Univ. Nouvelle-Calédonie, Univ. La Réunion, CNRS, UMR 9220 ENTROPIE, New Caledonia

5: Université de Nantes, Mer Molécules Santé, EA 2160, France

6: BioISI – Biosystems & Integrative Sciences Institute, Campo Grande, University of Lisbon, Faculty of Sciences, Portugal

Correspondence to: Constance Choquel (constance.choquel@gmail.com or constance.choquel@univ-angers.fr)

Abstract.

Oxygen and nitrate availabilities impact the marine nitrogen cycle at a range of spatial and temporal scales. Here, we demonstrate the impact of denitrifying foraminifera on the nitrogen cycle at two oxygen and nitrate contrasting stations in a fjord environment (Gullmar Fjord, Sweden). The foraminifera contribution to benthic denitrification was estimated by coupling living foraminifera microhabitat, denitrification rate measurement and sedimentary nitrate 2D distribution, combining diffusive equilibrium in thin films (DET) colorimetry and hyperspectral imagery. Oxygenated bottom waters with high nitrate content in sediment porewaters were dominated by the non-indigenous species (NIS) *Nonionella* sp. T1 which could denitrify up to 50-100 % of nitrate porewater. Contrastingly, hypoxic bottom waters where sediment porewaters were nitrate low, denitrifying foraminifera were scarce and did not contribute to nitrogen removal (~ 5 %). Our study showed that benthic foraminifera can be a major contributor of nitrogen mitigation in oxic coastal ecosystems and should be included in ecological and diagenetic models aiming at understanding biogeochemical cycles coupled to nitrogen.

1 Introduction

Hypoxic water occurs frequently in bottom-waters of shallow coastal seas, due to remineralization of organic matter and water stratification. In this study we used the hypoxia threshold of $63 \mu\text{mol L}^{-1}$ (e.g. Diaz et al., 2008; Breitburg et al., 2018). Hypoxia may have large ecological effects (Levin et al., 2009; Rabalais et al., 2010; Zhang et al., 2010), such as an increase of fauna mortality (Diaz et al., 2001). However, certain microorganisms, e.g. bacteria and foraminifera, can perform denitrification by respiring nitrate (Risgaard-Petersen et al., 2006) and thereby survive in depleted oxygen environments. The effects of decreasing dissolved oxygen availability at spatial and temporal scales will impact biogeochemical cycles such as the nitrogen cycle (Childs et al., 2002; Kemp et al., 2005; Conley et al., 2007; Diaz et al., 2008; Neubacher et al., 2013; Breitburg et al., 2018). The nitrogen cycle in marine sediments is a perpetual balance between nitrogen inputs (e.g. terrestrial

runoff, atmospheric precipitations) and outputs (e.g. denitrification from sediment and water column) (Galloway et al., 2004;

35 Sigman et al., 2009). In most semi-enclosed marine environments as the Baltic Sea, the nitrogen loss through benthic denitrification exceeds the inputs of nitrogen through nitrogen fixation. These sink regions of the ocean are mostly associated with anoxic regions (Gruber and Sarmiento 1997). This study focuses on how one important compartment of the marine meiofaunal community - the benthic foraminifera - is coupled to the nitrogen cycle during contrasted dissolved [O₂] conditions at two different stations, focusing on the impact of a non-indigenous species (NIS).

40

The nitrogen cycle occurring in marine sediments is dependent on the bottom-water oxygenation. In oxic bottom water conditions (Fig. 1a), ammonium (NH₄⁺) produced from remineralization of particulate organic nitrogen (PON) in sediments, diffuses toward the oxic sediment-superficial layer and through the water-sediment interface. Nitrification can occur in the oxic sediment and in the oxic water column through the conversion of NH₄⁺ to nitrate (NO₃⁻) (Rysgaard et al., 1994; 45 Thamdrup and Dalsgaard, 2008). Conversely, denitrification occurs in sediment when oxygen is scarce (below 5 μmol L⁻¹, Devol et al., 2008) and organic carbon and nitrate are available. Denitrification named “canonical denitrification” (NO₃⁻ → NO₂⁻ → NO → N₂O → N₂) is an anaerobic process whereby nitrate is used as the terminal electron acceptor in the oxidation of organic matter by facultative anaerobic metabolisms when oxygen is exhausted. Denitrification participates in the loss of the fixed Nitrogen to N₂ gas (Brandes et al., 2007 and references within). Another process can contribute to this loss of N₂ gas: 50 Anammox (anaerobic ammonia oxidation) (Engström et al., 2005; Brandma et al., 2011). According to Brandes et al. (2007 and references within) the “total denitrification” can be defined as the sum of the canonical denitrification plus the anammox. Nitrification and denitrification are thus strongly coupled, and denitrification can be enhanced by adjacent sedimentary nitrification zones or by direct NO₃⁻ diffusion from the overlying water towards the sediment (Kemp et al., 1990; Cornwell et al., 1999). When bottom water turns hypoxic, the nitrogen cycle occurring in the sediment is strongly affected (Fig. 1 b). 55 Nitrate production is reduced since nitrification cannot process under low oxygen conditions (~ 0 μmol L⁻¹; Rysgaard et al., 1994; Mortimer et al., 2004). However, deeper into reduced sediment, nitrification can occur through secondary reactions with NH₄⁺ oxidation by Mn and Fe oxides (Luther et al., 1997; Mortimer et al., 2004). Denitrification is the dominant process of nitrate reduction in coastal marine sediments (Thamdrup and Dalsgaard, 2008; Herbert, 1999). However, dissimilatory nitrate

reduction to ammonium (DNRA) can also contribute to nitrate depletion in reduced sediment leading to NO_3^- conversion into

60 NH_4^+ instead of nitrogen (N_2) (Christensen et al., 2000) and compete denitrification.

Benthic foraminifera were the first marine eukaryotes found to perform denitrification (Risgaard-Petersen et al., 2006), but not all foraminifera species can denitrify (Piña-Ochoa et al., 2010). Denitrifying foraminifera species are defined in our study as species able to perform denitrification proved by denitrification rate measurements. These denitrifying species
65 have a facultative anaerobic metabolism and nitrate-storing foraminifera can use either environmental oxygen or nitrate to respire (Piña-Ochoa et al., 2010). *Nonionella cf. stella* (Charrieau et al., 2019 and references therein) and *Globobulimina turgida* were identified as the first denitrifying foraminifera species (Risgaard-Petersen et al., 2006). Currently, nineteen denitrifying species are known (Glock et al., 2019). Foraminifera denitrification rates show a large range from $7 \pm 1 \text{ pmol N indiv.}^{-1} \text{ d}^{-1}$ to $2241 \pm 1825 \text{ pmol N indiv.}^{-1} \text{ d}^{-1}$ (Glock et al., 2019).

70 Recently, *Nonionella stella* was described as invasive in the North Sea region and reported in the Gullmar Fjord (Sweden) (< 5 %, Polovodova Asteman and Schönfeld, 2015). However, *Nonionella stella* sampled in the Santa Barbara Basin (California USA) differs morphologically (Charrieau et al., 2018) and genetically (Deldicq et al., 2019) from the specimens sampled in Kattegat and Oslofjord (Norway), respectively. Deldicq et al. (2019) describe these specimens as the *Nonionella* sp. T1 morphotype, a non-indigenous and suspected invasive species in the Oslofjord. The genus *Nonionella* is potentially
75 capable to denitrify as demonstrated with *Nonionella cf. stella* by Risgaard-Petersen et al. (2006). Denitrification rates of two species from the Gullmar Fjord have been measured: *Globobulimina turgida* (Risgaard-Petersen et al., 2006) and *Globobulimina auriculata* (Woehle et al., 2018). Additionally, *Stainforthia fusiformis* and *Bolivina pseudopunctata* are two dominant species in the deepest part of the fjord (Gustafsson and Nordberg, 2001; Filipsson and Nordberg, 2004). These species are also potential candidates for denitrification. Indeed, the denitrification rates of *Stainforthia fusiformis* from Perú
80 were measured by Piña-Ochoa et al. (2010) and several species of *Bolivina* from Perú, Bay of Biscay and Santa Barbara were measured by Glock et al. (2019); Piña-Ochoa et al. (2010) and Bernhard et al. (2012), respectively. On the other hand, other typical fjord species such as *Bulimina marginata*, *Cassidulina laevigata*, *Hyalinea balthica* are considered as non-denitrifying species by Piña-Ochoa et al. (2010) as their intracellular nitrate reserves are almost absent. The anaerobic metabolism of some

other species commonly found in the fjord such as *Leptohalysis scotti*, *Liebusella goesi*, *Nonionellina labradorica* and

85 *Textularia earlandi* is not documented in previous studies.

A high abundance of denitrifying foraminifera in both oxic and anoxic marine environments play an important role in the nitrogen cycle (Risgaard-Petersen et al., 2006; Piña-Ochoa et al., 2010; Bernhard et al., 2012; Glock et al., 2013; Xu et al., 2017). Previous estimates of foraminifera contributions to denitrification range from 1 to 90 % (Dale et al., 2016; Xu et al., 2017). Estimates of foraminifera contribution to benthic denitrification are limited by the high spatial and temporal 90 variability of sediment geochemistry and distribution of denitrifying foraminifera, which poses particular methodological challenges. Marine sediments often include chemical micro-heterogeneities (Aller et al., 1998; Stockdale et al., 2009), which can be averaged within the volume of a sediment slice. Moreover, sediment core slicing or centrifugation can induce cell lysis, which can induce a bias in porewater nitrate concentrations (Risgaard-Petersen et al., 2006). To characterize these microenvironments at submillimeter/ millimeter scales, new approaches have to be used. Recently, a 2D-DET (two Dimensions 95 Diffusive Equilibrium in Thin-film) technique combining colorimetry and hyperspectral imagery was developed to obtain the distribution of nitrite and nitrate in sediment porewater at millimeter resolution in two dimensions (Metzger et al., 2016). This method avoids mixing of intracellular nitrate and nitrate contained in the sediment porewater.

The present study aims to examine how the NIS *Nonionella* sp. T1 and the other denitrifying species affect the 100 nitrogen cycle by comparing two stations with contrasting oxygen and nitrate environments subjected to hypoxic events. The objectives of the paper are: (1) to characterize the density of the living benthic foraminifera at two contrasted stations; (2) to measure the denitrification rate of the NIS *Nonionella* sp. T1 and (3) to quantify its contributions to benthic denitrification; (4) to discuss the probable future impact of the NIS *Nonionella* sp. T1 on the foraminifera fauna and the nitrogen cycle in the Gullmar Fjord.

105

2 Material and Methods

2.1 Site description and sampling conditions

The Gullmar Fjord is 28 km long, 1-2 km wide and located on the Swedish West coast (Fig. 2). The fjord undergoes fluctuations between cold and temperate climates (Svansson, 1975; Nordberg, 1991; Polovodova Asteman and Nordberg, 110 Nordberg, 2013; Polovodova Asteman et al., 2018). The fjord is stratified (Fig. 2 d) in four water masses (Svansson, 1984; Arneborg, 2004). Hypoxia events in the fjord have been linked to the influence of the North Atlantic Oscillation (NAO) (Nordberg et al., 2000; Björk and Nordberg, 2003; Filipsson and Nordberg, 2004). Several monitoring stations are located in the fjord: Släggö (65 m depth), Björkholmen (70 m depth) and Alsbäck (117 m depth), the hydrographic and nutrient data were obtained from the SMHI's publically available data-base SHARK (Svenskt Havsarkiv, www.smhi.se). Since 2010, the threshold of hypoxia 115 ($[O_2] < 2 \text{ mg L}^{-1}$, i.e. $63 \mu\text{mol L}^{-1}$) in Alsbäck station (red squares, Fig. 3) is reached typically in late autumn and winter. Deep-water exchanges usually occur in late water-early spring. However, the duration of hypoxia varies between years and hypoxia events also occurred in the summer 2014 and 2015, due to lack of deep-water exchange. The frequency of hypoxic events has increased in the fjord (Nordberg et al., 2000; Filipsson and Nordberg, 2004).

Two sampling cruises were conducted in the Gullmar Fjord on board R/V *Skagerak* and *Oscar von Sydow*, 120 respectively. The 2017 cruise (GF17) took place between 14th and 15th November 2017 and two stations were sampled (GF17-3 and GF17-1, Fig. 2 c and d) to define the living foraminifera fauna and the sediment geochemistry at two contrasted stations. The 2018 cruise (GF18) took place on the 5th September 2018 with the focus to collect living *Nonionella* sp. T1 for O_2 respiration and denitrification rates measurements. Only one station (at the same position as GF17-3) was sampled.

GF17-3 (50 m water depth) is located closest to the mouth of the fjord ($58^{\circ}16'50.94''\text{N}/11^{\circ}30'30.96''\text{E}$) with bottom 125 waters from Skagerrak (blue diamond, Fig. 3) and GF17-1 (117 m depth) close to the deepest part of the fjord ($58^{\circ}19'41.40''\text{N}/11^{\circ}33'8.40''\text{E}$) near Alsbäck monitoring station in the middle of the stagnant basin (red square, Fig. 3). In November 2017, CTD profiles indicated the water mass structures at both stations (Fig. S1). Bottom water at GF17-3 station was oxic with a dissolved oxygen content of $234 \mu\text{mol L}^{-1}$. The dissolved oxygen content decreased strongly with depth at the GF17-1 station reaching $9 \mu\text{mol L}^{-1}$ at the seafloor, which is below the severe hypoxia threshold.

130

2.2 Foraminifera sampling and processing

During the 2017 cruise, two sediment cores per station (1A, 1C and 3A, 3C for GF17-1 and GF17-3 stations respectively) were immediately subsampled with a smaller cylindrical core (\varnothing 8.2 cm) and sliced every 2 mm up to 2 cm and every 5 mm from 2 to 5 cm to study living foraminifera distribution. The samples were incubated without light for 10–19 hours in ambient
135 seawater with Cell Tracker Green (CMFDA, 1 mM final concentration) at *in situ* temperatures (Bernhard et al., 2006) and then fixed with ethanol 96°. Fixed samples were sieved (>355, 150, 125 and 100 μm) and the >100 μm fraction, the most commonly fraction used for foraminiferal analyses in the Gullmar Fjord (see Charrieau et al., 2018 and references therein) was examined using an epifluorescence microscope equipped for fluorescein detection (i.e., 470 nm excitation; Olympus SZX13). In the present study, the foraminifera distribution will be described highlighting the NIS *Nonionella* sp. T1.

140

2.3 Geochemical sampling and processing

One core from the shallow GF17-3 station was reserved for O₂ microelectrode profiling. Oxygen concentration was measured in the dark with a Clark electrode (50 μm tip diameter, Unisense ®, Denmark) within the first 5 mm depth at a 100 μm vertical resolution. Due to technical problems, no oxygen profiling was done at the GF17-1 station.

145

One core per station was dedicated for geochemical analyses, they were carefully brought to Lund University (Sweden) and stored at the sampling site temperature (10°C) until further analysis the next day. Overlaying water of the GF17-3 core was gently air bubbled to maintain the oxygenated conditions recorded at this station. Overlaying water of the GF17-1 core was bubbled with N₂ gas passed through a solution of carbonate/bicarbonate to avoid pH rise due to degassing of CO₂ by N₂ bubbling.

150

Nitrite/Nitrate were analyzed using the 2D-DET method from Metzger et al. (2016). In brief, for each core, a DET (Diffusive Equilibrium in Thin films) gel probe (16 cm x 6.5 cm and 0.1 cm thickness) was hand-made prepared. The gel probe was inserted into the sediment and left for 5 hours to allow for a diffusive equilibration time between the gel and porewaters; After equilibration, the gel was removed of the core and laid on a first NO₂⁻ reagent gel. After 15 mn at ambient temperature the pink coloration must appear were nitrite is detected. A reflectance analysis photograph of the nitrite gels fauna was taken
155 with a hyperspectral camera (HySpex VNIR 1600). The next step was to convert existing nitrate into nitrite with the addition of a reagent gel of vanadium chloride (VCl₃). After 20 min at 50°C, additional pink is interpreted as porewater nitrate

concentration. Followed by the acquisition of another hyperspectral image and converted into false colours through a calibrated scale of concentrations, the final image was cropped to avoid border effects. Each pixel (190 μm x 190 μm) was decomposed as a linear combination of the logarithm of the different end-member spectra using ENVI software (unmixing function)
160 (Cesbron et al., 2014; Metzger et al., 2016). Nitrite and nitrate detection limit is 1.7 $\mu\text{mol L}^{-1}$ (Metzger et al., 2016). Nitrate production/consumption zones for each station were estimated by extracting the average and standard deviation of the 290 vertical 1D profiles ((5.5 cm width x 1 pixel) / 0.019 cm for 1-pixel size) on the 2D gels and modelling using PROFILE software (Berg et al., 1998)).

165 **2.4 Oxygen respiration and denitrification rates measurements of the NIS *Nonionella* sp. T1**

The two cores sampled in the 2018 cruise (GF18) at the shallower GF17-3 station were carefully transported at *in situ* temperature (8 °C) and stored for three days at the Department of Geosciences, Aarhus University (Denmark). *Nonionella* sp. T1 specimens were picked under *in situ* temperature and collected in a Petri dish, containing a thin layer of sediment (32 μm) to check their vitality. Only living, active *Nonionella* sp. T1 specimens were picked and cleaned several times using a brush
170 with micro-filtered, nitrate-free artificial seawater.

Oxygen respiration rates were measured, following the method developed by Høglund et al. (2008) using a Clark type oxygen microsensors (50 μm tip diameter, Unisense ®, Denmark) (Revsbech, 1989) calibrated by a two-point calibration using air-saturated water at *in situ* temperature (8 °C) and sodium ascorbate solution (to strip O₂ out of the system) as zero. Then, a pool of 5 living *Nonionella* sp. T1 was transferred into a glass microtube (inner diameter 0.5 mm, height 7.5 mm) that was
175 fixed inside a 20 ml test tube mounted in a glass-cooling bath (8 °C). A motorized micromanipulator was used to measure O₂ concentration profiles along a distance gradient that ranged from 200 μm of the foraminifera to 1200 μm using 100 μm steps. Seven O₂ concentration profiles were generated with one incubation containing the pool of *Nonionella* sp. T1. Negative controls were done by measuring O₂ rates from microtube with empty foraminifera shells and blanks with empty microtube. Oxygen respiration rates were calculated with Fick's first law of diffusion, $J = -D * \frac{dC}{dx}$, where J is the flux, dC/dx is the
180 concentration gradient obtained by profiles and D is the free diffusion coefficient of oxygen at 8 °C for a salinity of 34 (1.382 x 10⁻⁵ cm⁻² s⁻¹, Ramsing and Gundersen, 1994). The seven O₂ respiration rates were calculated as the product of the flux by

the cross section area of the microtube (0.196 mm^2). Then, the average O_2 respiration rate was divided by the 5 *Nonionella* sp. T1 presented in the microtube to obtain the respiration rate per individual.

185 The same pool of *Nonionella* sp. T1 specimens as for the O_2 respiration rates was used for denitrification rate measurements. Denitrification rates were measured as it is described in Risgaard-Petersen et al., (2006). In this method, denitrification is stopped at the N_2O production by acetylene inhibition that can be measured with a N_2O microprobe ($50 \mu\text{m}$ tip diameter, Unisense ®, Denmark). Thus, N_2O was measured as the end product instead of N_2 (Risgaard-Petersen et al., 2006).

190 Nitrous oxide flux was estimated from the chemical gradient profiled from the pool of *Nonionella* sp. T1 inserted in a microchamber. The N_2O production was multiplied by two because two moles of NO_3^- are required for the production of one mole of N_2O (Risgaard-Petersen et al., 2006). The microchamber is porous to gases and is bathed in a sodium ascorbate solution that maintains oxygen concentration at zero within the microchamber. The microchamber was filled with an oxygen/nitrate-free solution of artificial seawater saturated with acetylene (to inhibit N_2O transformation into N_2) containing 5 mM of Hepes
195 buffer (to maintain the pH stable). Calibration was performed using the standard addition method by successive injections of a N_2O saturated solution in order to have $14 \mu\text{M}$ steps of final concentration. Negative controls were done by checking the absence of O_2 from microchamber with empty foraminifera shells and blanks with empty microchamber. Then, the pool of
Nonionella sp. T1, was transferred to the microchamber with a micropipette. The N_2O concentration profiles were repeated seven times on the pool of *Nonionella* sp. T1. The source of nitrate during denitrification comes from intracellular nitrate
200 storage of *Nonionella* sp. T1 (not measured in this study).

Since O_2 respiration and denitrification rates are linked to cytoplasmic volume or biovolume (BV) (Geslin et al., 2011; Glock et al., 2019), the specimens from the pool of *Nonionella* sp. T1 were measured (width (a) and length (b) Fig. 4) using a micrometer mounted on a Leica stereomicroscope (MZ 12.5) to estimate the average BV. The volume of the shells was estimated by using the best resembling geometric shape, a spheroid prolate ($V = \frac{4}{3}\pi \left(\frac{a}{2}\right)^2 \left(\frac{b}{2}\right)$). Then, according to Hannah et
205 al., (1994) 75 % of the measured entire volume of the shell was used corresponding to the estimated cytoplasmic volume. To compare the size of the *Nonionella* sp. T1 sampled in the 2017 cruise (GF17, study of the fauna) with the *Nonionella* sp. T1

samples in the 2018 cruise (GF18, denitrification rate measurements), 5 specimens sampled in the 2017 cruise were also measured.

210 2.5 Contributions of the NIS *Nonionella* sp. T1 to diffusive oxygen and nitrate uptake

The following estimated contributions to sediment diffusive oxygen and nitrate uptake were performed mainly on the dominant denitrifying species, *Nonionella* sp. T1. The size of the *Nonionella* sp. T1 specimens sampled during the two cruises differed markedly (Table 1). Thus, we need to correct the denitrification rate of *Nonionella* sp. T1 specimens from the 2017 cruise to take into account the difference of shell size. Thus, the measured *Nonionella* sp. T1 denitrification rate (2018 cruise) 215 was normalized by specimen BV (2017 cruise) using the relationship: $\ln(y) = 0.68 \ln(x) - 5.57$, where y is the denitrification rate ($\text{pmol ind}^{-1} \text{d}^{-1}$) and x is the shell BV (μm^3) (Geslin et al., 2011; Glock et al., 2019; Equation S1). The corrected *Nonionella* sp. T1 denitrification rate is multiplied by the *Nonionella* sp. T1 specimens counted found in each denitrifying zones defined by PROFILE modelling. Then, two calculation approaches were discussed to estimate *Nonionella* sp. T1 contributions to benthic denitrification: (A) to divide the *Nonionella* sp. T1 denitrification rate by the nitrate porewater denitrification rate 220 estimated from PROFILE modelling, then the second calculation (B) to divide the *Nonionella* sp. T1 denitrification rate by the total denitrification from PROFILE plus the *Nonionella* sp. T1 denitrification rate. In the first approach (A) we suggest *Nonionella* sp. T1 use only the nitrate in the sediment porewater. In the second approach (B) we suggest that the foraminifera use both intracellular and porewater nitrate pools for denitrification.

225 3 Results

3.1 The NIS *Nonionella* sp. T1 oxygen respiration and denitrification rates in the Gullmar Fjord

The O_2 respiration rates measured in the pool of *Nonionella* sp. T1 specimens collected in the 2018 cruise (GF18) were $169 \pm 11 \text{ pmol O}_2 \text{ indiv}^{-1} \text{ d}^{-1}$ with an average BV of $1.3 \pm 0.7 \cdot 10^{+06} \mu\text{m}^3$ (BV details, Table 1). The denitrification rate, measured on the same pool of specimens, was $21 \pm 9 \text{ pmol N indiv}^{-1} \text{ d}^{-1}$.

230 The *Nonionella* sp. T1 average BV collected in the 2017 cruise (GF17-3) was $4.0 \pm 0.6 \text{ } 10^{+06} \mu\text{m}^3$, i.e. more than three times larger the *Nonionella* sp. T1 average BV from the 2018 cruise ($1.3 \pm 0.7 \text{ } 10^{+06} \mu\text{m}^3$). As denitrification rates and foraminifera BV are linked (see method), the measured denitrification rate was corrected using the BV of *Nonionella* sp. T1 from the 2017 cruise. Thus, the *Nonionella* sp. T1 corrected denitrification rate was $38 \pm 8 \text{ pmol N indiv}^{-1} \text{ d}^{-1}$ (Equation S1).

235 **3.2 The NIS *Nonionella* sp. T1 and foraminifera fauna regarding porewater nitrate micro-distribution**

The bottom water at GF17-3 station was oxic (Fig. S1, $[\text{O}_2] = 234 \mu\text{mol L}^{-1}$) and the measured oxygen penetration depth (OPD) in the sediment was $4.7 \pm 0.2 \text{ mm}$ ($n = 3$). No nitrite was revealed on the gel ($< 1.7 \mu\text{mol L}^{-1}$), only nitrate was detected. Bottom water average NO_3^- concentration was $14.6 \pm 2.3 \mu\text{mol L}^{-1}$ and nitrate concentration decreased with depth in the sediment (Fig. 5 c, d). Nitrate concentration ranged between 13.1 ± 3.2 to $11.7 \pm 3.4 \mu\text{mol L}^{-1}$, from the water-sediment 240 interface to the OPD. Nitrate concentration decreased strongly after the OPD from 11.7 ± 3.4 to $2.8 \pm 0.9 \mu\text{mol L}^{-1}$ until 4.0 cm depth. From 4.0 to 5.0 cm depth, NO_3^- concentration was very low with an average value of $2.7 \pm 0.9 \mu\text{mol L}^{-1}$ (Fig. 5 c, d). The PROFILE parameters (Berg et al., 1998) used on laterally averaged nitrate porewater vertical distribution of both stations are available in Table S1. Thus, the PROFILE modelling of the averaged nitrate porewater profiles revealed one nitrification zone from 0 to 1.2 cm depth and two denitrifying zones (red line, Fig. 5 d). The first denitrification zone occurred 245 between 1.2 to 3.6 cm depth with a nitrate consumption of $3.92 \text{ E}^{-05} \text{ nmol cm}^{-3} \text{ s}^{-1}$ and the second smaller consumption zone was from 3.6 to 5 cm depth ($1.53 \text{ E}^{-06} \text{ nmol cm}^{-3} \text{ s}^{-1}$). The total denitrification rate from 1.2 to 5 cm depth was $4.07 \text{ E}^{-05} \text{ nmol cm}^{-3} \text{ s}^{-1}$ (Fig. 5 d).

The total densities of living foraminifera were similar between the cores GF17-3A and 3C ($\varnothing 8.2 \text{ cm}$, 5 cm depth) with 1256 individuals and 1428 individuals, respectively (Fig. 5 a and b; Table S2, GF17-3A and 3C). *Nonionella* sp. T1 was the 250 main denitrifying species, accounting for 34 % of the total living fauna in the core GF17-3A and 74 % in GF17-3C (Fig. 5 a, b; Table S3). One other candidate to denitrification, *Stainforthia fusiformis*, was in minority: 1 % of the total fauna in both cores (Fig. 5 a, b; Table S3, GF17-3A and 3C). The other known denitrifying species previously reported in the Gullmar Fjord, *Globobulimina turgida* (Risgaard-Petersen et al., 2006) and *Globobulimina auriculata* (Whoole et al., 2018) were absent. Three non-denitrifying species (Piña-Ochoa et al., 2010; Xu et al., 2017; Glock et al., 2019) were dominant in the cores GF17-

255 3A and 3C: *Bulimina marginata* (37 and 5 %, respectively), *Cassidulina laevigata* (9 and 5 %) and *Leptoanalysis scotti* (11 and 9 %).

The density and the micro-distribution of *Nonionella* sp. T1 differed between the two cores (Fig. 5 a and b; Table S2, GF17-3A and 3C). In the core GF17-3A and 3C respectively, *Nonionella* sp. T1 density showed large variability from the water-sediment interface to 1.2 cm depth (Table S2) where *Nonionella* sp. T1 relative abundance accounted for 18 % and 50 % of the fauna in the nitrification zone (Table S3, GF17-3A and 3C). In the first denitrifying zone from 1.2 cm to 3.6 cm the *Nonionella* sp. T1 relative abundance represented 27 % and 78 % of the fauna. In the second denitrifying zone, the *Nonionella* sp. T1 relative abundance increased from 3.6 to 5 cm depth and dominated the fauna by 60 % and 98 %. The relative abundance of the denitrifying candidate, *Stainforthia fusiformis*, was a minor component in each zones of both cores and did not exceed 2 % (Table S3, GF17-3A and 3C). The three non-denitrifying species (e.g. *B. marginata*, *C. laevigata* and *L. scotti*) also dominated the fauna of both cores GF17-3A and 3C (Table S2 and S4). From the water-sediment interface to 1.2 cm depth *B. marginata* accounted for 42 % and 12 %, *C. laevigata* 16 % and 13 % and *L. scotti* 6 % and 11 %, respectively. In the first denitrifying zone (1.2-3.6 cm depth) *B. marginata* accounted for 34 % and 2 %, *C. laevigata* 7 % and 2% and *L. scotti* 25 % and 13 %, respectively. In the second denitrifying zone (3.6-5 cm depth) *B. marginata* accounted for 34 % and 0 %, *C. laevigata* was absent and *L. scotti* 5 % and 1 %, respectively.

270

Due to severe hypoxia at the GF17-1 station, oxygen was assumed to be below detection limit within the sediment. No nitrite was detected at this station ($< 1.7 \mu\text{mol L}^{-1}$). Average NO_3^- concentration in the bottom water reached $5.7 \pm 1.0 \mu\text{mol L}^{-1}$ (Fig. 5 g and h). Nitrate concentrations decreased from the sediment surface ($4.2 \pm 1.0 \mu\text{mol L}^{-1}$) to 1.6 cm ($1.8 \pm 0.6 \mu\text{mol L}^{-1}$) and then average nitrate concentration remained below the detection limit ($1.7 \mu\text{mol L}^{-1}$). However, a patch with higher nitrate concentration was visible on the left part of the gel between 2.0 and 3.0 cm depth. A 1D vertical profile passing through this patch (white line, Fig. 5 g) was extracted from the 2D image and the maximal nitrate concentration of the patch was above the detection limit with a value of $6.5 \mu\text{mol L}^{-1}$ at 2.3 cm depth (blue squares profile, Fig. 5 h). The PROFILE modelling (Table S1) of the laterally averaged nitrate vertical distribution revealed at the sampling time one denitrifying zone from the surface to 1.6 cm depth with a nitrate consumption of $2.71 \text{ E}^{-5} \text{ nmol cm}^{-3} \text{ s}^{-1}$ (red line, Fig. 5 h). Below 1.6 cm depth, nitrate

280 concentration was below the detection limit (hatched grey zone, Fig. 5 h), thus no PROFILE modelling was done after this depth.

Living foraminifera showed different total densities and a large difference in species distribution between the two cores GF17-1A and 1C (Fig. 5 e, f; Table S2), with 1457 individuals and 786 individuals, respectively (\varnothing 8.2, 5 cm depth). *Nonionella* sp. T1 represented a low relative abundance of the total fauna with 5 % in the core GF17-1A and was almost absent (1 %) in GF17-1 C (Table S3). The known denitrifying *G. auriculata* was minor in the fauna 1 % and 2%. The denitrifying candidate *S. fusiformis* was also found in the cores GF17-1A and 1C reaching only 3% of the total fauna (Figure 5 e, f; Table S3). The other denitrifying candidate *B. pseudopunctata*, was almost absent of the total fauna 0 % and 2 % (Table S3). The same three non-denitrifying species as for the oxic station were also dominant in both cores GF17-1A and 1C: *B. marginata* (64 and 30 %), *C. laevigata* (16 and 15 %) and *L. scotti* (4 and 36 %).

290 In the denitrifying zone (0-1.6 cm) *Nonionella* sp. T1 relative abundance was low, with 2 % in the core GF17-1A and was almost absent from the fauna in GF17-1C. In the core GF17-1A, *Nonionella* sp. T1 relative abundance reached 26 % of the fauna between 1.4 and 2.5 cm depth (Fig. 5 e, GF17-1A), whereas it was almost absent from the rest of the core GF17-1A and was absent from the core GF17-1C (Table S3). In the cores GF17-1A and 1C, *S. fusiformis* reached respectively 2 % and 3 % in the denitrifying zone (0-1.6 cm). In the rest of the cores from 1.6 to 5 cm depth, *S. fusiformis* represented 4 and 1 % of the fauna, respectively. The three other non-denitrifying species dominated both cores GF17-1A and 1C. In the denitrifying zone (0-1.6 cm depth) *B. marginata* accounted for 66 % and 35 %, *C. laevigata* 19 % and 19 % and *L. scotti* 4 % and 24 %. From 1.6 to 5 cm depth, *B. marginata* dominated the fauna by 61 % and 11 %, *C. laevigata* 5 % and 2 % and *L. scotti* 6 % and 75 %, respectively.

300 4 Discussion

4.1 The NIS *Nonionella* sp. T1 density in comparison with other species from the Gullmar Fjord

The presence and relative abundance of NIS *Nonionella* sp. T1 in the Gullmar Fjord and in the Skagerrak-Kattegat strait has been documented during the last decades. The earliest SEM observations of specimens resembling *Nonionella* sp. T1 morphotype in the deepest part of the fjord date back to summer 1993 (identified as *Nonionella turgida*, Gustafsson and

305 Nordberg, 2001). The invasive characteristics of *Nonionella stella* was firstly revealed by Polovodova Asteman and Schönfeld, (2015). Then, *Nonionella stella* was identified as *Nonionella* sp. T1 morphotype also described as NIS and potentially invasive species in the Oslofjord by Deldicq et al. (2019). The estimated introduction date of *Nonionella* sp. T1 into the deepest part of the Gullmar Fjord is 1985 according to Polovodova Asteman and Schönfeld, (2015). The relative abundance of *Nonionella* sp. T1 in the deepest fjord station was less than 5 % between 1985 and 2007 (Polovodova Asteman and Schönfeld, 2015 and 310 references within). At the GF17-1 hypoxic station, the *Nonionella* sp. T1 relative abundance was between 1-5 % (Table S3, GF17-1A and 1C). Thus, the *Nonionella* sp. T1 relative abundance in the deepest part of the fjord seems to remain stable. In contrast to station GF17-1, the GF17-3 oxic station was sampled for the first time in this study. In this station closer to the mouth of the fjord than GF17-1, the relative abundance of *Nonionella* sp. T1 varied between 34 and 74 % (Table S3, GF17-3A and 3C). Previous studies showed an increase in the relative abundance of *Nonionella* sp. T1 morphotype in the Skagerrak-315 Kattegat region (near the entrance of the Gullmar Fjord). The *Nonionella* sp. T1 represented 10 % of the fauna in June 2013 (Polovodova Asteman and Schönfeld, 2015). The Öresund strait linking the North Skagerrak, the Kattegat and the Baltic Sea, showed an increase in *Nonionella* sp. T1 relative abundance from 1 % to 14 % observed between 1998 and 2009 (Charrieau et al., 2019). The foraminifera fauna in the Gullmar Fjord has changed over the last decennium and *Nonionella* sp. T1 seemed to become an invasive species in the Gullmar Fjord oxic shallow water area.

320 The foraminifera fauna found at the GF17-1 station in the deepest part of the fjord differed from previous studies (Nordberg et al., 2000; Filipsson and Nordberg, 2004; Risgaard-Petersen et al., 2006; Polovodova Asteman and Nordberg, 2013; Polovodova Asteman and Schönfeld, 2015). Indeed, until the early 1980s, the foraminifera fauna in the deepest part of the fjord was dominated by a typical Skagerrak – Kattegat fauna (*Bulimina marginata*, *Cassidulina laevigata*, *Hyalinea balthica*, *Liebusella goësi*, *Nonionellina labradorica* and *Textularia earlandi*) (Nordberg et al., 2000). However, the fauna 325 changed. *S. fusiformis* and *B. pseudopunctata* became the major species (Nordberg et al., 2000; Filipsson and Nordberg, 2004). Further studies by Polovodova Asteman and Nordberg, (2013) demonstrated that at least until 2011 *S. fusiformis*, *B. pseudopunctata* and *T. earlandi* dominated the fauna. Foraminifera fauna described in the present study differs, it is the consequence of the occurrence of numerous severe hypoxic events in the fjord (Fig. 3) due to lack of deep-water exchange. In November 2017 *S. fusiformis* did not exceed 3 % of the fauna (Table S3, GF17-1A and 1C), *B. pseudopunctata* reached only

330 2 % in the core GF17-1C (Table S3, GF17-1C) and *T. earlandi* was a minor species < 1 %. Then, in November 2017 *B. marginata*, *C. laevigata* and *L. scotti* were the dominant species in the fjord. The *Elphidium clavatum-selseyensis* species complex (following the definition from Charrieau et al., 2018), *H. baltica*, *N. labradorica*, and *T. earlandi* were present in low relative abundance (< 5 %, Table S3). Namely, *G. turgida* reached 37 % of the foraminifera fauna in August 2005 at the deepest station (Risgaard-Petersen et al., 2006); whereas in November 2017 this species was minor. The decreasing in relative 335 abundance of *S. fusiformis* and *B. pseudopunctata* must be interpreted with caution since our study used the > 100 μm fraction whereas some of the previous studies used > 63 μm . We also wet picked the specimens and used Cell Tracker Green to identify living foraminifera, which might affect the results compared to Rose Bengal studies of dry sediment residuals. The relative abundance of the potential invasive *Nonionella* sp. T1 has increased according to the study of Polovodova Asteman and Schönfeld, (2015) in the oxic part of the fjord. The two non-denitrifying species *B. marginata* and *C. laevigata* described as 340 typical species of the Skagerrak-Kattegat fauna (Filipsson and Nordberg, 2004) have again increased markedly in the fjord. It is evident that the foraminifera fauna in the Gullmar Fjord is presently very dynamic with considerable species composition shifts.

4.2 Foraminifera ecology considering nitrate micro-distribution

345 Our study showed, for the first time, *Nonionella* sp. T1 dominated the foraminifera fauna in the Gullmar Fjord, this at the GF17-3 oxic station despite some spatial variability (Fig. 5 a, b; Table S2; S4). *Nonionella* sp. T1 density increased with sediment depth below the oxic zone (Fig. 5 a – d; Table S2), which could be explained by its preference to respire nitrate rather than oxygen. This would be following the hypothesis of using nitrate as a preferred electron acceptor suggested by Glock et al., (2019). *Nonionella* sp. T1 distributions could be explained by its capacity to store nitrate intracellularly before porewater 350 nitrate was denitrified by other organisms such as bacteria. At this station, *Nonionella* sp. T1 distributions may be explained as: following the oxic zone (Fig. 5 c, d; from the surface to OPD) *Nonionella* sp. T1 respire oxygen ($169 \pm 11 \text{ pmol O}_2 \text{ indiv}^{-1} \text{ d}^{-1}$). Deeper in the hypoxic zone containing nitrate (Fig. 5 c, d; from OPD to 3.6 cm depth), *Nonionella* sp. T1 accumulates intracellular nitrate and respire nitrate ($38 \pm 8 \text{ pmol N indiv}^{-1} \text{ d}^{-1}$). In the hypoxic zone where the nitrate porewater is depleted (Fig. 5 c, d; from 3.6 to 5 cm depth) *Nonionella* sp. T1 respire on its intracellular nitrate reserves to survive (Fig. 5 a, b; from

355 3.5 to 5 cm depth). When the intracellular nitrate reserve runs out, *Nonionella* sp. T1 can migrate to an upper zone where
nitrate is still present in the sediment to regenerate its intracellular nitrate reserve (Fig. 5 a, b; from 1.2 to 3.5 cm depth).

Hypoxia occurred approximately at least one month before the sampling cruise in the deepest part of the fjord (Fig. 3).
When hypoxia is extended to the water column, nitrification both in the water column and the sediment is reduced or even
360 stopped, as oxygen is almost absent (Fig. 1 b; Childs et al., 2002; Kemp et al., 2005; Conley et al., 2007; Jäntti and Hietanen,
2012). Under this condition, the coupled nitrification-denitrification processes are strongly reduced (Kemp et al., 1990). At the
GF17-1 station, no nitrification in superficial sediment was showed by our data and nitrate was low but still detectable in the
bottom water. Nitrate can diffuse from the water column into the sediment, and thereby generate the denitrification zone as
modelled by PROFILE between the surface and 1.6 cm depth (Fig. 5 h).

365 The rare presence of the NIS *Nonionella* sp. T1 and other denitrifying species as *Globobulimina auriculata*, *Bolivina*
pseudopunctata and *Stainforthia fusiformis* in the hypoxic station indicates that sediment chemical conditions turned
unfavorable towards denitrification during prolonged hypoxia. Instead, the non-denitrifying species *Bulimina marginata*,
Cassidulina laevigata, and *Leptohalysis scotti* dominated in this hypoxic environment. Their survival could be due to seasonal
dormancy (Ross and Hallock, 2016; LeKieffre et al., 2017) and propagules which can disperse and reproduce when
370 environmental conditions turn favorable again (Alve and Goldstein, 2003). The suspected deep nitrification zone (blue square
profile, Fig. 5 h) could explain the presence of nitrate micro-niches deeper in the sediment and might explain the patchy
distribution of *Nonionella* sp. T1 also at the hypoxic site (see Fig. 5 e; Table S2, GF17-1A). Therefore, deep nitrate production
in these micro-environments could favor the presence of *Nonionella* sp. T1, which can be attracted by this nitrate source of
electron acceptor to respire (Nomaki et al., 2015; Koho et al., 2011). This deep nitrification zone could be a result of an aerobic
375 or anaerobic process. An aerobic nitrification zone in deep sediment can be formed by macrofaunal activity (burrowing
activity) that introduce some oxygen deeper into anoxic sediment (Aller, 1982; Karlson et al., 2007; Nizzoli et al., 2007; Stief,
2013; Maire et al., 2016). This nitrification zone could also be due to an anaerobic process. The Gullmar Fjord is Mn-rich
(Goldberg et al., 2012) and metal-rich particles can be bio-transported into the anoxic sediment, thus allowing ammonium
oxidation into NO_3^- by Mn and Fe-oxides in the absence of oxygen deeper in the sediment (Aller, 1994; Luther et al., 1997).

4.3 Contributions and potential impacts of the NIS *Nonionella* sp. T1 to benthic denitrification in the Gullmar Fjord

If we consider that *Nonionella* sp. T1 is denitrifying the nitrate from sediment porewater (approach A, Table 2; see method 2.5) its contribution to benthic denitrification in the oxic station would be 47 % in the core GF17-3A and would reach 100 % in the core GF17-3C. If we consider that *Nonionella* sp. T1 also uses its intracellular nitrate pool for denitrification 385 (approach B), its contribution to benthic denitrification would be 32 % in the core GF17-3A and would reach 50 % in the core GF17-3C (Table 2). These two calculation approaches highlight the difficulties and the importance of knowing the concentration of environmental nitrate and foraminifera intracellular nitrate at the same time to estimate the contributions of foraminifera to benthic denitrification. Moreover, in this study there is no data on anammox process which contributes also in the total denitrification (Brandes et al., 2007). The results reported in previous studies as Engström et al., (2005) do not allow 390 us to extrapolate their data at our oxic station, located at the entrance of the fjord. Thus, we assume that our estimate of denitrification is conservative, since the possible contribution of anammox is not included in the calculation. However, despite these uncertainties *Nonionella* sp. T1 contribution to benthic denitrification supports the hypothesis that this non-indigenous denitrifying foraminifer play a major role in the benthic nitrogen cycle for sediments.

At the hypoxic station, the opposite was shown where the estimated contribution of *Nonionella* sp. T1 to benthic 395 denitrification was below 1 % whatever the calculation approach. The estimated contributions of the other denitrifying foraminifera found in the hypoxic station were low. Foraminifera contributed to almost 5 % of benthic denitrification in the hypoxic station. Compared to the oxic station, the NIS *Nonionella* sp. T1 and the other denitrifying species contributions to benthic denitrification were small in a prolonged hypoxic station of the Gullmar Fjord.

400 Overall, the Gullmar Fjord is well oxygenated except for the deepest basin where oxygen goes down when there is no deep water exchange (Fig. 3 c). Therefore, the GF17-3 oxic station could be considered representative of the Gullmar Fjord benthic ecosystem. *Nonionella* sp. T1 is not the most efficient denitrifying species compared to *Globobulimina turgida* (42 pmol N ind⁻¹ d⁻¹, with BV = 1.3 10⁺⁰⁶ µm³) and also less efficient than *Nonionella* cf. *stella* from Perú. However, *Nonionella* sp. T1 high density could accelerate sediment denitrification and participate to increase the contrast between the two

405 hydrographic conditions. Indeed, an increase in contrast due to oxygenation conditions: oxic vs severe hypoxia induced a gap in the availability of nitrate for anaerobic facultative metabolisms in the sediment. In the oxygenated part of the fjord, high contribution to benthic denitrification (estimated between 50 and 100%) by *Nonionella* sp. T1 could contribute to a potential de-eutrophication of the system by increasing the nitrogen loss. Primary production (PP) of the Gullmar Fjord is dominated by diatoms bloom in spring and autumn (Lindahl and Hernroth, 1983). Since the 1990s, Lindahl et al. (2003) observed an 410 increase in PP of the Gullmar Fjord, therefore a potential eutrophication. This increase in PP also shown in the adjacent Kattegat could be related to the nitrogen input loading from the land and atmosphere (Carstensen et al., 2003). Lindahl et al. (2003), argued that PP of the Gullmar Fjord was due to climatic forces resulting from a strong positive North Atlantic 415 Oscillation (NAO) index, which increased the availability of deep-water nutrients (Kattegat nitrate-rich) and due to warmer ocean. The benthic denitrification of the Gullmar Fjord produces nitrogen unassimilable by primary producers. Moreover, denitrifying foraminifera intracellular nitrate becomes unavailable to the system and can be bio-transported and permanently 420 sequestered in sediments (Glock et al., 2013; Prokopenko et al., 2011). Thus, denitrifying foraminifera including *Nonionella* sp. T1 could help counterbalance a potential eutrophication of the system via nitrogen loss (Seitzinger, 1988).

Whereas, in the hypoxic parts of the fjord, nitrate and nitrite rapidly exhausted become scarce, resulting in a decrease in denitrification. The consequence is a decrease of denitrifying foraminifera fauna. The increase of ammonium in a noxic 420 sediment resulting by a decrease in nitrification, denitrification and anammox processes does not allow the nitrogen elimination from the sediment to the water column. Thus, potentially promoting an ammonium accumulation in the deep fjord parts subjected to prolonged severe hypoxia (Fig. 1). Moreover, the low availability of nitrate in the sediment would possibly increase the benthic transfer towards the water column of reduced compounds such as manganese and iron produced deeper in the sedimentary column by other anaerobic metabolisms (Hulth et al., 1999). These new results demonstrate that the role of 425 denitrifying foraminifera is underestimated in the nitrogen cycle and overlooking this part of the meiofauna may lead to a misunderstanding of environments subject to hydrographic changes.

5 Conclusion


This study revealed a drastic change in living foraminifera fauna due to several hypoxic events that occurred in the last
430 decennium in the Gullmar Fjord. For the first time, the non-indigenous species (NIS) *Nonionella* sp. T1 dominated up to 74 %
the foraminifera fauna at a station with oxygenated bottom waters and high nitrate content in sediment porewater. This NIS
can denitrify up to 50-100 % of the nitrate porewater sediment under oxic conditions in the fjord. Whereas, under prolonged
hypoxia, nitrate depletion turns environmental conditions unfavorable for foraminifera denitrification, resulting in a low
density of *Nonionella* sp. T1 and other denitrifying species. Thus, foraminifera contribution to benthic denitrification was
435 negligible (~ 5 %) during prolonged seasonal hypoxia in the fjord. Moreover, the potential invasive denitrifying *Nonionella*
sp. T1 could impact the nitrogen cycle under oxic conditions by increasing the sediment denitrification and could
counterbalance potential eutrophication of the Gullmar Fjord. Thus, our study demonstrated that the role of denitrifying
foraminifera is underestimated in the nitrogen cycle especially in oxic environments.

440

445

Figures list

(a) Oxic bottom water



(b) Hypoxic bottom water




Figure 1. Simplified nitrogen cycling in marine sediments when the bottom water is oxic (a) and hypoxic (b). Chemical

450 formulae: PON (particulate organic nitrogen), NH_4^+ (ammonium), NO_3^- (nitrate), NO_2^- (nitrite), NO (nitrogen oxide),
N₂O (nitrous oxide), N₂ (nitrogen). The bold/dotted arrows indicate reactions advanced/reduced by oxygen and
nitrate presence/depletion. See text for more details. Modified from Jantti and Hietanen, (2012).

455

460






Figure 2. (a-c) Location of studied stations in the Gullmar Fjord (Sweden); blue diamond: GF17-3 oxic station (50 m depth); red square: GF17-1 hypoxic station (117 m depth); dark circles: monitoring stations Släggö (65 m depth) and Björkholmen (70 m depth). (d) Transect from the sill with four Gullmar Fjord water masses and studied stations (modified from Arneborg et al., 2004).



470

Figure 3. Record from January 2010 to September 2018 of bottom water oxygen ($[O_2]$) and nitrite + nitrate ($[NO_3^- + NO_2^-]$) measurements from the monitoring stations Släggö (65 m depth; black dot), Björkholmen (70 m depth; white triangle) and the sampling stations GF17-1 (Alsåk, 117 m depth; red square) and GF17-3 (50 m depth; blue diamond). The arrows indicate the date of the two sampling cruises: the 2017 cruise (14th, 15th November 2017) and the 2018 cruise

475 (5th September 2018). The grey zones indicate hypoxia threshold ($[O_2] < 63 \mu\text{mol L}^{-1}$).



480 **Figure 4. Scanning Electronic Microscope images of a *Nonionella* sp. T1 from the GF17-3 oxic station in the Gullmar Fjord. White lines (a, b) correspond to measured distances serving for a spheroid prolate volume model.**

485

490

Table 1. Total shell volume (μm^3) and the biovolume (BV, μm^3) corresponding to 75% of the total shell volume measured on the pool of five *Nonionella* sp. T1 from the 2017 and the 2018 cruises in the Gullmar Fjord. Abbreviations: sd (standard deviation), ind. (individual).

<i>Nonionella</i> sp. T1	1 st cruise total shell volume	1 st cruise BV	2 nd cruise total shell volume	2 nd cruise BV
ind. 1	$6.7 \cdot 10^{+06}$	$5.0 \cdot 10^{+06}$	$3.1 \cdot 10^{+06}$	$2.3 \cdot 10^{+06}$
ind. 2	$4.5 \cdot 10^{+06}$	$3.4 \cdot 10^{+06}$	$2.4 \cdot 10^{+06}$	$1.8 \cdot 10^{+06}$
ind. 3	$5.1 \cdot 10^{+06}$	$3.8 \cdot 10^{+06}$	$1.4 \cdot 10^{+06}$	$1.0 \cdot 10^{+06}$
ind. 4	$4.9 \cdot 10^{+06}$	$3.7 \cdot 10^{+06}$	$9.2 \cdot 10^{+05}$	$6.9 \cdot 10^{+05}$
ind. 5	$5.8 \cdot 10^{+06}$	$4.4 \cdot 10^{+06}$	$6.2 \cdot 10^{+05}$	$4.7 \cdot 10^{+05}$
Average (μm^3)	$5.4 \cdot 10^{+06}$	$4.0 \cdot 10^{+06}$	$1.7 \cdot 10^{+06}$	$1.3 \cdot 10^{+06}$
sd (μm^3)	$0.8 \cdot 10^{+06}$	$0.6 \cdot 10^{+06}$	$1.0 \cdot 10^{+06}$	$0.7 \cdot 10^{+06}$




Figure 5. Micro-distributions of living foraminifera densities in GF17-3oxic station (a, b) and in GF17-1hypoxic station (c, d). *Nonionella* sp. T1 specimens are in black, the sum of the non-denitrifying species in grey colors and the small dots (e, f) show the other denitrifying species (known and potential candidates). The maps of porewater nitrate 2D gels are presented for stations GF17-3 (c) and GF17-1 (g). The sediment-water interface is represented by a black line at 0 cm depth (c, g) and the Oxygen Penetration Depth (OPD) is represented by the dashed line in bold at 4.7 ± 0.2 mm depth (c). Nitrate 1D profiles (d and h, black dots) are calculated using the average value of each pixel line of the nitrate distribution image (290 pixels wide), the standard deviation is represented by two fine dotted lines (c and g respectively). The corresponding best-fitting concentration profiles (red dots, d and h) and the production zones (red line) are modelled with PROFILE. The 1D profile corresponding to $x = 1$ mm (white line, g) is represented with a blue square profile (h) and the deep nitrate spot is indicated by a black arrow. The hatched greyzone (h) represents the detection limit of the nitrate 2D gel ($<1.7 \mu\text{mol L}^{-1}$).

Table 2. Summary of the NIS *Nonionella* sp. T1 contributions to benthic denitrification in the Gullmar Fjord. The porewater denitrifications zones come from PROFILE modelling (Fig. 5 d, h). To estimate the contributions of *Nonionella* sp. T1 the counted specimens per zones was used. Two different approaches were used to estimate the contribution of *Nonionella* sp. T1: (A) divided the *Nonionella* sp. T1 denitrification rate by the nitrate porewater denitrification rate estimated from PROFILE modelling, then the second approach (B) divided the *Nonionella* sp. T1 denitrification rate by the denitrification rate from PROFILE plus the *Nonionella* sp. T1 denitrification rate. The calculations are detailed in Equation S2.

Stations	Sediment depth interval of denitrification (cm)	<i>Nonionella</i> sp. T1 (counted specimens per zone)	Nitrate porewater denitrification rates (nmol cm ⁻³ s ⁻¹)	<i>Nonionella</i> sp. T1 denitrification rates (nmol cm ⁻³ s ⁻¹)	<i>Nonionella</i> sp. T1 contribution (%), approach A	<i>Nonionella</i> sp. T1 contribution (%), approach B
GF17-3A	1.2 to 5	841	4.07 E ⁻⁰⁷	1.90 E ⁻⁰⁵	47	32
GF17-3C	1.2 to 5	1807	4.07 E ⁻⁰⁷	4.06 E ⁻⁰⁵	100	50
GF17-1A	0 to 1.6	3	2.71 E ⁻⁰⁵	6.72 E ⁻⁰⁸	0	0
GF17-1C	0 to 1.6	12	2.71 E ⁻⁰⁵	2.69 E ⁻⁰⁷	1	0

Team list

Constance Choquel
 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
constance.choquel@gmail.com

525 Emmanuelle Geslin
 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
emmanuelle.geslin@univ-angers.fr

Edouard Metzger
 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
 530 edouard.metzger@univ-angers.fr

Helena L. Filipsson
 Department of Geology, Lund University, Sweden
Helena.Filipsson@geol.lu.se

Nils Risgaard-Petersen
 535 Department of Biology, Aquatic Biology, Aarhus University, Denmark
nils.risgaard-petersen@bios.au.dk

Patrick Launeau
 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
patrick.launeau@univ-nantes.fr

540 Manuel Giraud
 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
Manuel.Giraud@univ-nantes.fr

Thierry Jauffrais
 Ifremer, IRD, Univ. Nouvelle-Calédonie, Univ. La Réunion, CNRS, UMR 9220 ENTROPIE, New Caledonia
 545 Thierry.Jauffrais@ifremer.fr

Bruno Jesus
 Université de Nantes, Mer Molécules Santé, EA 2160, France
bruno.jesus@univ-nantes.fr

Aurélia Mouret
 550 UMR 6112 LPG BIAF, Univ. Angers, Univ. Nantes, CNRS, France
aurelia.mouret@univ-angers.fr

Author contributions

555 C.C. participated in the sampling cruise, did the foraminifera taxonomy, contributed to 2D gel experiments and analyses by hyperspectral camera. C.C. did the nitrate and oxygen respiration measurements and wrote the present manuscript. E.G. participated in the sampling cruise, contributed to foraminifera analysis, scientific discussions. E.M. participated in the sampling cruise, managed with A.M. the 2D gels experiments, and contributed to hyperspectral camera treatments and scientific discussions and manuscript rewriting. H.L.F managed with A.M the sampling cruise and contributed to foraminifera

560 taxonomy and scientific discussions and manuscript rewriting. N.R.P. managed the oxygen and nitrate respiration measurements and contributed to the scientific discussions. P.L. managed hyperspectral treatments for 2D gels and contributed to scientific discussion. M.G. participated in the 2D gel lab experiments and hyperspectral treatments. T.J. participated to the sampling cruise, contributed to 2D gels experiments and scientific discussions and manuscript rewriting. B.J. contributed to scientific discussion and manuscript rewriting. A.M. managed the sampling cruise and 2D gels experiments and contributed
565 to hyperspectral camera treatments and scientific discussions and manuscript rewriting.

Author information

Corresponding author

Phone +33(0)2 41 73 53 82; fax: +33(0)2 41 73 53 52; e-mail: constance.choquel@gmail.com

570 **Competing interests**

The authors declare no competing interest.

Acknowledgements

The authors gratefully acknowledge the crews of the R/V Skagerak and Oscar von Sydow and the Kristineberg Marine
575 Research Station, the hydrographic data used in the project are from SMHI's database – SHARK. The collection of data for SHARK is organized by the Swedish environmental monitoring program and funded by the Swedish Agency for Marine and Water Management (SWAM). Thanks to Charlotte LeKieffre who helped during the sampling cruise, the SCIAM (Service Commun d'Imagerie et d'Analyses Microscopiques) of Angers University for the SEM images. HLF acknowledges funding from the Swedish Research Council VR (grant number 2017-04190). This project was funded by the French National Program
580 MANGA-2D (CNRS-INSU) and by the FRESCO project supported by the Region Pays de la Loire and by University of Angers.

References

- Aller, R. C., Hall, P. O. J., Rude, P. D. & Aller, J. Y. Biogeochemical heterogeneity and suboxic diagenesis in hemipelagic sediments of the Panama Basin. *Deep Sea Research Part I: Oceanographic Research Papers* 45, 133–165 (1998).
- Aller, R.C. The Effects of Macrobenthos on Chemical Properties of Marine Sediment and Overlying Water. *Animal-Sediment Relations*, 53-102. (1982).
- Aller, R.C. The sedimentary Mn cycle in Long Island Sound: Its role as intermediate oxidant and the influence of bioturbation, O₂, and Corg flux on diagenetic reaction balances. *Journal of Marine Research* 52, 259-295 (1994).
- Alve, E. & Murray, J. W. Marginal marine environments of the Skagerrak and Kattegat: a baseline study of living (stained) benthic foraminiferal ecology. *Palaeogeography, Palaeoclimatology, Palaeoecology* 146, 171–193 (1999).
- Alve, E. & Goldstein, S. T. Propagule transport as a key method of dispersal in benthic foraminifera (Protista). *Limnology and Oceanography*, 48(6), 2163-2170 (2003).
- Arneborg, L. Turnover times for the water above sill level in Gullmar Fjord. *Continental Shelf Research* 24, 443–460 (2004).
- Brandes, J.A., Devol, A.H. & Deutsch, C. « New Developments in the Marine Nitrogen Cycle ». *Chemical Reviews* 107, n° 2: 577-89 (2007).
- Berg, P., Risgaard-Petersen, N., & Rysgaard, S. Interpretation of Measured Concentration Profiles in Sediment Pore Water. *Limnology and Oceanography* 43, 1500-1510 (1998).
- Bernhard, J. M., Edgcomb, V. P., Casciotti, K. L., McIlvin, M. R. & Beaudoin, D. J. Denitrification likely catalyzed by endobionts in an allogromiid foraminifer. *The ISME Journal* 6, 951–960 (2012).
- Bernhard, J. M., Ostermann, D. R., Williams, D. S. & Blanks, J. K. Comparison of two methods to identify live benthic foraminifera: A test between Rose Bengal and CellTracker Green with implications for stable isotope paleoreconstructions: FORAMINIFERA VIABILITY METHOD COMPARISON. *Paleoceanography* 21, (2006).
- Björk, G. & Nordberg, K. Upwelling along the Swedish west coast during the 20th century. *Continental Shelf Research* 23, 1143–1159 (2003).

- Brandsma, J., van de Vossenberg, J., Risgaard-Petersen, N., Schmid, M. C., Engström, P., K, Eurenius, K., Hulth, S., Jaeschke, 610 Abbas, B., Hopmans, E.C., Strous, M., Schouten, S., Jetten, M. S. M., and Sinninghe Damsté, J. S. A multi-proxy study of anaerobic ammonium oxidation in marine sediments of the Gullmar Fjord, Sweden. *Environmental Microbiology Reports* 3, 360–366 (2011).
- Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert, D., Gutiérrez, D., 615 Isensee, K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C., Rabalais, N. N., Roman, M. R., Rose, K. A., Seibel, B. A., ... Zhang, J. Declining oxygen in the global ocean and coastal waters. *Science*, 359 (6371), (2018).
- Carstensen, J., Conley, D., & Müller-Karulis, B. Spatial and temporal resolution of carbon fluxes in a shallow coastal ecosystem, the Kattegat. *Marine Ecology Progress Series*, 252, 35-50 (2003).
- Cesbron, F., Metzger, E., Launeau, P., Deflandre, B., Delgard, M.-L., Thibault de Chanvalon, A., Geslin, E., Anschutz, P., & 620 Jézéquel, D. Simultaneous 2D Imaging of Dissolved Iron and Reactive Phosphorus in Sediment Porewaters by Thin-Film and Hyperspectral Methods. *Environmental Science & Technology*, 48(5), 2816-2826, (2014).
- Charrieau, L., Filipsson, H. L., Ljung, K., Chierici, M., Knudsen, K. L., & Kritzberg, E. The effects of multiple stressors on the distribution of coastal benthic foraminifera: A case study from the Skagerrak-Baltic Sea region. *Marine Micropaleontology*, 139 (Supplement C), 42-56, (2018).
- Charrieau, L., Ljung, K., Schenck, F., Daewel, U., Kritzberg, E., & Filipsson, H. L. Rapid environmental responses to climate- 625 induced hydrographic changes in the Baltic Sea entrance. *Biogeosciences*, 16, 3835-3852 (2019).
- Childs, C. R., Rabalais, N. N., Eugene, R. & Proctor, T. and L. M. Sediment denitrification in the Gulf of Mexico zone of hypoxia. (2002).
- Christensen, P. B., Rysgaard, S., Sloth, N. P., Dalsgaard, T. & Schwärter, S. Sediment mineralization, nutrient fluxes, 630 denitrification and dissimilatory nitrate reduction to ammonium in an estuarine fjord with sea cage trout farms. *Aquatic Microbial Ecology* 21, 73–84 (2000).
- Conley, D. J., Carstensen, J., Ærtebjerg, G., Christensen, P. B., Dalsgaard, T., Hansen, J. L. S., & Josefson, A. B. Long-Term Changes and Impacts of Hypoxia in Danish Coastal Waters. *Ecological Applications*, 17(sp5), S165-S184. (2007).

- Cornwell, J. C., Kemp, W. M. & Kana, T. M. Denitrification in coastal ecosystems: methods, environmental controls, and
635 ecosystem level controls, a review. *Aquatic Ecology* 33, 41–54 (1999).
- Cushman, J.A. & Moyer, D.A. Some Recent foraminifera from off San Pedro, California. Cushman Laboratory for
Foraminiferal Research Contributions, 6, 49–62 (1999).
- Dale, A. W., Sommer, S., Lomnitz, U., Bourbonnais, A. & Wallmann, K. Biological nitrate transport in sediments on the
Peruvian margin mitigates benthic sulfide emissions and drives pelagic N loss during stagnation events. *Deep Sea*
640 *Research Part I: Oceanographic Research Papers* 112, 123–136 (2016).
- Deldicq, N., Alve, E., Schweizer, M., Asteman, I. P., Hess, S., Darling, K., & Bouchet, V. M. P. History of the introduction of
a species resembling the benthic foraminifera *Nonionella stella* in the Oslofjord (Norway): Morphological, molecular
and paleo-ecological evidences. *Aquatic Invasions* 14, (2019).
- Devol, A.H. Denitrification including Anammox. Chapter 6 from *Nitrogen in the Marine Environment*, p263-292, edited by
645 Elsivier Inc, (2008).
- Diaz, R. J. Overview of Hypoxia around the World. *Journal of Environmental Quality* 30, 275–281 (2001).
- Diaz, R. J. & Rosenberg, R. Spreading Dead Zones and Consequences for Marine Ecosystems. *Science* 321, 926–929 (2008).
- Engström, Pia., Dalsgaard, T., Hulth, S., & Aller, R.C. Anaerobic ammonium oxidation by nitrite (anammox): Implications
for N₂ production in coastal marine sediments. *Geochimica et Cosmochimica Acta* 69, n° 8: 2057-65 (2005).
- 650 Filipsson, H. L. & Nordberg, K. Climate variations, an overlooked factor influencing the recent marine environment. An
example from Gullmar Fjord, Sweden, illustrated by benthic foraminifera and hydrographic data. *Estuaries* 27, 867–
881 (2004).
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C.
655 C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend, A. R., & Vöosmarty, C. J.
Nitrogen Cycles : Past, Present, and Future. *Biogeochemistry*, 70(2), 153-226 (2004).
- Geslin, E., Risgaard-Petersen, N., Lombard, F., Metzger, E., Langlet, D., & Jorissen, F. Oxygen respiration rates of benthic
foraminifera as measured with oxygen microsensors. *Journal of Experimental Marine Biology and Ecology*, 396(2),
108-114 (2011).

- Glock, N., Schönfeld, J., Eisenhauer, A., Hensen, C., Mallon, J., & Sommer, S. The role of benthic foraminifera in the benthic
660 nitrogen cycle of the Peruvian oxygen minimum zone. *Biogeosciences*, 10(7), 4767-4783, (2013).
- Glock, N., Roy, A.-S., Romero, D., Wein, T., Weissenbach, J., Revsbech, N. P., Høgslund, S., Clemens, D., Sommer, S., &
Dagan, T. Metabolic preference of nitrate over oxygen as an electron acceptor in foraminifera from the Peruvian
oxygen minimum zone. *Proceedings of the National Academy of Sciences*, 116(8), 2860-2865 (2019).
- Goldberg, T., Archer, C., Vance, D., Thamdrup, B., McAnena, A., & Poulton, S. W. Controls on Mo isotope fractionations in
665 a Mn-rich anoxic marine sediment, Gullmar Fjord, Sweden. *Chemical Geology*, 296-297, 73-82, (2012).
- Gruber, N., & Sarmiento, J. L. Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles*,
11(2), 235-266 (1997).
- Gustafsson, M. & Nordberg, K. Living (stained) benthic foraminiferal response to primary production and hydrography in the
deepest part of the Gullmar Fjord, Swedish West Coast, with comparisons to Höglund's 1927 material. *Journal of
670 Foraminiferal Research* 31, 2-11 (2001).
- Hannah, F., Rogerson, R. & Laybourn-Parry, J. Respiration rates and biovolumes of common benthic Foraminifera (Protozoa).
Journal of the Marine Biological Association of the United Kingdom 74, 301-312 (1994).
- Herbert, R. A. Nitrogen cycling in coastal marine ecosystems. *FEMS Microbiol Rev* 23, 563-590 (1999).
- Höglund, H. Foraminifera in the Gullmar Fjord and the Skagerrak. *Zoologiska Bidrag* 26, 1-328 (1947).
- 675 Høgslund, S., Revsbech, N. P., Cedhagen, T., Nielsen, L. P. & Gallardo, V. A. Denitrification, nitrate turnover, and aerobic
respiration by benthic foraminiferans in the oxygen minimum zone off Chile. *Journal of Experimental Marine Biology
and Ecology* 359, 85-91 (2008).
- Hulth, S., Aller, R.C. & Gilbert, F. Coupled anoxic nitrification/manganese reduction in marine sediments. *Geochimica et
Cosmochimica Acta* 63: 49-66 (1999).
- 680 Jäntti, H., & Hietanen, S. The Effects of Hypoxia on Sediment Nitrogen Cycling in the Baltic Sea. *AMBI* 41, 161-169 (2012).
- Karlson, K., Bonsdorff, E. & Rosenberg, R. The Impact of Benthic Macrofauna for Nutrient Fluxes from Baltic Sea Se diments.
ambi 36, 161-167 (2007).

Kemp, W. M., Sampou, P., Caffrey, J., Mayer, M., Henriksen, K., & Boynton, W. R. Ammonium recycling versus denitrification in Chesapeake Bay sediments. *Limnology and Oceanography*, 35(7), 1545-1563 (1990).

685 Kemp, W. M., Boynton, W. R., Adolf, J. E., Boesch, D. F., Boicourt, W. C., Brush, G., Cornwell, J. C., Fisher, T. R., Glibert, P. M., Hagy, J. D., Harding, L. W., Houde, E. D., Kimmel, D. G., Miller, W. D., Newell, R. I. E., Roman, M. R., Smith, E. M., & Stevenson, J. C. Eutrophication of Chesapeake Bay: Historical trends and ecological interactions. *Marine Ecology Progress Series*, 303, 1-29 (2005).

Koho, K. A., Piña-Ochoa, E., Geslin, E. & Risgaard-Petersen, N. Vertical migration, nitrate uptake and denitrification: survival mechanisms of foraminifers (*Globobulimina turgida*) under low oxygen conditions. *FEMS Microbiology Ecology* 75, 273-283 (2011).

Levin, L. A., Ekau, W., Gooday, A. J., Jorissen, F., Middelburg, J. J., Naqvi, S. W. A., Neira, C., Rabalais, N. N., & Zhang, J. Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* 6, 2063-2098 (2009).

LeKieffre, C., Spangenberg, J. E., Mabilleau, G., Escrig, S., Meibom, A., & Geslin, E. Surviving anoxia in marine sediments: 695 The metabolic response of ubiquitous benthic foraminifera (*Ammonia tepida*). *PLOS ONE*, 12(5), e0177604 (2017).

Lindahl, O., & Hernroth, L. Phyto-Zooplankton Community in Coastal Waters of Western Sweden -An Ecosystem Off Balance? *Marine Ecology Progress Series*, 10, 119-126 (1983).

Lindahl, O., Belgrano, A., Davidsson, L., & Hernroth, B. (1998). Primary production, climatic oscillations, and physico-chemical processes : The Gullmar Fjord time-series data set (1985–1996). *ICES Journal of Marine Science*, 55(4), 723-729 (2003).

700 Luther, G. W., Sundby, B., Lewis, B. L., Brendel, P. J. & Silverberg, N. Interactions of manganese with the nitrogen cycle: Alternative pathways to dinitrogen. *Geochimica et Cosmochimica Acta* 61, 4043–4052 (1997).

Maire, O., Barras, C., Gestin, T., Nardelli, M., Romero-Ramirez, A., Duchêne, J., & Geslin, E. How does macrofaunal 705 bioturbation influence the vertical distribution of living benthic foraminifera? *Marine Ecology Progress Series*, 561, 83-97 (2016).

Metzger, E., Thibault de Chanvalon, A., Cesbron, F., Barbe, A., Launeau, P., Jézéquel, D., & Mouret, A. Simultaneous Nitrite/Nitrate Imagery at Millimeter Scale through the Water–Sediment Interface. *Environmental Science & Technology*, 50(15), 8188-8195 (2016).

710 Mortimer, R. J. G., Harris, S. J., Krom, M. D., Freitag, T. E., Prosser, J. I., Barnes, J., Anschutz, P., Hayes, P. J., & Davies, I. M. Anoxic nitrification in marine sediments. *Marine Ecology Progress Series*, 276, 37-51 (2004).

Neubacher, E. C., Parker, R. E. & Trimmer, M. The potential effect of sustained hypoxia on nitrogen cycling in sediment from the southern North Sea: a mesocosm experiment. *Biogeochemistry* (2013).

715 Nizzoli, D., Bartoli, M., Cooper, M., Welsh, D. T., Underwood, G. J. C., & Viaroli, P. Implications for oxygen, nutrient fluxes and denitrification rates during the early stage of sediment colonisation by the polychaete *Nereis* spp. In four estuaries. *Estuarine, Coastal and Shelf Science*, 75(1), 125-134 (2007).

Nomaki, H., Chikaraishi, Y., Tsuchiya, M., Toyofuku, T., Suga, H., Sasaki, Y., Uematsu, K., Tame, A., & Ohkouchi, N. Variation in the nitrogen isotopic composition of amino acids in benthic foraminifera: Implications for their adaptation to oxygen-depleted environments. *Limnology and Oceanography*, 60(6), 1906-1916 (2015).

Nordberg, K. Oceanography in the Kattegat and Skagerrak Over the Past 8000 Years. *Paleoceanography* 6, 461–484 (1991).
720 Nordberg, K., Gustafsson, M. & Krantz, A.-L. Decreasing oxygen concentrations in the Gullmar Fjord, Sweden, as confirmed by benthic foraminifera, and the possible association with NAO. *Journal of Marine Systems* 23, 303–316 (2000).

Piña-Ochoa, E., Hogslund, S., Geslin, E., Cedhagen, T., Revsbech, N. P., Nielsen, L. P., Schweizer, M., Jorissen, F., Rysgaard, S., & Risgaard-Petersen, N. Widespread occurrence of nitrate storage and denitrification among Foraminifera and Gromiida. *Proceedings of the National Academy of Sciences*, 107(3), 1148-1153 (2010).

725 Polovodova Asteman, Filipsson, H. L. & Nordberg, K. Tracing winter temperatures over the last two millennia using a north-east Atlantic coastal record. *Climate of the Past*, 14, 1097-1118 (2018).

Polovodova Asteman, I. & Nordberg, K. Foraminiferal fauna from a deep basin in Gullmar Fjord: The influence of seasonal hypoxia and North Atlantic Oscillation. *Journal of Sea Research* 79, 40–49 (2013).

- 730 Polovodova Asteman, I. & Schönfeld, J. Recent invasion of the foraminifer *Nonionella stella* Cushman & Moyer, 1930 in northern European waters: evidence from the Skagerrak and its fjords. *Journal of Micropalaeontology* 35, 20–25 (2015).
- 735 Prokopenko, M. G., Sigman, D. M., Berelson, W. M., Hammond, D. E., Barnett, B., Chong, L., & Townsend-Small, A. (2011). Denitrification in anoxic sediments supported by biological nitrate transport. *Geochimica et Cosmochimica Acta*, 75(22), 7180-7199 (2011).
- Rabalais, N. N., Díaz, R. J., Levin, L. A., Turner, R. E., Gilbert, D., & Zhang, J. Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences*, 7(2), 585-619 (2010).
- Ramsing N., & Gundersen J. Seawater and gases: tabulated physical parameters of interest to people working with microsensors in marine systems. *Techn Rep MPI Mar Microbiology Bremen*, (1994).
- Revsbech, N. P. An oxygen microsensor with a guard cathode. *Limnology and Oceanography* 34, 474–478 (1989).
- 740 Risgaard-Petersen, N., Langezaal, A. M., Ingvarsson, S., Schmid, M. C., Jetten, M. S. M., Op den Camp, H. J. M., Derkzen, J. W. M., Piña-Ochoa, E., Eriksson, S. P., Peter Nielsen, L., Peter Revsbech, N., Cedhagen, T., & van der Zwaan, G. J. Evidence for complete denitrification in a benthic foraminifer. *Nature*, 443 (7107), 93-96 (2006).
- Ross, B. J. & Hallock, P. Dormancy in the Foraminifera: a review. *Journal of Foraminiferal Research* 46, 358–368 (2016).
- 745 Rysgaard, S., Risgaard-Petersen N., Sloth N. P., Jensen K., and L. P., Nielsen L.P. Oxygen Regulation of Nitrification and Denitrification in Sediments. *Limnology and Oceanography*. 39: (1643-1652) (1994).
- Seitzinger, S. P. Denitrification in freshwater and coastal marine ecosystems: Ecological and geochemical significance. *Limnology and Oceanography*, 33(4part2), 702-724 (1988).
- Sigman, D. M., Karsh, K. L., & Casciotti, K. L. Nitrogen Isotopes in the Ocean. *Encyclopedia of Ocean Sciences*, 40-54 (2009).
- 750 Stief, P. Stimulation of microbial nitrogen cycling in aquatic ecosystems by benthic macrofauna: mechanisms and environmental implications. *Biogeosciences* 10, 7829–7846 (2013).
- Stockdale, A., Davison, W., Zhang, H. Micro-scale biogeochemical heterogeneity in sediments: A review of available technology and observed evidence. *Earth-Science Reviews* 92, 81-97 (2009).

- Svansson, A. Long-term variations in the Kattegat hydrography. *ICES, CM*, (1984).
- 755 Svansson, A. Physical and chemical oceanography of the Skagerrak and the Kattegat. *Institute of Marine Research, Report No. 1.* (1975).
- Thamdrup, B. & Dalsgaard, T. *Microbial Ecology of the Oceans*. (John Wiley & Sons, Ltd, 2008).
- Thamdrup, B. New Pathways and Processes in the Global Nitrogen Cycle. *Annual Review of Ecology, Evolution, and Systematics* 43, 407–428 (2012).
- 760 Woehle, C., Roy, A.-S., Glock, N., Wein, T., Weissenbach, J., Rosenstiel, P., Hiebenthal, C., Michels, J., Schönfeld, J., & Dagan, T. A Novel Eukaryotic Denitrification Pathway in Foraminifera. *Current Biology* (2018).
- Xu, Z., Liu, S., Xiang, R. & Song, G. Live benthic foraminifera in the Yellow Sea and the East China Sea: vertical distribution, nitrate storage, and potential denitrification. *Marine Ecology Progress Series* 571, 65–81 (2017).
- Zhang, J., Gilbert, D., Gooday, A., Levin, L., Naqvi, S. W. A., Middelburg, J. J., Scranton, M., Ekau, W., Pena, A., Dewitte, B., Oguz, T., Monteiro, P. M. S., Urbán, E., Rabalais, N. N., Ittekkot, V., Kemp, W. M., Ulloa, O., Elmgren, R., Escobar-Briones, E., & Van Der Plas, A. K. Natural and human-induced hypoxia and consequences for coastal areas: Synthesis and future development. *Biogeosciences*, 7, 1443-1467 (2010).