

## Resistance and resilience of stream metabolism to high flow disturbances

Brynn O'Donnell<sup>1</sup> and Erin R. Hotchkiss<sup>1</sup>

5 <sup>1</sup>Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia

**Correspondence:** Erin R. Hotchkiss (ehotchkiss@vt.edu)

**Abstract.** Streams are ecosystems organized by disturbance. One of the most frequent and variable disturbances in running waters is elevated flow. Yet, we still have few estimates of how ecosystem processes, such as stream metabolism (gross primary production and ecosystem respiration; GPP and ER), respond to high flow events. Furthermore, we lack a predictive framework for understanding controls on within-site metabolic responses to flow disturbances. Using five years of high-frequency dissolved oxygen data from an urban- and agriculturally-influenced stream, we estimated daily GPP and ER and analyzed metabolic changes across 15 isolated high flow events. Metabolism was variable from day to day, even during lower flows; median and ranges for GPP and ER over the full measurement period were 3.7 (0.0, 17.3) and -9.6 (-2.2, -20.5) g O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup>. We calculated metabolic resistance as the magnitude of departure (M<sub>GPP</sub>, M<sub>ER</sub>) from the mean daily metabolism during antecedent lower flows (lower values of M represent higher resistance) and estimated resilience as the time until GPP and ER returned to the prior range of ambient equilibrium. We evaluated correlations between metabolic resistance and resilience with characteristics of each high flow event, antecedent conditions, and time since last flow disturbance. ER was more resistant and resilient than GPP. Median M<sub>GPP</sub> and M<sub>ER</sub> were -0.38 and -0.09, respectively. GPP was typically suppressed following flow disturbances, regardless of disturbance intensity. The magnitude of departure from baseflow ER during isolated storms increased with disturbance intensity. Additionally, GPP was less resilient and took longer to recover (0 to >9 days, mean = 2.5) than ER (0 to 6 days, mean = 1.1). Prior flow disturbances set the stage for how metabolism responds to later high flow events: the percent change in discharge during the most recent high flow event was significantly correlated with M of both GPP and ER as well as the recovery intervals for GPP. Given the flashy nature of streams draining human-altered landscapes and the variable consequences of flow for GPP and ER, testing how ecosystem processes respond to flow disturbances is essential to an integrative understanding of ecosystem function.

Deleted: range

Deleted: of

Deleted: 14

Deleted: 00

Deleted: In contrast, t

Deleted: ER

Deleted: 2

Deleted: 2

Deleted: 0.6

35 *Copyright statement.*

## 1 INTRODUCTION

40 Disturbances can alter stream ecosystem function by changing flow while influencing carbon and nutrient inputs, transformations, and exports (Stanley et al., 2010). Stream biogeochemical cycles are altered by long-term 'press' disturbances, such as land use change (e.g., Plont et al. 2020), and by episodic 'pulse' disturbances, such as transitory changes in allochthonous inputs (e.g., Bender et al. 1984; Dodds et al. 2004; Seybold and McGlynn 2018). Here, we use the definition of disturbance from White and Pickett (1985): "any relatively discrete event in time that disrupts the ecosystem. . . and changes resources,

55 substrate availability, or the physical environment". Frequent disturbances generate oscillations that form  
a dynamic ambient equilibrium (sensu Odum et al. 1995) that includes variability in processes (Resh et  
al., 1988; Stanley et al., 2010). Stream disturbances come in many forms, including: rapid increases in the  
volume and velocity of water, drought, substrate movement, and anthropogenic alterations of channel  
morphology, flow, or solute chemistry (Resh et al., 1988).

60 Elevated flow is one of the most pervasive, frequent disturbances to streams. Flow disturbances  
can scour the benthos, increase turbidity, and reduce light – all of which can change stream function (Hall  
et al., 2015; Blaszcak et al., 2019). However, flow is an inherent characteristic of streams and may  
influence stream function along a "subsidy-stress" gradient (sensu Odum et al. 1979; Figure 1). Extreme  
high flows can stress stream biota and induce conditions unfavorable for biotic processes, whereas more  
'normal', frequent high flows can stimulate internal biogeochemical transformations by bringing in  
65 limiting nutrients or organic matter subsidies (Lamberti and Steinman, 1997; Roley et al., 2014; Demars,  
2019). How changes in flow subsidize or stress stream functions will depend on a variety of factors,  
including the ecosystem process of interest.

Stream metabolism is an integrative whole-ecosystem estimate of the carbon fixed and respired  
by autotrophs and heterotrophs. Metabolism is most commonly estimated via diel changes in dissolved  
70 oxygen (Hall and Hotchkiss, 2017): autotrophs produce oxygen during gross primary production (GPP);  
autotrophs and heterotrophs consume oxygen during respiration, which we refer to as ecosystem  
respiration (ER) when measured at the whole-reach scale. Together, ER and GPP can elucidate whether a  
stream is a net producer (autotrophic;  $GPP > ER$ ) or consumer (heterotrophic;  $ER > GPP$ ) of carbon.  
Ecosystem metabolism is coupled with other ecosystem processes (e.g., nitrogen uptake, Hall and Tank  
75 2003) and is used to monitor stream health (Young et al., 2008; Jankowski et al., 2021) as well as  
ecosystem responses to disturbance and restoration (e.g., Arroita et al. 2019; Blersch et al. 2019; Palmer  
and Ruhi 2019).

Metabolism on any given day is influenced by current and past environmental factors. GPP can  
increase with light (Mulholland et al., 2001; Roberts and Mulholland, 2007), nutrients (Grimm and  
80 Fisher, 1986; Mulholland et al., 2001), temperature (Acuña et al., 2004), and transient storage  
(Mulholland et al., 2001). ER is controlled by organic carbon availability (e.g., Demars 2019), as well as  
the same physicochemical conditions as GPP, and consequently often mirrors GPP (e.g., Roberts et al.  
2007; Griffiths et al. 2013; Roley et al. 2014). Antecedent conditions may also play a role in the  
variability of ecosystem responses to flow (McMillan et al., 2018; Uehlinger and Naegeli, 1998). GPP  
85 and ER respond differently to flow disturbances (O'Donnell and Hotchkiss, 2019), likely influenced by  
where the microbes contributing to GPP and ER reside on or within the heterogeneous stream benthos  
(e.g., Uehlinger 2000, 2006). Autotroph reliance on light for energy creates a stream bed commonly  
dominated by photoautotrophic algal communities and associated heterotrophs. Many heterotrophs, on the  
other hand, are established within the substrata and hyporheic zone, which can increase resistance and  
90 resilience of ER relative to GPP (Uehlinger, 2000; Qasem et al., 2019). Environmental drivers of  
metabolism fluctuate in response to disturbances (e.g., Uehlinger 2000) but also vary sub-daily to  
seasonally, thus inducing temporal variation in GPP and ER during base flows that are best characterized  
as a pulsing steady state or dynamic equilibrium (e.g., Roberts et al. 2007).

The subsidy-stress relationship between flow and ecosystem function likely induces a range of  
95 metabolic responses to and recovery from flow changes (Figure 1). Both GPP and ER may decline due to  
disturbance during higher flows (Uehlinger, 2006; Roley et al., 2014; Reisinger et al., 2017); however,  
flow changes can also stimulate metabolism (Roberts et al., 2007; Demars, 2019). Ultimately, resistance

is reflected in the capacity of microbial assemblages to withstand a flow disturbance, with metabolic processes not reduced or stimulated outside of a dynamic ambient equilibrium. Resistance captures the instantaneous response of ecosystem metabolism to a flow disturbance. We can also quantify post-disturbance ecosystem responses by estimating resilience: the time it takes for a process returns to equilibrium following a disturbance (Carpenter et al., 1992). The resilience of ER and GPP following a flow disturbance may take anywhere from days to weeks (e.g., Uehlinger and Naegeli 1998; Smith and Kaushal 2015; Reisinger et al. 2017), and likely varies with season and the magnitude of disturbance (Uehlinger, 2006; Roberts et al., 2007). A flow event of lesser magnitude may yield higher resistance and resilience for both GPP and ER, by supplying subsidizing, limiting nutrients and organic matter from the terrestrial landscape without inducing extreme scour. Stream metabolism appears to have low resistance to disturbance but high resilience (Uehlinger and Naegeli, 1998; Reisinger et al., 2017). Understanding how different attributes of flow events (e.g., magnitude, timing) control resistance and recovery trajectories is a critical next step in characterizing metabolic responses to flow changes within and among ecosystems.

We quantified ecosystem resistance and resilience over several years of isolated, higher flow events to examine controls on and patterns of stream metabolic responses to disturbance. We had four hypotheses (Figure 1): (H1) ER will be more resistant than GPP to flow disturbances, given the protection of many heterotrophs within the streambed; (H2) there will be a stimulation of GPP and ER at intermediate flow disturbances due to an influx of limiting carbon and nutrients; (H3) metabolic resistance and resilience will change with the size of the event, with larger flow disturbances inducing more stress due to enhanced scour; and (H4) some flow events will not push GPP and ER outside of their ambient dynamic equilibrium. In addition to testing the subsidy stress hypotheses and differences in how GPP and ER may respond to and recover from higher flow events (Figure 1), we also analyzed the relationships between environmental variables and metabolic responses, including those prior to flow disturbances that may influence how stream microbial communities respond to flow changes. We predicted recent disturbances might make microbes more vulnerable and less resistant to the next high flow disturbance. We analyzed response and recovery dynamics (i.e., resistance and resilience) relative to a dynamic ambient equilibrium for 15 isolated flow events across 5 years in a flashy urban- and agriculturally-influenced stream. Our methods were chosen to address a lingering knowledge gap in our understanding of ecosystem processes, which motivated the three overall objectives of this work: (1) quantify how biological processes (GPP and ER) respond to and recover from discrete higher flow disturbances during storms (Figure 1, H2-H4), (2) test how the response and recovery of GPP and ER differ (Figure 1, H1), and (3) identify which environmental drivers best explain metabolic resistance and recovery.

## 2 METHODS

### 2.1 Study site

Stroubles Creek is a third-order, urban- and agriculturally-influenced stream draining a 15 km<sup>2</sup> sub-watershed of the New River in Southwest Virginia in the United States (Figure A1, O'Donnell and Hotchkiss 2019). The mean annual precipitation of Stroubles Creek's catchment is 1006 mm, with more than half (54%) of that precipitation falling from May-October (PRISM Climate Group, 2013). Annual mean air temperature is 11.3°C (0.4-22.0°C monthly mean minimum and maximum; PRISM Climate

Group 2013). The catchment draining into Stroubles Creek at our study location is 85.5% developed, 11.6% agriculture (pasture and crops), and 2.9% forested (Homer et al., 2015). Stroubles Creek has been designated an impaired waterway due to high sediment loading and has  $\text{NO}_3^-$  concentrations that typically exceed 1 mg/L  $\text{N-NO}_3^-$  (O'Donnell and Hotchkiss, 2019); biological oxygen demand in Stroubles Creek appears to be limited by organic carbon availability more so than inorganic nutrients (O'Donnell and Hotchkiss, unpublished data). Our study site is part of the Stream Research, Education, and Management Lab (StREAM Lab, [www.bse.vt.edu/research/facilities/StREAM\\_Lab.html](http://www.bse.vt.edu/research/facilities/StREAM_Lab.html)), and has been monitored by Virginia Tech researchers for over 10 years.

Deleted: on Stroubles

Deleted: a

## 2.2 Sensor data collection

High temporal resolution sensor data were collected from 2013-01-08 through 2018-04-14. Dissolved oxygen (DO) ( $\text{mg L}^{-1}$ ), turbidity (nephelometric turbidity unit, NTU), conductivity ( $\text{ms cm}^{-1}$ ), pH, and temperature ( $^{\circ}\text{C}$ ) data were logged at 15- 90 minute intervals by an in situ YSI 6920V2 sonde (Hession et al., 2020; O'Donnell and Hotchkiss, 2019). Because a freeze event impaired DO measurements from the YSI sonde, we gap-filled missing data with calibration-checked and comparable data from an adjacent PME MiniDOT from 2017-09-01 to 2018-04-14 (Figure A2; O'Donnell and Hotchkiss 2019). We obtained the data needed to model the relative change in light over 24-hours (Equation 1) from a nearby weather station (Figure A1), which also provided estimates of barometric pressure. A Campbell Scientific CS451 pressure transducer recorded stage measurements every 10 minutes. Velocity ( $v$ ) and width ( $w$ ) measurements were taken over multiple years to create site-specific relationships between stage, velocity, wetted width, and discharge ( $Q$ ). A stage-discharge relationship was created in 2013 and updated in 2018 to allow for daily estimates of depth ( $z$ ) from  $Q = vwz$ . Sensors were calibrated every 2-4 weeks according to best practice recommendations from the manufacturer (Hession et al. 2020) or, in the case of the PME DO sensor, with Winkler titration checks of our 100% and 0% calibration solutions (Hall and Hotchkiss 2017, O'Donnell and Hotchkiss 2019).

To remove lower-quality sensor data due to sensor error or periods of low flow, we used data cleaning and quality checks as in O'Donnell and Hotchkiss (2019). Briefly, we excluded values below the 1% and above the 99% quantile for physicochemical parameters that were heavily skewed (i.e., turbidity and conductivity). We removed physicochemical values we knew to be unreasonable (e.g., turbidity was cut off at zero). We calculated daily medians of physicochemical parameters for all days that had at least 80% of measurements over the course of the day after confirming the 80% cutoff as one that would not bias daily medians from dates without gaps in sensor measurements. Data from lower flow periods when individual sensors may have been out of water (Hession et al., 2020) were excluded when values were out of range of grab sample calibration checks.

## 2.3 Estimating ecosystem metabolism

We estimated GPP, ER, and K (the air-water gas exchange coefficient) from diel  $\text{O}_2$  (DO), light, and temperature sensor data using the same inverse modeling approach and data as O'Donnell and Hotchkiss (2019). Conservative tracer additions (Hotchkiss & O'Donnell, unpublished data) suggested there are no substantial groundwater inputs to this study reach that would otherwise bias our estimates of GPP and ER (Hall and Hotchkiss 2017). We selected the streamMetabolizer R package for our analyses (Appling et al., 2018a), which uses Bayesian parameter estimation and a hierarchical state space modeling framework to

190 generate daily estimates of GPP, ER, and K that create the best fit between modeled and observed DO data (Appling et al. 2018b; Equation 1; Table 1). GPP is multiplied by the proportion of light (PAR) at the previous measurement over total daily light.

Deleted: We note that

Deleted: , so the light term in the metabolism model function is unitless

[Eqn 1 – code copied from LaTeX version for track change doc]

$$195 \quad mDO_i = mDO_{i-\Delta t} + \frac{GPP \times PAR_{i-\Delta t}}{\sigma_{PAR}} + \frac{ER}{\sigma_{ER}} \Delta t + K_O(DO_{sat(i-\Delta t)} - mDO_{i-\Delta t}) \Delta t$$

200 We modeled GPP, ER, and K with both observation error and process error. We used most of the default model specs for streamMetabolizer. Model convergence was visualized via traceplot in the rstan package (Stan Development Team, 2019) to identify the proper number of burn-in steps (500); we saved 2000 Markov chain Monte Carlo (mcmc) steps from four chains after burn-in. We calculated credible intervals for posterior estimates of GPP and ER derived from the mcmc-derived distributions of GPP and ER.

Moved (insertion) [1]

Moved up [1]: We modeled GPP, ER, and K with both observation error and process error.

Deleted: mean daily standard error

205 Additionally, to decrease the chances of equifinality between GPP, ER, and K estimates (Appling et al., 2018b), we constrained day-to-day variability in K by binning the range of possible K estimates according to discharge (O'Donnell and Hotchkiss, 2019). We divided yearly discharge into six bins, which the hierarchical modeling framework of streamMetabolizer then used to create K-Q relationships to constrain model K estimates (O'Donnell and Hotchkiss, 2019). We used nighttime linear regression of DO as another way to estimate the range in K in Stroubles (Hall and Hotchkiss, 2017) and used regression-derived estimates of K to quality-check values of modeled K from streamMetabolizer (e.g., Figure A3). We quality-checked metabolism model output as in O'Donnell and Hotchkiss (2019). We removed all metabolism estimates that were biologically impossible, such as negative GPP or positive ER (ER is modeled as a negative flux of O<sub>2</sub> consumption). Next, we used diagnostics from fit() in stan to remove values resulting from a poor model fit or lack of chain convergence (Stan Development Team, 2019). We removed dates with poor model convergence when Rhat exceeded 1.1 and poor model fit when N<sub>eff</sub> (effective sample size) ended at or exceeded the product of the number of chains (4) and the number of saved mcmc steps (2000) specified for our model. Additionally, to avoid using biased estimates of metabolism, we removed K values below the 1% (< 3.38 d<sup>-1</sup>) and above the 99% (> 27.21 d<sup>-1</sup>) quantile of model estimates. 246 days of metabolism estimates were ultimately removed due to these model output evaluation criteria, resulting in 1375 days (of 1621 total from 2013-01-08 to 2018-04-14) of quality-checked GPP and ER for further analyses.

Deleted: these

Deleted: quality

Deleted: ¶

Deleted: ¶

Deleted: QA/QC

Deleted:

## 2.4 Selection of isolated flow events

225 To identify flow events for our analyses of metabolic resistance and resilience, we calculated the percent change in cumulative daily discharge (Q) relative to the day prior (Equation 2).

[Eqn 2 – code copied from LaTeX version for track change doc]

$$\% \Delta Q = \frac{Q_i - Q_{i-1}}{Q_{i-1}} \times 100$$

230 where  $Q_i$  is the discharge of the day of interest and  $Q_{i-1}$  is the discharge during the prior day. We selected isolated flow events that had a greater than 50% Q change relative to the antecedent cumulative daily Q.

245 We defined isolation as a period of three days before and three days after a high flow event where no  
other flow events exceeding 10% Q change occurred. In total, there were 15 isolated flow events across  
all 5 years that met our criteria for isolated flow events and had quality-checked metabolism estimates  
(Figure 2). Hydrograph and metabolism time series for each isolated flow event are available in Appendix  
Figures 4 - 18.

250 The designation of 50% change in flow for high flow events ensured analyzed events were  
outside of the range of baseline flows. We defined a flow event as >10% change in Q when comparing  
the high flow changes to prior metabolic rates, as smaller changes in Q may still influence metabolism. In  
testing different thresholds of flow change and different discharge metrics, we settled on our current  
method to optimize the thresholds for a change in Q that resulted in the highest number of quality-  
checked events while ensuring differences between classifications of ambient stream flow and higher  
255 flow events. The goal of this work was to assess how metabolism responded to and recovered from higher  
flow events that were also isolated flow events. We focused on quality over quantity when selecting for  
and analyzing stream metabolism results before, during, and after high flow events. After all appropriate  
quality-checking measures, we had 1375 days of metabolism estimates over five years (as reported in  
260 O'Donnell & Hotchkiss, 2019). To calculate resistance and recovery, we needed consecutive days of  
high-quality metabolism estimates, which further limited the number of high flow events appropriate for  
our analyses. For example, in 2016: there were 52 (out of 352) days with quality-checked sensor data that  
had a 50% flow change relative to the day prior. After looking at these 52 storms and selecting those that  
had three days before and three days after without any other flow events, we had 12 that were isolated.  
265 After quality-checking our metabolism estimates for all of those days, we had four high flow events from  
2016 that passed all quality-checking steps required for this analysis.

### 2.5 Characterizing metabolic resistance and resilience

270 To acknowledge the ambient day-to-day variability of GPP and ER, we used metabolism estimates from  
three days prior ~~each isolated flow event~~ to ~~calculate a mean value~~ of antecedent metabolism. We  
quantified metabolic responses to flow disturbances by comparing the pre-event metabolic ~~means~~ with  
event and post-event metabolism rates. To assess resistance, we estimated ~~the~~ metabolic magnitude of  
departure (M) during events to quantify the resistance of GPP and ER to ~~higher flow~~ disturbances. We  
275 calculated M per isolated flow event by comparing the difference between GPP and ER to the nearest  
value of the antecedent range (Equation 3; Figure 3),

[Eqn 3 – code copied from LaTeX version for track change]

$$280 \quad M = 1 - \frac{X_{\text{event}}}{X_{\text{prior}}}$$

where  $X_{\text{event}}$  is either GPP or ER ( $\text{g O}_2 \text{ m}^{-2} \text{ d}^{-1}$ ) on the day of the isolated flow event.  $X_{\text{prior}}$  is the ~~mean~~  
value of GPP or ER from the antecedent range, ~~and whether M is positive or negative depends~~ on if the  
isolated flow event resulted in a stimulated (increased) or suppressed (reduced) metabolic response. For  
instance, if GPP declined during a flow event, M was calculated as the difference between GPP for the  
285 isolated flow event and the ~~mean GPP~~ from the antecedent 3-day range (Figure 3). ~~If GPP or ER on the~~  
event day did not fall ~~above or below~~ the antecedent ~~mean~~, M was zero, thus indicating high resistance. A

Deleted: ¶  
¶

Deleted: quality

Deleted: define

Deleted: range

Deleted: for each isolated flow event

Deleted: range

Deleted: maximum or minimum

Deleted: depending

Deleted: minimum

Deleted: value

Deleted: We use the maximum or minimum values instead of the median or mean because this approach allowed us to better capture the full range of average metabolism estimates in ways that summarizing pre-storm rates to means or medians would exclude.

Deleted: the estimate of

Deleted: outside

Deleted: of

Deleted: range

Deleted: the magnitude of departure

310 negative  $M$  represents a suppression, and a positive  $M$  a stimulation, of GPP or ER relative to the antecedent mean.

315 To quantify the resilience of GPP and ER, we estimated recovery intervals (RI) by counting the number of days until metabolic rates returned to or exceeded pre-event mean GPP or ER, signifying a return to antecedent conditions (Figure 3). If metabolism (mean and 2.5-97.5% credible intervals) during the isolated flow event did not fall outside of the antecedent mean, the RI was zero days (metabolism cannot recover if it never shifts outside ambient values). To ensure additional flow events did not obscure the recovery interval of GPP or ER, we stopped counting RI the day before the next event (i.e., if another flow event happened four days later, we stopped counting RI at three days). To test for statistically significant differences between ER and GPP recovery intervals ( $RI_{ER}$  and  $RI_{GPP}$ ) and ER and GPP magnitude of departure ( $M_{ER}$  and  $M_{GPP}$ ), we ran Welch's t-tests in R (R Core Team, 2018).

320 While limiting our assessment to isolated flow events decreased the number of suitable events for analysis, our choice of methods allowed us to focus on metabolic response and recovery to discrete disturbances and avoid biased comparisons of multiple high flow (but not isolated) events that encompass time periods long enough (e.g., weeks) where pre/post comparisons are less meaningful. Because flow was so variable, we chose three days to balance best practices from past work on metabolic responses to storms (e.g., four days of prior stable baseflow, Reisinger et al., 2017) while ensuring we could analyze as many events with appropriately quality-checked data as possible.

## 2.6 Testing controls on metabolic resistance and resilience

330 We assessed three categories of potential predictors of metabolic resistance and resilience: antecedent conditions, characteristics of the isolated flow event, and characteristics of the most recent prior flow event. Antecedent conditions included GPP, ER, turbidity, water temperature, and light. Antecedent medians for turbidity were estimated from seven days prior due to missing sensor data. We had to remove poor-quality data from the turbidity dataset and chose to set methods that would accommodate inclusion of the most storms for our analysis. We compared the outcome of changing the number of days prior for events with turbidity data available for both three- and seven-day analyses and found no difference in the results. For all other variables, we estimated values from three days prior to the flow event for correlations between metabolism  $M$  and RI. Flow event characteristics included flow magnitude (% change of cumulative daily discharge; Equation 2), time of peak discharge, and environmental conditions (e.g., light, temperature, turbidity, season) on the event day. Characteristics of the most recent flow event included the magnitude of and days since the last flow event. We visually identified the most recent flow event (% change in cumulative daily discharge > 50) prior to each isolated flow event. We ran bivariate correlation analyses to quantify the strength and directions of linear relationships between predictor variables and metabolic resistance and resilience using the R `cor.test` function (R Core Team, 2018). We interpreted correlation strengths as: negligible ( $r = 0.0-0.3$ ), low ( $0.3-0.5$ ), moderate ( $0.5-0.7$ ), or high ( $0.7-1.0$ ) (Hinkle et al., 2003). All modeling and analyses were conducted in R (R Core Team, 2018).

## 3 RESULTS

### 3.1 Flow and metabolism

350 Stroubles Creek is a hydrologically dynamic stream, with frequent high flow events (Figure 2). Cumulative daily discharge for days with quality-checked metabolism estimates ranged from 66 to

Deleted: magnitude of departure (

Deleted: )

Deleted: equilibrium

Deleted: within the range of

Deleted: values

Deleted: ambient equilibrium

Deleted: from

Deleted: range

Deleted: and  $M$  is zero, there was no

Deleted: the

Deleted: range

Deleted: 3

Deleted: , and have noted this in our results as days+.

Deleted: significantly

Deleted: median

Deleted: flashy

114,408 m<sup>3</sup> d<sup>-1</sup>, with a median of 6,230 m<sup>3</sup> d<sup>-1</sup>. The 15 isolated flow events selected for analyses were within the mid-high range of all cumulative daily discharge values, and were of magnitudes that occurred multiple times a year (Table 2, Figure 2). We identified isolated flow events of interest based on percent changes in flow, so changes in cumulative daily discharge are proportional across seasons. During the entire measurement period, GPP ranged from 0.00 to 17.3 g O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (median = 3.7); ER ranged from -2.2 to -20.5 g O<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> (median = -9.6) (Figure 4; O'Donnell and Hotchkiss, 2019). Stroules was heterotrophic (|ER| > GPP), except for 38 days (3%) where GPP > ER, all of which occurred in spring except for one day in the fall.

### 3.2 Metabolic resistance and resilience

GPP most often declined following an isolated flow event (11 of 15 events had suppressed GPP on the high flow day), whereas ER was less likely to deviate from the antecedent equilibrium during higher flows (10 of 15 events had ER credible intervals that overlapped with antecedent mean ER). The magnitude of departure for GPP (M<sub>GPP</sub>) ranged from -0.95 to 0.34, with a mean of -0.38 (Table 3; Figure 5). GPP was inhibited during 11 and slightly stimulated (credible intervals still overlapped prior mean GPP) during three of 15 isolated flow events. The magnitude of departure for ER (M<sub>ER</sub>) ranged from -0.74 to 0.45, with a mean of -0.09 (Table 3; Figure 5). ER (mean and credible intervals) did not deviate from the antecedent mean for ten events (i.e., M<sub>ER</sub> was close to zero). ER responses to elevated flow were more variable than that of GPP; ER was both stimulated and suppressed during different high flow periods.

Although GPP exhibited stronger responses across isolated flow events than ER, M<sub>GPP</sub> and M<sub>ER</sub> were positively correlated (R<sup>2</sup> = 0.25, p = 0.03, Figure 5) and significantly different (t(26.3) = 2.15, p = 0.04). M<sub>GPP</sub> was lower than M<sub>ER</sub> for nearly all flow events, except for three in which M<sub>GPP</sub> and M<sub>ER</sub> were near zero (Table 3, Figure 5, Figure A19). The isolated flow event that induced the greatest stimulation of GPP (M<sub>GPP</sub> = 0.34) also stimulated ER (M<sub>ER</sub> = 0.08) but the credible intervals of GPP and ER on the high flow day overlapped with prior GPP and ER (Figure A06). The one high flow event that stimulated ER (M<sub>ER</sub> = 0.45) had no GPP response (M<sub>GPP</sub> = 0.04). Similarly, the only other event that stimulated GPP (M<sub>GPP</sub> = 0.32) had a minor ER response (M<sub>ER</sub> = -0.09), suggesting many flow disturbances may decouple GPP and ER.

Both GPP and ER typically recovered from flow-related stimulation or reduction in less than three days (Table 3). There were many isolated flow events where GPP took multiple days to recover but ER never departed from the antecedent dynamic equilibrium (i.e., RI = 0; Figure 5). When M<sub>GPP</sub> and M<sub>ER</sub> were both greater than zero, ER almost always recovered faster than GPP. RI<sub>GPP</sub> ranged from 0-9+ d, with an average of 2.5 d (Table 3). RI<sub>ER</sub> ranged from 0-6 d, with an average of 1.1 d (Table 3). There were only two isolated flow events where GPP recovered before ER. While ER always recovered before another flow event occurred, GPP did not recover before another flow event for two of 15 analyzed events. The recovery intervals for GPP and ER were not significantly different across all isolated high flow events (t(25.8) = -1.22, p = 0.23).

### 3.3 Controls on metabolic resistance and resilience after a flow disturbance

Although GPP and ER are linked processes, the variables that were moderate or strong predictors of resistance or resilience (r > 0.5) differed between ER and GPP (Table 4). The two predictors with

**Deleted:** rarely ... as less likely to deviated ... from the antecedent equilibrium during higher flows (10 of 15 events had ER credible intervals that overlapped with antecedent mean ER). The magnitude of departure for GPP (M<sub>GPP</sub>) ranged from -0.92 ... 5 to 0.09 ... 4, with a median ... mean of -0.14 ... 8 (Table 3; Figure 5). GPP was inhibited during 11 and slightly stimulated (credible intervals still overlapped prior mean GPP) during two ... hree of 15 isolated flow events. The magnitude of departure for ER (M<sub>ER</sub>) ranged from -0.59 ... 4 to 0.22 ... 5, with a median ... mean of -0.00 ... 9 (Table 3; Figure 5). Three of 15 flow events stimulated ER, five reduced ER, and ... R (mean and credible intervals) did not deviate from the antecedent equilibrium ... ean for seven ten events (i.e., M<sub>ER</sub> was close to zero). Overall, the direction of ... R responses to elevated flow was ... ere more variable than that of GPP. ... [1]

**Deleted:** 39 ... 5, p = 0.007 ... 3, Figure 5) and not significantly different (p = 0.06, α = 0.05 ... (26.3) = 2.15, p = 0.04). M<sub>GPP</sub> was less ... ower than M<sub>ER</sub> for nearly all flow events, except for one ... hree in which M<sub>GPP</sub> and M<sub>ER</sub> were both zero ... ere near zero and two where M<sub>GPP</sub> and M<sub>ER</sub> were both small ... Table 3, Figure 5, Figure A19). The isolated flow event that induced the greatest stimulation of GPP (M<sub>GPP</sub> = 0.09 ... 4) also stimulated ER (M<sub>ER</sub> = 0.05 ... 8) but the credible intervals of GPP and ER on the high flow day overlapped with prior GPP and ER (Figure A06). Of the other two events that ... [2]

**Deleted:** one ... ad no GPP response (M<sub>GPP</sub> = 0.04). and the other had a minor GPP reduction (M<sub>GPP</sub> = -0.06). ... imilarly, the only other event that stimulated GPP (M<sub>GPP</sub> = 0.03 ... 2) had no ... [3]

**Formatted:** Font: (Default) Times New Roman, Subscript

**Formatted:** Font: (Default) Times New Roman, Subscript

**Deleted:** M ... I = 0; Figure 5). When M<sub>GPP</sub> and M<sub>ER</sub> were both greater than zero, ER almost always recovered faster than GPP. RI<sub>GPP</sub> ranged from 1 ... -9+ d, with an average of 2.5 d (Table 3). RI<sub>ER</sub> ranged from 1 ... -2 ... d, with an average of 1.1 d (Table 3). There was ... ere only one ... wo isolated flow events where GPP recovered before ER. While ER always recovered before another flow event occurred, GPP did not recover before another flow event for four ... wo of 15 analyzed events. Excluding the four of 15 events when GPP did not recover, ... he recovery intervals for GPP and ER were not not correlated or ... ignificantly different across all isolated high flow events (p = 0.12, α = 0.05 ... [4]

**Deleted:** ; Figure A21. ... There ... he two were no ... [5]



505 moderate relationships with both  $M_{GPP}$  and  $RI_{GPP}$  were the percent change in Q during the most recent high flow event and antecedent mean GPP. The percent change in Q during the most recent high flow event was positively correlated with  $M_{ER}$  ( $r = -0.51$ ) and the only variable evaluated with  $r > 0.5$ . The magnitude of each disturbance, characterized by the % change in cumulative daily discharge, was negatively correlated with  $M_{GPP}$  ( $r = -0.40$ ,  $p = 0.14$ ) and  $M_{ER}$  ( $r = -0.49$ ,  $p = 0.07$ ) (Figure 6), and 510 positively correlated with  $RI_{GPP}$  ( $r = 0.71$ ,  $p = 0.003$ ). Overall, there were multiple environmental controls on metabolic resistance or resilience that had low correlations with either GPP or ER, but no significant drivers of both GPP and ER resistance and resilience (Table 4).

## 4 DISCUSSION

### 515 4.1 Metabolic resistance and resilience

GPP and ER responded differently to flow events in a heterotrophic stream draining a heterogeneous urban-agricultural landscape. Notably, ER was more resistant than GPP to metabolic changes induced by 520 higher flow (Figures 5,6). Of the fifteen isolated flow events analyzed here, two events stimulated ER and there were two instances of minor GPP stimulation (Table 3). The flow disturbance of heterotrophic activity was likely balanced by increased inputs of organic carbon from terrestrial sources that stimulated respiration (Roberts et al., 2007; Demars, 2019). The potential balance between subsidy and stress that buffered changes in ER during higher flows (10 of 15 events had  $M_{ER}$  near zero) was a pattern that 525 emerged in an analysis of metabolism at Stroubles Creek across all flows: GPP decreased but ER was relatively constant on days with higher than median flow (O'Donnell and Hotchkiss, 2019). While we do not discuss net ecosystem production results in the context of this work because the patterns mirror those for ER (O'Donnell and Hotchkiss, 2019), we note that during the time periods of different GPP and ER responses and recovery to higher flow, Stroubles Creek was even more heterotrophic due to the higher 530 resistance and resilience of ER relative to GPP. How often and when GPP and ER respond similarly to flow disturbances may differ among ecosystems as a function of their metabolic balance (GPP:ER), nutrient limitation status, and history of flow disturbance. Ultimately, flow-induced changes disproportionately disturbed GPP relative to ER, even in a stream like Stroubles Creek with frequently dynamic flows and relatively short recovery times.

535 ER was also more resilient than GPP. Differences in ER and GPP resilience were likely a result of flow-induced changes to physicochemical parameters (e.g., increasing turbidity with higher flows) that can also enhance the physical disturbance of flow on GPP (O'Donnell and Hotchkiss, 2019). For instance, sustained periods of high turbidity following a flow disturbance can prolong the recovery of GPP by inhibiting light attenuation (Blaszczak et al., 2019). In contrast, higher resilience of ER is likely a function of greater resistance of ER to disturbances (i.e., smaller M; Table 3) as well as flow-induced ER 540 stimulation. The correlation of  $M_{GPP}$  and  $M_{ER}$ , but a lack of correlation between  $RI_{GPP}$  and  $RI_{ER}$  (Figure 5, Table 3), suggests GPP and ER were temporarily decoupled while recovering, despite similar initial responses of GPP and ER to flow disturbances.

The dynamic nature of stream metabolism, even during low flow periods, must guide how we 545 quantify metabolic responses to disturbance. While we estimated resistance as a deviation from an antecedent average (e.g., as in Reisinger et al. 2017; Roley et al. 2014), which may limit how well we capture the dynamic antecedent equilibrium of GPP and ER during baseflow periods (Figure 1), we also leveraged posterior information about our certainty in GPP and ER estimates (i.e., Bayesian credible

Deleted: or stronger

Deleted: for

Deleted: .

Deleted: Because the median  $RI_{ER}$  was zero, bivariate correlations could not be used to determine potential predictors of ER resilience.

Deleted: Peak discharge of the flow event

Deleted: negatively

Deleted:

Deleted: 59

Deleted: 46

Deleted: 08

Deleted: and

Deleted: 66

Deleted: 01

Deleted: ,  $\alpha = 0.05$

Deleted: but was

Deleted: not significantly

Deleted: 0.54

Deleted: ,  $\alpha = 0.05$

Deleted: were strongly correlated

Deleted: isolated

Deleted: high

Deleted: three

Formatted: Font: (Default) Times New Roman, Subscript

Deleted: When

Deleted: (e.g., Reisinger et al. 2017; Roley et al. 2014), only a subset of analyzed flow events yielded 100% resistance and every metabolism estimate was classified as a reduction or stimulation

580 intervals). By assigning RIs of zero days when the mean and credible intervals of high flow GPP or ER  
were overlapping the mean of GPP or ER from three days prior to the high flow event, we reduced  
potential bias of assuming more discrete differences between day to day metabolism estimates that may  
come with using means or medians instead of the full posterior distributions provided by Bayesian  
parameter estimation. Without acknowledging the dynamic ambient equilibrium of metabolism in many  
585 streams and rivers, we may overestimate disturbances in ecosystem function. In assessing metabolic  
responses and recovery from smaller flow events relative to dynamic metabolism during ambient flows  
and acknowledging the uncertainty of metabolism model estimates, we found some of the shortest  
metabolic recovery intervals recorded in the literature (Figure 8; Table A1). Incorporating the dynamic  
nature of metabolism and standardizing calculations of metabolic recovery dynamics will enable more  
robust, cross-site comparisons of complex ecosystem response to changes in flow.

590 Our analysis of only 15 isolated flow events provided examples of all four hypothesized changes  
in metabolism with flow (Figure 1; H1-H4). ER was more resistant than GPP to most flow disturbances  
(H1). At small-to-intermediate sized flow disturbances, the response of metabolism was variable (H2,H3),  
with the greatest range of metabolic stimulation or reduction (i.e., subsidy or stress) observed at smaller  
flow changes (Figure 6). ER and GPP also did not increase or decrease relative to their ambient values  
600 during several high flow events (H4). With increasing intensity of flow disturbance, stress and  
replacement may indeed scale with intensity (H3). We note that many smaller streams, even those  
draining heavily modified landscapes, may continue to play an important role in carbon cycling and  
nutrient removal, especially during smaller flow disturbances. Further work exploring when and why  
metabolism-flow dynamics adhere to predicted disturbance responses is critical for a predictive  
understanding of disturbances and ecosystem health.

#### 4.2 Controls on metabolic resistance and resilience after a flow disturbance

605 In addition to testing potential subsidy-stress responses of metabolism to higher flow disturbances, a  
major objective of this work was to identify potential controls metabolic resistance and resilience. While  
GPP responded similarly to flows regardless of magnitude, ER was more resistant to smaller magnitude  
isolated flow events. Our prediction that isolated flow events of greater magnitudes (i.e., larger % change  
in cumulative daily discharge) would result in less resistance and higher  $M_r$  due to increased scouring,  
was supported only marginally for  $M_{GPP}$  and  $M_{ER}$  (Table 4). GPP appears to have low resistance to flow  
610 disturbances, regardless of flow magnitude (Table 4, Figure 7; Reisinger et al. 2017; Roley et al. 2014).  
Of the other stream metabolism studies that provided results suitable to include in our comparison of %  
reduction in GPP or ER and metabolic recovery intervals ( $RI_{GPP}$ ,  $RI_{ER}$ ; Figure 7), two were from streams  
draining more heavily urbanized watersheds (Reisinger et al. 2017, Qasem et al. 2019), and one was from  
a stream draining an agriculturally-dominated landscape (Roley et al. 2014). It appears streams draining  
615 more urbanized landscapes have larger reductions in metabolism and longer recovery intervals after  
higher flow disturbances; additional analyses at sites covering a range of land cover types and flow  
regimes will provide exciting opportunities to see if the trends in Figure 7 are more broadly applicable.

The different responses of GPP and ER to variable flow may be attributed to differences in  
energy sources and locations of autotrophs and heterotrophs (Uehlinger, 2000, 2006). Primary producers  
reside in exposed areas on the streambed to access light required for photosynthesis, and are thus more  
620 vulnerable to scour than heterotrophic biofilms tucked within, and protected by, substrates in the  
streambed, sediments, and hyporheic zone (Uehlinger, 2000). At some threshold of higher flows that

Deleted: .

Deleted: the

Deleted: equilibrium of

Deleted: equilibrium

Deleted:  $M_{GPP}$  and  $M_{ER}$

Deleted:  $M_{ER}$

Deleted: .

630 disturb more protected areas within and below streambeds, we expect ER will decline as flow-induced stress exceeds flow-induced carbon and nutrient subsidies. Analyses of the interactions between flow-induced changes in shear stress, water depth, and light availability may provide additional insights to tests of predicted subsidy-stress dynamics related to stream metabolism. Future analyses that include event duration may also provide new insights into flow-metabolism dynamics: Do sustained, higher flows change GPP and ER in the same way as a more instantaneous, intense flow event? As is common of long-term characterizations of metabolism in streams, many high flow days had metabolism model outputs that did not hold up to quality checks and thus were not included in our analyses. Overcoming the logistic and computational challenges of estimating metabolism during extreme flows that disturb deeper substrates will also allow us to better test predictions relating flow magnitude with ecosystem functions.

640 Quantifying how different antecedent conditions induce variable responses from GPP and ER is critical to furthering our understanding of stream ecosystem responses to flow disturbances. ~~Contrary to our prediction that past scouring might reduce future resistance to disturbances, the size of the most recent antecedent flow disturbance had a positive relationship with both  $M_{GPP}$  and  $M_{ER}$  (Table 4, Figure A19).  $M_{GPP}$  was smaller and GPP was more resistant when the most recent flow events were larger. Similarly, the % change in cumulative daily discharge from the last event was positively correlated with  $M_{GPP}$  and  $M_{ER}$ .~~ Stream biota still recovering and regenerating biomass lost from scour might respond differently to flow events depending on successional stage (Peterson and Stevenson, 1992). Furthermore, biomass growth initially stimulated by a preceding event may have been limited by one or more nutrients later supplied by the isolated flow event. ~~Antecedent GPP and  $RI_{GPP}$  were positively correlated, while  $M_{GPP}$  was negatively correlated with antecedent GPP.~~ We ultimately do not know what caused the unexpected relationships between the magnitude of the most recent event and  $M_{GPP}$  as well as  $M_{ER}$  in Stroubles Creek; quantifying the interactions between ~~recovery of biofilm communities~~ and changes in nutrient limitation across multiple flow events may provide improved insights into the mechanisms ~~jinking metabolism responses to higher flows with antecedent flow and GPP.~~

655 Environmental conditions on the day of isolated flow events that promote biomass growth, such as high light and temperature, were not significant predictors of ER or GPP recovery intervals. Metabolic recovery trajectories often increase with temperature and light (Uehlinger and Naegeli, 1998; Uehlinger, 2000), and consequently may change seasonally, with faster recoveries in spring and slower recoveries in winter (Uehlinger, 2000, 2006). ~~While we did not find any strong predictors of  $RI_{ER}$  among the environmental variables in our dataset, changes in the source, magnitude, and biological reactivity of organic matter inputs may alter  $RI_{ER}$  (Roberts et al., 2007). Combining high-frequency nutrient and organic matter quality measurements with metabolic resistance and resilience estimates will offer an improved understanding of how changing nutrients and organic matter mediate metabolic responses to flow changes.~~

## 665 5 CONCLUSIONS

670 Metabolic regimes are punctuated by high flow events that create frequent pulses of stimulated or reduced GPP or ER (e.g., Uehlinger 2006; Beaulieu et al. 2013; Bernhardt et al. 2018). As such, changes in flow play an influential role in the trends and variability in metabolism. While geomorphology and disturbance regimes may control metabolic resistance across sites (Uehlinger, 2000; Blaszcak et al., 2019), within-site variability of M and RI may be controlled by the characteristics of each flow event ~~as well as prior flow disturbances~~. Differences between ER and GPP response and recovery to flow disturbances at our

Deleted: Contrary to our predictions

Deleted: , A21

Deleted:  $M_{ER}$

Formatted: Font: (Default) Times New Roman, Subscript

Formatted: Font: (Default) Times New Roman, Subscript

Deleted: successional

Deleted: behind this relationship

Deleted: We found a negative relationship between  $RI_{GPP}$  and both temperature and season (Figure A20), suggesting that under colder conditions and in winter months, GPP takes longer to recover.

Deleted: How do changes in nutrients and organic matter govern the resistance and resilience of GPP and ER?

Deleted: resources

685 study site were controlled by higher resistance and resilience of ER relative to GPP. Within this study, our  
prediction that ER would be more resistant than GPP to flow disturbances was supported, as ER  
frequently did not even deviate from the antecedent ambient equilibrium. However, ER had less  
resistance to events of greater magnitude. ~~Indeed, both  $M_{ER}$  and  $M_{GPP}$  were negatively correlated with the~~  
690 ~~% change in discharge of flow event, but  $M_{ER}$  had a stronger negative relationship with the % change in~~  
~~discharge than  $M_{GPP}$ .~~ Metabolic responses to small and intermediate flow disturbances were variable: GPP  
and ER were both stimulated and suppressed. We suggest there may be a resistance threshold to flow  
disturbances, where controls other than flow magnitude (e.g., season, light, turbidity) might regulate  
metabolic responses to lower flow changes. Using segmented process-discharge relationships to quantify  
a resistance threshold of processes to flow disturbances (O'Donnell and Hotchkiss, 2019) may support a  
695 more predictive understanding of metabolic response to flow disturbances, as it provided insights on how  
patterns of water quality parameters and metabolism changed across the full range of flow, thus  
supporting the inferences we were able to make from storm-specific analyses in this paper.

One motivation of our work was to better understand metabolic dynamics in less pristine  
ecosystems, where ~~more dynamic~~ hydrology makes estimating metabolism more challenging, and further  
700 decreased the number of events with appropriate data for our analysis. Despite only analyzing 15 high  
flow events ~~in this study~~, many of the past analyses on related topics included a similar or fewer number  
of events over a shorter time period. Our work fills in substantial knowledge gaps: we analyzed across  
seasons (not only summer months or a short sensor deployment period) and high flow magnitudes (not  
only base flow or the highest flow disturbances), which allowed us to show a suite of different metabolic  
705 responses to changing flow. We are also left with questions about how ecosystem processes respond to  
discrete changes: how might environmental drivers of metabolic subsidy or stress determine thresholds of  
resistance and timelines of recovery? How do recent high flow events facilitate improved resistance to  
flow disturbances? What is the role of flow duration in altering metabolism within and after high flow  
events? Ultimately, we are entering an era of metabolic data opportunity (e.g., Bernhardt et al. 2018). As  
710 time series of metabolism lengthen and modeling tools improve, we envision exciting opportunities to  
better assess the consequences of isolated flow events as well as the impacts of multiple, sequential high  
flow disturbances that did not meet our criteria for analyzing isolated flow events in this paper. While the  
short time periods between high flow events in many streams and rivers make isolating and quantifying  
functional resistance and resilience an ongoing challenge, including dynamic flow in our assessment of  
715 metabolic regimes is a critical next step toward a more holistic understanding of frequently disturbed  
ecosystems.

**Code and data availability.** All data and results are included in the Appendices of this paper.  
Supplemental data files and metadata (not embedded in this document) can be downloaded at  
720 <https://tinyurl.com/RMetab>

**Author contributions.** BO developed the ideas for this manuscript with ERH. BO led the data analyses  
~~for the first draft of the manuscript~~ and developed the resistance/resilience indices with ERH. ~~BO and~~  
725 ~~ERH wrote and edited the manuscript. ERH led all analyses and edits during the manuscript revision~~  
~~process.~~

**Competing interests.** No competing interests

Deleted: :

Deleted: was

Deleted: , whereas  $M_{GPP}$  was not, suggesting that GPP responded similarly to changes in flow regardless of flow magnitude

Deleted: flashy

Deleted: ment of

Moved (insertion) [2]

Moved up [2]: BO and ERH wrote and edited the manuscript.

Deleted: BO and ERH wrote and edited the manuscript.

740 **Acknowledgements.** We thank W.C. Hession for sharing StREAM Lab sensor data and his knowledge of  
Stroubles Creek with us and L. Lehmann for assistance with database access and questions. S. Rahman  
assisted with fieldwork and preliminary data analyses. We acknowledge D. McLaughlin for helpful  
conversations and edits throughout this project. This work was supported by [an Endowment Award to BO](#)  
745 [from the Society for Freshwater Science as well as through funding received by ERH from Virginia](#)  
[Tech's Department of Biological Sciences and College of Science.](#)

Deleted: Virginia Tech's Department of Biological Sciences and an

Deleted: .

## References

- 750 Acuña, V., Giorgi, A., Muñoz, I., Uehlinger, U., and Sabater, S.: Flow extremes and benthic organic  
matter shape the metabolism of a head water Mediterranean stream, *Freshwater Biology*, 49, 960–971,  
<https://doi.org/10.1111/j.1365-2427.2004.01239.x>, <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2427.2004.01239.x>, 2004.
- 755 Appling, A. P., Hall, R. O., Arroita, M., and Yackulic, C. B.: streamMetabolizer: Models for Estimating  
Aquatic Photosynthesis and Respiration, <https://github.com/USGS-R/streamMetabolizer>, R package  
version 0.10.9, 2018a.
- 760 Appling, A. P., Hall Jr., R. O., Yackulic, C. B., and Arroita, M.: Overcoming Equifinality: Leveraging  
Long Time Series for Stream Metabolism Estimation, *Journal of Geophysical Research: Biogeosciences*,  
123, 624–645, <https://doi.org/10.1002/2017JG004140>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JG004140>, 2018b.
- 765 Arroita, M., Elosegi, A., and Hall Jr., R. O.: Twenty years of daily metabolism show riverine recovery  
following sewage abatement, *Limnology and Oceanography*, 64, S77–S92,  
<https://doi.org/10.1002/lno.11053>, <https://aslopubs.onlinelibrary.wiley.com/doi/abs/10.1002/lno.11053>,  
2019.
- 770 Beaulieu, J. J., Arango, C. P., Balz, D. A., and Shuster, W. D.: Continuous monitoring reveals multiple  
controls on ecosystem metabolism in a suburban stream, *Freshwater Biology*, 58, 918–937,  
<https://doi.org/10.1111/fwb.12097>, <https://onlinelibrary.wiley.com/doi/abs/10.1111/fwb.12097>, 2013.
- Bender, E. A., Case, T. J., and Gilpin, M. E.: Perturbation Experiments in Community Ecology: Theory  
and Practice, *Ecology*, 65, 1–13, <http://www.jstor.org/stable/1939452>, 1984.
- 775 Bernhardt, E. S., Heffernan, J. B., Grimm, N. B., Stanley, E. H., Harvey, J. W., Arroita, M., Appling, A.  
P., Cohen, M. J., McDowell, W. H., Hall Jr., R. O., Read, J. S., Roberts, B. J., Stets, E. G., and Yackulic,  
C. B.: The metabolic regimes of flowing waters, *Limnology and Oceanography*, 63, S99–S118,  
<https://doi.org/10.1002/lno.10726>, <https://aslopubs.onlinelibrary.wiley.com/doi/abs/10.1002/lno.10726>, 2018.
- 780 Blaszcak, J. R., Delesantro, J. M., Urban, D. L., Doyle, M. W., and Bernhardt, E. S.: Scoured or  
suffocated: Urban stream ecosystems oscillate between hydrologic and dissolved oxygen extremes,

- 785 Limnology and Oceanography, 64, 877–894, <https://doi.org/10.1002/lno.11081>,  
<https://aslopubs.onlinelibrary.wiley.com/doi/abs/10.1002/lno.11081>, 2019.
- Blersch, S. S., Blersch, D. M., and Atkinson, J. F.: Metabolic Variance: A Metric to Detect Shifts in Stream Ecosystem Function as a 340 Result of Stream Restoration, JAWRA Journal of the American  
790 Water Resources Association, 55, 608–621, <https://doi.org/10.1111/1752-1688.12753>,  
<https://onlinelibrary.wiley.com/doi/abs/10.1111/1752-1688.12753>, 2019.
- Carpenter, S. R., Kraft, C. E., Wright, R., He, X., Soranno, P. A., and Hodgson, J. R.: Resilience and Resistance of a Lake Phosphorus Cycle Before and After Food Web Manipulation, The American  
795 Naturalist, 140, 781–798, <https://doi.org/10.1086/285440>, <https://doi.org/10.1086/285440>, PMID:  
19426043, 1992.
- Demars, B. O. L.: Hydrological pulses and burning of dissolved organic carbon by stream respiration,  
800 Limnology and Oceanography, 64, 406–421, <https://doi.org/10.1002/lno.11048>,  
<https://aslopubs.onlinelibrary.wiley.com/doi/abs/10.1002/lno.11048>, 2019.
- Dodds, W. K., Martí, E., Tank, J. L., Pontius, J., Hamilton, S. K., Grimm, N. B., Bowden, W. B.,  
McDowell, W. H., Peterson, B. J., Valett, H. M., Webster, J. R., and Gregory, S.: Carbon and nitrogen  
805 stoichiometry and nitrogen cycling rates in streams, *Oecologia*, 140, 458–467,  
<https://doi.org/10.1007/s00442-004-1599-y>, <https://doi.org/10.1007/s00442-004-1599-y>, 2004.
- Griffiths, N. A., Tank, J. L., Royer, T. V., Roley, S. S., Rosi-Marshall, E. J., Whiles, M. R., Beaulieu, J.  
J., and Johnson, L. T.: Agricultural land use alters the seasonality and magnitude of stream metabolism,  
810 Limnology and Oceanography, 58, 1513–1529, <https://doi.org/10.4319/lo.2013.58.4.1513>, 2013.
- Grimm, N. B. and Fisher, S. G.: Nitrogen Limitation in a Sonoran Desert Stream, *Journal of the North  
American Benthological Society*, 5, 2–15, <https://doi.org/10.2307/1467743>, 1986.
- Hall, R. O. and Hotchkiss, E. R.: Chapter 34 - Stream Metabolism, in: *Methods in Stream Ecology* (Third  
815 Edition), edited by Lamberti, G. A. and Hauer, F. R., pp. 219 – 233, Academic Press,  
<https://doi.org/https://doi.org/10.1016/B978-0-12-813047-6.00012-7>, 2017.
- Hall, R. O. and Tank, J. L.: Ecosystem metabolism controls nitrogen uptake in streams in Grand Teton  
National Park, Wyoming, *Limnology and Oceanography*, 48, 1120–1128,  
820 <https://doi.org/10.4319/lo.2003.48.3.1120>, 2003.
- Hall, R. O., Yackulic, C. B., Kennedy, T. A., Yard, M. D., Rosi-Marshall, E. J., Voichick, N., and Behn,  
K. E.: Turbidity, light, temperature, and hydropeaking control primary productivity in the Colorado River,  
Grand Canyon, *Limnology and Oceanography*, 60, 512–526, <https://doi.org/10.1002/lno.10031>, 2015.  
825
- Hession, W., Lehmann, L., Wind, L., and Lofton, M.: High-frequency time series of stage height, stream  
discharge, and water quality 365 (specific conductivity, dissolved oxygen, pH, temperature, turbidity) for

- 830 Stroubles Creek in Blacksburg, Virginia, USA 2013-2018 ver 1, Environmental Data Initiative,  
<https://doi.org/10.6073/pasta/42727d38837cb4bdf04ce4e0d158ea92>, 2020.
- Homer, C., Dewitz, J., Yang, L., Jin, S., Danielson, P., Xian, G., Coulston, J., Herold, N., Wickham, J.,  
and Megown, K.: Completion of the 2011 National Land Cover Database for the Conterminous United  
States Representing a Decade of Land Cover Change Information, *Photogrammetric Engineering and  
Remote Sensing*, 81, 345–354, 2015.
- 835 Jankowski, [K.J.](#), [Mejia, F.H.](#), [Blaszczak, J.R.](#), [Holtgrieve, G.W.](#): [Aquatic ecosystem metabolism as a tool  
in environmental management](#). *WIREs Water*, e1521, <https://doi.org/10.1002/wat2.1521>, 2021.
- 840 Lambert, G. A. and Steinman, A. D.: A Comparison of Primary Production in Stream Ecosystems,  
*Journal of the North American Benthological Society*, 16, 95–104, <http://www.jstor.org/stable/1468241>,  
1997.
- 845 McMillan, S. K., Wilson, H. F., Tague, C. L., Hanes, D. M., Inamdar, S., Karwan, D. L., Loecke, T.,  
Morrison, J., Murphy, S. F., and Vidon, P.: Before the storm: antecedent conditions as regulators of  
hydrologic and biogeochemical response to extreme climate events, *Biogeochemistry*, 141, 487–501,  
<https://doi.org/10.1007/s10533-018-0482-6>, 2018.
- 850 Mulholland, P. J., Fellows, C. S., Tank, J. L., Grimm, N. B., Webster, J. R., Hamilton, S. K., Martí, E.,  
Ashkenas, L., Bowden, W. B., Dodds, W. K., McDowell, W. H., Paul, M. J., and Peterson, B. J.: Inter-  
biome comparison of factors controlling stream metabolism, *Freshwater Biology*, 46, 1503–1517,  
<https://doi.org/10.1046/j.1365-2427.2001.00773.x>, 2001.
- 855 O'Donnell, B. and Hotchkiss, E. R.: Coupling Concentration- and Process-Discharge Relationships  
Integrates Water Chemistry and 380 Metabolism in Streams, *Water Resources Research*, 55, 10 179–10  
190, <https://doi.org/10.1029/2019WR025025>, 2019.
- 860 Odum, E. P., Finn, J. T., and Franz, E. H.: Perturbation Theory and the Subsidy-Stress Gradient,  
*BioScience*, 29, 349–352, <http://www.jstor.org/stable/1307690>, 1979.
- 865 Odum, W. E., Odum, E. P., and Odum, H. T.: Nature's pulsing paradigm, *Estuaries*, 18, 547,  
<https://doi.org/10.2307/1352375>, <https://doi.org/10.2307/1352375>, 1995.
- Palmer, M. and Ruhí, A.: Linkages between flow regime, biota, and ecosystem processes: Implications  
for river restoration, *Science*, 365, <https://doi.org/10.1126/science.aaw2087>, 2019.
- 870 Peterson, C. G. and Stevenson, R. J.: Resistance and Resilience of Lotic Algal Communities: Importance  
of Disturbance Timing and Current, *Ecology*, 73, 1445–1461, <http://www.jstor.org/stable/1940689>, 1992.
- Plont, S., O'Donnell, B. M., Gallagher, M. T., and Hotchkiss, E. R.: Linking Carbon and Nitrogen  
Spiraling in Streams, *Freshwater Science*, 390 pp. 126–136, <https://doi.org/10.1086/707810>, 2020.


Deleted:

- PRISM Climate Group: "PRISM spatial climate AN81m dataset, 1981-2010",  
http://prism.oregonstate.edu, 2013.
- 875 Qasem, K., Vitousek, S., O'Connor, B., and Hoellein, T.: The effect of floods on ecosystem metabolism in suburban streams, *Freshwater Science*, 38, 412–424, <https://doi.org/10.1086/703459>, <https://doi.org/10.1086/703459>, 2019.
- 880 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org/>, 2018.
- Reisinger, A. J., Rosi, E. J., Bechtold, H. A., Doody, T. R., Kaushal, S. S., and Groffman, P. M.: Recovery and resilience of urban stream metabolism following Superstorm Sandy and other floods, *Ecosphere*, 8, e01 776, <https://doi.org/10.1002/ecs2.1776>, 2017.
- 885 Resh, V. H., Brown, A. V., Covich, A. P., Gurtz, M. E., Li, H. W., Minshall, G. W., Reice, S. R., Sheldon, A. L., Wallace, J. B., 400 and Wissmar, R. C.: The Role of Disturbance in Stream Ecology, *Journal of the North American Benthological Society*, 7, 433–455, <https://doi.org/10.2307/1467300>, <https://doi.org/10.2307/1467300>, 1988.
- 890 Roberts, B. J. and Mulholland, P. J.: In-stream biotic control on nutrient biogeochemistry in a forested stream, West Fork of Walker Branch, *Journal of Geophysical Research: Biogeosciences*, 112, <https://doi.org/10.1029/2007JG000422>, 2007.
- 895 Roberts, B. J., Mulholland, P. J., and Hill, W. R.: Multiple Scales of Temporal Variability in Ecosystem Metabolism Rates: Results from 2 Years of Continuous Monitoring in a Forested Headwater Stream, *Ecosystems*, 10, 588–606, <https://doi.org/10.1007/s10021-007-9059-2>, 2007.
- 900 Roley, S. S., Tank, J. L., Griffiths, N. A., Hall, R. O., and Davis, R. T.: The influence of floodplain restoration on whole-stream metabolism in an agricultural stream: insights from a 5-year continuous data set, *Freshwater Science*, 33, 1043–1059, <http://www.jstor.org/stable/10.4101/1086/677767>, 2014.
- 905 Seybold, E. and McGlynn, B.: Hydrologic and biogeochemical drivers of dissolved organic carbon and nitrate uptake in a headwater stream network, *Biogeochemistry*, 138, 23–48, <https://doi.org/10.1007/s10533-018-0426-1>, <https://doi.org/10.1007/s10533-018-0426-1>, 2018.
- Smith, R. M. and Kaushal, S. S.: Carbon cycle of an urban watershed: exports, sources, and metabolism, *Biogeochemistry*, 126, 173–195, <https://doi.org/10.1007/s10533-015-0151-y>, 2015.
- 910 Stan Development Team: RStan: the R interface to Stan, <http://mc-stan.org/>, r package version 2.19.2, 2019.
- 915 Stanley, E. H., Powers, S. M., and Lottig, N. R.: The evolving legacy of disturbance in stream ecology: concepts, contributions, and coming challenges, *Journal of the North American Benthological Society*, 29, 67 – 83, <https://doi.org/10.1899/08-027.1>, 2010.



- Uehlinger, U.: Resistance and resilience of ecosystem metabolism in a flood-prone river system, *Freshwater Biology*, 45, 319–332, 420 <https://doi.org/10.1111/j.1365-2427.2000.00620.x>, 2000.
- 920 Uehlinger, U.: Annual cycle and inter-annual variability of gross primary production and ecosystem respiration in a floodprone river during a 15-year period, *Freshwater Biology*, 51, 938–950, <https://doi.org/10.1111/j.1365-2427.2006.01551.x>, 2006.
- 925 Uehlinger, U. and Naegeli, M. W.: Ecosystem Metabolism, Disturbance, and Stability in a Prealpine Gravel Bed River, *Journal of the North American Benthological Society*, 17, 165–178, <http://www.jstor.org/stable/1467960>, 1998.
- White, P. and Pickett, S.: Chapter 1 - Natural Disturbance and Patch Dynamics: An Introduction, in: *The Ecology of Natural Disturbance and Patch Dynamics*, edited by Pickett, S. and White, P., pp. 3 – 13, Academic Press, San Diego, <https://doi.org/https://doi.org/10.1016/B978-0-08-050495-7.50006-5>, 1985.
- 930
- Young, R. G., Matthaei, C. D., and Townsend, C. R.: Organic matter breakdown and ecosystem metabolism: functional indicators for assessing river ecosystem health, *Journal of the North American Benthological Society*, 27, 605 – 625, <https://doi.org/10.1899/07-121.1>, 2008.
- 935

940




945



**Figure 1.** Potential metabolic responses along a subsidy-stress gradient of stream flow (adopted from Odum et al. 1979). Flow is on the x-axis. The y-axis represents ecosystem metabolism (i.e., gross primary production and ecosystem respiration; GPP and ER), scaled to the same "normal" starting values for comparison, and is broken into four categories as proposed by Odum et al. (1979): (1) subsidy (when flow replenishes carbon and nutrients and metabolism increases), (2) normal (periods of dynamic equilibrium under ambient flow), (3) stress (when ecosystem processes are suppressed by disturbance), and (4) replacement (when there is a severe reduction in metabolism and communities are scoured or replaced). H1-H4 labels correspond to different hypotheses about how GPP and ER may respond differently to flow (H1) and how metabolism might change with flow (H2-H4), and are described further in the main text of the introduction. The inset graph next to the 'normal' bracket depicts how ambient process rates are best represented by a dynamic ambient equilibrium rather than a fixed point of stability (sensu Odum et al. 1995).

950

955



960 **Figure 2.** (A) Time series of cumulative daily discharge ( $\text{m}^3 \text{d}^{-1}$ ) on all days with quality-checked  
 965 metabolism estimates from 2013-01-08 to 2018-04-14. The 15 isolated flow events analyzed for  
 metabolic responses to higher flow are represented by open squares. (B) Frequency distribution of  
 cumulative daily discharge for days with quality-checked metabolism estimates. Vertical dashed lines  
 denote the cumulative daily discharge values of the 15 different isolated flow events. (C) Box plots of  
 cumulative daily discharge ( $\text{m}^3 \text{d}^{-1}$ ) for all days with metabolism estimates versus from isolated flow event  
 days that fit our criteria for analyzing metabolic resistance and resilience.




970


**Figure 3.** Example calculations of metabolic resistance (M) and resilience (RI). Daily gross primary production (GPP) was estimated for the three days before, one day during, and two days following an isolated flow event that occurred on 2017-02-09. The solid (prior to the flow event) and dashed (during and after the flow event) horizontal line represents the average GPP estimates from three days prior to the flow event. In this case, GPP declined with higher flow, and the magnitude of departure (M with arrow) is the difference between mean prior GPP estimate from the antecedent range (dashed line) and GPP during the event. After this flow event, GPP recovered to its prior average on day two.

975

- Deleted: (grey square),
- Deleted: Dashed
- Deleted: s
- Deleted: maximum and minimum
- Deleted: minimum
- Deleted: bottom
- Deleted: antecedent range




990 **Figure 4.** Gross primary production (GPP, top) and ecosystem respiration (ER, bottom) in Stroubles Creek, VA from 2013-01-08 to 2018-04-14. ER is represented here as a negative rate because it is the consumption of oxygen. Dashed vertical lines mark the isolated flow events that fit our criteria for analyzing metabolic responses to flow change (Figure 2).




995 **Figure 5.** Resistance (i.e., magnitude of departure) of gross primary production (GPP) versus ecosystem respiration (ER) in Stroubles Creek, VA. Dashed line is the 1:1 line; solid line is the linear model fit through all data (p-value = 0.03,  $R^2 = 0.25$ ).

1000


- Deleted: s
- Deleted: are
- Deleted: s
- Deleted: 007
- Deleted: 39



**Figure 6.** Flow event magnitude (% change in cumulative daily discharge ( $Q$ ) relative to the day prior) was negatively correlated with magnitude of departure ( $M$ ) for gross primary production (GPP;  $R^2=0.10$ ,  $p=0.143$ ) and ecosystem respiration (ER;  $R^2=0.18$ ,  $p=0.066$ ). The solid purple line is the regression line for the relationship between  $M_{GPP}$  and % change in discharge, while the dashed orange line is the regression line for the relationship between  $M_{ER}$  and % change in discharge.



**Figure 7.** A synthesis of metabolic recovery intervals (days) and % reduction of gross primary production (GPP) and ecosystem respiration (ER) in response to flow disturbances. Open symbols represent GPP response, and closed symbols signify ER response. A negative % reduction is a stimulation. Included in Table A1 are additional studies that have reported either recovery intervals, or % metabolic reduction in response to flow disturbances, but not both, and consequently could not be included here.



Deleted:


Deleted: 15...0,  $r=-0.46$ , ... [6]

Deleted: 08...43) and ecosystem respiration (ER;  $R^2=0.39$ ...8,  $r=-0.66$ , ... [7]

Deleted: 01...66,  $\alpha=0.05$ ... The solid black...purple line is the regression line for the relationship between  $M_{GPP}$  and % change in discharge, while the dashed red...range line is the regression line for the relationship between  $M_{ER}$  and % change in discharge. % change in cumulative daily discharge was significantly negatively correlated with  $M_{ER}$ , but not significantly correlated with  $M_{GPP}$ . ... [8]

Formatted: Font: (Default) Times New Roman, Highlight

Formatted: Font: (Default) Times New Roman, Highlight



Deleted:

Deleted: ¶

**Table 1.** Parameter symbols, descriptions, and units used in Equation 1

Parameter symbol	Parameter description (units)
$mDO$	Modeled $O_2$ ( $g O_2 m^{-3}$ )
$\Delta t$	Measurement interval (d)
$GPP$	Gross primary production ( $g O_2 m^{-2} d^{-1}$ )
$ER$	Ecosystem respiration ( $g O_2 m^{-2} d^{-1}$ )
$z$	Mean stream channel depth (m)
$K_O$	Air-water gas exchange of $O_2$ ( $d^{-1}$ )
$O_{sat}$	DO at saturation ( $g O_2 m^{-3}$ )
$PAR$	Photosynthetically active radiation ( $\mu mol m^{-2} s^{-1}$ )

**Table 2.** Cumulative daily discharge (CDQ), % change in CDQ relative to the prior day, metabolism (gross primary production (GPP) ecosystem respiration (ER)), and air-water gas exchange (K) of isolated flow events analyzed for metabolic recovery

Date	CDQ ( $m^3 d^{-1}$ )	% $\Delta$ CDQ	GPP ( $g O_2 m^{-2} d^{-1}$ )	ER ( $g O_2 m^{-2} d^{-1}$ )	K ( $d^{-1}$ )
2013-03-12	33,970	713	1.5	-4.8	9.0
2013-03-31	13,849	188	2.4	-8.0	13.0
2013-05-23	11,923	69	3.7	-12.6	15.2
2013-06-02	6,545	93	3.2	-10.6	13.1
2015-02-02	18,842	210	1.4	-4.7	20.9
2015-05-17	19,683	94	7.2	-13.5	15.9
2015-09-03	4,447	120	6.5	-11.2	12.8
2016-04-01	13,869	67	4.8	-7.6	13.9
2016-04-07	12,478	53	5.0	-9.7	19.2
2016-04-22	18,340	114	1.9	-10.4	13.0
2016-08-21	9,418	94	0.3	-2.6	4.8
2017-02-09	20,383	149	2.2	-7.4	17.6
2017-08-21	44,543	1,105	2.5	-4.3	4.1
2017-09-06	11,600	269	0.6	-12.1	17.3
2017-10-16	8,761	54	3.4	-11.4	17.8

**Table 3.** Magnitude of departure (M, unitless) and recovery intervals (RI, days) of gross primary production (ER) during and after fifteen isolated flow events between 2013-01-08 and 2018-04-14. A negative M represents a simulation, where GPP or ER increase relative to the antecedent equilibrium. If GPP did not recover noted with a "\*" and those events are not included in recovery interval correlation analyses.

Date	$M_{GPP}$	RI <sub>GPP</sub> (d)	$M_{ER}$	RI <sub>ER</sub> (d)
2013-03-12	-0.76	3a*	-0.31	1*
2013-03-31	-0.44	1	0.00	na
2013-05-23	0.09	1	0.05	1
2013-06-02	-0.05	1	0.00	na
2015-02-02	-0.14	1	0.00	na
2015-05-17	0.00	na	0.22	2
2015-09-03	0.00	na	0.00	na
2016-04-01	-0.14	1	-0.05	1
2016-04-07	-0.03	2	0.00	na
2016-04-22	-0.83	4a*	-0.07	1*
2016-08-21	-0.92	1	-0.58	1
2017-02-09	-0.06	2	0.00	1
2017-08-21	-0.64	9a*	-0.59	1*
2017-09-06	-0.87	1	0.00	na
2017-10-16	0.03	6a*	0.00	na*

Deleted:

1050

1055

1060

1065

**Table 3.** Magnitude of departure (M, unitless) and recovery intervals (RI, days) of gross primary production (GPP) and ecosystem respiration (ER) during and after fifteen isolated flow events between 2013-01-08 and 2018-04-14. A negative M represents a suppression, and a positive M a stimulation, where GPP or ER increase relative to the prior mean GPP or ER calculated over three days. Estimates of M differed between GPP and ER ( $t(26.3)=2.15$ ,  $p=0.04$ ), while the RI for GPP and ER were not significantly different ( $t(25.8)=-1.22$ ,  $p=0.23$ ). The two instances where GPP did not recover during the isolated flow event analyzed are noted with an “NA” and the number of days without recovery (X+) that could be counted before the next high flow event occurred.

<u>Date</u>	<u>M<sub>GPP</sub></u>	<u>RI<sub>GPP</sub> (d)</u>	<u>M<sub>ER</sub></u>	<u>RI<sub>ER</sub> (d)</u>
<u>2013-03-12</u>	<u>-0.78</u>	<u>NA (6+)</u>	<u>-0.34</u>	<u>6</u>
<u>2013-03-31</u>	<u>-0.60</u>	<u>2</u>	<u>0.14</u>	<u>0</u>
<u>2013-05-23</u>	<u>0.34</u>	<u>0</u>	<u>0.08</u>	<u>0</u>
<u>2013-06-02</u>	<u>-0.34</u>	<u>2</u>	<u>0.27</u>	<u>0</u>
<u>2015-02-02</u>	<u>-0.30</u>	<u>1</u>	<u>0.05</u>	<u>0</u>
<u>2015-05-17</u>	<u>0.04</u>	<u>0</u>	<u>0.45</u>	<u>6</u>
<u>2015-09-03</u>	<u>-0.27</u>	<u>2</u>	<u>-0.17</u>	<u>0</u>
<u>2016-04-01</u>	<u>-0.38</u>	<u>4</u>	<u>-0.29</u>	<u>2</u>
<u>2016-04-07</u>	<u>-0.28</u>	<u>5</u>	<u>0.01</u>	<u>0</u>
<u>2016-04-22</u>	<u>-0.87</u>	<u>6</u>	<u>-0.23</u>	<u>0</u>
<u>2016-08-21</u>	<u>-0.95</u>	<u>2</u>	<u>-0.74</u>	<u>1</u>
<u>2017-02-09</u>	<u>-0.12</u>	<u>0</u>	<u>0.11</u>	<u>0</u>
<u>2017-08-21</u>	<u>-0.67</u>	<u>NA (9+)</u>	<u>-0.63</u>	<u>1</u>
<u>2017-09-06</u>	<u>-0.90</u>	<u>2</u>	<u>-0.01</u>	<u>0</u>
<u>2017-10-16</u>	<u>0.32</u>	<u>0</u>	<u>-0.10</u>	<u>0</u>
<u>Average</u>	<u>-0.38</u>	<u>2.5</u>	<u>-1.09</u>	<u>1.1</u>

1070

1075

1080



**Table 4.** Pearson correlations ( $r$ ) between predicted drivers of gross primary production (GPP) and ecosystem respiration (ER) magnitudes of departure (M) and recovery intervals (RI) of isolated flow events. Predictor variables with moderate or stronger relationships ( $r > 0.5$ ; Hinkle et al. 2003) are bolded. p-values are included in parentheses.

Predictor variable	$r, RI_{GPP}$	$r, M_{GPP}$	$r, RI_{ER}$	$r, M_{ER}$
<b>Isolated flow event of interest</b>				
Daily median light	0.19 (0.51)	0.17 (0.55)	-0.10 (0.74)	-0.06 (0.84)
Daily peak discharge	<b>-0.65 (0.01)</b>	-0.23 (0.42)	0.36 (0.18)	-0.39 (0.15)
Daily median temperature	0.10 (0.72)	-0.02 (0.94)	0.00 (1.00)	-0.29 (0.30)
Event median discharge (Q)	0.14 (0.63)	-0.13 (0.65)	0.50 (0.06)	0.09 (0.75)
% change in Q during event	<b>0.71 (0.00)</b>	-0.40 (0.14)	0.30 (0.28)	-0.49 (0.07)
Season	0.02 (0.93)	-0.10 (0.73)	-0.19 (0.50)	-0.27 (0.34)
Time of peak Q	0.14 (0.61)	-0.06 (0.82)	0.07 (0.81)	-0.04 (0.89)
Turbidity	0.46 (0.13)	-0.41 (0.19)	0.26 (0.41)	-0.07 (0.83)
<b>Most recent flow event</b>				
Days since last event	0.05 (0.86)	-0.07 (0.82)	-0.12 (0.67)	-0.08 (0.78)
Last event cumulative daily Q	-0.40 (0.14)	0.49 (0.06)	-0.21 (0.45)	0.14 (0.62)
% change in Q during last event	<b>-0.56 (0.03)</b>	<b>0.63 (0.01)</b>	0.38 (0.16)	<b>0.51 (0.05)</b>
<b>Antecedent conditions</b>				
Antecedent GPP	<b>0.62 (0.01)</b>	<b>-0.54 (0.04)</b>	0.13 (0.64)	-0.29 (0.29)
Antecedent ER	-0.21 (0.46)	0.00 (1.00)	0.21 (0.46)	0.33 (0.23)
Antecedent median gas exchange	0.26 (0.36)	-0.07 (0.81)	-0.11 (0.70)	-0.41 (0.13)
Antecedent median light	0.06 (0.82)	0.03 (0.92)	0.06 (0.83)	0.26 (0.36)
Antecedent median Q	-0.22 (0.41)	0.44 (0.10)	0.21 (0.44)	0.47 (0.08)
Antecedent median water temperature	0.07 (0.79)	-0.02 (0.95)	-0.09 (0.75)	-0.29 (0.29)
Antecedent median turbidity	0.12 (0.69)	-0.02 (0.95)	0.11 (0.71)	-0.29 (0.34)

Deleted:

**Table 4.** Pearson correlations ( $r$ ) between predicted drivers of gross primary production (GPP) and ecosystem respiration (ER) departure (M) and recovery intervals (RI) of isolated flow events. Predictor variables with moderate or stronger relationships ( $r > 0.5$ ; Hinkle et al. 2003) are bolded. p-values are included in parentheses.


Predictor variable (units)	$r, RI_{GPP}$	$r, M_{GPP}$	$r, RI_{ER}$	$r, M_{ER}$
<b>Isolated flow event of interest</b>				
Daily median light	-0.26 (0.51)	0.16 (0.57)	n/a	-0.17 (0.55)
Daily peak discharge	-0.03 (0.94)	-0.32 (0.25)	n/a	<b>-0.59 (0.02)</b>
Daily median temperature	-0.46 (0.21)	-0.11 (0.71)	n/a	-0.28 (0.31)
Event median discharge	0.00 (1.0)	-0.21 (0.45)	n/a	0.03 (0.91)
% change in discharge during event	-0.24 (0.54)	-0.46 (0.08)	n/a	<b>-0.66 (0.01)</b>
Season	-0.42 (0.26)	-0.16 (0.57)	n/a	-0.18 (0.53)
Time of peak discharge	<b>-0.65 (0.06)</b>	-0.12 (0.68)	n/a	-0.10 (0.73)
Turbidity	-0.21 (0.59)	-0.41 (0.18)	n/a	-0.18 (0.57)
<b>Most recent flow event</b>				
Days since last event	0.23 (0.55)	0.15 (0.58)	n/a	0.10 (0.71)
Last event cumulative daily discharge	<b>0.59 (0.09)</b>	0.40 (0.14)	n/a	0.21 (0.45)
% change in discharge during last event	<b>0.55 (0.12)</b>	<b>0.51 (0.05)</b>	n/a	0.47 (0.08)
<b>Antecedent conditions</b>				
Antecedent median gas exchange	0.26 (0.50)	-0.04 (0.88)	n/a	-0.08 (0.77)
Antecedent gross primary production	0.02 (0.96)	-0.48 (0.07)	n/a	-0.12 (0.68)
Antecedent ecosystem respiration	0.16 (0.67)	0.26 (0.34)	n/a	0.14 (0.62)
Antecedent median light	-0.31 (0.41)	0.12 (0.68)	n/a	0.04 (0.89)
Antecedent median discharge	0.08 (0.83)	0.34 (0.22)	n/a	0.40 (0.14)
Antecedent median water temperature	-0.32 (0.40)	-0.14 (0.63)	n/a	-0.29 (0.29)
Antecedent median turbidity	-0.23 (0.58)	0.13 (0.67)	n/a	0.00 (0.99)



1095


**Figure A1.** Stroubles Creek watershed and land cover types in the area that drains to the StREAM Lab monitoring site at Bridge 1, Blacksburg, VA, U.S.A. The nearby weather station used for these analyses is just west of the watershed boundary and within the same valley. We created this map using ArcGIS, NHDplus version 2.1, and the U.S. Geological Survey's 2011 National Land Cover Database.

1100




**Figure A2.** Dissolved oxygen measurements from the two sensors – YSI Sonde and PME Minidot - at Bridge 1 on Stroubles Creek. The spread of YSI Sonde values spanning from the end of January to mid-April was likely a result of a freeze event. We used PME data during the period of record when YSI data did not pass our quality assurance checks.

1105




**Figure A3.** Example of data used to confirm modeled  $K_{600}$  ( $d^{-1}$ ) using a regression of the nighttime dissolved oxygen saturation deficit versus changes in saturation (as in Hall and Hotchkiss 2017). These data are from 2017-09-04, when the estimated value for  $K_{600}$  was  $22 d^{-1}$ .

1110




Deleted:



**Figure A4.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2013-03-12 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.


Deleted: Hydrograph




**Figure A5.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2013-03-31 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.

Deleted: Hydrograph


Deleted: .



Deleted:




Deleted:




**Figure A6.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2013-05-23 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.

Deleted: Hydrograph  
Deleted: .




**Figure A7.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2013-06-02 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.


Deleted: Hydrograph  
Deleted: .



Deleted:




1155 **Figure A8.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2015-02-02 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph




1160 **Figure A9.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2015-05-17 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph

Deleted:




**Figure A10.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2015-09-03 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph

Deleted: .




**Figure A11.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2016-04-01 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph

Deleted: .




**Figure A12.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2016-04-07 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph

Deleted: .



**Figure A13.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2016-04-22 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:


Deleted: Hydrograph

Deleted: .






**Figure A14.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2016-08-21 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph

Deleted: .



**Figure A15.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2017-02-09 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.



Deleted:


Deleted: Hydrograph

Deleted: .


1215

1220

1225



**Figure A16.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2017-08-21 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:


Deleted: Hydrograph

Deleted: .

1235



**Figure A17.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2017-09-06 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




Deleted:

Deleted: Hydrograph


Deleted: .

1240


1245



Deleted:




**Figure A18.** Cumulative daily discharge and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2017-10-16 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals.




**Figure A19.** The magnitude of the previous high flow event (% change in cumulative daily discharge) had a positive relationship with  $M_{GPP}$  and  $M_{ER}$ . GPP is represented by filled black circles; ER by red crosses. The black, solid regression line reflects the relationship between magnitude of the last event and  $M_{GPP}$ , whereas the dashed, red regression line represents the relationship between the magnitude of the last event and  $M_{ER}$ .


**Deleted:** Hydrograph and metabolism (gross primary production; GPP and ecosystem respiration; ER) time series for the Stroubles Creek flow event on 2017-10-16 (noted with a dashed vertical red line in all three panels). The dashed horizontal black lines are mean values of GPP and ER prior to the high flow event. Error bars for posterior estimates of GPP and ER are 2.5% and 97.5% Bayesian credible intervals. ... [9]



**Deleted:**



**Deleted:**



**Deleted:**

**Table A1.** Literature review of published reduction and recovery intervals (RI) of stream gross primary production (GPP) and ecosystem respiration (ER) after high flow events. If not enough information was given to calculate reduction or RI, we listed as "n/a". \*=approximated days of recovery from figure in publication. \*\*=approximation given in publication.

Source	Reduction in GPP (%)	Reduction in ER (%)	RI <sub>GPP</sub> (days)	RI <sub>ER</sub> (days)
Uehlinger and Naegeli 1998	0.53	0.24	n/a	n/a
Uehlinger 2000	0.53	0.37	n/a	n/a
Uehlinger 2000	0.37	0.14	n/a	n/a
Uehlinger 2006	0.49	0.19	n/a	n/a
Roberts et al. 2007	0.90**	n/a	5.0	5.0
Roberts et al. 2007	n/a	n/a	4*	4.0
Roley et al. 2014	-1.10	-1.10	3.8	2.8
Roley et al. 2014	-0.10	-1.50	5.2	1.8
Roley et al. 2014	0.50	-0.80	16.9	1.4
Roley et al. 2014	-0.10	-1.20	7.6	4.0
Smith and Kaushal 2015	0.50**	n/a	14-21	n/a
Resinger et al. 2017	0.92	0.86	18.2	15.7
Resinger et al. 2017	0.84	0.72	7.2	10.3
Resinger et al. 2017	0.99	0.88	5.4	6.9
Resinger et al. 2017	0.99	0.81	10.1	14.1
Resinger et al. 2017	0.53	0.89	7.1	11.2
Resinger et al. 2017	0.94	0.79	7.6	13.1
Resinger et al. 2017	0.71	0.11	4.3	no recovery
Resinger et al. 2017	0.88	0.70	6.9	11.2
Resinger et al. 2017	0.97	0.84	9.0	8.8
Resinger et al. 2017	0.83	0.20	13.8	9.9
Resinger et al. 2017	0.17	0.50	11.3	11.7
Qasem et al. 2019	0.06	-0.49	3.8	1.7
Qasem et al. 2019	-0.25	-0.68	6.7	3.7
Qasem et al. 2019	0.01	-0.80	4.5	2.0
Qasem et al. 2019	0.25	-0.10	2.0	5.4
Qasem et al. 2019	0.11	-1.43	2.6	9.5
Qasem et al. 2019	-1.20	-1.02	2.3	1.6
This study (mean)	-0.38	-0.09	2.5	1.1

**Deleted:** Figure A21. Pearson correlation coefficients for tested controls on ecosystem respiration magnitude of departure ( $M_{ER}$ , gross primary production magnitude of departure ( $M_{GPP}$ ), and recovery interval of GPP ( $RI_{GPP}$ ). Variables are divided into three groups (left-right): median antecedent conditions, disturbance characteristics, and ... [10]

**Formatted:** Font: Bold

**Formatted:** Centered, Line spacing: Multiple 1.15 li

**Formatted:** Font: (Default) Times New Roman, 11 pt, Subscript

**Formatted:** Font: (Default) Times New Roman, 11 pt, Subscript

**Formatted Table**

**Formatted:** Centered, Line spacing: Multiple 1.15 li

**Formatted:** Centered, Line spacing: Multiple 1.15 li

**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [1] Deleted** **Erin Hotchkiss** **11/30/21 3:51:00 PM**



**Page 8: [2] Deleted** **Erin Hotchkiss** **11/30/21 3:58:00 PM**



Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [2] Deleted	Erin Hotchkiss	11/30/21 3:58:00 PM
▼		
Page 8: [3] Deleted	Erin Hotchkiss	11/30/21 4:08:00 PM
▼		
Page 8: [3] Deleted	Erin Hotchkiss	11/30/21 4:08:00 PM
▼		
Page 8: [3] Deleted	Erin Hotchkiss	11/30/21 4:08:00 PM
▼		
Page 8: [3] Deleted	Erin Hotchkiss	11/30/21 4:08:00 PM
▼		
Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
▼		
Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
▼		
Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
▼		
Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [4] Deleted	Erin Hotchkiss	11/30/21 4:10:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [5] Deleted	Erin Hotchkiss	11/30/21 5:06:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [5] Deleted	Erin Hotchkiss	11/30/21 5:06:00 PM
---------------------	----------------	---------------------

▼  

Page 8: [5] Deleted	Erin Hotchkiss	11/30/21 5:06:00 PM
---------------------	----------------	---------------------

▼  

Page 22: [6] Deleted	Erin Hotchkiss	11/30/21 11:55:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [6] Deleted	Erin Hotchkiss	11/30/21 11:55:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [7] Deleted	Erin Hotchkiss	11/30/21 11:55:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [7] Deleted	Erin Hotchkiss	11/30/21 11:55:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [7] Deleted	Erin Hotchkiss	11/30/21 11:55:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [8] Deleted	Erin Hotchkiss	11/30/21 11:56:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [8] Deleted	Erin Hotchkiss	11/30/21 11:56:00 AM
----------------------	----------------	----------------------

▼  

Page 22: [8] Deleted	Erin Hotchkiss	11/30/21 11:56:00 AM
----------------------	----------------	----------------------

▼

**Page 22: [8] Deleted** **Erin Hotchkiss** **11/30/21 11:56:00 AM**



**Page 22: [8] Deleted** **Erin Hotchkiss** **11/30/21 11:56:00 AM**



**Page 35: [9] Deleted** **Erin Hotchkiss** **11/30/21 12:51:00 PM**



**Page 35: [9] Deleted** **Erin Hotchkiss** **11/30/21 12:51:00 PM**



**Page 36: [10] Deleted** **Erin Hotchkiss** **11/30/21 12:42:00 PM**

