



# ***Interactive comment on “Contribution of the nongrowing season to annual N<sub>2</sub>O emissions from the continuous permafrost region in Northeast China” by Weifeng Gao et al.***

**Weifeng Gao et al.**

gaodw\_hit@163.com

Received and published: 6 June 2021

Dear Lutz Merbold:

Thank you for your letter and the constructive comments on our manuscript entitled “Contribution of the nongrowing season to annual N<sub>2</sub>O emissions from the continuous permafrost region in Northeast China” (No.: bg-2020-305). Those comments are very helpful for revising and improving our paper, as well as the important guiding significance to our research. We have carefully checked the paper and revised it base on your comments, which we hope meet with approval. Major modifications in the revised

[Printer-friendly version](#)

[Discussion paper](#)

manuscript were noted as yellow. All the responses to your comments are as following:

General remarks and major comments:

The authors present a study that investigates N<sub>2</sub>O emissions from three swamp forests in permafrost region of NorthEast China. Specifically the authors aim at addressing the contribution of non-growing season N<sub>2</sub>O emissions to the annual budget - clearly a challenge to derive reliable data with sufficient temporal resolution in permafrost regions.

While the aim of the study become clear, the authors tend to their results in various places in the manuscript. The fact that they focus on the swamp forests only comes out at a late stage while the title suggests something very different. While the could be of potential interest to the readers of BG, the current version of the manuscript can not accepted for publication for various reasons.

Overall, the results cover both, growing and non-growing season data. The nongrowing season data has considerably less temporal coverage - while it also remains unclear how long the actual growing season lasts - yet no uncertainty estimates given the accumulated numbers. At the same time, the analysis of driver variables is superficial and needs considerable work. The discussion is a loose list, sometimes a chaotic list of studies and what these found, thus it is extremely difficult for the reader to know, whether the numbers and facts presented are part of this study or another study. The actual discussion of the results however is lacking.

Some more minor comments, which need to be addressed nevertheless: No hypothesis given, The conclusion is a repetition of the results, Gapfilling procedures to derive annual budgets are not explained in detail, and many more which can be found in the commented PDF file.

Thank you for your meaningful suggestion. According to your suggestion, we have carefully revised the manuscript.

[Printer-friendly version](#)[Discussion paper](#)

First, indeed, obtaining reliable data with sufficient temporal resolution in permafrost regions is clearly a challenge. Voigt et al., (2020) summarized the studies on the N<sub>2</sub>O emissions from permafrost-affected soils, which found that the measurements of N<sub>2</sub>O emissions in permafrost regions were sparse and lacking during the nongrowing season, making the magnitude of N<sub>2</sub>O fluxes across the vast permafrost regions uncertain. Lack of wintertime measurements from permafrost regions adds to the uncertainty of the estimate contribution of nongrowing season emissions to annual budget (Voigt et al., 2020). Thus, we hope to fill in the data on the temporal variation of N<sub>2</sub>O emissions during the nongrowing season, the contribution of N<sub>2</sub>O emissions from the nongrowing season to annual budget, and the key regulatory factors on N<sub>2</sub>O emissions during the nongrowing season in this study. Indeed, sampling frequency of N<sub>2</sub>O emissions was lack during the winter in our study. Only one study reported the contribution of nongrowing season N<sub>2</sub>O emission to annual budget in the permafrost region, which was also lack of sampling frequency during the nongrowing season (Marushchak et al., 2011). According to previous studies, we clearly explained the question of the legitimacy of the integrated release estimate caused by the lack of frequency of measurements. Please see lines 469-486.

#### Reference

Marushchak, M. E., Pitkamaki, A., Koponen, H., Biasi, C., Seppala, M., and Martikainen, P. J.: Hot spots for nitrous oxide emissions found in different types of permafrost peatlands, *Global Change Biol*, 17, 2601-2614, <https://doi.org/10.1111/j.1365-2486.2011.02442.x>, 2011.

Voigt, C., Marushchak, M. E., Abbott, B. W., Biasi, C., Elberling, B., Siciliano, S. D., Sonnentag, O., Stewart, K. J., Yang, Y., and Martikainen, P. J.: Nitrous oxide emissions from permafrost-affected soils, *Nature Reviews Earth & Environment*, 1, 420-434, <https://doi.org/10.1038/s43017-020-0063-9>, 2020.

Second, we focused on the N<sub>2</sub>O emissions from the permafrost region. Thus, three

typical swamp forests in the permafrost region of Daxing'an Mountains were selected as our research target (Gao et al., 2019a, 2019b). The typical permafrost ecosystems in the Daxing'an mountains were different with other permafrost region, such as peatland in the arctic, tundra in Ny-Ålesund, forest in Eastern Siberia, alpine meadow in Qinghai-Tibetan Plateau. So, we thought that the N<sub>2</sub>O emissions from swamp forests were representative in the permafrost regions of Daxing'an Mountains. The results were "overall" in various places in the manuscript. According to your suggestion, we have revised it.

#### Reference

Gao, W. F., Yao, Y. L., Gao, D. W., Wang, H., Song, L. Q., Sheng, H. C., Cai, T. J., and Liang, H.: Responses of N<sub>2</sub>O emissions to spring thaw period in a typical continuous permafrost region of the Daxing'an Mountains, northeast China, *Atmos Environ*, 214, 116822, <https://doi.org/10.1016/j.atmosenv.2019.116822>, 2019a.

Gao, W. F., Yao, Y. L., Liang, H., Song, L. Q., Sheng, H. C., Cai, T. J., and Gao, D. W.: Emissions of nitrous oxide from continuous permafrost region in the Daxing'an Mountains, Northeast China, *Atmos Environ*, 198, 34-45, <https://doi.org/10.1016/j.atmosenv.2018.10.045>, 2019b.

Third, according to your meaningful suggestion, we defined the nongrowing season (include winter and spring thaw period) and growing season in the statistical analysis section. Please see lines 219-226. The length of nongrowing season and growing season were shown in the table 2. Please see lines 321-324. We explained the legitimacy of the integrated release estimate caused by the less temporal coverage during the nongrowing season. Please see lines 469-486. The cumulative N<sub>2</sub>O emissions were linearly and sequentially accumulated from the emissions between every two adjacent intervals of the measurements with each collar. And then the standard deviation is used to express the degree of dispersion of the cumulative N<sub>2</sub>O emissions. We re-analyzed the relationship between N<sub>2</sub>O emissions and driver variables during the growing sea-

son. We used stepwise multiple linear regression replace the multivariate regression analysis during the growing season. Please see table 5. We also rewritten the relevant abstract (Please see lines 42-45), result (Please see lines 401-410), discussion (Please see lines 559-573), and conclusion section (Please see lines 598-599). According to your suggestion, we have made major revised of the discussion. We have rewritten the 4.1 section of the discussion. We also modified the logical structure of the 4.2 and 4.3 section. Meanwhile, we clearly described the numbers and facts presented are this study or cited papers.

Fourth, according to your minor comments, we have carefully revised the manuscript. The reply is as follows:

Miner comments

Abstract

1. L32, nongrowing season N<sub>2</sub>O emissions, how is this defined?

The authors' answer:

The nongrowing season N<sub>2</sub>O emissions were defined as the N<sub>2</sub>O emissions during the nongrowing season, which were the release of N<sub>2</sub>O emissions started the daily mean air temperature remained below 0 °C for at least five consecutive days and ended at soil thawing to a depth of 20 cm. We have added the definition. Please see lines 220-226.

2. L35 1.76-2.86 times, thats very accurate, does your methods allow such accurate estimate?

The authors' answer:

Thank you for your comments. We calculated the mean N<sub>2</sub>O emissions during the winter, spring thaw period, and growing season for two years. And then we compared the mean N<sub>2</sub>O emissions among winter, spring thaw period, and growing season. We

think the data was accurate.

3. L38-39, The nongrowing season N<sub>2</sub>O emissions ranged from 0.89 to 1.44 kg ha<sup>-1</sup>, which contributed to 41.96–53.73% of the annual budget. if your growing season fluxes are that much higher, why would the contribution of nongrowing season fluxes be so large? definition and length of both seasons is essential!

The authors' answer:

Thank you for your meaningful suggestion. We defined the nongrowing season (include winter and spring thaw period) and growing season in the statistical analysis section. Please see lines 219-226. The nongrowing season lasted for about 230 days, which were 1.8-fold than that of the growing season (130 days, Table 2). The cumulative N<sub>2</sub>O emissions during the short spring thaw period were contributed to 15.63–33.00% of the annual emissions. Although the rates of N<sub>2</sub>O during the growing season were higher than that of nongrowing season, the longer nongrowing season period made the cumulative nongrowing season N<sub>2</sub>O emissions close to the emissions during the growing season.

4. L40, different among during, wording.

The authors' answer:

Thank you for your comments. We have revised it.

5. L41-44, The N<sub>2</sub>O emissions from total two-year observation period and nongrowing season were mainly affected by soil temperature, while the N<sub>2</sub>O emissions from growing season were controlled by soil temperature, water table level, and their interactions. can you be more precisely? how much of the variability of N<sub>2</sub>O fluxes were expl. by T<sub>soil</sub>?

The authors' answer:

Thank you for your meaningful suggestion. The N<sub>2</sub>O emissions from nongrowing sea-

son and total two-year observation period were mainly affected by soil temperature, which could explain 3.01-9.54% and 6.07-14.48% temporal variation of N<sub>2</sub>O emissions, respectively. While the N<sub>2</sub>O emissions from growing season were controlled by soil temperature, water table level, pH, NH<sub>4</sub><sup>+</sup>-N, NO<sub>3</sub><sup>-</sup>-N, total nitrogen, total organic carbon, and C/N ratio, which could explain 14.51–45.72% temporal variation of N<sub>2</sub>O emissions. We have revised it. Please see lines 40-45.

6. L45, is an important, are an important.

The authors' answer:

We have revised it.

Introduction

1. L64, tropical soil, soils.

The authors' answer:

We have revised it.

2. L78, 40 kg ha<sup>-1</sup>, I presume you mean N per year - please adjust.

The authors' answer:

Thank you for your suggestion. We replace the kg ha<sup>-1</sup> with kg N ha<sup>-1</sup>.

3. L116, hypothesis are missing

The authors' answer:

We have added the hypothesis. We hypothesis that: (i) N<sub>2</sub>O emissions from permafrost region had significantly temporal variation on an annual scale, and mean N<sub>2</sub>O emissions during the nongrowing season were lower than that of growing season; (ii) the nongrowing season N<sub>2</sub>O emissions had significance contribute to annual budget, which cannot be ignored in the permafrost region; (iii) there had difference key regulatory factors for N<sub>2</sub>O emissions during the nongrowing season, growing season, and

entire period. Please see lines 109-114.

## Material and Methods

1. L128, primarily brown forest soil, WRB Soil type classification.

The authors' answer:

Thank you for your suggestion. According to the SC1 comments, the USDA soil type classification was used in the manuscript. The zonal soil at the study site is Gelisols (according to USDA soil classification). Please see lines 131-132.

2. L157, units are missing.

The authors' answer:

Thank you for your meaningful comments. We had added the units. Please see table 1.

3. L171, inserted 20 cm into the soil, thats quite deep - why?

The authors' answer:

The base collar was inserted 20 cm into the soil to prevent the exchange of N<sub>2</sub>O gas inside and outside of base collar, which could better evaluate the release of N<sub>2</sub>O inside the base collar.

4. L185, gas N<sub>2</sub>O concentration, N<sub>2</sub>O concentration.

The authors' answer:

We have revised it.

5. L198-199, In the nongrowing season, soil moisture was determined by the oven-drying method. gravimetric water content.

The authors' answer:



Yes, gravimetric water content was determined by the oven-drying method. And then volumetric water content was converted by gravimetric water content and bulk density. We have revised it. Please see lines 185-186 and 192-198.

6. L233, reported by (Hou et al., 2012), formatting.

The authors' answer:

We have revised it (by Hou et al., (2012)).

Results

1. L254, temporal pattern, patterns.

The authors' answer:

We have replaced pattern with patterns.

2. L260-261, Negative emissions mainly occurred during the winter and spring thaw period. Negative values were real and not part due to leaks? What were your tests to ensure chamber closure?

The authors' answer:

The negative values were real. We checked the air tightness of each chamber before gas sampling to ensure that the chamber was closure.

3. L264, Fig 1, only site three is in the end significantly different, isn't it? there is a low coverage of the winter fluxes.

The authors' answer:

Yes. The site three in the end is significantly different. In the revised manuscript, we increased the discussion of sampling frequency of N<sub>2</sub>O emissions during the winter. Please see lines 469-486.

4. L297, overall, how confident are you in the cumulative winter emissions? much lower

sampling frequency and thus larger uncertainty - see Barton et al. Scientific reports on sampling frequency vs annual/seasonal budgets. I am missing a thorough uncertainty analysis.

The authors' answer:

Thank you for your meaningful suggestion. Indeed, the sampling frequency would influence the assess of the cumulative N<sub>2</sub>O emissions. According your suggestion, we increased the discussion of N<sub>2</sub>O emission during the winter. According to your meaningful suggestion, we clearly explained the question of the legitimacy of the integrated release estimate caused by the lack of frequency during the winter. Please see lines 469-486.

5. L307, STP: spring thaw period. what does this number mean?

The authors' answer:

The number means the contribution of spring thaw period to the annual N<sub>2</sub>O emissions. Spring thaw period have a significant effect on the N<sub>2</sub>O emissions. During spring thaw period, the N<sub>2</sub>O emissions reached 15106  $\mu\text{g m}^{-2} \text{h}^{-1}$  from agricultural ecosystems (Flesch et al., 2018) and the cumulative N<sub>2</sub>O emissions reached 26 kg N ha<sup>-1</sup> over a 35 d spring thaw period (Dunmola et al., 2010), which could account for more than 70% of annual N<sub>2</sub>O emissions (Fu et al., 2018; Wolf et al., 2010; Wu et al., 2010; Yang et al., 2015). Thus, the cumulative N<sub>2</sub>O emission from spring thaw period was an important part of the annual budget. However, the cumulative N<sub>2</sub>O emissions during the nongrowing season were contribution 15.63-33.00% to annual N<sub>2</sub>O emissions in the permafrost region, which was not significantly as previous studies.

Reference

Dunmola, A. S., Tenuta, M., Moulin, A. P., Yapa, P., and Lobb, D. A.: Pattern of greenhouse gas emission from a Prairie Pothole agricultural landscape in Manitoba, Canada, *Can J Soil Sci*, 90, 243-256, <https://doi.org/10.4141/cjss08053>, 2010.

Printer-friendly version

Discussion paper

Flesch, T. K., Baron, V. S., Wilson, J. D., Basarab, J. A., Desjardins, R. L., Worth, D., and Lemke, R. L.: Micrometeorological measurements reveal large nitrous oxide losses during spring thaw in Alberta, *Atmosphere*, 9, 128, <https://doi.org/10.3390/atmos9040128>, 2018.

Fu, Y. F., Liu, C. Y., Lin, F., Hu, X. X., Zheng, X. H., Zhang, W., and Ca, G. M.: Quantification of year-round methane and nitrous oxide fluxes in a typical alpine shrub meadow on the Qinghai-Tibetan Plateau, *Agr Ecosyst Environ*, 255, 27-36, [10.1016/j.agee.2017.12.003](https://doi.org/10.1016/j.agee.2017.12.003), 2018.

Wolf, B., Zheng, X. H., Brueggemann, N., Chen, W. W., Dannenmann, M., Han, X. G., Sutton, M. A., Wu, H. H., Yao, Z. S., and Butterbach-Bahl, K.: Grazing-induced reduction of natural nitrous oxide release from continental steppe, *Nature*, 464, 881-884, [10.1038/nature08931](https://doi.org/10.1038/nature08931), 2010.

Wu, X., Brueggemann, N., Gasche, R., Shen, Z. Y., Wolf, B., and Butterbach-Bahl, K.: Environmental controls over soil-atmosphere exchange of N<sub>2</sub>O, NO, and CO<sub>2</sub> in a temperate Norway spruce forest, *Global Biogeochem Cy*, 24, GB2012, <https://doi.org/10.1029/2009GB003616>, 2010.

Yang, X. M., Chen, H. Q., Gong, Y. S., Zheng, X. H., Fan, M. S., and Kuzyakov, Y.: Nitrous oxide emissions from an agro-pastoral ecotone of northern China depending on land uses, *Agr Ecosyst Environ*, 213, 241-251, [10.1016/j.agee.2015.08.011](https://doi.org/10.1016/j.agee.2015.08.011), 2015.

6. L315-317, positively correlated with air temperature ( $P < 0.05$ ) and C/N<sub>0-10</sub> ( $P < 0.05$ ) and negatively correlated with pH<sub>0-10</sub> ( $P < 0.01$ ), pH<sub>10-20</sub> ( $P < 0.05$ ), NO<sub>3</sub><sup>-</sup>-N<sub>0-10</sub> ( $P < 0.05$ ), NO<sub>3</sub><sup>-</sup>-N<sub>10-20</sub> ( $P < 0.05$ ), and TN<sub>0-10</sub> ( $P < 0.05$ ). how did you test the effects? individually or combined?

The authors' answer:

The Pearson's correlation analysis was used to test the correlations between the N<sub>2</sub>O emissions and environmental factors. The data were individually, not combined.

Printer-friendly version

Discussion paper

7. L351, Fig. 2, still very low explanatory power.... what about the very low values, which time period is this?

The authors' answer:

Yes, the adjusted R<sup>2</sup> were low. The soil temperature could explain 3.01-3.27% temporal variation of N<sub>2</sub>O emissions from LL swamp forest during the nongrowing season.

8. L365, Table 5, wouldn't a season specific analysis be useful? lots of parameters but low explanatory power, which driver is missing...?

The authors' answer:

Thank you for your meaningful suggestion. According to your suggestion, we used the stepwise multiple linear regression analyses to quantify the relationship between N<sub>2</sub>O emissions and environmental factors during the growing season. The explanatory power of the stepwise multiple linear regression was higher than that of multivariate regression. Please see table 5. We also revised the relevant abstract (Please see lines 42-45), result (Please see lines 401-410), discussion (Please see lines 559-573), and conclusion section (Please see lines 598-599).

Discussion

1. L370, 4.1 Soil temperature controls the mean N<sub>2</sub>O emissions from the different periods, wouldn't a season specific analysis be useful?

The authors' answer:

Thank you for your meaningful suggestion. The anonymous referee 1 agrees with your comments that the soil temperature controls the mean N<sub>2</sub>O emissions from the different periods; there is no need for discussion. We have restructured the discussion. According to anonymous referee 1 comments, the 4.1 section discussed the effects of swamp forest types on N<sub>2</sub>O emissions during the nongrowing season. We have rewritten the 4.1 section. According to your comments, the season specific analysis was discussed

in the 4.3 section.

2. L389, The annual N<sub>2</sub>O emissions showed significant temporal variations in grasslands. here is the focus on grasslands not very clear as you investigated swamp forests - I see that there aren't many other studies, yet you would are to explain on why this is comparable or not?

The authors' answer:

Thank you for your suggestion. To our knowledge, the N<sub>2</sub>O emissions from permafrost region were reported from peatland (Marushchak et al., 2011; Palmer & Horn, 2012; Repo et al., 2009), tundra (Lamb et al., 2011), forest (Takakai et al., 2008), grassland (Takakai et al., 2008), alpine meadow (Chen et al., 2017), thermokarst (Mu et al., 2017), beach (Zhu et al., 2012), and lake edge (Gregorich et al., 2006). Except for our previous studies, N<sub>2</sub>O emissions from swamp forests were not reported in other permafrost region (Gao et al, 2019a, 2019b). Thus, we selected the permafrost ecosystems in the permafrost region, which could be not the swamp forest.

#### Reference

Chen, X. P., Wang, G. X., Zhang, T., Mao, T. X., Wei, D., Hu, Z. Y., and Song, C. L.: Effects of warming and nitrogen fertilization on GHG flux in the permafrost region of an alpine meadow, *Atmos Environ*, 157, 111-124, <https://doi.org/10.1016/j.atmosenv.2017.03.024>, 2017.

Gao, W. F., Yao, Y. L., Gao, D. W., Wang, H., Song, L. Q., Sheng, H. C., Cai, T. J., and Liang, H.: Responses of N<sub>2</sub>O emissions to spring thaw period in a typical continuous permafrost region of the Daxing'an Mountains, northeast China, *Atmos Environ*, 214, 116822, <https://doi.org/10.1016/j.atmosenv.2019.116822>, 2019a.

Gao, W. F., Yao, Y. L., Liang, H., Song, L. Q., Sheng, H. C., Cai, T. J., and Gao, D. W.: Emissions of nitrous oxide from continuous permafrost region in the Daxing'an Mountains, Northeast China, *Atmos Environ*, 198, 34-45,

BGD

Interactive  
comment

Printer-friendly version

Discussion paper

<https://doi.org/10.1016/j.atmosenv.2018.10.045>, 2019b.

Gregorich, E. G., Hopkins, D. W., Elberling, B., Sparrow, A. D., Novis, P., Greenfield, L. G., and Rochette, P.: Emission of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from lakeshore soils in an Antarctic dry valley, *Soil Biol Biochem*, 38, 3120-3129, [10.1016/j.soilbio.2006.01.015](https://doi.org/10.1016/j.soilbio.2006.01.015), 2006.

Lamb, E. G., Han, S., Lanoil, B. D., Henry, G. H. R., Brummell, M. E., Banerjee, S., and Siciliano, S. D.: A High Arctic soil ecosystem resists long-term environmental manipulations, *Global Change Biol*, 17, 3187-3194, <https://doi.org/10.1111/j.1365-2486.2011.02431.x>, 2011.

Marushchak, M. E., Pitkamaki, A., Koponen, H., Biasi, C., Seppala, M., and Martikainen, P. J.: Hot spots for nitrous oxide emissions found in different types of permafrost peatlands, *Global Change Biol*, 17, 2601-2614, <https://doi.org/10.1111/j.1365-2486.2011.02442.x>, 2011.

Mu, C. C., Abbott, B. W., Zhao, Q., Su, H., Wang, S. F., Wu, Q. B., Zhang, T. J., and Wu, X. D.: Permafrost collapse shifts alpine tundra to a carbon source but reduces N<sub>2</sub>O and CH<sub>4</sub> release on the northern Qinghai-Tibetan Plateau, *Geophys Res Lett*, 44, 8945-8952, <https://doi.org/10.1002/2017GL074338>, 2017.

Palmer, K., and Horn, M. A.: Actinobacterial nitrate reducers and proteobacterial denitrifiers are abundant in N<sub>2</sub>O-metabolizing tundra peat, *Appl Environ Microb*, 78, 5584-5596, <https://doi.org/10.1128/aem.00810-12>, 2012.

Repo, M. E., Susiluoto, S., Lind, S. E., Jokinen, S., Elsakov, V., Biasi, C., Virtanen, T., and Martikainen, P. J.: Large N<sub>2</sub>O emissions from cryoturbated peat soil in tundra, *Nat Geosci*, 2, 189-192, <https://doi.org/10.1038/NGEO434>, 2009.

Takakai, F., Desyatkin, A. R., Lopez, C. M. L., Fedorov, A. N., Desyatkin, R. V., and Hatano, R.: CH<sub>4</sub> and N<sub>2</sub>O emissions from a forest-alas ecosystem in the permafrost taiga forest region, eastern Siberia, Russia, *J Geophys Res-Biogeosci*, 113, G02002,

**BGD**

Interactive  
comment

Printer-friendly version

Discussion paper

<https://doi.org/10.1029/2007JG000521>, 2008.

Zhu, R. B., Chen, Q. Q., Ding, W., and Xu, H.: Impact of seabird activity on nitrous oxide and methane fluxes from High Arctic tundra in Svalbard, Norway, *J Geophys Res-Biogeosci*, 117, 1703-1705, <https://doi.org/10.1029/2012jg002130>, 2012.

3. L392-393, The N<sub>2</sub>O was taken up during the freezing period, taken up by what - is it that primarily due to diffusion into snow or similar?

The authors' answer:

The mechanism of N<sub>2</sub>O absorption in marsh during the freezing period probably results from frozen soil absorbing atmospheric N<sub>2</sub>O through frost-induced cracks and perhaps the anaerobic environment was too strong and reduce N<sub>2</sub>O to N<sub>2</sub> by reductase enzymes (Du et al., 2006).

Reference

Hao, Q. J., Wang, Y. S., Song, C. C., and Huang, Y.: Contribution of winter fluxes to the annual CH<sub>4</sub>, CO<sub>2</sub> and N<sub>2</sub>O emissions from freshwater marshes in the Sanjiang Plain, *J Environ Sci-China*, 18, 270-275, <https://doi.org/10.3321/j.issn:1001-0742.2006.02.012> 2006.

4. L394, specified periods, here you mean seasons?

The authors' answer:

The specified periods means that the freezing period, thawing period, and growing season.

5. L395, These trends were also observed in the permafrost region. irrelevant, because you focus on permafrost regions

The authors' answer:

Thank you for your suggestion. We have deleted it.

BGD

Interactive  
comment

Printer-friendly version

Discussion paper

6. L395-396, In the “hot spots” of N<sub>2</sub>O emission from permafrost region, which are those?

The authors' answer:

The “hot spots” of N<sub>2</sub>O emission from permafrost region were bare peat circles in the arctic and subarctic (Marushchak et al., 2011; Repo et al., 2009). The nongrowing season contributed by 20–69% to the annual emissions from the peat circles, which were  $1.40 \pm 0.15$  g N<sub>2</sub>O m<sup>-2</sup>.

Reference

Marushchak, M. E., Pitkamaki, A., Koponen, H., Biasi, C., Seppala, M., and Martikainen, P. J.: Hot spots for nitrous oxide emissions found in different types of permafrost peatlands, *Global Change Biol*, 17, 2601-2614, <https://doi.org/10.1111/j.1365-2486.2011.02442.x>, 2011.

Repo, M. E., Susiluoto, S., Lind, S. E., Jokinen, S., Elsakov, V., Biasi, C., Virtanen, T., and Martikainen, P. J.: Large N<sub>2</sub>O emissions from cryoturbated peat soil in tundra, *Nat Geosci*, 2, 189-192, <https://doi.org/10.1038/NGEO434>, 2009.

7. L396-398, high N<sub>2</sub>O emissions were observed during the nongrowing season in the bare peatland, which contributed 20–69% to the annual emissions from the bare peatland (Marushchak et al., 2011). okay but again you are not looking at peatlands but swamp forest

The authors' answer:

Thank you for your suggestion. Except for our previous studies, N<sub>2</sub>O emissions from swamp forests were not reported in other permafrost region. Thus, we select the permafrost ecosystems from the permafrost region, which could be not the swamp forest.

8. L400, were mainly negligible, everytime and would this also be relevant for your study?



The authors' answer:

Thank you for your suggestion. We have deleted it.

9. L403-405, However, the drivers of N<sub>2</sub>O emissions between the nongrowing season and growing season were not clear in the permafrost region. says who?

The authors' answer:

This was what we concluded from previous studies. We summarized the studies on N<sub>2</sub>O emissions from permafrost regions. We found that the difference of N<sub>2</sub>O emissions between nongrowing season and growing season and their drivers was not reported before in the permafrost region.

10. L405-410, Our results showed that the N<sub>2</sub>O emissions were the highest during the growing season and lowest during the nongrowing season. The mean N<sub>2</sub>O emissions from the growing season were significantly higher than the emissions from the winter in the LL and B sites, whereas the N<sub>2</sub>O emissions during the spring thaw period was not significantly different from growing season and winter in the three swamp forests (Fig. 3). this is a repetition of the results.

The authors' answer:

Thank you for your suggestion. We deleted this part in the discussion section and integrated it to the results section. Please see lines 308-311.

11. L418-421, We found that the mean N<sub>2</sub>O emissions of the three specified periods were significantly positively correlated with soil temperature at 5, 10, and 15 cm, which could explain 91.36–94.07, 91.97–95.92, and 81.71–92.85% of the temporal variation of mean N<sub>2</sub>O emissions in the LL, LC, and B sites, respectively (Fig. 4). how or why is this different from figure 2?

The authors' answer:

In the figure 2, the linear models were used to quantify the relationship between N<sub>2</sub>O

[Printer-friendly version](#)[Discussion paper](#)

emission and soil temperature. We found that the temporal variation of N<sub>2</sub>O emissions were significantly positively correlation with soil temperature during the nongrowing season and entire period. Thus, the soil temperature controlled the temporal variation of soil N<sub>2</sub>O emissions in the permafrost region. The Figure 4 focused on the drivers of mean N<sub>2</sub>O emissions among the three specific periods (the winter, spring thaw period, and growing season). The soil temperature drivers the difference of N<sub>2</sub>O emissions among winter, spring thaw period, and growing season. Thus, the Figure 2 and the Figure 4 was different.

12. L442, Figure 4, Which is which period? panel identifiers are mission, are these aggregated values and if so how were these aggregated? are all three soil temperature levels necessary? Which depth is the most relevant?

The authors' answer:

Thank you for your meaningful suggestion. The legend of Figure 4a should be growing season, winter, and spring thaw period. The spots with black, light gray, and gray represent the growing season, winter, and spring thaw period, respectively. We have revised it. Please see Figure 4. Yes, these were aggregated values. We calculated the average N<sub>2</sub>O emission during the growing season, winter, and spring thaw period in each base collar. And then the average N<sub>2</sub>O emission from each collar was used to compare the difference of three specific period. According to the adjust R<sup>2</sup>, the soil temperature at 5 cm depth were the most relevant with N<sub>2</sub>O emissions in the LC and B sites. While, soil temperature at 10 cm depth was the most relevant with N<sub>2</sub>O emissions in the LL sites. Thus, we think that all three soil temperature levels are necessary.

13. L448, was, were.

The authors' answer:

We have revised it.

14. L450, 14 kg ha<sup>-1</sup>, units.

The authors' answer:

Thank you for your suggestion. We replace the kg ha<sup>-1</sup> with kg N ha<sup>-1</sup>.

15. L451-452, released approximately 0.1 Tg yr<sup>-1</sup> N<sub>2</sub>O emissions to the atmosphere in the bare peat region of the Arctic, accounting for up to 0.6% of the global annual N<sub>2</sub>O emissions. how was the upscaling done?

The authors' answer:

Global N<sub>2</sub>O emissions from the peat circles were estimated using the mean 4% coverage of peat circles in the peat plateau, a 20% coverage of the peat plateaus in the Arctic (Walker et al., 2005; Virtanen et al., 2004) and the total land area of the tundra zone of 7:34 × 10<sup>6</sup> km<sup>2</sup> (Matthews, 1983). The finally compared the global annual N<sub>2</sub>O estimate with the Arctic methane emissions (20 Tg CH<sub>4</sub> yr<sup>-1</sup>) using the GWP approach (Christensen, 1993).

#### Reference

Christensen, T. R.: Methane emission from arctic tundra, *Biogeochemistry*, 21, 117-139, [10.1007/BF00000874](https://doi.org/10.1007/BF00000874), 1993.

Matthews, E.: Global vegetation and land use: new high-resolution data bases for climate studies, *Journal of Applied Meteorology*, 22, 474-487, [https://doi.org/10.1175/1520-0450\(1983\)022<0474:GVALUN>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<0474:GVALUN>2.0.CO;2), 1983.

Virtanen, T., Mikkola, K., and Nikula, A.: Satellite image based vegetation classification of a large area using limited ground reference data: a case study in the Usa Basin, north-east European Russia, *Polar Res*, 23, 51-66, <https://doi.org/10.3402/polar.v23i1.6266>, 2004.

Walker, D., Raynolds, M., Daniëls, F., Einarsson, E., Elvebakk, A., Gould, W., Katenin, A., Kholod, S., Markon, C., Melnikov, E., Moskalenko, N., Talbot, S., Yurtsev, B.,

and Team, T.: The Circumpolar Arctic vegetation map, *J Veg Sci*, 16, 267-282, <https://doi.org/10.1111/j.1654-1103.2005.tb02365.x>, 2005.

16. L455-456, However, the contribution of the nongrowing season to the annual N<sub>2</sub>O budget is uncertain in the permafrost region. is or was - after your study I thought it would be less uncertain.

The authors' answer:

Thank you for your suggestion. We have rewritten it. Please see lines 465-467.

17. L458-460, The cumulative N<sub>2</sub>O emissions during the nongrowing season ranged from 0.89 to 1.44 kg ha<sup>-1</sup>, which contributed to 41.96–53.73% of the annual budget in the permafrost region of the Daxing'an Mountains. redundant information.

The authors' answer:

Thank you for your suggestion. We have deleted the redundant information.

18. L470, frigid terrestrial ecosystems, lots of repetitions.

The authors' answer:

Thank you for your comments. We have deleted the repetition.

19. L470-472, When the pulse of N<sub>2</sub>O emissions occurred during the spring thaw period, emissions from the non-growing seasons dominated (67–74%) the annual total emissions. is this now this study or the study by Fu et al - unclear

The authors' answer:

Thank you for your suggestion. This study was by Fu et al., (2018). We have revised it.

20. L476-480, The cumulative N<sub>2</sub>O emissions during the spring thaw period ranged from 0.35 to 0.66 kg ha<sup>-1</sup> and contributed 15.61 to 33.00% of the annual budget in the permafrost region of the Daxing'an Mountains, and these ranges were generally lower than the emissions during the winter and growing season. you state many dif-

ferent numbers concerning the contribution of non growing season N<sub>2</sub>O emissions to the annual budget - please clarify what is what. Also you always state the Daxing'an Mountains - however to my understanding you looked at swamp forests "only" in the region, correct?

The authors' answer:

Thank you for your suggestion. According to previous studies, the pulse of N<sub>2</sub>O emissions during the spring thaw period had significantly influence on the contribution of the nongrowing season to the total annual N<sub>2</sub>O budget (Fu et al., 2018; Li et al., 2012). We want to state that there was no pulse of N<sub>2</sub>O emissions were occurred in the present study. Thus, the contribution of nongrowing season N<sub>2</sub>O emissions to the annual budget were lower. We have rewritten it. Please see lines 504-508. Yes, it was swamp forests. We have revised it.

#### Reference

Fu, Y., Liu, C., Lin, F., Hu, X., Zheng, X., Zhang, W., and Ca, G.: Quantification of year-round methane and nitrous oxide fluxes in a typical alpine shrub meadow on the Qinghai-Tibetan Plateau, *Agr Ecosyst Environ*, 255, 27-36, 636 <https://doi.org/10.1016/j.agee.2017.12.003>, 2018.

Li, K. H., Gong, Y. M., Song, W., Lv, J. L., Chang, Y. H., Hu, Y. K., Tian, C. Y., Christie, P., and Liu, X. J.: No significant nitrous oxide emissions during spring thaw under grazing and nitrogen addition in an alpine grassland, *Global Change Biol*, 18, 2546-2554, <https://doi.org/10.1111/j.1365-2486.2012.02704.x>, 2012.

21. L483-484, which were not the same as the cumulative N<sub>2</sub>O emissions in the three swamp forests. why? what do you mean?

The authors' answer:

Thank you for your suggestion. We mean that the cumulative N<sub>2</sub>O emissions were not lowest in the winter and highest in the growing season as the mean N<sub>2</sub>O emissions.

Because of the length of the season had significantly affected on the cumulative N<sub>2</sub>O emissions. We have revised it. Please see lines 511-512.

22. L495-499, Our results confirmed that half of the N<sub>2</sub>O emissions were released during the nongrowing season, indicating that the N<sub>2</sub>O emissions during the nongrowing season cannot be ignored in the permafrost regions. In the future, the N<sub>2</sub>O emissions of the nongrowing season should be emphasized in the permafrost region, especially in the context of global climate change. this is the key message and the parts before are quite chaotic I must admit - thus kindly structure logically.

The authors' answer:

Thank you for your meaningful suggestion. According to your suggestion, we have revised the logical structure before this part.

23. L504, by air temperature, soil temperature, before it was stated that soil temperature was the most important driving factor

The authors' answer:

Yes, the soil temperature was the most important driving factor in our study. In this part, we summarized the driving factors of N<sub>2</sub>O emissions from the permafrost region during the growing season. We found that the N<sub>2</sub>O emissions from permafrost region were influenced by air temperature, soil temperature, water table level, soil moisture, precipitation, pH, NH<sub>4</sub><sup>+</sup>-N, NO<sub>3</sub><sup>-</sup>-N, TOC, gross N mineralization, N content, and C/N ratio during the growing season. We have revised it. Please see line 530-532.

24. L508-509, However, the drivers of nongrowing season N<sub>2</sub>O emission remain unknown in the permafrost region. why?

The authors' answer:

Previous studies mainly focused on the N<sub>2</sub>O emissions and their driving factors during the growing season. However, seldom studies reported the N<sub>2</sub>O emissions during

nongrowing seasons in the permafrost region. The relationship between nongrowing season N<sub>2</sub>O emissions and environment factors was not reported. Thus, the drivers of nongrowing season N<sub>2</sub>O emission remain unknown in the permafrost region.

25. L511-512, are very complex and can be produced by nitrification, nitrifier denitrification, and denitrification in the permafrost region. not just in the permafrost region. please look at Butterbach-Bahl et al 2013 PToRS for a review on the processes of N<sub>2</sub>O.

The authors' answer:

Thank you for your meaningful suggestion. We summarized the produce pathway of N<sub>2</sub>O emissions from permafrost regions. We found that the N<sub>2</sub>O emissions from permafrost region were mainly came from nitrification, denitrification, and nitrifier denitrification (Ma et al., 2007; Gil et al., 2017; Siljanen et al., 2019). Indeed, the N<sub>2</sub>O emissions could produce by nitrification, denitrification, nitrifier denitrification, coupled nitrification-denitrification, co-denitrification, and dissimilatory nitrate reduction to ammonium (Butterbach-Bahl et al., 2013). However, except for nitrification, denitrification, and nitrifier denitrification, other pathway of N<sub>2</sub>O emissions was not reported in the permafrost region. Reference

Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philosophical Transactions of the Royal Society B Biological Sciences*, 368, 20130122, 10.1098/rstb.2013.0122 2013.

Gil, J., Pérez, T., Boering, K., Martikainen, P. J., and Biasi, C.: Mechanisms responsible for high N<sub>2</sub>O emissions from subarctic permafrost peatlands studied via stable isotope techniques, *Global Biogeochem Cy*, 31, 172-189, <https://doi.org/10.1002/2015GB005370>, 2017.

Ma, W. K., Schautz, A., Fishback, L. A. E., Bedard-Haughn, A., Farrell, R. E., and Siciliano, S. D.: Assessing the potential of ammonia oxidizing bacteria to produce nitrous

oxide in soils of a high arctic lowland ecosystem on Devon Island, Canada, *Soil Biol Biochem*, 39, 2001-2013, <https://doi.org/10.1016/j.soilbio.2007.03.001>, 2007.

Siljanen, H., J.E. Alves, R., Ronkainen Jussi, G., Lamprecht, R., Bhattarai, H. R., Bagnoud, A., Marushchak, M., Martikainen, P., Schleper, C., and Biasi, C.: Archaeal nitrification is a key driver of high nitrous oxide emissions from arctic peatlands, *Soil Biol Biochem*, 137, 107539, <https://doi.org/10.1016/j.soilbio.2019.107539>, 2019.

26. L538-539, The N<sub>2</sub>O emissions were major produced by denitrification in the permafrost region. how did you determine this?

The authors' answer:

According to previous studies, the denitrification can be the predominant pathway for N<sub>2</sub>O production during the soil moisture content exceed 70% (Butterbach-Bahl et al., 2013). We have cited the reference.

Reference

Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philosophical Transactions of the Royal Society B Biological Sciences*, 368, 20130122, [10.1098/rstb.2013.0122](https://doi.org/10.1098/rstb.2013.0122) 2013.

27. L549-552, Soil temperature was the major limiting factor related to annual N<sub>2</sub>O emissions in the permafrost region. Except for soil temperature, the N<sub>2</sub>O emissions from the permafrost region were also affected by pH, NO<sub>3</sub><sup>-</sup>-N, TN, and C/N ratio. this section is again quite unstructured or is more like a loose list of information without the logical interconnection - thus substantial rewriting is needed

The authors' answer:

Thank you for your meaningful suggestion. We have rewritten this part. Except for soil temperature, the N<sub>2</sub>O emissions from the permafrost region were also affected by pH,



NO<sub>3</sub>–N, TN, and C/N ratio. Voigt et al., (2020) summary the studies on nitrous oxide emissions from permafrost-affected soils, which found that the high N<sub>2</sub>O emissions in pristine permafrost regions can be expected in areas with a sparse to absent vegetation cover, a low to intermediate C:N ratio, high C and high mineral N content (in particular, NO<sub>3</sub>–N), in combination with relatively high temperatures and a favourable moisture content. We also found that the high NO<sub>3</sub>–N content, high TOC, and relatively high temperatures would increase the N<sub>2</sub>O emissions in the permafrost region of Daxing'an Mountains. Please see lines 579-587. Reference

Voigt, C., Marushchak, M. E., Abbott, B. W., Biasi, C., Elberling, B., Siciliano, S. D., Sonnentag, O., Stewart, K. J., Yang, Y., and Martikainen, P. J.: Nitrous oxide emissions from permafrost-affected soils, *Nature Reviews Earth & Environment*, 1, 420-434, <https://doi.org/10.1038/s43017-020-0063-9>, 2020.

### Conclusions

1. L555-556, The N<sub>2</sub>O emissions from the nongrowing season were quantified in the permafrost region. its not representative for the whole permafrost region I am afraid.

The authors' answer:

Thank you for your meaningful suggestion. We have revised it. The permafrost region was limited to the Daxing'an Mountains.

2. L560-561, which cannot be ignored. In the different periods studied, N<sub>2</sub>O emissions had different key limiting factors in the permafrost region. distinguish between factor and driver variables, again its not the whole permafrost region

The authors' answer:

Thank you for your meaningful suggestion. We have replaced the factors with driver variables. We also defined the permafrost region as the permafrost region of Daxing'an Mountains.

3. L562-564, and the annual N<sub>2</sub>O emissions were driven by soil temperature, whereas the growing season N<sub>2</sub>O emissions were affected by soil temperature, water table level, and their interaction. this is not a conclusion? and whats next?

The authors' answer:

Thank you for your meaningful suggestion. Yes, this was not the conclusion. We added the conclusion after this part. Our results indicated that nongrowing season N<sub>2</sub>O emissions were an important part of the annual budget, which were of great significance for the accurate assessment of regional climate change and the impact of global climate change on permafrost region. Please see lines 600-602.

Thank you for your comments; we are happy to make additional revisions if needed.

Sincerely,

The authors,

Dr. Da-Wen Gao,

School of Environment and Energy Engineering, Beijing University of Civil Engineering and Architecture, Beijing 100044, China

E-mail address: gaodw@hit.edu.cn

---

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-305>, 2020.

Printer-friendly version

Discussion paper