

Reviewer 1:

- Overall, I found the paper interesting but perhaps better suited to a more remote-sensing oriented journal as it is difficult to distill a strong biological story in the comparisons presented.
 - We appreciate that the reviewer thinks that this article may be better suited in a remote-sensing journal, but think that the biogeosciences community would be interested in a new way to monitor phenological change. We have already published the methodological details of our GOES algorithm in a remote sensing journal, and to make this paper more accessible to non-remote-sensors we will reduce the remote-sensing jargon. Ultimately our interest here is in the biological process of phenology and not the satellite specifics.
- Equation 3: I think that the top case should be for $t \geq k$ and the bottom for $t < k$
 - Changed. (Line 174).
- Did you investigate the 0.6040 artefact in the raw bands rather than just the indices? I wonder if it is also creating less obvious errors in other parts of the data - e.g. in S2 there are some outliers just after the series of values that were removed. Are both the red and NIR bands fixed at some value for these and are those invalid NIR/red values wherever they occur or only when they contribute to a ratio of 0.6040?
 - We thank the reviewer for the suggestion on how to further eliminate noise by looking at the raw red and near-infrared (NIR) values for the observations that had an NDVI value of 0.6040. We updated the manuscript (line 135) to further clarify that all of the NDVI values of 0.6040 only occurred in the early morning and the evening. Based on the days and sites that we looked at, the NDVI value of 0.6040 seems to occur for multiple different combinations of red and NIR values (e.g., $R=1.87e-5$, $NIR=7.57e-5$; $R=-2.80e-4$, $NIR=-1.14e-3$; $R=3.18e-4$, $NIR=1.29e-3$; $R=1.95e-5$, $NIR=7.90e-5$). It seems that when these reflectances occur they are usually in pairs that result in an NDVI value of 0.6040. When we saw them occurring elsewhere they were still limited to the noisy morning and evening sections. We already filtered most of this out by removing the NDVI values that occurred before 1.5 hours after sunrise or 1.5 hours before sunset. In future work, we will consider further filtering based on the reflectance bands, but almost all of this noise appeared to be already filtered out by removing the morning and evening values.
- Would there be a way to automate the removal of the 68 days to further automate the algorithm?
 - Added sentence on lines 153-155.
- Better legends on the supplemental material would improve readability.
 - We added axes to the supplementary figures and updated legends.
- The authors assert the GOES data can provide real-time estimates of phenological transitions whereas MODIS cannot due to the temporal frequency. However as I understand it the algorithm was fit on the entire time series at once, not subsetting down particular portions of the year. It is possible that the higher temporal frequency of the GOES data would allow for real-time estimation, but I think to demonstrate this one would need to iteratively refit using data only up to Jan, Feb, March (for example) and compare the estimated transition dates between the two datasets/algorithms. It is pos-

sible none would fit, though adding an informed prior on c and d based on the year(s) prior data might(?) provide enough information for them to converge without the full season of data.

- We thank the reviewer for pointing this out – it is definitely true that this specific curve-fitting approach would not be the best option for real-time applications. We clarified this by slightly changing abstract (line 21) and adding a sentence to the discussion (lines 357-360) that points out alternative approaches, such as detecting the passage of predefined thresholds or more sophisticated iterative data assimilation and forecasting approaches, like we have previously employed in Viskari et al. 2015 Ecological Applications. Indeed, adding GOES to our iterative phenology forecast was the original motivation for much of this work.
- What would implications be of NOT using the 16 composited NDVI/EVI from modis and instead using the native temporal resolution data and preprocessing as per the GOES? Would using the same snowcover mask on both datasets align the results? Is there any way to make the raw data more comparable to disentangle the cause of the differences (bands vs temporal resolution vs pre-processing)?
 - The raw data are not comparable between GOES and MODIS because GOES is geostationary and MODIS is on sun-synchronous satellites. Unlike GOES, MODIS changes viewing angles between measurements. The advantage of geostationary data is not just the higher frequency of measurements, but also the consistent viewing angle. The MODIS composite product is created not just to average to reduce noise, but to consider the multiple viewing angles of MODIS observations. To the best of our knowledge there are also no good error estimates for the daily MODIS data, which would also affect our ability to compare them. There has been a lot of calibration and validation that has gone into the MODIS products so returning to the raw MODIS data would be difficult and very time consuming – at the moment we don't think this would be worth the effort.

Reviewer 2:

- The authors stated that the benefit of the GOES series is the high temporal resolution (sub-daily). The authors used the comparison to the PhenoCam data (and better performance of the GOES data) to support this claim. This claim to me is mostly correct with one exception: the comparison between GOES time-series and MODIS 16-day time-series is not fair because there is a daily MODIS reflectance product at 500 m resolution that takes into account of the impact of sun-sensor geometry – the MOD43A4 product. The results have demonstrated that GOES ABI is likely better than MODIS 16-day products in capturing spring phenology. Still, proper discussion (and some texts in the introduction) on the daily MODIS product is necessary. The downside of the MODIS 16-day product could potentially be addressed with a daily MODIS product.
 - We updated the abstract (lines 14-15) to emphasize that these results are for the studied MODIS products. We added more to the introduction (lines 60-65) on why we're focusing on the 16-day product and to the discussion (lines 291-295).
- Line 51: Adding a sentence on why changing viewing angle limits the temporal resolution could be useful for non-remote-sensing readers.
 - Added at lines 54-59.

- Line 65: “subject to the same temporal limitations” – but both are subject to cloud impact. No data when there is cloud cover.
 - We added a sentence pointing out that because GOES makes measurements throughout the day it is better poised to be able to capture times within a day that may be cloud free (lines 87-88). In our previous work, which is mentioned here, we showed that we could estimate daily NDVI reliably on most days even if there was cloud cover at a specific time (e.g. noon) by exploiting the predictable diurnal pattern of GOES NDVI. Obviously, there are still days with complete cloud cover, but these are much fewer for GOES than MODIS.
- Line 144: This approach misses two main features in the GCC curve: The peak in the later spring and early summer, and the gradual decline in GCC in the summer to fall. Similarly, the equation does not fit the NDVI data well (misses the summer decline in NDVI). See the following paper and the Klosterman et al. that was cited:
 - Elmore, A.J., Guinn, S.M., Minsley, B.J. and Richardson, A.D., 2012. Landscape controls on the timing of spring, autumn, and growing season length in mid-Atlantic forests. *Global Change Biology*, 18(2), pp.656-674.
 - Using the methods from Elmore et al. or Klosterman et al. could potentially improve the late spring transition date estimation (and the mismatch between satellite the Phe- noCam data).
 - We added sentences to discussion (lines 288-291) pointing to this as a future direction, but also noting that any biases are likely shared across the different sensors.
- Line 230: the use of “prematurely” gives a sense of that MODIS incorrectly estimates the start of spring. However, another possibility is the green-up of the understory may cause MODIS vegetation indices to increase even when Phenocam data do not show any changes. Richardson and O’Keefe (2009) showed that understory spring is about 10-20 days earlier at Harvard Forest:
 - We changed “prematurely” to “earlier” (line 257) and further clarified this in lines 258-259.
- Line 264: It is unclear what the difference is between “canopy greenness” and “leaf presence and canopy structure”. Both GCC and vegetation indices (EVI and NDVI) are affected by these factors, but in different ways: GCC (R, G, B) can be affected by these factors differently compared with EVI and NDVI (which has a NIR band). I suggest rewriting this sentence.
 - We tried to clarify this in lines 301-303.
- Section 4.2.: the uncertainties in the fall phenology estimation could also be attributed to the heterogeneity in the timing of fall phenology (compared with much-synchronized spring phenology). Worth some discussion.
 - Added sentences to lines 305-307.

Improving the monitoring of deciduous broadleaf phenology using the Geostationary Operational Environmental Satellite (GOES) 16 and 17

Kathryn I. Wheeler¹, Michael C. Dietze¹

¹Department of Earth and Environment, Boston University, Boston, MA, 02215, USA

5 Correspondence to: Kathryn I. Wheeler (kiwheel@bu.edu)

Abstract. Monitoring leaf phenology allows for tracking the progression of climate change and seasonal variations in a variety of organismal and ecosystem processes. Networks of finite-scale remote sensing, such as the PhenoCam Network, provide valuable information on phenological state at high temporal resolution, but have limited coverage. To more broadly remotely sense phenology, satellite-based data that has lower temporal resolution has primarily been used (*e.g.*, 16-day MODIS NDVI product). Recent versions of the Geostationary Operational Environmental Satellites (GOES-16 and -17) allow the monitoring of NDVI at temporal scales comparable to that of PhenoCam throughout most of the western hemisphere. Here we [begin to](#) examine the current capacity of this new data to measure the phenology of deciduous broadleaf forests for the first two full calendar years of data (2018 and 2019) by fitting double-logistic Bayesian models and comparing the start, middle, and end of season transition dates to those obtained from PhenoCam and MODIS 16-day NDVI and EVI products. Compared to the [se](#) MODIS [products](#), GOES was more correlated with PhenoCam at the start and middle of spring, but had a larger bias (3.35 ± 0.03 days later than PhenoCam) at the end of spring. Satellite-based autumn transition dates were mostly uncorrelated with those of PhenoCam. PhenoCam data produced significantly more certain (all p -values ≤ 0.013) estimates of all transition dates than any of the satellite sources did. GOES transition date uncertainties were significantly smaller than those of MODIS EVI for all transition dates (all p -values ≤ 0.026), but were only smaller (based on p -value < 0.05) than those from MODIS NDVI for the beginning and middle of spring estimates. GOES will improve the monitoring of phenology at large spatial coverages and is able to provide real-time indicators of phenological change even [when the entire, spring transition period, occurs](#) within the 16-day resolution of these MODIS products.

Deleted: indices

Deleted: for

Deleted: s

Deleted: that might

1 Introduction

The influence of leaf phenology is ubiquitous across many processes and relationships in ecology, local and regional climates, and weather – ranging from leaf-trait relationships, nutrients in leaf litter leachate, surface roughness, transpiration, leaf-spectra relationships, albedo and energy budgets, and annual primary productivity (Alekseychik et al., 2017; Hudson et al., 2018; McKown et al., 2013; Piao et al., 2019; Richardson et al., 2012; Schwartz et al., 2002; Xue et al., 1996; Zhu and Zeng, 2017). In addition, since phenology is often highly sensitive to climatic variables such as temperature and precipitation (Killingbeck, 2004), it has been a primary ecological indicator of climate change (Parmesan and Yohe, 2003). Overall, spring in deciduous

35 forests has been found to advance and autumn has been found to delay (Gao et al., 2019; Liu et al., 2016), but the results are heterogeneous, especially for autumn (Gill et al., 2015; Richardson et al., 2013). These trends in changes are usually on the magnitude of days (*e.g.*, Keenan et al., 2014b, found an advancement in spring of 0.48 ± 0.2 days/year; Parmesan and Yohe, 2003, an advancement of 0.23 days/year). However, trends are often dependent on the phenology index used (Keenan et al., 2014b). This is particularly problematic for autumn where leaf color change often precedes leaf abscission, affecting the similarity of autumn change in greenness indices (*e.g.*, Green Chromatic Coordinate; GCC), the Normalized Difference
40 Vegetation Index (NDVI), and the Enhanced Vegetation Index (EVI). Similarly, observation frequency can be particularly important in spring, where the trend, the interannual variability, and time required for green-up can all be smaller than common satellite data product frequencies.

The longest vegetation phenological records date back to the monitoring of the flowering of Japanese cherry trees in the ninth century (Richardson et al., 2013). Since then many naturalists have tracked phenology in a variety of ecosystems,
45 such as deciduous broadleaf (DB) forests. These human observations, though, are limited in scale and also rely on extensive manpower, time, and consistency. The United States National Phenology Network is able to circumvent many of these challenges by relying on citizen scientist data, but is still limited by the timeliness of observation uploads and the inability to provide full, consistent coverage. Remote sensing techniques, both near-surface and satellite-based, are able to monitor temporal changes in vegetation reflectance at a near-real time and consistent frequency. Near-surface techniques include digital
50 cameras, such as those that are part of the PhenoCam Network, that take repeated imagery of canopies and track how the ratios of red, green, and blue digital numbers change throughout the year (Richardson et al., 2007). The PhenoCam Network includes over 750 site-years of data across different biomes (Richardson et al., 2018b), but it has inherently limited spatial coverage.

Satellites such as Aqua, Terra, Sentinel-2, and Landsat provide full coverage observations of phenological-sensitive indices such as NDVI and EVI. However, in addition to being sensitive to clouds, these satellites are sun-synchronous (*i.e.*,
55 their orbits are set to pass over a specific local time rather than being fixed over a specific location) and, thus, while observations have near global coverage, at any given site they are at a limited frequency. In addition, to be able to cover the earth these orbits are not exactly the same each day and, thus, images are taken from varying viewing angles, which can add considerably complexity to the analysis and interpretation of data (*i.e.*, one needs to deconvolve changes in vegetation state from changes in view angle). Because of this and challenges from frequent clouds, MODIS (Moderate Resolution Imaging
60 Spectroradiometer, which is on Aqua and Terra) NDVI and EVI products are created by compositing data over multi-day periods (*e.g.*, 16 days). This can be interpolated into daily estimates of NDVI and EVI or can be provided as 16-day composites. While the daily interpolated products are sometimes used in phenology (*e.g.*, Ju et al., 2010; Keenan et al., 2014a; Liu et al., 2017), the 16-day composite NDVI and EVI products are also widely used (*e.g.*, Ahl et al., 2006; Hmimina et al., 2013; Richardson et al., 2018b; Zheng and Zhu, 2017) and can be easily accessed through the MODIS web API and the MODISTools
65 R Package (Tuck et al., 2014). This lower temporal resolution results in MODIS NDVI- and EVI- based estimates of phenological transition dates having larger uncertainties than those derived from PhenoCam (Klosterman et al., 2014). Additionally, while this temporal resolution may be adequate for some applications of NDVI, spring transitions and climate

Formatted: Font: Italic

Deleted: , which results in the trade-off of limited temporal resolution due to changes in viewing angles at each pass over.

Deleted: the 16-Day

Formatted: Font: Italic

Formatted: Font: Italic

Moved (insertion) [1]

Moved up [1]: (*e.g.*, Ahl et al., 2006; Hmimina et al., 2013; Richardson et al., 2018b; Zheng and Zhu, 2017)

Deleted: serve as two of the most commonly used sources of satellite phenology data (*e.g.*, Ahl et al., 2006; Hmimina et al., 2013; Richardson et al., 2018b; Zheng and Zhu, 2017)(Ju et al., 2010; Liu et al., 2017).

change induced changes, as already mentioned, can happen at time scales much shorter than this 16-day resolution, which has weakened how correlated the MODIS NDVI- and EVI- observed start of spring estimates are with those from PhenoCam (Filippa et al., 2018; Hufkens et al., 2012; Klosterman et al., 2014; Richardson et al., 2018a). To be able to accurately track phenological transitions and changes at large scales, satellite-based data at a finer temporal resolution is needed.

The United States' National Oceanic and Atmospheric Administration's Geostationary Operational Environmental Satellite (GOES) -16 and -17 are the first satellites in the long-standing GOES series that possess a new sensor, the Advanced Baseline Imager (ABI), that includes the necessary bands to calculate NDVI (Schmit et al., 2016). As the name implies, these satellites (one assumed the position of GOES-East at 75°W in December 2018 and one the position of GOES-West at 137°W in February 2019; Schmit et al., 2016) are geostationary and are thus not subject to many of the same limitations as sun-synchronous satellites because they are able to take frequent measurements across their view with constant viewing angles. While geostationary satellites are still subject to clouds, the higher temporal resolution of potential measurements allows them to have a greater number of non-cloudy measurements than sun-synchronous satellites. GOES collects data every five minutes for the continental U.S. and every ten minutes for much of the western hemisphere. This high frequency data can be noisy in deciduous forests, however, but daily midday NDVI values can be estimated with uncertainty quantifications from statistical models that utilize the characteristic diurnal pattern of NDVI (Wheeler and Dietze, 2019). Additional geostationary satellites that possess the ability to monitor NDVI over other parts of the world include Meteosat over Africa and Europe, and Himawari over east Asia and Oceania.

In this study, we investigated how GOES-16 (and by association GOES-17) compares to commonly-used 16-day MODIS NDVI and EVI products in relation to PhenoCam through estimations of phenology transition dates for DB forests in the eastern U.S. We selected sites within the PhenoCam Network and fit phenological curves for the different data sources (PhenoCam, MODIS NDVI, MODIS EVI, and GOES NDVI) in a Bayesian context for the first full calendar years of data (2018 and 2019). We calculated start, middle, and end of season transition dates and compared those estimates between the different data sources. We hypothesized (1) GOES's higher measurement frequency would generate spring transition date estimates that are more similar than MODIS to PhenoCam; (2) since DB canopy spring transitions often occur faster than autumn, and changes in leaf color and area are more synchronous, spring transition dates would be more similar across the different data sources than the autumn ones; (3) since there exist differences in the sensitivities of different sensors to leaf color versus leaf presence, GOES autumn transition dates would be most similar to MODIS NDVI; and (4) because of the higher data volumes, GOES would produce transition date estimates with lower uncertainties than MODIS.

2 Methods

2.1 Site selection

From the PhenoCam Network, fifteen DB sites were selected to be compared to their associated MODIS and GOES pixels. To attempt to maintain homogeneity in the associated pixels of different spatial resolution (especially since the MODIS pixels do

Deleted: temporal

110 not necessarily fall completely within the GOES pixels), we used Google Earth to exclude PhenoCam sites that were within
the width of a GOES pixel (~1 km) from another land cover type (e.g., grassland, urban, or large water body). Distinguishing
evergreen species using Google Earth is more difficult and, thus, several of the sites do likely have nearby evergreen species
that are included in the same satellite pixels. However, these sites still display predominantly DB phenology curves and, thus,
were still included in this comparison. Specific site locations can be found in Fig. 1 and Table 1. Additional metadata on the
115 sites is available on the PhenoCam Network website (<https://phenocam.sr.unh.edu/webcam/>).

2.2 Data processing

2.2.1 GOES data download and quality control

The study time period of 1 January 2018 through 31 December 2019 was selected due to the availability of new GOES data
for the first two calendar years.

120 ABI L1b radiance values were downloaded from NOAA's Comprehensive Large Array-data Stewardship System for
the GOES channel 3 (near infrared) and channel 2 (red) for the study period (GOES-R Calibration Working Group and GOES-
R Series Program, 2017). After the ABI L2+ clear sky mask (ACM) and data quality flags were applied (GOES-R Algorithm
Working Group and GOES-R Series Program, 2018), radiance values were converted to reflectance factors under the guidance
of the GOES R Product Definition and Users' Guide (<https://www.goes-r.gov/users/docs/PUG-L1b-vol3.pdf>,
125 pp 27-28). Additional information on accessing and processing the data can be found in (Wheeler and Dietze, 2019) and
through our Github repository https://github.com/k-wheeler/NEFI_pheno/tree/master/GOESDiurnalNDVI. To account for
differences in spatial resolution, the four 0.5 km red reflectance factors that fell within a near infrared (NIR) pixel were
averaged together and NDVI was calculated on a 1 km spatial resolution calculated following Eq. (1):

$$NDVI = \frac{\rho_{NIR} - \rho_{Red}}{\rho_{NIR} + \rho_{Red}}, \quad (1)$$

130 where ρ_{NIR} and ρ_{Red} refer to the reflectance factor at the NIR band and red bands, respectively. While GOES does provide a
blue band, which would allow for the calculation of EVI, additional calibration would likely need to be conducted to establish
coefficients needed in the EVI equation, which is outside of the scope of this study. NDVI values that occurred before 1.5
hours after sunrise and after 1.5 hours before sunset (calculated using the Sunalc R package; Thieumel and Elmarhraoui,
2019) were removed due to high noise. Additionally, the NDVI values of 0.6040 were regularly and abnormally present in the
135 dataset [early in the morning or in the evening](#) throughout the study period and, thus, were removed as noise (e.g.,
Supplementary Fig. S1). All calculations were performed in R (R Core Team, 2017).

Deleted:

2.2.2 Daily GOES NDVI estimates

Daily midday GOES NDVI values were estimated using the Bayesian statistical model described in (Wheeler and Dietze,
2019). In summary, this model relies on the characteristic diurnal NDVI pattern for DB pixels of increasing in the morning

(represented with an inverted exponential decrease function) and decreasing in the afternoon (represented with an exponential decrease function), with a changepoint parameter between the two exponential functions (Supplementary Fig. S2). The error model accounts for negative bias in noise due to atmospheric attenuation (*e.g.*, from clouds and aerosols) by calculating the probability that each observation is clear or cloudy and the amount of atmospheric transmissivity. Daily midday NDVI values with 95% credible intervals (CI) were obtained from the GOES data for all days with at least ten observations (*i.e.*, observations where both radiance values had data quality flags of “acceptable” and had an “acceptable” non-cloudy value from the ACM product). We changed the prior on the parameter c (the midday maximum NDVI estimate) from that reported in Wheeler and Dietze (2019) to an uninformative Beta(1,1) instead of Beta(2,1.5). With more data, it was clear that the original prior was incorrectly pulling fits for winter days too high. We also filtered out days that had < 25 observations and did not have observations in at least five different hours. These thresholds were determined by examining various combinations to minimize noise and maintain most of the fitted days. Additionally, after a visual inspection of the diurnal fits and data, 68 days of the total 3645 days ($< 2\%$) that remained after the previous filtering were removed due to poor fits that were heavily influenced by one outlier point. [As this research scales up, implementing and testing automatic quality control methods will become more important, but that is beyond the scope of this paper as here we want to focus on assessing phenological patterns and not remote sensing quality control algorithms.](#)

2.2.3 MODIS and PhenoCam data download

The 250m 16-day NDVI and EVI bands from the MODIS product MOD13Q1 were used for the study period (Didan, 2015). This product was selected because it is easily accessible through the MODISTools R Package (Tuck et al., 2014) and the same temporal resolution for MODIS products has been used in numerous other comparisons between phenology data sources (*e.g.*, Ahl et al., 2006; Hmimina et al., 2013; Richardson et al., 2018b; Zheng and Zhu, 2017). Product data quality flags were applied to MODIS data. Daily midday PhenoCam GCC and standard deviation values were downloaded directly from the PhenoCam website archive (PhenoCam, 2018 and 2019; Seyednasrollah et al., 2019). The PhenoCam at Russell Sage did not collect data for most of 2019 and, thus, we only fit one year of data (2018) to this site.

2.3 Phenology model fitting

Phenological curves were fit for each source of data (GOES NDVI, MODIS NDVI, MODIS EVI, and PhenoCam GCC). Based on the highly cited Zhang et al. (2003) paper, spring and autumn phenological changes were both modeled using a logistic curve calculated following Eq. (2):

$$\mu_t = \frac{c}{1 + e^{a+bt}} + d, \quad (2)$$

where t is the time in days, μ_t is the phenology metric, a and b are fitting parameters, $c + d$ is the maximum value, and d is the winter background value for the metric. The interpretation and realistic limits of the parameters a and b are somewhat obscure,

so we reparametrized the model in terms of the midpoint date (50% change), $M = -a/b$. This gives a double-logistic curve calculated following Eq. (3):

$$\mu_t = \begin{cases} \frac{c}{1+\exp[b_A(t-M_A)]} + d & t > k \\ \frac{c}{1+\exp[b_S(t-M_S)]} + d & t \leq k \end{cases} \quad (3)$$

175 where b_A and b_S indicate the b parameters (rate of change) for the autumn green-down and the spring green-up, respectively; M_A and M_S are the autumn and spring midpoints, respectively; and k is the change-point day in the summer where the function switches from the green-down logistic curve to the green-up logistic curve, which we assumed to be the 182nd day of the year (1 July) in order to separate the year into two, which has been done elsewhere (e.g., Fu et al., 2016). Spring green-up was completed by the end of June in all of our sites; thus, the model fits were not heavily sensitive to this assumption. We assumed
180 that the minimum (d) and the maximum ($c + d$) phenological index values (GCC, NDVI, EVI) would not change during this one year for all sites and, thus, both parts of the change-point function fit the same c and d values. Years were fit independently for each site and, thus, we did not assume that the c and d values were the same between years. To be able to compare the influence of the different data sources on the uncertainty in the posterior, we used relatively uninformative Gaussian priors for the parameters (Table 2), which were created through simulating reasonable data. Since the motivation for this study was to
185 illustrate the ability of GOES to monitor phenological change by comparing it to other remotely-sensed data sources, we focused on only one transition date estimation method; though, additional methods are explored elsewhere (e.g., Klosterman et al., 2014)

190 Additionally, since the diurnal fit method (Wheeler and Dietze, 2019) produces estimates of uncertainty on daily NDVI, the means and precisions were incorporated in a normally-distributed errors-in-variables model within the phenology model. Likewise, errors-in-variables models were applied for the PhenoCam sites using the provided daily GCC standard deviation. Since the MOD13Q1 product does not provide a daily quantification of uncertainty (other than the quality flags that we separately applied), generic values were used based on the standard deviations given in Miura et al., (2000) of 0.01 and 0.02 for NDVI and EVI observations, respectively.

195 The phenology models were fit in JAGS (Plummer, 2003; version 4.3.0) using standard Markov Chain Monte Carlo (MCMC) approaches. JAGS was called from R (R Core Team, 2017; version 3.4.1) using the rjags (Plummer, 2018; version 4.7) and runjags (Denwood, 2016) packages. Five chains were run and all models converged as assessed using Gelman-Brooks-Rubin statistic (GBR < 1.05) and all had effective sample sizes > 5000 after burn-in was removed. From the posterior outputs, 95% CI's were calculated. To plot and compare the different data sources, which inherently have different ranges, the predicted phenological curves for all joint parameter posteriors were rescaled to have a range of 0 to 1.

200 As explained in Zhang et al. (2003), the start and end of season transition dates for the double logistic fits can be calculated as the roots of the third derivative of Eq. (3), which is illustrated in Fig. 2. The roots of the third derivative of our reparametrized function (Eq. 3) were calculated following Eqs. 4 and 5:

Deleted: ≤
Deleted: >

$$205 \quad \text{Root1} = \frac{b \times M + \log(\sqrt{3}+2)}{b}, \quad (4)$$

$$\text{Root2} = \frac{b \times M + \log(2-\sqrt{3})}{b}, \quad (5)$$

where *Root1* signifies the start of season and end of season transition dates for spring and autumn, respectively. *Root2* signifies the end of season and start of season for spring and autumn, respectively. 95% CI's were calculated from the 50% midpoint transition date posteriors and from the sample-specific *Root1* and *Root2* values.

210 2.4 Model comparison

Transition dates were compared between the different data sources with an emphasis on how the transition date estimates from the satellite-based data compared to PhenoCam. We calculated the coefficient of determination (R^2) and root mean square error (RMSE) by comparing the means of the transition dates from one data source with the means of the transition dates from another source for all possible combinations. It is important to note that these calculations are based on the deviation from the one-to-one line and not the line-of-best fit, which is often different from the predicted line. Since we are testing the similarity of the transition date estimates from the two sources, the predicted line is the one-to-one line. Additionally, bias of the transition dates was assessed by subtracting samples from the joint parameter posteriors of one source from those of another source. The medians of these differences (one median for each site) were then averaged across sites for each comparison. Width of 95% CI's were also compared between the different data sources for each transition date using paired *t*-tests.

220 3 Results

3.1 Overall fits

The selected phenological model fit well to the GOES daily data (Fig. 3a and b, Supplementary Figs. S3 and S4). Credible interval widths in the rescaled phenology fits were noticeably narrower for PhenoCam models, than GOES, and similar between the two MODIS products (Fig. 3a and b, Supplementary Figs. S5 and S6). Several sites (*e.g.*, Green Ridge 2018; Fig. 225 3e and g) had spring green-up periods that were shorter than the 16-day temporal resolution of the MODIS products, but were shown in the GOES measurements (Fig. 3e). Based on the output from the paired *t*-tests, PhenoCam transition date uncertainties were statistically narrower than those of all satellite data for all transition dates (*p*-value < 0.015 for all comparisons; Table S2). All GOES transition dates were statistically more certain (based on CI width) than the corresponding MODIS EVI estimates (*p*-value < 0.03), but were only significantly more certain than MODIS NDVI for middle and end of 230 spring (Fig. 4; Supplementary Table S2).

Deleted: ¶

3.2 Spring transition dates

235 GOES was correlated with PhenoCam for the start of spring transition, where MODIS possessed an early bias (Fig. 5; Table
3). GOES vs. PhenoCam (*i.e.*, PhenoCam median transition dates were the independent variable and GOES median transition
dates were the dependent variable) had the highest R^2 values (0.62 versus 0.00 and 0.00) and the lowest RMSE and average
bias (5.18 ± 0.03 earlier than PhenoCam; Table 3). Both MODIS NDVI and EVI, on the other hand, were biased earlier 10.18
 ± 0.1 and 11.66 ± 0.03 days on average, respectively. This bias was consistent amongst most sites (Fig. 6a). GOES was also
240 most correlated with PhenoCam for the middle of spring estimate, with the highest R^2 value (0.72), lowest RMSE (7.06 days)
and lowest average bias of 0.92 ± 0.03 days earlier than PhenoCam (Table 3). Most of the MODIS NDVI and EVI models
were biased early (Fig. 6b). Both MODIS data products were slightly more correlated with PhenoCam for the end of spring
than GOES. MODIS NDVI had the highest R^2 value of 0.80, but had a slightly earlier bias than MODIS EVI (1.75 ± 0.04 days
vs 0.79 ± 0.05 days). MODIS EVI and GOES had the same R^2 values of 0.71, but GOES had a later bias of 3.35 ± 0.03 days
(Table 3). There existed less correlation between GOES and the MODIS products (Supplementary Table S3).

245 3.3 Autumn transition dates

Autumn transition dates agreed less across all sources of data than in Spring. Except for MODIS EVI at the end of autumn (R^2
of 0.36), none of the satellite-based data sources were able to explain any variation in PhenoCam transition date estimates
(Table 3). Both NDVI sources (*i.e.*, GOES and MODIS) were consistently biased later (Table 3). MODIS EVI was biased
earlier than PhenoCam for the beginning and middle of autumn transition dates. Except for the middle of autumn where
250 MODIS NDVI and GOES had a slight correlation (R^2 of 0.34), satellite data sources were overall uncorrelated with each other
for all autumn transition dates (all had R^2 values = 0.00). The biases were the smallest between MODIS NDVI and GOES for
the first two transition dates (GOES was 5.25 ± 0.24 days earlier and 2.89 ± 0.05 days later than MODIS NDVI in the start
and middle of Autumn, respectively; Supplementary Table S3).

4 Discussion

255 4.1 Spring transition dates similarity

Our hypothesis that GOES spring transition dates would be more similar than MODIS to PhenoCam ones was supported by
our results for the start and middle of spring. While MODIS earlier sensing start of spring compared to PhenoCam has also
been attributed to a variety of reasons ranging from different viewing angles that sense the heterogeneity of phenological
change between different canopy layers (Ahl et al., 2006; Andrew D. Richardson and O'Keefe, 2009; Keenan et al., 2014b;
260 Ryu et al., 2014; Schwartz et al., 2002), spatial scaling affected by significant topography (Fisher et al., 2007), and snow melt
(Delbart et al., 2006), all these issues should affect the GOES-PhenoCam comparison as well. Based off of the similarity found
here between GOES and PhenoCam at marking the start and middle of spring, the mismatch between MODIS and PhenoCam

Deleted: prematurely

Deleted:

265 is likely largely due to the temporal resolution of the MODIS products. Hufkens et al. (2012) also points out that the temporal resolution of MODIS products cannot be expected to precisely track rapid leaf emergence in the spring due to the longer temporal resolution. By averaging over a relatively long time period (compared to the length of spring green-up), such techniques likely prematurely inflate NDVI and EVI values, giving the false impression that green-up occurred earlier. GOES, on the other hand, allows for daily NDVI estimates that are inherently more capable of tracking the initial spring increase.

270 Contrary to our hypothesis, though, we found that both MODIS indices were slightly less biased than GOES with PhenoCam at the end of spring, even with a higher or similar R^2 value to MODIS NDVI and EVI, respectively. Bias is likely more reliable as a measure of similarity than R^2 and RMSE because it includes the uncertainties in the transition dates for each site by utilizing the MCMC posterior samples instead of just comparing the median transition dates for each site. The later bias of GOES compared to PhenoCam and the MODIS products could potentially be due to an early bias in both PhenoCam and
275 the MODIS products. PhenoCam GCC has been found previously to reach its end of spring before many physiological traits such as total chlorophyll concentrations, leaf area and mass, leaf nitrogen and carbon concentrations, and leaf area index (Keenan et al., 2014b; Yang et al., 2014). Thus, even though GOES is biased slightly later than PhenoCam for the end of spring, it could be due to inherent differences between GCC and NDVI and should not affect GOES's ability to monitor interannual variability in the transition date as more years become available. MODIS could be biased early for the end of spring
280 in a similar way as that discussed in the previous paragraph related to its temporal resolution. Klosterman et al. (2014) found, however, that on average MODIS had a later end of spring estimate than PhenoCam. One primary component of their study was to investigate the impacts of land-cover heterogeneity on transition date comparisons between different sources and, thus, they included more heterogeneous landscapes in their dataset. They concluded that due to scaling issues, MODIS pixels that have smaller proportions of deciduous forests have MODIS end of spring estimates that are later than near-surface estimates.
285 Thus, it is reasonable to assume that since we attempted to not include any sites with substantial non-forest land cover types within the MODIS and GOES pixels, that our average bias should be lower. The effects of land-cover heterogeneity on the estimates of end of spring transition should be kept in mind when using GOES to monitor more heterogeneous sites.

290 As with many studies, the results and conclusions of this study could be dependent on the methods used. It is possible that a different transition date estimation method (e.g., the one proposed by Klosterman et al., 2014, that also accounted for summer green-down, which we were not focused on) would result in different conclusions. If there is bias in the methods here to estimate transition dates, it is shared across data sources, sites, and years. Similarly, using a different MODIS product might also result in different conclusions. While a daily MODIS product does exist, it is also created using multi-day periods of measurements (Ju et al., 2010) and, thus, is also possibly subject to similar constraints. We encourage others to consider more complex models and other phenology products, but the primary aim of this study is to demonstrate the value of GOES for studying phenology with an initial comparison to PhenoCams and MODIS.

Formatted: Font: Italic

4.2 Spring and autumn compared

As we hypothesized, spring transition dates were more similar across data sources than autumn ones. This mismatch between PhenoCam GCC autumn transition dates and NDVI and EVI (low R^2 values and high biases) has been found in numerous other studies (e.g., Hufkens et al., 2012; Keenan et al., 2014a; Klosterman et al., 2014; Richardson et al., 2018b; Zhang et al., 2018). This is most likely due to physiological differences between the different metrics (i.e., GCC, NDVI, and EVI) that become more apparent in the autumn with changes in color and canopy structure often occurring separately. While all three metrics measure some combination of greenness and canopy structure, only GCC directly considers green reflectance in its calculation and NDVI and EVI have been found to be more impacted by leaf presence and canopy structure (Kobayashi et al., 2007; Pettorelli et al., 2005). The higher uncertainties in the autumn transition dates, compared to spring ones, across all data sources were expected given the longer season length and the higher heterogeneity in autumn compared to spring. For example, triggers of autumn phenology are less understood and consistent than spring (Piao et al., 2019) and the timing of autumn phenological events differs greater between species than in spring (Richardson et al., 2006). Like the other data sources, GOES spring transition date estimates were most certain and most similar to those derived from other data sources.

Deleted: GCC is by definition a measure of greenness, but NDVI and EVI are also

4.3 Autumn transition dates similarity

We hypothesized that in the autumn, the transition dates derived from GOES NDVI data would be most similar to those from MODIS NDVI data, which was mostly true. The low biases that existed between the two at the start and middle of autumn (MODIS NDVI was 5.25 ± 0.24 days later and 2.89 ± 0.05 earlier than GOES, respectively; Supplementary Table S3) are promising while the high end of autumn bias (MODIS NDVI was 11.03 ± 0.24 days earlier than GOES) could potentially be due to the high amount of noise in the GOES data that remains in the winter in many sites (Supplementary Figs. S2 and S3). Future directions for GOES that should help decrease these biases include developing a snow cover mask (which is a planned GOES product), developing a more sophisticated atmospheric correction algorithm for GOES reflectance data, and developing methodology for correcting for seasonal variations in solar angle. GOES will inherently be better at establishing a winter baseline at sites with less snowy days than sites that consistently have a layer of snow obstructing accurate satellite measurements. This could also be improved by developing multi-year phenology models that assume the winter NDVI baseline is the same (or changes slowly) between years and, thus, are improved by a higher number of winter observations. Furthermore, more informative priors in the diurnal fit model for estimating daily NDVI values that change seasonally would also improve the ability to estimate winter NDVI values with more certainty. These will likely help improve the correlation between GOES NDVI and MODIS NDVI.

4.4 Uncertainty in transition date estimates

We hypothesized that the increased temporal frequency in GOES data would produce more certain estimates of transition dates than MODIS. In practice, was found differences between MODIS indices, with GOES transition dates being significantly more

certain than MODIS EVI for all transition dates, but only significantly more certain than MODIS NDVI for the start and middle
330 of spring. However, as previously discussed, there are nontrivial differences between *what* NDVI and GCC are measuring in
the autumn and end of spring that transcend simple issues of data quality and quantity, which suggests GOES is providing
important new information about vegetation phenology. Once future work further improves the GOES products, reducing
noise due to factors such as snow and atmospheric attenuation, the widths of the CI's are expected to improve. Additionally,
the lack of a spatially- and temporally-varying MODIS uncertainty product, as we have produced for GOES, provided a
335 limitation to this comparison and it is possible that specific daily MODIS uncertainties, congruent to that we used from the
GOES data, would affect this conclusion. In particular, many of the MODIS validation efforts have focused on within-season
comparisons (*e.g.*, Miura et al., 2000) not periods of phenological transition and, thus, MODIS uncertainties are likely
underestimated. Furthermore, the differences in results between comparing GOES transition dates' CI widths to those from
MODIS NDVI and EVI is likely partially due to the differences in observational error applied. Based on Miura et al. (2000),
340 a smaller observational error was applied to MODIS NDVI than MODIS EVI, which likely was enough to allow for all GOES
transition date estimates to be significantly more certain than the respective MODIS EVI estimates, but not always MODIS
NDVI. This emphasizes the importance of providing uncertainty estimates with remotely sensed phenology data, which fitting
diurnal curves to GOES data allows (Wheeler and Dietze, 2019).

4.5 Future phenological applications of GOES NDVI data

345 With its full coverage and high temporal resolution, GOES-16 and GOES-17 have the potential to revolutionize the study of
leaf phenology and allow for a variety of studies that previously would not have been possible at the extent they are now. First,
many studies have found that climate change is altering phenology on the scale of days per decade (Cleland et al., 2007;
Keenan et al., 2014a; Parmesan and Yohe, 2003; Root et al., 2003). Because of the long temporal scale of the studied MODIS
NDVI and EVI products, their ability to precisely and accurately monitor both these trends and interannual variability is
350 inherently limited. While a daily MODIS NDVI product is becoming more readily available (and MODIS measurements are
taken sub-daily, but at varying viewing angles), it still remains more inaccessible than many of the lower-frequency MODIS
data products because it is relatively new. It is important to have additional remotely-sensed data sources, especially ones that
are not affected by changing viewing angles.

Second, GOES is able to provide real-time data of spring green-up even for those springs that occur quicker than the
355 16-day resolution of this MODIS product (*e.g.*, Green Ridge 2018 in Fig. 3). These sites possessed no 16-day MODIS NDVI
nor EVI measurements during the green-up period. This limitation would become even more severe when monitoring of green-
up in real-time, as the transition would only be detected after the fact. [The seasonal curve fitting methodology used here to
estimate transition dates would not be suitable in real-time, but alternative methods, such as determining that spring has started
if the phenological index is 10% greater than the winter baseline or iteratively assimilating data into a process model](#) (Viskari
360 et al., 2015) [could be used to provide insight into whether or not certain transition dates have occurred in real-time.](#) With the
data that GOES is able to supply, it becomes more possible to monitor and forecast the start and progression of green-up at

large scales using near-real-time data, instead of having to wait for the next reliable MODIS product value, which might be 15 days away.

Deleted:

A third beneficial future application of the high temporal GOES NDVI data is the ability to monitor the effects of storms (*e.g.*, hurricanes), droughts, and frosts on phenology and NDVI. It is possible that the effects of some of these disturbances are only present for less than the temporal resolution of MODIS. For example, Richardson et al. (2018b) found that the effect of a spring frost event was visible within PhenoCam data, but not clearly visible within the 16-day MODIS data. By providing higher temporal spring NDVI data, it is more likely that the effects of similar frost events could be observed at more areas that do not have a PhenoCam present.

A fourth benefit is that by combining data from multiple high-temporal sources (*i.e.*, PhenoCam and GOES), we now may be able to ask questions related to differentiating the physiological impacts of phenological change in different indices and at different spatial scales. For example, with high-temporal NDVI data, we now can start asking questions about what specific phenological processes control the rate of spring increase (*i.e.*, budburst vs. leaf expansion) and how these are affected by spatial scale. Similarly, combining PhenoCam and GOES data has the potential to help us better disentangle different autumn phenological processes (*i.e.*, leaf color change vs. leaf fall).

We are not suggesting that GOES should replace other phenology data sources, but that a combination of different data sources, which each have their own strengths and weaknesses, is beneficial. MODIS has a much longer record of data than GOES and still remains an important source of information. Additionally, it has a different spatial scale that is between PhenoCam and GOES and a combination of the three could help answer spatial questions related to NDVI (*e.g.*, how does NDVI scale between canopy level to landscape level and how does this change seasonally?).

In conclusion, we have shown that GOES-16 and -17 possess great potential at enhancing the monitoring of leaf phenology, which will allow us to ask and answer new questions and improve our knowledge of this complicated, but important aspect of ecology and environmental science.

Code availability

All code will be available soon on Github at https://github.com/k-wheeler/NEFI_pheno and https://github.com/k-wheeler/NEFI_pheno/tree/master/PhenologyBayesModeling.

Data availability

All data was publicly available and can be accessed as described in the methods section.

390 **Author contribution**

KW and MD designed the study and KW executed the modeling and analysis. KW wrote the manuscript with inputs and suggestions from MD throughout the writing process.

Competing interests

The authors declare that they have no conflict of interest.

395 **Acknowledgements**

This work was made possible by the U.S. National Science Foundation grant 1638577. KIW also acknowledges support under the U.S. National Science Foundation Graduate Research Fellowship grant number 1247312. Special thanks to the Dietze lab members for feedback on the manuscript. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. We thank our many collaborators, including site PIs and technicians, for their efforts in support of PhenoCam. The development of PhenoCam has been funded by the Northeastern States Research Cooperative, NSF's Macrosystems Biology program (awards EF-1065029 and EF-1702697), and DOE's Regional and Global Climate Modeling program (award DE-SC0016011). We acknowledge additional support from the US National Park Service Inventory and Monitoring Program and the USA National Phenology Network (grant number G10AP00129 from the United States Geological Survey), and from the USA National Phenology Network and North Central Climate Science Center (cooperative agreement number G16AC00224 from the United States Geological Survey). We also thank the USDA Forest Service Air Resource Management program and the National Park Service Air Resources program for contributing their camera imagery to the PhenoCam archive. Research at the Bartlett Experimental Forest tower is supported by the National Science Foundation (grant DEB-1114804) and the USDA Forest Service's Northern Research Station. Research at the Coweeta flux tower is funded through the USDA Forest Service, Southern Research Station; USDA Agriculture and Food Research Initiative Foundational Program, award number 2012-67019-19484; EPA agreement number 13-IA-11330140-044; and the National Science Foundation, Long-Term Ecological Research (LTER) program, award #DEB-0823293. Research at Harvard Forest is partially supported through the National Science Foundation's LTER program (DEB-1237491), and Dept. of Energy Office of Science (BER). The Hubbard Brook Ecosystem Study is a collaborative effort at the Hubbard Brook Experimental Forest, which is operated and maintained by the USDA Forest Service, Northern Research Station, Newtown Square, PA. Research at the MOFLUX site is supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research Program, Climate and Environmental Sciences Division. ORNL is managed by UT-Battelle, LLC, for the U.S. Department of Energy under contract DE-AC05-00OR22725. U.S. Department of Energy support for the University of Missouri (Grant DE-FG02-03ER63683) is gratefully acknowledged. Research at the Morgan-Monroe Ameriflux site is supported by the US Department of Energy, Office of Science, Office of

420 [Biological and Environmental Research through the Ameriflux Management Project administered by Lawrence Berkeley National Lab. Funding for the Shenandoah PhenoCam and related research has been provided by the U.S. Geological Survey Land Change Science Program \(Shenandoah National Park Phenology Project\) with logistical support from the National Park Service in collaboration with the University of Virginia Department of Environmental Sciences. Camera images from the Shining Rock Wilderness are provided courtesy of the USDA Forest Service Air Resources Management Program. Primary support for the University of Michigan AmeriFlux Core Site \(US-UMd\) provided by the Department of Energy Office of Science. Infrastructure support provided by the University of Michigan Biological Station. Research at the Willow Creek Ameriflux core site is provided by the Dept. Of Energy Office of Science to the ChEAS Cluster.](#)

Formatted: English (US)

References


- 430 Ahl, D. E., Gower, S. T., Burrows, S. N., Shabanov, N. V., Myneni, R. B. and Knyazikhin, Y.: Monitoring spring canopy phenology of a deciduous broadleaf forest using MODIS, *Remote Sensing of Environment*, 104(1), 88–95, doi:10.1016/j.rse.2006.05.003, 2006.
- Alekseychik, P. K., Korrensalo, A., Mammarella, I., Vesala, T. and Tuittila, E.-S.: Relationship between aerodynamic roughness length and bulk sedge leaf area index in a mixed-species boreal mire complex, *Geophysical Research Letters*, 44(11), 5836–5843, doi:10.1002/2017GL073884, 2017.
- 435 Andrew D. Richardson and O’Keefe, J.: Phenological differences between understory and overstory, in *Phenology of Ecosystem Processes*, pp. 87–117, Springer, New York, NY., 2009.
- Cleland, E., Chuine, I., Menzel, A., Mooney, H. and Schwartz, M.: Shifting plant phenology in response to global change, *Trends in Ecology & Evolution*, 22(7), 357–365, doi:10.1016/j.tree.2007.04.003, 2007.
- 440 Delbart, N., Le Toan, T., Kergoat, L. and Fedotova, V.: Remote sensing of spring phenology in boreal regions: A free of snow-effect method using NOAA-AVHRR and SPOT-VGT data (1982–2004), *Remote Sensing of Environment*, 101(1), 52–62, doi:10.1016/j.rse.2005.11.012, 2006.
- Denwood, M. J.: runjags: An R Package Providing Interface Utilities, Model Templates, Parallel Computing Methods and Additional Distributions for MCMC Models in JAGS, *Journal of Statistical Software*, 71(9), 1–25, doi:10.18637/jss.v071.i09, 2016.
- 445 Didan: MOD13Q1 MODIS/Terra Vegetation Indices 16-Day L3 Global 250m SIN Grid V006 [NDVI and pixel_reliability], NASA EOSDIS LP DAAC., 2015.
- Filippa, G., Cremonese, E., Migliavacca, M., Galvagno, M., Sonnentag, O., Humphreys, E., Hufkens, K., Ryu, Y., Verfaillie, J., Morra di Cella, U. and Richardson, A. D.: NDVI derived from near-infrared-enabled digital cameras: Applicability across different plant functional types, *Agricultural and Forest Meteorology*, 249, 275–285, doi:10.1016/j.agrformet.2017.11.003, 450 2018.
- Fisher, J. I., Richardson, A. D. and Mustard, J. F.: Phenology model from surface meteorology does not capture satellite-based greenup estimations, *Global Change Biology*, 13(3), 707–721, doi:10.1111/j.1365-2486.2006.01311.x, 2007.

- Fu, Y., Zheng, Z., Shi, H. and Xiao, R.: A Novel Large-Scale Temperature Dominated Model for Predicting the End of the Growing Season, edited by W. Yuan, PLoS ONE, 11(11), e0167302, doi:10.1371/journal.pone.0167302, 2016.
- 455 Gao, M., Piao, S., Chen, A., Yang, H., Liu, Q., Fu, Y. H. and Janssens, I. A.: Divergent changes in the elevational gradient of vegetation activities over the last 30 years, *Nat Commun*, 10(1), 2970, doi:10.1038/s41467-019-11035-w, 2019.
- Gill, A. L., Gallinat, A. S., Sanders-DeMott, R., Rigden, A. J., Gianotti, D. J. S., Mantooh, J. A. and Templer, P. H.: Changes in autumn senescence in northern hemisphere deciduous trees: a meta-analysis of autumn phenology studies, *Annals of Botany*, 116(6), 875–888, doi:10.1093/aob/mcv055, 2015.
- 460 GOES-R Algorithm Working Group and GOES-R Series Program: NOAA GOES-R Series Advanced Baseline Imager (ABI) Level 2 Clear Sky Mask, NOAA National Centers for Environmental Information, doi:doi:10.7289/V5SF2TGP, 2018.
- GOES-R Calibration Working Group and GOES-R Series Program: NOAA GOES-R Series Advanced Baseline Imager (ABI) Level 1b Radiances. [Channels 2 and 3], NOAA National Centers for Environmental Information, doi:doi:10.7289/V5BV7DSR, 2017.
- 465 Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *International Journal of Climatology*, 25(15), 1965–1978, doi:10.1002/joc.1276, 2005.
- Hmimina, G., Dufrêne, E., Pontailleur, J.-Y., Delpierre, N., Aubinet, M., Caquet, B., de Grandcourt, A., Burban, B., Flechard, C., Granier, A., Gross, P., Heinesch, B., Longdoz, B., Moureaux, C., Ourcival, J.-M., Rambal, S., Saint André, L. and Soudani, K.: Evaluation of the potential of MODIS satellite data to predict vegetation phenology in different biomes: An investigation using ground-based NDVI measurements, *Remote Sensing of Environment*, 132, 145–158, doi:10.1016/j.rse.2013.01.010, 2013.
- 470 Hudson, J. E., Levia, D. F., Wheeler, K. I., Winters, C. G., Vaughan, M. C. H., Chace, J. F. and Sleeper, R.: American beech leaf-litter leachate chemistry: Effects of geography and phenophase, *Journal of Plant Nutrition and Soil Science*, 181(2), 287–295, doi:10.1002/jpln.201700074, 2018.
- 475 Hufkens, K., Friedl, M., Sonnentag, O., Braswell, B. H., Milliman, T. and Richardson, A. D.: Linking near-surface and satellite remote sensing measurements of deciduous broadleaf forest phenology, *Remote Sensing of Environment*, 117, 307–321, doi:10.1016/j.rse.2011.10.006, 2012.
- Jin, S., Homer, C., Yang, L., Danielson, P., Dewitz, J., Li, C., Zhu, Z., Xian, G. and Howard, D.: Overall Methodology Design for the United States National Land Cover Database 2016 Products, *Remote Sensing*, 11(24), 2971, doi:10.3390/rs11242971, 2019.
- 480 Ju, J., Roy, D. P., Shuai, Y. and Schaaf, C.: Development of an approach for generation of temporally complete daily nadir MODIS reflectance time series, *Remote Sensing of Environment*, 114(1), 1–20, doi:10.1016/j.rse.2009.05.022, 2010.
- Keenan, T. F., Gray, J., Friedl, M. A., Toomey, M., Bohrer, G., Hollinger, D. Y., Munger, J. W., O’Keefe, J., Schmid, H. P., Wing, I. S., Yang, B. and Richardson, A. D.: Net carbon uptake has increased through warming-induced changes in temperate forest phenology, *Nature Climate Change*, 4(7), 598–604, doi:10.1038/nclimate2253, 2014a.
- 485 Keenan, T. F., Darby, B., Felts, E., Sonnentag, O., Friedl, M. A., Hufkens, K., O’Keefe, J., Klosterman, S., Munger, J. W., Toomey, M. and Richardson, A. D.: Tracking forest phenology and seasonal physiology using digital repeat photography: a critical assessment, *Ecological Applications*, 24(6), 1478–1489, doi:10.1890/13-0652.1, 2014b.
- Killingbeck, K. T.: Nutrient Resorption, in *Plant Cell Death Processes*, Elsevier, London., 2004.


- 490 Klosterman, S. T., Hufkens, K., Gray, J. M., Melaas, E., Sonnentag, O., Lavine, I., Mitchell, L., Norman, R., Friedl, M. A. and Richardson, A. D.: Evaluating remote sensing of deciduous forest phenology at multiple spatial scales using PhenoCam imagery, *Biogeosciences*, 11(16), 4305–4320, doi:10.5194/bg-11-4305-2014, 2014.
- Kobayashi, H., Suzuki, R. and Kobayashi, S.: Reflectance seasonality and its relation to the canopy leaf area index in an eastern Siberian larch forest: Multi-satellite data and radiative transfer analyses, *Remote Sensing of Environment*, 106(2), 238–252, doi:10.1016/j.rse.2006.08.011, 2007.
- 495 Liu, Q., Fu, Y. H., Zhu, Z., Liu, Y., Liu, Z., Huang, M., Janssens, I. A. and Piao, S.: Delayed autumn phenology in the Northern Hemisphere is related to change in both climate and spring phenology, *Global Change Biology*, 22(11), 3702–3711, doi:10.1111/gcb.13311, 2016.
- Liu, Y., Hill, M. J., Zhang, X., Wang, Z., Richardson, A. D., Hufkens, K., Filippa, G., Baldocchi, D. D., Ma, S., Verfaillie, J. and Schaaf, C. B.: Using data from Landsat, MODIS, VIIRS and PhenoCams to monitor the phenology of California oak/grass savanna and open grassland across spatial scales, *Agricultural and Forest Meteorology*, 237–238, 311–325, doi:10.1016/j.agrformet.2017.02.026, 2017.
- 500 McKown, A. D., Guy, R. D., Azam, M. S., Drewes, E. C. and Quamme, L. K.: Seasonality and phenology alter functional leaf traits, *Oecologia*, 172(3), 653–665, doi:10.1007/s00442-012-2531-5, 2013.
- 505 Miura, T., Huete, A. R. and Yoshioka, H.: Evaluation of sensor calibration uncertainties on vegetation indices for MODIS, *IEEE Transactions on Geoscience and Remote Sensing*, 38(3), 1399–1409, doi:10.1109/36.843034, 2000.
- Parmesan, C. and Yohe, G.: A globally coherent fingerprint of climate change impacts across natural systems, *Nature*, 421(6918), 37–42, doi:10.1038/nature01286, 2003.
- Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J.-M., Tucker, C. J. and Stenseth, N. Chr.: Using the satellite-derived NDVI to assess ecological responses to environmental change, *Trends in Ecology & Evolution*, 20(9), 503–510, doi:10.1016/j.tree.2005.05.011, 2005.
- 510 PhenoCam: Provisional data for site harvard, ROI DB_0001; hubbardbrooksfws, ROI: DB_3000; umichbiological, ROI: DB_2000; coweeta, ROI: DB_0001; bartlettir, ROI: DB_1000; missouriozarks, ROI: DB_0001; morganmonroe, ROI: DB_1000; russellsage, ROI: DB_0001; willowcreek, ROI: DB_1000; bullshoals, ROI: DB_1000; dukehw, ROI: DB_1000, greenridge1, ROI: DB_1000; shenandoah, ROI: DB_0001; marcel, ROI: DB_1000; shiningrock, DB_0003., Downloaded from <http://phenocam.sr.unh.edu/> on 30 June 2020., 2018.
- 515 Piao, S., Liu, Q., Chen, A., Janssens, I. A., Fu, Y., Dai, J., Liu, L., Lian, X., Shen, M. and Zhu, X.: Plant phenology and global climate change: Current progresses and challenges, *Global Change Biology*, 25(6), 1922–1940, doi:10.1111/gcb.14619, 2019.
- 520 Plummer, M.: rjags: Bayesian Graphical Models using MCMC. [online] Available from: <https://CRAN.R-project.org/package=rjags>, 2018.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: <https://www.R-project.org/>, 2017.
- Richardson, A. D., Bailey, A. S., Denny, E. G., Martin, C. W. and O'keefe, J.: Phenology of a northern hardwood forest canopy, *Global Change Biology*, 12(7), 1174–1188, doi:10.1111/j.1365-2486.2006.01164.x, 2006.

- 525 Richardson, A. D., Jenkins, J. P., Braswell, B. H., Hollinger, D. Y., Ollinger, S. V. and Smith, M.-L.: Use of digital webcam images to track spring green-up in a deciduous broadleaf forest, *Oecologia*, 152(2), 323–334, doi:10.1007/s00442-006-0657-z, 2007.
- Richardson, A. D., Anderson, R. S., Arain, M. A., Barr, A. G., Bohrer, G., Chen, G., Chen, J. M., Ciais, P., Davis, K. J., Desai, A. R., Dietze, M. C., Dragoni, D., Garrity, S. R., Gough, C. M., Grant, R., Hollinger, D. Y., Margolis, H. A., McCaughey, H.,
530 Migliavacca, M., Monson, R. K., Munger, J. W., Poulter, B., Raczka, B. M., Ricciuto, D. M., Sahoo, A. K., Schaefer, K., Tian, H., Vargas, R., Verbeeck, H., Xiao, J. and Xue, Y.: Terrestrial biosphere models need better representation of vegetation phenology: results from the North American Carbon Program Site Synthesis, *Global Change Biology*, 18(2), 566–584, doi:10.1111/j.1365-2486.2011.02562.x, 2012.
- Richardson, A. D., Keenan, T. F., Migliavacca, M., Ryu, Y., Sonnentag, O. and Toomey, M.: Climate change, phenology, and
535 phenological control of vegetation feedbacks to the climate system, *Agricultural and Forest Meteorology*, 169, 156–173, doi:10.1016/j.agrformet.2012.09.012, 2013.
- Richardson, A. D., Hufkens, K., Milliman, T. and Frolking, S.: Intercomparison of phenological transition dates derived from the PhenoCam Dataset V1.0 and MODIS satellite remote sensing, *Scientific Reports*, 8(1), 5679, doi:10.1038/s41598-018-23804-6, 2018a.
- 540 Richardson, A. D., Hufkens, K., Milliman, T., Aubrecht, D. M., Chen, M., Gray, J. M., Johnston, M. R., Keenan, T. F., Klosterman, S. T., Kosmala, M., Melaas, E. K., Friedl, M. A. and Frolking, S.: Tracking vegetation phenology across diverse North American biomes using PhenoCam imagery, *Scientific Data*, 5, 180028, doi:10.1038/sdata.2018.28, 2018b.
- Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C. and Pounds, J. A.: Fingerprints of global warming on wild animals and plants, *Nature*, 421(6918), 57–60, doi:10.1038/nature01333, 2003.
- 545 Ryu, Y., Lee, G., Jeon, S., Song, Y. and Kimm, H.: Monitoring multi-layer canopy spring phenology of temperate deciduous and evergreen forests using low-cost spectral sensors, *Remote Sensing of Environment*, 149, 227–238, doi:10.1016/j.rse.2014.04.015, 2014.
- Schmit, T. J., Griffith, P., Gunshor, M. M., Daniels, J. M., Goodman, S. J. and Lehair, W. J.: A Closer Look at the ABI on the GOES-R Series, *Bull. Amer. Meteor. Soc.*, 98(4), 681–698, doi:10.1175/BAMS-D-15-00230.1, 2016.
- 550 Schwartz, M. D., Reed, B. C. and White, M. A.: Assessing satellite-derived start-of-season measures in the conterminous USA, *International Journal of Climatology*, 22(14), 1793–1805, doi:10.1002/joc.819, 2002.
- Seyednasrollah, B., Young, A. M., Hufkens, K., Milliman, T., Friedl, M. A., Frolking, S. and Richardson, A. D.: Tracking vegetation phenology across diverse biomes using Version 2.0 of the PhenoCam Dataset, *Scientific Data*, 6(1), 222, doi:10.1038/s41597-019-0229-9, 2019.
- 555 Thieurmel, B. and Elmarhraoui, A.: *suncalc*: Compute Sun Position, Sunlight Phases, Moon Position and Lunar Phase. [online] Available from: <https://CRAN.R-project.org/package=suncalc>, 2019.
- Tuck, S. L., Phillips, H. R. P., Hintzen, R. E., Scharlemann, J. P. W., Purvis, A. and Hudson, L. N.: MODISTools – downloading and processing MODIS remotely sensed data in R, *Ecology and Evolution*, 4(24), 4658–4668, doi:10.1002/ece3.1273, 2014.
- 560 Viskari, T., Hardiman, B., Desai, A. R. and Dietze, M. C.: Model-data assimilation of multiple phenological observations to constrain and predict leaf area index, *Ecological Applications*, 25(2), 546–558, doi:10.1890/14-0497.1, 2015.


- Wheeler, K. I. and Dietze, M. C.: A Statistical Model for Estimating Midday NDVI from the Geostationary Operational Environmental Satellite (GOES) 16 and 17, *Remote Sensing*, 11(21), 2507, doi:10.3390/rs11212507, 2019.
- 565 Xue, Y., Fennessy, M. J. and Sellers, P. J.: Impact of vegetation properties on U.S. summer weather prediction, *Journal of Geophysical Research: Atmospheres*, 101(D3), 7419–7430, doi:10.1029/95JD02169, 1996.
- Yang, L., Jin, S., Danielson, P., Homer, C., Gass, L., Bender, S. M., Case, A., Costello, C., Dewitz, J., Fry, J., Funk, M., Granneman, B., Liknes, G. C., Rigge, M. and Xian, G.: A new generation of the United States National Land Cover Database: Requirements, research priorities, design, and implementation strategies, *ISPRS Journal of Photogrammetry and Remote Sensing*, 146, 108–123, doi:10.1016/j.isprsjprs.2018.09.006, 2018.
- 570 Yang, X., Tang, J. and Mustard, J. F.: Beyond leaf color: Comparing camera-based phenological metrics with leaf biochemical, biophysical, and spectral properties throughout the growing season of a temperate deciduous forest, *Journal of Geophysical Research: Biogeosciences*, 119(3), 181–191, doi:10.1002/2013JG002460, 2014.
- 575 Zhang, X., Friedl, M. A., Schaaf, C. B., Strahler, A. H., Hodges, J. C. F., Gao, F., Reed, B. C. and Huete, A.: Monitoring vegetation phenology using MODIS, *Remote Sensing of Environment*, 84(3), 471–475, doi:10.1016/S0034-4257(02)00135-9, 2003.
- Zhang, X., Jayavelu, S., Liu, L., Friedl, M. A., Henebry, G. M., Liu, Y., Schaaf, C. B., Richardson, A. D. and Gray, J.: Evaluation of land surface phenology from VIIRS data using time series of PhenoCam imagery, *Agricultural and Forest Meteorology*, 256–257, 137–149, doi:10.1016/j.agrformet.2018.03.003, 2018.
- 580 Zheng, Z. and Zhu, W.: Uncertainty of Remote Sensing Data in Monitoring Vegetation Phenology: A Comparison of MODIS C5 and C6 Vegetation Index Products on the Tibetan Plateau, *Remote Sensing*, 9, 1288, doi:10.3390/rs9121288, 2017.
- Zhu, J. and Zeng, X.: Influences of the seasonal growth of vegetation on surface energy budgets over middle to high latitudes, *International Journal of Climatology*, 37(12), 4251–4260, doi:10.1002/joc.5068, 2017.




585 Figure 1: Map of 2016 National Land Cover Database (NLCD) classification (Jin et al., 2019; Yang et al., 2018) for the study region showing the locations of the selected sites. Selected sites are located throughout the deciduous forested area in the United States. Specific locations are given in Table 1.




590 Figure 2: Schematic based off of Fig. 2 in Zhang et al. (2003) describing the selection of transition dates (shown using the vertical lines). (a) illustrates an example phenological curve of green-down and green-up. (b) illustrates the first derivative. (c) illustrates the second derivative where the root (second derivative equals 0) gives the value of the middle of season date. (d) illustrates the third derivative where the roots give the start and end of both seasons.



595 **Figure 3:** Timeseries of data from GOES NDVI daily data (a and b), PhenoCam GCC (c and d), MODIS NDVI (e and f), MODIS EVI (g and h) and the rescaled credible intervals (i and j) for Green Ridge 2018 (left column) and Harvard Forest 2019 (right column). (a–h) include mean observations in black dots with 95% confidence intervals shown with vertical gray lines. 95% credible intervals (CI) are given with the different shading specific to each data source. The Green Ridge 2018 spring occurred quicker than the MODIS temporal resolution, but was captured by the daily resolution of the GOES data.




600 Figure 4: The 95% credible interval (CI) widths for the different data sources for spring start (a), middle (b), and end (c); and autumn start (d), middle (e), and end (f). Colors denote the different data sources, which are labeled on the x-axis. PhenoCam had the most certain transition date estimates and GOES was always more certain than MODIS EVI, but only more certain than MODIS NDVI for the middle and end of spring.



605 Figure 5: Scatter plots showing how the different data sources compare for their estimation of spring start (a), middle (b), and end (c); and autumn start (d), middle (e), and end (f). Median transition dates are indicated by the point and 95% credible intervals are indicated by the lines. The x-axis gives the day of year of the PhenoCam (PC) transition and the y-axis indicates the day of year of the different satellite data sources, which are color-coded as indicated in the legend. Spring correlations are much higher than autumn ones and GOES dates are more correlated at the start and middle of spring (a and b), but are slightly biased late at the end of spring (c).

610



615 Figure 6: The different biases for the different satellite-based data sources compared to PhenoCam for their estimation of spring start (a), middle (b), and end (c); and autumn start (d), middle (e), and end (f). A negative bias indicates the given data source was earlier than PhenoCam. The red line denotes zero bias. Boxes are color coded by the data source as indicated on the x-axis. The biases are larger in the autumn than in the spring. There are some with little median bias.

Table 1: Characteristics of selected sites including the coordinates and some climate data from WorldClim (Hijmans et al., 2005), which was assessed from the PhenoCam website (<https://phenocam.sr.unh.edu/webcam/>).

Site name	Latitude	Longitude	Mean Annual temperature (°C)	Mean Annual precipitation (mm)
-----------	----------	-----------	------------------------------	--------------------------------

Marcell	47.514	-93.469	2.9	687.0
Willow Creek	45.806	-90.079	3.9	820.0
University of Michigan	45.560	-84.714	5.9	797.0
Biological Station (UMBS)				
Bartlett	44.065	-71.288	5.5	1224.0
Hubbard Brook	43.927	-71.741	4.6	1190.0
Harvard Forest	42.538	-72.172	6.8	1139.0
Green Ridge	39.691	-78.407	10.5	935.0
Morgan Monroe	39.323	-86.413	11.2	1087.0
Missouri Ozarks	38.744	-92.200	12.4	974.0
Shenandoah	38.617	-78.350	8.4	1222.0
Bull Shoals	36.563	-93.067	13.9	1084.0
Duke	35.974	-79.100	14.6	1166.0
Shining Rock	35.390	-82.775	9.3	1835.0
Coweeta	35.060	-83.428	12.5	1722.0
Russell Sage*	32.457	-91.974	18.1	1341.0

*Note: Due to the cessation of PhenoCam data collection in 2019, Russell Sage was only included in the 2018 analysis.

620

Table 2: Parameter priors. “Satellites” refers to all GOES, MODIS NDVI and MODIS EVI. The priors on when the middle of spring and autumn occurred was set as the day of year (DOY). Normal distribution is abbreviated with N.

Parameter Name	Parameter Abbreviation	Data Source	Distribution (Mean, Standard Deviation)
Middle of spring (DOY)	M_S	All	N (110, 40)
Middle of autumn (DOY)	M_A	All	N (300, 40)
Spring rate of change	b_S	All	N (-0.10, 0.05)
Autumn rate of change	b_A	All	N (0.10, 0.05)
Minimum of phenological curve	d	Satellites	N (0.6, 0.2)
Minimum of phenological curve	d	PhenoCam	N (0.35, 0.15)
Range of phenological data	c	Satellites	N (0.4, 0.2)
Range of phenological data	c	PhenoCam	N (0.3, 0.15)

625 **Table 3. Summary statistics for comparisons with PhenoCam transition dates. *Negative indicates the data source is earlier than PhenoCam. The widths of the 95% credible intervals (CI) of the biases are given.**

Data Source	R²	RMSE (days)	Average Bias* (days; 95% CI)
<i>Start of Spring:</i>			
GOES	0.62	9.06	-5.18 ± 0.03
MODIS NDVI	0.00	13.53	-10.18 ± 0.1
MODIS EVI	0.00	13.4	-11.66 ± 0.03
<i>Middle of Spring:</i>			
GOES	0.77	7.06	-0.92 ± 0.03
MODIS NDVI	0.00	10.86	-5.56 ± 0.04
MODIS EVI	0.5	9.42	-6.61 ± 0.03
<i>End of Spring:</i>			
GOES	0.71	7.99	3.35 ± 0.03
MODIS NDVI	0.12	10.49	-0.95 ± 0.1
MODIS EVI	0.71	7.7	-1.57 ± 0.03
630 <i>Start of Autumn:</i>			
GOES	0.00	32.3	12.59 ± 0.11
MODIS NDVI	0.00	34.2	17.84 ± 0.26
MODIS EVI	0.00	26.67	-7.5 ± 0.11
<i>Middle of Autumn:</i>			
GOES	0.00	25.61	17.04 ± 0.07
MODIS NDVI	0.00	23.43	14.16 ± 0.08
MODIS EVI	0.00	15.64	-2.7 ± 0.07
<i>End of Autumn:</i>			
GOES	0.00	26.58	21.5 ± 0.07
MODIS NDVI	0.00	16.17	10.47 ± 0.24
MODIS EVI	0.36	11.23	2.11 ± 0.06