

1 **Sub-alpine grassland productivity increased with warmer and**
2 **drier conditions, but not with higher N-deposition, in an**
3 **altitudinal transplantation experiment**

4
5
6 Matthias Volk¹, Matthias Suter², Anne-Lena Wahl¹, Seraina Bassin^{1,3}

7 ¹Climate and Agriculture, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

8 ²Forage Production and Grassland Systems, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

9 ³Pädagogische Hochschule Schaffhausen, Ebnetstrasse 80, 8200 Schaffhausen, Switzerland

10

11 Corresponding author: Matthias Volk (matthias.volk@agroscope.admin.ch)

[2] verschoben (Einfügung)

Gelöscht: reveals increased

Gelöscht: under

Gelöscht: The rising productivity of alpine grassland under warming, drought and N-deposition treatments

[2] nach oben verschoben: Sub-alpine grassland reveals increased productivity under warmer and drier conditions, but not with higher N-deposition, in an altitudinal transplantation experiment¶

Gelöscht: ¶

21 **Abstract**

22 Multiple global change drivers affect plant productivity of grasslands, and thus ecosystem services like forage
23 production and the soil carbon sink. Subalpine grasslands seem particularly affected and may serve as a proxy
24 for the cold, continental grasslands of the northern hemisphere. Here, we conducted a four-year field-experiment
25 (AlpGrass) with 216 turf monoliths, subjected to three global change drivers: warming, moisture and N-
26 deposition. Monoliths from six different sub-alpine pastures were transplanted to a common location with six
27 climate scenario sites (CS). CS' were located along an altitudinal gradient from 2360 to 1680 m a.s.l.,
28 representing an April - October mean temperature change of -1.4 °C to +3.0 °C, compared to CS_{reference} with no
29 temperature change and with climate conditions comparable to the sites of origin. To uncouple temperature
30 effects along the altitudinal gradient from soil moisture and soil fertility effects, an irrigation treatment (+12-
31 21 % of ambient precipitation) and an N-deposition treatment (+3 kg and +15 kg N ha⁻¹ a⁻¹) was applied in a
32 factorial design, the latter simulating a fertilizing air pollution effect.
33 Moderate warming led to increased productivity. Across the four-year experimental period, the mean annual
34 yield peaked at intermediate CSs (+43 % at +0.7 °C and +44 % at +1.8 °C), coinciding with c. 50 % of days
35 with less than 40 % soil moisture during the growing season. The yield increase was smaller at the lowest,
36 warmest CS (+3.0 °C), but was still 12 % larger than at CS_{reference}. These yield-differences among CSs were well
37 explained by differences in soil moisture and received thermal energy. Irrigation had a significant effect on yield
38 (+16-19 %) in dry years, whereas atmospheric N-deposition did not result in a significant yield response. We
39 conclude that productivity of semi-natural, highly diverse subalpine grassland will increase in the near future.
40 Despite increasingly limiting soil water content, plant growth will respond positively to up to +1.8 °C warming
41 during the growing period, corresponding to +1.3 °C annual mean warming.

- Gelöscht: , the
- Gelöscht: largest single terrestrial biome.
- Gelöscht: E
- Gelöscht: ,
- Gelöscht: soil protection
- Gelöscht: depend on this plant productivity
- Gelöscht: good
- Gelöscht: vast
- Gelöscht: Thus
- Gelöscht: W
- Gelöscht: warming × moisture × N-deposition
- Gelöscht: factorially
- Gelöscht: ×
- Gelöscht: ,
- Gelöscht: ×
- Gelöscht: from six different subalpine pastures (sites of origin)
- Gelöscht: At a common location,
- Gelöscht: T
- Gelöscht: t
- Gelöscht: he m
- Gelöscht: re
- Gelöscht: at
- Gelöscht: at
- Gelöscht: We further applied
- Gelöscht: dry
- Gelöscht: (growing-season-days with soil moisture <40 %).
- Gelöscht: Days with dry less soil moisture explained the average

71 **1 Introduction**

72 The present period of global warming is most pronounced in the cold regions of high altitude and high latitude
73 (Core writing team, IPCC 2014). The productivity of these ecosystems is temperature-limited, and even though
74 the temporal distribution of total annual radiation differs, they share many similarities. After the temperature
75 decline following the Holocene climate optimum (ca. 9000 - 6000 a BP; Vinther et al., 2009), they are now
76 experiencing a rapid rewarming.

77 In cold environments, the perspective on climate change is different compared to temperate and warm
78 environments. First, mitigation of the thermal growth limitation is likely to have beneficial effects on plant
79 growth. Second, the warming-associated drought-risk is lower. The evaporative demand is much lower and at
80 least the initial water supply for plant growth is granted, because even a small winter snowpack supplies a large
81 soil moisture resource in spring. Third, in many regions the warming comes along with rising atmospheric
82 nitrogen (N) deposition, originating from agriculture and fossil fuel burning. Atmospheric N deposition can be
83 as little as <5 kg N ha⁻¹ a⁻¹ at remote mountain sites (Rihm and Kurz, 2001), but can reach rates >40 kg N ha⁻¹ a⁻¹
84 elsewhere in Switzerland (Rihm and Achermann, 2016). This fertilizing air pollution agent promotes plant
85 growth and has the potential to reduce plant species diversity by favoring fast growing species (Vitousek et al.,
86 1997; Bobbink et al., 2010; Phoenix et al., 2012). Alone and in interaction, all three factors increase the
87 ecosystem plant productivity potential in cold regions. Given that essential ecosystem services strongly co-
88 depend on plant productivity (e.g., forage supply for livestock and wildlife, soil erosion control and support of
89 the biological carbon sink), an improved knowledge on how climate warming affects productivity of colder
90 grassland systems is required.

91 However, previous warming experiments on plant productivity have shown inconsistent results. For example,
92 tundra vegetation showed an up to twofold productivity increase in response to increased summer temperature
93 (Van der Wal and Stien, 2014). In contrast, Liu et al. (2018) combined long-term observations with a
94 manipulative experiment to find that total net primary productivity (NPP) in Tibetan grassland remained
95 unaffected, though the relative abundance of grasses was increased at the expense of forbs and sedges by
96 drought and warmth. In yet another meta-analysis, only 13 out of 20 experimental grassland sites revealed small
97 increases of plant productivity due to warming (Rustad et al., 2001); while grassland ecosystems in general
98 showed both positive and negative responses, the colder tundra systems (high latitude or altitude) with lower
99 precipitation had positive and larger productivity responses to warming.

100 To make matters more complicated, evapotranspiration will increase in warming experiments. The resulting,
101 temperature-confounded lower soil moisture makes it impossible to determine the proper temperature effect on
102 plant growth. Only comparing the plant growth response at warming-confounded, reduced soil moisture and at
103 experimentally mitigated soil moisture allows to distinguish warming effects from moisture effects.

104 A common restriction for the usability of climate change experiments for ecosystem productivity projections
105 lies in the low number of concurrently manipulated environmental factors (Rustad 2008; but see Dukes et al.,
106 2005 for an exception). This potentially leads to an overestimation of effects when data from several, single
107 factor experiments are combined in meta-analyses or models (Leuzinger et al., 2011). Indeed, productivity
108 responses to combined factors are usually less than additive in size, compared to single treatment responses
109 (Dieleman et al., 2012; Xu et al., 2013). Not only can a low number of treatment factors, but also a low number
110 of treatment levels invite overly simplistic interpretation of experimental results, if only a short or linear

Gelöscht: ,

Gelöscht: ¶

Gelöscht: ,

Gelöscht: driven by

Gelöscht: were favored over

Gelöscht:

Gelöscht: Given that essential ecosystem services strongly co-depend on plant productivity (e.g., forage supply for livestock and wildlife, soil erosion control and support of the biological carbon sink), an improved knowledge on how climate warming affects productivity of colder grassland - systems is required.¶

123 segment out of a larger range of biologically possible responses is represented in the data. For example, a hump-
 124 shaped response curve (2-dimensional) under atmospheric N-deposition best described the properties of a soil
 125 C-sink in subalpine grassland (Volk et al., 2016). Similarly, a ridge-shaped response surface (3-dimensional),
 126 driven by temperature and precipitation during 17 experimental years, was needed to explain NPP data (Zhu et
 127 al., 2016). These findings suggest that the outcome of a global change productivity-experiment depends to some
 128 degree on the chosen treatment levels and their interaction with the ambient climate during the experiment.
 129 Combining multiple treatments with many levels might thus improve interpretation of experimental outcomes
 130 and related climate change predictions.
 131 Here, we present four years of results from a field experiment in the Swiss Alps. Turf monoliths from a variety
 132 of grassland communities at six different sites of origin were transplanted to one common experimental site to
 133 test for a plant productivity response that is not restricted to a specific species composition. At the common site,
 134 transplanted turf monoliths were distributed over six levels of altitude to generate a climate gradient. Doing so,
 135 we included not only the temperature change, but also the changing length of the growing period. The between-
 136 year weather variability created a large variety of climate situations within the range of potential growth
 137 conditions. Additionally, to uncouple temperature effects along the altitudinal gradient from soil moisture and
 138 soil fertility effects, a two-level irrigation treatment and a three-level atmospheric N-deposition treatment were
 139 set up in a factorial design. We hypothesized that

- 140 1) The effect of warming on plant growth would be beneficial at moderate warming levels, but detrimental at
- 141 high warming levels.
- 142 2) Increased soil water content would mitigate the detrimental effects of excessive warming levels.
- 143 3) N-deposition would exhibit a generally favorable effect on plant growth. This effect would further
- 144 increase with higher temperatures and irrigation due to their mitigating effect on thermal and water co-
- 145 limitations.

- Gelöscht:** a large
- Gelöscht:** factor levels
- Gelöscht:** s
- Gelöscht:** factors
- Gelöscht:** over larger gradients
- Gelöscht:** -
- Gelöscht:** treatment
- Gelöscht:** We used a variety of grassland communities by transplanting t
- Gelöscht:** ,
- Gelöscht:** observe
- Gelöscht:** T

158 **2 Materials and Methods**

159 This experiment (AlpGrass experiment) used grassland monoliths to investigate climate change effects on
160 subalpine pasture ecosystems in the central Alps. At six different sites in the Canton Graubünden, Switzerland,
161 areas of 1 ha on southerly exposed, moderate slopes were selected at an altitude of ca. 2150 m a.s.l. to serve as
162 'sites of origin'. All six sites were mountain grassland used for summer livestock grazing, within ≤ 55 km
163 distance of each other, but their soil (typical depth 20-30 cm) developed either on basic or on acidic bedrock.
164 Thus, the sites of origin shared very similar climatic conditions, but represented a wide range of soil properties
165 and plant communities. Plant communities at the sites of origin were generally dominated by grass and sedge
166 species, but comprised also a substantial share of forbs and a few legume species. Two of the summer pastures
167 were characterized by Sesleretalia vegetation (with *Sesleria caerulea*, *Anthyllis vulneraria*, *Helianthemum*
168 *nummularium* present at both); the other four were dominated by Nardetalia vegetation (with typical species
169 *Nardus stricta*, *Leontodon helveticus*, and *Potentilla aurea*). *Nardus stricta*, *Polygonum viviparum*, and *Carex*
170 *sempervirens* were present in almost all monoliths, regardless of grassland type. Detailed information on soil
171 properties and species composition of the different origins can be found in Wüst-Galley et al. (2020).
172 In June 2012 a total of 252 monoliths (6 sites of origin × 42 monoliths) of 0.1 m² surface area (L × W × H = 37
173 × 27 × 22 cm) were excavated at the sites of origin. Randomly generated X-Y-coordinates were used to choose
174 the location of excavation. If a distinct location had sufficiently deep soil and no rocks, if bare soil and woody
175 species were < 10 %, and if there was no apparent dominance of single plant species, then monoliths were
176 extracted. Else, the next pair of coordinates was probed. Monoliths were placed into precisely-fitting, well-
177 drained plastic boxes to facilitate future transport and avoid potential side effects of experimental treatments
178 applied later (Appendix Fig. A1). To minimize the disturbance of temperature and moisture conditions,
179 monoliths were immediately reinserted into the ground at their respective site of origin.

180 Half a year later, in November 2012, 36 monoliths were transported from each site of origin to the common
181 AlpGrass experimental site, while 6 monoliths each remained at their original site to allow for an assessment of
182 the transplanting effect. Standardizing harvests were done in 2012 and 2013, to homogenize the canopy of the
183 previously grazed monoliths that had more heterogeneous canopies than mown grassland.

184
185 **2.1 Experimental site and treatment design**

186 The AlpGrass experimental site is located on the south slope of Piz Cotschen (3029 m), above Ardez in the
187 Lower Engadine valley (Graubünden, Switzerland). The site as a whole covers a 680 m altitudinal gradient,
188 characterized by a vegetation change from montane forest (WGS 84 N 46.77818°, E 10.17143°) to subalpine
189 grassland (WGS 84 N 46.79858°, E 10.17843°). Along the gradient, six separate climate scenario sites (CS)
190 were located at different altitudes (CS1: 2360 m, CS2: 2170 m, CS3: 2040 m, CS4: 1940 m, CS5: 1830 m, CS6:
191 1680 m a.s.l.). Photographs of the environment can be found in the Appendix (Photographs A 1-4). Because
192 CS2 had a similar altitude as the sites of origin, it was chosen as a reference site (hereafter CS2_{reference}).
193 CS2_{reference} and sites of origin are all characterized by cold winters with permanent snow cover. The snow-free
194 period lasts approximately from May to October, with a mean April – October air temperature of 6.5 °C during
195 the experiment (Tab. 1). Annual mean temperature at CS2_{reference} was 3.2 °C and mean precipitation sum was
196 748 mm (Tab. 2).

Gelöscht: (MLs)

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Gelöscht:

Formatiert: Schriftart: Kursiv

Gelöscht: More Eextensive

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: ,

Gelöscht: Standardizing harvests were done in 2012 (zero-year) and 2013 (acclimation), while quantitative harvests used in this analysis continued from 2014 to 2017.

211 At each of the 6 CS, 6 **monoliths**, from each of the six sites of origin were installed in the ground within their
 212 drained plastic boxes, flush with the surrounding grassland surface, resulting in 36 **monoliths**, per CS and a total
 213 of 216 transplanted **monoliths** (Appendix Figs. A2-A4). **Monoliths in their containers** were arranged side by side
 214 without a separating gap or buffer zone. The grassland surrounding the **monoliths**, was frequently mown to
 215 prevent the introduction of new species/genotypes by seed dispersal.
 216 At each CS, an irrigation and an N-deposition treatment were set up in a **full-factorial design**. One half of the 36
 217 **monoliths** (3 **monoliths**, per site of origin) received only ambient precipitation and no additional water, the other
 218 half received additional water during the growing season. Within each irrigation treatment, **monoliths**, were
 219 subjected to an N treatment representing three levels of atmospheric N-deposition (treatment details below, and
 220 see Appendix Tab. A1 for a schematic description). At each CS, irrigation and N treatments were arranged in a
 221 randomized complete block design (six blocks each containing all six irrigation × N treatment combinations).
 222 Moreover, **monoliths**, of the six sites of origin were assigned to the six blocks by restricted randomization so that
 223 an equal distribution of sites of origin to all blocks was ensured.

- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: MLs.
- Gelöscht:
- Gelöscht: ; occasional gaps between monolithsMLs and the surrounding turf were filled with soil to prevent air flow
- Gelöscht: MLs
- Gelöscht: cross
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: MLs

Site	Alt. (m)	Air temp. (Mean, °C) ±1SE		Δ T (°C)	DD0°C _{total}	Pre-harvest period	
		Apr. – Oct.	annual			Apr. – Oct.	Mean ±1SE
CS1	2360	5.1 ±0.17	1.6 ±0.20	-1.4	1156 ±50	78	±4.3
CS2 _{reference}	2170	6.5 ±0.17	3.2 ±0.23	0.0	1440 ±43	91	±3.8
CS3	2040	7.2 ±0.17	3.7 ±0.20	0.7	1649 ±67	107	±4.4
CS4	1940	8.0 ±0.16	4.7 ±0.25	1.5	1746 ±71	104	±2.8
CS5	1830	8.3 ±0.17	4.6 ±0.21	1.8	1829 ±10	97	±3.4
CS6	1680	9.5 ±0.17	5.8 ±0.21	3.0	2095 ±14	104	±3.5

225 **Table 1** Climatic parameter means across years (±1SE) at the climate scenario sites (CS) during the experiment:
 226 Mean air temperature from April to October and for the whole year, April – Oct. air temperature difference (Δ
 227 T) of respective CS' compared to CS2_{reference}. Degree days above 0 °C for the snow free period between annual
 228 harvests (DD0°C_{total}). Pre-harvest period length is the number of days between snow-melt and harvest.

- Formatiert: Einzug: Hängend: 0.05 cm
- Formatiert: Rechts: 0.38 cm
- Formatiert: Einzug: Hängend: 0.05 cm
- Formatiert: Rechts: 0.38 cm
- Formatiert: Einzug: Hängend: 0.05 cm
- Formatiert: Rechts: 0.38 cm
- Formatiert: Einzug: Hängend: 0.05 cm
- Formatiert: Rechts: 0.38 cm
- Formatiert: Einzug: Hängend: 0.05 cm
- Formatiert: Rechts: 0.38 cm

Site	Alt. (m)	Precipitation (sum, mm)		Dry days (%)		Harvest Date (Ø)
		Apr. – Oct.	annual	not irrigated	irrigated	
CS1	2360	674 ±18	752 ±20	27 ±5.3	17 ±5.1	12 Aug
CS2 _{reference}	2170	656 ±27	748 ±27	31 ±1.7	20 ±2.7	26 July
CS3	2040	629 ±26	732 ±21	42 ±5.2	24 ±4.3	22 July
CS4	1940	614 ±20	739 ±22	33 ±2.2	24 ±3.5	14 July
CS5	1830	628 ±20	780 ±17	55 ±4.4	41 ±5.0	09 July
CS6	1680	570 ±19	687 ±21	73 ±3.1	53 ±4.5	05 July

231 **Table 2** Precipitation sums for the climate scenario sites, aggregated from April to October and annually. For
 232 comparison: The closest Swiss Federal Office for Meteorology station (Scuol, 1303 m a.s.l., 9 km distance)
 233 reported 662 mm mean annual precipitation during the experiment. Dry days (%) indicates the percentage of

246 days during the pre-harvest period with SWC <40 %. The phenology triggered harvest date reflects the delayed
247 vegetation development at higher altitudes.

248 **2.2 Climate scenario site (CS) climate change treatment**

249 The different altitudes of the CSs created a climate change scenario treatment, commencing in November 2012,
250 when the monoliths were installed at the AlpGrass site, and ending in 2017 with the final harvest. The difference
251 in altitude between the sites of origin and the respective CS at the AlpGrass experimental site determined the
252 change of climatic conditions that the transplanted monoliths experienced. These conditions include the mean
253 growing period temperature, from April to October. We assumed the evenly moderate temperature (ca. 0 °C)
254 under the winter snow cover to be of little importance for differences in ecosystem productivity. The CS
255 temperature treatment was specified as the deviation from CS_{reference} temperature. The available thermal energy
256 was expressed as degree days (DD) above a threshold of 0 °C (DD0°C). To derive DD, the sum of hourly
257 temperature means above 0°C during one day was calculated and, then, divided by 24 hours. To quantify the total
258 thermal energy available for growth, DD during the snow-free period between the annual harvests (DD0°C_{total})
259 was summed up, considering that the perennial vegetation continues to grow after mowing.
260 Differences in volumetric soil water content (SWC) were quantified as the proportion of days during the
261 growing period with a SWC < 40 % (hereafter 'dry days'). This < 40 %-threshold does not necessarily imply
262 plant growth limitation, but it was developed to reliably contrast the soil moisture status between the CSs and
263 between years. Thus, more time below the threshold indicates a 'drier period' in relative terms.

265 **2.3 Irrigation treatment**

266 An irrigation treatment with two levels was set up to distinguish the warming effect from the soil moisture
267 effect, driven by warming. In several applications throughout the growing period, precipitation equivalents of 20
268 mm were applied to the monoliths under the irrigation treatment. The total amount of water added per monolith
269 was 80, 120, 120 and 80 mm in 2014, 2015, 2016 and 2017, respectively. These amounts were equivalent to 12 -
270 21 % of the recorded precipitation sum during the growing periods.

272 **2.4 N-deposition treatment**

273 The N-deposition treatment consisted of three levels. Atmospheric N-deposition from air pollution was
274 simulated to amount to a deposition of 3 and 15 kg N ha⁻¹ a⁻¹, on top of the present background deposition. We
275 used a 200 ml ammonium nitrate (NH₄⁺ NO₃⁻)/water solution per monolith, which was applied in twelve, ca. bi-
276 weekly fractions, covering the growing period. Monoliths without additional N-deposition received water
277 without ammonium nitrate.

279 **2.5 Meteorology**

280 At all six CS we measured air temperature, relative humidity (Hygroclip 2 in an unspirated radiation shield,
281 Rotronic, Switzerland), and precipitation (ARG100 tipping bucket raingauge, Campbell Scientific, UK). Soil
282 temperature and SWC were measured at 8 cm depth in 6 monoliths, each at topmost CS1 and intermediate CS3,
283 CS4 and CS5, using a SWC reflectometer with 12 cm rods (CS655, Campbell Scientific, UK). At CS_{reference} and
284 lowest CS6 these values were measured in 18 monoliths, and two points in the surrounding grassland. The
285 measurement interval for all parameters was 10 minutes originally and was later integrated for longer periods as
286 necessary.

Gelöscht: MLs

Gelöscht: MLs

Gelöscht: we calculated

Gelöscht: ,

Gelöscht: we

Gelöscht: this sum

Gelöscht: we summed degree days

Gelöscht: The thermal energy was expressed as degree day values (DD0°C), resulting from hourly air temperature means above a threshold of 0 °C, added for one day, then divided by 24. To quantify the total thermal energy available we summed degree days during the snow-free period between the annual harvests (DD0°C_{total}), considering that the perennial vegetation continues to grow after mowing.

Gelöscht: We quantified d

Formatiert: Nicht Hervorheben

Gelöscht: D

Formatiert: Nicht Hervorheben

Formatiert: Nicht Hervorheben

Gelöscht: dubbed

Gelöscht: were quantified as 'percent (%) dry days'. This represents the proportion of days during the growing period with a SWC < 40 %.

Gelöscht: e

Gelöscht: strong

Gelöscht: , while SWC is actually close to 40

Gelöscht: %

Gelöscht: provided a good

Gelöscht: for

Gelöscht: differences in the

Formatiert: Englisch (Vereinigte Staaten)

Gelöscht: means

Gelöscht: MLs

Gelöscht: ML

Gelöscht: MLs

Gelöscht: MLs

319 At each site of origin, we installed Hobo U12-008 data loggers with TMC-HD sensors (Onset Computer
320 Corporation, USA) in three monoliths and one spot in the undisturbed, surrounding grassland for comparison
321 with the reference climate scenario site CS_{2reference}.
322 Ambient wet N-deposition was measured at CS_{2reference} and lowest CS6 using bulk samplers (VDI 4320 Part 3,
323 2017; c.f. Thimonier et al., 2019) between April 2013 and April 2015. Nitrate (NO₃⁻) in rainwater and melted
324 snow was analyzed by ion chromatography (ICS-1600, Dionex, USA) and NH₄⁺ was analyzed using a flow
325 injection analyzer (FIAstar 5000, Foss, Denmark) with gas diffusion membrane, detection was completed with
326 UV/VIS photometry (SN EN ISO 11732).

327 2.6 Plant productivity

329 All plant material (including mosses and lichens) of the monoliths, was cut 2 cm above the soil surface once per
330 year at canopy maturity. This plant removal serves as a proxy for the short, but intensive summer grazing period
331 of the traditional management. As a result of the phenology-triggered harvests (anthesis of *Festuca rubra*), the
332 topmost CS1 was cut on average 38 days later than the lowest CS6. Plants were dried at 60 °C, allowed to cool
333 in a desiccator and weighed to determine dry matter yield (hereafter biomass yield).

334 2.7 Data analyses

336 Data were analyzed by linear mixed-effects models. First, we were interested in the overall response of biomass
337 yield over years as affected by the treatment factors. To this aim, biomass yield was averaged across the four
338 experimental years (2014-2017) and was modeled as function of CS (factor of 6 levels), irrigation (factor of 2
339 levels), and N-deposition (factor of 3 levels), including all interactions. ‘Site of origin’ (6 sites) and block (36
340 levels: 6 CS × 6 blocks) were modeled as random factors (random intercepts). Restricted maximum likelihood
341 was used for parameter estimation. For the inference on fixed effects, the Kenward–Roger method was applied
342 to determine the approximate denominator degrees of freedom (Kenward and Roger 1997), and the marginal
343 and conditional R² of the model were computed following Nakagawa and Schielzeth (2013). Differences in
344 biomass yield between single CSs and the CS_{reference} were tested based on the model contrasts (post-hoc *t*-tests,
345 without using multiple comparisons). To receive additional insight into within year treatment effects, this very
346 same model was also applied to data of each of the four individual years.

347 Second, to consider the time effect and the repeated structure of the data, biomass yield of all four years was
348 modeled as function of year (factor of 4 levels), CS, irrigation, and N-deposition (factor levels as described),
349 including all interactions. Here, random factors consisted of an identifier for monolith (216 levels) to consider
350 the potential correlation of monoliths’ biomass yield over years (modeled as random intercept). In addition, the
351 model included the random factor ‘site of origin’ and allowed for a separate block term at each of the four years
352 (details as described). Residuals of all models were evaluated for normality and homoscedasticity and fulfilled
353 assumptions of linear mixed-effects models. Finally, to gain insight into effects of thermal energy and drought
354 on plant productivity, biomass yield was modeled as function of each DD0°C_{total} and percent days with less soil
355 moisture (‘dry days’) using generalized additive models (GAM). Generalized additive models had to be used as
356 simple linear models could not appropriately handle these relationships. The GAMs included the fixed factor
357 irrigation and a smooth term for the continuous variables DD0°C_{total} and percent dry days, respectively, for both
358 levels of the irrigation treatment. Model validation revealed that the assumptions of GAMs were met : more

Gelöscht: MLs

Gelöscht: regression

Gelöscht: regressed

Gelöscht: on

Gelöscht: regressed on

Gelöscht: MLs

Gelöscht: MLs’

Gelöscht: regression

Gelöscht: dry

368 [information on the model specification is given in the Appendix.](#) All data was analyzed with the statistics
369 software R, version 4.0.2 (R Core Team 2020) and packages lme4 for linear-mixed effect models (Bates et al.,
370 2015) and mgcv for GAMs (Wood, 2017).

Gelöscht: .

372 **3. Results**

373

374 **3.1 Climate scenario site (CS) environmental conditions**

375 **3.1.1 Low atmospheric background N-deposition**

376 Total N-deposition was 3.3 kg N ha⁻¹ a⁻¹ at CS2_{reference} and 4.3 kg N ha⁻¹ a⁻¹ at the lowest CS6. The seasonal
377 distribution showed peak deposition rates in June and July.

378

379 **3.1.2 Consistent temperature, precipitation and drought changes with altitude**

380 The mean Apr. – Oct. temperature gradient of up to +3 °C compared to CS2_{reference}, distributed over four
381 altitudinal levels (CS3 – CS6), constituted the warming treatment. Conversely, temperature at the topmost CS1
382 constituted a cooling treatment (Δ temp. -1.4 °C), extending the range of temperature responses tested (Δ temp.,
383 Tab. 1). As intended, the DD0°C_{total} steadily increased from CS2_{reference} to lowest CS6. The pre-harvest period
384 (PHP) length was fairly similar among CSs, because the early snow-melt at the lower CS was compensated by
385 an early harvest (Tab. 1).

386 We observed a small, non-linear increase of precipitation with altitude during April – October. The recorded
387 annual precipitation sum was somewhat larger than the sum for the growing period (Tab. 2).

388 The length of the period with less soil moisture (% dry days) doubled along the altitudinal gradient: At the two
389 top CSs only one third of the pre-harvest period was dry, compared to two thirds of the time at the lowest site
390 CS6 (compare Tables 1 & 2). The irrigation treatment reduced the incidence of days with < 40 % soil moisture
391 to 60-80 % of the non-irrigated situation (Tab. 2).

392

393 **3.1.3 Small transplantation effects on soil temperature and moisture**

394 At the sites of origin, the mean April – October soil temperatures in the undisturbed grassland were 8.8 ° (±0.3)
395 compared to 8.9 ° (±0.3) in the monoliths. At CS2_{reference} this difference was 9.2 ° vs. 9.5 °C. Thus, the
396 surrounding grassland at CS2_{reference} site was on average 0.4 °C warmer than at the sites of origin, and monoliths
397 at CS2_{reference} were 0.3 °C warmer than the undisturbed grassland surrounding the experiment. Volumetric SWC
398 in the undisturbed grassland was 1 % lower on average compared to SWC in the monoliths at CS2_{reference} and
399 lowest CS6.

400

401 **3.2. Yield**

402 **3.2.1 Insignificant transplantation effect**

403 The mean annual yield was 20 % larger at CS2_{reference} (control treatment monoliths), compared to the origins
404 (162 g m⁻²; ±12.7), but not significantly different (P = 0.19; paired, two-sided t-test). Equally important, the
405 difference showed no trend, as in some years the yield at CS2_{reference} was higher, in some years it was lower
406 compared to the sites of origin.

407

408 **3.2.2 Strongest climate scenario site effect at intermediate CS**

409 Across the four years, we found a highly significant effect of the CS on aboveground biomass yield (Tab. 3). At
410 intermediate sites, yields increased by +43 %, +18 % and +44 % (CS3, CS4 and CS5, respectively; Tab. 4, P ≤
411 0.05 at least), related to +0.7, +1.5, and +1.8 °C of the warming component of the respective CS (compare Table

Gelöscht: continuous

Gelöscht: dry

Gelöscht: dry

Gelöscht: MLs

Gelöscht: I

Gelöscht: warming

Gelöscht: yields

Gelöscht: at sites

Gelöscht: .

Gelöscht: (

Gelöscht: increases being

Gelöscht: and an increase between 209 and 389 of DD0°C_{total}

425 1). Even at the warmest site CS6 the yield was still +12 % larger compared to the CS2_{reference} site ($\Delta T = +3 \text{ }^\circ\text{C}$,
 426 $\text{ADD0}^\circ\text{C}_{\text{total}} = 655$). The coldest site CS1 was not less productive than CS2_{reference}. In the year of the overall
 427 maximum productivity (2016), also, the coldest site CS1 and the warmest site CS6 produced their respective
 428 record yield (Tab. 4). Overall, the yields of the 24 combinations of year \times CS varied by a factor of 2.1 (yields
 429 averaged across irrigation and N-deposition treatments). The yield response to CSs differed between years
 430 (Appendix Tab. A2, year \times CS interaction: $P < 0.001$) in that the CS effect became weaker towards the end of
 431 the experiment (Appendix Tab. A3).

Gelöscht: both

Variable	df _{num}	df _{den}	F-value	P
Climate Scenario (CS)	5	29.1	14.9	< 0.001
Irrigation	1	145.2	6.5	0.012
N	2	145.2	1.3	0.287
CS \times Irrigation	5	145.2	1.1	0.352
CS \times N	10	145.2	0.5	0.864
Irrigation \times N	2	145.2	1.1	0.348
CS \times Irrigation \times N	10	145.2	1.3	0.241

df_{num}: degrees of freedom of term; df_{den}: degrees of freedom of error (which can be fractional in restricted maximum likelihood analysis)

434 **Table 3** Summary of analyses for the effects of climate scenario site (CS), irrigation and N deposition on
 435 aboveground biomass yield of subalpine grassland. Data were averaged across the four experimental years (total
 436 n = 216). F-tests refer to the fixed effects of the linear mixed-effects model. The marginal and conditional R^2
 437 were 0.41 and 0.50, respectively.

439
440
441

Year	CS2 _{reference}		Aboveground biomass yield (g m ⁻² , means ±1SE)									
	% dry days	DD0°C _{total}	CS1	CS2 _{reference}	CS3	CS4	CS5	CS6	CS mean			
2014	30	1353	149 ^{ns} ±8.0	170 ±11.0	238 ^{***} ±8.8	203* ±11.6	255 ^{***} ±15.2	152 ^{ns} ±10.5	194 ±5			
2015	38	1359	147 ^{ns} ±8.1	138 ±5.8	248 ^{***} ±12.1	171† ±8.9	310 ^{***} ±13.6	198 ^{***} ±8.8	202 ±5			
2016	22	1509	230 ^{ns} ±8.7	222 ±9.1	297 ^{***} ±10.2	247 ^{ns} ±11.1	271 ^{**} ±15.3	250† ±9.8	253 ±4			
2017	34	1541	152 ^{ns} ±8.5	166 ±7.8	208* ±10.0	201* ±11.7	169 ^{ns} ±9.1	176 ^{ns} ±8.3	178 ±4			
Mean	36	1440	170 ^{ns} ±7.1	174 ±6.9	248 ^{***} ±7.9	205* ±9.0	251 ^{***} ±11.5	194 ^{ns} ±6.9				

*** $P \leq 0.001$, ** $P \leq 0.01$, * $P \leq 0.05$, † $P \leq 0.1$, ^{ns} $P > 0.1$

Table 4 Aboveground biomass yield (means ±1SE) per CS and year, averaged across irrigation, N-deposition treatments, and site of origin. Within each year, significance tests are against CS2_{reference}, based on contrasts derived from linear mixed-effects models (see Table 3 and Appendix Tab. A1, for the respective model summaries). Shaded values indicate the CS with the greatest aboveground biomass yield per year and across years.


- Formatiert: Hervorheben
- Formatiert: Hervorheben
- Formatiert: Hervorheben
- Formatiert: Hervorheben
- Formatiert: Hervorheben
- Gelöscht: the model
- Gelöscht: regression
- Gelöscht: y

449 **3.2.3 Irrigation effect in dry years**

450 Despite a mere +7.7 % average yield increase (Fig. 1 b), irrigation turned out to be a significant factor across years
 451 (Table 3). Yet, the effect of irrigation differed between years (Appendix Tab. A2, year × irrigation interaction: $P <$
 452 0.001), and single years analysis detected positive effects of irrigation only in 2015 (+15.8 %) and 2017 (+18.8 %)
 453 (Appendix Tab. A3). In these years, the percentage of days with $< 40\%$ soil moisture was highest.

454
 455 **3.2.4 No nitrogen deposition effect**

456 Five years of experimentally increased atmospheric nitrogen deposition (+3 and +15 kg N ha⁻¹ a⁻¹) did not cause a
 457 significant response of biomass yield. Moreover, there was no significant interaction detected between the N-
 458 treatment and the factors CS or irrigation (Fig. 1 a; Tab. 3). Single years analysis, to test for a late response to
 459 accumulating amounts of N, revealed a marginally significant effect only in 2016 (Appendix Tab. A3).



460 **Figure 1 a, b** Aboveground biomass yield as a function of the altitude of CSs. Data were averaged across years;
 461 circles denote means ±1 SE. Warming and dry days (%) increase with decreasing altitude from left to right. a) Yield
 462 values grouped by N-deposition treatment (0, 3 and 15 kg N ha⁻¹ a⁻¹, in addition to 4-5 kg N background deposition).
 463

Gelöscht: B)

Gelöscht: dry

[1] nach unten verschoben: (Fig. 1 A; Tab. 3)

[1] verschoben (Einfügung)

Gelöscht: A

Formatiert: Englisch (Vereinigte Staaten)

Gelöscht: A

Gelöscht: B

Gelöscht: A

471 **b)** Yield values grouped by irrigation treatment. Overlapping means and SEs are shifted horizontally to improve
 472 their visibility.

474 **3.2.5 Yield at climate scenario sites strongly relates to changes in thermal energy and soil moisture.**

475 Biomass yield at the different CSs was significantly related to thermal energy, expressed as growing $DD0^{\circ}C_{total}$.
 476 Here, intermediate CSs (CS3, CS5) had greatest yields at intermediate values of $DD0^{\circ}C_{total}$, indicated by the
 477 curvature of the predicted line in particular under irrigated conditions (Fig. 2 **a**, Appendix Tab. A4, smooth term for
 478 $DD0^{\circ}C_{total}$: $P < 0.001$).

479 Similarly, biomass yield was significantly related to $\%$ days with soil moisture $< 40\%$ ('dry days') during the growing
 480 season, with intermediate CS3 and CS5 having highest yields at around 50 % of dry days under no irrigation and at
 481 around 30 % dry days under additional irrigation (Fig. 2 **b**, Appendix Tab. A5, smooth term for dry days: $P <$
 482 0.001). Under unirrigated conditions, in parallel with a doubling of dry days (from 27 % at topmost CS1 to 55 % at
 483 intermediate CS5), yield consistently rose and only fell at the driest and warmest site CS6, with 73 % dry days.

485 **Figure 2 a, b.** Aboveground biomass yield at the six CS as **a)** a function of total received thermal energy
 486 ($DD0^{\circ}C_{total}$), and **b)** percent of days with dry soil (SWC $< 40\%$) during the growing season (dry days %). Data were
 487 averaged across years; circles denote means ± 1 SE per CS and irrigation treatment. The predicted line is based on a
 488 generalized additive model using all data (± 1 SE light grey shaded). Dark grey indicates the cross-section of the two
 489 SE bands. Overlapping means and SEs in **(a)** are shifted horizontally to improve their visibility.

Gelöscht: B

Gelöscht: Climate scenario yields

Gelöscht: temperature

Gelöscht: changes

Gelöscht: A

Gelöscht: soil moisture content, expressed as percent of

Gelöscht: dry

Gelöscht: (SWC

Gelöscht:)

Gelöscht: B


Gelöscht: A

Gelöscht: B

Gelöscht: A

Gelöscht: B

Gelöscht: A



505 **4 Discussion**

506 **We found a substantial and significant positive effect of climate scenarios, equivalent to warming of up to + 1.8 °C**
507 **(Apr. – Oct. mean), on aboveground biomass of subalpine grasslands (up to +44 % yield).** Contrary to expectation,
508 additional resource supply through irrigation and N-deposition had only marginal (water) or no effects (N) on yield,
509 respectively. Our transplanting experiment proved to be efficient in assessing several linked climate change drivers
510 in their effect on plant growth.

511

512 **4.1 Climate scenario temperature effects**

513 The phenology-triggered harvest opened the possibility to extend the growing period in cool years and shorten the
514 exposure to drought stress in warm years. Thus, beneficial thermal effects were maximized, while detrimental
515 drought effects were minimized. As a consequence, we displayed the yield over a continuous x-axis of degree days
516 between harvests ($DD0^{\circ}C_{total}$, Fig. 2 A). This represents the available thermal resource, associated with a particular
517 yield, much better than mean temperatures of CS, or categorical values for CS1-CS6.

518 In cold environments, the warming is so important because the metabolic growth processes, which utilize the
519 assimilated energy, are strongly temperature dependent, much more so than the assimilation process *per se* (Körner
520 2003). In a meta-analysis of grassland responses to warming that included 32 sites, distinctly positive warming
521 effects on growth were found in the colder portion of those ecosystems (Rustad et al., 2001), very similar to
522 responses in the subalpine grassland of the current study. Interestingly, also the response size of our effects is in the
523 same range as that reported by Rustad et al., (2001).

524 Plant growth at the intermediate climate scenarios that represented a warming of 0.7 °C, 1.5 °C and 1.8 °C (Apr.-
525 Oct.) clearly benefitted from greater warmth. However, the increase of responses was somewhat inconsistent (CS4
526 ca. +18 %, CS3 and CS5 both > +40 %), matching only partly our first hypothesis. The erratic response of
527 intermediate CS4 is likely the result of an interaction of micro-topography effects on climate that were not detected
528 by our meteorological measurements, cockchafer **infestation** (*Melolontha melolontha*: **bug whose larvae fed on**
529 **roots**), or the occurrence of mast years in some species at that CS. In the extreme treatment at lowest CS6 (+3 °C
530 Apr.-Oct., +2.4 °C annual mean) the positive response to warming finally ceased to increase, but yield was still

531 somewhat larger than at CS2_{reference}. **This comparatively low growth response suggests that the water supply at CS6**
532 **had already reached critically low levels; yet, the larger thermal energy resource must have partly compensated for**
533 **the radically smaller soil water resource, leading also to some growth benefit at CS6 (compare Figure 2 a & b).**

534 Despite substantial cooling at topmost CS1, coinciding with a temperature decline of -1.4 °C, the mean yields for
535 CS1 and CS2_{reference} were very similar (Tab. 4, Fig. 1). This is indicative of a plant community that is well cold-
536 adapted. Indeed, local historical records from the Swiss Federal Office for Meteorology (MeteoSwiss) show that
537 only 100 years ago the local April-October mean air temperature was 1.4-1.5 °C lower than today (30 a running
538 mean, courtesy P. Calanca using MeteoSwiss data from Segl-Maria site at 1804 m a.s.l.). In effect, the cooling
539 upward-transplantation represented a climatic time travel of 100 years into the past, **and the similar yield responses**
540 **between CS1 and CS2_{reference} indicate that subalpine grassland productivity may not have changed much during the**
541 **past century. Moreover, the dramatic temperature dynamics during the past 12,000 years of the Holocene**

Gelöscht: We found a substantial and significant positive effect of climate scenario warming (up to + 1.8 °C) on aboveground biomass yield of subalpine grasslands (up to +44 %).

Gelöscht: infestation

Gelöscht: er

Gelöscht: This demonstrates that the growth benefit from

Gelöscht: a

Gelöscht: This demonstrates that the increased thermal growth resource compensated for a radically reduced soil water resource (compare Figure 2 A & B).

Gelöscht: .

Gelöscht: Also

Gelöscht: that occurred

556 interglacial, suggest that temperature adaptations in modern plant genotypes may actually match not only today's
557 weather, but also warmer and cooler climate conditions. From this perspective, and with respect to the genetic
558 memory of plants, the undiminished productivity at topmost CS1 illustrates that assumed 'control' temperatures in
559 warming experiments only represent the most recent point of an extremely dynamic climatic history.

561 4.2 Climate scenario soil moisture effects

562 The differences in soil moisture content that resulted naturally from 24 different climatic situations (6 CS/altitude
563 levels × 4 years) created a hump-shaped response curve of yield over drought (Fig. 2 b). This does imply that, with
564 decreasing altitude and increasing warmth, productivity rose despite more days with soil moisture < 40 %.

565 The importance of soil moisture for plant growth has been shown predominantly in much drier grasslands, e.g., in
566 warmer prairie (Xu et al., 2013) or cold alpine grassland (Wang et al., 2013), where release from drought stress
567 benefitted growth. For example, along a temperature and altitude gradient in semiarid Tibetan alpine grassland,
568 productivity increased with altitude due to reduced drought stress, but despite decreasing temperatures. Only after an
569 800 m rise in altitude, productivity eventually became smaller, and further reduced drought stress did not constitute
570 a further advantage on plant growth (Wang et al., 2016).

571 In our experiment, soil moisture values and its proxies integrate information on moisture and temperature. Thus, the
572 two-dimensional growth response curve along the altitudinal gradient, peaking at the least detrimental situation
573 between moisture limitation and thermal limitation (Fig. 2 b), is analogous to the three-dimensional response surface
574 found in the Jasper Ridge experiment (Zhu et al., 2016). Unfortunately, our experiment did not produce a sufficient
575 number of data points for a 3-D presentation. Based on these results, we infer that a joint evaluation of soil moisture
576 and temperature is mandatory to assess reliably warming effects of climate change on plant growth in the subalpine
577 environment.

579 4.3 Irrigation treatment

580 We had assumed that increased SWC would mitigate detrimental effects of excessive warming. Surprisingly
581 however, the overall irrigation effect on yield was not very substantial, despite large differences in the percentages
582 of days with soil moisture < 40% during the growing season (Table 2, Fig. 2 b). Moreover, the positive responses
583 did not increase consistently with warmth, but were strongest at the intermediate CS3 and CS4 (Fig. 1 b). Analyses
584 of individual years showed that the two significant responses of annual yield to irrigation coincided with the two
585 driest years. This evidence suggests that maximum mitigation of (low) temperature limitation requires simultaneous
586 release of water limitation, while at the same time the amount of water applied in our study was insufficient to
587 compensate for increased evapotranspiration at CS5 and the warmest site CS6.

589 4.4 N-deposition treatment

590 We hypothesized a generally positive effect of N-deposition on plant growth, but found no significant overall effect
591 of N-deposition on yield after four years and only a marginal effect in one year. Historically, the responsiveness of
592 (sub-)alpine vegetation to improved nutrient supply was considered to be restricted due to an overriding effect of

Gelöscht: ,

Gelöscht: , that are still contained

Gelöscht: ,

Gelöscht: Also the dramatic temperature dynamics during the 12,000 a of the present Holocene interglacial suggest that temperature adaptations, contained in modern plant genotypes, may actually match not only today's climate conditions.

Gelöscht: is not surprising. Instead, it

Gelöscht: , with respect to the genetic memory of plants

Gelöscht: B

Gelöscht: dry

Gelöscht: B

Gelöscht: reliably

Gelöscht: dry

Gelöscht: B

Gelöscht: B

Gelöscht: .

Gelöscht:

612 thermal energy limitation. Yet, studies with very high rates of N application (40-100 kg N ha⁻¹ a⁻¹; Körner et al.,
613 1997; Heer and Körner 2002) showed substantial yield responses, also at alpine sites. Low N-dose responses of total
614 plant yield may require N-accumulation over years or a compound interest effect in plant biomass. For example,
615 only in the seventh treatment year a strong, +31 % total yield growth response to 5 kg N ha⁻¹ a⁻¹ was reported by
616 Volk et al., (2014) from subalpine grassland.

617 Low dose experiments (5-30 kg N ha⁻¹ a⁻¹), however, can induce a species composition change (Bowman et al.,
618 2012), indicating a growth benefit for some species at the expense of others. Yet, such single species responses may
619 be only transient: a strong *Carex* species response to as little as 5 kg N ha⁻¹ a⁻¹ in similar subalpine vegetation was
620 recently found to cease after five years (Bassin et al., 2009 and 2013). Taken together, we conclude that the cold-
621 adapted, mature and low productivity grassland either responds with a >4 year time lag, or that the N-deposition
622 treatment was below the critical load for aboveground biomass responses.

624 4.5 Transplantation

625 The turf monoliths at CS2_{reference} were only slightly warmer and moister compared to the sites of origin, suggesting a
626 low transplantation impact (we have found no transplantation effect data from other experiments to compare with).
627 However, within the experimental site similar temperature increases between CS2_{reference} and CS3 caused a much
628 larger productivity increase (+43 %). We reason that this incongruence can be explained by the difference in melt-
629 out time, which was on average only 3 days earlier at CS2_{reference} (julian day 118) than at the sites of origin, but 21
630 days earlier at CS3 than at CS2_{reference}. We thus assume that the substantially earlier start of the growing season
631 caused the stronger growth response, despite a similar temperature change. This effect, induced by the
632 transplantation of the grassland monoliths along the altitudinal gradient, demonstrates the importance of integrating
633 multiple drivers in climate change experiments to allow for a multi-factor driven plant response.

634 In our study, the effect of altitude on photosynthesis substrate limitation was considered negligible, compared to the
635 climate effects. The assimilation conditions of alpine plants have been the subject of investigation for decades. Since
636 the theoretical considerations of Gale (1972) and the field studies by Körner and Diemer (1987) and Körner et al.,
637 (1988), a predominant 'altitude-tolerance' of photosynthesis is widely accepted. Relevant environmental parameters
638 that change with altitude (temperature, CO₂ and O₂ partial pressure, vapor-pressure deficit and photosynthetic
639 photon flux density) have antagonistic effects on assimilation efficiency (see Wang et al., (2016) for a recent
640 discussion on the topic).

Gelöscht: ¶

Single key species on the other hand showed immediate positive responses to realistic N-deposition rates (20-25 kg N ha⁻¹ a⁻¹; Bowman et al., 2006; Bassin et al., 2007; Inauen et al., 2012). Similarly, a l

Gelöscht: found no total aboveground biomass response

Gelöscht: but

Gelöscht: However

Gelöscht:

Gelöscht: ¶

Indeed, *Carex spp.* can support a positive N-deposition growth response, but only until warming and drought create a competitive advantage for grasses over sedges (Liu et al., 2018; Wüst-Galley et al., 2020). Thus, the latest studies suggest that there is a positive N-deposition × warming interaction on the response of *Carex spp.*¶
In our experiment, we found no significant overall effect of N-deposition on yield after five years and only a marginal effect in one year. We thus

Gelöscht: 5

Gelöscht: MLs

662 **5 Conclusions**

663 Despite dwindling soil water content, the subalpine grassland growth increased to up to +1.8 °C warming during the
664 growing period (corresponding to +1.3 °C annual mean), compared to present temperatures. Even at the maximum
665 warming (corresponding to +2.4 °C annual mean) the yield was larger than at the reference site. At the same time -
666 1.4 °C cooling during the growing period (corresponding to -1.7 °C annual mean) did not reduce plant growth. This
667 implies that subalpine grassland productivity has likely not increased during the past century warming, but, despite
668 growing soil moisture deficits, will do so with continued warming in the near future.

669 **Author contribution**

670 MV and SB designed the experiment, MV, ALW and SB conducted field work. MV and MS analyzed the data. MV
671 led the writing of the manuscript, with significant contribution from MS. All authors contributed critically to the
672 drafts and gave final approval for publication.

673

674 **Data availability**

675 [The data analyzed for the current study are available in the Dryad Digital Repository at ...](#)

676

677 **Competing interests**

678 The authors declare that they have no conflict of interest.

679

680 **Acknowledgements**

681 We received essential financial support through the Federal Office for the Environment (contract No. 00.5100.PZ /
682 R442-1499). The Federal Office for Meteorology (MeteoSwiss) is kindly acknowledged for providing access to
683 meteorological data. We thank Pierluigi Calanca for handling these data. The Gemeinde Ardez and Alpmeister
684 Claudio Franziscus generously allowed us to work on the Allmend. N-concentration analyses courtesy of
685 Forschungsstelle für Umweltbeobachtung (FUB-AG, Rapperswil, Switzerland). We are grateful to Robin Giger for
686 his untiring support in the field and the lab, and to the scientific site manager Andreas Gauer, who was in charge of
687 the field sites.

Gelöscht: Data will be made available immediately to individuals upon request by the corresponding author. At a later point data will be deposited upon request at a publicly accessible repository, according to Swiss federal research institution guidelines.¶

693 **References**

- 694 Bassin, S., Volk, M., Suter, M., Buchmann, N., Fuhrer, J.: Nitrogen deposition but not ozone affects productivity
695 and community composition of subalpine grassland after 3 yr of treatment. *New Phytologist* 175, 3, 523-534,
696 2007.
- 697 Bassin, S., Werner, R. A., Sörgel, K., Volk, M., Buchmann, N., Fuhrer, J.: Effects of combined ozone and nitrogen
698 deposition on the in situ properties of eleven key plant species of a subalpine pasture. *Oecologia* 158, 4, 747-756,
699 2009.
- 700 Bassin, S., Volk, M., Fuhrer, J.: Species composition of subalpine grassland is sensitive to nitrogen deposition, but
701 not to ozone, after seven years of treatment. *Ecosystems* 16, 6, 1105-1117, 2013.
- 702 Bates, D., Maechler, M., Bolker, B., Walker, S.: lme4: Linear mixed-effects models using Eigen and S4. Version
703 1.1-10. <https://CRAN.R-project.org/package=lme4>, 2015.
- 704 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S.,
705 Davidson, E., Dentener, F., Emmett, B., Erisman, J. W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L., De Vries,
706 W.: Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecological*
707 *applications* 20, 1, 30-59, 2010.
- 708 Bowman, W. D., Gartner, J. R., Holland, K., Wiedermann, M.: Nitrogen Critical Loads For Alpine Vegetation And
709 Terrestrial Ecosystem Response: Are We There Yet? *Ecological Applications* 16, 1183-1193, 2006.
- 710 Bowman, W. D., Murgel, J., Blett, T., Porter, E.: Nitrogen critical loads for alpine vegetation and soils in Rocky
711 Mountain National Park. *Journal of Environmental Management* 103, 165-171, 2012.
- 712 Core Writing Team, Pachauri, R. K., Meyer, L. A. editors: IPCC, 2014: Climate change 2014: Synthesis Report.
713 Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on
714 Climate Change. IPCC, Geneva, Switzerland. 151p, 2014.
- 715 Dieleman, W. I., Vicca, S., Dijkstra, F. A., Hagedorn, F., Hovenden, M.J., Larsen, K.S., ... & King, J.: Simple
716 additive effects are rare: a quantitative review of plant biomass and soil process responses to combined
717 manipulations of CO₂ and temperature. *Global Change Biology* 18, 9, 2681-2693, 2012.
- 718 Dukes, J. S., Chiariello, N. R., Cleland, E. E., Moore, L. A., Shaw, M. R., Thayer, S., Tobeck, T., Mooney, H. A.,
719 Field, C. B.: Responses of grassland production to single and multiple global environmental changes. *PLoS*
720 *Biology* 3, 10, e319, 2005.
- 721 Heer, C. and Körner, C.: High elevation pioneer plants are sensitive to mineral nutrient addition. *Basic and Applied*
722 *Ecology*, 3, 1, 39-47, 2002.
- 723 Gale, J.: Availability of carbon dioxide for photosynthesis at high altitudes: theoretical considerations. *Ecology*, 53,
724 3, 494-497, 1972.
- 725 Inauen, N., Körner, C., Hiltbrunner, E.: No growth stimulation by CO₂ enrichment in alpine glacier forefield plants.
726 *Global Change Biology*, 18, 3, 985-999, 2012.
- 727 Kenward, M. G., Roger, J. H.: Small sample inference for fixed effects from restricted maximum likelihood.
728 *Biometrics*, 53, 3, 983-997, 1997.

729 Körner, C. and Diemer, M.: In situ photosynthetic responses to light, temperature and carbon dioxide in herbaceous
730 plants from low and high altitude. *Functional Ecology*, 179–194, 1987.

731 Körner, C., Farquhar, G. D., Roksandic, Z.: A global survey of carbon isotope discrimination in plants from high
732 altitude. *Oecologia*, 74, 623-632, 1988.

733 Körner, C., Diemer, M., Schächli, B., Niklaus, P., Arnone III J.: The responses of alpine grassland to four seasons of
734 CO₂ enrichment: a synthesis. *Acta Oecologica* 18, 3, 165-175, 1997.

735 Körner, C.: *Alpine plant life: functional plant ecology of high mountain ecosystems*. Springer Science & Business
736 Media. 344p., 2003.

737 Leuzinger, S., Luo, Y., Beier, C., Dieleman, W., Vicca, S., Körner, C.: Do global change experiments overestimate
738 impacts on terrestrial ecosystems? *Trends in ecology and evolution* 26, 5, 236-241, 2011.

739 Liu, H., Mi, Z., Lin, L., Wang, Y., Zhang, Z., Zhang, F., ... & Zhao, X.: Shifting plant species composition in
740 response to climate change stabilizes grassland primary production. *Proceedings of the National Academy of*
741 *Sciences* 115, 16, 4051-4056, 2018.

742 Nakagawa, S. and Schielzeth, H.: A general and simple method for obtaining R² from generalized linear mixed-
743 effects models. *Methods in ecology and evolution*, 4, 2, 133-142, 2013.

744 Phoenix, G. K., Emmett, B. A., Britton, A. J., Caporn, S. J.M., Dise, N. B., Helliwell, R., Jones, L., Leake, J. R.,
745 Leith, I. D., Sheppard, L. J., Sowerby, A., Pilkington, M. G., Rowe, E. C., Ashmore, M. R., Power, S. A.:
746 Impacts of atmospheric nitrogen deposition: responses of multiple plant and soil parameters across contrasting
747 ecosystems in long-term field experiments. *Global Change Biology*, 18, 1197–1215, 2012.

748 R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing,
749 Vienna, Austria, 2020. <http://www.R-project.org>.

750 Rihm, B. and Kurz, D.: Deposition and critical loads of nitrogen in Switzerland. In *Acid rain 2000*. pp. 1223-1228,
751 Springer, Dordrecht, 2001.

752 Rihm, B. and Achermann, B. Critical Loads of Nitrogen and their Exceedances. Swiss contribution to the effects-
753 oriented work under the Convention on Long-range Transboundary Air Pollution (UNECE). Federal Office for
754 the Environment, Bern. *Environmental studies* no. 1642, 78p., 2016.

755 Rustad, L. E., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A. ... & Gurevitch, J.: A meta-analysis of
756 the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental
757 ecosystem warming. *Oecologia* 126, 4, 543-562, 2001.

758 Rustad, L. E.: The response of terrestrial ecosystems to global climate change: towards an integrated approach.
759 *Science of the total environment* 404, 2-3, 222-235, 2008.

760 Thimonier, A., Kosonen, Z., Braun, S., Rihm, B., Schleppi, P., Schmitt, M., ... & Thöni, L.: Total deposition of
761 nitrogen in Swiss forests: Comparison of assessment methods and evaluation of changes over two decades.
762 *Atmospheric Environment* 198, 335-350, 2019.

763 Van Der Wal, R. and Stien, A.: High-arctic plants like it hot: A long-term investigation of between-year variability
764 in plant biomass. *Ecology* 95, 12, 3414-3427, 2014.

765 Vinther, B. M., Buchardt, S. L., Clausen, H. B., Dahl-Jensen, D., Johnsen, S. J., Fisher, D. A. ... & Blunier, T.:
766 Holocene thinning of the Greenland ice sheet. *Nature* 461, 7262, 385, 2009.

767 Vitousek, P. M., Aber, J., Howarth, R. W., Likens, G. E., Matson, P. A., Schindler, D. W., Schlesinger, W. H.,
768 Tilman, G. D.: Human alteration of the global nitrogen cycle: causes and consequences. *Ecological Applications*
769 7, 737–750, 1997.

770 Volk, M., Wolff, V., Bassin, S., Ammann, C., Fuhrer, J.: High tolerance of subalpine grassland to long-term ozone
771 exposure is independent of N input and climatic drivers. *Environmental Pollution* 189, 161-168, 2014.

772 Volk, M., Enderle, J., Bassin, S.: Subalpine grassland carbon balance during 7 years of increased atmospheric N
773 deposition. *Biogeosciences* 13, 12, 3807-3817, 2016.

774 Wang, Z., Luo, T., Li, R., Tang, Y., Du, M.: Causes for the unimodal pattern of biomass and productivity in alpine
775 grasslands along a large altitudinal gradient in semi-arid regions. *Journal of Vegetation Science* 24, 1, 189-201,
776 2013.

777 Wang, H., Prentice, I. C., Davis, T. W., Keenan, T. F., Wright, I. J., Peng, C.: Photosynthetic responses to altitude:
778 an explanation based on optimality principles. *New Phytologist* 213, 3, 976-982, 2016.

779 Wood, S. N.: *Generalized Additive Models: An Introduction with R*. 2nd edition. Chapman and Hall/CRC. London,
780 2017.

781 Wüst-Galley, C., Volk, M., Bassin, S.: Interaction of climate change and nitrogen deposition on subalpine pastures.
782 (in revision) 2020.

783 Xu, X., Sherry, R. A., Niu, S., Li, D., Luo, Y.: Net primary productivity and rain-use efficiency as affected by
784 warming, altered precipitation, and clipping in a mixed-grass prairie. *Global Change Biology* 19, 9, 2753-2764,
785 2013.

786 Zhu, K., Chiariello, N. R., Tobeck, T., Fukami, T., Field, C. B.: Nonlinear, interacting responses to climate limit
787 grassland production under global change. *Proceedings of the National Academy of Sciences* 113, 38, 10589-
788 10594, 2016.

789 **Appendix for**

790 **Sub-alpine grassland productivity increased with warmer and**
791 **drier conditions, but not with higher N-deposition, in an**
792 **altitudinal transplantation experiment**

793

794 Matthias Volk¹, Matthias Suter², Anne-Lena Wahl¹, Seraina Bassin^{1,3}

795 ¹Climate and Agriculture, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

796 ²Forage Production and Grassland Systems, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

797 ³Pädagogische Hochschule Schaffhausen, Ebnetstrasse 80, 8200 Schaffhausen, Switzerland

798

799 Corresponding author: Matthias Volk (matthias.volk@agroscope.admin.ch)

Gelöscht: ¶

Gelöscht: unter

Formatiert: MS title, Links, Abstand Vor: 0 Pt., Abstand zwischen Absätzen gleicher Formatierung einfügen

Gelöscht: reveals increased

Gelöscht: The rising productivity of alpine grassland under warming, drought and N-deposition treatments

Formatiert: Schriftart: Nicht Fett, Englisch (Vereinigte Staaten)

805 **Appendix Tables**

806 **Table A1** Schematic layout of monolith arrangement at each CS of the AlpGrass experimental site. At each CS, six
 807 monoliths, from each of six sites of origin were transplanted, resulting in 36 monoliths. Two irrigation and three N-
 808 deposition treatments were set up in a factorial design, resulting in six irrigation × N treatment combinations, which
 809 were assigned to each of the six monoliths per site of origin. The six irrigation × N treatment combinations were
 810 arranged in a randomized complete block design of six blocks. Regarding sites of origin, the monoliths were assigned
 811 to the six blocks in a restricted randomization, so that an equal distribution of sites of origin to all blocks was ensured.
 812 It follows that the six monoliths, from each site of origin received all irrigation × N treatment combinations and were
 813 evenly distributed on the site. Displayed is a possible randomization of irrigation and N treatments per block; at each
 814 CS separate randomizations were performed.

- Gelöscht: (ML)
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: cross-
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: MLs
- Gelöscht: calculated

Block 1			Block 2			Block 3		
W0.N15	W1.N0	W0.N3	W0.N3	W0.N0	W1.N15	W1.N15	W0.N0	W1.N0
W0.N0	W1.N3	W1.N15	W1.N3	W0.N15	W1.N0	W1.N3	W0.N15	W0.N3

W1.N15	W1.N0	W0.N0	W1.N3	W0.N3	W1.N0	W0.N0	W1.N0	W0.N15
W0.N3	W0.N15	W1.N3	W0.N0	W1.N15	W0.N15	W1.N3	W1.N15	W0.N3
Block 4			Block 5			Block 6		

815 W0: no additional water (ambient precipitation only), W1: additional water during
 816 growing period; N0: no N fertilizer, N3: 3 kg N ha⁻¹ a⁻¹, N15: 15 kg N ha⁻¹ a⁻¹

825 **Table A2** Summary of analyses for the effects of climate scenario (CS), irrigation, and N deposition on
 826 aboveground biomass yield of subalpine grassland over four experimental years. *F*-tests refer to the fixed
 827 effects of a linear mixed-effects model that included all four years for a repeated measures analysis. The
 828 marginal and conditional R^2 were 0.68 and 0.80, respectively.

829

Variable	df _{num}	df _{den}	<i>F</i> -value	<i>P</i>
Year	3	45.5	66.2	< 0.001
Climate Scenario (CS)	5	198.0	18.3	< 0.001
Irrigation	1	166.6	6.2	0.014
N	2	166.6	1.2	0.304
Year × CS	15	63.0	9.6	< 0.001
Year × Irrigation	3	450.5	13.6	< 0.001
Year × N	6	450.5	0.9	0.492
CS × Irrigation	5	166.6	1.1	0.380
CS × N	10	166.6	0.5	0.882
Irrigation × N	2	166.6	1.0	0.365
Year × CS × Irrigation	15	450.5	2.9	< 0.001
Year × CS × N	30	450.5	0.8	0.749
Year × Irrigation × N	6	450.5	1.4	0.199
CS × Irrigation × N	10	166.6	1.2	0.275
Year × CS × Irrigation × N	30	450.5	1.4	0.066

830 df_{num}: degrees of freedom of term; df_{den}: degrees of freedom of error (which can be fractional
 831 in restricted maximum likelihood analysis)

832 **Table A3** Summary of analyses for the effects of climate scenario (CS), irrigation, and N deposition on aboveground biomass yield of subalpine grassland at each of
 833 four experimental years (2014 – 2017). *F*-tests refer to the fixed effects of a linear mixed-effects model to each of the four years.
 834

Variable	df _{num}	2014			2015			2016			2017		
		df _{den}	<i>F</i> -value	<i>P</i>	df _{den}	<i>F</i> -value	<i>P</i>	df _{den}	<i>F</i> -value	<i>P</i>	df _{den}	<i>F</i> -value	<i>P</i>
Climate Scenario (CS)	5	28.9	17.2	< 0.001	29.5	24.9	< 0.001	29.3	4.5	0.004	29.4	4.0	0.006
Irrigation	1	145.2	1.5	0.224	145.1	21.6	< 0.001	145.3	1.1	0.290	145.4	19.2	< 0.001
N	2	145.2	0.7	0.481	145.1	0.5	0.610	145.3	2.6	0.078	145.4	0.3	0.728
CS × Irrigation	5	145.2	2.3	0.048	145.1	2.0	0.080	145.3	1.8	0.126	145.4	0.8	0.563
CS × N	10	145.2	0.5	0.912	145.1	0.7	0.751	145.3	0.9	0.531	145.4	0.5	0.896
Irrigation × N	2	145.2	1.9	0.151	145.1	0.8	0.448	145.3	0.7	0.509	145.4	1.2	0.290
CS × Irrigation × N	10	145.2	1.5	0.157	145.1	1.0	0.429	145.3	1.5	0.157	145.4	1.3	0.226

835 df_{num}: degrees of freedom of term; df_{den}: degrees of freedom of error (which can be fractional in restricted maximum likelihood analysis)
 836

837 **Table A4** Summary of analyses for the effects of total received thermal energy ($DD0^{\circ}C_{total}$) on aboveground
 838 biomass yield of subalpine grassland under two irrigation treatments. Data were averaged across the four
 839 experimental years (total n = 216). *F*-values and approximate *P*-values refer to a generalized additive model that
 840 used a smooth term for each irrigation treatment.

841

Parametric terms	df	<i>F</i> -value	<i>P</i>
Irrigation	2	1613.0	< 0.001
Smooth terms	edf	<i>F</i> -value	<i>P</i>
s($DD0^{\circ}C_{total}$) – No irrigation	1.72	7.7	< 0.001
s($DD0^{\circ}C_{total}$) – Additional irrigation	2.34	10.2	< 0.001

842 df: degrees of freedom; edf: effective degrees of freedom (which can be fractional
 843 in smooth terms of generalized additive models)

844 s: smoothing function applied on term

845

846

847

848

849

850 **Table A5** Summary of analyses for the effects of percent days with soil moisture < 40% during the growing
 851 season (dry days %) on aboveground biomass yield of subalpine grassland under two irrigation treatments. Data
 852 were averaged across the four experimental years (total n = 216). *F*-values and approximate *P*-values refer to a
 853 generalized additive model that used a smooth term for each irrigation treatment.

Gelöscht: dry

854

Parametric terms	df	<i>F</i> -value	<i>P</i>
Irrigation	2	402.9	< 0.001
Smooth terms	edf	<i>F</i> -value	<i>P</i>
s(Dry days %) – No irrigation	2.55	11.3	< 0.001
s(Dry days %) – Additional irrigation	2.59	8.1	< 0.001

855 df: degrees of freedom; edf: effective degrees of freedom (which can be fractional
 856 in smooth terms of generalized additive models)

857 s: smoothing function applied on term

859 **Appendix Photographs**

Formatiert: Englisch (Vereinigte Staaten)

Gelöscht: Figures

860 **Photograph A1** Monolith produced at site of origin “Alp Nova” (46.72786°N, 9.72609°E) in June 2012. After
861 monoliths were excavated in the close surroundings, they were fitted tightly into plastic containers. A total of
862 216 monoliths was produced at six such sites of origin, for later use at the altitudinal transplantation site of the
863 AlpGrass Experiment.



864
865
866 **Photograph A2** Topmost climate scenario site CS1 (2360 m, 46.79859°N, 10.17840°E). Monoliths are in the
867 right part of the fenced area. Monolith arrangement in two double rows of nine allows easy access and equal
868 distribution of edge-effects.



870 **Photograph A3** Control climate scenario site CS2_{reference} (2170 m, 46.79264°N, 10.17714°E). Along the
871 altitudinal transplantation gradient this CS is representative of the sites of origin, because the share the same
872 altitude.



873
874
875 **Photograph A4** Lowest climate scenario site CS6 (1680 m, 46.77818°N, 10.17143°E). Due to its low altitude
876 this CS is the warmest and driest site along the altitudinal transplantation gradient.



```

877 Appendix R codes
878 #####
879 #Linear mixed-effects model to analyze effects of the climate scenario treatment (CS), irrigation
880 and N deposition (N_Treat) on aboveground biomass yield.
881
882 #Package to load
883 library(lme4)
884
885 #####
886 #Reading in the data
887 d.data <- read.table("C:/Volk_etal_2021_AlpsGrass.csv", header=TRUE, sep=";")
888
889 #####
890 #Define factors
891 d.data$CS <- as.factor(d.data$CS)
892 d.data$Irrigation <- as.factor(d.data$Irrigation)
893 d.data$N_Treat <- as.factor(d.data$N_Treat)
894 d.data$Origin <- as.factor(d.data$Origin)
895 d.data$Block <- as.factor(d.data$Block)
896
897
898 #####
899 #Full model, including all interactions, as described in the first paragraph of the 'Data analyses'
900 section
901
902 Model.A <- lmer(DM ~ CS + Irrigation + N_Treat +
903               CS:Irrigation + CS:N_Treat + Irrigation:N_Treat + CS:Irrigation:N_Treat +
904               (1 | Origin) + (1 | Block), REML=TRUE, data=d.data)
905
906 #####
907 #The model summary, given in Table 3, is received by
908
909 library(lmerTest)
910 anova(Model.A, ddf="Kenward-Roger", type=1)
911
912
913 #####
914 #Contrasts to test for differences in biomass yield between single CSs and the CSreference (across
915 irrigation and the N treatments)
916
917 summary( update(Model.A, contrasts=list(CS=contr.treatment(levels(d.data$CS),base=2),
918 Irrigation="contr.sum", N_Treat="contr.sum")), ddf="Kenward-Roger")
919
920
921 #####
922 #This very same model and contrast code was applied to data of each individual year.

```

Formatiert: Englisch (Vereinigte Staaten)

Gelöscht :

Formatiert: Englisch (Vereinigte Staaten)

```

924 #####
925 Generalized additive models to test for the effects of thermal energy (DD0Ctot) and percent days
926 with less soil moisture (PercDryDays) on aboveground biomass yield.
927
928 #Package to load
929 library(mgcv)
930
931 #####
932 #GAM for the effect of thermal energy on yield
933
934 Model.B <- gam(DM ~ -1 + Irrigation + s(DD0Ctot, by=Irrigation), gamma=3.6,
935             knots=list(DD0Ctot=rep(seq(from=min(d.data$DD0Ctot)+100,
936             to=max(d.data$DD0Ctot)-100, length.out=12), each=18)),
937             method="REML", data=d.data)
938
939 #####
940 #The model summary, given in Table A4, is received by
941
942 anova(Model.B)
943
944 #####
945 #GAM for the effect of percent dry days on yield
946
947 Model.C <- gam(DM ~ -1 + Irrigation + s(PercDryDays, by=Irrigation), gamma=1.7,
948             knots=list(PercDryDays=rep(seq(from=min(d.data$PercDryDays)+5,
949             to=max(d.data$PercDryDays)-5, length.out=12), each=18)),
950             method="REML", data=d.data)
951
952 #####
953 #The model summary, given in Table A5, is received by
954
955 anova(Model.C)
956
957 #####
958 Note: The fitted lines in Figure 2a) & b) are based on the predicted values from Model.B and
959 Model.C, respectively.

```