

1 **Sub-alpine grassland productivity increased with warmer and**  
2 **drier conditions, but not with higher N-deposition, in an**  
3 **altitudinal transplantation experiment**

4

5

6 Matthias Volk<sup>1</sup>, Matthias Suter<sup>2</sup>, Anne-Lena Wahl<sup>1</sup>, Seraina Bassin<sup>1,3</sup>

7 <sup>1</sup>Climate and Agriculture, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

8 <sup>2</sup>Forage Production and Grassland Systems, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

9 <sup>3</sup>Pädagogische Hochschule Schaffhausen, Ebnetstrasse 80, 8200 Schaffhausen, Switzerland

10

11 Corresponding author: Matthias Volk ([matthias.volk@agroscope.admin.ch](mailto:matthias.volk@agroscope.admin.ch))

12 **Abstract**

13 Multiple global change drivers affect plant productivity of grasslands and thus ecosystem services like forage  
14 production and the soil carbon sink. Subalpine grasslands seem particularly affected and may serve as a proxy  
15 for the cold, continental grasslands of the northern hemisphere. Here, we conducted a four-year field-experiment  
16 (AlpGrass) with 216 turf monoliths, subjected to three global change drivers: warming, moisture and N-  
17 deposition. Monoliths from six different sub-alpine pastures were transplanted to a common location with six  
18 climate scenario sites (CS). CS' were located along an altitudinal gradient from 2360 to 1680 m a.s.l.,  
19 representing an April - October mean temperature change of  $-1.4\text{ }^{\circ}\text{C}$  to  $+3.0\text{ }^{\circ}\text{C}$ , compared to  $\text{CS}_{\text{reference}}$  with no  
20 temperature change and with climate conditions comparable to the sites of origin. To uncouple temperature  
21 effects along the altitudinal gradient from soil moisture and soil fertility effects, an irrigation treatment ( $+12\text{-}$   
22  $21\%$  of ambient precipitation) and an N-deposition treatment ( $+3\text{ kg}$  and  $+15\text{ kg N ha}^{-1}\text{ a}^{-1}$ ) was applied in a  
23 factorial design, the latter simulating a fertilizing air pollution effect.

24 Moderate warming led to increased productivity. Across the four-year experimental period, the mean annual  
25 yield peaked at intermediate CSs ( $+43\%$  at  $+0.7\text{ }^{\circ}\text{C}$  and  $+44\%$  at  $+1.8\text{ }^{\circ}\text{C}$ ), coinciding with c.  $50\%$  of days  
26 with less than  $40\%$  soil moisture during the growing season. The yield increase was smaller at the lowest,  
27 warmest CS ( $+3.0\text{ }^{\circ}\text{C}$ ), but was still  $12\%$  larger than at  $\text{CS}_{\text{reference}}$ . These yield-differences among CSs were well  
28 explained by differences in soil moisture and received thermal energy. Irrigation had a significant effect on yield  
29 ( $+16\text{-}19\%$ ) in dry years, whereas atmospheric N-deposition did not result in a significant yield response. We  
30 conclude that productivity of semi-natural, highly diverse subalpine grassland will increase in the near future.  
31 Despite increasingly limiting soil water content, plant growth will respond positively to up to  $+1.8\text{ }^{\circ}\text{C}$  warming  
32 during the growing period, corresponding to  $+1.3\text{ }^{\circ}\text{C}$  annual mean warming.

## 33 **1 Introduction**

34 The present period of global warming is most pronounced in the cold regions of high altitude and high latitude  
35 (Core writing team, IPCC 2014). The productivity of these ecosystems is temperature-limited, and even though  
36 the temporal distribution of total annual radiation differs, they share many similarities. After the temperature  
37 decline following the Holocene climate optimum (ca. 9000 - 6000 a BP; Vinther et al., 2009), they are now  
38 experiencing a rapid rewarming.

39 In cold environments, the perspective on climate change is different compared to temperate and warm  
40 environments. First, mitigation of the thermal growth limitation is likely to have beneficial effects on plant  
41 growth. Second, the warming-associated drought-risk is lower. The evaporative demand is much lower and at  
42 least the initial water supply for plant growth is granted because even a small winter snowpack supplies a large  
43 soil moisture resource in spring. Third, in many regions the warming comes along with rising atmospheric  
44 nitrogen (N) deposition, originating from agriculture and fossil fuel burning. Atmospheric N deposition can be  
45 as little as  $<5 \text{ kg N ha}^{-1} \text{ a}^{-1}$  at remote mountain sites (Rihm and Kurz, 2001), but can reach rates  $>40 \text{ kg N ha}^{-1} \text{ a}^{-1}$   
46 elsewhere in Switzerland (Rihm and Achermann, 2016). This fertilizing air pollution agent promotes plant  
47 growth and has the potential to reduce plant species diversity by favoring fast growing species (Vitousek et al.,  
48 1997; Bobbink et al., 2010; Phoenix et al., 2012). Alone and in interaction, all three factors increase the  
49 ecosystem plant productivity potential in cold regions. Given that essential ecosystem services strongly co-  
50 depend on plant productivity (e.g., forage supply for livestock and wildlife, soil erosion control and support of  
51 the biological carbon sink), an improved knowledge on how climate warming affects productivity of colder  
52 grassland systems is required.

53 However, previous warming experiments on plant productivity have shown inconsistent results. For example,  
54 tundra vegetation showed an up to twofold productivity increase in response to increased summer temperature  
55 (Van der Wal and Stien, 2014). In contrast, Liu et al. (2018) combined long-term observations with a  
56 manipulative experiment to find that total net primary productivity (NPP) in Tibetan grassland remained  
57 unaffected, though the relative abundance of grasses was increased at the expense of forbs and sedges by  
58 drought and warmth. In yet another meta-analysis, only 13 out of 20 experimental grassland sites revealed small  
59 increases of plant productivity due to warming (Rustad et al., 2001): while grassland ecosystems in general  
60 showed both positive and negative responses, the colder tundra systems (high latitude or altitude) with lower  
61 precipitation had positive and larger productivity responses to warming.

62 To make matters more complicated, evapotranspiration will increase in warming experiments. The resulting,  
63 temperature-confounded lower soil moisture makes it impossible to determine the proper temperature effect on  
64 plant growth. Only comparing the plant growth response at warming-confounded, reduced soil moisture and at  
65 experimentally mitigated soil moisture allows to distinguish warming effects from moisture effects.

66 A common restriction for the usability of climate change experiments for ecosystem productivity projections  
67 lies in the low number of concurrently manipulated environmental factors (Rustad 2008; but see Dukes et al.,  
68 2005 for an exception). This potentially leads to an overestimation of effects when data from several, single  
69 factor experiments are combined in meta-analyses or models (Leuzinger et al., 2011). Indeed, productivity  
70 responses to combined factors are usually less than additive in size, compared to single treatment responses  
71 (Dieleman et al., 2012; Xu et al., 2013). Not only can a low number of treatment factors, but also a low number  
72 of treatment levels invite overly simplistic interpretation of experimental results, if only a short or linear

73 segment out of a larger range of biologically possible responses is represented in the data. For example, a hump-  
74 shaped response curve (2-dimensional) under atmospheric N-deposition best described the properties of a soil  
75 C-sink in subalpine grassland (Volk et al., 2016). Similarly, a ridge-shaped response surface (3-dimensional),  
76 driven by temperature and precipitation during 17 experimental years, was needed to explain NPP data (Zhu et  
77 al., 2016). These findings suggest that the outcome of a global change productivity-experiment depends to some  
78 degree on the chosen treatment levels and their interaction with the ambient climate during the experiment.  
79 Combining multiple treatments with many levels might thus improve interpretation of experimental outcomes  
80 and related climate change predictions.

81 Here, we present four years of results from a field experiment in the Swiss Alps. Turf monoliths from a variety  
82 of grassland communities at six different sites of origin were transplanted to one common experimental site to  
83 test for a plant productivity response that is not restricted to a specific species composition. At the common site,  
84 transplanted turf monoliths were distributed over six levels of altitude to generate a climate gradient. Doing so,  
85 we included not only the temperature change, but also the changing length of the growing period. The between-  
86 year weather variability created a large variety of climate situations within the range of potential growth  
87 conditions. Additionally, to uncouple temperature effects along the altitudinal gradient from soil moisture and  
88 soil fertility effects, a two-level irrigation treatment and a three-level atmospheric N-deposition treatment were  
89 set up in a factorial design. We hypothesized that

- 90 1) The effect of warming on plant growth would be beneficial at moderate warming levels, but detrimental at  
91 high warming levels.
- 92 2) Increased soil water content would mitigate the detrimental effects of excessive warming levels.
- 93 3) N-deposition would exhibit a generally favorable effect on plant growth. This effect would further  
94 increase with higher temperatures and irrigation due to their mitigating effect on thermal and water co-  
95 limitations.

## 96 2 Materials and Methods

97 This experiment (AlpGrass experiment) used grassland monoliths to investigate climate change effects on  
98 subalpine pasture ecosystems in the central Alps. At six different sites in the Canton Graubünden, Switzerland,  
99 areas of 1 ha on southerly exposed, moderate slopes were selected at an altitude of ca. 2150 m a.s.l. to serve as  
100 ‘sites of origin’. All six sites were mountain grassland used for summer livestock grazing, within  $\leq 55$  km  
101 distance of each other, but their soil (typical depth 20-30 cm) developed either on basic or on acidic bedrock.  
102 Thus, the sites of origin shared very similar climatic conditions, but represented a wide range of soil properties  
103 and plant communities. Plant communities at the sites of origin were generally dominated by grass and sedge  
104 species, but comprised also a substantial share of forbs and a few legume species. Two of the summer pastures  
105 were characterized by Sesleretalia vegetation (with *Sesleria caerulea*, *Anthyllis vulneraria*, *Helianthemum*  
106 *nummularium* present at both); the other four were dominated by Nardetalia vegetation (with typical species  
107 *Nardus stricta*, *Leontodon helveticus*, and *Potentilla aurea*). *Nardus stricta*, *Polygonum viviparum*, and *Carex*  
108 *sempervirens* were present in almost all monoliths, regardless of grassland type. Detailed information on soil  
109 properties and species composition of the different origins can be found in Wüst-Galley et al. (2020).  
110 In June 2012 a total of 252 monoliths (6 sites of origin  $\times$  42 monoliths) of 0.1 m<sup>2</sup> surface area ( $L \times W \times H = 37$   
111  $\times 27 \times 22$  cm) were excavated at the sites of origin. Randomly generated X-Y-coordinates were used to choose  
112 the location of excavation. If a distinct location had sufficiently deep soil and no rocks, if bare soil and woody  
113 species were  $< 10\%$ , and if there was no apparent dominance of single plant species, then monoliths were  
114 extracted. Else, the next pair of coordinates was probed. Monoliths were placed into precisely-fitting, well-  
115 drained plastic boxes to facilitate future transport and avoid potential side effects of experimental treatments  
116 applied later (Appendix Fig. A1). To minimize the disturbance of temperature and moisture conditions,  
117 monoliths were immediately reinserted into the ground at their respective site of origin.  
118 Half a year later, in November 2012, 36 monoliths were transported from each site of origin to the common  
119 AlpGrass experimental site, while 6 monoliths each remained at their original site to allow for an assessment of  
120 the transplanting effect. Standardizing harvests were done in 2012 and 2013 to homogenize the canopy of the  
121 previously grazed monoliths that had more heterogeneous canopies than mown grassland.

122

### 123 2.1 Experimental site and treatment design

124 The AlpGrass experimental site is located on the south slope of Piz Cotschen (3029 m), above Ardez in the  
125 Lower Engadine valley (Graubünden, Switzerland). The site as a whole covers a 680 m altitudinal gradient,  
126 characterized by a vegetation change from montane forest (WGS 84 N 46.77818°, E 10.17143°) to subalpine  
127 grassland (WGS 84 N 46.79858°, E 10.17843°). Along the gradient, six separate climate scenario sites (CS)  
128 were located at different altitudes (CS1: 2360 m, CS2: 2170 m, CS3: 2040 m, CS4: 1940 m, CS5: 1830 m, CS6:  
129 1680 m a.s.l.). Photographs of the environment can be found in the Appendix (Photographs A 1-4). Because  
130 CS2 had a similar altitude as the sites of origin, it was chosen as a reference site (hereafter CS2<sub>reference</sub>).  
131 CS2<sub>reference</sub> and sites of origin are all characterized by cold winters with permanent snow cover. The snow-free  
132 period lasts approximately from May to October, with a mean April – October air temperature of 6.5 °C during  
133 the experiment (Tab. 1). Annual mean temperature at CS2<sub>reference</sub> was 3.2 °C and mean precipitation sum was  
134 748 mm (Tab. 2).

135 At each of the 6 CS, 6 monoliths from each of the six sites of origin were installed in the ground within their  
 136 drained plastic boxes, flush with the surrounding grassland surface, resulting in 36 monoliths per CS and a total  
 137 of 216 transplanted monoliths (Appendix Figs. A2-A4). Monoliths in their containers were arranged side by side  
 138 without a separating gap or buffer zone. The grassland surrounding the monoliths was frequently mown to  
 139 prevent the introduction of new species/genotypes by seed dispersal.

140 At each CS, an irrigation and an N-deposition treatment were set up in a full-factorial design. One half of the 36  
 141 monoliths (3 monoliths per site of origin) received only ambient precipitation and no additional water, the other  
 142 half received additional water during the growing season. Within each irrigation treatment, monoliths were  
 143 subjected to an N treatment representing three levels of atmospheric N-deposition (treatment details below, and  
 144 see Appendix Tab. A1 for a schematic description). At each CS, irrigation and N treatments were arranged in a  
 145 randomized complete block design (six blocks each containing all six irrigation × N treatment combinations).  
 146 Moreover, monoliths of the six sites of origin were assigned to the six blocks by restricted randomization so that  
 147 an equal distribution of sites of origin to all blocks was ensured.

148

Site	Alt. (m)	Air temp. (Mean, °C) ±1SE		Δ T (°C)	DD0°C <sub>total</sub>	Pre-harvest period	
		Apr. – Oct.	annual	Apr. – Oct.	Mean ±1SE	# Days	±1SE
CS1	2360	5.1 ±0.17	1.6 ±0.20	-1.4	1156 ±50	78	±4.3
CS2 <sub>reference</sub>	2170	6.5 ±0.17	3.2 ±0.23	0.0	1440 ±43	91	±3.8
CS3	2040	7.2 ±0.17	3.7 ±0.20	0.7	1649 ±67	107	±4.4
CS4	1940	8.0 ±0.16	4.7 ±0.25	1.5	1746 ±71	104	±2.8
CS5	1830	8.3 ±0.17	4.6 ±0.21	1.8	1829 ±10	97	±3.4
CS6	1680	9.5 ±0.17	5.8 ±0.21	3.0	2095 ±14	104	±3.5

149 **Table 1** Climatic parameter means across years (±1SE) at the climate scenario sites (CS) during the experiment:  
 150 Mean air temperature from April to October and for the whole year, April – Oct. air temperature difference (Δ  
 151 T) of respective CS' compared to CS2<sub>reference</sub>. Degree days above 0 °C for the snow free period between annual  
 152 harvests (DD0°C<sub>total</sub>). Pre-harvest period length is the number of days between snow-melt and harvest.

153

154

Site	Alt. (m)	Precipitation (sum, mm)		Dry days (%)		Harvest
		Apr. – Oct.	annual	not irrigated	irrigated	Date (Ø)
CS1	2360	674 ±18	752 ±20	27 ±5.3	17 ±5.1	12 Aug
CS2 <sub>reference</sub>	2170	656 ±27	748 ±27	31 ±1.7	20 ±2.7	26 July
CS3	2040	629 ±26	732 ±21	42 ±5.2	24 ±4.3	22 July
CS4	1940	614 ±20	739 ±22	33 ±2.2	24 ±3.5	14 July
CS5	1830	628 ±20	780 ±17	55 ±4.4	41 ±5.0	09 July
CS6	1680	570 ±19	687 ±21	73 ±3.1	53 ±4.5	05 July

155 **Table 2** Precipitation sums for the climate scenario sites, aggregated from April to October and annually. For  
 156 comparison: The closest Swiss Federal Office for Meteorology station (Scuol, 1303 m a.s.l., 9 km distance)  
 157 reported 662 mm mean annual precipitation during the experiment. Dry days (%) indicates the percentage of

158 days during the pre-harvest period with SWC <40 %. The phenology triggered harvest date reflects the delayed  
159 vegetation development at higher altitudes.

160 **2.2 Climate scenario site (CS) climate change treatment**

161 The different altitudes of the CSs created a climate change scenario treatment, commencing in November 2012,  
162 when the monoliths were installed at the AlpGrass site, and ending in 2017 with the final harvest. The difference  
163 in altitude between the sites of origin and the respective CS at the AlpGrass experimental site determined the  
164 change of climatic conditions that the transplanted monoliths experienced. These conditions include the mean  
165 growing period temperature, from April to October. We assumed the evenly moderate temperature (ca. 0 °C)  
166 under the winter snow cover to be of little importance for differences in ecosystem productivity. The CS  
167 temperature treatment was specified as the deviation from CS2<sub>reference</sub> temperature. The available thermal energy  
168 was expressed as degree days (DD) above a threshold of 0 °C (DD0°C). To derive DD, the sum of hourly  
169 temperature means above 0°C during one day was calculated and then divided by 24 hours. To quantify the total  
170 thermal energy available for growth, DD during the snow-free period between the annual harvests (DD0°C<sub>total</sub>)  
171 was summed up, considering that the perennial vegetation continues to grow after mowing.

172 Differences in volumetric soil water content (SWC) were quantified as the proportion of days during the  
173 growing period with a SWC < 40 % (hereafter 'dry days'). This < 40 %-threshold does not necessarily imply  
174 plant growth limitation, but it was developed to reliably contrast the soil moisture status between the CSs and  
175 between years. Thus, more time below the threshold indicates a 'drier period' in relative terms.

176

177 **2.3 Irrigation treatment**

178 An irrigation treatment with two levels was set up to distinguish the warming effect from the soil moisture  
179 effect, driven by warming. In several applications throughout the growing period, precipitation equivalents of 20  
180 mm were applied to the monoliths under the irrigation treatment. The total amount of water added per monolith  
181 was 80, 120, 120 and 80 mm in 2014, 2015, 2016 and 2017, respectively. These amounts were equivalent to 12-  
182 21 % of the recorded precipitation sum during the growing periods.

183

184 **2.4 N-deposition treatment**

185 The N-deposition treatment consisted of three levels. Atmospheric N-deposition from air pollution was  
186 simulated to amount to a deposition of 3 and 15 kg N ha<sup>-1</sup> a<sup>-1</sup>, on top of the present background deposition. We  
187 used a 200 ml ammonium nitrate (NH<sub>4</sub><sup>+</sup> NO<sub>3</sub><sup>-</sup>)/water solution per monolith, which was applied in twelve, ca. bi-  
188 weekly fractions, covering the growing period. Monoliths without additional N-deposition received water  
189 without ammonium nitrate.

190

191 **2.5 Meteorology**

192 At all six CS we measured air temperature, relative humidity (Hygroclip 2 in an un aspirated radiation shield,  
193 Rotronic, Switzerland), and precipitation (ARG100 tipping bucket raingauge, Campbell Scientific, UK). Soil  
194 temperature and SWC were measured at 8 cm depth in 6 monoliths each at topmost CS1 and intermediate CS3,  
195 CS4 and CS5, using a SWC reflectometer with 12 cm rods (CS655, Campbell Scientific, UK). At CS2<sub>reference</sub> and  
196 lowest CS6 these values were measured in 18 monoliths and two points in the surrounding grassland. The  
197 measurement interval for all parameters was 10 minutes originally and was later integrated for longer periods as  
198 necessary.



199 At each site of origin, we installed Hobo U12-008 data loggers with TMC-HD sensors (Onset Computer  
200 Corporation, USA) in three monoliths and one spot in the undisturbed, surrounding grassland for comparison  
201 with the reference climate scenario site CS<sub>2reference</sub>.  
202 Ambient wet N-deposition was measured at CS<sub>2reference</sub> and lowest CS6 using bulk samplers (VDI 4320 Part 3,  
203 2017; c.f. Thimonier et al., 2019) between April 2013 and April 2015. Nitrate (NO<sub>3</sub><sup>-</sup>) in rainwater and melted  
204 snow was analyzed by ion chromatography (ICS-1600, Dionex, USA) and NH<sub>4</sub><sup>+</sup> was analyzed using a flow  
205 injection analyzer (FIAstar 5000, Foss, Denmark) with gas diffusion membrane, detection was completed with  
206 UV/VIS photometry (SN EN ISO 11732).

207

## 208 **2.6 Plant productivity**

209 All plant material (including mosses and lichens) of the monoliths was cut 2 cm above the soil surface once per  
210 year at canopy maturity. This plant removal serves as a proxy for the short, but intensive summer grazing period  
211 of the traditional management. As a result of the phenology-triggered harvests (anthesis of *Festuca rubra*), the  
212 topmost CS1 was cut on average 38 days later than the lowest CS6. Plants were dried at 60 °C, allowed to cool  
213 in a desiccator and weighed to determine dry matter yield (hereafter biomass yield).

214

## 215 **2.7 Data analyses**

216 Data were analyzed by linear mixed-effects models. First, we were interested in the overall response of biomass  
217 yield over years as affected by the treatment factors. To this aim, biomass yield was averaged across the four  
218 experimental years (2014-2017) and was modeled as function of CS (factor of 6 levels), irrigation (factor of 2  
219 levels), and N-deposition (factor of 3 levels), including all interactions. ‘Site of origin’ (6 sites) and block (36  
220 levels: 6 CS × 6 blocks) were modeled as random factors (random intercepts). Restricted maximum likelihood  
221 was used for parameter estimation. For the inference on fixed effects, the Kenward–Roger method was applied  
222 to determine the approximate denominator degrees of freedom (Kenward and Roger 1997), and the marginal  
223 and conditional  $R^2$  of the model were computed following Nakagawa and Schielzeth (2013). Differences in  
224 biomass yield between single CSs and the CS<sub>reference</sub> were tested based on the model contrasts (post-hoc  $t$ -tests,  
225 without using multiple comparisons). To receive additional insight into within year treatment effects, this very  
226 same model was also applied to data of each of the four individual years.

227 Second, to consider the time effect and the repeated structure of the data, biomass yield of all four years was  
228 modeled as function of year (factor of 4 levels), CS, irrigation, and N-deposition (factor levels as described),  
229 including all interactions. Here, random factors consisted of an identifier for monolith (216 levels) to consider  
230 the potential correlation of monoliths’ biomass yield over years (modeled as random intercept). In addition, the  
231 model included the random factor ‘site of origin’ and allowed for a separate block term at each of the four years  
232 (details as described). Residuals of all models were evaluated for normality and homoscedasticity and fulfilled  
233 assumptions of linear mixed-effects models. Finally, to gain insight into effects of thermal energy and drought  
234 on plant productivity, biomass yield was modeled as function of each DD0°C<sub>total</sub> and percent days with less soil  
235 moisture (‘dry days’) using generalized additive models (GAM). Generalized additive models had to be used as  
236 simple linear models could not appropriately handle these relationships. The GAMs included the fixed factor  
237 irrigation and a smooth term for the continuous variables DD0°C<sub>total</sub> and percent dry days, respectively, for both  
238 levels of the irrigation treatment. Model validation revealed that the assumptions of GAMs were met; more

239 information on the model specification is given in the Appendix. All data was analyzed with the statistics  
240 software R, version 4.0.2 (R Core Team 2020) and packages lme4 for linear-mixed effect models (Bates et al.,  
241 2015) and mgcv for GAMs (Wood, 2017).

## 242 3. Results

243

### 244 3.1 Climate scenario site (CS) environmental conditions

#### 245 3.1.1 Low atmospheric background N-deposition

246 Total N-deposition was  $3.3 \text{ kg N ha}^{-1} \text{ a}^{-1}$  at CS<sub>2reference</sub> and  $4.3 \text{ kg N ha}^{-1} \text{ a}^{-1}$  at the lowest CS6. The seasonal  
247 distribution showed peak deposition rates in June and July.

248

#### 249 3.1.2 Consistent temperature, precipitation and drought changes with altitude

250 The mean Apr. – Oct. temperature gradient of up to  $+3 \text{ }^\circ\text{C}$  compared to CS<sub>2reference</sub>, distributed over four  
251 altitudinal levels (CS3 – CS6), constituted the warming treatment. Conversely, temperature at the topmost CS1  
252 constituted a cooling treatment ( $\Delta \text{ temp. } -1.4 \text{ }^\circ\text{C}$ ), extending the range of temperature responses tested ( $\Delta \text{ temp.}$ ,  
253 Tab. 1). As intended, the  $\text{DD}0^\circ\text{C}_{\text{total}}$  steadily increased from CS<sub>2reference</sub> to lowest CS6. The pre-harvest period  
254 (PHP) length was fairly similar among CSs, because the early snow-melt at the lower CS was compensated by  
255 an early harvest (Tab. 1).

256 We observed a small, non-linear increase of precipitation with altitude during April – October. The recorded  
257 annual precipitation sum was somewhat larger than the sum for the growing period (Tab. 2).

258 The length of the period with less soil moisture (% dry days) doubled along the altitudinal gradient: At the two  
259 top CSs only one third of the pre-harvest period was dry, compared to two thirds of the time at the lowest site  
260 CS6 (compare Tables 1 & 2). The irrigation treatment reduced the incidence of days with  $< 40 \%$  soil moisture  
261 to 60-80 % of the non-irrigated situation (Tab. 2).

262

#### 263 3.1.3 Small transplantation effects on soil temperature and moisture

264 At the sites of origin, the mean April – October soil temperatures in the undisturbed grassland were  $8.8 \text{ }^\circ (\pm 0.3)$   
265 compared to  $8.9 \text{ }^\circ\text{C} (\pm 0.3)$  in the monoliths. At CS<sub>2reference</sub> this difference was  $9.2 \text{ }^\circ$  vs.  $9.5 \text{ }^\circ\text{C}$ . Thus, the  
266 surrounding grassland at CS<sub>2reference</sub> site was on average  $0.4 \text{ }^\circ\text{C}$  warmer than at the sites of origin, and monoliths  
267 at CS<sub>2reference</sub> were  $0.3 \text{ }^\circ\text{C}$  warmer than the undisturbed grassland surrounding the experiment. Volumetric SWC  
268 in the undisturbed grassland was 1 % lower on average compared to SWC in the monoliths at CS<sub>2reference</sub> and  
269 lowest CS6.

270

## 271 3.2. Yield

### 272 3.2.1 Insignificant transplantation effect

273 The mean annual yield was 20 % larger at CS<sub>2reference</sub> (control treatment monoliths), compared to the origins  
274 ( $162 \text{ g m}^{-2}; \pm 12.7$ ), but not significantly different ( $P = 0.19$ ; paired, two-sided *t*-test). Equally important, the  
275 difference showed no trend, as in some years the yield at CS<sub>2reference</sub> was higher, in some years it was lower  
276 compared to the sites of origin.

277

### 278 3.2.2 Strongest climate scenario site effect at intermediate CS

279 Across the four years, we found a highly significant effect of the CS on aboveground biomass yield (Tab. 3). At  
280 intermediate sites, yields increased by +43 %, +18 % and +44 % (CS3, CS4 and CS5, respectively; Tab. 4,  $P \leq$   
281 0.05 at least), related to  $+0.7$ ,  $+1.5$ , and  $+1.8 \text{ }^\circ\text{C}$  of the warming component of the respective CS (compare Table

282 1). Even at the warmest site CS6 the yield was still +12 % larger compared to the CS2<sub>reference</sub> site ( $\Delta T = +3$  °C,  
 283  $\Delta DD0^{\circ}C_{total} = 655$ ). The coldest site CS1 was not less productive than CS2<sub>reference</sub>. In the year of the overall  
 284 maximum productivity (2016), also the coldest site CS1 and the warmest site CS6 produced their respective  
 285 record yield (Tab. 4). Overall, the yields of the 24 combinations of year  $\times$  CS varied by a factor of 2.1 (yields  
 286 averaged across irrigation and N-deposition treatments). The yield response to CSs differed between years  
 287 (Appendix Tab. A2, year  $\times$  CS interaction:  $P < 0.001$ ) in that the CS effect became weaker towards the end of  
 288 the experiment (Appendix Tab. A3).

289  
 290

Variable	df <sub>num</sub>	df <sub>den</sub>	F-value	P
Climate Scenario (CS)	5	29.1	14.9	< 0.001
Irrigation	1	145.2	6.5	0.012
N	2	145.2	1.3	0.287
CS $\times$ Irrigation	5	145.2	1.1	0.352
CS $\times$ N	10	145.2	0.5	0.864
Irrigation $\times$ N	2	145.2	1.1	0.348
CS $\times$ Irrigation $\times$ N	10	145.2	1.3	0.241

df<sub>num</sub>: degrees of freedom of term; df<sub>den</sub>: degrees of freedom of error (which can be fractional in restricted maximum likelihood analysis)

291 **Table 3** Summary of analyses for the effects of climate scenario site (CS), irrigation and N deposition on  
 292 aboveground biomass yield of subalpine grassland. Data were averaged across the four experimental years (total  
 293  $n = 216$ ). *F*-tests refer to the fixed effects of the linear mixed-effects model. The marginal and conditional  $R^2$   
 294 were 0.41 and 0.50, respectively.

295  
296  
297

Year	CS <sub>2reference</sub>		Aboveground biomass yield (g m <sup>-2</sup> , means ±1SE)										
	% dry days	DD0°C <sub>total</sub>	CS1	CS <sub>2reference</sub>	CS3	CS4	CS5	CS6	CS mean				
2014	30	1353	149 <sup>ns</sup> ±8.0	170 ±11.0	238 <sup>***</sup> ±8.8	203* ±11.6	255 <sup>***</sup> ±15.2	152 <sup>ns</sup> ±10.5	194 ±5.3				
2015	38	1359	147 <sup>ns</sup> ±8.1	138 ±5.8	248 <sup>***</sup> ±12.1	171 <sup>†</sup> ±8.9	310 <sup>***</sup> ±13.6	198 <sup>***</sup> ±8.8	202 ±5.7				
2016	22	1509	230 <sup>ns</sup> ±8.7	222 ±9.1	297 <sup>***</sup> ±10.2	247 <sup>ns</sup> ±11.1	271 <sup>**</sup> ±15.3	250 <sup>†</sup> ±9.8	253 ±4.7				
2017	34	1541	152 <sup>ns</sup> ±8.5	166 ±7.8	208* ±10.0	201* ±11.7	169 <sup>ns</sup> ±9.1	176 <sup>ns</sup> ±8.3	178 ±4.0				
Mean	36	1440	170 <sup>ns</sup> ±7.1	174 ±6.9	248 <sup>***</sup> ±7.9	205* ±9.0	251 <sup>***</sup> ±11.5	194 <sup>ns</sup> ±6.9					

298 \*\*\*  $P \leq 0.001$ , \*\*  $P \leq 0.01$ , \*  $P \leq 0.05$ , †  $P \leq 0.1$ , <sup>ns</sup>  $P > 0.1$


299 **Table 4** Aboveground biomass yield (means ±1SE) per CS and year, averaged across irrigation, N-deposition treatments, and site of origin. Within each year, significance  
300 tests are against CS<sub>2reference</sub>, based on contrasts derived from linear mixed-effects models (see Table 3 and Appendix Tab. A1, for the respective model summaries). Shaded  
301 values indicate the CS with the greatest aboveground biomass yield per year and across years.

302 **3.2.3 Irrigation effect in dry years**

303 Despite a mere +7.7 % average yield increase (Fig. 1 b), irrigation turned out to be a significant factor across years  
304 (Table 3). Yet, the effect of irrigation differed between years (Appendix Tab. A2, year × irrigation interaction:  $P <$   
305 0.001), and single years analysis detected positive effects of irrigation only in 2015 (+15.8 %) and 2017 (+18.8 %)  
306 (Appendix Tab. A3). In these years, the percentage of days with  $< 40$  % soil moisture was highest.

307  
308 **3.2.4 No nitrogen deposition effect**

309 Five years of experimentally increased atmospheric nitrogen deposition (+3 and +15 kg N ha<sup>-1</sup> a<sup>-1</sup>) did not cause a  
310 significant response of biomass yield. Moreover, there was no significant interaction detected between the N-  
311 treatment and the factors CS or irrigation (Fig. 1 a; Tab. 3). Single years analysis, to test for a late response to  
312 accumulating amounts of N, revealed a marginally significant effect only in 2016 (Appendix Tab. A3).



313

314 **Figure 1 a, b** Aboveground biomass yield as a function of the altitude of CSs. Data were averaged across years;  
315 circles denote means  $\pm 1$  SE. Warming and dry days (%) increase with decreasing altitude from left to right. **a)** Yield  
316 values grouped by N-deposition treatment (0, 3 and 15 kg N ha<sup>-1</sup> a<sup>-1</sup>, in addition to 4-5 kg N background deposition).

317 **b)** Yield values grouped by irrigation treatment. Overlapping means and SEs are shifted horizontally to improve  
 318 their visibility.

319

320 **3.2.5 Yield at climate scenario sites strongly relates to changes in thermal energy and soil moisture**


321 Biomass yield at the different CSs was significantly related to thermal energy, expressed as growing  $DD0^{\circ}C_{total}$ .

322 Here, intermediate CSs (CS3, CS5) had greatest yields at intermediate values of  $DD0^{\circ}C_{total}$ , indicated by the  
 323 curvature of the predicted line in particular under irrigated conditions (Fig. 2 a, Appendix Tab. A4, smooth term for  
 324  $DD0^{\circ}C_{total}$ :  $P < 0.001$ ).

325 Similarly, biomass yield was significantly related to days with soil moisture  $< 40\%$  ('dry days') during the growing  
 326 season, with intermediate CS3 and CS5 having highest yields at around 50% of dry days under no irrigation and at  
 327 around 30% dry days under additional irrigation (Fig. 2 b, Appendix Tab. A5, smooth term for dry days:  $P <$   
 328  $0.001$ ). Under unirrigated conditions, in parallel with a doubling of dry days (from 27% at topmost CS1 to 55% at  
 329 intermediate CS5), yield consistently rose and only fell at the driest and warmest site CS6, with 73% dry days.

330

331 **Figure 2 a, b** Aboveground biomass yield at the six CS as **a)** a function of total received thermal energy  
 332 ( $DD0^{\circ}C_{total}$ ), and **b)** percent of days with dry soil ( $SWC < 40\%$ ) during the growing season (dry days %). Data were  
 333 averaged across years; circles denote means  $\pm 1$  SE per CS and irrigation treatment. The predicted line is based on a  
 334 generalized additive model using all data ( $\pm 1$  SE light grey shaded). Dark grey indicates the cross-section of the two  
 335 SE bands. Overlapping means and SEs in (a) are shifted horizontally to improve their visibility.



## 336 4 Discussion

337 We found a substantial and significant positive effect of climate scenarios, equivalent to warming of up to + 1.8 °C  
338 (Apr. – Oct. mean), on aboveground biomass of subalpine grasslands (up to +44 % yield). Contrary to expectation,  
339 additional resource supply through irrigation and N-deposition had only marginal (water) or no effects (N) on yield,  
340 respectively. Our transplanting experiment proved to be efficient in assessing several linked climate change drivers  
341 in their effect on plant growth.

342

### 343 4.1 Climate scenario temperature effects

344 The phenology-triggered harvest opened the possibility to extend the growing period in cool years and shorten the  
345 exposure to drought stress in warm years. Thus, beneficial thermal effects were maximized, while detrimental  
346 drought effects were minimized. As a consequence, we displayed the yield over a continuous x-axis of degree days  
347 between harvests ( $DD0^{\circ}C_{total}$ , Fig. 2 A). This represents the available thermal resource, associated with a particular  
348 yield, much better than mean temperatures of CS, or categorical values for CS1-CS6.

349 In cold environments, the warming is so important because the metabolic growth processes, which utilize the  
350 assimilated energy, are strongly temperature dependent, much more so than the assimilation process *per se* (Körner  
351 2003). In a meta-analysis of grassland responses to warming that included 32 sites, distinctly positive warming  
352 effects on growth were found in the colder portion of those ecosystems (Rustad et al., 2001), very similar to  
353 responses in the subalpine grassland of the current study. Interestingly, also the response size of our effects is in the  
354 same range as that reported by Rustad et al., (2001).

355 Plant growth at the intermediate climate scenarios that represented a warming of 0.7 °C, 1.5 °C and 1.8 °C (Apr.-  
356 Oct.) clearly benefitted from greater warmth. However, the increase of responses was somewhat inconsistent (CS4  
357 ca. +18 %, CS3 and CS5 both > +40 %), matching only partly our first hypothesis. The erratic response of  
358 intermediate CS4 is likely the result of an interaction of micro-topography effects on climate that were not detected  
359 by our meteorological measurements, cockchafer infestation (*Melolontha melolontha*: bug whose larvae fed on  
360 roots), or the occurrence of mast years in some species at that CS. In the extreme treatment at lowest CS6 (+3 °C  
361 Apr.-Oct., +2.4 °C annual mean) the positive response to warming finally ceased to increase, but yield was still  
362 somewhat larger than at CS2<sub>reference</sub>. This comparatively low growth response suggests that the water supply at CS6  
363 had already reached critically low levels; yet, the larger thermal energy resource must have partly compensated for  
364 the radically smaller soil water resource, leading also to some growth benefit at CS6 (compare Figure 2 a & b).

365 Despite substantial cooling at topmost CS1, coinciding with a temperature decline of -1.4 °C, the mean yields for  
366 CS1 and CS2<sub>reference</sub> were very similar (Tab. 4, Fig. 1). This is indicative of a plant community that is well cold-  
367 adapted. Indeed, local historical records from the Swiss Federal Office for Meteorology (MeteoSwiss) show that  
368 only 100 years ago the local April-October mean air temperature was 1.4-1.5 °C lower than today (30 a running  
369 mean, courtesy P. Calanca using MeteoSwiss data from Segl-Maria site at 1804 m a.s.l.). In effect, the cooling  
370 upward-transplantation represented a climatic time travel of 100 years into the past, and the similar yield responses  
371 between CS1 and CS2<sub>reference</sub> indicate that subalpine grassland productivity may not have changed much during the  
372 past century. Moreover, the dramatic temperature dynamics during the past 12,000 years of the Holocene



373 interglacial suggest that temperature adaptations in modern plant genotypes may actually match not only today's  
374 weather, but also warmer and cooler climate conditions. From this perspective, and with respect to the genetic  
375 memory of plants, the undiminished productivity at topmost CS1 illustrates that assumed 'control' temperatures in  
376 warming experiments only represent the most recent point of an extremely dynamic climatic history.

377

#### 378 **4.2 Climate scenario soil moisture effects**

379 The differences in soil moisture content that resulted naturally from 24 different climatic situations (6 CS/altitude  
380 levels  $\times$  4 years) created a hump-shaped response curve of yield over drought (Fig. 2 b). This does imply that, with  
381 decreasing altitude and increasing warmth, productivity rose despite more days with soil moisture  $< 40\%$ .

382 The importance of soil moisture for plant growth has been shown predominantly in much drier grasslands, e.g., in  
383 warmer prairie (Xu et al., 2013) or cold alpine grassland (Wang et al., 2013), where release from drought stress  
384 benefitted growth. For example, along a temperature and altitude gradient in semiarid Tibetan alpine grassland,  
385 productivity increased with altitude due to reduced drought stress, but despite decreasing temperatures. Only after an  
386 800 m rise in altitude, productivity eventually became smaller, and further reduced drought stress did not constitute  
387 a further advantage on plant growth (Wang et al., 2016).

388 In our experiment, soil moisture values and its proxies integrate information on moisture *and* temperature. Thus, the  
389 two-dimensional growth response curve along the altitudinal gradient, peaking at the least detrimental situation  
390 between moisture limitation and thermal limitation (Fig. 2 b), is analogous to the three-dimensional response surface  
391 found in the Jasper Ridge experiment (Zhu et al., 2016). Unfortunately, our experiment did not produce a sufficient  
392 number of data points for a 3-D presentation. Based on these results, we infer that a joint evaluation of soil moisture  
393 and temperature is mandatory to assess reliably warming effects of climate change on plant growth in the subalpine  
394 environment.

395

#### 396 **4.3 Irrigation treatment**

397 We had assumed that increased SWC would mitigate detrimental effects of excessive warming. Surprisingly  
398 however, the overall irrigation effect on yield was not very substantial, despite large differences in the percentages  
399 of days with soil moisture  $< 40\%$  during the growing season (Table 2, Fig. 2 b). Moreover, the positive responses  
400 did not increase consistently with warmth, but were strongest at the intermediate CS3 and CS4 (Fig. 1 b). Analyses  
401 of individual years showed that the two significant responses of annual yield to irrigation coincided with the two  
402 driest years. This evidence suggests that maximum mitigation of (low) temperature limitation requires simultaneous  
403 release of water limitation, while at the same time the amount of water applied in our study was insufficient to  
404 compensate for increased evapotranspiration at CS5 and the warmest site CS6.

405

#### 406 **4.4 N-deposition treatment**

407 We hypothesized a generally positive effect of N-deposition on plant growth, but found no significant overall effect  
408 of N-deposition on yield after four years and only a marginal effect in one year. Historically, the responsiveness of  
409 (sub-)alpine vegetation to improved nutrient supply was considered to be restricted due to an overriding effect of

410 thermal energy limitation. Yet, studies with very high rates of N application (40-100 kg N ha<sup>-1</sup> a<sup>-1</sup>; Körner et al.,  
411 1997; Heer and Körner 2002) showed substantial yield responses, also at alpine sites. Low N-dose responses of total  
412 plant yield may require N-accumulation over years or a compound interest effect in plant biomass. For example,  
413 only in the seventh treatment year a strong, +31 % total yield growth response to 5 kg N ha<sup>-1</sup> a<sup>-1</sup> was reported by  
414 Volk et al., (2014) from subalpine grassland.  
415 Low dose experiments (5-30 kg N ha<sup>-1</sup> a<sup>-1</sup>), however, can induce a species composition change (Bowman et al.,  
416 2012), indicating a growth benefit for some species at the expense of others. Yet, such single species responses may  
417 be only transient: a strong *Carex* species response to as little as 5 kg N ha<sup>-1</sup> a<sup>-1</sup> in similar subalpine vegetation was  
418 recently found to cease after five years (Bassin et al., 2009 and 2013). Taken together, we conclude that the cold-  
419 adapted, mature and low productivity grassland either responds with a >4 year time lag, or that the N-deposition  
420 treatment was below the critical load for aboveground biomass responses.

421

#### 422 **4.5 Transplantation**

423 The turf monoliths at CS2<sub>reference</sub> were only slightly warmer and moister compared to the sites of origin, suggesting a  
424 low transplantation impact (we have found no transplantation effect data from other experiments to compare with).  
425 However, within the experimental site similar temperature increases between CS2<sub>reference</sub> and CS3 caused a much  
426 larger productivity increase (+43 %). We reason that this incongruence can be explained by the difference in melt-  
427 out time, which was on average only 3 days earlier at CS2<sub>reference</sub> (julian day 118) than at the sites of origin, but 21  
428 days earlier at CS3 than at CS2<sub>reference</sub>. We thus assume that the substantially earlier start of the growing season  
429 caused the stronger growth response, despite a similar temperature change. This effect, induced by the  
430 transplantation of the grassland monoliths along the altitudinal gradient, demonstrates the importance of integrating  
431 multiple drivers in climate change experiments to allow for a multi-factor driven plant response.  
432 In our study, the effect of altitude on photosynthesis substrate limitation was considered negligible, compared to the  
433 climate effects. The assimilation conditions of alpine plants have been the subject of investigation for decades. Since  
434 the theoretical considerations of Gale (1972) and the field studies by Körner and Diemer (1987) and Körner et al.,  
435 (1988), a predominant ‘altitude-tolerance’ of photosynthesis is widely accepted. Relevant environmental parameters  
436 that change with altitude (temperature, CO<sub>2</sub> and O<sub>2</sub> partial pressure, vapor-pressure deficit and photosynthetic  
437 photon flux density) have antagonistic effects on assimilation efficiency (see Wang et al., (2016) for a recent  
438 discussion on the topic).

439 **5 Conclusions**

440 Despite dwindling soil water content, the subalpine grassland growth increased to up to +1.8 °C warming during the  
441 growing period (corresponding to +1.3 °C annual mean), compared to present temperatures. Even at the maximum  
442 warming (corresponding to +2.4 °C annual mean) the yield was larger than at the reference site. At the same time -  
443 1.4 °C cooling during the growing period (corresponding to -1.7 °C annual mean) did not reduce plant growth. This  
444 implies that subalpine grassland productivity has likely not increased during the past century warming, but, despite  
445 growing soil moisture deficits, will do so with continued warming in the near future.

446 **Author contribution**

447 MV and SB designed the experiment, MV, ALW and SB conducted field work. MV and MS analyzed the data. MV  
448 led the writing of the manuscript, with significant contribution from MS. All authors contributed critically to the  
449 drafts and gave final approval for publication.

450

451 **Data availability**

452 The data analyzed for the current study are available in the Dryad Digital Repository at ...

453

454 **Competing interests**

455 The authors declare that they have no conflict of interest.

456

457 **Acknowledgements**

458 We received essential financial support through the Federal Office for the Environment (contract No. 00.5100.PZ /  
459 R442-1499). The Federal Office for Meteorology (MeteoSwiss) is kindly acknowledged for providing access to  
460 meteorological data. We thank Pierluigi Calanca for handling these data. The Gemeinde Ardez and Alpmeister  
461 Claudio Franziscus generously allowed us to work on the Allmend. N-concentration analyses courtesy of  
462 Forschungsstelle für Umweltbeobachtung (FUB-AG, Rapperswil, Switzerland). We are grateful to Robin Giger for  
463 his untiring support in the field and the lab, and to the scientific site manager Andreas Gauer, who was in charge of  
464 the field sites.

465 **References**

- 466 Bassin, S., Volk, M., Suter, M., Buchmann, N., Fuhrer, J.: Nitrogen deposition but not ozone affects productivity  
467 and community composition of subalpine grassland after 3 yr of treatment. *New Phytologist* 175, 3, 523-534,  
468 2007.
- 469 Bassin, S., Werner, R. A., Sörgel, K., Volk, M., Buchmann, N., Fuhrer, J.: Effects of combined ozone and nitrogen  
470 deposition on the in situ properties of eleven key plant species of a subalpine pasture. *Oecologia* 158, 4, 747-756,  
471 2009.
- 472 Bassin, S., Volk, M., Fuhrer, J.: Species composition of subalpine grassland is sensitive to nitrogen deposition, but  
473 not to ozone, after seven years of treatment. *Ecosystems* 16, 6, 1105-1117, 2013.
- 474 Bates, D., Maechler, M., Bolker, B., Walker, S.: lme4: Linear mixed-effects models using Eigen and S4. Version  
475 1.1-10. <https://CRAN.R-project.org/package=lme4>, 2015.
- 476 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S.,  
477 Davidson, E., Dentener, F., Emmett, B., Erisman, J. W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L., De Vries,  
478 W.: Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecological*  
479 *applications* 20, 1, 30-59, 2010.
- 480 Bowman, W. D., Gartner, J. R., Holland, K., Wiedermann, M.: Nitrogen Critical Loads For Alpine Vegetation And  
481 Terrestrial Ecosystem Response: Are We There Yet? *Ecological Applications* 16, 1183-1193, 2006.
- 482 Bowman, W. D., Murgel, J., Blett, T., Porter, E.: Nitrogen critical loads for alpine vegetation and soils in Rocky  
483 Mountain National Park. *Journal of Environmental Management* 103, 165-171, 2012.
- 484 Core Writing Team, Pachauri, R. K., Meyer, L. A. editors: IPCC, 2014: Climate change 2014: Synthesis Report.  
485 Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on  
486 Climate Change. IPCC, Geneva, Switzerland. 151p, 2014.
- 487 Dieleman, W. I., Vicca, S., Dijkstra, F. A., Hagedorn, F., Hovenden, M.J., Larsen, K.S., ... & King, J.: Simple  
488 additive effects are rare: a quantitative review of plant biomass and soil process responses to combined  
489 manipulations of CO<sub>2</sub> and temperature. *Global Change Biology* 18, 9, 2681-2693, 2012.
- 490 Dukes, J. S., Chiariello, N. R., Cleland, E. E., Moore, L. A., Shaw, M. R., Thayer, S., Tobeck, T., Mooney, H. A.,  
491 Field, C. B.: Responses of grassland production to single and multiple global environmental changes. *PLoS*  
492 *Biology* 3, 10, e319, 2005.
- 493 Heer, C. and Körner, C.: High elevation pioneer plants are sensitive to mineral nutrient addition. *Basic and Applied*  
494 *Ecology*, 3, 1, 39-47, 2002.
- 495 Gale, J.: Availability of carbon dioxide for photosynthesis at high altitudes: theoretical considerations. *Ecology*, 53,  
496 3, 494-497, 1972.
- 497 Inauen, N., Körner, C., Hiltbrunner, E.: No growth stimulation by CO<sub>2</sub> enrichment in alpine glacier forefield plants.  
498 *Global Change Biology*, 18, 3, 985-999, 2012.
- 499 Kenward, M. G., Roger, J. H.: Small sample inference for fixed effects from restricted maximum likelihood.  
500 *Biometrics*, 53, 3, 983-997, 1997.

501 Körner, C. and Diemer, M.: In situ photosynthetic responses to light, temperature and carbon dioxide in herbaceous  
502 plants from low and high altitude. *Functional Ecology*, 179–194, 1987.

503 Körner, C., Farquhar, G. D., Roksandic, Z.: A global survey of carbon isotope discrimination in plants from high  
504 altitude. *Oecologia*, 74, 623-632, 1988.

505 Körner, C., Diemer, M., Schächli, B., Niklaus, P., Arnone III J.: The responses of alpine grassland to four seasons of  
506 CO<sub>2</sub> enrichment: a synthesis. *Acta Oecologica* 18, 3, 165-175, 1997.

507 Körner, C.: *Alpine plant life: functional plant ecology of high mountain ecosystems*. Springer Science & Business  
508 Media. 344p., 2003.

509 Leuzinger, S., Luo, Y., Beier, C., Dieleman, W., Vicca, S., Körner, C.: Do global change experiments overestimate  
510 impacts on terrestrial ecosystems? *Trends in ecology and evolution* 26, 5, 236-241, 2011.

511 Liu, H., Mi, Z., Lin, L., Wang, Y., Zhang, Z., Zhang, F., ... & Zhao, X.: Shifting plant species composition in  
512 response to climate change stabilizes grassland primary production. *Proceedings of the National Academy of*  
513 *Sciences* 115, 16, 4051-4056, 2018.

514 Nakagawa, S. and Schielzeth, H.: A general and simple method for obtaining R<sup>2</sup> from generalized linear mixed-  
515 effects models. *Methods in ecology and evolution*, 4, 2, 133-142, 2013.

516 Phoenix, G. K., Emmett, B. A., Britton, A. J., Caporn, S. J.M., Dise, N. B., Helliwell, R., Jones, L., Leake, J. R.,  
517 Leith, I. D., Sheppard, L. J., Sowerby, A., Pilkington, M. G., Rowe, E. C., Ashmore, M. R., Power, S. A.:  
518 Impacts of atmospheric nitrogen deposition: responses of multiple plant and soil parameters across contrasting  
519 ecosystems in long-term field experiments. *Global Change Biology*, 18, 1197–1215, 2012.

520 R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing,  
521 Vienna, Austria, 2020. <http://www.R-project.org>.

522 Rihm, B. and Kurz, D.: Deposition and critical loads of nitrogen in Switzerland. In *Acid rain 2000*. pp. 1223-1228,  
523 Springer, Dordrecht, 2001.

524 Rihm, B. and Achermann, B. Critical Loads of Nitrogen and their Exceedances. Swiss contribution to the effects-  
525 oriented work under the Convention on Long-range Transboundary Air Pollution (UNECE). Federal Office for  
526 the Environment, Bern. Environmental studies no. 1642, 78p., 2016.

527 Rustad, L. E., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A. ... & Gurevitch, J.: A meta-analysis of  
528 the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental  
529 ecosystem warming. *Oecologia* 126, 4, 543-562, 2001.

530 Rustad, L. E.: The response of terrestrial ecosystems to global climate change: towards an integrated approach.  
531 *Science of the total environment* 404, 2-3, 222-235, 2008.

532 Thimonier, A., Kosonen, Z., Braun, S., Rihm, B., Schleppei, P., Schmitt, M., ... & Thöni, L.: Total deposition of  
533 nitrogen in Swiss forests: Comparison of assessment methods and evaluation of changes over two decades.  
534 *Atmospheric Environment* 198, 335-350, 2019.

535 Van Der Wal, R. and Stien, A.: High-arctic plants like it hot: A long-term investigation of between-year variability  
536 in plant biomass. *Ecology* 95, 12, 3414-3427, 2014.

537 Vinther, B. M., Buchardt, S. L., Clausen, H. B., Dahl-Jensen, D., Johnsen, S. J., Fisher, D. A. ... & Blunier, T.:  
538 Holocene thinning of the Greenland ice sheet. *Nature* 461, 7262, 385, 2009.

539 Vitousek, P. M., Aber, J., Howarth, R. W., Likens, G. E., Matson, P. A., Schindler, D. W., Schlesinger, W. H.,  
540 Tilman, G. D.: Human alteration of the global nitrogen cycle: causes and consequences. *Ecological Applications*  
541 7, 737–750, 1997.

542 Volk, M., Wolff, V., Bassin, S., Ammann, C., Fuhrer, J.: High tolerance of subalpine grassland to long-term ozone  
543 exposure is independent of N input and climatic drivers. *Environmental Pollution* 189, 161-168, 2014.

544 Volk, M., Enderle, J., Bassin, S.: Subalpine grassland carbon balance during 7 years of increased atmospheric N  
545 deposition. *Biogeosciences* 13, 12, 3807-3817, 2016.

546 Wang, Z., Luo, T., Li, R., Tang, Y., Du, M.: Causes for the unimodal pattern of biomass and productivity in alpine  
547 grasslands along a large altitudinal gradient in semi-arid regions. *Journal of Vegetation Science* 24, 1, 189-201,  
548 2013.

549 Wang, H., Prentice, I. C., Davis, T. W., Keenan, T. F., Wright, I. J., Peng, C.: Photosynthetic responses to altitude:  
550 an explanation based on optimality principles. *New Phytologist* 213, 3, 976-982, 2016.

551 Wood, S. N.: *Generalized Additive Models: An Introduction with R*. 2<sup>nd</sup> edition. Chapman and Hall/CRC. London,  
552 2017.

553 Wüst-Galley, C., Volk, M., Bassin, S.: Interaction of climate change and nitrogen deposition on subalpine pastures.  
554 (in revision) 2020.

555 Xu, X., Sherry, R. A., Niu, S., Li, D., Luo, Y.: Net primary productivity and rain-use efficiency as affected by  
556 warming, altered precipitation, and clipping in a mixed-grass prairie. *Global Change Biology* 19, 9, 2753-2764,  
557 2013.

558 Zhu, K., Chiariello, N. R., Tobeck, T., Fukami, T., Field, C. B.: Nonlinear, interacting responses to climate limit  
559 grassland production under global change. *Proceedings of the National Academy of Sciences* 113, 38, 10589-  
560 10594, 2016.

561 **Appendix for**

562 **Sub-alpine grassland productivity increased with warmer and**  
563 **drier conditions, but not with higher N-deposition, in an**  
564 **altitudinal transplantation experiment**

565

566 Matthias Volk<sup>1</sup>, Matthias Suter<sup>2</sup>, Anne-Lena Wahl<sup>1</sup>, Seraina Bassin<sup>1,3</sup>

567 <sup>1</sup>Climate and Agriculture, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

568 <sup>2</sup>Forage Production and Grassland Systems, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland

569 <sup>3</sup>Pädagogische Hochschule Schaffhausen, Ebnetstrasse 80, 8200 Schaffhausen, Switzerland

570

571 Corresponding author: Matthias Volk (matthias.volk@agroscope.admin.ch)



572 **Appendix Tables**

573 **Table A1** Schematic layout of monolith arrangement at each CS of the AlpGrass experimental site. At each CS, six  
 574 monoliths from each of six sites of origin were transplanted, resulting in 36 monoliths. Two irrigation and three N-  
 575 deposition treatments were set up in a factorial design, resulting in six irrigation × N treatment combinations, which  
 576 were assigned to each of the six monoliths per site of origin. The six irrigation × N treatment combinations were  
 577 arranged in a randomized complete block design of six blocks. Regarding sites of origin, the monoliths were assigned  
 578 to the six blocks in a restricted randomization, so that an equal distribution of sites of origin to all blocks was ensured.  
 579 It follows that the six monoliths from each site of origin received all irrigation × N treatment combinations and were  
 580 evenly distributed on the site. Displayed is a possible randomization of irrigation and N treatments per block; at each  
 581 CS separate randomizations were performed.

Block 1			Block 2			Block 3		
W0.N15	W1.N0	W0.N3	W0.N3	W0.N0	W1.N15	W1.N15	W0.N0	W1.N0
W0.N0	W1.N3	W1.N15	W1.N3	W0.N15	W1.N0	W1.N3	W0.N15	W0.N3

W1.N15	W1.N0	W0.N0	W1.N3	W0.N3	W1.N0	W0.N0	W1.N0	W0.N15
W0.N3	W0.N15	W1.N3	W0.N0	W1.N15	W0.N15	W1.N3	W1.N15	W0.N3
Block 4			Block 5			Block 6		

582 W0: no additional water (ambient precipitation only), W1: additional water during  
 583 growing period; N0: no N fertilizer, N3: 3 kg N ha<sup>-1</sup> a<sup>-1</sup>, N15: 15 kg N ha<sup>-1</sup> a<sup>-1</sup>

584 **Table A2** Summary of analyses for the effects of climate scenario (CS), irrigation, and N deposition on  
 585 aboveground biomass yield of subalpine grassland over four experimental years. *F*-tests refer to the fixed  
 586 effects of a linear mixed-effects model that included all four years for a repeated measures analysis. The  
 587 marginal and conditional  $R^2$  were 0.68 and 0.80, respectively.

588

Variable	df <sub>num</sub>	df <sub>den</sub>	<i>F</i> -value	<i>P</i>
Year	3	45.5	66.2	< 0.001
Climate Scenario (CS)	5	198.0	18.3	< 0.001
Irrigation	1	166.6	6.2	0.014
N	2	166.6	1.2	0.304
Year × CS	15	63.0	9.6	< 0.001
Year × Irrigation	3	450.5	13.6	< 0.001
Year × N	6	450.5	0.9	0.492
CS × Irrigation	5	166.6	1.1	0.380
CS × N	10	166.6	0.5	0.882
Irrigation × N	2	166.6	1.0	0.365
Year × CS × Irrigation	15	450.5	2.9	< 0.001
Year × CS × N	30	450.5	0.8	0.749
Year × Irrigation × N	6	450.5	1.4	0.199
CS × Irrigation × N	10	166.6	1.2	0.275
Year × CS × Irrigation × N	30	450.5	1.4	0.066

589 df<sub>num</sub>: degrees of freedom of term; df<sub>den</sub>: degrees of freedom of error (which can be fractional  
 590 in restricted maximum likelihood analysis)

591 **Table A3** Summary of analyses for the effects of climate scenario (CS), irrigation, and N deposition on aboveground biomass yield of subalpine grassland at each of  
592 four experimental years (2014 – 2017). *F*-tests refer to the fixed effects of a linear mixed-effects model to each of the four years.  
593

Variable	df <sub>num</sub>	2014			2015			2016			2017		
		df <sub>den</sub>	<i>F</i> -value	<i>P</i>	df <sub>den</sub>	<i>F</i> -value	<i>P</i>	df <sub>den</sub>	<i>F</i> -value	<i>P</i>	df <sub>den</sub>	<i>F</i> -value	<i>P</i>
Climate Scenario (CS)	5	28.9	17.2	< 0.001	29.5	24.9	< 0.001	29.3	4.5	0.004	29.4	4.0	0.006
Irrigation	1	145.2	1.5	0.224	145.1	21.6	< 0.001	145.3	1.1	0.290	145.4	19.2	< 0.001
N	2	145.2	0.7	0.481	145.1	0.5	0.610	145.3	2.6	0.078	145.4	0.3	0.728
CS × Irrigation	5	145.2	2.3	0.048	145.1	2.0	0.080	145.3	1.8	0.126	145.4	0.8	0.563
CS × N	10	145.2	0.5	0.912	145.1	0.7	0.751	145.3	0.9	0.531	145.4	0.5	0.896
Irrigation × N	2	145.2	1.9	0.151	145.1	0.8	0.448	145.3	0.7	0.509	145.4	1.2	0.290
CS × Irrigation × N	10	145.2	1.5	0.157	145.1	1.0	0.429	145.3	1.5	0.157	145.4	1.3	0.226

594 df<sub>num</sub>: degrees of freedom of term; df<sub>den</sub>: degrees of freedom of error (which can be fractional in restricted maximum likelihood analysis)  
595

596 **Table A4** Summary of analyses for the effects of total received thermal energy ( $DD0^{\circ}C_{total}$ ) on aboveground  
 597 biomass yield of subalpine grassland under two irrigation treatments. Data were averaged across the four  
 598 experimental years (total  $n = 216$ ).  $F$ -values and approximate  $P$ -values refer to a generalized additive model that  
 599 used a smooth term for each irrigation treatment.

600

Parametric terms	df	$F$ -value	$P$
Irrigation	2	1613.0	< 0.001
Smooth terms	edf	$F$ -value	$P$
s( $DD0^{\circ}C_{total}$ ) – No irrigation	1.72	7.7	< 0.001
s( $DD0^{\circ}C_{total}$ ) – Additional irrigation	2.34	10.2	< 0.001

601 df: degrees of freedom; edf: effective degrees of freedom (which can be fractional  
 602 in smooth terms of generalized additive models)

603 s: smoothing function applied on term

604

605

606

607

608

609 **Table A5** Summary of analyses for the effects of percent days with soil moisture < 40% during the growing  
 610 season (dry days %) on aboveground biomass yield of subalpine grassland under two irrigation treatments. Data  
 611 were averaged across the four experimental years (total  $n = 216$ ).  $F$ -values and approximate  $P$ -values refer to a  
 612 generalized additive model that used a smooth term for each irrigation treatment.

613

Parametric terms	df	$F$ -value	$P$
Irrigation	2	402.9	< 0.001
Smooth terms	edf	$F$ -value	$P$
s(Dry days %) – No irrigation	2.55	11.3	< 0.001
s(Dry days %) – Additional irrigation	2.59	8.1	< 0.001

614 df: degrees of freedom; edf: effective degrees of freedom (which can be fractional  
 615 in smooth terms of generalized additive models)

616 s: smoothing function applied on term

617 **Appendix Photographs**

618 **Photograph A1** Monolith produced at site of origin “Alp Nova” (46.72786°N, 9.72609°E) in June 2012. After  
619 monoliths were excavated in the close surroundings, they were fitted tightly into plastic containers. A total of  
620 216 monoliths was produced at six such sites of origin, for later use at the altitudinal transplantation site of the  
621 AlpGrass Experiment.



622

623

624 **Photograph A2** Topmost climate scenario site CS1 (2360 m, 46.79859°N, 10.17840°E). Monoliths are in the  
625 right part of the fenced area. Monolith arrangement in two double rows of nine allows easy access and equal  
626 distribution of edge-effects.



627 **Photograph A3** Control climate scenario site CS2<sub>reference</sub> (2170 m, 46.79264°N, 10.17714°E). Along the  
628 altitudinal transplantation gradient this CS is representative of the sites of origin, because they share the same  
629 altitude.



630

631

632 **Photograph A4** Lowest climate scenario site CS6 (1680 m, 46.77818°N, 10.17143°E). Due to its low altitude  
633 this CS is the warmest and driest site along the altitudinal transplantation gradient.



634 **Appendix R codes**

```
635 #####
636 #Linear mixed-effects model to analyze effects of the climate scenario treatment (CS), irrigation
637 and N deposition (N_Treat) on aboveground biomass yield.
638
639 #Package to load
640 library(lme4)
641
642 #####
643 #Reading in the data
644 d.data <- read.table("C:/Volk_etal_2021_AlpsGrass.csv", header=TRUE, sep=";")
645
646 #####
647 #Define factors
648 d.data$CS <- as.factor(d.data$CS)
649 d.data$Irrigation <- as.factor(d.data$Irrigation)
650 d.data$N_Treat <- as.factor(d.data$N_Treat)
651 d.data$Origin <- as.factor(d.data$Origin)
652 d.data$Block <- as.factor(d.data$Block)
653
654
655 #####
656 #Full model, including all interactions, as described in the first paragraph of the 'Data analyses'
657 section
658
659 Model.A <- lmer(DM ~ CS + Irrigation + N_Treat +
660               CS:Irrigation + CS:N_Treat + Irrigation:N_Treat + CS:Irrigation:N_Treat +
661               (1 | Origin) + (1 | Block), REML=TRUE, data=d.data)
662
663 #####
664 #The model summary, given in Table 3, is received by
665
666 library(lmerTest)
667 anova(Model.A, ddf="Kenward-Roger", type=1)
668
669
670 #####
671 #Contrasts to test for differences in biomass yield between single CSs and the CSreference (across
672 irrigation and the N treatments)
673
674 summary( update(Model.A, contrasts=list(CS=contr.treatment(levels(d.data$CS),base=2),
675 Irrigation="contr.sum", N_Treat="contr.sum")), ddf="Kenward-Roger")
676
677
678 #####
679 #This very same model and contrast code was applied to data of each individual year.
```

```

680 #####
681 Generalized additive models to test for the effects of thermal energy (DD0Ctot) and percent days
682 with less soil moisture (PercDryDays) on aboveground biomass yield.
683
684 #Package to load
685 library(mgcv)
686
687
688 #####
689 #GAM for the effect of thermal energy on yield
690
691 Model.B <- gam(DM ~ -1 + Irrigation + s(DD0Ctot, by=Irrigation), gamma=3.6,
692             knots=list(DD0Ctot=rep(seq(from=min(d.data$DD0Ctot)+100,
693             to=max(d.data$DD0Ctot)-100, length.out=12), each=18)),
694             method="REML", data=d.data)
695
696 #####
697 #The model summary, given in Table A4, is received by
698
699 anova(Model.B)
700
701
702 #####
703 #GAM for the effect of percent dry days on yield
704
705 Model.C <- gam(DM ~ -1 + Irrigation + s(PercDryDays, by=Irrigation), gamma=1.7,
706             knots=list(PercDryDays=rep(seq(from=min(d.data$PercDryDays)+5,
707             to=max(d.data$PercDryDays)-5, length.out=12), each=18)),
708             method="REML", data=d.data)
709
710 #####
711 #The model summary, given in Table A5, is received by
712
713 anova(Model.C)
714
715
716 #####
717 Note: The fitted lines in Figure 2a) & b) are based on the predicted values from Model.B and
718 Model.C, respectively.

```