

Radium-228-derived ocean mixing and trace element inputs in the South Atlantic

Yu-Te Hsieh¹, Walter Geibert², E. Malcolm S. Woodward³, Neil J. Wyatt⁴, Maeve C. Lohan⁴, Eric P. Achterberg^{4,5}, Gideon M. Henderson¹

¹Department of Earth Sciences, University of Oxford, UK
²Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany
³Plymouth Marine Laboratory, Plymouth, UK
⁴Ocean and Earth Sciences, National Oceanography Centre, Southampton, UK
⁵GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

Correspondence to: Yu-Te Hsieh (yu-te.hsieh@earth.ox.ac.uk)

Abstract. Trace elements play important roles as micronutrients in modulating marine productivity in the global ocean. The South Atlantic around 40°S is a prominent region of high productivity and a transition zone between the nitrate-depleted Subtropical Gyre and the iron-limited Southern Ocean. However, the sources and fluxes of trace elements to this region remain unclear. In this study, the distribution of the naturally occurring radioisotope ²²⁸Ra in the water column of the South Atlantic (Cape Basin and Argentine Basin) has been investigated along a 40°S zonal transect to estimate ocean mixing and trace element supply to the surface ocean. Ra-228 profiles have been used to determine the horizontal and vertical mixing rates in the near-surface open ocean. In the Argentine Basin, horizontal mixing from the continental shelf to the open ocean shows an eddy diffusion of $K_x = 1.8 \pm 1.4$ ($10^6 \text{ cm}^2 \text{ s}^{-1}$) and an integrated advection velocity $w = 0.6 \pm 0.3 \text{ cm s}^{-1}$. In the Cape Basin, horizontal mixing is $K_x = 2.7 \pm 0.8$ ($10^7 \text{ cm}^2 \text{ s}^{-1}$) and vertical mixing $K_z = 1.0 - 1.7 \text{ cm}^2 \text{ s}^{-1}$ in the upper 600 m layer. Three different approaches (²²⁸Ra-diffusion, ²²⁸Ra-advection and ²²⁸Ra/TE-ratio) have been applied to estimate the dissolved trace-element fluxes from shelf to open ocean. These approaches bracket the possible range of off-shelf fluxes from the Argentine margin to be: $4 - 21$ ($\times 10^3$) nmol Co m⁻² d⁻¹, $8 - 19$ ($\times 10^4$) nmol Fe m⁻² d⁻¹ and $2.7 - 6.3$ ($\times 10^4$) nmol Zn m⁻² d⁻¹. Off-shelf fluxes from the Cape margin are: $4.3 - 6.2$ ($\times 10^3$) nmol Co m⁻² d⁻¹, $1.2 - 3.1$ ($\times 10^4$) nmol Fe m⁻² d⁻¹ and $0.9 - 1.2$ ($\times 10^4$) nmol Zn m⁻² d⁻¹. On average, at 40°S in the Atlantic, vertical mixing supplies $0.1 - 1.2$ nmol Co m⁻² d⁻¹, $6 - 9$ nmol Fe m⁻² d⁻¹, and $5 - 7$ nmol Zn m⁻² d⁻¹ to the euphotic zone. Compared with atmospheric dust and continental shelf inputs, vertical mixing is a more important source for supplying dissolved trace elements to the surface 40°S Atlantic. It is insufficient, however, to provide the trace elements removed by biological uptake, particularly for Fe. Other inputs (e.g. particulate, or from winter deep-mixing) are required to balance the trace element budgets in this region.

Deleted: 7

Deleted: 5

Deleted: 3.8

Deleted: 2

Deleted: 7.9

Deleted: 20

Deleted: 5

Deleted: 4

Deleted: 3.6

Deleted: 11

Deleted: 13

Deleted: 16

1 Introduction

Trace elements (TEs) play important roles as micronutrients for marine productivity in the surface ocean (Morel and Price, 2003; Lohan and Tagliabue, 2018). For example, iron, zinc and cobalt are known to be essential micronutrients for the cellular metabolic enzymes in marine phytoplankton and hence they co-limit primary productivity in some ocean regions. The South Subtropical Convergence (SSTC) in the South Atlantic, near 40°S (Fig. 1), is a prominent high productivity region (0.2-0.3 mg chlorophyll a m⁻³, Longhurst, 2007) and a transition zone between the nitrate-depleted Subtropical Gyre and the iron-limited Southern Ocean, creating one of the most dynamic nutrient environments in the global oceans (Moore et al., 2004). However, the trace element sources and fluxes that fuel this region remain poorly constrained. Modeling and experimental studies have both suggested that this region is iron limited or co-limited (Moore et al., 2004; Browning et al., 2014; 2017). It also has the lowest reported dissolved zinc concentrations in the global oceans (Wyatt et al., 2014), and the replacement for zinc by cobalt is crucial for phytoplankton, practically in low zinc regions (Price and Morel, 1990). Thus, knowing the sources and fluxes of iron, zinc and cobalt can improve our understanding of the limiting factors for productivity in this highly productive region.

Oceanic mixing and advection facilitate the transport of nutrients to the euphotic zone (Oschlies, 2002). The distribution of TEs in the surface ocean is primarily controlled by the inputs from the continental shelves (i.e. rivers, submarine groundwater discharge (SGD) and sediments), deep ocean waters (regeneration, continental slopes, and hydrothermal vents) and aeolian inputs, with these mediated by lateral and vertical mixing (diffusive/turbulent mixing), advection, particle scavenging and biological uptake. In particular, deep winter mixing has been shown to be an important mechanism bringing TEs from below the mixed layer to the surface ocean (Tagliabue et al., 2014; Achterberg et al., 2018; 2020; Rigby et al., 2020). Geochemical tracers for ocean mixing can therefore be used to indirectly estimate TE inputs and outputs in the upper ocean, e.g. tritium-³He (Jenkins, 1988; Schlitzer, 2016) and radium isotopes-²²⁸Ra (Cai et al., 2002; Ku et al., 1995; Nozaki and Yamamoto, 2001; Sarmiento et al., 1990; Moore, 2000; Charette et al., 2007; Sanial et al. 2018).

The four naturally occurring radium isotopes cover a wide range of half-lives (²²⁶Ra, $T_{1/2}$ = 1600 years; ²²⁸Ra, $T_{1/2}$ = 5.75 years; ²²³Ra, $T_{1/2}$ = 11.4 days; ²²⁴Ra, $T_{1/2}$ = 3.66 days), which enables us to study oceanic processes at different time scales. Ra-228 is continuously produced through the decay of ²³²Th in shelf sediments, released into seawater, and then transported into the surface open ocean by mixing or advection. The half-life of ²²⁸Ra is much shorter than the estimated Ra residence time by removal of ~500 years (Moore and Dymond, 1991). The distribution of ²²⁸Ra in the ocean is therefore mainly controlled by ocean transport and radioactive decay, and can be used to estimate lateral mixing from the coastal shelf or continental slope to the open ocean (Kaufman et al., 1973; Knauss et al., 1978; Yamada and Nozaki, 1986; Sanial et al. 2018). Subsequent downward mixing from the surface can also be used to assess vertical mixing in the upper water column (Charette et al., 2007; Sarmiento et al., 1976; van Beek et al., 2008). Radium-228 has also been used as a conservative tracer to estimate submarine groundwater discharge (SGD) (Windom et al., 2006; Moore et al., 2008; Kwon et al., 2014; Rodellas

Deleted: While t

Deleted: ,

Moved (insertion) [2]

Deleted: their sources and fluxes to the ocean remain poorly constrained.he

Moved up [2]: their sources and fluxes to the ocean remain poorly constrained. For example,

Deleted: For example, the South Subtropical Convergence (SSTC) in the South Atlantic, near 40°S, is a prominent high productivity region and an important zone for the dynamic inter-ocean exchange between the Atlantic, Pacific, Indian, and Southern Oceans (Fig. 1). This region is also a transition zone between the nitrate-depleted Subtropical Gyre and the iron-limited Southern Ocean, creating one of the most dynamic nutrient environments in the world oceans (Moore et al., 2004). The chlorophyll concentration is generally high (0.2-0.3 mg chlorophyll a m⁻³) in this region (Longhurst, 2007). Modeling and experimental studies have both suggested that this region is iron limited or co-limited (Moore et al., 2004; Browning et al., 2014; 2017). However, the sources and fluxes of TEs to this region are not well quantified. ...Previous work has assessed TE input to the wider Previous work has assessed TE input to the wider South Atlantic from rivers (Vieira et al., 2020), atmospheric dust (Gaiero et al., 2013), shelf sediments (Graham et al., 2015), the Agulhas current (Paul et al., 2015) and hydrothermal vents (Saito et al., 2013). There are also studies of TE distributions in some areas of the South Atlantic (e.g. Chever et al., 2010; Bown et al., 2011). Two UK-GEOTRACES cruises in 2010-2012 provided a significant increase to such observations, focusing particularly on 40°S (Homoky et al., 2013; Browning et al., 2014; Wyatt et al., 2014, 2020; Chance et al., 2015; Menzel Barraqueta et al., 2019). These studies did not assess the fluxes of TEs by ocean mixing and transport. In this study, we address this issue using samples taken on the UK-GEOTRACES cruises.

Deleted: The chlorophyll concentration is generally high (0.2-0.3 chlorophyll a m⁻³) in this region (Longhurst, 2007). Modeling and experimental studies have both suggested that this region is iron limited or co-limited (Moore et al., 2004; Browning et al., 2014; 2017). However, the sources and fluxes of TEs to this region are not well quantified. ...Previous work has assessed TE input to the wider Previous work has assessed TE input to the wider South Atlantic from rivers (Vieira et al., 2020), atmospheric dust (Gaiero et al., 2013), shelf sediments (Graham et al., 2015), the Agulhas current (Paul et al., 2015) and hydrothermal vents (Saito et al., 2013). There are also studies of TE distributions in some areas of the South Atlantic (e.g. Chever et al., 2010; Bown et al., 2011). Two UK-GEOTRACES cruises in 2010-2012 provided a significant increase to such observations, focusing particularly on 40°S (Homoky et al., [1]

Formatted: Font colour: Text 1

Deleted: Previous work has assessed TE input to the wider South from rivers (Vieira et al., 2020), atmospheric dust (Gaiero et al., 2013), shelf sediments (Graham et al., 2015), the Agulhas current (Paul et al., 2015) and hydrothermal vents (Saito et al., 2013). The [2]

Deleted: play an important role in

Deleted: transporting

Deleted: s

et al., 2015; Le Gland et al., 2017), river inputs (Vieira et al., 2020), continental shelf (Rutgers van der Loeff et al., 1995; Charette et al., 2016; Sanial et al. 2018; Kipp et al., 2018a) and hydrothermal inputs (Kipp et al., 2018b).

Previous work has assessed TE inputs to the wider South Atlantic from rivers (Vieira et al., 2020), atmospheric dust (Gaiero et al., 2013), shelf sediments (Graham et al., 2015), the Agulhas current (Paul et al., 2015) and hydrothermal vents (Saito et al., 2013). There are also studies of TE distributions and basin-scale inputs in some areas of the South Atlantic (e.g. Chever et al., 2010; Bown et al., 2011; Noble et al., 2012). Two UK-GEOTRACES cruises in 2010-2012 provided a significant increase in such observations, focusing particularly on 40°S (Homoky et al., 2013; Browning et al., 2014; Wyatt et al., 2014, 2020; Chance et al., 2015; Menzel Barraqueta et al., 2019). These published studies did not assess the fluxes of TEs by ocean mixing and transport. In this study, we address this issue using samples taken on the two 40°S Atlantic UK-GEOTRACES cruises. We investigate the distributions of ²²⁸Ra, as well as ²²⁶Ra, in both the Argentine and Cape Basins of a 40°S latitudinal transect in the Atlantic Ocean. This is also the first exploration of the vertical and horizontal ²²⁸Ra distributions reported for the Cape Basin. We investigate the application of seawater ²²⁸Ra as a tracer for vertical and horizontal mixing in the surface South Atlantic, to provide estimates of the dissolved TE fluxes, with a focus on cobalt, iron and zinc, in the micronutrient-depleted euphotic zone.

2 Study sites and methods

2.1 Hydrographic setting

The study was conducted from the RRS *Discovery* and RRS *James Cook* during two UK GEOTRACES cruises, D357 and JC068, along the GA10 40°S transect of the Atlantic Ocean (Fig. 1). The surface currents show dynamic interaction and mixing on both the western and eastern sides of this transect. In the Argentine Basin, the Rio de la Plata estuary is located on the western margin of the transect. The boundary currents Brazil Current (BrC) and Malvinas Current (MC) meet between 33°S and 45°S along the continental margin of South America and these become the South Atlantic Current (SAC) transporting water eastwards along 40°S. The water from the BrC is captured between Stn20 and Stn21 with a strong west to east gradient in salinity and other chemical and physical properties, which also suggests a limited exchange of water across the continental shelf break (see below). In the Cape Basin, the SAC turns northeast before reaching the continent of Africa, and the Agulhas Current (AC) adds warm water eddies from the Indian Ocean. The warm and salty water of the AC was sampled in the top 500 m at Stn2 (Fig. 1b). These currents meet and become the Benguela Current (BeC) flowing through the Cape Basin and into the South Atlantic.

2.2 Water sampling

Seawater samples for Ra analysis were collected from 14 stations on the D357 and JC068 cruises (Fig. 1). The first cruise (D357) took place in the SE Atlantic (Cape Basin) between October and November 2010; the second cruise (JC068) took place along the whole 40°S transect between December 2011 and January 2012. For radium isotope analyses, the Cape Basin

Deleted: ¶
In this study, w

Deleted: M

Deleted: Study sites and sampling

Deleted: Seawater samples were collected from 14 stations on the RRS *Discovery* and RRS *James Cook* during two UK GEOTRACES cruises, GA10E (D357) and GA10W (JC068), along the 40°S transect of the Atlantic Ocean (Fig. 1). The first cruise (D357) took place in the SE Atlantic (Cape Basin) between October and November 2010; the second cruise (JC068) took place along the whole 40°S transect between December 2011 and January 2012. For radium isotope analyses, the Cape Basin samples were only taken from D357, and the Argentine Basin samples from JC068. Stations from these cruises are shown, along with those from previous Ra studies (e.g. GEOSECS and the Transient Tracers in the Ocean, TTO) in this region (Fig. 1a). ¶

Deleted: the Atlantic basins at 40°S

Formatted: Font: Bold

210 samples were only taken from D357, and the Argentine Basin samples from JC068. For trace elements (TEs), samples were taken from both D357 and JC068. Stations from these cruises are shown, along with those from previous Ra studies (e.g. GEOSECS and the Transient Tracers in the Ocean, TTO) in this region (Fig. 1a).

215 A total of 48 samples were analysed for $^{228}\text{Ra}/^{226}\text{Ra}$ ratio and ^{228}Ra concentration, and 33 samples for ^{226}Ra concentration, collected using three different sampling techniques during the cruise. Surface seawater samples (80 – 100 L) at 5 m depth were collected via a trace-metal clean seawater supply (fish) using a Teflon bellow pump (Almatec-A15) and acid-cleaned tubing. Samples between 50 m and 400 m were collected using a standard Conductivity, Temperature and Depth (CTD) rosette, typically sampling from four 20 litre Niskin bottles. These samples were stored briefly in low-density polyethylene (LDPE) cubitainers and then filtered through Mn-fibre cartridges by gravity (flow rate $< 0.5 \text{ L min}^{-1}$) on-board for the extraction of radium isotopes (Moore et al., 1985; Reid et al., 1979). In addition, large-volume seawater sampling (300 – 600 L) was carried out using in-situ stand-alone pump systems (SAPs), pumping sea water over Mn-fibres in polypropylene cartridges (van Beek et al., 2008) at flow rate of $2\text{--}5 \text{ L min}^{-1}$ at three selected sampling stations (Stn1, Stn3 and Stn4.5). As the cartridge Ra collection efficiency varies hugely, ranging from 70 to 128% (Geibert et al., 2013), all samples collected by these collection techniques (pump, CTD, and SAP) were only used for measurement of $^{228}\text{Ra}/^{226}\text{Ra}$ ratios. To address the efficiency issue, for most samples, a separate sample of 250 mL seawater was also collected for measurement of ^{226}Ra concentration. This method provides the advantage of allowing a correction for variable efficiencies during the sample preparation.

220 Trace element samples were collected using a titanium CTD rosette fitted with trace metal clean Teflon-coated Niskin bottles and filtered on-board through $0.8/0.2 \mu\text{m}$ polyethersulfone (PES) membrane cartridge filters (AcroPak500TM, Pall), before analysis. All the TE data and fluxes reported and discussed in this study refer to the dissolved fraction only. The data of zinc and cobalt were measured and published by Wyatt et al. (2014 and 2020). Some of the iron data were determined and published by Browning et al. (2014) and Clough et al. (2016). All the TE data are available on the GEOTRACES Intermediate Data Product (IDP) 2017 (Schlitzer et al., 2018).

230 All the trace-metal cleaning procedures followed the GEOTRACES sampling protocols (Cutter et al., 2010). In brief, the sample tubing and bottles were rinsed with Milli-Q water and filled with 0.1M HCl for one day. After emptying the acid, the tubing and bottles were rinsed thoroughly with Milli-Q water. The tubing and bottles were also rinsed with open-ocean seawater before sampling.

235 **2.3 Ra isotopes analysis**

Mn-fibre samples were first counted on-board using a 4-channel radium delayed coincidence counting (RaDeCC) system for ^{224}Ra , ^{223}Ra , ^{228}Th and ^{227}Ac (Moore and Arnold, 1996), but the data for ^{224}Ra , ^{223}Ra , ^{228}Th , and ^{227}Ac are not discussed in this paper. After the counting, Ra was purified using the following procedure for precisely measuring $^{228}\text{Ra}/^{226}\text{Ra}$ ratios and ^{226}Ra concentrations by a Nu Instrument multi-collector inductively-coupled-plasma mass-spectrometry (MC-ICP-MS) at the

Deleted: A

Deleted: by mass spectrometry, and ^{224}Ra , ^{223}Ra , ^{228}Th , and ^{227}Ac activities using a delayed coincidence RaDeCC device. ...

Deleted: F

Deleted: by mass spectrometry

Deleted: .

Deleted: (TE)

Deleted: 2

Deleted: radium delayed coincidence counting

Deleted: (RaDeCC)

University of Oxford following the procedures established by Hsieh and Henderson (2011). Mn-fibres were ashed at 550°C for 6 hours and then leached with distilled 6N HCl to remove Ra from the ashed fibres. Ra was then co-precipitated with Sr(Ra)SO₄ in the leached solution, centrifuged and cleaned with 3N HCl and pure H₂O a few times until the pH was > 4. To increase the dissolution rate, Sr(Ra)SO₄ was converted to Sr(Ra)CO₃ by adding 2 mL 1M Na₂CO₃ solution and heated on a hotplate for 3 hours. After centrifuging and discarding the supernatant, Sr(Ra)CO₃ was finally dissolved in 2 ml 6N HCl for ion exchange column chemistry, using Bio-Rad AG50-X8 cation exchange resin (to separate Ra and Ba from ²²⁸Th and other matrix elements, e.g. Ca, Sr and Mn) and Eichrom Sr-Spec resin (to separate Ra from Ba to avoid molecular interferences during MC-ICP-MS analysis).

Smaller seawater samples (250 mL) collected for ²²⁶Ra were spiked with a ²²⁸Ra-spike (Hsieh and Henderson, 2011) and the Ra was purified by the precipitation of CaCO₃ and processing with ion exchange columns of AG1-X8, AG50-X8 and Sr-Spec resin for the measurement of ²²⁶Ra concentrations by MC-ICP-MS (Foster et al., 2004). In general, the contribution of seawater ²²⁸Ra (< 0.05 attomole) is negligible, compared to the spiked ²²⁸Ra signal (≈ 70 attomole). Assessments of overall chemical blanks were conducted the same way as for the samples throughout the whole chemical procedures, except that there was no added seawater. The blanks were found to contribute less than 1% of the ²²⁶Ra in the sample and were not detectable for ²²⁸Ra.

During the MC-ICP-MS analyses, ²²⁸Ra and ²²⁶Ra were measured simultaneously on two ion counters, and the uranium standard CRM-145 was used to bracket each sample for the mass bias and ion counter gain corrections. Instrumental memories of ²²⁸Ra and ²²⁶Ra were also detected on ion counters before each measurement. The machine memory was about 0.2 ± 0.1 cps (counts per second) (n = 16, 2S.E.) The memory correction was insignificant for ²²⁶Ra, because the ratio of memory to sample signal is small (< 10⁻⁴). However, the memory correction could be significant for samples with low ²²⁸Ra activities and count rates. For instance, the count-rate during analysis of ²²⁸Ra on the sample collected at 4741 m at Stn4.5 was only 0.5 cps. At this low count-rate, instrumental memory contributed ~40% of the signal to the sample ²²⁸Ra signal and the uncertainty of memory correction becomes substantial. In this study, most of the surface and deep waters in the South Atlantic were measured at count rates > 2 cps ²²⁸Ra, which provides assurance that the contribution of the instrumental memory uncertainty to the total uncertainty is < 10%.

For samples without accompanied ²²⁶Ra measurements, silica data (Table 1) are used to assess ²²⁶Ra activities (Appendix A). The Atlantic ocean ²²⁶Ra-Si relationship is based on the GEOTRACES (GA03), GEOSECS and TTO datasets (Fig. A1, Ku and Lin, 1976; Key et al., 1990; 1992a,b; Charette et al., 2015). This relationship is used to determine ²²⁶Ra activities in the 27 cases where no subsamples were collected for separate ²²⁶Ra analysis (such ²²⁶Ra estimates are shown in brackets in Table 1). The relationship has a slope of 0.119 dpm 100 L⁻¹ of ²²⁶Ra per μmol L⁻¹ of Si and an intercept of 8.8 dpm 100 L⁻¹, which is comparable with the average slope of 0.1 observed by Broecker et al. (1976) in the Atlantic. The Si-extrapolated ²²⁶Ra and measured ²²⁶Ra activities (data from this study and the TTO) show a relatively consistent result although the extrapolated ²²⁶Ra has a larger uncertainty (±11% 2S.E.) than the measured ²²⁶Ra uncertainty (±4%, 2S.E.) The paired t-test

Deleted: (

Deleted: ,

Deleted: convert

Deleted: ,

Deleted: was added

Deleted:

Formatted: Superscript

Deleted: O

Formatted: Superscript

Deleted: monitored throughout the whole chemical procedures, and

Deleted: Measurements of ²²⁸Ra and ²²⁶Ra were performed using a Nu Instrument MC-ICP-MS using the protocols described in Hsieh and Henderson (2011).²²

Deleted: Ra-228

Deleted: ,

Deleted: T

Deleted: ,

Deleted: global

Deleted: from the ²²⁶Ra and silica data from

Deleted: 43

Deleted: 0

shows a p-value of 0.55 (> 0.05), suggesting that the differences between the extrapolated ^{226}Ra and the measured ^{226}Ra data are not statistically significant. The uncertainties of ^{226}Ra activity have been used in the error propagation of ^{228}Ra activity; the total uncertainty of ^{228}Ra is typically about 6 – 12% (2S.E.)

2.4 ^{228}Ra -derived 1-D mixing models

The distribution of seawater ^{228}Ra in the ocean is mainly controlled by mixing, advection, radioactive decay and additional removal/input. It has been widely used as a tracer for measuring diffusion coefficients and advection rates on a basin-wide scale in the surface or at intermediate depths in the ocean (e.g. Cochran, 1992; Ku and Luo, 2008; Sanial et al., 2018). The one-dimensional (1-D) ^{228}Ra advection-diffusion model is commonly expressed by the formula (e.g. Moore, 2015):

$$\frac{\partial A}{\partial t} = K_x \frac{\partial^2 A}{\partial x^2} - w \frac{\partial A}{\partial x} - \lambda A \pm J \tag{1}$$

where A is activity of ^{228}Ra , t is time, K_x is horizontal eddy diffusion coefficient, w is advection velocity, x is offshore distance, λ is decay constant ($\lambda_{\text{Ra}228} = 3.82 \times 10^{-9} \text{ s}^{-1}$), and J is additional input or removal of ^{228}Ra .

To use the model to accurately calculate the mixing rates, several assumptions need to be made:

(1) Steady state ($\partial A / \partial t = 0$)

This assumption requires long-term monitoring of ^{228}Ra activities in the ocean due to the long half-life of ^{228}Ra . Charette et al. (2015) compared the ^{228}Ra data in the North Atlantic from the US GEOTRACES and the TTO programs, and found that the upper ocean ^{228}Ra inventories have remained constant over the past 30 years. Although ^{228}Ra seasonality in coastal areas may introduce uncertainty to the mixing model, the assumption of steady state is likely to be valid for ^{228}Ra on decadal timescales and at ocean-basin scales. The comparison between our ^{228}Ra data and the limited data from the TTO in this region show good agreement (Fig. 2b), supporting this assumption for the South Atlantic.

(2) No additional input or removal of ^{228}Ra ($\pm J = 0$)

Based on particle removal, Ra residence time is estimated to be ~500 years in the surface ocean (Moore and Dymond, 1991). However, there is no measurable particle removal of Ra in the surface open ocean at the time scale of ^{228}Ra half-life (5.75 years) (Moore, 2015). In theory, the distribution of seawater ^{228}Ra is controlled by both vertical and horizontal mixing. For example, vertical mixing could potentially introduce additional “removal” of ^{228}Ra from the surface water and affect the horizontal distribution of ^{228}Ra in the surface ocean, which would require 2-D models and a high sample resolution dataset to resolve the problem. However, vertical mixing ($\sim 0.1 - 1 \text{ cm}^2 \text{ s}^{-1}$) is typically 5 to 8 orders of magnitude smaller than horizontal mixing ($\sim 10^5 - 10^7 \text{ cm}^2 \text{ s}^{-1}$), and the sample resolution is not good enough for precise 2-D modeling in this study. Therefore, we only apply the 1-D model to estimate the maximum horizontal mixing rates by neglecting the term of downward mixing.

Deleted: 3

Deleted: the decade

Formatted: Superscript

Deleted: v

Formatted: Superscript

Deleted: water

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Deleted: is not good enough for precise 2-D modeling

Deleted: . Therefore

Deleted: ,

Similar assumptions need to be made for the ^{228}Ra -derived vertical mixing - the gradient of ^{228}Ra with depth mainly reflects the vertical mixing rates, and the vertical ^{228}Ra distribution is not affected by additional input of horizontal ^{228}Ra below the mixed layer. These assumptions are mostly true in the upper open ocean, where horizontal ^{228}Ra mainly comes from the continental margins (shelf and slope sediments) and there is a lack of other important ^{228}Ra sources in the middle of the ocean. However, if the lateral input becomes important and starts to interfere with the ^{228}Ra vertical profiles (e.g. receiving strong advective shelf water or profiles closer to the seafloor), the 1-D ^{228}Ra -derived vertical mixing rates can be significantly overestimated.

Formatted: Superscript
Formatted: Superscript
Formatted: Superscript

Deleted:

(3) Boundary conditions ($A = A_0$ at $x = 0$ and $A = 0$ at $x \rightarrow \infty$)

It has been pointed out that the boundary condition of $A = 0$ at $x \rightarrow \infty$ is incorrect for using ^{228}Ra to determine coastal mixing as the distribution of ^{228}Ra could be controlled by water mass mixing within the coastal distance scale (< 50 km) rather than eddy diffusion (Moore, 2000). In theory, this boundary condition may be valid on the ocean-basin scale, as the major sink of ^{228}Ra in the ocean is radioactive decay. However, observed seawater ^{228}Ra is still not completely zero in the remote ocean. To avoid this problem, we follow the suggestion of Moore (2015) and define the ^{228}Ra excess ($^{228}\text{Ra}_{\text{ex}} = ^{228}\text{Ra} - ^{228}\text{Ra}_{\text{bg}}$) by subtracting the background value in the middle of the South Atlantic, $^{228}\text{Ra}_{\text{bg}}: 0.23 \pm 0.06$ dpm 100 L^{-1} (ISE, $n=6$). This value is determined by the average of the observed values at (1) the remote surface waters in a previous study around the 40°S transect (Hanfland, 2002; Fig. 1, ANT XV/4 station S8, S10 and S11); and (2) the water depth between 1000 and 3500 m in this study (except for 2580 m at Stn1 on the continental slope). The mid-depth ^{228}Ra background (1000 – 3500 m) shares a similar background as the remote surface waters (~ 0.2 dpm 100 L^{-1}), suggesting that the ^{228}Ra background needs to be corrected in both horizontal and vertical mixing calculations. For comparison, the central North Atlantic shows a similar mid-depth ^{228}Ra background value (~ 0.16 dpm 100 L^{-1}) between 1000 and 3000 m depth from the GEOTRACES GA03 transect (Stn12 ~ 20, Charette et al., 2015). However, the surface value in the central North Atlantic (~ 2.2 dpm 100 L^{-1}) is significantly higher than that in the South Atlantic (~ 0.2 dpm 100 L^{-1}), which has also been observed by Moore et al. (2008). We therefore use a background value of $^{228}\text{Ra}_{\text{bg}}: 0.23 \pm 0.06$ dpm 100 L^{-1} determined from the South Atlantic.

Deleted: T

Deleted: Although,
Deleted: on the ocean-basin scale,
Deleted: ,

Deleted: the water depth between 1000 and 3500 m (except 2580 m at Stn1 on the continental slope) in this study and the

Deleted: This mid-water ^{228}Ra background cannot be supported by in-situ production from ^{232}Th in the water column, because the activity of ^{232}Th is ~ 100 times lower than that of ^{228}Ra ($^{232}\text{Th} = 1\sim 20 \times 10^{-4}$ dpm 100 L^{-1} , Deng et al., 2014). ...

Formatted: Superscript
Formatted: Superscript

The boundary condition can now be rewritten as $A_{\text{ex } 0} = A_0 - A_{\text{bg}}$ at $x = 0$ and $A_{\text{ex}} = 0$ at $x = \infty$. Considering the assumptions discussed above, Eqn. (1) can be written as:

Deleted: In this case, t

$$0 = K_x \frac{\partial^2 A_{\text{ex}}}{\partial x^2} - w \frac{\partial A_{\text{ex}}}{\partial x} - \lambda A_{\text{ex}} \quad (2)$$

In this study, we consider two scenarios in the horizontal ^{228}Ra calculations: (1) mixing only ($w = 0$); and (2) advection only ($K_x = 0$). We use these two scenarios to provide independent assessments of chemical fluxes in the surface ocean, to bracket the range of possible TE fluxes that are consistent with the ^{228}Ra data regardless of the combination of mixing and advection in the real ocean. The advection model is only applied to the Argentine Basin data after the shelf break where the advection signal is strong because of the Brazil Current. Although the advection only scenario is an unrealistic one, it provides an end-

member for comparisons of TE fluxes under different settings. Using the boundary conditions, $A_{ex,0} = A_0 - A_{bg}$ at $x = 0$ and $A_{ex} = 0$ at $x = \infty$, Eqn. (2) can be solved for diffusive mixing only:

$$A_{ex} = A_{ex,0} \exp(-ax), \text{ where } a = \sqrt{\lambda/K_x} \quad (3)$$

and for advection only:

$$w = \lambda x / \ln(A_{ex,0}/A_{ex}) = X_{1/2}/T_{1/2} \quad (4)$$

where $A_{ex,0}$ is the activity of $^{228}\text{Ra}_{ex}$ at $x = 0$ (i.e. Stn0 or Stn25); $X_{1/2}$ is the distance at which $A_{ex} = 0.5A_{ex,0}$; and $T_{1/2}$ is the half-life of ^{228}Ra (5.75 years). The exponential fit of the surface ^{228}Ra data provides the estimate of the maximum diffusion coefficients (K_x) at both ends of the transect, and the linear fit of the surface ^{228}Ra data after the shelf break in the Argentine basin provides the minimum estimate of the advection water transport (w) along the west end of the transect (see discussion below).

For vertical mixing, the calculation of K_z is based on a situation in which ^{228}Ra is mixed horizontally away from the coast in the surface mixed layer and then down into the subsurface. The 1-D mixing model can therefore be applied to the depth profiles of ^{228}Ra to calculate K_z near the surface ocean. Under similar boundary conditions $A_{ez,0} = A_0 - A_{bg}$ at $z = 0$ and $A_{ez} = 0$ at $z = \infty$, the diffusion equation can be solved for vertical mixing to fit the vertical ^{228}Ra profiles:

$$A_{ez} = A_{ez,0} \exp(-az), \text{ where } a = \sqrt{\lambda/K_z} \quad (5)$$

where $A_{ez,0}$ is the activity of $^{228}\text{Ra}_{ex}$ at $z = 0$ (i.e. the mixed layer). In the 1-D mixing model, the term for diapycnal advection is generally negligible, as the oceanic vertical advection velocity is usually very small, i.e. $10^{-3} \sim 10^{-5} \text{ cm s}^{-1}$ (Liang et al., 2017).

2.5 Trace element flux calculations

In this study, we use three different ^{228}Ra approaches to quantify the horizontal and diffusive vertical dTE fluxes in the Cape Basin and Argentine. More details of the calculations are provided in Appendix D.

(1) ^{228}Ra -derived-diffusive TE fluxes

To calculate both lateral and vertical TE fluxes, the ^{228}Ra -derived diffusion coefficients (K_x or K_z) are applied to Fick's first law of molecular diffusion in the following equation:

$$F_{TE-d} = K_{x \text{ or } z} (\Delta TE / \Delta x \text{ or } \Delta z) \quad (6)$$

where F_{TE-d} is the diffusive flux of the TEs, and $\Delta TE / \Delta x$ or Δz is the gradient of TE concentration over either the horizontal distance x to the coasts or the vertical depth z below the mixed layer, which can be obtained from the linear regression of horizontal and vertical TE profiles (section 3.2).

Formatted: Font: Not Bold

410 (2) ²²⁸Ra-derived-advective TE fluxes

Formatted: Font: Not Bold

Surface water after the boundary of the shelf break and the Brazil Current in the Argentine Basin carries strong offshore advection signals along the SAC towards the open ocean (Fig. 1). Assuming that the mixing of TE is conservative, the advective TE fluxes can be calculated using the following equation:

$$F_{TE-a} = w \cdot [TE]_{ave-0} \quad (7)$$

415 where F_{TE-a} is the offshore advective flux of TE, w is the net offshore advection velocity along the SAC in the Argentine Basin, and $[TE]_{ave-0}$ is the average concentrations of dissolved TEs in the initial advective waters around where the Brazil Current merges into the SAC (around Stn21).

(3) TE/²²⁸Ra-ratio-derived TE fluxes

Formatted: Font: Not Bold

420 Previous studies have combined the use of the shelf ²²⁸Ra fluxes with the ratios of TE/²²⁸Ra in the surface waters between continental shelves and open oceans to estimate the shelf-ocean TE inputs from the continental margins to the open oceans (Charette et al., 2016; Sanial et al., 2018; Vieira et al., 2020). This method provides the integrated net fluxes of TEs, considering all the possible inputs (e.g. rivers, SGD and sediments) and outputs (e.g. particle scavenging, biological uptake and radioactive decay) of ²²⁸Ra and TEs during water mixing between the continental shelf and the open ocean. More details of the method are given in Charette et al. 2016. In brief, assuming that the net shelf-ocean exchange is mainly driven by eddy diffusion, the cross-shelf TE fluxes can be calculated using the following equation:

$$F_{TE} = F_{228Ra} \cdot \left(\frac{\Delta TE}{\Delta 228Ra} \right) = F_{228Ra} \cdot \left(\frac{TE_{shelf} - TE_{ocean}}{228Ra_{shelf} - 228Ra_{ocean}} \right) \quad (8)$$

425 where F_{228Ra} is the cross-shelf ²²⁸Ra flux (Appendix D); TE_{shelf} and $228Ra_{shelf}$ are the average concentrations of the TE and ²²⁸Ra in the surface waters on the shelf (GA10W: between Stn23 and Stn25; GA10E: between Stn0 and Stn1) respectively; and TE_{ocean} and $228Ra_{ocean}$ are the average concentrations in the open ocean (GA10W: between Stn18 and Stn19; GA10E: between Stn4 and Stn4.5). The ratios of $\Delta TE/\Delta 228Ra$ are reported in Table D1.

Deleted: s

3 Results

3.1 Ra isotope concentrations

435 The results of ²²⁶Ra and ²²⁸Ra activities and the activity ratios of ²²⁸Ra/²²⁶Ra are presented in Table 1 (with \pm 2S.E.) The vertical profiles of measured ²²⁶Ra show good agreement with the data from the closest GEOSECS and TTO stations in this region (e.g. Ku and Lin, 1976; Key et al., 1990; 1992a,b) (Fig. 2a). Ra-226 activities range from 8.2 to 22.4 dpm 100 L⁻¹. In this study, the results of ²²⁶Ra are mainly used for calculating ²²⁸Ra activities, and will not be discussed in further detail.

The activity ratios of $^{228}\text{Ra}/^{226}\text{Ra}$ range from 0.017 to 1.599 in the surface water and are comparable with the ratios from 0.080 to 2.810 observed in previous studies in this region (TTO data, Windom et al., 2006 and Hanfland, 2002). The vertical profiles of ^{228}Ra activity are shown in Fig. 2b. In surface waters, the activities of ^{228}Ra ranging from 1.02 to 17.66 dpm 100 L⁻¹ in the Argentine Basin are consistent with the observed values from 0.07 to 24.0 dpm 100 L⁻¹ from the previous studies (TTO data and Windom et al. 2006), and the activities of ^{228}Ra ranging from 0.99 to 3.22 dpm 100 L⁻¹ in the Cape Basin are consistent with the observed values from 0.67 to 4.23 dpm 100 L⁻¹ from the previous study (Hanfland, 2002). Between 600 m and 4000 m, $^{228}\text{Ra}/^{226}\text{Ra}$ ratios decrease to 0.015 – 0.030 and ^{228}Ra activities decrease to 0.29 – 0.32 dpm 100 L⁻¹. In the 100 m closest to the ocean floor, the ratios of $^{228}\text{Ra}/^{226}\text{Ra}$ increase to 0.048 – 0.088 and the activities of ^{228}Ra also increase to 1.07 – 1.94 dpm 100 L⁻¹. This is the first dataset of seawater ^{228}Ra reported in the intermediate and deep waters in the Cape Basin. ~~The ^{228}Ra values in the Cape Basin are noticeably higher~~ than those observed in intermediate (< 0.1 dpm 100 L⁻¹) and deep waters (0.22 – 0.28 dpm 100 L⁻¹) to the south in the Southern Ocean (Charette et al., 2007; van Beek et al., 2008). The vertical profiles of seawater ^{228}Ra in the Cape Basin are consistent ~~with~~ GEOSECS and TTO observations elsewhere in the South Atlantic Ocean (Moore et al., 1985). ~~The samples collected by pump (fish), CTD and SAP within the mixed layer at each station show consistent ^{228}Ra and ^{226}Ra results, suggesting that there is no significant difference in the Ra results between these three sampling methods.~~

Deleted: It shows higher

Deleted: , however, with expectations based on

Formatted: Superscript

Formatted: Superscript

3.2 Micronutrient concentrations

Dissolved Co, Fe and Zn concentration data (Wyatt et al., 2014 and 2020; Browning et al., 2014; Clough et al., 2016; Schlitzer et al., 2018) are summarised in Appendix B (Table B1 and B2). At the continental margins, surface shelf waters show much higher trace element concentrations (Co: 146.2 pM, Fe: 1.53 nM, and Zn: 0.59 nM in the Argentine margin; Co: 46.9 pM, Fe: 0.35 nM, and Zn: 0.14 nM in the Cape margin) than observed in the open ocean surface waters along the 40°S transect (Fig. 3). ~~In the Argentine Basin, the distribution of trace elements generally follows the salinity in the surface waters. The low salinity waters (< 29 psu) around 200 km from the South America coast show high TE concentrations (Co: > 80 pM, Fe: > 1 nM, and Zn: > 0.5 nM). In contrast, the high salinity waters (> 35 psu) in the Brazil Current show much lower TE concentrations (Co: < 60 pM, Fe: < 0.5 nM, and Zn: < 0.2 nM). Similar correlations between trace elements and salinity have been observed along other western boundaries of the Atlantic as well (e.g. the North American shelf, Bruland and Franks 1984; Noble et al., 2017). In the upper ocean (< 600 m), these trace element concentrations are generally low in the surface mixed layer and increase with depth below the mixed layer (Fig. 4). Co concentrations range from 1.6 to 61.2 pM, and Fe and Zn concentrations range from 0.05 to 0.54 and 0.01 to 0.53 nM, respectively. In general, these trace element concentrations (< 600 m) are slightly higher in Stn1 than other stations further away from the continental shelf.~~

Deleted: T

Deleted: (Fig. 4)

Deleted: In the upper 500 m,

4 Discussion

4.1 ^{228}Ra -derived horizontal mixing and advection

4.1.1 Argentine Basin

The distribution of $^{228}\text{Ra}_{\text{ex}}$ in the Argentine Basin (GA10W) is controlled by the Rio de la Plata river plume and the Brazil Current (Fig. 5a), and this is supported by a good correlation with salinity (linear regression $R^2 = 0.96$; Fig. C1 in Appendix C). If we apply the 1-D mixing model (Eqn.3) to fit the $^{228}\text{Ra}_{\text{ex}}$ data between Stn25 and Stn21, across the boundary of the Brazil Current, the gradient of the exponential fit (a) is 0.0047 ± 0.0027 and the estimate of the offshore horizontal diffusion coefficient K_x is $1.8 \pm 1.4 \times 10^6 \text{ cm}^2 \text{ s}^{-1}$, which is likely to be an overestimate due to the influence of the river plume and the advection of the boundary current. Nevertheless, this estimate is still within the range of other estimates of K_x between 10^5 and $10^8 \text{ cm}^2 \text{ s}^{-1}$ in a variety of margin and open ocean settings (e.g. Kaufman et al., 1973; Knauss et al., 1978; Yamada and Nozaki, 1986).

Across the boundary of the Brazil Current, the advection of the eastward flowing SAC shows a significant impact on the distribution of $^{228}\text{Ra}_{\text{ex}}$ in the surface Argentine Basin (Fig. 5a). If we apply the 1-D advection model (Eqn.4) to fit the data between Stn21 and Stn18 (Fig. 5a), the $X_{1/2}$ is $11 \pm 600 \text{ km}$ (from Stn 21) and the estimate of average advection velocity w is $0.6 \pm 0.3 \text{ cm s}^{-1}$. Although the ^{228}Ra -derived velocity is smaller than the typical velocities ($2 \sim 4 \text{ cm s}^{-1}$) around the South Atlantic Subtropical Gyre (Schlitzer, 1996), similar advective ^{228}Ra signals have been previously observed in other surface ocean current systems, including the Peru and Kuroshio currents in the Pacific (Knauss et al. 1978 Yamada and Nozaki 1986).

4.1.2 Cape Basin

Ra-228 data from the Cape transect (GA10E) are used to calculate the offshore horizontal diffusion coefficient (K_x) in the Cape Basin (Fig. 5b). Applying the simple 1-D mixing model, assuming $w = 0$, to fit the surface $^{228}\text{Ra}_{\text{ex}}$ data in the Cape Basin (Fig. 5b), the gradient of the exponential fit (a) is 0.0012 ± 0.0003 and the estimate of K_x is $2.7 \pm 0.8 \times 10^7 \text{ cm}^2 \text{ s}^{-1}$, which is an order of magnitude higher than the value observed in the Argentine Basin (see above), but within the range of other observed values in the oceans ($10^5 \sim 10^8 \text{ cm}^2 \text{ s}^{-1}$).

The horizontal diffusion coefficient K_x is likely to be overestimated in the Cape Basin due to the influence of episodic water advection. The Agulhas Current Leakage (ACL) is known for transporting water from the Indian Ocean into the South Atlantic and episodically introduces eddies (Agulhas Rings) into the Cape Basin (Beal et al., 2011). However, the signals of mixing and advection cannot be easily separated with the ^{228}Ra data alone (Fig. 5b). For example, the distribution of $^{228}\text{Ra}_{\text{ex}}$ in the surface Cape Basin shows elevated values (Stn2 and Stn4.5) above the fitted curve and coincides with the elevated salinity and temperature data (Fig. 5d), which indicates that the elevated $^{228}\text{Ra}_{\text{ex}}$ is likely to come from an advective signal (e.g. ACL). The ACL signal has also been identified with a distinct Pb isotope signature in the upper water column at Stn2

Deleted: 7

Deleted: 5

Deleted: 1

Deleted: 13

(Paul et al., 2015). The application of a 1-D mixing model may actually be biased by the addition of these high $^{228}\text{Ra}_{\text{ex}}$ waters, therefore, the horizontal diffusion coefficient K_x is likely to be a maximum estimate for the Cape Basin. Nevertheless, the overall gradient of $^{228}\text{Ra}_{\text{ex}}$, decreasing along the distance away from the shore, is driven by the loss of ^{228}Ra through both water mixing and radioactive decay.

Despite uncertainty in the diffusion coefficients due to advection of other sources, the ^{228}Ra data do place bounds on maximum horizontal mixing in the surface ocean away from the eastern and western boundaries of the Atlantic at 40°S. These bounds can be used to quantify the trace element inputs from the continental margins to the South Atlantic (see Discussion 4.3).

4.2 ^{228}Ra -derived vertical mixing

Vertical diffusion coefficients (K_z) are calculated at six stations where the depth profiles of $^{228}\text{Ra}_{\text{ex}}$ are available (Stn1, Stn2, Stn3, Stn4.5, Stn18 and Stn21). The best-fit exponential curve gradients (a) from the depth (z) profiles of $^{228}\text{Ra}_{\text{ex}}$ activities below the surface mixed layer are used in the same 1-D mixing model (Eqn.5) to calculate the vertical mixing coefficient K_z for the upper ≈ 600 m of each station in both the Argentine and Cape Basins (Fig. 6), and resulting K_z values range from 1 to $53 \text{ cm}^2 \text{ s}^{-1}$ at these stations. It should be noted that Stn2 only has one Ra data point below the mixed layer, and hence it is not considered in the vertical TE input calculations (section 4.3), but the estimate of K_z shows a similar value as other stations with more data points. The high K_z values of $53 \text{ cm}^2 \text{ s}^{-1}$ and $7 \text{ cm}^2 \text{ s}^{-1}$ at Stn18 and Stn21, respectively, are most likely biased by the lateral inputs of ^{228}Ra below the mixed layer (see later discussion). Excluding the values of Stn18 and Stn21, the range of K_z values from 1.0 to $1.7 \text{ cm}^2 \text{ s}^{-1}$ is broadly comparable to the average K_z of $1.5 \text{ cm}^2 \text{ s}^{-1}$ assessed from tritium measurements in the South Atlantic (Li et al., 1984). These estimates are also consistent with the range of observed vertical mixing from 0.1 to $10 \text{ cm}^2 \text{ s}^{-1}$ using different methods (e.g. ^7Be , SF₆ dye release and microstructure shear probe methods) in different ocean-basin settings (Kunze and Sanford, 1996; Ledwell et al., 1993; Martin et al., 2010; Painter et al., 2014; Kadko et al., 2020).

Given that the calculation of K_z is based on the vertical gradients of ^{228}Ra driven by vertical mixing and radioactive decay only, it therefore relies on the assumption that the vertical gradients are not dominated by lateral input of ^{228}Ra at depths below the surface mixed layer. This assumption is supported by an inspection of horizontal ^{228}Ra gradients at depths below the mixed layer (Fig. 7). Due to the sample resolution, detailed inspection is only available for the Cape Basin. Here, unlike the exponential change seen in the surface layer, $^{228}\text{Ra}_{\text{ex}}$ activities at 50 m and deeper do not show an increasing gradient towards the continental margin. This therefore argues against lateral mixing away from the shore as the major mechanism driving sub-surface ^{228}Ra concentrations in the Cape Basin. In addition, the distribution of ^{228}Ra on an isopycnal surface (≈ 200 m depth) is largely constant and shows no lateral gradient (Fig. 7). Studies using theoretical models to simulate seawater ^{228}Ra distribution have also shown that the horizontal eddy mixing (K_x) has little effect on the vertical distribution of ^{228}Ra (Lamontagne and Webster, 2019).

Deleted: ,

Deleted: at its

Deleted: The calculation of vertical (diapycnal) mixing (K_z) is based on a situation in which ^{228}Ra is mixed horizontally away from the coast in the surface mixed layer and then down into the subsurface. The 1-D mixing model (Eqn.3) can therefore be applied to the depth profiles of $^{228}\text{Ra}_{\text{ex}}$ in the upper 600 m of the water columns to calculate K_z near the surface ocean (Fig. 6). In the 1-D mixing model, the term of diapycnal advection is generally negligible, as the oceanic vertical advection velocity is usually very small, i.e. $10^{-3} \sim 10^{-5} \text{ cm s}^{-1}$ (Liang et al., 2017). The best-fit exponential curves in $^{228}\text{Ra}_{\text{ex}}$ activities to depth (z) below the surface mixed layer are used in the same 1-D mixing model (Eqn.3) to calculate vertical mixing K_z for the upper ≈ 600 m of each station in both the Argentine and Cape Basins (Fig. 6).

Deleted: ,

Deleted: 65

Deleted: (Fig. 6)

Formatted: Font: Italic

Formatted: Font: Italic, Subscript

Formatted: Font: Italic

Deleted: 65

Deleted: 5

Deleted: sigma-t of $\sim 26.7 \text{ kg m}^{-3}$;

Formatted: Font: Italic

560 The depth profiles of $^{228}\text{Ra}_{\text{ex}}$ in the Argentine Basin show evidence of advective ^{228}Ra below the mixed layer, potentially from the nearer shore shelf waters. For example, elevated $^{228}\text{Ra}_{\text{ex}}$ values are seen around 400 m at Stn18 and 200 m at Stn21 (Fig. 6a and b) and these may explain the extremely high K_z values at these stations. Although the possibility of lateral inputs cannot be entirely excluded, particularly in the Argentine Basin, the vertical variation of ^{228}Ra near the surface mixed layer still provides the first estimates of maximum vertical mixing and the upper limits of trace element inputs from vertical mixing to the surface ocean along the 40°S transect.

4.3 Trace element inputs in the South Atlantic

Trace elements (TEs) are important micronutrients for marine productivity in the surface ocean. Atmospheric dust deposition is an important source of TEs to the surface ocean. The soluble atmospheric dust deposition fluxes to the surface 40°S Atlantic have been assessed from the same cruise as this study: these are $0.02 - 0.05 \text{ nmol Co m}^{-2} \text{ d}^{-1}$, $1.6 - 5.2 \text{ nmol Fe m}^{-2} \text{ d}^{-1}$ and $0.6 - 6 \text{ nmol Zn m}^{-2} \text{ d}^{-1}$ (Chance et al., 2015). However, other inputs of these TEs to the euphotic zone in the South Atlantic are still unknown (e.g. shelf-ocean and vertical mixing). In this study, we consider three different ^{228}Ra approaches to quantify the horizontal and vertical TE fluxes in the Cape Basin and Argentine Basin along the 40°S transect: (1) ^{228}Ra -derived-diffusive; (2) ^{228}Ra -derived-advective; and (3) TE/ ^{228}Ra -ratio-derived TE fluxes (section 2.5 and Appendix D). The results of the TE fluxes are summarized in Table 2 (shelf-ocean, horizontal) and Table 3 (vertical). For comparison, the horizontal TE fluxes are normalised to the areas of shelf-ocean cross-section (Table D2, Urien and Ewing, 1974; Nelson et al., 1998; Emery, 1966; Windom et al., 2006; Carr and Botha, 2012; Hooker et al., 2013; Vieira et al., 2020) (illustrated in Fig. 8) unless otherwise specified.

Surprisingly, the estimates of shelf-ocean TE fluxes show relatively good agreements (within uncertainties) between these three approaches (Table 2), given the limitations of the 1-D ^{228}Ra -mixing model (Moore, 2015). A similar observation has also been found in the ^{228}Ra study in the Peruvian continental shelf (Sanial et al., 2018), which suggests that the assumptions made for the 1-D ^{228}Ra -mixing model are reasonable. In addition, the TE fluxes show consistent results between the D357 and JC068 data in the Cape Basin. These observations are likely to be a result of the gradients of ^{228}Ra and TEs representing a long-term average at an ocean basin scale and being closer to a steady-state condition in the upper water column (e.g. the ^{228}Ra and TEs in the North Atlantic, Charette et al., 2015).

585 The ^{228}Ra -derived shelf-ocean Co fluxes range from 4 to $2.1 \times 10^3 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Argentine margin and from 4.3 to $6.2 \times 10^3 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Cape margin of the 40°S transect in the South Atlantic. In comparison, previous studies have applied the TE/ ^{228}Ra approach to estimate the shelf Co fluxes in several continental margins: the western North Atlantic ($1.6 \times 10^5 \text{ nmol m}^{-2} \text{ d}^{-1}$, Charette et al., 2016), the Peruvian shelf ($1.4 \times 10^5 \text{ nmol m}^{-2} \text{ d}^{-1}$, Sanial et al., 2018) and the Congo-offshelf 3°S ($2.8 \times 10^6 \text{ nmol m}^{-2} \text{ d}^{-1}$, Vieira et al., 2020). Although these fluxes are about one and two orders of magnitude respectively higher than the estimates in the South Atlantic, these regions are also associated with low oxygen which increases dissolution of Mn and Fe oxides in sediments and is prone to result in higher Co fluxes (e.g. Hawco et al., 2016). A low

Deleted: the

Deleted: , following the methods in Charette et al. (2016), Sanial et al. (2018) and Vieira et al. (2020); and the shelf ^{228}Ra fluxes, based on the global seawater ^{228}Ra inverse models (Kwon et al., 2014; Le Gland et al., 2017). The details of calculations are provided in ...

Moved (insertion) [1]

Deleted: Carr and Botha, 2012

Deleted: -

Deleted: -

Deleted: may be

Deleted: not un

Deleted: Comparing

Deleted: the shelf-ocean TE fluxes from this study with previous studies, all the TE fluxes are normalised to the area of the shelf-ocean cross section. The areas of shelf-ocean cross section from the reference regions are summarised in Table D2 (Emery, 1966; Windom et al., 2006; Hooker et al., 2013; Vieira et al., 2020). In this study, the

Moved up [1]: Emery, 1966; Windom et al., 2006; Hooker et al., 2013; Vieira et al., 2020). In this study, t

Deleted: 3.8

Deleted: 2

shelf-ocean Co flux has been reported in the eastern South Atlantic continental shelf ($11 \sim 18 \times 10^3 \text{ nmol m}^{-2} \text{ d}^{-1}$, Bown et al., 2011), which is ~~very~~ close to this study region in the Cape Basin.

615 Along the 40°S transect, the ^{228}Ra -derived shelf-ocean Fe fluxes range from ~~8~~ to ~~19~~ $\times 10^4 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Argentine margin and from $1.2 \text{ to } 3.1 \times 10^4 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Cape margin, which ~~are slightly lower than~~ the estimates of the shelf-ocean Fe flux ($4.5 \times 10^5 \text{ nmol m}^{-2} \text{ d}^{-1}$) in the western North Atlantic (Charette et al., 2016). However, these fluxes are significantly lower than ~~those high Fe fluxes observed in regions with river plumes (e.g., River Congo; $4.1 \times 10^8 \text{ nmol m}^{-2} \text{ d}^{-1}$, Vieira et al., 2020), submarine groundwater discharge ($1.3 \times 10^8 \text{ nmol m}^{-2} \text{ d}^{-1}$, Windom et al., 2006) and the oxygen~~

620 minimum zone ($2.1 \times 10^6 \text{ nmol m}^{-2} \text{ d}^{-1}$, Sanial et al., 2018).

Lastly, the ^{228}Ra -derived shelf-ocean Zn fluxes range from 2.7 to ~~6.3~~ $\times 10^4 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Argentine and from 0.9 to $1.2 \times 10^4 \text{ nmol m}^{-2} \text{ d}^{-1}$ in the Cape margins. When compared with the only available shelf-ocean Zn flux value ($1.8 \times 10^6 \text{ nmol m}^{-2} \text{ d}^{-1}$) in the western North Atlantic (Charette et al., 2016), the Zn fluxes from this study indicate ~~low Zn inputs~~ in the South Atlantic. ~~The different shelf-ocean Zn inputs between the North and South Atlantic require more detailed study to understand the processes supplying and removing Zn in the shelf waters and the influence of anthropogenic Zn. Nevertheless, the low Zn inputs support the previous observation that surface water along the 40°S transect has some of the lowest reported dissolved Zn concentrations in the global oceans (Wyatt et al., 2014).~~

From below the mixed layer, the vertical dissolved TE fluxes range from ~~0.1~~ to $1.2 \text{ nmol Co m}^{-2} \text{ d}^{-1}$, from ~~6~~ to ~~9~~ $\text{nmol Fe m}^{-2} \text{ d}^{-1}$, and from ~~5~~ to ~~7~~ $\text{nmol Zn m}^{-2} \text{ d}^{-1}$ along the 40°S transect (Table 3). These fluxes are consistent with previous estimates of

630 Co in the South Atlantic ($0.04 - 0.46 \text{ nmol m}^{-2} \text{ d}^{-1}$, Bown et al., 2011; $0.1 - 4 \text{ nmol m}^{-2} \text{ d}^{-1}$, Rigby et al., 2020) and in the high latitude North Atlantic ($0.15 - 0.5 \text{ nmol m}^{-2} \text{ d}^{-1}$, Achterberg et al., 2020), of Fe in the North Atlantic ($0.14 - 21.1 \text{ nmol m}^{-2} \text{ d}^{-1}$, Painter et al., 2014), South Atlantic ($1 - 27 \text{ nmol m}^{-2} \text{ d}^{-1}$, Rigby et al., 2020) and the Southern Ocean (~~3~~ $- 31 \text{ nmol m}^{-2} \text{ d}^{-1}$, Blain et al., 2007; $2.3 - 14 \text{ nmol m}^{-2} \text{ d}^{-1}$, Charette et al., 2007), and of Zn in the Atlantic ($2.7 - 137 \text{ nmol m}^{-2} \text{ d}^{-1}$, Rigby et al., 2020; Achterberg et al., 2020). However, the vertical diffusive Fe fluxes ~~are smaller than some other vertical fluxes estimated in the Southern Ocean ($27 - 135 \text{ nmol m}^{-2} \text{ d}^{-1}$, Dulaiova et al., 2009) and the winter mixing fluxes in the high latitude North Atlantic (e.g. $27.3 - 103 \text{ nmol m}^{-2} \text{ d}^{-1}$, Achterberg et al., 2018).~~ It is also worth mentioning that the TE fluxes estimated by Blain et al. (2007), Bown et al. (2011) and Painter et al. (2014) use the K_2 values derived from the vertical density profiles instead of ^{228}Ra .

4.4 Mass-balance budgets for trace elements in the South Atlantic

640 The mass-balance budgets of dissolved TEs from different sources (horizontal shelf inputs, vertical upward mixing, and atmospheric dust deposition) and sinks (exported fluxes) in the surface South Atlantic (40°S transect) are calculated and summarised in Table 4 and Fig. 8. The vertical upward mixing appears to be a more important source supplying TEs to the surface water at 40°S compared to atmospheric dust and continental shelf inputs. However, the dominant source or seasonal variation of the vertical TE inputs cannot be identified in this study. Apart from the internal regeneration, TEs from a

Deleted: in fact

Deleted: 7.9

Deleted: 20

Deleted: compares well with

Deleted: those

Deleted: values of shelf-ocean Fe

Deleted: in the presence of

Deleted: 5

Deleted: the likely lower boundary of the values

Deleted: T

Deleted: is

Deleted: also

Deleted: s

Deleted: 4

Deleted: 3.6

Deleted: 11

Deleted: 13

Deleted: 16

Deleted: $27 - 135 \text{ nmol m}^{-2} \text{ d}^{-1}$, Dulaiova et al., 2009;

Deleted: seem generally smaller than

Deleted: Fe

Deleted: in the high latitude North Atlantic

Deleted: ¶

Atmospheric dust deposition is another important source supplying TEs to the surface ocean. The soluble atmospheric dust deposition fluxes to the surface 40°S Atlantic are $0.02 - 0.05 \text{ nmol Co m}^{-2} \text{ d}^{-1}$, $1.6 - 5.2 \text{ nmol Fe m}^{-2} \text{ d}^{-1}$ and $0.6 - 6 \text{ nmol Zn m}^{-2} \text{ d}^{-1}$ as reported by Chance et al. (2015) from the same cruise.¶

subsurface lateral input from the continental margin can subsequently be brought to the surface by vertical mixing (e.g. Rijkenberg et al., 2014). Deep winter convective mixing has also been shown as an important source of TEs to the surface ocean (e.g. Achterberg et al., 2018; 2020; Rigby et al., 2020).

A bulk estimate of the dissolved TE exported fluxes from the surface ocean, supported by new production biological uptake, can be made using a sinking particulate organic carbon (POC) flux and an estimated TE/C uptake ratio. In this study region, previous studies have reported the ^{234}Th -derived POC fluxes ranging from 3.1 to 9.7 $\text{mmol C m}^{-2} \text{d}^{-1}$ in the top 100 m integration depth ($7.0 \pm 2.2 \text{ mmol C m}^{-2} \text{d}^{-1}$, Thomalla et al., 2006; $6.4 \pm 3.3 \text{ mmol C m}^{-2} \text{d}^{-1}$, Owens et al., 2015). The estimates of cellular TE/C ratios are 0.3 ~ 3, 10 ~ 100, and 5 ~ 10 $\mu\text{mol/mol}$ for Co/C, Fe/C and Zn/C respectively, based on the measurements in marine phytoplankton under different TE concentrations in surface waters (e.g. Co: 10^{-12} to $10^{-11} \text{ mol L}^{-1}$, Sunda and Huntsman, 1995; Fe: 10^{-8} to $10^{-7} \text{ mol L}^{-1}$, Sunda and Huntsman, 1997; Zn: 10^{-11} to $10^{-10} \text{ mol L}^{-1}$, Sunda and Huntsman, 2000). These concentrations were chosen to represent the ranges of TE concentrations found in the surface waters of this region. Multiplying the POC fluxes with the TE/C ratios, the exported fluxes of Co, Fe, and Zn are 1 ~ 29, 31 ~ 970, and 16 ~ 97 $\text{nmol m}^{-2} \text{d}^{-1}$ respectively. The results agree with the estimates of particulate TE removal fluxes in the North Atlantic (e.g. Co: 0.27 ~ 6.8 and Fe: 274 ~ 2740 $\text{nmol m}^{-2} \text{d}^{-1}$, Hayes et al., 2018), even though the N Atlantic assessments have accounted for both biological uptake and particle scavenging fluxes by directly using TE/ ^{234}Th ratios.

In general, the exported TE fluxes are higher than the net dissolved TE inputs that we have identified in this study (Table 4). Taking Fe as an example, the total dissolved Fe inputs ($22 - 45 \times 10^6 \text{ mol yr}^{-1}$) only contribute 1 ~ 56 % of the biological consumption of dissolved Fe, which is not enough to balance the iron budget in the surface ocean. This could imply that (1) the spatial and temporal variability in ^{234}Th -derived POC flux is crucial (given the mean-life of ^{234}Th is 35 days); (2) much lower TE/C ratios are required; or (3) other sources of TEs need to be considered (e.g. lateral-transport particulate TE or winter deep-mixing).

In the calculations of TE removal fluxes, we have considered a reasonable range of ^{234}Th -derived POC fluxes and the TE/C ratios, which helps to bring the lower end of the TE removal fluxes closer to the upper end of the total TE input fluxes. This may be enough to explain the offsets of Zn and Co budgets, considering the uncertainties, but it is still not enough to explain the offset of Fe. The range of observed phytoplankton TE/C ratios in the global oceans can vary widely (e.g. Co/C: 0.00047 ~ 25.6 $\mu\text{mol/mol}$; Fe/C: 2.1 ~ 258 $\mu\text{mol/mol}$; Zn/C: 0.02 ~ 110 $\mu\text{mol/mol}$, Moore et al., 2013). Direct measurements of the TE/C ratios in the suspended particles in the South Atlantic are required to constrain the removal TE fluxes. The Fe released from lateral-transport particles has been suggested as a potential source to explain the high dissolved Fe concentration observed in the upper 800 m waters in the Southwest Atlantic (Rijkenberg et al., 2014). It is not possible to evaluate this flux or to provide more discussion at this stage. Further studies are needed to understand the sources and the TE concentrations of these particles and the mechanism releasing TEs from these particles in the ocean.

Deleted: If we assume that the ^{234}Th -derived POC fluxes ranging from 3.1 to 9.7 $\text{mmol C m}^{-2} \text{d}^{-1}$ in the top 100 m integration depth, the estimates of cellular TE/C ratios are 0.3 ~ 3, 10 ~ 100, and 5 ~ 10 $\mu\text{mol/mol}$ for Co/C, Fe/C and Zn/C respectively, based on the measurements in marine phytoplankton under different TE concentrations in surface waters (e.g. Co: 10^{-12} to $10^{-11} \text{ mol L}^{-1}$, Sunda and Huntsman, 1995; Fe: 10^{-8} to $10^{-7} \text{ mol L}^{-1}$, Sunda and Huntsman, 1997; Zn: 10^{-11} to $10^{-10} \text{ mol L}^{-1}$, Sunda and Huntsman, 2000). These concentrations were chosen to represent the ranges of TE concentrations found in the surface waters of this region. Multiplying the POC fluxes with the TE/C ratios, the exported fluxes of Co, Fe, and Zn are 1 ~ 29, 31 ~ 970, and 16 ~ 97 $\text{nmol m}^{-2} \text{d}^{-1}$ respectively. The results agree with the estimates of particulate TE removal fluxes in the North Atlantic (e.g. Co: 0.27 ~ 6.8 and Fe: 274 ~ 2740 $\text{nmol m}^{-2} \text{d}^{-1}$, Hayes et al., 2018), even though the N Atlantic assessments have accounted for both biological uptake and particle scavenging fluxes by directly using TE/ ^{234}Th ratios.

Formatted ... [4]

Deleted: , These concentrations were chosen to represent the ranges of TE concentrations found in the surface waters of this region.

Deleted: tT...e exported TE fluxes are much ... higher than the net dissolved TE inputs that we have identified in this study (Table 4).

Deleted: Previous studies have shown that the ^{234}Th -derived POC fluxes can vary between 1.5 and 7 $\text{mmol C m}^{-2} \text{d}^{-1}$ in the South Atlantic subtropical gyre (Charette and Moran, 1999; Thomalla et al., 2006). The difference is only a factor of 4 to 5... which helps to bring the lower end of the TE removal fluxes closer to the upper end of the total TE input fluxes.

5 Conclusions

This study investigates the distribution of ^{226}Ra in the 40°S Atlantic and provides constraints on ocean mixing and dissolved TE fluxes (Co, Fe and Zn) to the high productivity region in the South Atlantic. Although the ^{226}Ra 1-D mixing model shows some limitations in the assumptions, the ^{226}Ra data do place bounds on maximum mixing rates in this study and the estimates are within the range of observed values in the global oceans. Three different ^{226}Ra approaches (1-D diffusion, advection and $^{226}\text{Ra}/\text{TE}$ ratio) have been applied to estimate the dissolved TE fluxes to the 40°S Atlantic, and the results are comparable to each other. The net dissolved TE fluxes suggest that vertical upward mixing is more important than atmospheric dust deposition and continental shelf supply as the main source supplying dissolved TEs to the surface 40°S Atlantic. However, considering the biological uptake, these dissolved TE inputs are generally not enough to balance the TE budgets in the surface ocean of this region, particularly for Fe. Apart from vertical upward mixing, continental shelves and atmospheric dust inputs, other TE inputs (e.g. particulate or winter deep-mixing) may need to be considered to improve our understanding of micronutrient limitations in the high productivity region in the South Atlantic.

Appendix A. Seawater ^{226}Ra and silica concentrations




Figure A1: (a) Relationship between the Atlantic seawater ^{226}Ra activity and silica concentration (GEOTRACES GA03, GEOSECS and TTO datasets: Ku and Lin, 1976; Key et al., 1990; 1992a,b; Charette et al., 2015). The dashed line shows a linear regression through all the data (b) Plots of Si-extrapolated ^{226}Ra against measured ^{226}Ra in this study and the TTO program. The paired t-test shows a p-value of 0.55 (> 0.05). The solid line is 1:1, and the purple dashed lines show the linear regressions with the error bars of ±11% (2S.E.)

Appendix B. Trace element (TE) data

Table B1 Surface water (< 10m) dissolved trace element data

Cruise	Stn	Lon	Lat	dCo (pM)	dFe (nM)	dZn (nM)
D357	Stn0.5	17.606	-34.336		0.03	
D357	Stn0.5_r	17.586	-34.307	30.7	0.15	
D357	Stn1	17.054	-34.620	42.2		
D357	Stn1_r	17.010	-34.617	33.4		
D357	Stn1.5	16.060	-34.918	10.8		
D357	Stn2.5	14.074	-35.941	7.3		
D357	Stn3	13.270	-36.466	17.4	0.06	
D357	Stn3.5	11.582	-37.457	16.1		
D357	Stn4	10.052	-38.416	16.5		
D357	Stn4.5	7.729	-39.255	4.7	0.06	
D357	Stn4.5	7.729	-39.255		0.02	
D357	FISH_1B	16.967	-34.629	21.0		
D357	FISH_2B	16.084	-34.823	16.2		
D357	FISH_3A	15.091	-35.442	21.9		
D357	FISH_3B	13.256	-36.435	8.5		
D357	FISH_3_Co	13.129	-36.499	17.0		
D357	FISH_4B	7.688	-39.302	6.5		
D357	FISH_4_Co	10.084	-38.350	17.3		
D357	FISH_10B	17.599	-34.340	19.7		
D357	FISH_12_Co	11.717	-37.366	8.4		
JC068	FISH_1001	18.128	-34.044	27.7	0.30	0.14
JC068	FISH_1002	17.921	-34.154	29.3	0.30	0.09
JC068	FISH_1003	17.726	-34.237	46.9	0.35	0.07
JC068	FISH_1004	17.452	-34.392	27.3	0.13	0.06
JC068	Stn1	17.054	-34.612	24.6	0.18	0.09
JC068	FISH_1005	17.057	-34.611	26.6	0.13	0.05
JC068	FISH_1006	16.667	-34.774	16.0	0.12	0.16
JC068	FISH_1007	16.274	-34.937	17.8	0.12	0.07
JC068	FISH_1008	15.889	-35.097	15.2	0.09	0.00
JC068	FISH_1009	15.398	-35.312	19.5	0.10	0.01
JC068	FISH_1010	14.996	-35.468	15.7	0.11	0.00
JC068	FISH_1011	14.585	-35.659	16.3	0.11	0.00
JC068	FISH_1012	14.190	-35.836	11.5	0.14	
JC068	FISH_1013	13.825	-35.990	17.5	0.11	0.04
JC068	FISH_1015	13.188	-36.302	18.9	0.12	0.01
JC068	Stn3	13.104	-36.348	29.7	0.18	
JC068	FISH_1016	12.335	-36.935	22.8	0.14	

Commented [YTH2]: Stn2 TE data have been removed from the tables.

JC068	FISH_1017	11.733	-37.395	14.6	0.13	0.10
JC068	FISH_1018	11.086	-37.890	15.2	0.16	0.01
JC068	FISH_1019	10.443	-38.368	8.1	0.16	0.03
JC068	FISH_1020	9.778	-38.603	6.0	0.13	0.04
JC068	FISH_1021	8.606	-38.992	8.3	0.15	0.04
JC068	FISH_1022	8.378	-39.069	5.1	0.20	
JC068	FISH_1023	7.705	-39.289	7.4	0.10	0.00
JC068	FISH_1024	7.116	-39.485	5.2	0.10	0.04
JC068	FISH_1114	-54.017	-35.988	83.9		
JC068	Stn24	-54.000	-36.000	65.5	1.53	0.58
JC068	FISH_1110	-53.335	-36.335	146.2	1.35	0.59
JC068	FISH_1109	-53.166	-36.522	71.4	0.50	0.25
JC068	Stn22	-53.102	-36.536	36.7	1.05	0.17
JC068	FISH_1108	-53.007	-36.595	111.4	0.44	0.16
JC068	FISH_1107	-52.850	-36.720	59.8	0.42	0.31
JC068	FISH_1106	-52.721	-36.831	47.6	0.54	0.14
JC068	FISH_1105	-52.598	-36.943	10.5	0.39	0.19
JC068	Stn21	-52.497	-37.019	43.2	0.22	0.03
JC068	Stn21	-52.427	-37.049	59.7	0.31	0.11
JC068	FISH_1103	-52.103	-37.265	70.9	0.37	0.09
JC068	FISH_1102	-51.405	-37.769	51.8	0.33	0.26
JC068	Stn20	-50.992	-38.042	19.9	0.30	0.05
JC068	FISH_1101	-51.114	-38.145	16.5	0.22	0.22
JC068	FISH_1100	-50.632	-38.208	18.4	0.20	0.29
JC068	FISH_1099	-49.954	-38.588	10.1	0.21	0.15
JC068	FISH_1098	-49.286	-38.965	17.1	0.57	
JC068	FISH_1097	-48.464	-39.422	28.2	0.62	0.24
JC068	FISH_1096	-47.540	-39.923	12.8	0.66	0.26
JC068	Stn19	-47.417	-39.992	21.7	0.21	0.10
JC068	FISH_1095	-44.824	-40.001	14.9	0.50	0.18
JC068	FISH_1094	-43.768	-40.003	12.3	0.23	0.17
JC068	Stn18	-42.417	-40.000	16.3	0.23	0.22
JC068	Stn18	-42.297	-40.000	9.2	0.23	0.25

Table B2 Upper ocean (< 600m) dTE data

Cruise	Stn	Lat	Lon	Depth (m)	dCo (pM)	dFe (nM)	dZn (nM)
D357	Stn1	17.054	-34.620	5	41.5		
D357	Stn1	17.054	-34.620	20	40.0		
D357	Stn1	17.054	-34.620	34	38.2		
D357	Stn1	17.054	-34.620	49	48.8		

D357	Stn1	17.054	-34.620	49	42.6	
D357	Stn1	17.054	-34.620	74	54.6	
D357	Stn1	17.054	-34.620	99	61.2	
D357	Stn1	17.054	-34.620	148	58.8	
D357	Stn1	17.054	-34.620	197	48.5	
D357	Stn1	17.054	-34.620	347	59.3	
D357	Stn1	17.054	-34.620	495	60.8	
D357	Stn1	17.054	-34.620	624	54.4	
D357	Stn3	13.117	-36.333	5		0.07
D357	Stn3	13.117	-36.333	10	17.7	
D357	Stn3	13.117	-36.333	23		0.06
D357	Stn3	13.117	-36.333	29	17.0	
D357	Stn3	13.117	-36.333	47		0.05
D357	Stn3	13.117	-36.333	50	17.5	
D357	Stn3	13.117	-36.333	50		0.06
D357	Stn3	13.117	-36.333	69	28.0	
D357	Stn3	13.117	-36.333	97		0.09
D357	Stn3	13.117	-36.333	99	34.1	
D357	Stn3	13.117	-36.333	119	17.2	
D357	Stn3	13.117	-36.333	196		0.14
D357	Stn3	13.117	-36.333	197	42.7	
D357	Stn3	13.117	-36.333	218	37.1	
D357	Stn3	13.117	-36.333	395		0.30
D357	Stn3	13.117	-36.333	496	53.8	
D357	Stn3	13.117	-36.333	594		0.44
D357	Stn4.5	7.800	-39.217	10		0.07
D357	Stn4.5	7.800	-39.217	20	1.7	
D357	Stn4.5	7.800	-39.217	23		0.06
D357	Stn4.5	7.800	-39.217	47		0.05
D357	Stn4.5	7.800	-39.217	50	7.7	
D357	Stn4.5	7.800	-39.217	74	3.5	
D357	Stn4.5	7.800	-39.217	97		0.07
D357	Stn4.5	7.800	-39.217	99	8.7	0.06
D357	Stn4.5	7.800	-39.217	124	8.4	
D357	Stn4.5	7.800	-39.217	149	14.1	
D357	Stn4.5	7.800	-39.217	173	16.1	
D357	Stn4.5	7.800	-39.217	197		0.12
D357	Stn4.5	7.800	-39.217	198	30.1	
D357	Stn4.5	7.800	-39.217	302	28.8	
D357	Stn4.5	7.800	-39.217	400		0.25
D357	Stn4.5	7.800	-39.217	500	36.9	

JC068	Stn1	17.054	-34.612	18	26.9	0.22	0.21
JC068	Stn1	17.054	-34.612	29	17.5	0.14	0.13
JC068	Stn1	17.054	-34.612	41	22.3	0.28	0.01
JC068	Stn1	17.054	-34.612	48	31.9	0.08	0.01
JC068	Stn1	17.054	-34.612	58	34.9	0.16	0.02
JC068	Stn1	17.054	-34.612	78	32.6	0.29	0.39
JC068	Stn1	17.054	-34.612	98	43.9	0.37	0.34
JC068	Stn1	17.054	-34.612	197	45.4	0.35	0.34
JC068	Stn1	17.054	-34.612	247	44.8	0.45	0.25
JC068	Stn1	17.054	-34.612	296	50.8	0.48	0.28
JC068	Stn1	17.054	-34.612	398	45.2	0.54	0.49
JC068	Stn3	13.104	-36.348	19	24.9	0.13	0.21
JC068	Stn3	13.104	-36.348	34	36.4	0.17	0.27
JC068	Stn3	13.104	-36.348	48	27.8	0.25	0.10
JC068	Stn3	13.104	-36.348	74	30.7	0.14	0.21
JC068	Stn3	13.104	-36.348	98	36.6	0.17	0.08
JC068	Stn3	13.104	-36.348	149	31.1		0.13
JC068	Stn3	13.104	-36.348	199	34.5	0.29	0.06
JC068	Stn3	13.104	-36.348	298	34.2	0.33	0.20
JC068	Stn3	13.104	-36.348	398	38.1	0.41	0.53
JC068	Stn3	13.104	-36.348	498	36.1	0.46	0.35

780 **Appendix C. Relationship between seawater ²²⁸Ra_{ex} and salinity in the Argentine Basin**




Figure C1: Relationship between seawater ²²⁸Ra_{ex} activity and salinity in the Argentine Basin. The dashed line shows the linear regression. The error bars of ²²⁸Ra_{ex} activity are 2S.E.

Commented [YTH3]: Fig. C1 has been updated.

Appendix D. Trace element (TE) flux calculations

(1) ²²⁸Ra-derived-diffusive TE fluxes

To calculate both lateral and vertical TE fluxes, the ²²⁸Ra-derived diffusion coefficients (K_z or K_x) are applied to Eqn. (6) with the gradient of TE concentration ($\Delta TE/\Delta x$ or Δz) over either the horizontal distance x to the coasts or the vertical depth z below the mixed layer, which can be obtained from the linear regression of horizontal and vertical TE profiles in Fig. 3 and Fig. 4 respectively. In the Argentine Basin, the cross-shelf horizontal gradients of TEs (Fig. 3a, c and e) and ²²⁸Ra-derived K_x (Fig. 5a) have been used to assess the TE fluxes cross the shelf break and the Brazil Current. In the Cape Basin, where the TE gradients (Co and Fe) from both cruises D357 and JC068 are used for comparison (Fig. 3b and d), the differences are generally less than 20% (except for the vertical gradient of Co at Stn3). As the vertical gradients and ²²⁸Ra-derived K_z in the Argentine Basin are likely to be biased by lateral inputs, the vertical TE fluxes are not assessed in the western transect here. The ²²⁸Ra-derived-diffusion TE fluxes are summarized in Table 2 (horizontal) and Table 3 (vertical).

(2) ²²⁸Ra-derived-advective TE fluxes

The advective TE fluxes after the boundary of the shelf break and the Brazil Current in the Argentine Basin can be calculated using Eqn. (7) with the net offshore advection velocity ($w = 0.6 \pm 0.3 \text{ cm s}^{-1}$) along the SAC in the Argentine Basin and the average concentrations of dissolved TEs in the initial advective waters around where the Brazil Current merges into the SAC (around Stn21, Fig. 3): $[\text{Co}]_{\text{ave-0}} = 40 \pm 21 \text{ pM}$; $[\text{Fe}]_{\text{ave-0}} = 0.36 \pm 0.13 \text{ nM}$; $[\text{Zn}]_{\text{ave-0}} = 0.12 \pm 0.07 \text{ nM}$ (1 S.D., $n = 4$). The calculated advective TE fluxes are summarised in Table 2 for comparison. As the advective signals cannot be easily separated from the mixing in the Cape Basin, the advective TE fluxes are not assessed in the eastern transect here.

(3) TE/²²⁸Ra-ratio-derived TE fluxes

The cross-shelf TE/²²⁸Ra-ratio-derived TE fluxes can be calculated using Eqn. (8) with the $\Delta TE/\Delta^{228}\text{Ra}$ ratios reported in Table D1 and the estimates of shelf ²²⁸Ra flux ($F_{228\text{Ra}}$) based on the inverse models using the global seawater ²²⁸Ra database and inventory (Kwon et al., 2014; Le Gland et al., 2017). In the South Atlantic, the average shelf ²²⁸Ra flux is $1.7 \pm 0.3 \times 10^{10} \text{ atoms m}^2 \text{ yr}^{-1}$ around the Uruguayan and South African continental margins (normalised to shelf area; Charette et al., 2016). For comparison, the flux is converted to the shelf-ocean cross sectional flux by multiplying the average continental shelf widths (Cape Basin: 80 km; Argentine Basin: 120 km) and then dividing by the water depths at the shelf break (Cape Basin: 150 m; Argentine Basin: 160 m) (Urien and Ewing, 1974; Nelson et al., 1998; Carr and Botha, 2012). The shelf length is shared between the two surfaces (the shelf horizontal plane and vertical section in Fig. 8) and cancelled out during the calculation (Table D2). The cross-shelf ²²⁸Ra flux ($F_{228\text{Ra}}$) becomes $1.3 \pm 0.2 \times 10^{13} \text{ atoms m}^2 \text{ yr}^{-1}$ in the Argentine Basin and $0.9 \pm 0.2 \times 10^{13} \text{ atoms m}^2 \text{ yr}^{-1}$ in the Cape Basin. The calculated TE fluxes are summarised in Table 2.

It is important to remember that when considering the sources of ²²⁸Ra and TEs in the ocean, TEs may have their maximum source term at a different depth than ²²⁸Ra. Whereas ²²⁸Ra has a clear maximum from the continental shelf in the surface

Deleted: In this study, we use three different ²²⁸Ra approaches to quantify the horizontal and diffusive vertical dTE fluxes in the Cape Basin and Argentine.

Deleted: Fick's first law of molecular diffusion in the following equation: Eqn. (

Deleted: $F_{TE-a} = K_x \text{ or } z \left(\frac{\Delta TE}{\Delta x \text{ or } \Delta z} \right)$ (D1)
where F_{TE-a} is the diffusive flux of the TEs, and $\Delta TE/\Delta x$ or Δz is

Deleted: ,

Deleted: ,

Deleted: As discussed above (in Section 4.1.2), surface water after the boundary of the shelf break and the Brazil Current in the Argentine Basin carries strong offshore advection signals along the SAC towards the open ocean. Assuming that the mixing of TE is conservative, tT

Deleted: the following equation: $F_{TE-a} = w \cdot [TE]_{\text{ave-0}}$ (D2)
where F_{TE-a} is the offshore advective flux of TE

Deleted: ,

Deleted: w is t

Deleted: ,

Deleted: $[TE]_{\text{ave-0}}$ is

Deleted: Previous studies have combined the use of the shelf ²²⁸Ra fluxes with the ratios of TE/²²⁸Ra in the surface waters between continental shelves and open oceans to estimate the shelf-ocean TE inputs from the continental margins to the open oceans (Charette et al., 2016; Sanial et al., 2018; Vieira et al., 2020). This method provides the integrated net fluxes of TEs, considering all the possible inputs (e.g. rivers, SGD and sediments) and outputs (e.g. particle scavenging, biological uptake and radioactive decay) of ²²⁸Ra and TEs during water mixing between the continental shelf and the open ocean. More details of the method are given in Charette et al. 2016. In brief, assuming that the net shelf-ocean exchange is mainly driven by eddy diffusion, tT

Formatted: Superscript

Deleted: the following equation

Deleted: $F_{TE} = F_{228\text{Ra}} \cdot \left(\frac{\Delta TE}{\Delta^{228}\text{Ra}} \right) = F_{228\text{Ra}} \cdot \left(\frac{TE_{\text{shelf}} - TE_{\text{ocean}}}{^{228}\text{Ra}_{\text{shelf}} - ^{228}\text{Ra}_{\text{ocean}}} \right)$ (D3)

where $F_{228\text{Ra}}$ is the cross-shelf ²²⁸Ra flux; TE_{shelf} and $^{228}\text{Ra}_{\text{shelf}}$ are the average concentrations of the TE and ²²⁸Ra in the surface waters on the shelf (GA10W: between Stn23 and Stn25; GA10E: between Stn0 and Stn1) respectively; and TE_{ocean} and $^{228}\text{Ra}_{\text{ocean}}$ are the average concentrations in the open ocean (GA10W: between Stn18 and Stn19; GA10E: between Stn4 and Stn4.5). The ratios of $\Delta TE/\Delta^{228}\text{Ra}$ are reported in Table D1. The shelf ²²⁸Ra flux is

Deleted: Fig. 8 and

mixed layer, redox-sensitive, more particle-bound or hydrothermal-related TEs may see a maximum at deeper levels, due to particle resuspension, low oxygen saturation or hydrothermal activity. In this sense, our model calculations provide the estimates of lateral TE inputs in the surface mixed layer with a clear source from the continental shelf. The vertical TE inputs, however, do not separate the TE inputs between internal cycling, hydrothermal or lateral transport from the continental margin at greater depths. Nevertheless, all these inputs would only become relevant for productivity if they reach the surface mixed layer later, e.g. by vertical mixing as quantified here.

Table D1 Shelf and open ocean average dTE and ²²⁸Ra concentrations and ΔTE/Δ²²⁸Ra ratios^a

		²²⁸ Ra (10 ⁵ atoms L ⁻¹)	dCo (pmol L ⁻¹)	dFe (nmol L ⁻¹)	dZn (nmol L ⁻¹)
Argentina	Shelf	6.4 ± 1.1	99 ± 42	1.44 ± 0.12	0.58 ± 0.01
	Open ocean	1.3 ± 0.5	15 ± 5	0.28 ± 0.12	0.18 ± 0.05
Cape	Shelf	1.3 ± 0.1	30 ± 8	0.22 ± 0.09	0.08 ± 0.03
	Open ocean	0.5 ± 0.1	9 ± 5	0.11 ± 0.06	0.05 ± 0.05
ΔTE/Δ ²²⁸ Ra (10 ⁻⁷ nmol atom ⁻¹)					
Argentina margin			1.6 ± 1.1	22 ± 14	7.7 ± 3.9
Cape margin			2.5 ± 1.5	12 ± 8	3.6 ± 4.2

^aOnly JC068 TE data are used. All errors are ±1S.D.

Table D2 Average shelf width and shelf-break water depth for shelf-ocean dTE flux normalisation^a

Location	Shelf width (km)	Water depth (m)	References
South African margin (Cape Basin)	80	150	Nelson et al., 1998; Carr and Botha, 2012
Uruguayan margin (Argentine Basin)	120	160	Urien and Ewing, 1974
Western North Atlantic margin	135	132	Emery, 1966
Peruvian margin	100	200	Hooker et al., 2013
	Cross section width (km)	Mixed layer depth (m)	
Congo River margin*	300	15	Vieira et al., 2020
Brazilian margin**	240	10	Windom et al., 2006

^aShelf-ocean TE or ²²⁸Ra fluxes presented in this study are normalised to the area of shelf-ocean cross section (by default, the cross section at shelf-break = shelf length x shelf-break water depth). To convert the shelf TE or ²²⁸Ra flux (usually normalised by shelf area) from previous studies, the shelf fluxes are multiplied by the shelf width and length, and then divided by the area of cross section. Shelf length should drop off from the flux conversion. *Congo River margin TE fluxes are divided by a defined cross section (the width of river plume x mixed layer depth). **Brazilian margin Fe flux is divided by a defined cross section (the width of a defined coastline x mixed layer depth).

870

Data availability. All the original and supporting data are shown in the manuscript. The data are also available publicly at the GEOTRACES IDP2017 (<https://www.bodc.ac.uk/geotraces/data/idp2017/>).

Author contributions. YTH, WG and GMH designed the radium projects. YTH conducted the Ra-228 and Ra-226 analyses. EMSW conducted the Si measurements. NJW, MCL and EPA contributed the TE data and interpretation. YTH prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors wish to thank the officers and crew of *RRS Discovery* and *RRS James Cook* for their assistance on the UK-GEOTRACES GA10 cruises (D357 and JC068). We would like to thank Alex Thomas, Andrew Mason and Steve Wyatt for assistance with mass spectrometry and laboratory support. We would also like to thank Willard Moore, Will Homoky, Yves Plancherel and Christian Schlosser for feedback and discussion at different stages of the manuscript preparation. This study was funded by grants from the UK Natural Environment Research Council for the UK-GEOTRACES GA10 cruises NE/H008497/1 to WG, NE/F017316/1 to GMH, NE/H004475/1 to MCL and NJW, and NE/H004394/1 to EPA.

References

885 Achterberg, E.P., Steigenberger, S., Marsay, C.M., LeMoigne, F.A.C., Painter, S.C., Baker, A.R., et al.: Iron biogeochemistry in the high latitude North Atlantic Ocean, Scientific Reports, 8, 1283, <https://doi.org/10.1038/s41598-018-19472-1>, 2018.

Achterberg, E.P., Steigenberger, S., Klar, J.K., Browning, T.J., Marsay, C.M., Painter, S.C., Vieira, L., Baker, A.R., Hamilton, D.S., Tanhua, T., Moore, C.M.: Trace element biogeochemistry in the high latitude North Atlantic Ocean; seasonal variations and volcanic inputs, Global Biogeochemical Cycles, <https://doi.org/10.1029/2020GB006674>, 2020.

890 Beal, L.M., De Ruijter, W.P.M., Biastoch, A., Zahn, R. and 136, S.W.I.W.G.: On the role of the Agulhas system in ocean circulation and climate, Nature, 472, 429-436, <https://doi.org/10.1038/nature09983>, 2011.

Blain, S., Quéguiner, B., Armand, L. et al.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, Nature, 446, 1070-1074, <https://doi.org/10.1038/nature05700>, 2007.

895 Bown J., Boye, M., Baker, A., Duvieilbourg, E., Lacan, F., Le Moigne, F., Planchon, F., Speich, S., Nelson, D.M.: The biogeochemical cycle of dissolved cobalt in the Atlantic and the Southern Ocean south off the coast of South Africa, Marine Chemistry, 126, 193-206, <https://doi.org/10.1016/j.marchem.2011.03.008>, 2011.

Broecker, W.S., Goddard, J. and Sarmiento, J.L.: Distribution of Ra-226 in Atlantic Ocean, Earth and Planetary Science Letters, 32, 220-235, [https://doi.org/10.1016/0012-821X\(76\)90063-7](https://doi.org/10.1016/0012-821X(76)90063-7), 1976.

Deleted: , in press

Field Code Changed

Formatted: Font: Not Bold

- Browning, T.J., Bouman, H.A., Moore, C.M., Schlosser, C., Tarran, G.A., Woodward, E.M.S., and Henderson, G.M.: Nutrient regimes control phytoplankton ecophysiology in the South Atlantic, *Biogeosciences*, 11, 463-479, <https://doi.org/10.5194/bg-11-463-2014>, 2014.
- 905 Browning, T.J., Achterberg, E.P., Yong, J.C., Rapp, I., Utermann, C., and Moore, C.M.: Iron limitation of microbial phosphorus acquisition in the tropical North Atlantic, *Nature Communications*, 8, 15465, <https://doi.org/10.1038/ncomms15465>, 2017.
- Bruland, K.W., and Franks, R.P.: Mn, Ni, Cu, Zn and Cd in the Western North Atlantic, in: *Trace Metals in Sea Water*, edited by Wong, C.S., Boyle, E., Bruland, K.W., Burton, J.D., Goldberg, E.D., Springer, 395-414, https://doi.org/10.1007/978-1-4757-6864-0_23, 1983.
- 910 Cai, P., Huang, Y., Chen, M., Guo, L., Liu, G., and Qiu, Y.: New production based on Ra-228-derived nutrient budgets and thorium-estimated POC export at the intercalibration station in the South China Sea, *Deep-Sea Research Part I-Oceanographic Research Papers*, 49, 53-66, [https://doi.org/10.1016/S0967-0637\(01\)00040-1](https://doi.org/10.1016/S0967-0637(01)00040-1), 2002.
- Carr, A.S. and Botha, G.A.: Coastal Geomorphology, in: *Southern African Geomorphology: Recent Trends and New Directions*, edited by Holmes, P. and Meadows, M. E., Sun Press, Bloemfontein, 269-303, 2012.
- 915 Chance, R., Jickells, T.D., and Baker, A.R.: Atmospheric trace metal concentrations, solubility and deposition fluxes in remote marine air over the south-east Atlantic, *Marine Chemistry*, 177, 45-56, <https://doi.org/10.1016/j.marchem.2015.06.028>, 2015.
- Charette, M.A., Gonneea, M.E., Morris, P.J., Statham, P., Fones, G., Planquette, H., Salter, I., and Garabato, A.N.: Radium isotopes as tracers of iron sources fueling a Southern Ocean phytoplankton bloom, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 54, 1989-1998, <https://doi.org/10.1016/j.dsr2.2007.06.003>, 2007.
- 920 Charette, M.A., Morris, P.J., Henderson, P.B., and Moore, W.S.: Radium isotope distributions during the US GEOTRACES North Atlantic cruises, *Marine Chemistry*, 177, 184-195, <https://doi.org/10.1016/j.marchem.2015.01.001>, 2015.
- Charette, M.A., Lam, P.J., Lohan, M.C., Kwon, E.Y., Hatje, V., Jeandel, C., Shiller, A.M., Cutter, G.A., Thomas, A., Boyd, P.W., Homoky, W.B., Milne, A., Thomas, H., Andersson, P.S., Porcelli, D., Tanaka, T., Geibert, W., Dehairs, F., and
- 925 Garcia-Orellana, J.: Coastal ocean and shelf-sea biogeochemical cycling of trace elements and isotopes: lessons learned from GEOTRACES, *Phil. Trans. R. Soc. A*, 374, 20160076, <https://doi.org/10.1098/rsta.2016.0076>, 2016.
- Chever, F., Bucciarelli, E., Sarthou, G., Speich, S., Arhan, M., Penven, P., Tagliabue, A.: Physical speciation of iron in the Atlantic sector of the Southern Ocean along a transect from the subtropical domain to the Weddell Sea Gyre, *Journal of Geophysical Research-Oceans*, 115, <https://doi.org/10.1029/2009JC005880>, 2010.
- 930 Clough, R., Floor, G.H., Quétel, C.R., Milne, A., Lohan, M.C., Worsfold, P.J.: Measurement uncertainty associated with shipboard sample collection and filtration for the determination of the concentration of iron in seawater, *Anal. Methods*, 8, 6711-6719, <https://doi.org/10.1039/C6AY01551D>, 2016.

Deleted: Charette, M.A. and Moran, S.B.: Rates of particle scavenging and particulate organic carbon export estimated using ²³⁴Th as a tracer in the subtropical and equatorial Atlantic Ocean, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 46, 885-906, [https://doi.org/10.1016/S0967-0645\(99\)00006-5](https://doi.org/10.1016/S0967-0645(99)00006-5), 1999.

- Cochran, J.K.: The oceanic chemistry of the uranium- and thorium-series nuclides, in: Uranium-Series Disequilibrium: Applications to Earth, Marine, and Environmental Sciences, edited by Ivanovich, M. and Harmon R.S., Clarendon Press, Oxford, 334-391, 1992.
- Cutter, G., Andersson, P., Codispoti, L., Croot, P., Francois, R., Lohan, M.C., Obata, H. and Rutgers van der Loeff, M.: Sampling and sample-handling protocols for GEOTRACES cruises, 2010, <http://www.geotraces.org/libraries/documents/Intercalibration/Cookbook.pdf>
- Dulaiova, H., Ardelan, M.V., Henderson, P.B., and Charette, M.A.: Shelf-derived iron inputs drive biological productivity in the southern Drake Passage, *Global Biogeochem. Cycles*, 23, GB4014, <https://doi.org/10.1029/2008GB003406>, 2009.
- Emery, K.O.: Atlantic continental shelf and slope of the United States: U.S. Geological Survey Professional Paper 529-A, 1-23, 1966.
- Foster, D.A., Staubwasser, M. and Henderson, G.M.: Ra-226 and Ba concentrations in the Ross Sea measured with multicollector ICP mass spectrometry, *Marine Chemistry*, 87, 59-71, <https://doi.org/10.1016/j.marchem.2004.02.003>, 2004.
- Gaiero, D.M., Simonella, L., Gassó, S., Gili, S., Stein, A.F., Sosa, P., Becchio, R., Arce, J. and Marelli H.: Ground/satellite observations and atmospheric modeling of dust storms originating in the high Puna- Altiplano deserts (South America), implications for the interpretation of paleo-climatic archives, *Journal of Geophysical Research, Atmospheres*, 118, 3817-3831, <https://doi.org/10.1002/jgrd.50036>, 2013.
- Geibert, W., Rodellas, V., Annett, A., van Beek, P., Garcia-Orellana, J., Hsieh, Y.-T., Masque, P.: 226Ra determination via the rate of 222Rn ingrowth with the Radium Delayed Coincidence Counter (RaDeCC), *Limnology and Oceanography Methods*, 11, 594-603, <https://doi.org/10.4319/lom.2013.11.594>, 2013.
- Graham, R.M., De Boer, A.M., van Sebille, E., Kohfeld, K.E., and Schlosser, C.: Inferring source regions and supply mechanisms of iron in the Southern Ocean from satellite chlorophyll data, *Deep Sea Res. Part I*, 104, 9-25, <https://doi.org/10.1016/j.dsr.2015.05.007>, 2015.
- Hanfland, C.: Radium-226 and Radium-228 in the Atlantic sector of the Southern Ocean, Ph.D. thesis, Alfred Wegener Institut für Polar und Meeresforschung, Bremerhaven, Germany, 135 pp., 2002.
- Hayes, C.T., Black, E.E., Anderson, R.F., Baskaran, M., Buesseler, K.O., Charette, M.A., Cheng, H., Cochran, J.K., Edwards, R.L., Fitzgerald, P., Lam, P.J., Lu, Y., Morris, S.O., Ohnemus, D.C., Pavia, F.J., Stewart, G., and Tang, Y.: Flux of Particulate Elements in the North Atlantic Ocean Constrained by Multiple Radionuclides, *Global Biogeochemical Cycles*, 32, 1738-1758, 2018.
- Ho, T.-Y., Quigg, A., Finkel, Z.V., Milligan, A.J., Wyman, K., Falkowski, P.G., and Morel, F.M.M.: The elemental composition of some marine phytoplankton, *Journal of Phycology*, 39, 1145-1159, <https://doi.org/10.1111/j.0022-3646.2003.03-090.x>, 2003.
- Hawco, N.J., Ohnemus, D.C., Resing, J.A., Twining, B.S., and Saito, M.A.: A dissolved cobalt plume in the oxygen minimum zone of the eastern tropical South Pacific, *Biogeosciences*, 13, 5697-5716, <https://doi.org/10.5194/bg-13-5697-2016>, 2016.

Deleted: Deng, F., Thomas, A.L., Rijkenberg, M.J.A., and Henderson, G.M.: Controls on seawater ²³¹Pa, ²³⁰Th and ²³²Th concentrations along the flow paths of deep waters in the Southwest Atlantic, *Earth and Planetary Science Letters*, 390, 93-102, <https://doi.org/10.1016/j.epsl.2013.12.038>, 2014.

Field Code Changed

- Homoky, W., John, S.G., Conway, T.M., and Mills, R.A.: Distinct iron isotopic signatures and supply from marine sediment dissolution, *Nat. Commun.*, 4, 2143, <https://doi.org/10.1038/ncomms3143>, 2013.
- Hooker, Y., Prieto-Rios, E., and Solis-Marín, F.A.: Echinoderms of Peru, in: *Echinoderm research and diversity in Latin America*, edited by Alvarado-Barrientos, J.J. and Solis-Marín, F.A., Springer-Verlag, Berlin, 277–299, 2013.
- Hsieh, Y.-T. and Henderson, G.M.: Precise measurement of $^{228}\text{Ra}/^{226}\text{Ra}$ ratios and Ra concentrations in seawater samples by multi-collector ICP mass spectrometry, *Journal of Analytical Atomic Spectrometry*, 26, 1338–1346, <https://doi.org/10.1039/C1JA10013K>, 2011.
- Jenkins, W.J.: Nitrate Flux into the Euphotic Zone near Bermuda. *Nature*, 331, 521–523, <https://doi.org/10.1038/331521a0>, 1988.
- Kadko, D., Aguilar-Islas, A., Buck, C.S., Fitzsimmons, J.N., Landing, W.M., Shiller, A., Till, C.P., Bruland, K.W., Boyle, E.A. and Anderson, R.F.: Sources, fluxes and residence times of trace elements measured during the U.S. GEOTRACES East Pacific Zonal Transect, *Marine Chemistry*, 222, 103781, <https://doi.org/10.1016/j.marchem.2020.103781>, 2020.
- Kaufman, A., Trier, R., Broecker, W.S. and Feely, H.W.: Distribution of ^{228}Ra in the world ocean, *Journal of Geophysical Research*, 78, 8827–8848, <https://doi.org/10.1029/JC078i036p08827>, 1973.
- Key, R.M., Rotter, R.J., McDonald, G.J., and Slater, R.D.: Western Boundary Exchange Experiment Final Data Report for large volume samples ^{228}Ra , ^{226}Ra , ^9Be , and ^{10}Be Results; Technical Report 90-1, Ocean Tracer Laboratory, Dept. Geology and Geophysics, Princeton University, Princeton, 1990.
- Key, R.M., Moore, W.S., and Sarmiento, J.L.: Transient Tracers in the Ocean North Atlantic Study Final Data Report for ^{228}Ra and ^{226}Ra ; Technical Report 92-2, Ocean Tracer Laboratory, Dept. Geology and Geophysics, Princeton University, Princeton, 1992a.
- Key, R.M., Moore, W.S., and Sarmiento, J.L. Transient Tracers in the Ocean Tropical Atlantic Study Final Data Report for ^{228}Ra and ^{226}Ra , Technical Report 92-3, Ocean Tracer Laboratory, Dept. Geology and Geophysics, Princeton University, Princeton, 1992b.
- Kipp, L.E., Charette, M.A., Moore, W.S., Henderson, P.B., and Rigor, I.G.: Increased fluxes of shelf-derived materials to the central Arctic Ocean, *Science Advances*, 4(1), eaao1302, DOI: 10.1126/sciadv.aao1302, 2018a.
- Kipp, L.E., Sanial, V., Henderson, P.B., van Beek, P., Reyss, J.-L., Hammond, D.E., Moore, W.S., and Charette M.A.: Radium isotopes as tracers of hydrothermal inputs and neutrally buoyant plume dynamics in the deep ocean, *Marine Chemistry*, 201, 51–65, <https://doi.org/10.1016/j.marchem.2017.06.011>, 2018b.
- Knauss, K.G., Ku, T.L. and Moore, W.S.: Radium and Thorium Isotopes in Surface Waters of East Pacific and Coastal Southern-California, *Earth and Planetary Science Letters*, 39, 235–249, [https://doi.org/10.1016/0012-821X\(78\)90199-1](https://doi.org/10.1016/0012-821X(78)90199-1), 1978.
- Ku, T.L. and Lin, M.C.: Ra-226 Distribution in Antarctic Ocean, *Earth and Planetary Science Letters*, 32, 236–248, [https://doi.org/10.1016/0012-821X\(76\)90064-9](https://doi.org/10.1016/0012-821X(76)90064-9), 1976.

- 1010 Ku, T.L. and Luo, S.: Ocean circulation/mixing studies with decay-series isotopes, in: U-Th Series Nuclides in Aquatic Systems, edited by Krishnaswami, S. and Cochran, J.K., Elsevier, 307-344, 2008.
- Ku, T.L., Luo, S.D., Kusakabe, M. and Bishop, J.K.B.: Ra-228-Derived Nutrient Budgets in the Upper Equatorial Pacific and the Role of New Silicate in Limiting Productivity, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 42, 479-497, [https://doi.org/10.1016/0967-0645\(95\)00020-Q](https://doi.org/10.1016/0967-0645(95)00020-Q), 1995.
- 1015 Kunze, E., and Sanford, T.B.: Abyssal mixing: Where it is not, *J. Phys. Oceanogr.*, 26, 2286-2296, [https://doi.org/10.1175/1520-0485\(1996\)026<2286:AMWIIN>2.0.CO;2](https://doi.org/10.1175/1520-0485(1996)026<2286:AMWIIN>2.0.CO;2), 1996.
- Kwon, E.Y., Kim, G., Primeau, F., Moore, W.S., Cho, H.-M., DeVries, T., Sarmiento, J.L., Charette, M.A., and Cho, Y.-K.: Global estimate of submarine groundwater discharge based on an observationally constrained radium isotope model, *Geophys. Res. Lett.*, 41, 8438-8444, <https://doi.org/10.1002/2014GL061574>, 2014.
- 1020 Lamontagne, S. and Webster, I.T.: Theoretical assessment of the effect of vertical dispersivity on coastal seawater radium distribution, *Front. Mar. Sci.*, 6, 357, <https://doi.org/10.3389/fmars.2019.00357>, 2019.
- Le Gland, G., Mémery, L., Aumont, O., and Resplandy, L.: Improving the inverse modeling of a trace isotope: how precisely can radium-228 fluxes toward the ocean and submarine groundwater discharge be estimated? *Biogeosciences*, 14, 3171-3189, <https://doi.org/10.5194/bg-14-3171-2017>, 2017.
- 1025 Ledwell, J.R., Watson, A.J., and Law, C.S.: Evidence for slow mixing across the pycnocline from an open-ocean tracer-release experiment, *Nature*, 364, 701-703, <https://doi.org/10.1038/364701a0>, 1993.
- Li, Y.H., Peng, T.H., Broecker, W.S. and Ostlund, H.G.: The Average Vertical Mixing Coefficient for the Oceanic Thermocline, *Tellus Series B-Chemical and Physical Meteorology*, 36, 212-217, <https://doi.org/10.3402/tellusb.v36i3.14905>, 1984.
- 1030 Liang, X., Spall, M., and Wunsch, C.: Global ocean vertical velocity from a dynamically consistent ocean state estimate, *Journal of Geophysical Research: Oceans*, 122, 8208-8224, <https://doi.org/10.1002/2017JC012985>, 2017.
- Lohan, M.C. and Tagliabue A.: Oceanic Micronutrients: trace metals that are essential for marine life. *Elements*, 14, 385-90, <https://doi.org/10.2138/gselements.14.6.385>, 2018.
- Longhurst, A.R.: *Ecological Geography of the Sea*, Academic Press, <https://doi.org/10.1016/B978-0-12-455521-1.X5000-1>, 2007.
- 1035 Martin, A.P., Lucas, M.I., Painter, S.C., Pidcock, R., Prandke, H., Prandke, H., and Stinchcombe, M.C.: The supply of nutrients due to vertical turbulent mixing: a study at the Porcupine Abyssal Plain study site (49°50' N 16°30' W) in the Northeast Atlantic, *Deep-Sea Res. Pt. II*, 57, 1293–1302, <https://doi.org/10.1016/j.dsr2.2010.01.006>, 2010.
- Menzel Barraqueta, J.L., Klar, J.K., Gledhill, M., Schlosser, C., Shelley, R., Planquette, H.F., Wenzel, B., Sarthou, G., and
- 1040 Achterberg, E.P.: Atmospheric deposition fluxes over the Atlantic Ocean: a GEOTRACES case study, *Biogeosciences*, 16, 1525-1542, <https://doi.org/10.5194/bg-16-1525-2019>, 2019.
- Moore, C.M., Mills, M.M., Arrigo, K.R., Berman-Frank, I., Bopp, L., Boyd, P.W., Galbraith, E.D., Geider, R.J., Guieu, C., Jaccard, S.L., Jickells, T.D., La Roche, J., Lenton, T.M., Mahowald, N.M., Earanon, E., Marinov, I., Moore, J.K.,

- Nakatsuka, T., Oschlies, A., Saito, M.A., Thingstad, T.F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nature Geoscience*, 6, 701-710, <https://doi.org/10.1038/ngeo1765>, 2013.
- Moore, J.K., Doney, S.C. and Lindsay, K.: Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model. *Global Biogeochemical Cycles*, 18, GB4028, <https://doi.org/10.1029/2004GB002220>, 2004.
- Moore, W.S. and Arnold, R.: Measurement of Ra-223 and Ra-224 in coastal waters using a delayed coincidence counter, *Journal of Geophysical Research-Oceans*, 101, 1321-1329, <https://doi.org/10.1029/95JC03139>, 1996.
- Moore, W.S. and Dymond, J.: Fluxes of Ra-226 and Barium in the Pacific-Ocean - the Importance of Boundary Processes, *Earth and Planetary Science Letters*, 107, 55-68, [https://doi.org/10.1016/0012-821X\(91\)90043-H](https://doi.org/10.1016/0012-821X(91)90043-H), 1991.
- Moore, W.S., Key, R.M. and Sarmiento, J.L.: Techniques for Precise Mapping of Ra-226 and Ra-228 in the Ocean, *Journal of Geophysical Research-Oceans*, 90, 6983-6994, <https://doi.org/10.1029/JC090iC04p06983>, 1985.
- Moore, W.S.: Determining coastal mixing rates using radium isotopes, *Cont. Shelf Res.*, 20, 1993-2007, [https://doi.org/10.1016/S0278-4343\(00\)00054-6](https://doi.org/10.1016/S0278-4343(00)00054-6), 2000.
- Moore, W.S., Sarmiento, J.L., and Key, R.M.: Submarine groundwater discharge revealed by ^{228}Ra distribution in the upper Atlantic Ocean, *Nature Geosciences*, 1, 309-311, <https://doi.org/10.1038/ngeo183>, 2008.
- Moore, W.S.: Inappropriate attempts to use distributions of ^{228}Ra and ^{226}Ra in coastal waters to model mixing and advection rates, *Cont. Shelf Res.*, 105, 95-100, <https://doi.org/10.1016/j.csr.2015.05.014>, 2015.
- Morel, F.M.M. and Price, N.M.: The biogeochemical cycles of trace metals in the oceans, *Science*, 300, 944-947, DOI: 10.1126/science.1083545, 2003.
- Nelson, G., Boyd, A.J., Agenbag, J.J., and Duncombe Rae, C.M.: An upwelling filament north-west of Cape Town, South Africa, *South African Journal of Marine Science*, 19, 75-88, <https://doi.org/10.2989/025776198784126953>, 1998.
- Noble, A.E., Lamborg, C.H., Ohnemos, D.C., Lam, P.J., Goepfert, T.J., Measures, C.I., Frame, C.H., Casciotti, K.L., DiTullio, G.R., Jennings, J., and Saito, M.A.: Basin-scale inputs of cobalt, iron, and manganese from the Benguela-Angola front to the South Atlantic Ocean, *Limnology and Oceanography*, 57, 989-1010, <https://doi.org/10.4319/lo.2012.57.4.0989>, 2012.
- Noble, A.E., Ohnemos, D.C., Hawco, N.J., Lam, P.J., and Saito, M.A.: Coastal sources, sinks and strong organic complexation of dissolved cobalt within the US North Atlantic GEOTRACES transect GA03, *Biogeosciences*, 14, 2715-2739, <https://doi.org/10.5194/bg-14-2715-2017>, 2017.
- Nozaki, Y. and Yamamoto, Y.: Radium 228 based nitrate fluxes in the eastern Indian Ocean and the South China Sea and a silicon-induced "alkalinity pump" hypothesis, *Global Biogeochemical Cycles*, 15, 555-567, <https://doi.org/10.1029/2000GB001309>, 2001.
- Oschlies, A.: Nutrient supply to the surface waters of the North Atlantic: A model study, *Journal of Geophysical Research-Oceans*, 107(C5), <https://doi.org/10.1029/2000JC000275>, 2002.
- Owens, S.A., Pike, S., and Buesseler, K.O.: Thorium-234 as a tracer of particle dynamics and upper ocean export in the Atlantic Ocean, *Deep Sea Research, Part II*, 116, 42-59, <https://doi.org/10.1016/j.dsr2.2014.11.010>, 2015.

Painter, S.C., Henson, S.A., Forryan, A., Steigenberger, S., Klar, J., Stinchcombe, M.C., Rogan, N., Baker, A.R.,
 1080 Achterberg, E.P., and Moore, C.M.: An assessment of the vertical diffusive flux of iron and other nutrients to the surface
 waters of the subpolar North Atlantic Ocean, *Biogeosciences*, 11, 2113-2130, <https://doi.org/10.5194/bg-11-2113-2014>,
 2014.

Paul, M., Van De Flierdt, T., Rehkämper, M., Khondoker, R. Weiss, D., Lohan, M.C., Homoky, W.B.: Tracing the Agulhas
 leakage with lead isotopes, *Geophys. Res. Lett.*, 42, 8515-8521, <https://doi.org/10.1002/2015GL065625>, 2015.

1085 [Price, N.M. and Morel F.M.M.: Cadmium and cobalt substitution for zinc in a marine diatom, *Nature*, 344, 658-660, 1990.](#)

Reid, D.F., Key, R.M. and Schink, D.R.: Radium, Thorium and Actinium extraction from seawater using an improved
 manganese-oxide-coated fiber, *Earth and Planetary Science Letters*, 43: 223-226, [https://doi.org/10.1016/0012-](https://doi.org/10.1016/0012-821X(79)90205-X)
[821X\(79\)90205-X](https://doi.org/10.1016/0012-821X(79)90205-X), 1979.

Rigby, S.J., Williams, R.G., Achterberg, E.P., and Tagliabue, A.: Resource availability and entrainment are driven by offsets
 between nutriclines and winter mixed-layer depth, *Global Biogeochemical Cycles*, 34, e2019GB006497,
 1090 <https://doi.org/10.1029/2019GB006497>, 2020.

Rijkenberg, M.J.A., Middag, R., Laan, P., Gerringa, L.J.A., van Aken, H.M., Schoemann, V., de Jong, J.T.M., and de Baar,
 H.J.W.: The distribution of dissolved iron in the west Atlantic Ocean, *PLoS One*, 9, e101323,
<https://doi.org/10.1371/journal.pone.0101323>, 2014.

Rodellas, V., Garcia-Orellana, J., Masqué, P., Feldman, M., and Weinstein, Y.: Submarine groundwater discharge as a major
 1095 source of nutrients to the Mediterranean Sea, *Proc. Natl. Acad. Sci.*, 12, 3926-3930,
<https://doi.org/10.1073/pnas.1419049112>, 2015.

Rutgers van der Loeff, M. M., Key, R. M., Scholten, J., Bauch, D. and Michel, A.: 228Ra as a tracer for shelf water in the
 arctic ocean, *Deep. Res. Part II*, 42, 1533–1553, [https://doi.org/10.1016/0967-0645\(95\)00053-4](https://doi.org/10.1016/0967-0645(95)00053-4), 1995.

Saito, M.A., Noble, A.E., Tagliabue, A., Goepfert, T.J., Lamborg, C.H., and Jenkins, W.J.: Slow-spreading submarine ridges
 1100 in the South Atlantic as a significant oceanic iron source, *Nature Geoscience*, 6, 775-779, <https://doi.org/10.1038/ngeo1893>,
 2013.

Sanial, V., Kipp, L.E., Henderson, P.B., van Beek, P., Reyss, J.-L., Hammond, D.E., Hawco, N., Saito, M.A., Resing, J.A.,
 Sedwick, P.N., Moore, W.S., and Charette, M.A.: Radium-228 as a tracer of dissolved trace element inputs from the
 Peruvian continental margin, *Marine Chemistry*, 201, 20-34, <https://doi.org/10.1016/j.marchem.2017.05.008>, 2018.

1105 Sarmiento, J.L., Feely, H.W., Moore, W.S., Bainbridge, A.E. and Broecker, W.S.: Relationship between Vertical Eddy
 Diffusion and Buoyancy Gradient in Deep-Sea, *Earth and Planetary Science Letters*, 32, 357-370,
[https://doi.org/10.1016/0012-821X\(76\)90076-5](https://doi.org/10.1016/0012-821X(76)90076-5), 1976.

Sarmiento, J.L., Thiele, G., Key, R.M. and Moore, W.S.: Oxygen and Nitrate New Production and Remineralization in the
 North-Atlantic Subtropical Gyre, *Journal of Geophysical Research-Oceans*, 95, 18303-18315,
 1110 <https://doi.org/10.1029/JC095iC10p18303>, 1990.

Deleted: V

Deleted: D

- Schlitzer, R.: Mass and Heat Transport in the South Atlantic Derived from Historical Hydrographic Data, in: The South Atlantic, Springer, Berlin, Heidelberg, 305-323, https://doi.org/10.1007/978-3-642-80353-6_17, 1996.
- 1115 Schlitzer, R.: Quantifying He fluxes from the mantle using multi-tracer data assimilation, Phil. Trans. R. Soc. A, 374, 20150288, <https://doi.org/10.1098/rsta.2015.0288>, 2016.
- Schlitzer, R., Anderson, R.F., Masferrer Dodas, E., et al.: The GEOTRACES Intermediate Data Product 2017, Chemical Geology, 493, 210-223, <https://doi.org/10.1016/j.chemgeo.2018.05.040>, 2018.
- 1120 [Sunda, W.G. and Huntsman S.A.: Cobalt and zinc interreplacement in marine phytoplankton: Biological and geochemical implications, Limnology and Oceanography, 40, 1404-1417, https://doi.org/10.4319/lo.1995.40.8.1404, 1995.](#)
- [Sunda, W.G. and Huntsman S.A.: Interrelated influence of iron, light and cell size on marine phytoplankton growth, Nature, 390, 389-392, https://doi.org/10.1038/37093, 1997.](#)
- [Sunda, W.G. and Huntsman S.A.: Effect of Zn, Mn and Fe on Cd accumulation in phytoplankton: Implications for oceanic Cd cycling, Limnology and Oceanography, 45, 1501-1516, https://doi.org/10.4319/lo.2000.45.7.1501, 2000.](#)
- 1125 Tagliabue, A., Sallee, J. B., Bowie, A. R., Levy, M., Swart, S., & Boyd, P. W.: Surface-water iron supplies in the Southern Ocean sustained by deep winter mixing, Nature Geoscience, 7, 314–320, <https://doi.org/10.1038/Ngeo2101>, 2014.
- Thomalla, S., Turnewitsch, R., Lucas, M., and Poulton, A.: Particulate organic carbon export from the North and South Atlantic gyres: the ²³⁴Th/²³⁸U disequilibrium approach, Deep Sea Research Part II: Topical Studies in Oceanography, 53, 1629-1648, <https://doi.org/10.1016/j.dsr2.2006.05.018>, 2006.
- 1130 Urien C.M. and Ewing M.: Recent Sediments and Environment of Southern Brazil, Uruguay, Buenos Aires, and Rio Negro Continental Shelf, in: The Geology of Continental Margins, edited by Burk, C.A. and Drake, C.L., Springer, Berlin, Heidelberg, 157-177, https://doi.org/10.1007/978-3-662-01141-6_12, 1974.
- van Beek, P., Bourquin, M., Reyss, J.-L., Souhaut, M., Charette, M.A., and Jeande, C.: Radium isotopes to investigate the water mass pathways on the Kerguelen Plateau (Southern Ocean), Deep-Sea Research Part II, 55, 622-637, <https://doi.org/10.1016/j.dsr2.2007.12.025>, 2008.
- 1135 Vieira, L.H., Krisch, S., Hopwood, M.J., Beck, A.J., Scholten, J., Liebetrau, V., and Achterberg, E.P.: Unprecedented Fe delivery from the Congo River margin to the South Atlantic Gyre, Nat. Commun. 11: 556, <https://doi.org/10.1038/s41467-019-14255-2>, 2020.
- Windom, H.L., Moore, W.S., Niencheski, L.F.H., and Jahnke, R.A.: Submarine groundwater discharge: a large, previously unrecognized source of dissolved iron to the South Atlantic Ocean, Marine Chemistry, 102, 252-266, <https://doi.org/10.1016/j.marchem.2006.06.016>, 2006.
- 1140 Wyatt, N.J., Milne, A., Woodward, E.M.S., Rees, A.P., Browning, T.J., Bouman, H.A., Worsfold, P.J., and Lohan, M.C., Biogeochemical cycling of dissolved zinc along the GEOTRACES South Atlantic transect GA10 at 40°S, Global Biogeochem. Cy., 28, 44-56, <https://doi.org/10.1002/2013GB004637>, 2014.

Wyatt, N.J., Milne, A., Achterberg, E.P., Browning, T.J., Bouman, H.A., Woodward, E.M.S., and Lohan, M.C.: Seasonal cycling of zinc and cobalt in the Southeast Atlantic along the GEOTRACES GA10 section, Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-42>, in review, 2020.

Yamada, M. and Nozaki, Y.: Radium Isotopes in Coastal and Open Ocean Surface Waters of the Western North Pacific, Marine Chemistry, 19, 379-389, [https://doi.org/10.1016/0304-4203\(86\)90057-5](https://doi.org/10.1016/0304-4203(86)90057-5), 1986.

Commented [YTH4]: All tables and figures below have been updated.

Table 1 ²²⁶Ra and ²²⁸Ra activities, ²²⁸Ra/²²⁶Ra activity ratios and silica concentration

Cruise	Stn	Lon	Lat	Depth	Sal	Temp	²²⁶ Ra ^a	²²⁸ Ra ^b	²²⁸ Ra/ ²²⁶ Ra	Si
				(m)	(psu)	(°C)	(dpm 100 L ⁻¹)	(dpm 100 L ⁻¹)	(activity ratio)	(μM)
D357	0	17.967	-34.183	5	35.49	17.80	8.3 ± 0.3	3.22 ± 0.23	0.390 ± 0.028	1.42
D357	1	17.050	-34.617	5	35.49	17.95	8.3 ± 0.3	2.94 ± 0.19	0.354 ± 0.023	2.58
D357	1	17.050	-34.617	20	35.49	17.92	8.6 ± 0.3	2.68 ± 0.16	0.312 ± 0.019	2.41
D357	1	17.050	-34.617	50	35.47	17.50	8.6 ± 0.3	2.94 ± 0.17	0.342 ± 0.020	3.64
D357	1	17.050	-34.617	100	35.29	14.27	9.4 ± 0.4	1.19 ± 0.09	0.127 ± 0.010	6.02
D357	1	17.050	-34.617	200	35.01	11.66	9.6 ± 0.4	0.80 ± 0.08	0.084 ± 0.008	6.77
D357	1	17.050	-34.617	400	34.60	7.91	11.6 ± 0.4	0.29 ± 0.05	0.025 ± 0.005	17.97
D357	1	17.050	-34.617	1600	34.82	2.73	(15.3 ± 1.7)	0.23 ± 0.06	0.015 ± 0.004	54.95
D357	1	17.050	-34.617	2580	34.86	2.35	(15.5 ± 1.8)	0.58 ± 0.11	0.037 ± 0.006	56.31
D357	2	15.000	-35.467	5	35.58	17.94	8.3 ± 0.3	2.71 ± 0.18	0.327 ± 0.021	2.75
D357	2	15.000	-35.467	50	35.58	17.94	8.2 ± 0.3	2.85 ± 0.19	0.347 ± 0.023	3.45
D357	2	15.000	-35.467	100	35.57	17.78	8.4 ± 0.3	2.73 ± 0.19	0.326 ± 0.022	3.40
D357	2	15.000	-35.467	400	35.25	13.31	9.3 ± 0.3	0.76 ± 0.08	0.082 ± 0.008	5.31
D357	3	13.117	-36.333	5	35.08	13.00	9.1 ± 0.3	1.52 ± 0.13	0.167 ± 0.014	2.01
D357	3	13.117	-36.333	10	35.08	13.00	9.1 ± 0.3	1.65 ± 0.22	0.182 ± 0.024	2.34
D357	3	13.117	-36.333	20	35.08	13.00	8.8 ± 0.3			2.46
D357	3	13.117	-36.333	50	35.00	12.62	(9.1 ± 1.0)	2.40 ± 0.37	0.264 ± 0.028	2.37
D357	3	13.117	-36.333	100	34.73	11.08	9.2 ± 0.3	1.24 ± 0.09	0.135 ± 0.010	2.71
D357	3	13.117	-36.333	200	34.65	10.26	9.3 ± 0.3	0.61 ± 0.05	0.065 ± 0.005	3.20
D357	3	13.117	-36.333	400	34.56	8.20	9.7 ± 0.3	0.41 ± 0.04	0.042 ± 0.004	9.52
D357	3	13.117	-36.333	1410	34.59	3.01	(16.5 ± 1.8)	0.29 ± 0.05	0.018 ± 0.002	64.46
D357	3	13.117	-36.333	1500	34.65	2.87	16.6 ± 0.6			66.06
D357	3	13.117	-36.333	4335	34.74	1.16	(21.2 ± 2.3)	0.74 ± 0.11	0.035 ± 0.003	104.23
D357	3	13.117	-36.333	4425	34.74	1.13	22.2 ± 0.7			105.61
D357	3	13.117	-36.333	4706	34.73	1.10	(21.5 ± 2.4)	0.65 ± 0.10	0.030 ± 0.003	106.90
D357	3	13.117	-36.333	4776	34.73	1.09	(21.6 ± 2.4)	1.89 ± 0.31	0.088 ± 0.011	107.65

D357	3	13.117	-36.333	4823	34.73	1.10	22.2	±	0.7								123.30
D357	3	13.117	-36.333	4895	34.73	1.10	22.1	±	0.7								125.26
D357	4	10.400	-38.400	5	34.84	11.78	9.2	±	0.3	0.99	±	0.11	0.108	±	0.012	1.92	
D357	4	10.400	-38.400	700	34.25	4.59	13.4	±	0.5	0.26	±	0.03	0.019	±	0.003	20.07	
D357	4.5	7.800	-39.217	5	35.17	13.85	8.6	±	0.3	1.26	±	0.10	0.146	±	0.011	1.64	
D357	4.5	7.800	-39.217	10	35.17	13.78	(9.0	±	1.0)	1.08	±	0.15	0.121	±	0.010	1.67	
D357	4.5	7.800	-39.217	20	35.17	13.71	8.6	±	0.3							1.71	
D357	4.5	7.800	-39.217	90	35.17	13.56	7.4	±	0.3							1.59	
D357	4.5	7.800	-39.217	200	34.97	11.87	10.6	±	0.4	0.91	±	0.07	0.086	±	0.007	2.57	
D357	4.5	7.800	-39.217	400	34.69	9.22	10.0	±	0.3	0.69	±	0.10	0.069	±	0.010	6.92	
D357	4.5	7.800	-39.217	600	34.36	6.17	(10.3	±	1.1)	0.31	±	0.05	0.030	±	0.003	13.09	
D357	4.5	7.800	-39.217	2500	34.83	2.55	16.6	±	0.6							56.06	
D357	4.5	7.800	-39.217	3241	34.84	2.22	(16.3	±	1.8)	0.45	±	0.08	0.027	±	0.004	63.43	
D357	4.5	7.800	-39.217	3500	34.83	2.09	18.9	±	0.6							67.39	
D357	4.5	7.800	-39.217	4241	34.76	1.33	(20.1	±	2.2)	0.44	±	0.07	0.022	±	0.003	95.28	
D357	4.5	7.800	-39.217	4500	34.74	1.16	22.2	±	0.8							101.64	
D357	4.5	7.800	-39.217	4741	34.74	1.16	(21.0	±	2.3)	0.35	±	0.08	0.017	±	0.003	102.38	
D357	4.5	7.800	-39.217	5000	34.73	1.15	22.2	±	0.8							103.12	
D357	4.5	7.800	-39.217	5141	34.73	1.16	(21.6	±	2.4)	0.99	±	0.15	0.046	±	0.005	107.45	
D357	4.5	7.800	-39.217	5211	34.73	1.17	(21.5	±	2.4)	1.04	±	0.15	0.048	±	0.005	107.29	
D357	4.5	7.800	-39.217	5231	34.73	1.18	22.4	±	0.8							107.06	
JC068	18	-42.416	-40.001	5	34.66	18.19	(8.8	±	1.0)	2.15	±	0.24	0.268	±	0.006	0.15	
JC068	18	-42.416	-40.001	170	34.43	7.83	(9.5	±	1.0)	1.18	±	0.14	0.134	±	0.007	5.76	
JC068	18	-42.416	-40.001	420	34.14	4.52	(9.9	±	1.0)	1.52	±	0.18	0.163	±	0.005	9.15	
JC068	19	-47.417	-39.994	5	34.72	18.59	(8.9	±	1.0)	3.74	±	0.42	0.463	±	0.008	0.59	
JC068	20	-51.029	-37.983	5	35.09	22.18	(9.0	±	1.0)	3.95	±	0.44	0.480	±	0.004	1.53	
JC068	21	-52.503	-37.026	5	36.03	23.91	(8.9	±	1.0)	3.65	±	0.40	0.446	±	0.005	1.22	
JC068	21	-52.503	-37.026	100	36.36	19.69	(8.9	±	1.0)	2.12	±	0.25	0.260	±	0.010	1.16	
JC068	21	-52.503	-37.026	200	35.81	16.43	(9.0	±	1.0)	2.84	±	0.33	0.344	±	0.012	1.86	
JC068	21	-52.503	-37.026	600	34.56	7.85	(10.1	±	1.1)	1.02	±	0.12	0.107	±	0.004	10.66	
JC068	22	-53.102	-36.538	5	30.26	23.00	(10.0	±	1.1)	13.55	±	1.49	1.441	±	0.008	9.80	
JC068	23	-53.337	-36.338	5	29.62	23.35	(10.0	±	1.1)	14.08	±	1.55	1.469	±	0.008	11.04	
JC068	24	-54.000	-36.000	5	28.48	23.06	(11.3	±	1.2)	17.66	±	1.95	1.599	±	0.011	21.22	
JC068	25	-54.560	-35.493	5	30.61	23.04	(9.2	±	1.0)	12.76	±	1.41	1.497	±	0.013	3.68	

⁸Ra-226 activity in bracket is extrapolated from the ²²⁶Ra-silica relationship in Fig. A1. ⁸Ra-228 activity is calculated from the activity ratio of ²²⁸Ra/²²⁶Ra multiplied by ²²⁶Ra activity. All errors are 2 standard errors.

1155

Table 2 ²²⁸Ra-derived shelf-ocean dTE fluxes along the 40°S Atlantic transect^a

dTE fluxes	dCo	dFe	dZn
	(10 ³ nmol m ⁻² d ⁻¹)	(10 ⁴ nmol m ⁻² d ⁻¹)	(10 ⁴ nmol m ⁻² d ⁻¹)
(1) ²²⁸ Ra-derived diffusive fluxes			
Argentine margin (JC068)	4.0 ± 4.2	11 ± 9	4.4 ± 3.7
Cape margin (JC068)	4.7 ± 1.6	1.5 ± 0.9	1.2 ± 0.7
Cape margin (D357)	4.3 ± 1.8	1.2 ± 1.3	
(2) ²²⁸ Ra-derived advective fluxes			
Argentine margin (JC068)	21 ± 16	19 ± 13	6.3 ± 4.9
(3) TE/ ²²⁸ Ra-ratio-derived fluxes			
Argentine margin (JC068)	5.7 ± 4.0	7.9 ± 5.1	2.7 ± 1.5
Cape margin (JC068)	6.2 ± 4.0	3.1 ± 2.2	0.9 ± 1.1

^aFluxes are normalised to the area of cross-shelf section. All errors are ±1S.D.

1160

Table 3 ²²⁸Ra-derived vertical dTE fluxes along the 40°S Atlantic transect^a

dTE fluxes	dCo	dFe	dZn
Station	(nmol m ⁻² d ⁻¹)	(nmol m ⁻² d ⁻¹)	(nmol m ⁻² d ⁻¹)
Stn1 (JC068)	0.6 ± 0.3	11 ± 6	7 ± 5
Stn1 (D357)	0.3 ± 0.2		
Stn3 (JC068)	0.1 ± 0.1	6 ± 2	5 ± 3
Stn3 (D357)	0.7 ± 0.3	6 ± 2	
Stn4.5 (D357)	1.2 ± 0.6	7 ± 4	

^aFluxes are normalised to the surface area. All errors are ±1S.D.

Table 4 Net dTE fluxes in the 40°S Atlantic open ocean high productivity zone^a

dTE fluxes	dCo	dFe	dZn
	(10 ⁵ mol year ⁻¹)	(10 ⁶ mol year ⁻¹)	(10 ⁶ mol year ⁻¹)
(1) Shelf-ocean inputs			
Argentine Basin	1.6 – 8.7	3.2 – 7.8	1.1 – 2.6
Cape Basin	1.7 – 2.5	0.5 – 1.3	0.4 – 0.5
Argentine + Cape	3.3 – 11.2	3.7 – 9.1	1.5 – 2.1
(2) Vertical mixing	4 – 30	15 – 23	12 – 19
(3) Atmospheric inputs	0.5 – 1.3	4.1 – 13	1.6 – 16
(1)+(2)+(3) Total dTE fluxes	8 – 43	22 – 45	15 – 37
(4) Removal dTE fluxes	24 – 752	80 – 2507	40 – 251

^aThe high productivity zone is illustrated in Fig.8. The net dTE fluxes of (1) shelf-ocean inputs (Table 2) are multiplied by the area of cross-shelf section above the removal flux integration depth (100 m). The net TE fluxes of (2) vertical mixing (Table 3), (3) atmospheric inputs (Chance et al., 2015) and (4) the exported fluxes (assessed from the ²³⁴Th-POC fluxes and TE/C ratios) are multiplied by the surface area.




Figure 1: (a) Map of cruise tracks, station locations and surface currents. GA10, cruises, D357 and JC068 stations are labelled with green and red circles respectively. Stations of previous Ra studies are labelled with open symbols (TTO: Key et al., 1990; 1992a,b; GEOSECS: Ku and Lin, 1976; ANT XV/4: Hanfland, 2002). Surface currents are shown with arrows. (b) Salinity profiles along the cruise track of UK-GEOTRACES-GA10, labelled with water masses. Vertical lines indicate the stations where vertical Ra water profiles are available.

Deleted: E

Deleted: (

Deleted:) and GA10W (

Deleted:)




Figure 2: Depth profiles of (a) ^{226}Ra and (b) ^{228}Ra activities. The grey squares show Ra data from the previous GEOSECS study (Ku and Lin, 1976); the triangles show Ra data from the TTO program (Key et al., 1990; 1992a,b). Different water masses are characterised in (a) ^{226}Ra profile (see details in text). Error bars are $\pm 2\text{S.E.}$




Figure 3: Dissolved trace elements (dCo, dFe and dZn) and salinity in the surface water (< 10 m) along the 40°S Argentine and Cape transects. Red squares show data from cruise JC068 and green circles show data from cruise D357. The orange band indicates the boundary of Brazil Current (BrC) in the Argentine transect, highlighted by high salinity and changing TE gradients. $[\text{TE}]_{\text{ave-0}}$ is the average concentrations of dissolved TEs in the initial advective waters around where the Brazil Current merges into the SAC (around Stn21; $\pm 1\text{S.D.}$, $n = 4$). The dashed lines show linear regression trends through the TE data (Argentine transect: only data from the shelf to BrC; Cape transect: the whole transect) and the gradient ($\Delta\text{TE}/\Delta x$) errors are $\pm 1\text{S.D.}$.

Deleted: .

Deleted: T

Formatted: Font: Symbol

Formatted: Font: Symbol




Figure 4: Depth profiles of dissolved trace elements (dCo, dFe and dZn) in the upper ocean (< 600 m). Red squares show data from cruise JC068 and green circles show data from cruise D357. The dashed lines show linear regression trends, and the vertical gradient ($\Delta TE/\Delta z$) errors are ± 1 S.D.

Deleted: with

Deleted: uncertainty

Deleted: (

Deleted:)




Figure 5: Plots of $^{228}\text{Ra}_{\text{ex}}$ in the surface ocean ($< 10 \text{ m}$) along the (a) Argentine and (b) Cape transects of 40°S Atlantic, with the distributions of salinity and temperature shown in (c) and (d). The orange band indicates the boundary of Brazil Current (BrC) in the Argentine transect, highlighted by high salinity and temperature. The red and green dashed lines show exponential regression trends through the $^{228}\text{Ra}_{\text{ex}}$ data (Argentine transect: only to BrC; Cape transect: the whole transect). The gradients of the exponential fit (the a values) are used in Eqn. (3) for the K_x calculation. The errors of a and K_x are $\pm 1\text{S.D.}$ The grey dashed line shows a linear regression trend through the $^{228}\text{Ra}_{\text{ex}}$ data from BrC to the open ocean in the Argentine transect, which is used in Eqn. (4) to estimate the advection water transport velocity (w).

Deleted: equations

Deleted: rearranged to match

Formatted: Font: Italic

210




Figure 6: Depth profiles of (a) to (f) seawater $^{228}\text{Ra}_{\text{ex}}$ activity in red (Argentine Basin) and green (Cape Basin) circles and density (sigma-t) shown in grey dashed lines, and (g) to (l) density and temperature in the upper ocean shown at Stn1, 2, 3, 4.5, 18 and 21. Depths of mixed layer are labelled with the orange horizontal dashed lines, defined by the sigma-t and temperature profiles. The red and green dashed lines show exponential regression trends through the $^{228}\text{Ra}_{\text{ex}}$ data below the mixed layer (including the average value of the mixed layer). The gradients of the exponential fit (the a values) are used in Eqn. (5) for the K_e calculation (errors $\pm 1\text{S.D.}$)

Deleted: equations

Deleted: .

Deleted: 3

Formatted: Font: Italic




Figure 7: Plots of $^{228}\text{Ra}_{\text{ex}}$ activity at water depth 5, 50, 100, 200 and 400 m versus distance to the coast of Cape Town in South Africa. The dashed line shows an exponential regression line through the data at 5 m.




Figure 8: Schematic diagram of dissolved trace element inputs and outputs in the high productivity zone in the open ocean along the surface 40°S Atlantic. The approximate dimensions of the high productivity zone and continental shelves are labelled, assuming that the zone spans across the latitude from 35°S to 45°S (~1110 km) and the longitude from 55°W to 20°E (~6375 km) with a **removal flux integration** depth of 100 m. The arrows indicate different TE inputs and outputs in this region. The TE fluxes from Table 4 are shown and colour-coded according to the sources. The vertical red cross sections are used to normalise the shelf-ocean TE and ^{228}Ra fluxes. The net shelf-ocean TE fluxes represent the total inputs from rivers, SGD and shelf sediments, and the outputs by particle or biological removal between the shelf-ocean mixing from both sides of the continental margins. The blue surface area is used to estimate the net TE inputs from dust and vertical mixing, and the red cross section above the mixed layer is used to estimate the net shelf-ocean TE fluxes (Table 4).

Deleted: n
Deleted: annual average mixed layer
Deleted: ~
Deleted: 5
Deleted: , and the units are $10^5 \text{ mol year}^{-1}$ for dCo and $10^6 \text{ mol year}^{-1}$ for dFe and dZn. ...

▼

▲

Page 2: [1] Deleted

Yu-Te Hsieh

07/01/2021 18:57:00

▼

▲

Page 2: [2] Deleted

Yu-Te Hsieh

07/01/2021 19:01:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [3] Deleted

Yu-Te Hsieh

12/01/2021 15:34:00

▼

▲

Page 15: [4] Formatted

Yu-Te Hsieh

12/01/2021 15:45:00

▼

▲

Page 15: [4] Formatted

Yu-Te Hsieh

12/01/2021 15:45:00

Superscript

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [5] Deleted Yu-Te Hsieh 12/01/2021 15:56:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲ Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00 ▼

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [6] Deleted Yu-Te Hsieh 12/01/2021 16:07:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00

▲
▼.....◀
Page 15: [7] Deleted Yu-Te Hsieh 12/01/2021 17:32:00