

Variability of North Atlantic CO₂ fluxes for the 2000–2017 period estimated from atmospheric inverse analyses

Zhaohui Chen¹, Parvatha Suntharalingam¹, Andrew J. Watson², Ute Schuster², Jiang Zhu³, and Ning Zeng⁴

5 ¹School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK.

²College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4RJ, UK.

³International Center for Climate and Environment Sciences, Institute of Atmospheric Physics, Chinese Academy of Science, Beijing, 10029, China.

10 ⁴Department of Atmospheric and Oceanic Science, and Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland, 20742, USA

Correspondence to: Zhaohui Chen (Zhaohui.chen@uea.ac.uk)

Abstract. We present new estimates of the regional North Atlantic (15°N–80°N) CO₂ flux for the 2000–2017 period using atmospheric CO₂ measurements from the NOAA long term surface site network in combination with an atmospheric carbon cycle data assimilation system (GEOSChem–LETKF). We assess the sensitivity of flux estimates to alternative ocean CO₂ prior flux distributions and to the specification of uncertainties associated with ocean fluxes. We present a new scheme to characterize uncertainty in ocean prior fluxes, derived from a set of eight surface pCO₂–based ocean flux products, and which reflects uncertainties associated with measurement density and pCO₂–interpolation methods. This scheme provides improved model performance, in comparison to fixed prior uncertainty schemes, based on metrics of model–observation differences at the network of surface sites. Long term average posterior flux estimates for the 2000–2017 period from our GEOSChem–LETKF analyses are -0.255 ± 0.037 PgC y⁻¹ for the subtropical basin (15°N–50°N), and -0.203 ± 0.037 PgC y⁻¹ for the subpolar region (50°N–80°N, eastern boundary at 20°E). Our basin–scale estimates of interannual variability (IAV) are 0.036 ± 0.006 PgC y⁻¹ and 0.034 ± 0.009 PgC y⁻¹ for subtropical and subpolar regions respectively. We find statistically significant trends in carbon uptake for the subtropical and subpolar North Atlantic of -0.064 ± 0.007 and -0.063 ± 0.008 PgC y⁻¹ decade⁻¹; these trends are of comparable magnitude to estimates from surface ocean pCO₂–based flux products, but larger, by a factor of 3–4, than trends estimated from global ocean biogeochemistry models.

1 Introduction

The ocean plays a key role in the global carbon budget, accounting for 2.5 ± 0.6 PgC y⁻¹ of net CO₂ uptake from the atmosphere during the last decade (period 2009–2019), a level equivalent to ~26% of global fossil CO₂ emissions (Friedlingstein et al., 2020). The North Atlantic ocean has been identified a region of significant net oceanic CO₂ uptake in a range of recent analyses (Schuster et al., 2013, Landschützer et al., 2013, Lebehof et al., 2019), and also the location of the largest Northern Hemisphere uptake of anthropogenic CO₂ in recent decades (Gruber et al., 2019, Khatiwala et al., 2013, Sabine et al., 2004). Recent

estimates of net air–sea CO₂ fluxes derived from sea surface partial pressure CO₂ measurements (pCO₂) indicate net annual uptake for the North Atlantic over the past decade (2009–2018) with a range of 0.35–0.55 PgC y⁻¹ (Landschutzer et al., 2016; Rodenbeck et al., 2013; Zeng et al., 2015; Watson et al., 2020), and equivalent to about 14%–22% of the global net ocean carbon ocean sink reported for this period. Regionally aggregated air–sea CO₂ fluxes over the North Atlantic basin also display significant variability on interannual (Watson et al., 2009) and decadal timescales (Landschützer et al., 2016, 2019). Based on analyses of surface pCO₂ measurements, variations in regional pCO₂ trends were observed in the subtropical and subpolar regions, potentially associated with large–scale climate oscillations such as the North Atlantic Oscillation and the Atlantic Multi–decadal Variation (McKinley et al., 2011, Landschützer et al., 2019, Macovei et al., 2020). Devries et al. (2019) estimated a negative trend (i.e., a strengthening ocean sink) in North Atlantic CO₂ uptake for the 2000–2009 period based on analysis of pCO₂–based estimates and ocean models. Lebehot et al. (2019) find statistically significant trends in surface ocean CO₂ fugacity (fCO₂) for the 1992–2014 period from both observation–based surface mapping methods and from the CMIP5 Earth System models.

Recent analyses of North Atlantic air–sea CO₂ fluxes have primarily been based on ‘bottom–up’ methods of varying complexity which use interpolated surface ocean pCO₂ distributions (derived from in situ pCO₂ measurements) in combination with parameterizations of air–sea gas exchange (e.g., Landschützer et al., 2013; Rödenbeck et al., 2015; Takahashi et al., 2002, 2009). Estimates of air–sea CO₂ fluxes have also been derived by alternative methods such as global ocean biogeochemical models (e.g., Buitenhuis et al., 2013; Law et al., 2017), and ‘top–down’ methods which involve the application of inverse analyses or data assimilation methods to atmospheric and oceanic CO₂ measurements (e.g., Gruber et al., 2009, Mikaloff–Fletcher et al., 2006, Gurney et al., 2003, Peylin et al., 2013). ‘Top–down’ analyses estimate surface CO₂ fluxes by using information on observed gradients in atmospheric CO₂ together with atmospheric transport constraints (typically from 3 D atmospheric models) and prior information on the magnitude and associated uncertainties of surface CO₂ flux distributions (Rödenbeck et al., 2003; van der Laan–Luijkx et al., 2017; Peters et al., 2005; Peylin et al., 2013; Chevallier et al., 2014; Gaubert et al., 2019).

Previous studies also note that estimates of carbon fluxes from the atmospheric inverse method are sensitive to the specification of the prior flux distribution and its associated uncertainty distribution (Carouge et al., 2010; Chatterjee et al., 2013; Peylin et al., 2013). While there have been recent studies evaluating the sensitivity of land–based carbon flux estimates to specification of the prior flux and its uncertainty, there has been far less examination of ocean flux estimates from inverse methods. Several global inverse model assessments of the past decade have relied on the climatological ocean–atmosphere CO₂ flux database of Takahashi et al. (2009) to specify prior ocean fluxes. In view of the limited information available on the temporal and spatial variability of ocean carbon fluxes from this climatological ocean database, these inverse analyses have adopted different approaches to the specification of prior uncertainty for ocean fluxes, ranging from uncertainties derived from a separate ocean model inversion (in the case of Nassar et al., 2011), to a specified percentage of the prior flux magnitude (Feng et al., 2016, Liu et al., 2016).

65 In this study we present a new long term estimate of North Atlantic air–sea CO₂ fluxes for recent decades (period 2000–2017) using atmospheric inverse methods. We focus in particular on the specification of prior ocean fluxes (including sensitivity of flux estimates to alternative prior flux distributions) and on their associated flux uncertainties. To our knowledge these influences on inverse estimates of North Atlantic CO₂ flux have not been assessed previously. We use the carbon cycle data assimilation system GEOSChem–LETKF (GCL, described further in Sect. 2) which combines the global atmospheric CO₂ transport model GEOS–Chem (Nassar et al., 2010) with the Local Ensemble Transform Kalman Filter (LETKF) data assimilation system (Hunt et al., 2007; Miyoshi et al., 2007; Liu et al., 2019). In recent years several new global air–sea CO₂ flux products have been developed based on mappings of ocean surface pCO₂ measurements (e.g., Landschutzer et al., 2016, Rodenbeck et al., 2014, Watson et al., 2020, and products reported in the intercomparison of Roedenbeck et al., 2015). These ocean flux distributions are frequently derived from interpolations of surface ocean pCO₂ measurements from the SOCAT database (Bakker et al., 2016) together with parameterizations of air–sea gas exchange. Following recent updates, the surface ocean pCO₂ database SOCATv2020 (<https://www.socat.info/index.php/data-access/>), now includes over 28 million surface ocean carbon measurements. The SOCAT database provides a valuable resource towards the development of bottom–up estimates of ocean–atmosphere CO₂ fluxes, and a compilation of these flux products is reported in the recent Global Carbon Budget (Friedlingstein et al., 2020). The increased range of global air–sea CO₂ flux products available (beyond the Takahashi et al., 2009 climatology) provides a valuable opportunity to develop an improved representation of air–sea CO₂ flux variability and a more robust characterization of the uncertainties associated with ocean carbon fluxes. In this study we employ some of the recently developed ocean CO₂ flux products to provide a new method of characterizing the prior ocean flux uncertainty used for atmospheric inverse analyses. The methodology is based on the ensemble spread of the multiple ocean flux products, and reflects underlying uncertainties in these products, such as those associated with sampling density of the surface measurements and interpolation method employed. It provides a spatially and temporally variable specification of prior flux uncertainty that will be of value to the inverse modeling community.

The remainder of the paper is organized as follows: Section 2 covers the methodology of the atmospheric inverse analysis, outlining the carbon cycle data assimilation system (GEOSChem–LETKF), the atmospheric CO₂ observations, and specifications of prior fluxes and uncertainties. Further details of the methodology are presented in the Appendix. In Sect. 3 we present GEOSChem–LETKF assessments of alternative specifications of ocean prior fluxes and flux uncertainties, and then use these results to derive long term estimates of North Atlantic CO₂ fluxes for the 2000–2017 period. We also summarize specific characteristics of North Atlantic CO₂ fluxes derived from these analyses, namely, long term means, trends, and interannual variability of fluxes, and compare our results with other recent relevant studies.

2 Materials and Methods

95 2.1 Overview

Our analysis employs the global GEOS–Chem atmospheric chemistry transport model together with the Local Ensemble Transform Kalman Filter (LETKF) (described in Sect. 2.2) and atmospheric CO₂ observations from the NOAA–ESRL network of surface sites (Sect. 2.3). Section 2.4 describes the compilation of the set of air–sea CO₂ flux products and the derivation of the prior flux uncertainty specification for the North Atlantic based on the ensemble spread of these products. Section 3 presents model results, including sensitivity analyses assessing different prior flux representations and flux uncertainty schemes (Sect. 100 3.1), and regional CO₂ flux estimates for the 2000–2017 period from the GEOSChem–LETKF system (Sect. 3.2). Further details on model analyses, observations and uncertainty calculations are presented in the sections below and in the Appendix.

2.2 The GEOSChem–Local Ensemble Transform Kalman Filter (GCL) system

The GEOS–Chem atmospheric chemistry transport model has been used in a range of previous investigations into atmospheric CO₂ and applied in conjunction with inverse analyses to estimate surface carbon fluxes (Nassar et al., 2010, 2011; Suntharalingam et al., 2005; Liu et al., 2016). In this analysis we employ GEOSChem v11–01 at a horizontal resolution of 2° latitude by 2.5° longitude, with 47 levels in the vertical. Model transport fields are provided by GEOS–5 assimilated meteorological data from the NASA Global Modeling and Assimilation Office (GMAO, Rienecker et al., 2008). The GEOSChem configuration employed here primarily follows that of Nassar et al. (2011), but with updated representation of prior fluxes; more detail on the prior CO₂ fluxes and uncertainties implemented in this study is given in Sect. 2.4. 110

The Local Ensemble Transform Kalman Filter (LETKF) is a data assimilation system which provides an estimate given a prior (or “background”) estimate of the current state based on past and current data (in this case, the atmospheric CO₂ mole fraction observations). The general framework of the LETKF is described in Hunt et al. (2007); it has been adapted by Miyoshi et al. (2007) to provide gridscale localized analysis of flux estimates. The LETKF system has been used to estimate CO₂ fluxes in a range of previous studies (e.g, Kang et al., 2012; Liu et al., 2016, 2019). The LETKF provides iterative estimates of the time evolution of the system state, x , (here representing the grid–scale surface carbon fluxes). Each step involves a forecast stage (based on a physical model of the system evolution) and a state estimation stage (the ‘analysis’ step), which combines system observations, y , together with the background forecast, x^b , to derive the improved state estimate. The observation operator H provides the mapping from the state space to the observation space; in this study H is provided by the GEOS–Chem atmospheric model. 120

In this analysis we employ the complete GEOSChem–LETKF (GCL) data assimilation system to conduct sensitivity analyses on the ocean prior fluxes, and to provide a long term flux estimate of surface CO₂ fluxes for the North Atlantic for the period 2000–2017. We report a posteriori fluxes on monthly timescales for the 2000–2017 period; the optimized monthly fluxes are derived from four sequential weeks of assimilation cycles, as further described below. Our methods follow the implementation of the LETKF system by Liu et al. (2019), who have extended the previous carbon data assimilation system of Kang et al. 125

(2011, 2012). The study of Kang et al. (2011) assimilated meteorological data and atmospheric CO₂ concentrations to provide estimated atmospheric CO₂ concentrations as part of the state estimate. Kang et al. (2012) extended this method to also provide estimates of surface carbon fluxes. Both these LETKF studies assimilated meteorological data and atmospheric CO₂ concentrations and employed a short assimilation window of 6 hours in order to maintain linear behaviour of the ensemble perturbations (Kang et al., 2011, 2012). In addition, Kang et al. (2012) also tested longer assimilation windows (up to 3 weeks) for LETKF formulations that assimilated atmospheric CO₂ concentrations alone (eliminating the assimilation of the meteorological data). The LETKF system of Liu et al. (2019) extended the Kang et al. (2011, 2012) analyses by incorporating the GEOSChem atmospheric model as the forecast model, along with its representation of surface CO₂ fluxes which provide the prior flux specification for the forecast step. However, Liu et al. (2019) assimilate only atmospheric CO₂ measurements (i.e., no assimilation of meteorological measurements), and use an assimilation window of 7 days; the duration of the assimilation window was selected to maximize the correlation between observations and surface fluxes. The GEOSChem–LETKF system employed in our study follows the Liu et al. (2019) formulation; atmospheric CO₂ measurements are assimilated at 7 day timescales, with the LETKF analysis step providing updates of the surface fluxes and associated uncertainties required as initial conditions for the next weekly forecast step. We report monthly flux estimates following four assimilation cycles. Further details on the LETKF and the governing equations for flux estimation are provided in Appendix A.

2.3 Atmospheric CO₂ Observations

Atmospheric CO₂ observations used for this study are taken from the NOAA–ESRL GLOBALVIEWplus Observation Package v4.2 (Obspack, Cooperative Global Atmospheric Data Integration Project, 2018). CO₂ measurement records for the period 2000–2017 from 86 surface sites were used in this analysis. Further details on the measurement sites and the site–specific observation uncertainty characteristics are presented in Table A1 of the Appendix. The specification of observational uncertainty associated with incorporation of the atmospheric CO₂ measurements into the LETKF is derived using the methods of Chevallier et al. (2010); we use the standard deviation of measurement variability from detrended and deseasonalized CO₂ time series at each measurement site. The resulting specification of observational uncertainty varies between 0.16 ppm (for stations in and around the Southern Ocean) to over 5 ppm (for stations in continental interiors) (see Appendix Table A1 for more details).

2.4 Specification of Prior CO₂ Fluxes and Associated Flux Uncertainties

The GEOSChem model CO₂ simulation employed in this study includes representation of fossil fuel emissions, air–sea fluxes and exchange with the terrestrial biosphere. Details of the data sources used to specify the prior flux distributions are outlined here. Fossil fuel emissions are taken from Chevallier et al. (2019) (Global Atmospheric Research version 4.3.2, Crippa et al.,

2016, scaled globally and annually from Le Quéré et al., 2018), and land biosphere fluxes from the Joint UK Land Environment Simulator (JULES, Clark et al., 2011).

160 The focus of our study is on North Atlantic Ocean CO₂ fluxes, and we investigate the representation of ocean prior fluxes and prior flux uncertainty in more detail. Firstly, in Sect. 3.1, in a set of sensitivity analyses, we compare the implementation of three different representations of ocean CO₂ fluxes that have been used to specify prior fluxes in recent inverse analyses: (i) the widely used Takahashi et al. (2009) climatology (hereinafter Ta), (ii) the interannually varying flux product of Landschützer et al. (2016) derived from surface pCO₂ distributions (hereinafter La), and (iii) the interannual fluxes from the ocean mixed-layer scheme of Rödenbeck et al. (2014) (hereinafter Ro). We also evaluate in more detail, the impact of different specifications
165 of prior flux uncertainty for ocean fluxes. Many previous atmospheric inverse estimates of air–sea carbon fluxes have employed relatively simple characterizations of the prior ocean flux uncertainty, for example, based on a fixed proportion of the grid–scale or regional prior flux (Nassar et al., 2011, Liu et al., 2016, Feng et al., 2016). In Sect. 3.1, we employ both fixed flux uncertainties, and also present an alternative scheme derived from the ensemble spread of ocean CO₂ flux products, as described below.

170 The prior ocean flux distributions employed in atmospheric inversions are frequently derived from interpolations of the surface ocean pCO₂ database (e.g., SOCAT, Bakker et al., 2016) in combination with ocean–atmosphere gas exchange parameterizations. Uncertainties in the derived products stem from uncertainties in the input data (e.g., density of measurements), interpolation methods, and gas–transfer parameterizations (Landschützer et al., 2013). However, some ocean regions, the North Atlantic in particular, have a higher density of pCO₂ measurements and more consistent flux estimates from
175 pCO₂–based products (Schuster et al., 2013, Landschützer et al., 2013). Here we exploit the recent expansion of pCO₂–based ocean flux products to outline a new specification of ocean prior flux uncertainty based on the ensemble–spread of the different flux products (the “spread–based” uncertainty scheme). Towards the development of the spread–based scheme, we have compiled a set of eight global gridded interannually varying ocean–atmosphere CO₂ flux products. These are Landschützer et al., 2016, Rodenbeck et al., 2014, Denvil–Sommer et al., 2019, Iida et al., 2015, Zeng et al., 2015, Gregor et al., 2019, Chau
180 et al., 2020, and Watson et al., 2020.

The spread–based prior flux uncertainty scheme uses a diagnostic derived from the variation among the set of ocean atmosphere carbon flux products (see Eq. (1)). This scheme specifies lower uncertainty levels where alternative prior flux representations are in accord (e.g., when well–constrained by availability of surface pCO₂ measurements), and higher uncertainty levels where the prior flux distributions differ significantly (typically in under–sampled regions or those of
185 significant flux variability). This specification follows previously used methods to characterize uncertainties in ocean flux distributions (e.g., Bopp et al., 2013). For this spread–based uncertainty specification, the gridded prior flux uncertainty, $U(i,j)$ (for a gridcell with coordinates (i,j)) is specified as the standard deviation of the spread of the different prior flux products. Thus, the uncertainty $U(i,j)$ is calculated as:

$$U(i, j) = \text{sqrt}\left(\sum_k^K (f_k(i, j) - \overline{f(i, j)})^2\right)/(K - 1) \quad (1)$$

Here K is the total number of the prior ocean flux products considered, and subscript k refers to an individual flux product. $f_k(i, j)$ represents the gridded monthly flux for each prior ocean flux and $\overline{f(i, j)}$ is the gridded monthly mean across all prior ocean flux products. These prior flux uncertainties are estimated on monthly timescales and also account for interannual variations. The uncertainty statistics of the prior ocean flux distributions will be dependent on the uncertainties associated with the respective inputs and methods of constructing the flux products. Ocean–atmosphere carbon flux products derived from surface ocean pCO_2 measurements are generally subject to two main sources of uncertainty: (i) in the specification of the surface CO_2 partial pressure difference across the air–sea interface, and (ii) in the specification of the gas–exchange coefficient used to derive fluxes (e.g., see discussion of Landschutzer et al., 2013; Watson et al., 2020). In the extended database of 8 pCO_2 –based flux products that we present above, the majority of the flux products (seven of the eight) rely on the surface ocean pCO_2 data of the SOCAT database (Bakker et al., 2016). These flux products will be subject to similar uncertainties associated with data coverage in different ocean regions, although the uncertainties due to differences among surface interpolation methods may vary.

In this study we account for spatial correlations in the prior ocean fluxes, by inclusion of off diagonal elements in the background error covariance matrix P^b (Appendix A Eq. A3). We follow the recommendations of Jones et al. (2012) on autocorrelation length scales in the surface ocean. That study derived spatial autocorrelation functions for air–sea fluxes from an analysis of the surface ocean pCO_2 database reported in Takahashi et al. (2009), combined with a gas exchange parameterization. We currently do not account for spatial correlation in land fluxes, but will investigate this in future analyses.

3 Results and Discussion


We first present in Sect. 3.1 results of short–term sensitivity tests that compare the influence of different prior ocean flux distributions and prior ocean flux uncertainty schemes on GCL estimates of North Atlantic (NA) CO_2 fluxes. Using these analyses as a basis, in Sect. 3.2 we conduct a multi–year GCL analysis of North Atlantic CO_2 ocean fluxes for the 2000–2017 period. We also report on derived characteristics of regionally aggregated North Atlantic subtropical and subpolar fluxes (long term means, trends and interannual variability) and compare these GCL results with recent estimates from other methodologies, including global ocean biogeochemical models (GOBMs), other atmospheric inverse studies, and surface pCO_2 –based data products.

3.1 Sensitivity tests on specification of prior flux uncertainty

In this section we investigate, via sensitivity analyses, the application of the spread–based prior flux uncertainty scheme outlined in Sect. 2.4 in comparison to the fixed prior uncertainty levels commonly used in previous inverse estimates of ocean

CO₂ fluxes. The alternative specifications of prior flux uncertainty for ocean fluxes employed include (a) fixed percentage-based levels (U1:60% of prior flux, and U2:120% of prior flux), and (b) gridded flux uncertainties representing the variation or ‘spread’ of the different ocean flux data products at each location, and based on the standard deviation of the variation among the prior fluxes (U3: spread-based uncertainty; see Eq. (1)). The selection of the fixed percentage prior uncertainty levels used in the sensitivity analyses was based on the range of variability seen for the individual prior flux distributions (Fig. 1) for the sub-regions of the North Atlantic. These ranged from average levels of ~60% for the subtropical North Atlantic to levels greater than 120% for the subpolar North Atlantic, hence we have selected a level of U1:60% to characterize the lower sensitivity case, and U2:120% for the higher case. We apply the alternative flux uncertainty specifications to the three different ocean prior flux distributions discussed in Sect. 2.4, namely: (i) the Takahashi et al. (2009) climatology (Ta), (ii) the flux product of Landschützer et al. (2016) (La), and (iii) the flux product of Rödenbeck et al. (2014) (Ro).

Sensitivity analyses are conducted for the year 2003, following a 3 year GEOSChem model spin-up, starting from January 1st, 2000; the length of spin-up was determined by recommendations on the duration required for stabilization of tropospheric CO₂ gradients (e.g., Gurney et al., 2002), and following methods used for previous GEOSChem CO₂ analyses (e.g., Nassar et al., 2010). The year 2003 was selected for sensitivity tests as the first viable year following spin-up. Analyses of inter-annual variability in Atlantic CO₂ (e.g., Landschützer et al., 2013; Schuster et al., 2013) do not find 2003 to be an anomalous year for regional ocean fluxes. We evaluate the sensitivity of posterior ocean flux estimates with three different prior ocean uncertainty schemes U1, U2, and U3, described above; these are applied in turn for each of the three prior ocean flux distributions (Ta, La and Ro). Figure 1 presents the seasonal variation of the spatial distribution of the spread-based prior ocean flux uncertainty U3 (3 month averages for the year 2003). Figure 1 demonstrates that over the course of the year, and particularly in the Northern Hemisphere winter months, the spread-based uncertainty scheme (U3) provides a looser constraint on prior fluxes (i.e., levels of prior flux uncertainty > 120%) than the U1 and U2 schemes in the subpolar region, and a tighter constraint in the subtropical region (levels < 60%).



240 **Figure 1.** Distribution of the spread-based prior ocean flux uncertainty (U_3) (3 month averages for the year 2003). The distribution in this
 study is calculated from the following 8 air-sea CO_2 flux products: (1) Denvil-Sommer et al., 2019 (product LSCE-FFNN-v1); (2) Iida et
 al., 2015 (JMA); (3) Zeng et al., 2015 (NIES); (4) Gregor et al., 2019 (CSIR-ML6); (5) Chau et al., 2020 (CMEMS); (6) Watson et al.,
 2020; (7) Landschützer et al., 2016; (8) Rödenbeck et al., 2014. It is represented here as a percentage of the prior ocean flux for ease of
 245 comparison with U_1 and U_2 . The percentage shown for each grid-cell is derived from the ratio of spread-based prior ocean uncertainty
 divided by the prior ocean flux value at that grid cell. DJF represents the monthly average for December, January, February; MAM for
 March, April, May; JJA for June, July, August; SON for September, October, November.

Table 1 summarizes the prior and posterior ocean flux estimates for the global and North Atlantic region (sub-divided into
 250 subpolar and subtropical regions) from the respective sensitivity tests. The distribution of prior flux for the subtropical North
 Atlantic shows closer agreement among the three source representations (T_a , L_a and R_o), with regional variation of 0.05 PgC
 y^{-1} , in comparison to a regional variation of $\sim 0.1 \text{ PgC y}^{-1}$ for the subpolar region. Under the constraints provided by the
 atmospheric CO_2 observations all posterior flux estimates for the North Atlantic show increased uptake (Table 1), indicating

that all three representations of ocean prior flux underestimate the regional net atmosphere–ocean flux for the 2003 period. Largest changes in the regional posterior fluxes are estimated under the U3 specification of prior flux uncertainty. In addition, our estimates indicate a larger increase in CO₂ uptake in the subpolar basin (~0.05 PgC y⁻¹, changing from a prior flux range of -0.13 to -0.23 PgC y⁻¹ to posterior flux range of -0.18 to -0.27 PgC y⁻¹, for the U3 scenarios), in comparison to the smaller magnitude change for the subtropical North Atlantic basin (of ~ 0.04 PgC y⁻¹ from around -0.18 PgC y⁻¹ to -0.22 PgC y⁻¹ for the U3 scenarios)

Table 1. Global and North Atlantic CO₂ flux estimates from the GEOSChem–LETKF(GCL) system for year 2003 (PgC y⁻¹) summarizing sensitivity analyses on the prior ocean flux distribution and prior flux uncertainty. Prior flux references are Ta: Takahashi et al., 2009; La: Landschutzer et al., 2016; Ro: Rodenbeck et al., 2014. Prior flux uncertainty specifications are: U1: 60%; U2: 120%; U3: spread-based (following methods of Sect. 2.4).

Global Ocean CO ₂ Flux (PgC y ⁻¹)					
Ta	-1.37	La	-1.25	Ro	-2.09
TaU1	-1.63±0.13	LaU1	-1.52±0.13	RoU1	-2.31±0.16
TaU2	-2.05±0.26	LaU2	-1.96±0.26	RoU2	-2.68±0.31
TaU3	-1.97±0.17	LaU3	-1.83±0.19	RoU3	-2.60±0.18
North Atlantic subtropics [15°N–50°N]					
Ta	-0.22	La	-0.18	Ro	-0.17
TaU1	-0.23±0.02	LaU1	-0.19±0.02	RoU1	-0.18±0.02
TaU2	-0.25±0.05	LaU2	-0.21±0.04	RoU2	-0.20±0.04
TaU3	-0.26±0.03	LaU3	-0.22±0.03	RoU3	-0.23±0.03
North Atlantic subpolar [50°N–80°N], eastern boundary at 20°E					
Ta	-0.23	La	-0.13	Ro	-0.21
TaU1	-0.23±0.05	LaU1	-0.13±0.02	RoU1	-0.22±0.04
TaU2	-0.25±0.1	LaU2	-0.14±0.05	RoU2	-0.23±0.09
TaU3	-0.27±0.05	LaU3	-0.18±0.05	RoU3	-0.24±0.05


We note that the increases in estimated uptake for the North Atlantic basins are relatively smaller (on average in the range 10–20%) than the increased uptake estimated on the global scale (~30–50% changes, see Table 1), indicating that the prior flux representations of North Atlantic carbon uptake are more consistent with the constraints from atmospheric CO₂ measurements than the comparison on a global scale.

The U3 flux uncertainty specification is derived from the variation among a set of ocean–atmosphere carbon flux products (Eq. (1)). This scheme specifies lower uncertainty levels where alternative prior flux representations are in accord (e.g., when well constrained by availability of surface pCO₂ measurements, as in the subtropical North Atlantic), and higher uncertainty levels where the prior flux distributions differ significantly (typically in under-sampled regions or those of significant flux variability, such as the subpolar North Atlantic). We further assess the value of the U3 scheme using a metric of GCL modeled

atmospheric CO₂ concentration; specifically, estimates of the model–observation mismatch for the year 2003 at the NOAA network station sites in the North Atlantic using the a posteriori fluxes associated with the sensitivity analyses of this section (Appendix Table A2). The results summarized in Table A2 indicates that scheme U3 provides the smallest magnitude model–observation mismatch for the individual North Atlantic sites and for the global network average. For the long term analyses of the remainder of this study, therefore, we use the U3 spread–based flux uncertainty scheme in preference to the fixed level flux uncertainty schemes used in many previous inverse analyses.

3.2 Multi–year analyses of North Atlantic CO₂ fluxes

In this section we present results of a multi–year GCL analysis (for the period 2000–2017), calculating regional estimates of North Atlantic CO₂ fluxes on annual to decadal timescales. Prior flux distributions for fossil emissions, and exchange with the land biosphere fluxes are as described in Sect. 2.4. For ocean prior fluxes, we employ the distribution of Landschützer et al. (2016); this is an established surface pCO₂–based product and also provides inter–annually varying fluxes over the entire estimation period (2000–2017), in comparison to the climatology–only fluxes of Takahashi et al. (2009). Ocean prior flux uncertainties are specified by the spread–based scheme U3 described above and derived from the eight ocean–atmosphere pCO₂–based flux products summarized in Sect. 2.4.



290 **Figure 2.** Comparison of annual air–sea CO₂ fluxes for North Atlantic for the 2000–2017 period for: (a) North Atlantic Subtropics; and (b) North Atlantic Subpolar regions. The GCL posterior flux estimate from this study (red) is derived from the prior flux of Landschützer et al., 2016 (pCO₂La: black). The grey shaded area represents the uncertainty estimate on the GCL posterior flux (plotted at a 1 sigma level). Also shown are the flux estimates of (i) Chevallier et al., 2019 (CAMS: yellow); (ii) Jacobson et al., 2020 (CT: CarbonTracker2019: pink); and (iii) van der Laan–Luijkx et al., 2017 (CTE: Carbon Tracker Europe: blue). All time series shown have a 12 month running mean filter applied.

295 Figure 2 presents the variation of air–sea CO₂ flux for the North Atlantic subtropical and subpolar regions for the 2000–2017 period (represented as a 12 month running average). We also plot on Fig. 2 flux estimates from three other atmospheric inverse analysis studies including CAMS (v18r2, Chevallier et al., 2019), CT (CarbonTracker 2019, Jacobson et al., 2020) and CTE (Carbon Tracker Europe, van der Laan–Luijkx et al., 2017). All data are regridded to 2° latitude × 2.5° longitude to be consistent with the GCL model resolution.

300 For the North Atlantic subtropical region, the GCL posterior flux magnitude is close to that of the ocean prior flux employed (Landschützer et al., 2016), with differences of approximately 0.01 PgC y⁻¹ over the period. Variation among the other inverse flux estimates can reach up to 0.3 PgC y⁻¹ (e.g., between CT and CAMS in 2017), and these differences can be ascribed, in part, to the different underlying prior flux distributions used in the respective inverse analyses (see Sect. 3.2.2). For the North


Atlantic subpolar region, the GCL posterior flux estimate deviates more from the prior flux estimate (e.g., showing differences of up to 0.04 PgC y^{-1}), especially for some years (2012–2017) of the analysis. The majority of flux estimates for the North Atlantic subpolar region are in closer accord (Fig. 2b) with differences of less than 0.2 PgC y^{-1} (the CT estimate is an exception indicating variations of greater than 0.3 PgC y^{-1} from the other estimates). A potential reason for the anomalous behaviour of the CT estimate in the North Atlantic is the underlying prior flux uncertainties used in the analysis which give a loose constraint on the prior ocean fluxes and allow the ocean fluxes deviate far from the prior influenced by the atmospheric CO_2 signals (Jacobson et al., 2020).

We also note that Peylin et al. (2013) have suggested that significant inter-annual variability in atmospheric inverse estimates is a potential indicator of ‘flux leakage’, where significant variability of terrestrial carbon fluxes in combination with sparse atmospheric sampling can result in misattribution of carbon flux estimates between land and ocean. To assess the significance of flux leakage in our GCL analyses, we have calculated estimates of the diagnostic recommended by Peylin et al. (2013) (i.e., the correlation between the annual total land and total ocean fluxes) for the Northern Hemisphere as a whole (Equator to 90°N), and also by latitudinal region. Estimates of this diagnostic are relatively low for our GCL analyses (values of 0.2 and 0.5 for the subpolar and subtropical regions) indicating low potential for flux leakage. As a point of comparison, Peylin et al. (2013) note that six out of eleven atmospheric inverse analyses in their model inter-comparison reported correlation coefficients of greater than 0.5.

320

3.2.1 Long term mean

Figure 3 provides a comparison of the following GCL flux estimates and associated characteristics for the North Atlantic subtropical and subpolar regions for the period 2000–2017: (i) the long term mean of air–sea CO_2 flux estimates (the underlying data are tabulated in Table 2); (ii) the estimated inter-annual variability (IAV) of fluxes (Table 3); and (iii) the long term trends (Table 4). The IAV is calculated following methods of Rödenbeck et al. (2015) (i.e., derived from the standard deviation of the residuals of a 12 month running mean over the CO_2 flux time series).



330 **Figure 3.** Comparison of CO₂ ocean flux metrics for the 2000–2017 period for North Atlantic subtropics (left panels) and subpolar regions (right panels). Metrics shown are the long term mean (panels (a) and (b)); interannual variability (IAV) (panels (c) and (d)); and long term trend (panels (e) and (f)). The GCL estimates (red stars) are shown in comparison to other atmospheric inverse analyses (red symbols), surface ocean pCO₂ products (blue) and global ocean biogeochemistry models (GOBMs, purple). Also shown are the estimated mean values from each sub-group of analyses (filled cross symbols) with their minimum–maximum range. Circled symbols in panel (e) and (f) indicate a statistically significant trend.

335

340

345 **Table 2.** Summary metrics of GEOSChem–LETKF North Atlantic (NA) CO₂ flux estimates, and comparison with independent estimates (from atmospheric inverse analyses, surface pCO₂ mappings, and Global Ocean Biogeochemistry models (GOBMs)) for the period 2000–2017. Listed are estimates for the long term mean. The metrics listed in this table are plotted in Fig.3a and 3b.

Long term mean (PgC y⁻¹)		
NA Subtropics (15°N–50°N)	NA Subpolar (50°N–80°N; eastern boundary at 20°E)	
Atmospheric inversions		
-0.255±0.037	-0.203±0.037	<i>This study (GCL)</i> ^a
-0.203	-0.208	CAMS (Chevallier et al. 2019)
-0.315	-0.347	CTE (van der Laan–Luijkx et al. 2017)
-0.307	-0.340	CT (Jacobson et al. 2020)
[-0.315, -0.203]	[-0.347, -0.203]	Range of all atmospheric inverse studies (minimum to maximum)
Surface ocean pCO₂–based flux products		
-0.263	-0.230	pCO ₂ La (Landschutzer et al. 2016)
-0.284	-0.252	pCO ₂ Ro (Rodenbeck et al. 2014)
-0.264	-0.208	CMEMS (Chau et al. 2020)
-0.302	-0.248	CSIR (Gregor et al. 2019)
-0.295	-0.241	JMA (Iida et al. 2015)
-0.309	-0.192	LSCEFFNN (Denvil–Sommer et al. 2019)
-0.193	-0.171	NIES (Zeng et al. 2015)
-0.305	-0.259	Watson et al. (2020)
[-0.309, -0.193]	[-0.259, -0.171]	Range of all pCO ₂ –based representations (minimum to maximum)
Global ocean biogeochemistry models		
-0.150	-0.197	NEMO–PlankTOM5 (Buitenhuis et al. 2010)
-0.238	-0.217	CCSM–BEC (Doney et al. 2009)
-0.342	-0.341	NEMO–PISCES (CNRM) (Séférian et al. 2013)
-0.188	-0.321	MPIOM–HAMOCC (Ilyina et al. 2013)
-0.351	-0.316	NorESM–OC (Schwinger et al. 2016)
-0.205	-0.256	MITgcm–REcoM2 (Hauck et al. 2016)
[-0.351, -0.150]	[-0.341, -0.197]	Range of GOBM studies (minimum to maximum)

^a The uncertainty of the long term mean estimate from the GCL (this study) is calculated as the standard deviation of the annual flux estimates over the (2000–2017) period.

350 We also present in Fig. 3 the equivalent estimates from other independent assessments, including (i) other atmospheric inverse analyses, (ii) surface ocean pCO₂-based analyses, and (iii) analyses from global ocean biogeochemistry models (GOBMs). For the North Atlantic subtropical region, the long term mean of the GCL posterior flux estimate is -0.255 ± 0.037 PgC y⁻¹ (Figure 3a and Table 2). It lies in the range spanned by the other inverse analyses (-0.31 to -0.20 PgC y⁻¹, of slightly larger magnitude than CAMS, but smaller than CT and CTE), and is in good agreement with the mean-value estimates from surface
355 pCO₂-based methods and GOBMs. For the North Atlantic subpolar region, the GCL estimate of the long term mean uptake is -0.203 ± 0.037 PgC y⁻¹ (Table 2), close to the inverse estimate of the CAMS analysis, and of smaller magnitude (by ~ 0.1 PgC y⁻¹) than the inverse estimates of CT and CTE. The GCL estimate is consistent with the mean estimate from pCO₂-based products, and within the range of flux estimates from GOBMs (from -0.341 to -0.197 PgC y⁻¹).

360 3.2.2 Interannual variability

The interannual variability (IAV) of CO₂ flux estimates derived from the GCL is 0.036 ± 0.006 PgC y⁻¹ for the North Atlantic subtropics and 0.034 ± 0.009 PgC y⁻¹ for the North Atlantic subpolar region (Fig. 3c and 3d, Table 3). The IAV estimates from the different inverse analyses for both the North Atlantic subtropics and subpolar regions display a larger range (0.032 to 0.084 PgC y⁻¹ and 0.023 to 0.114 PgC y⁻¹ respectively), than the ranges displayed by GOBMs (0.014 to 0.027 PgC y⁻¹ and 0.015 to
365 0.024 PgC y⁻¹ respectively) and pCO₂-based fluxes (0.029 to 0.050 PgC y⁻¹ and 0.009 to 0.037 PgC y⁻¹ respectively). The larger range of IAV from atmospheric inverse analyses is influenced especially by high magnitude IAV estimates from the CarbonTracker (CT) inverse analysis. Potential causes of the differences among atmospheric inversion between the GCL and CAMS IAV estimates and those of the CarbonTracker estimates are the different prior ocean fluxes employed by the inverse analyses, and the relative weighting assigned to the influence of atmospheric CO₂ observations (Jacobson et al., 2020). The
370 GCL and CAMS estimates use the prior flux of Landschützer et al. (2016), CTE uses the prior flux of Rodenbeck et al. (2014) and the CarbonTracker inversions use the prior flux of Jacobson et al. (2007).

Recent synthesis studies of global ocean carbon fluxes have noted that GOBMs underestimate the magnitude of IAV in comparison to estimates from pCO₂-based mappings and inverse analyses (DeVries et al. 2019, Hauck et al. 2020). An important driver of IAV is the variability in biological carbon export; the lower variability observed in the GOBMs could result
375 from opposing changes in biological vs. circulation impacts on carbon export, which potentially reduces the sensitivity of the GOBM air-sea carbon fluxes to climate variability (Landschützer et al., 2013, DeVries et al., 2019).

380

Table 3. Summary metrics of GEOSChem–LETKF North Atlantic (NA) CO₂ flux estimates, and comparison with independent estimates (from atmospheric inverse analyses, surface pCO₂ mappings, and Global Ocean Biogeochemistry models (GOBMs)) for the period 2000–2017. Listed are estimates for the interannual variability of the regional fluxes over the period. The metrics listed in this table are plotted in Fig. 3 c, d.

385

Interannual Variability (IAV) (PgC y⁻¹)		
NA Subtropics (15°N–50°N)	NA Subpolar (50°N–80°N; eastern boundary at 20°E)	
Atmospheric inversions		
<i>0.036±0.006</i>	<i>0.034±0.009</i>	<i>This study (GCL)^a</i>
0.032	0.023	CAMS (Chevallier et al. 2019)
0.065	0.064	CTE (van der Laan–Luijkx et al. 2017)
0.084	0.114	CT (Jacobson et al. 2020)
[0.032, 0.084]	[0.023, 0.114]	Range of all atmospheric inverse studies (minimum to maximum)
Surface ocean pCO₂–based flux products		
0.038	0.036	pCO ₂ La (Landschutzer et al. 2016)
0.050	0.035	pCO ₂ Ro (Rodenbeck et al. 2014)
0.031	0.018	CMEMS (Chau et al. 2020)
0.035	0.019	CSIR (Gregor et al. 2019)
0.028	0.009	JMA (Iida et al. 2015)
0.037	0.016	LSCEFFNN (Denvil–Sommer et al. 2019)
0.029	0.019	NIES (Zeng et al. 2015)
0.037	0.037	Watson et al. (2020)
[0.029, 0.050]	[0.009, 0.037]	Range of all pCO ₂ –based representations (minimum to maximum)
Global ocean biogeochemistry models		
0.018	0.018	NEMO–PlankTOM5 (Buitenhuis et al. 2010)
0.014	0.015	CCSM-BEC (Doney et al. 2009)
0.027	0.024	NEMO-PISCES (CNRM) (Séférian et al. 2013)
0.016	0.019	MPIOM-HAMOCC (Ilyina et al. 2013)
0.021	0.016	NorESM-OC (Schwinger et al. 2016)
0.017	0.016	MITgcm-REcoM2 (Hauck et al. 2016)
[0.014, 0.027]	[0.015, 0.024]	Range of GOBM studies (minimum to maximum)

^a The uncertainty of the estimated IAV from the GCL (this study) is calculated as the standard deviation of the ensemble posterior fluxes

3.2.3 Estimated Trends of North Atlantic CO₂ Fluxes

390 Our calculations of estimated trends for the 2000–2017 period are presented in Table 4 and Fig. 3e and 3f. We also highlight
in the table and figure panels, the trend estimates that are statistically significant (significant at the 95% confidence level,
Montgomery et al., 2012). Our GCL analyses indicate statistically significant trends for the 2000–2017 period of -0.064 ± 0.007
PgC y⁻¹ decade⁻¹ in the North Atlantic subtropical basin, and 0.063 ± 0.008 PgC y⁻¹ decade⁻¹ in the subpolar region. These
estimated trends are of similar magnitude to those estimated from surface ocean pCO₂ products, but of much larger magnitude
395 (by a factor of 3–4) than decadal trends estimated from the GOBMs (Fig. 3e, Table 4). Our findings are similar to those of
Devries et al. (2019), who noted that decadal trend estimates of North Atlantic CO₂ uptake for the 2000s from the SOCOM
inter-comparison of pCO₂-based flux products were larger than those from the GOBMS in their analysis (see Fig. 3 of their
study).

400

405

410

415

420 **Table 4.** Summary metrics of GEOSChem–LETKF North Atlantic (NA) CO₂ flux estimates, and comparison with independent estimates (from atmospheric inverse analyses, surface pCO₂ mappings, and Global Ocean Biogeochemistry models (GOBMs)) for the period 2000–2017. Listed are estimates for the trend of the regional fluxes over the period. The metrics listed in this table are plotted in Fig. 3 e, f of the main study.

Trend (PgC y⁻¹ decade⁻¹)		
NA Subtropics (15°N–50°N)	NA Subpolar (50°N–80°N; eastern boundary at 20°E)	
Atmospheric inversions		
-0.064±0.007 (S) ^a	-0.063±0.008(S) ^a	<i>This study (GCL)</i> ^b
-0.016	-0.023	CAMS (Chevallier et al. 2019)
-0.068	-0.028	CTE (van der Laan–Luijkx et al. 2017)
-0.067	-0.102	CT (Jacobson et al. 2020)
[-0.016, -0.085]	[-0.023, -0.126]	Range of all Atmospheric inverse studies (minimum to maximum)
Surface ocean pCO₂-based flux products		
-0.068(S)	-0.056(S)	pCO ₂ La (Landschutzer et al. 2016)
-0.070(S)	-0.029	pCO ₂ Ro (Rodenbeck et al. 2014)
-0.057(S)	-0.027(S)	CMEMS (Chau et al. 2020)
-0.063(S)	-0.034(S)	CSIR (Gregor et al. 2019)
-0.048(S)	-0.001	JMA (Iida et al. 2015)
-0.070(S)	-0.021	LSCEFFNN (Denvil–Sommer et al. 2019)
-0.057(S)	-0.037(S)	NIES (Zeng et al. 2015)
-0.069(S)	-0.064(S)	Watson et al. (2020)
[-0.048, -0.070]	[-0.001, -0.064]	Range of all pCO ₂ -based representations (minimum to maximum)
Global ocean biogeochemistry models		
-0.015	-0.023	NEMO–PlankTOM5 (Buitenhuis et al. 2010)
-0.010	0.0002	CCSM–BEC (Doney et al. 2009)
-0.021	-0.002	NEMO–PISCES (CNRM) (Séférian et al. 2013)
-0.014	-0.011	MPIOM–HAMOCC (Ilyina et al. 2013)
-0.036(S)	-0.023	NorESM–OC (Schwinger et al. 2016)
-0.013	-0.014	MITgcm–REcoM2 (Hauck et al. 2016)
[-0.010, -0.036]	[-0.0002, -0.023]	Range of GOBM studies (minimum to maximum)

425 ^a The symbol (S) indicates that the calculated trend is statistically significant (at the 95% confidence interval).

^b The uncertainty of the fitted trend from the GCL estimates is reported as 1 standard deviation of the OLS fitted slope (Montgomery et al. 2012).

4 Summary

In this study we present a new long term estimate of North Atlantic air–sea CO₂ fluxes for recent decades (period 2000–2017) using the atmospheric carbon cycle data assimilation system GEOSChem–LETKF. We focus, in particular, on the specification of prior ocean fluxes, including sensitivity of flux estimates to alternative prior flux distributions, and on the specification of uncertainties associated with ocean fluxes. Towards this we have developed the ‘spread-based’ flux uncertainty scheme which represents the variability among a set of different prior ocean CO₂ flux representations. The scheme ascribes higher levels of uncertainty to regions with larger discrepancies among ocean CO₂ prior flux representation that arise from uncertainties associated with measurement density and pCO₂–interpolation methods (Sect. 2.4). The spread-based flux uncertainty scheme provides improved performance in comparison to schemes with fixed prior flux uncertainty levels, based on an assessment metric of differences in model–observation values for atmospheric CO₂ at North Atlantic measurement sites of the NOAA–GLOBALVIEWCO₂ network (Sect. 3.1). It provides a valuable new means of specifying prior flux uncertainties for atmospheric inverse analyses of ocean CO₂ fluxes.

We have used the spread–based flux uncertainty scheme in the GEOSChem–LETKF to derive estimates of CO₂ fluxes in the North Atlantic for the 2000–2017 period. Long term mean estimates of the regional ocean CO₂ uptake are -0.255 ± 0.037 PgC y⁻¹ for the North Atlantic subtropics and -0.203 ± 0.037 PgC y⁻¹ for North Atlantic subpolar region, and are consistent with recent regional flux estimates from surface pCO₂–based methods and global ocean biogeochemistry models (GOBMs). The GEOSChem–LETKF estimates of interannual variability in air–sea CO₂ fluxes are 0.036 ± 0.006 PgC y⁻¹ (North Atlantic subtropics) and 0.034 ± 0.009 PgC y⁻¹ (North Atlantic subpolar). In common with estimates from other atmospheric CO₂ inverse studies, the magnitude of IAV derived from the GEOSChem–LETKF is larger than corresponding estimates from GOBMs. Our GEOSChem–LETKF estimates also indicate statistically significant trends of increasing CO₂ uptake for the North Atlantic subtropical and subpolar region (estimated trend of -0.064 ± 0.007 and -0.063 ± 0.008 PgC y⁻¹ decade⁻¹ respectively). These trends are of comparable magnitude to those estimated from surface pCO₂–based flux products, but much larger than those derived from global ocean biogeochemistry models for the 2000–2017 period. Estimates of inter–annual variability and long term trends derived from our GEOSChem–LETKF analyses are generally more robust for the North Atlantic subtropics than for the subpolar region, and characterized by smaller uncertainty bounds. Limiting factors affecting estimates for the North Atlantic subpolar region include higher levels of uncertainty associated with specification of prior fluxes (Fig. 1), and the observational uncertainty at the atmospheric measurement CO₂ sites in these high northern latitudes (Table A1). The number of regional atmospheric CO₂ measurement sites available to constrain North Atlantic subpolar fluxes are also relatively few in comparison to the subtropical region. Improved ocean CO₂ flux estimates and associated metrics for this North Atlantic region will be obtained by provision of additional high accuracy marine boundary layer CO₂ measurements for the region from fixed surface sites and from ships and buoys (Wanninkhof et al., 2019).

460 Appendix A: The Local Ensemble Transform Kalman Filter (LETKF) system

Here we briefly describe the LETKF system used for estimation of surface CO₂ fluxes. The methodology follows that of Hunt et al. (2007) and Miyoshi et al. (2007), and additional detail is provided in these publications. The LETKF has been previously used in meteorological forecasting, and more recently in atmospheric CO₂ data assimilation (e.g., Liu et al. 2019, 2016; Kang et al. 2012). The LETKF provides iterative estimates of the time evolution of the system state, x , (here representing the
 465 gridscale surface carbon fluxes, of dimension m). Each step involves a forecast stage (based on a physical model of the system evolution) and a state estimation stage (the ‘analysis’ step), which combines system observations, y (of dimension n), together with the background forecast, x^b , to derive the improved state estimate. The observation operator H provides the mapping from the state space to the observation space; in this study H is provided by the GEOSChem atmospheric model. In the analysis step, the surface carbon flux estimates are obtained by minimization of a cost function (Eq. A1) which accounts for deviations
 470 of the system state x , from the background forecast, x^b , and for the mismatch between observations (y) and their modeled representations (Hx):

$$J(x) = (x - x^b)^T P^{-1} (x - x^b) + (y - Hx)^T R^{-1} (y - Hx) \quad (\text{A1})$$

B represents the background flux covariance matrix, and R represents the observation covariance matrix.

In the LETKF system, an ensemble of model simulations is used to calculate the sample mean and covariance of the system state; thus, the background state x^b is given by $(x^{b(i)}: i = 1, 2, \dots, k)$ for k ensemble members. The sample mean \bar{x}^b and
 475 covariance P^b of the background state vector given by:

$$\bar{x}^b = k^{-1} \sum_{i=1}^k x^{b(i)} \quad (\text{A2})$$

$$\begin{aligned} P^b &= (k - 1)^{-1} \sum_{i=1}^k (x^{b(i)} - \bar{x}^b)(x^{b(i)} - \bar{x}^b)^T \\ &= (k - 1)^{-1} X^b (X^b)^T \end{aligned} \quad (\text{A3})$$

X^b is an $m \times k$ matrix whose i th column is $x^{b(i)} - \bar{x}^b$. P^b is the background state covariance matrix ($m \times m$).

Similarly the analysis state is represented by the ensemble $(x^{a(i)}: i = 1, 2, \dots, k)$ with its sample mean and covariance given by:

$$\bar{x}^a = k^{-1} \sum_{i=1}^k x^{a(i)} \quad (\text{A4})$$

$$\begin{aligned} P^a &= (k - 1)^{-1} \sum_{i=1}^k (x^{a(i)} - \bar{x}^a)(x^{a(i)} - \bar{x}^a)^T \\ &= (k - 1)^{-1} X^a (X^a)^T \end{aligned} \quad (\text{A5})$$

where X^a is the $m \times k$ matrix whose i th column is $x^{a(i)} - \bar{x}^a$.

The analysis state and covariance \bar{x}^a and P^a are updated based on the background information \bar{x}^b and observations y through the following equations:

$$\bar{x}^a = \bar{x}^b + P^a H^T R^{-1} (y - H \bar{x}^b) \quad (\text{A6})$$

$$P^a = (I + P^b H^T R^{-1} H)^{-1} P^b \quad (\text{A7})$$

The ensemble $y^{b(i)}$ of background observation vectors is defined by:

$$y^{b(i)} = H(x^{b(i)}) \quad (\text{A8})$$

$$H(\bar{x}^b + X^b w) \approx \bar{y}^b + Y^b w \quad (\text{A9})$$

where Y^b is the $n \times k$ matrix whose i th column is $(y^{b(i)} - \bar{y}^b)$, and w is a Gaussian random vector with mean $\bar{w}^b = 0$ and covariance $\tilde{P}^b = (k - 1)^{-1} I$. Then the analogues of analysis equations (6) and (7) are:

$$\bar{w}^a = \tilde{P}^a (Y^b)^T R^{-1} (y - \bar{y}^b) \quad (\text{A10})$$

$$\tilde{P}^a = [(k - 1)I + (Y^b)^T R^{-1} Y^b]^{-1} \quad (\text{A11})$$

Following Hunt et al. (2007) and Miyoshi et al. (2007) (refer to these publications for the complete LETKF derivation) the overall analysis equation is:

$$x = \bar{x}^b + X^b [\tilde{P}^a (Y^b)^T R^{-1} (y - \bar{y}^b) + [(k - 1)\tilde{P}^a]^{1/2}] \quad (\text{A12})$$

The LETKF allows flexibility in the choice of observations to be assimilated at each grid point, based on the distance (r) of the observations from the gridpoint. The localization weighting function $f(r)$ is given by:

$$f(r) = \exp\left(-\frac{r^2}{2L^2}\right) \quad (\text{A13})$$

where L is an observation localization length which can be predefined to determine the outer boundary of the influence of the observations; i.e., the localization weighting function drops to zero at a value of

$$r = 2. \sqrt{\frac{10}{3}} L \quad (\text{A14})$$

The observation localization is realized by multiplying the inverse of the localization function $f(r)$ with the observational error covariance R .

The parameter L represents the horizontal localization radius, and is set to 2000 km for this study, following Liu et al. (2016). The localization radius is used in the LETKF in a latitude-dependent weighting function which characterizes the spatial scale of the region within which atmospheric CO₂ observations are assimilated at each gridpoint (Miyoshi et al. 2007).

495

Table A1. Atmospheric CO₂ measurement sites^a

Site code	Longitude (degrees)	Latitude (degrees)	Altitude (m)	Site name	U ^b (ppm)
ABP	-38.16	-12.76	6	Arembepe, Bahia	1.04
ALT	-62.51	82.45	195	Alert, Nunavut	1.34
AMY	126.33	36.54	125	Anmyeon-do	8.88
ASC	-14.40	-7.97	90	Ascension Island	0.66
ASK	5.63	23.26	2715	Assekrem	0.80
AZR	-27.08	38.75	24	Terceira Island, Azores	2.26
BAL	16.67	55.50	28	Baltic Sea	5.50
BCS	-110.20	23.30	14	Baja California Sur	3.42
BGU	3.23	41.97	13	Begur	3.93
BHD	174.87	-41.41	90	Baring Head Station	1.12
BKT	100.32	-0.20	875	Bukit Kototabang	3.49
BME	-64.65	32.37	17	St. Davids Head, Bermuda	2.57
BMW	-64.88	32.27	60	Tudor Hill, Bermuda	2.12
BRW	-156.60	71.32	28	Barrow Atmospheric Baseline Observatory	1.88
BSC	28.67	44.18	5	Black Sea, Constanta	9.88
CBA	-162.72	55.20	25	Cold Bay, Alaska	2.41
CFA	147.06	-19.28	5	Cape Ferguson, Queensland	1.04
CGO	144.68	-40.68	164	Cape Grim, Tasmania	0.40
CHR	-157.15	1.70	5	Christmas Island	0.60
CIB	-4.93	41.81	850	Centro de Investigacion de la Baja Atmosfera (CIBA)	3.97
CPT	18.49	-34.35	260	Cape Point	0.74
CRI	73.83	15.08	66	Cape Rama	3.47
CRZ	51.85	-46.43	202	Crozet Island	0.49
CYA	110.52	-66.28	55	Casey, Antarctica	0.29
DRP	-64.91	-55.00	10	Drake Passage	0.41
DSI	116.73	20.70	8	Dongsha Island	3.46
EIC	-109.45	-27.15	55	Easter Island	1.80
ELL	0.96	42.58	2005	Estany Llong	2.41
ESP	-126.53	49.38	47	Estevan Point, British Columbia	1.49
FKL	25.67	35.34	150	Finokalia, Crete	3.34

GMI	144.66	13.39	6	Mariana Islands	2.22
GPA	131.05	-12.25	37	Gunn Point	2.02
HBA	-26.21	-75.61	35	Halley Station, Antarctica	0.16
HPB	11.02	47.80	990	Hohenpeissenberg	6.71
HSU	-124.73	41.05	8	Humboldt State University	5.78
HUN	16.65	46.95	344	Hegyhatsal	6.00
ICE	-20.29	63.40	127	Storhofdi, Vestmannaeyjar	2.03
IZO	-16.48	28.30	2378	Izana, Tenerife, Canary Islands	1.21
KEY	-80.20	25.67	6	Key Biscayne, Florida	4.14
KUM	-154.82	19.52	8	Cape Kumukahi, Hawaii	1.77
KZD	75.57	44.45	412	Sary Taukum	3.19
KZM	77.88	43.25	2524	Plateau Assy	3.00
LJO	-117.26	32.87	20	La Jolla, California	2.72
LLB	-112.45	54.95	546	Lac La Biche, Alberta	8.91
LLN	120.86	23.46	2867	Lulin	5.27
LMP	12.61	35.51	50	Lampedusa	2.08
MAA	62.87	-67.62	42	Mawson Station, Antarctica	0.32
MEX	-97.31	18.98	4469	High Altitude Global Climate Observation Center	1.33
MHD	-9.90	53.33	26	Mace Head, County Galway	3.23
MID	-177.37	28.22	8	Sand Island, Midway	1.39
MKN	37.30	-0.06	3649	Mt. Kenya	1.98
MLO	-155.58	19.53	3402	Mauna Loa, Hawaii	0.63
MQA	158.97	-54.48	13	Macquarie Island	0.33
NAT	-35.26	-5.52	20	Farol De Mae Luiza Lighthouse	1.44
NMB	15.03	-23.58	461	Gobabeb	1.13
NWR	-105.58	40.05	3526	Niwot Ridge, Colorado	1.88
OBN	36.60	55.12	484	Obninsk	6.49
OTA	142.82	-38.52	50	Otway, Victoria	17.45
OXX	11.81	50.03	1185	Ochsenkopf	8.18
PAL	24.12	67.97	570	Pallas-Sammaltunturi, GAW Station	3.72
PDM	0.14	42.94	2877	Pic Du Midi	2.71
POC	-145.13	14.97	20	Pacific Ocean	1.47
PSA	-64.00	-64.92	15	Palmer Station, Antarctica	0.23

PTA	-123.73	38.95	22	Point Arena, California	5.50
RK1	-177.90	-29.20	12	Kermadec Island	2.23
RPB	-59.43	13.17	20	Ragged Point	0.83
SDZ	117.12	40.65	298	Shangdianzi	9.57
SEY	55.53	-4.68	7	Mahe Island, Seychelles	0.98
SGP	-97.48	36.62	374	Southern Great Plains, Oklahoma	4.91
SHM	174.10	52.72	28	Shemya Island, Alaska	2.91
SIS	-1.26	60.09	33	Shetland Islands	2.87
SMO	-170.57	-14.25	47	Tutuila, American Samoa	1.19
STM	2.00	66.00	7	Ocean Station M	2.03
SUM	-38.42	72.60	3215	Summit	1.32
SYO	39.58	-69.00	16	Syowa Station, Antarctica	0.23
TAC	1.14	52.52	236	Tacolneston	6.78
TAP	126.13	36.73	21	Tae-ahn Peninsula	6.90
THD	-124.15	41.05	112	Trinidad Head, California	4.54
TIK	128.89	71.60	29	Hydrometeorological Observatory of Tiksi	2.64
USH	-68.31	-54.85	32	Ushuaia	1.41
UTA	-113.72	39.90	1332	Wendover, Utah	2.65
UUM	111.10	44.45	1012	Ulaan Uul	2.78
WIS	34.78	30.86	482	Weizmann Institute of Science at the Arava Institute, Ketura	2.39
WLG	100.92	36.27	3815	Mt. Waliguan	2.26
WPC	167.50	-29.86	10	Western Pacific Cruise	1.70
ZEP	11.89	78.91	479	Ny-Alesund, Svalbard	1.82

500

^a Source reference: Cooperative Global Atmospheric Data Integration Project, 2018. Version: obspack_co2_1_GLOBALVIEWplus_v4.2_2019-03-19 (<https://doi.org/10.25925/20190319>)

^b The specification of observational uncertainty U on atmospheric CO₂ measurements (and represented in matrix R of Eq. A1) is calculated as the standard deviation of measurement variability and using the detrended and deseasonalized CO₂ time series at each measurement site (following methods of Chevallier et al., 2010).

505

510 **Table A2:** Model–Observation mismatch in atmospheric CO₂ concentrations (unit: ppm) at North Atlantic sites (average over year 2003).
 GCL model values are derived from the a posteriori model analyses associated with the sensitivity analyses of Sect. 3.1. Atmospheric CO₂
 observations are from the NOAA–GLOBALVIEW network described in Sect. 2.3.

Sensitivity Analyses (Sect. 2.3)	North Atlantic Sites						Global Network Average
	BMW	KEY	AVI	AZR	IZO	ICE	
U1Ta	0.81	0.88	0.76	1.49	1.83	1.20	0.54
U2Ta	0.71	0.82	0.69	1.42	1.74	1.13	0.44
U3Ta	0.58	-0.20	0.36	0.85	1.64	0.40	0.01
U1La	0.92	0.93	0.76	1.60	1.94	1.44	0.61
U2La	0.84	0.87	0.68	1.53	1.86	1.38	0.51
U3La	0.52	-0.27	0.21	0.84	1.58	0.58	0.04
U1Ro	0.74	0.74	0.54	1.35	1.66	0.98	0.35
U2Ro	0.66	0.68	0.47	1.28	1.57	0.91	0.26
U3Ro	0.55	-0.28	0.21	0.73	1.48	0.27	-0.11

515 *Data Availability.*

Data sources: (i) Atmospheric CO₂ measurements were taken from obspack_co2_1_GLOBALVIEWplus_v4.2_2019-03-19 (<https://doi.org/10.25925/20190319>); (ii) Prior ocean flux oc_v1.7 from Rödenbeck et al. (2013) taken from <http://www.bgc-jena.mpg.de/CarboScope/>. Prior ocean flux Landschützer et al. (2016) taken from https://www.nodc.noaa.gov/ocads/oceans/SPCO2_1982_present_ETH_SOM_FFN.html. Prior ocean flux from Takahashi et al. (2009) taken from <ftp://ftp.as.harvard.edu/gcgrid/geos-chem>. (iii) CarbonTracker CT2019 results provided by NOAA ESRL, Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov>. CTE flux estimates downloaded from ftp://ftp.wur.nl/carbontracker/data/fluxes/data_flux1x1_monthly/ on 24 November 2020. The flux estimates from CAMS (v18r2) taken from <https://apps.ecmwf.int/datasets/data/cams-ghg-inversions/>. (iv) The model CO₂ fluxes for JULES (land) and GOBMs (ocean) taken from (Le Quéré et al., 2018). Time series of reconstructed surface ocean *p*CO₂ and CO₂ fluxes (LSCEFFNN) from Denvil–Sommer et al., 2019 are the first version of CMEMS, downloaded from <http://marine.copernicus.eu/services-portfolio/access-to-products/>. The products from Iida et al., 2015 downloaded from http://www.data.jma.go.jp/gmd/kaiyou/english/co2_flux/co2_flux_data_en.html. The products from Zeng et al., 2015 downloaded from <https://db.cger.nies.go.jp/DL/10.17595/20201020.001.html.en>. The products from CMEMS, CSIR, Watson taken from Friedlingstein et al., 2020.

Author contributions. ZC and PS designed the study. ZC, PS, JZ and NZ developed the model. ZC, PS, AW, and US discussed the design of simulations. ZC performed the simulations and analysis and wrote the initial manuscript. All authors contributed to the writing of the paper.

Competing interests. The authors declare that they have no conflict of interest.

535 *Disclaimer.* The work reflects only the author's view, the European Commission and their executive agency are not responsible for any use that may be made of the information the work contains.

Acknowledgements. This work was performed using the High Performance Computing Cluster at the University of East Anglia.

Financial support. It was supported through the UK Natural Environment Research Council grants NE/K002473/1(RAGNARoCC).

References

- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W.-
- 545 J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot,
- 550 D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multidecade record of high-quality fCO₂ data in version 3 of the Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., and Seferian,
- 555 R.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225-6245, <https://doi.org/10.5194/bg-10-6225-2013>, 2013.
- Buitenhuis, E. T., Rivkin, R. B., Sailley, S., and Le Quéré, C.: Biogeochemical fluxes through microzooplankton, *Global*

- Biogeochemical Cycles, 24, GB4015, <https://doi.org/10.1029/2009GB003601>, 2010.
- Carouge, C., Bousquet, P., Peylin, P., Rayner, P., and Ciais, P.: What can we learn from European continuous atmospheric CO₂ measurements to quantify regional fluxes--Part 1: Potential of the 2001 network, *Atmospheric Chemistry & Physics*, 10, 3107-3117, <https://doi.org/10.5194/acp-10-3107-2010>, 2010.
- Chatterjee, A., Engelen, R. J., Kawa, S. R., Sweeney, C., and Michalak, A. M.: Background error covariance estimation for atmospheric CO₂ data assimilation, *Journal of Geophysical Research: Atmospheres*, 118, 10140-10154, <https://doi.org/10.1002/jgrd.50564>, 2013.
- 565 Chau, T. T., Gehlen, M., and Chevallier, F.: Global Ocean Surface Carbon Product MULTIOBS_GLO_BIO_CARBON_SURFACE_REP_015_008, E.U. Copernicus Marine Service Information, available at: https://resources.marine.copernicus.eu/?option=com_cswandview=detailsandproduct_id=MULTIOBS_G, last access:16 November 2020.
- Chevallier, F., Ciais, P., Conway, T., Aalto, T., Anderson, B., Bousquet, P., Brunke, E., Ciattaglia, L., Esaki, Y., and Fröhlich, M.: CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements, *Journal of Geophysical Research: Atmospheres*, 115, D21307, <https://doi.org/10.1029/2010JD013887>, 2010.
- 570 Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., and Bousquet, P.: Toward robust and consistent regional CO₂ flux estimates from in situ and spaceborne measurements of atmospheric CO₂, *Geophysical Research Letters*, 41, 1065-1070, <https://doi.org/10.1002/2013GL058772>, 2014.
- 575 Chevallier, F., Remaud, M., O'Dell, C. W., Baker, D., Peylin, P., and Cozic, A.: Objective evaluation of surface-and satellite-driven carbon dioxide atmospheric inversions, *Atmospheric Chemistry and Physics*, 19, 14233-14251, <https://doi.org/10.5194/acp-19-14233-2019>, 2019.
- Clark, D., Mercado, L., Sitch, S., Jones, C., Gedney, N., Best, M., Pryor, M., Rooney, G., Essery, R., Blyth, E., Boucher, O., Harding, R., Huntingford, C., and Cox, P.: The Joint UK Land Environment Simulator (JULES), model description - Part 2: Carbon fluxes and vegetation dynamics, *Geoscientific Model Development*, 4, 701-722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R., and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmospheric Chemistry and Physics*, 16, 3825-3841, <https://doi.org/10.5194/acp-16-3825-2016>, 2016.
- 585 Denvil-Sommer, A., Gehlen, M., Vrac, M., and Mejia, C.: LSCEFFNN-v1: a two-step neural network model for the reconstruction of surface ocean pCO₂ over the global ocean, *Geosci. Model Dev.*, 12, 2091-2105, <https://doi.org/10.5194/gmd-12-2091-2019>, 2019.
- DeVries, T., Le Quéré, C., Andrews, O., Berthet, S., Hauck, J., Ilyina, T., Landschützer, P., Lenton, A., Lima, I. D., and Nowicki, M.: Decadal trends in the ocean carbon sink, *Proceedings of the National Academy of Sciences*, 116, 11646-11651, <https://doi.org/10.1073/pnas.1900371116>, 2019.
- 590 Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.:

- Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air–sea CO₂ fluxes: Physical climate and atmospheric dust, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 640-655, <https://doi.org/10.1016/j.dsr2.2008.12.006>, 2009.
- 595 Feng, L., Palmer, P. I., Parker, R. J., Deutscher, N. M., Feist, D., Kivi, R., Morino, I., and Sussmann, R.: Estimates of European uptake of CO₂ inferred from GOSAT XCO₂ retrievals: sensitivity to measurement bias inside and outside Europe, *Atmospheric Chemistry and Physics*, 16, 1289-1302, <https://doi.org/10.5194/acp-16-1289-2016>, 2016.
- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneeth, A., Arora, V., Bates,
600 N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, k., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson,
605 E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon budget 2020. *Earth System Science Data*, 12, 3269-3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- Gaubert, B., Stephens, B. B., Basu, S., Chevallier, F., Deng, F., Kort, E. A., Patra, P. K., Peters, W., Rödenbeck, C., and Saeki,
610 T.: Global atmospheric CO₂ inverse models converging on neutral tropical land exchange, but disagreeing on fossil fuel and atmospheric growth rate, *Biogeosciences*, 16, 117-134, <https://doi.org/10.5194/bg-16-117-2019>, 2019.
- Gregor, L., Lebehot A., Kok, S., Monteiro P.: A comparative assessment of the uncertainties of global surface ocean CO₂ estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) -have we hit the wall?. *Geoscientific Model Development*, 12, 5113–5136, <https://doi.org/10.5194/gmd-12-5113-2019>, 2019.
- 615 Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., and Lindsay, K.: Oceanic sources, sinks, and transport of atmospheric CO₂, *Global Biogeochemical Cycles*, 23, GB1005, <https://doi.org/10.1029/2008GB003349>, 2009.
- Gruber, N., Landschützer, P., and Lovenduski, N. S.: The variable Southern Ocean carbon sink, *Annual Review of Marine Science*, 11, 159-186, <https://doi.org/10.1146/annurev-marine-121916-063407>, 2019.
- 620 Gurney, K., Law, R., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y. H., Ciais, P., Fan, S., Fung, I. Y., Gloor, M., Heimann, M., Higuchi, K., John, J., Maki, T., Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B.C., Randerson, J., Sarmiento, J., Taguchi, S., Takahashi, T., and Yuan, C.: Towards robust regional estimates of CO₂ sources and sinks using atmospheric transport models, *Nature*, 415, 626–630, <https://doi.org/10.1038/415626a>, 2002.
- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y. H., Ciais, P., and
625 Fan, S.: TransCom 3 CO₂ inversion intercomparison: 1. Annual mean control results and sensitivity to transport and prior flux

- information, *Tellus B: Chemical and Physical Meteorology*, 55B, 555-579, <https://doi.org/10.3402/tellusb.v55i2.16728>, 2003.
- Hauck, J., Köhler, P., Wolf-Gladrow, D., and Völker, C.: Iron fertilisation and century-scale effects of open ocean dissolution of olivine in a simulated CO₂ removal experiment, *Environmental Research Letters*, 11, 024007, <https://doi.org/10.1088/1748-9326/11/2/024007>, 2016.
- 630 Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C., Bopp, L., Chau, T. T., Gürses, Ö., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R.: Consistency and challenges in the ocean carbon sink estimate for the Global Carbon Budget. *Frontiers in Marine Science*, 7, 852, <https://doi.org/10.3389/fmars.2020.571720>, 2020.
- Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter, *Physica D: Nonlinear Phenomena*, 230, 112-126, <https://doi.org/10.1016/j.physd.2006.11.008>, 2007.
- 635 Iida, Y., Kojima, A., Takatani, Y., Nakano, T., Midorikawa, T., and Ishii, M.: Trends in pCO₂ and sea-air CO₂ flux over the global open oceans for the last two decades. *Journal of Oceanography*, doi:10.1007/s10872-015-0306-4, 2015.
- Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287-315, <https://doi.org/10.1029/2012MS000178>, 2013.
- Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global Biogeochemical Cycles*, 21, GB1019, <https://doi.org/10.1029/2005GB002556>, 2007.
- 645 Jacobson, A. R., Schuldt, K. N., Miller, J. B., Oda, T., Tans, P., Andrews, A., Mund, J., Ott, L., Collatz, G. J., Aalto, T., Afshar, S., Aikin, K., Aoki, S., Apadula, F., Baier, B., Bergamaschi, P., Beyersdorf, A., Biraud, S. C., Bollenbacher, A., Bowling, D., Brailsford, G., Abshire, J. B., Chen, G., Chen, H., Chmura, L., Climadat, S., Colomb, A., Conil, S., Cox, A., Cristofanelli, P., Cuevas, E., Curcoll, R., Sloop, C. D., Davis, K., Wekker, S. D., Delmotte, M., DiGangi, J. P., Dlugokencky, E., Ehleringer, J., Elkins, J. W., Emmenegger, L., Fischer, M. L., Forster, G., Frumau, A., Galkowski, M., Gatti, L. V., Gloor, E., Griffiths, T., Hammer, S., Haszpra, L., Hatakka, J., Heliasz, M., Hensen, A., Hermanssen, O., Hints, E., Holst, J., Jaffe, D., Karion, A., Kawa, S. R., Keeling, R., Keronen, P., Kolari, P., Kominkova, K., Kort, E., Krummel, P., Kubistin, D., Labuschagne, C., Langenfelds, R., Laurent, O., Laurila, T., Lauvaux, T., Law, B., Lee, J., Lehner, I., Leuenberger, M., Levin, I., Levula, J., Lin, J., Lindauer, M., Loh, Z., Lopez, M., Lund Myhre, C., Machida, T., Mammarella, I., Manca, G., Manning, A., Manning, A., Marek, M. V., Marklund, P., Martin, M. Y., Matsueda, H., McKain, K., Meijer, H., Meinhardt, F., Miles, N., Miller, C. E.,
- 655 Mölder, M., Montzka, S., Moore, F., Morgui, J., Morimoto, S., Munger, B., Necki, J., Newman, S., Nichol, S., Niwa, Y., O'Doherty, S., Ottosson-Löfvenius, M., Paplawsky, B., Peischl, J., Peltola, O., Pichon, J., Piper, S., Plass-Dölmer, C., Ramonet, M., Reyes-Sanchez, E., Richardson, S., Riris, H., Ryerson, T., Saito, K., Sargent, M., Sasakawa, M., Sawa, Y., Say, D., Scheeren, B., Schmidt, M., Schmidt, A., Schumacher, M., Shepson, P., Shook, M., Stanley, K., Steinbacher, M., Stephens, B., Sweeney, C., Thoning, K., Torn, M., Turnbull, J., Tørseth, K., van den Bulk, P., van der Laan-Luijkx, I. T., van Dinter, D.,

- 660 Vermeulen, A., Viner, B., Vitkova, G., Walker, S., Weyrauch, D., Wofsy, S., Worthy, D., Young, D., and Zimnoch, M.: CarbonTracker CT2019, Model published by NOAA Earth System Research Laboratory, Global Monitoring Division, <https://doi.org/10.25925/39m3-6069>, 2020.
- Jones, S. D., Le Quéré, C., and Rödenbeck, C.: Autocorrelation characteristics of surface ocean pCO₂ and air-sea CO₂ fluxes. *Global biogeochemical cycles*, 26, 2, <https://doi.org/10.1029/2010GB004017>, 2012
- 665 Kang, J. S., Kalnay, E., Liu, J., Fung, I., Miyoshi, T., and Ide, K.: Variable localization” in an ensemble Kalman filter: Application to the carbon cycle data assimilation." *Journal of Geophysical Research: Atmospheres*, 116, D9 <https://doi.org/10.1029/2010JD014673>, 2011.
- Kang, J. S., Kalnay, E., Miyoshi, T., Liu, J., and Fung, I.: Estimation of surface carbon fluxes with an advanced data assimilation methodology, *Journal of Geophysical Research: Atmospheres*, 117, D24101, <https://doi.org/10.1029/2012JD018259>, 2012.
- 670 Khatiwala, S. P., Tanhua, T., Mikaloff Fletcher, S. E., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G., Murata, A., and Ríos, A.: Global ocean storage of anthropogenic carbon, *Biogeosciences*, 10, 2169-2191, <https://doi.org/10.5194/bg-10-2169-2013>, 2013.
- Landschützer, P., Gruber, N., Bakker, D. C., Schuster, U., Nakaoka, S. I., Payne, M. R., Sasse, T. P., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793-7815, <https://doi.org/10.5194/bg-10-7793-2013>, 2013.
- 675 Landschuetzer, P., Gruber, N., and Bakker, D. C.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochemical Cycles*, 30, 1396-1417, <https://doi.org/10.1002/2015GB005359>, 2016.
- Landschützer, P., Ilyina, T., and Lovenduski, N. S.: Detecting Regional Modes of Variability in Observation-Based Surface Ocean pCO₂, *Geophysical Research Letters*, 46, 2670-2679, <https://doi.org/10.1029/2018GL081756>, 2019.
- 680 Law, R. M., Ziehn, T., Matear, R. J., Lenton, A., Chamberlain, M. A., Stevens, L. E., Ying-Ping, W., Srbinovsky, J., Bi, D., and Yan, H.: The carbon cycle in the Australian Community Climate and Earth System Simulator (ACCESS-ESM1)–Part 1: Model description and pre-industrial simulation, *Geoscientific Model Development*, 10, 2567-2590, <https://doi.org/10.6194/gmd-10-2567-2017>, 2017.
- 685 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., Boden, T. A., Tans, P. P., Andrews, O. D., Arora, V. K., Bakker, D. C. E., Barbero, L., Becker, M., Betts, R. A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Cosca, C. E., Cross, J., Currie, K., Gasser, T., Harris, I., Hauck, J., Haverd, V., Houghton, R. A., Hunt, C. W., Hurtt, G., Ilyina, T., Jain, A. K., Kato, E., Kautz, M., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lima, I., Lombardozi, D., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., Padin, X. A., Peregon, A., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Reimer, J., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., van Heuven, S., Viovy, N., Vuichard, N., Walker, A. P., Watson, A. J., Wiltshire, A. J., Zaehle, S., and Zhu, D.: Global carbon budget 2017, *Earth System Science*

- Data, 10, 405-448, <https://doi.org/10.5194/essd-10-405-2018>, 2018.
- 695 Lebehot, A. D., Halloran, P. R., Watson, A. J., McNeill, D., Ford, D. A., Landschützer, P., Lauvset, S. K., and Schuster, U.: Reconciling observation and model trends in North Atlantic surface CO₂, *Global Biogeochemical Cycles*, 33, 1204-1222, <https://doi.org/10.1029/2019GB006186>, 2019.
- Liu, J., Bowman, K. W., and Lee, M.: Comparison between the Local Ensemble Transform Kalman Filter (LETKF) and 4D-Var in atmospheric CO₂ flux inversion with the Goddard Earth Observing System-Chem model and the observation impact diagnostics from the LETKF, *Journal of Geophysical Research: Atmospheres*, 121, 13,066-013,087, <https://doi.org/10.1002/2016JD025100>, 2016.
- 700 Liu, Y., Kalnay, E., Zeng, N., Asrar, G., Chen, Z., and Jia, B.: Estimating surface carbon fluxes based on a local ensemble transform Kalman filter with a short assimilation window and a long observation window: an observing system simulation experiment test in GEOS-Chem 10.1, *Geoscientific Model Development*, 12, 2899-2914, <https://doi.org/10.5194/gmd-12-2899-2019>, 2019.
- 705 Macovei, V. A., Hartman, S. E., Schuster, U., Torres-Valdés, S., Moore, C. M., and Sanders, R. J.: Impact of physical and biological processes on temporal variations of the ocean carbon sink in the mid-latitude North Atlantic (2002–2016), *Progress in Oceanography*, 180, 102223, <https://doi.org/10.1016/j.pocean.2019.102223>, 2020.
- McKinley, G. A., Fay, A. R., Takahashi, T., and Metzl, N.: Convergence of atmospheric and North Atlantic carbon dioxide trends on multidecadal timescales, *Nature Geoscience*, 4, 606-610, <https://doi.org/10.1038/ngeo1193>, 2011.
- 710 Mikaloff Fletcher, S., Gruber, N., Jacobson, A. R., Doney, S., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., and Menemenlis, D.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean, *Global Biogeochemical Cycles*, 20, GB2002, <https://doi.org/10.1029/2005GB002530>, 2006.
- Miyoshi, T., Yamane, S., and Enomoto, T.: Localizing the error covariance by physical distances within a local ensemble transform Kalman filter (LETKF), *SOLA*, 3, 089-092, <https://doi.org/10.2151/sola.2007-023>, 2007.
- 715 Montgomery, D. C., Peck, E. A., and Vining, G. G.: *Introduction to Linear Regression Analysis*, 5th Edition, John Wiley & Sons, 2012.
- Nassar, R., Jones, D. B., Suntharalingam, P., Chen, J. M., Andres, R. J., Wecht, K. J., Yantosca, R. M., Kulawik, S. S., Bowman, K. W., and Worden, J. R.: Modeling global atmospheric CO₂ with improved emission inventories and CO₂ production from the oxidation of other carbon species, *Geoscientific Model Development*, 3, 689-716, <https://doi.org/10.5194/gmd-3-689-2010>, 2010.
- 720 Nassar, R., Jones, D. B. A., Kulawik, S. S., Worden, J. R., Bowman, K. W., Andres, R. J., Suntharalingam, P., Chen, J. M., Brenninkmeijer, C., and Schuck, T.: Inverse modeling of CO₂ sources and sinks using satellite observations of CO₂ from TES and surface flask measurements, *Atmospheric Chemistry and Physics*, 11, 6029-6047, <https://doi.org/10.5194/acp-11-6029-2011>, 2011.
- 725 Peters, W., Miller, J., Whitaker, J., Denning, A., Hirsch, A., Krol, M., Zupanski, D., Bruhwiler, L., and Tans, P.: An ensemble data assimilation system to estimate CO₂ surface fluxes from atmospheric trace gas observations, *Journal of Geophysical*

- Research: Atmospheres, 110, D24304, <https://doi.org/10.1029/2005JD006157>, 2005.
- 730 Peylin, P., Law, R., Gurney, K., Chevallier, F., Jacobson, A., Maki, T., Niwa, Y., Patra, P., Peters, W., and Rayner, P.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO₂ inversions, *Biogeosciences*, 10, 6699–6720, <https://doi.org/10.5194/bg-10-6699-2013>, 2013.
- Rienecker, M. M., Suarez, M., Todling, R., Bacmeister, J., Takacs, L., Liu, H., Gu, W., Sienkiewicz, M., Koster, R., and Gelaro, R.: The GEOS-5 Data Assimilation System: Documentation of Versions 5.0. 1, 5.1. 0, and 5.2. 0, NASA Technical Report Series on Global Modelling and Data Assimilation, NASA/TM–2008–104606, 27, 2008.
- 735 Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: Time-dependent atmospheric CO₂ inversions based on interannually varying tracer transport, *Tellus*, 55B, 488–497, <https://doi.org/10.3402/tellusb.v55i2.16707>, 2003.
- Rödenbeck, C., Keeling, R. F., Bakker, D. C., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean pCO₂ and sea–air CO₂ flux variability from an observation-driven ocean mixed-layer scheme. *Ocean Science*, 9, 2, 193–216, <https://doi.org/10.5194/os-9-193-2013>, 2013.
- 740 Rödenbeck, C., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., Reum, F., Keeling, R. F., and Heimann, M.: Interannual sea-air CO₂ flux variability from an observation-driven ocean mixed–layer scheme, *Biogeosciences*, 11, 4599–4613, <https://doi.org/10.5194/bg-11-4599-2014>, 2014.
- Rödenbeck, C., Bakker, D. C., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P., Metzl, N., Nakaoka, S., and Olsen, A.: Data-based estimates of the ocean carbon sink variability—first results of the Surface Ocean pCO₂ Mapping intercomparison (SOCOM), *Biogeosciences*, 12, 7251–7278, <https://doi.org/10.5194/bg-12-7251-2015>, 2015.
- 745 Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C., Wallace, D. W., and Tilbrook, B.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, <https://doi.org/10.1126/science.1097403>, 2004.
- Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A., González-Dávila, M., Gruber, N., Jones, S., and Krijnen, J.: An assessment of the Atlantic and Arctic sea-air CO₂ fluxes, 1990–2009, *Biogeosciences*, 10, 607–627, <https://doi.org/10.5194/bg-10-607-2013>, 2013.
- 750 Schwinger, J., Goris, N., Tjiputra, J. F., Kriest, I., Bentsen, M., Bethke, I., Ilicak, M., Assmann, K. M., and Heinze, C.: Evaluation of NorESM-OC (versions 1 and 1.2), the ocean carbon–cycle stand–alone configuration of the Norwegian Earth System Model (NorESM1), *Geoscientific Model Development*, 9, 2589–2622, <https://doi.org/10.5194/gmd-9-2589-2016>, 2016.
- 755 Séférian, R., Bopp, L., Gehlen, M., Orr, J. C., Ethé, C., Cadule, P., Aumont, O., y Méliá, D. S., Voldoire, A., and Madec, G.: Skill assessment of three earth system models with common marine biogeochemistry, *Climate Dynamics*, 40, 2549–2573, <https://doi.org/10.1007/s00382-012-1362-8>, 2013.
- Suntharalingam, P., Randerson, J. T., Krakauer, N., Logan, J. A., and Jacob, D. J.: Influence of reduced carbon emissions and oxidation on the distribution of atmospheric CO₂: Implications for inversion analyses, *Global Biogeochemical Cycles*, 19, GB4003, <https://doi.org/10.1029/2005GB002466>, 2005.
- 760 Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R., Feely, R. A.,

- and Sabine, C.: Global sea–air CO₂ flux based on climatological surface ocean pCO₂, and seasonal biological and temperature effects, *Deep Sea Research Part II: Topical Studies in Oceanography*, 49, 1601-1622, [https://doi.org/10.1016/S0967-0645\(02\)00003-6](https://doi.org/10.1016/S0967-0645(02)00003-6), 2002.
- 765 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., and Sabine, C.: Climatological mean and decadal change in surface ocean pCO₂, and net sea–air CO₂ flux over the global oceans, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 554-577, <https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.
- Van Der Laan–Luijkx, I. T., Van Der Velde, I. R., Van Der Veen, E., Tsuruta, A., Stanislawski, K., Babenhauserheide, A.,
770 Zhang, H. F., Liu, Y.-N., He, W., and Chen, H.: The CarbonTracker Data Assimilation Shell (CTDAS) v1. 0: implementation and global carbon balance 2001-2015, *Geoscientific Model Development*, 10, 2785-2800, <https://doi.org/10.5194/gmd-10-2785-2017>, 2017.
- Wanninkhof, R., Pickers, P., Omar, A., Sutton, A., Murata, A., Olsen, A., Bb, S., Tilbrook, B., Munro, D., and Pierrot, D.: A
775 surface ocean CO₂ reference network, SOCONET and associated marine boundary layer CO₂ measurements, *Frontiers in Marine Science*, 6, <https://doi.org/10.3389/fmars.2019.00400>, 2019.
- Watson, A. J., Schuster, U., Bakker, D. C., Bates, N. R., Corbière, A., González-Dávila, M., Friedrich, T., Hauck, J., Heinze, C., and Johannessen, T.: Tracking the variable North Atlantic sink for atmospheric CO₂, *Science*, 326, 1391-1393, <https://doi.org/10.1126/science.1177394>, 2009.
- Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G. C., Landschützer, P., Woolf, D. K., and Goddijn-Murphy,
780 L.: Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon inventory, *Nat. Commun.*, 11, 1–6, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.
- Zeng, J., Nojiri, Y., Nakaoka, Shin-ichiro, Nakajima, H. and Shirai, T.: Surface ocean CO₂ in 1990–2011 modelled using a feed-forward neural network. *Geoscience Data Journal*, 2: 47-51. <https://doi.org/10.1002/gdj3.26>, 2015.