

1 **Different responses of ecosystem CO<sub>2</sub> and N<sub>2</sub>O emissions and CH<sub>4</sub> uptake to**  
2 **seasonally asymmetric warming in an alpine grassland of the Tianshan**  
3 **Mountains**

4 Yanming Gong<sup>1</sup>, Ping Yue<sup>2</sup>, Kaihui, Li<sup>1</sup>, Anwar Mohammat<sup>1\*</sup>, Yanyan Liu<sup>1\*</sup>

5 <sup>1</sup>State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and  
6 Geography, CAS, Urumqi, 830011, China

7 <sup>2</sup>Urat Desert-Grassland Research Station, Northwest Institute of Eco-Environment and  
8 Resources, CAS, Lanzhou, 730000, China

9 \*Correspondence: Anwar Mohammat and Yanyan Liu, Email: liuyany@ms.xjb.ac.cn

10 **Abstract:**

11 An experiment was conducted to investigate the effect of seasonally asymmetric  
12 warming on ecosystem respiration (*Re*), CH<sub>4</sub> uptake, and N<sub>2</sub>O emissions in alpine  
13 grassland of the Tianshan Mountains of Central Asia, from October 2016 to  
14 September 2019. Our results indicated that the annual mean of *Re*, CH<sub>4</sub>, and N<sub>2</sub>O  
15 fluxes in growing season were 42.83 mg C m<sup>-2</sup> h<sup>-1</sup>, -41.57 μg C m<sup>-2</sup> h<sup>-1</sup>, and 4.98 μg N  
16 m<sup>-2</sup> h<sup>-1</sup>, respectively. Furthermore, warming during the non-growing season increased  
17 *Re* and CH<sub>4</sub> uptake by 7.9% and 10.6% in growing season, 10.5% and 9.2% in  
18 non-growing seasons, respectively. However, the increase in N<sub>2</sub>O emission in the  
19 growing season was mainly caused by the warming during the growing season (by  
20 29.7%). the warming throughout the year and warming during the non-growing  
21 season increased N<sub>2</sub>O emissions by 101.9% and 192.3% in non-growing seasons,  
22 respectively. The *Re*, CH<sub>4</sub> uptake, and N<sub>2</sub>O emissions were positively correlated with

23 soil temperature. Our results suggested that *Re*, CH<sub>4</sub> uptake and N<sub>2</sub>O emissions were  
24 regulated by soil temperature, rather than soil moisture, in the case of seasonally  
25 asymmetric warming. In addition, the response rate was defined by the changes in  
26 greenhouse gas fluxes driven by warming. In our field experiment, we observed the  
27 stimulatory effect of warming during the non-growing season on *Re* and CH<sub>4</sub> uptake.  
28 In contrast, the response rates of *Re* and N<sub>2</sub>O emissions were gradually attenuated by  
29 long-term annual warming and the response rate of *Re* was also weakened by  
30 warming over the growing season. These findings highlight the importance of  
31 warming in the non-growing season in regulating greenhouse gas fluxes, a finding  
32 which is crucial for improving our understanding of C and N cycles under the  
33 scenarios of global warming.

34 **Keywords:** Alpine steppe; Extreme climatic event; Greenhouse gas fluxes;  
35 Warming of open-top chambers

## 36 **1. Introduction**

37 Since the industrial revolution, human activities have intensified global warming.  
38 The global surface temperature increased by about 0.85°C from 1880 to 2012 (IPCC,  
39 2013). Furthermore, the temperature is expected that the surface temperature will  
40 increase by about 1.1–6.4°C by the end of this century (IPCC, 2007, 2013). The rise  
41 in atmospheric temperature over the year is not continuous on the temporal scale but  
42 there is asymmetrical warming across the seasons (Xia et al., 2014). The 3rd and 4rd  
43 Assessment Report of the Inter-Governmental Panel on Climate Change (IPCC)  
44 proposed that, against the backdrop of global warming, the temperature change shows

45 that the warming amplitude in the winter is greater than that in the summer, with the  
46 warming amplitude at high latitude being greater than that at low latitude, and  
47 confirmed that the warming shows asymmetric trends on a seasonal scale (Easterling  
48 et al., 1997; IPCC, 2001, 2007).

49 Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) are three of the  
50 major greenhouse gases (GHGs) in the atmosphere that directly cause global climate  
51 warming, with their contributions to global warming being 60%, 20%, and 6%,  
52 respectively (IPCC, 2007, 2013). Experimental warming is known to influence  
53 ecosystem respiration (*Re*), CH<sub>4</sub> uptake, and N<sub>2</sub>O emission (Pärn et al., 2018; Treat et  
54 al., 2018; Wang et al., 2019). Information on *Re*, CH<sub>4</sub> uptake, and N<sub>2</sub>O emission will  
55 enhance our understanding of ecosystem C and N cycling processes and improve our  
56 predictions of the response of ecosystems to global climate change (Li et al., 2020;  
57 Wang et al., 2019).

58 At present, most studies focus on the influence of warming on GHG flux in  
59 terrestrial ecosystems during the summer months (Keenan et al., 2014; Li et al., 2011;  
60 Yang et al., 2014). Nevertheless, data on the influence of asymmetric warming on the  
61 GHG flux on a seasonal scale are scarce. A study of the Alaskan tundra found that  
62 summer warming (using open-top chambers to increase air temperatures in the  
63 growing season) significantly increased *Re* in the growing season by about 20%  
64 (Natali et al., 2011). Compared with the slight effect of winter warming on the  
65 ecosystem respiration in the growing season, warming increased ecosystem  
66 respiration during the snow-covered non-growing season by more than 50% (Natali et

67 al., 2011). Lin et al. (2015) reported that the response of soil CH<sub>4</sub> uptake rates to  
68 temperature increases in alpine meadows of the Qinghai-Tibet Plateau were not  
69 consistent seasonally, with CH<sub>4</sub> uptake in the non-growing season being more  
70 sensitive to temperature (increasing by 162%) than the corresponding value in the  
71 growing season. A study by Cantarel et al. (2012) in an alpine grassland ecosystem  
72 showed that the response of N<sub>2</sub>O emission to warming showed clear seasonal  
73 differences, with the N<sub>2</sub>O emission in the growing season showing significant  
74 differences between the warming treatments, whereas the response of N<sub>2</sub>O emission  
75 to the warming treatments in November was not obvious. A recent study showed that  
76 seasonal variations in carbon flux were more closely related to air temperature in the  
77 meadow steppe (Zhao et al., 2019). Another study found that experimental warming  
78 enhanced CH<sub>4</sub> uptake in the relatively arid alpine steppe, but had no significant effects  
79 on CH<sub>4</sub> emission in the moist swamp meadow (Li et al., 2020). Furthermore, soil CH<sub>4</sub>  
80 uptake was not significantly affected by warming in the alpine meadow of the Tibetan  
81 Plateau (Wu et al., 2020). In contrast, a global meta-analysis showed that  
82 experimental warming stimulates ecosystem respiration in grassland ecosystems, and  
83 the response of ecosystem respiration to warming strongly varies across the different  
84 grassland types, with greater warming responses in cold than in temperate and semi-  
85 arid grasslands (Wang et al., 2019). Across the data set, Li et al. (2020) demonstrated  
86 that N<sub>2</sub>O emissions were significantly enhanced by whole-year warming treatments.  
87 In contrast, no significant effects on soil N<sub>2</sub>O emissions were observed by in  
88 short-season warming.

89 In summary, the GHG fluxes in terrestrial ecosystems shows significant  
90 interannual, and seasonal variations, and its response to warming also varies over  
91 different temporal scales. After long-term uniform warming, the biotic and abiotic  
92 factors have adapted to the temperature increase, and the GHG fluxes response to  
93 increasing temperature is smaller than that in the early stages of warming. For  
94 example, over longer time periods of warming, accelerated carbon decomposition and  
95 increased plant N uptake may decrease soil organic C and N pools (Wu et al., 2012),  
96 and the microbial community with variable C use efficiency may reduce the  
97 temperature sensitivity of heterotrophic respiration (Zhou et al., 2012). Moreover,  
98 climate warming is often unstable, with most of it occurring as extreme events  
99 (Jentsch et al., 2007). The heterogeneity of warming may change the adaptability of  
100 GHG fluxes to warming, and thus affect the carbon and nitrogen cycles in terrestrial  
101 ecosystems. Therefore, we hypothesize the stimulatory effect of warming during the  
102 non-growing season on  $R_e$ ,  $\text{CH}_4$  uptake and  $\text{N}_2\text{O}$  emissions. However, the response  
103 rates of  $R_e$ ,  $\text{CH}_4$  uptake and  $\text{N}_2\text{O}$  emissions were gradually attenuated by long-term  
104 annual warming and warming over the growing season, respectively. Our results will  
105 help to assess this variation with respect to GHG fluxes response to increasing  
106 temperatures against the backdrop of global climate change, by carrying out  
107 seasonally asymmetrical warming studies in alpine grasslands.

## 108 **2. Materials and methods**

109 The experiment was conducted from October 2016 to September 2019 at the  
110 Bayinbuluk Grassland Ecosystem Research Station, Chinese Academy of Sciences

111 (42°52.76' ~ 42°53.17' N, 83°41.90' ~ 83°43.12' E, 2460 m above sea level), which is  
112 located in the southern Tianshan mountains of Central Asia, Xinjiang Uyghur  
113 Autonomous Region of China. Permafrost is present in the Bayinbuluk alpine  
114 grassland, with the average maximum frozen depth (from 2000 to 2011, Zhang et al.,  
115 2018) being more than 250 cm. The mean annual temperature was -4.8 °C per decade,  
116 with the lowest monthly temperature in January (-27.4 °C) and the highest in July  
117 (11.2 °C), and the mean annual precipitation amounted to 265.7 mm, with 78.1%  
118 occurring during the growing season, from June to September (Geng et al., 2019).  
119 Variations in soil temperature, soil moisture, air temperature and precipitation are  
120 shown in Figure S1, S2, S3 and S4, respectively. The site was fenced since 2005, all  
121 the plots were dominated by *Stipa purpurea*, *Festuca ovina*, *Oxytropis glabra*, and  
122 *Potentilla multifida*. The soil was sub-alpine steppe soil, the parent material of the soil  
123 was Loess, and the average annual soil moisture was 5.9% (2017-2019).

124 The open-top chambers (OTCs) were made of 5 mm thick tempered glass. To  
125 reduce the impact of precipitation and snow, the OTC was constructed with a  
126 hexagonal round table which was 100 cm high, and the diagonals of the bottom and  
127 top were 100 cm and 60 cm, respectively. Four treatments were simulated using OTCs:  
128 warming throughout the year (AW), warming in the non-growing season (October 1 to  
129 May 31 of the next year) only (NGW), warming in the growing season (June 1 to  
130 September 30) only (GW) and no warming (NW). After the warming in the NGW or  
131 GW, the tempered glass was removed and the frame was retained. Three replicate  
132 plots were established for each treatment, each plot measuring 1 m × 1 m, with a 3-m

133 wide buffer zone between adjacent plots, making a total of 12 plots. Soil temperature  
134 and soil moisture were measured at a frequency of every half an hour by an outdoor  
135 temperature and humidity data recorder (at 10 cm depth; HOBO U23-001; Onset  
136 Computer Corporation, Bourne, USA). The air temperature and humidity inside the  
137 OTCs is also recorded at a frequency of every half an hour using HOBO Pro  
138 RH/TEMP Data LOGGERS (hanged in the center of the OCTs, 50cm above the  
139 surface; Onset Computer Corporation, Bourne, USA). Soil temperature and air  
140 temperature were increased about 2.3 °C and 4 °C by the warming treatment,  
141 respectively (Figure S1 and S3). Soil moisture was reduced about 5% by the warming  
142 treatment (Figure S2).

143 *Re*, CH<sub>4</sub> and N<sub>2</sub>O fluxes were measured using static chambers, made of PVC  
144 tubing with diameter 0.25 m and height 0.17 m, with one chamber in each of the 12  
145 plots. Gas samples were taken 0, 10, 20 and 30 minutes after the lid of the static  
146 chamber was sealed in between 12:00 and 14:00 (GMT + 8), collecting once or twice  
147 a week. The rates of ecosystem respiration, CH<sub>4</sub> and N<sub>2</sub>O fluxes were calculated  
148 based on the change in concentration of CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> in each chamber over  
149 time by a linear or non-linear equation ( $P < 0.05$ ,  $r^2 > 0.95$ ) (the positive flux values  
150 represent emission, and the negative flux values represent uptake; Liu et al. 2012;  
151 Wang et al. 2013). Concentrations of individual gases in samples were measured using  
152 a gas chromatograph (GC) (Agilent 7890A; Agilent Technologies, Santa Clara, CA,  
153 USA).

154 Effects of seasonally asymmetric warming on *Re*, CH<sub>4</sub> uptake, and N<sub>2</sub>O

155 emissions were analyzed by two-way repeated-measures analysis of variance  
156 (ANOVA). One-way ANOVA was used to compare soil temperature, soil moisture  
157 and air temperature differences, respectively. Nonlinear regression analyses  
158 (Exponential Growth, Single, 3 Parameter) was used to identify the relationship  
159 between ecosystem respiration ( $R_e$ ) and soil temperature (at 10-cm depth) from  
160 October 2016 to September 2019. General linear analyses were used to identify  
161 significant linear correlations and regressions between soil temperature, or soil  
162 moisture variation and the responses of  $\text{CH}_4$  uptake, or  $\text{N}_2\text{O}$  emissions.  
163 Variation-partitioning analysis was used to disentangled the influence of soil  
164 temperature and soil moisture on  $R_e$ ,  $\text{CH}_4$  uptake and  $\text{N}_2\text{O}$  emission under the four  
165 treatments in the growing season and the non-growing season, respectively. The  
166 natural logarithm of the response ratio (RR) was used to reflect the effects of  
167 seasonally asymmetric warming on alpine grassland GHG fluxes (Hedges et al., 1999).  
168 The RR is the ratio of the mean value of the chosen variable in the warming group  
169 ( $\bar{W}_t$ ) to that in the control group (NW;  $\bar{W}_c$ ), and is an index of the effect of  
170 seasonally asymmetric warming on the corresponding variable (Eq. 1). All statistical  
171 analyses were conducted using SPSS (version 20.0) (IBM, Armonk, NY, USA) with  
172 the statistically significant difference threshold set at  $P < 0.05$ .

$$RR = \ln\left(\frac{\bar{W}_t}{\bar{W}_c}\right) = \ln(\bar{W}_t) - \ln(\bar{W}_c) \quad (1)$$

### 174 **3. Results**

175 Our study showed that the Bayinbuluk alpine grassland exhibited a low  $R_e$ , was a  
176 net  $\text{CH}_4$  sink, and a negligible  $\text{N}_2\text{O}$  source. The annual mean values of  $R_e$ ,  $\text{CH}_4$  uptake,



177 and N<sub>2</sub>O emissions in the growing season were 42.83 mg C m<sup>-2</sup> h<sup>-1</sup>, 41.57 μg C m<sup>-2</sup> h<sup>-1</sup>,  
178 and 4.98 μg N m<sup>-2</sup> h<sup>-1</sup>, respectively, from October 2016 to September 2019. One-way  
179 ANOVA results of *Re*, CH<sub>4</sub> uptake and N<sub>2</sub>O emissions among the four warming  
180 treatments were not significant, with the exception that the soil CH<sub>4</sub> uptake in the  
181 growing season 2019 under GW treatment was significantly higher than that of the  
182 AW and NGW treatments ( $P < 0.05$ ). Compared with the control group (NW), the *Re*  
183 was decreased by 7.5% and 4.0% in the growing season and non-growing season,  
184 respectively, under AW and decreased by 2.4% and 8.5% under GW in the growing  
185 season and non-growing season, respectively. However, compared with the control  
186 group, the *Re* under NGW increased by 7.9% and 10.5% in the growing season and  
187 non-growing season, respectively, averaged over the three years (Figure 2 a). The AW  
188 temperature change induced a 6.4% increase in CH<sub>4</sub> uptake in the growing season and  
189 a 3.8% decrease in the non-growing season. The GW treatment resulted in 7.1% and  
190 10.2% increases in CH<sub>4</sub> uptake in the growing season and non-growing season,  
191 respectively. On the contrary, the NGW generated a 10.6% and 9.2% decrease in CH<sub>4</sub>  
192 uptake in the growing season and non-growing season, respectively (Figure 2 b). The  
193 AW and NGW treatments resulted in 5.8% and 2.2% decreases, respectively, in N<sub>2</sub>O  
194 emission in the growing season, and 101.9% and 192.3% increases, respectively, in  
195 N<sub>2</sub>O emission in the non-growing season. Compared with the control, NW group, the  
196 N<sub>2</sub>O emission increased by 29.7% and decreased by 24.4% under GW in the growing  
197 season and non-growing season, respectively (Figure 2 c).

198 The results of two-way repeated measures ANOVA showed significant

199 interannual differences of  $Re$  in the growing season ( $P < 0.05$ , Figure 1 a), whereas  
200 the  $CH_4$  uptake under the warming treatment exhibited significant differences in the  
201 growing season ( $P < 0.01$ ; Figure 1 b), and the interannual  $N_2O$  emission showed  
202 significant differences in both the growing season and non-growing season ( $P < 0.05$ ,  
203 Figure 1 c). Therefore, interannual variation was larger than the impact of the  
204 warming treatment (for  $Re$  and  $N_2O$  emissions, Figure 1), whereas the warming  
205 treatment had a significant impact on  $CH_4$  uptake. Under the four warming treatments,  
206  $Re$  was significantly exponential growth correlated with soil temperature ( $P < 0.05$ ;  
207 Figure S5 a). we observed increasing  $CH_4$  uptake with increasing soil temperature ( $P$   
208  $< 0.05$ ; Figure S5 b). On the other hand, the  $N_2O$  emission showed a significantly  
209 positive linear correlation with soil temperature, but only under NGW ( $P < 0.05$ ;  
210 Figure S5 c).

211 The soil moisture was reduced by warming in the alpine grassland (Figure S2).  
212 However,  $Re$ ,  $CH_4$  uptake and  $N_2O$  emission were no significant linearly correlated  
213 with soil moisture, respectively ( $P \geq 0.05$ ; Figure S6). We disentangled the influence  
214 of soil temperature and soil moisture on  $Re$ ,  $CH_4$  uptake, and  $N_2O$  emission by  
215 variation-partitioning analysis under the four treatments in the growing season and the  
216 non-growing season (Figure 4). Our results showed that, under the NGW treatment,  
217  $Re$ ,  $CH_4$  uptake, and  $N_2O$  emission in the non-growing season were more influenced  
218 by soil temperature than by soil moisture. Under the GW treatment, there was the  
219 single effect of soil temperature on  $CH_4$  uptake and  $N_2O$  emission in the non-growing  
220 season. In contrast, there were the joint effects of soil temperature and moisture on  $Re$

221 in the non-growing season under the GW treatment.  $Re$  in the growing season was  
222 influenced more by soil moisture than soil temperature under the GW treatment.  
223 Annual  $Re$  under the AW treatment was influenced by the joint effects of soil  
224 temperature and moisture.

#### 225 **4. Discussion**

226 Our study found that the response rate of  $Re$  to temperature significantly  
227 decreased with the increase in soil temperature ( $\Delta ST_{AW}$  and  $\Delta ST_{GW}$ ) under AW and  
228 GW treatments, respectively (Figure 3 a, c;  $P < 0.05$ ). This finding indicated that the  
229 response of  $Re$  to soil temperature became less and less sensitive to soil temperature  
230 with warming throughout the year (or the growing season) in the alpine grasslands.  
231 On the contrary, NGW significantly increased the response rate of  $Re$  to temperature  
232 change ( $\Delta ST_{NGW}$ ), indicating that warming in the non-growing season amplified the  
233 sensitivity of  $Re$  to temperature change (Figure 3 b,  $P < 0.05$ ). In addition, Zou et al.  
234 (2018) showed that the soil fluxes of  $CO_2$  increased exponentially with increasing  
235 temperature, but warming decreased the temperature sensitivity by 23% in the  
236 grassland. Furthermore, Natali et al. (2011) also confirmed that, compared with the  
237  $CO_2$  flux in the growing season, the  $CO_2$  flux in the nongrowing season was more  
238 sensitive to the temperature increase.

239 Ecosystem  $CH_4$  flux is the net result of  $CH_4$  production and consumption,  
240 occurring simultaneously under the action of methanogenic archaea and  
241 methane-oxidizing bacteria (e.g., Mer and Roger, 2001). In addition, our results  
242 demonstrated that warming increased  $CH_4$  uptake in the growing season, but  
243 decreased  $CH_4$  uptake in the non-growing season in the alpine grassland, findings

244 similar to those from other grassland ecosystems (Lin et al., 2015; Wu et al., 2020;  
245 Zhu et al., 2015). Our results also demonstrated that seasonally asymmetric warming  
246 did not significantly affect the response rate of CH<sub>4</sub> uptake (Figure 3 d-f,  $P > 0.05$ ).  
247 CH<sub>4</sub> flux depended on temperature, pH, and the availability of substrate (e.g., Treat et  
248 al., 2015). The CH<sub>4</sub> uptake observed during the three growing season and  
249 non-growing season implied that the alpine grassland soil could act as an atmospheric  
250 CH<sub>4</sub> sink, a finding which agrees with the results of many previous studies in similar  
251 regions (Wei et al., 2015; Zhao et al., 2017). Hu et al. (2016) suggested that  
252 asymmetrical responses of CH<sub>4</sub> fluxes to warming and cooling should be taken into  
253 account when evaluating the effects of climate change on CH<sub>4</sub> uptake in the alpine  
254 meadow on the Tibetan plateau. Unlike CH<sub>4</sub> flux in alpine grasslands, Treat et al.  
255 (2018) confirmed that wetland was a small CH<sub>4</sub> source in the non-growing season,  
256 whereas uplands varied from CH<sub>4</sub> sinks to CH<sub>4</sub> sources. The latest research confirmed  
257 that warming in the Arctic had become more apparent in the non-growing season than  
258 in the typical growing season (Bao et al., 2020). Hereby, Bao et al. (2020) found that  
259 the CH<sub>4</sub> emissions during the spring thaw and the autumn freeze contributed  
260 approximately one-quarter of the annual total CH<sub>4</sub> emissions. That experimental  
261 warming is stimulating soil CH<sub>4</sub> uptake in the growing season implies that the  
262 grasslands of the Bayinbuluk may have the potential to remove more CH<sub>4</sub> from the  
263 atmosphere under future global warming conditions.

264 Furthermore, with the increased variation in soil temperature, the response rate of  
265 N<sub>2</sub>O emission gradually decreased under AW treatment (Figure 3 g,  $P < 0.05$ ). Our  
266 results suggested that the response of N<sub>2</sub>O emission to temperature increase was  
267 limited by the warming that occurred throughout the year. However, our results

268 displayed N<sub>2</sub>O emission peaks during the freeze–thaw periods (e.g., May 2017, June  
269 2018 and April 2019). Warming increased N<sub>2</sub>O emissions in the thawing period due to  
270 disruption of the gas diffusion barrier and greater C and N availability for microbial  
271 activity (Nyborg et al., 1997). Wagner-Riddle et al. (2017) also demonstrated that the  
272 magnitude of the freeze/thaw-induced N<sub>2</sub>O emissions was associated with the number  
273 of days with soil temperatures below 0°C. Pärn et al. (2018) found that N<sub>2</sub>O emission  
274 from organic soils increases with rising soil NO<sub>3</sub><sup>-</sup>, follows a bell-shaped distribution  
275 with soil moisture. Another study has shown that a whole - year warming treatment  
276 significantly increased N<sub>2</sub>O emissions, but daytime, night-time or short - season  
277 warming did not have significant effects (Li et al., 2020). In addition, Cantarel et al.  
278 (2010) suggested that the N<sub>2</sub>O flux from cool and upland grasslands may be driven  
279 primarily by response to elevated temperature under projected future climate  
280 conditions.

## 281 **5. Conclusions**

282 In summary, the effect of seasonally asymmetrical warming on *Re* and N<sub>2</sub>O  
283 emission was obvious, unlike the situation with CH<sub>4</sub> uptake. The *Re* and N<sub>2</sub>O  
284 emission were able to adapt to continuous warming, resulting in a reduced response  
285 rates of the *Re* and N<sub>2</sub>O emission to temperature increase. Warming in the  
286 non-growing season increased the temperature dependence of the *Re*. Thus, we  
287 believe that the study of climate change should pay greater attention to warming in the  
288 non-growing season, to avoid underestimating the greenhouse effect on *Re* in alpine  
289 grasslands.

290 **Data availability**

291 The measured CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes and soil temperature and soil water  
292 content data are available in Zenodo (<http://doi.org/10.5281/zenodo.4244207>).

293 **Author contributions**

294 GYM, LYY and MA conceive the research question, designed the study approach,  
295 led the field survey, ensured data curation and conducted formal analysis. YP and  
296 LKH assisted with data collection and analysis. GYM wrote the first draft of the paper,  
297 and all co-authors contributed to writing review and editing.

298 **Competing interests**

299 The authors declare that they have no conflicts of interest.

300 **Acknowledgments**

301 This work was supported by the NSFC Program (41603084, 41703131,  
302 41673079).

303 **References**

- 304 Bao, T., Xu, X. Y., Jia, G. S., Billesbach, D. P. and Sullivan, R. C.: Much stronger  
305 tundra methane emissions during autumn-freeze than spring-thaw. *Glob. Change*  
306 *Biol.*, 27, 376–387, doi: 10.1111/GCB.15421, 2020.
- 307 Cantarel, A. A. M., Bloor, J. M. G., Deltroy, N. and Soussana, J. -F.: Effects of  
308 Climate Change Drivers on Nitrous Oxide Fluxes in an Upland Temperate  
309 Grassland. *Ecosystems*, 14, 223–233, doi: 10.1007/s10021-010-9405-7, 2011.
- 310 Cantarel, A. A. M., Bloor, J. M. G., Pommier, T., Guillaumaud, N., Moiro, C.,  
311 Soussana, J. F. and Poly, F.: Four years of experimental climate change modifies the

312 microbial drivers of N<sub>2</sub>O fluxes in an upland grassland ecosystem. *Glob. Change*  
313 *Biol.*, 18, 2520–2531, doi: 10.1111/j.1365-2486.2012.02692.x, 2012.

314 Chen, W., Zheng, X., Chen, Q., Wolf, B., Butterbach-Bahl, K., Brueggemann, N., and  
315 Lin, S.: Effects of increasing precipitation and nitrogen deposition on CH<sub>4</sub> and N<sub>2</sub>O  
316 fluxes and ecosystem respiration in a degraded steppe in Inner Mongolia, China.  
317 *Geoderma*, 192, 335–340, doi: 10.1016/j.geoderma.2012.08.018, 2013.

318 Easterling, W. E., Hays, C. J., Easterling, M. M., and Brandle, J. R.: Modelling the  
319 effect of shelterbelts on maize productivity under climate change: An application of  
320 the EPIC model. *Agr. Ecosyst. Environ.*, 61, 163–176, doi:  
321 10.1016/S0167-8809(96)01098-5, 1997.

322 Geng, F. Z., Li, K. H., Liu, X. J., Gong, Y. M., Yue, P., Li, Y. G. and Han, W. X.:  
323 Long-term effects of N deposition on N<sub>2</sub>O emission in an alpine grassland of  
324 Central Asia. *Catena*, 182, 104100, doi: 10.1016/j.catena.2019.104100, 2019.

325 Hedges, L. V., Gurevitch, J. and Curtis, P. S.: The meta-analysis of response ratios in  
326 experimental ecology. *Ecology*, 80, 1150–1156, doi: 10.2307/177062, 1999.

327 Hu, Y. G., Wang, Q., Wang, S. P., Zhang, Z. H., Dijkstra, F. A., Zhang, Z. S., Xu, G. P.,  
328 Duan, J. C., Du, M. Y. and Niu, H. S.: Asymmetric responses of methane uptake to  
329 climate warming and cooling of a Tibetan alpine meadow assessed through a  
330 reciprocal translocation along an elevation gradient. *Plant Soil*, 402, doi:  
331 10.1007/s11104-016-2791-7, 263–275, 2016.

332 IPCC, 2001. *Climate change 2001: Impacts, adaptation and vulnerability*,  
333 *Contribution of Working Group II to the Third Assessment Report of the*

334 Intergovernmental Panel on Climate Change. Cambridge University Press,  
335 Cambridge, UK, and New York, USA, 2001. No. of pages: 1032.

336 IPCC, 2007. Climate Change 2007: The Physical Science Basis. Contribution of  
337 Working Group I to the Fourth Assessment Report of the Intergovernmental Panel  
338 on Climate Change. Cambridge University Press, Cambridge, United Kingdom and  
339 New York, NY, USA, pp. 996.

340 IPCC, 2013. Climate Change 2013: The Physical Science Basis. Contribution of  
341 Working Group I to the Fifth Assessment. Report of the Intergovernmental Panel on  
342 Climate Change. Cambridge University Press, Cambridge, pp. 1535.

343 Jentsch, A., Kreyling, J. and Beierkuhnlein, C.: A new generation of climate change  
344 experiments: events, not trends. *Front. in Ecol. Environ.*, 5, 315–324, doi:  
345 10.1890/1540-9295, 2007.

346 Keenan, T. F., Gray, J., Friedl, M. A., Toomey, M., Bohrer, G., Hollinger, D. Y.,  
347 Munger, J. W., O’Keefe, J., Schmid, H. P., Wing, I. S., Yang, B. and Richardson, A.  
348 D.: Net carbon uptake has increased through warming-induced changes in  
349 temperate forest phenology. *Nat. Clim. Change*, 4, 598–604, doi:  
350 10.1038/nclimate2253, 2014.

351 Li, F., Yang, G. B., Peng, Y. F., Wang, G. Q., Qin, S. Q., Song, Y. T., Fang, K., Wang,  
352 J., Yu, J. C., Liu, L., Zhang, D. Y., Chen, K. L., Zhou, G. Y., and Yang, Y. H.:  
353 Warming effects on methane fluxes differ between two alpine grasslands with  
354 contrasting soil water status. *Agr. Forest Meteorol.*, 290, 107988, doi:  
355 10.1016/j.agrformet.2020.107988, 2020.



356 Li, L. F., Zheng, Z. Z., Wang, W. J., Biederman, J. A., Xu, X. L., Ran, Q. W., Qian, R.  
357 Y., Xu, C., Zhang, B., Wang, F., Zhou, S. T., Cui, L. Z., Che, R. X., Hao, Y. B., Cui,  
358 X. Y., Xu, Z. H. and Wang, Y. F.: Terrestrial N<sub>2</sub>O emissions and related functional  
359 genes under climate change: A global meta-analysis. *Global Change Biol.*, 26, 931–  
360 943, doi: 10.1111/gcb.14847, 2020.

361 Li, N., Wang, G. X., Yang, Y., Gao, Y. H. and Liu, G. S.: Plant production, and carbon  
362 and nitrogen source pools are strongly intensified by experimental warming in  
363 alpine ecosystems in the Qinghai-Tibet Plateau. *Soil Biol. Biochem.*, 43, 942–953,  
364 doi: 10.1016/j.soilbio.2011.01.009, 2011.

365 Lin, X. W., Wang, S. P., Hu, Y. G., Luo, C. Y., Zhang Z. H., Niu, H. S. and Xie, Z. B.:  
366 Experimental warming increases seasonal methane uptake in an alpine meadow on  
367 the Tibetan Plateau. *Ecosystems*, 18, 274–286, doi: 10.1007/s10021-014-9828-7,  
368 2015.

369 Liu, C., Wang, K., and Zheng, X.: Responses of N<sub>2</sub>O and CH<sub>4</sub> fluxes to fertilizer  
370 nitrogen addition rates in an irrigated wheat-maize cropping system in northern  
371 China. *Biogeosciences*, 9, 839–850, doi: 10.5194/bg-9-839-2012, 2012.

372 Mer, J. L., and Roger, P.: Production, oxidation, emission and consumption of  
373 methane by soils: a review. *Eur. J. Soil Biol.*, 37, 25–50, doi:  
374 10.1016/S1164-5563(01)01067-6, 2001.

375 Natali, S. M., Schuur, E. A. G., Trucco, C., Pries, C. E. H., Crummer, K. G., and  
376 Lopez, A. F. B.: Effects of experimental warming of air, soil and permafrost on  
377 carbon balance in Alaskan tundra. *Glob. Change Biol.*, 17, 1394–1407, doi:

378 10.1111/j.1365-2486.2010.02303.x, 2011.

379 Nyborg, M., Laidlaw, J. W., Solberg, E. D. and Malhi, S. S.: Denitrification and  
380 nitrous oxide emissions from a Black Chernozemic soil during spring thaw in  
381 Alberta. *Can. J. Soil Sci.*, 77, 153–160, doi: 10.4141/S96-105, 1997.

382 Pärn, J., Verhoeven, J. T. A., Butterbach-Bahl, K., Dise, N. B., Ullah, S., Aasa, A.,  
383 Egorov, S., Espenberg, M., Jarveoja, J., Jauhiainen, J., Kasak, K., Klemedtsson, L.,  
384 Kull, A., Laggoun-Defarge, F., Lapshina, E. D., Lohila, A., Lohmus, K., Maddison,  
385 M., Mitsch, W. J., Muller, C., Niinemets, U., Osborne, B., Pae, T., Salm, J. O.,  
386 Sgouridis, F., Sohar, K., Soosaar, K., Storey, K., Teemusk, A., Tenywa, M. M.,  
387 Tournebize, J., Truu, J., Veber, G., Villa, J. A., Zaw, S. S., and ander, U.:  
388 Nitrogen-rich organic soils under warm well-drained conditions are global nitrous  
389 oxide emission hotspots. *Nat. Commun.*, 9, 1135, doi:  
390 10.1038/s41467-018-03540-1, 2018.

391 Treat, C. C., Bloom, A. A., and Marushchak, M. E.: Non-growing season methane  
392 emissions-a significant component of annual emissions across northern ecosystems.  
393 *Glob. Change Biol.*, 24, 3331–3343, doi: 10.1111/gcb.14137, 2018.

394 Treat, C. C., Natali, S. M., Ernakovich, J., Iversen, C. M., Lupascu, M., McGuire, A.  
395 D., Norby, R. J., Chowdhury, T. R., Richter, A., Santruckova, H., Schadel, C.,  
396 Schuur, E. A. G., Sloan, V. L., Turetsky, M. R., and Waldrop, M. P.: A pan-Arctic  
397 synthesis of CH<sub>4</sub> and CO<sub>2</sub> production from anoxic soil incubations. *Glob. Change*  
398 *Biol.*, 21, 2787–2803, doi: 10.1111/gcb.12875, 2015.

399 Wagner-Riddle, C., Congreves, K. A., Abalos, D., Berg, A. A., Brown, S. E.,

400 Ambadan, J. T., Gao, X. P., and Tenuta, M. Globally important nitrous oxide  
401 emissions from croplands induced by freeze-thaw cycles. *Nat. Geosci.*, 10, 279–286,  
402 doi: 10.1038/Ngeo2907, 2017.

403 Wang, K., Zheng, X. H., Pihlatie, M., Vesala, T., Liu, C. Y., Haapanala, S.,  
404 Mammarella, I., Rannik, U., Liu, H. Z.: Comparison between static chamber and  
405 tunable diode laser-based eddy covariance techniques for measuring nitrous oxide  
406 fluxes from a cotton field. *Agr. Forest Meteorol.*, 171, 9–19, doi:  
407 10.1016/j.agrformet.2012.11.009, 2013.

408 Wang, N., Quesada, B., Xia, L. L., Butterbach-Bahl, K., Goodale, C. L., and Kiese, R.:  
409 Effects of climate warming on carbon fluxes in grasslands- A global meta-analysis.  
410 *Glob. Change Biol.*, 25, 1839–1851, doi: 10.1111/gcb.14603, 2019.

411 Wei, D., Tenzin-Tarchen, X. -R., Wang, Y. S., and Wang, Y. H.: Considerable methane  
412 uptake by alpine grasslands despite the cold climate: in situ measurements on the  
413 central Tibetan Plateau, 2008–2013. *Glob. Chang. Biol.*, 21, 777–788, doi:  
414 10.1111/gcb.12690, 2015.

415 Wu, H. B., Wang, X. X., Ganjurjav, H., Hu, G. Z., Qin, X. B. and Gao, Q.: Effects of  
416 increased precipitation combined with nitrogen addition and increased temperature  
417 on methane fluxes in alpine meadows of the Tibetan Plateau. *Sci. Total Environ.*,  
418 705, 135818, doi: 10.1016/j.scitotenv.2019.135818, 2020.

419 Wu, Z., Dijkstra, P., Koch, G. W., and Hungate, B. A.: Biogeochemical and ecological  
420 feedbacks in grassland responses to warming. *Nat. Clim. Change*, 2, 458–461, doi:  
421 10.1038/nclimate1486, 2012.

422 Xia, J. Y., Chen, J. Q., Piao, S. L., Ciais P., Luo Y. Q. and Wan, S. Q.: Terrestrial  
423 carbon cycle affected by non-uniform climate warming. *Nat. Geosci.*, 7, 173–180,  
424 doi: 10.1038/ngeo2093 2014.

425 Yang, Y. H., Li, P., Ding, J. Z., Zhao, X., Ma, W. H., Ji, C. J. and Fang, J. Y.: Increased  
426 topsoil carbon stock across China's forests. *Glob. Change Biol.*, 20, 2687–2696, doi:  
427 10.1111/gcb.12536, 2014.

428 Zhang, Y. F., Hao, J. S., Huang, F. R., Li, L. H.: Controlling and Influencing Factors  
429 on the Basic Features of Seasonally Frozen Soil in Kaidu River Basin. *Journal of*  
430 *China Hydrology*, 38, 12–18, 2018, in Chinese.

431 Zhao, H. C., Jia, G. S., Wang, H. S., Zhang, A. Z., and Xu, X. Y.: Seasonal and  
432 interannual variations in carbon fluxes in East Asia semi-arid grasslands. *Sci. Total*  
433 *Environ.*, 668, 1128–1138, doi: 10.1016/j.scitotenv.2019.02.378, 2019.

434 Zhao, Z. Z., Dong, S. K., Jiang, X. M., Liu, S. L., Ji, H. Z., Li, Y., Han, Y. H., and Sha,  
435 W.: Effects of warming and nitrogen deposition on CH<sub>4</sub>, CO<sub>2</sub> and N<sub>2</sub>O emissions in  
436 alpine grassland ecosystems of the Qinghai-Tibetan Plateau. *Sci. Total Environ.*,  
437 592, 565–572, doi: 10.1016/j.scitotenv.2017.03.082, 2017.

438 Zhou, J. Z., Xue, K., Xie, J. P., Deng, Y., Wu, L. Y., Cheng, X. H., Fei, S. F., Deng, S.  
439 P., He, Z. L., Van Nostrand, J. D., and Luo, Y. Q.: Microbial mediation of carbon-  
440 cycle feedbacks to climate warming. *Nat. Clim. Change*, 2, 106–110, doi:  
441 10.1038/nclimate1331, 2012.

442 Zhu, X. X., Luo, C. Y., Wang, S. P., Zhang, Z. H., Cui, S. J., Bao, X. Y., Jiang, L. L.,  
443 Li, X. N., Wang, Q., and Zhou, Y.: Effects of warming, grazing/cutting and nitrogen

444 fertilization greenhouse gas fluxes during growing seasons in an alpine. Agric. For.


445 Meteorol., 214–215, 506–514, doi: 10.1016/j.agrformet.2015.09.008, 2015.

446 Zou, J. L., Tobin, B., Luo, Y. Q. and Osborne, B.: Differential responses of soil CO<sub>2</sub>

447 and N<sub>2</sub>O fluxes to experimental warming. Agr. Forest Meteorol., 259, 11–22, doi:

448 10.1016/j.agrformet.2018.04.006, 2018.


449



450

451 Figure 1 Monthly variation of a). ecosystem respiration (*Re*), b). CH<sub>4</sub> uptake and c).

452 N<sub>2</sub>O emissions under the four treatments from October 2016 to September 2019. AW,  
453 warming throughout the year; NGW, warming in the non-growing season only; GW,  
454 warming in the growing season only; NW, non-warming. The blue arrows indicate  
455 warming effects. The data points represent mean  $\pm$  standard error, SE. The tables  
456 illustrate the tests of significance for year (Y) and warming (W) on *Re*, CH<sub>4</sub> uptake  
457 and N<sub>2</sub>O emission by two-way repeated-measures analysis of variance (ANOVA) in  
458 the growing season (GS) and the non-growing season (NGS), respectively; \**P* < 0.05;  
459 \*\**P* < 0.01; ns, non-significant.



460

461

462

463

464

465

466

467

468

Figure 2 Boxplot presentation of variations in ecosystem respiration ( $Re$ ),  $CH_4$

uptake, and  $N_2O$  emission under four treatments in the growing season and

non-growing season from October 2016 to September 2019. The median and mean are

represented by the black and red lines in the box, respectively. The box (the


interquartile range) represents the middle 50% of the data, whereas the whiskers

represent the ranges for the bottom 25% and the top 25% of the data values, excluding

outliers. GS, growing season; NGS, non-growing season; AW, warming throughout

the year; NGW, warming in the non-growing season only; GW, warming in the

469 growing season only; NW, non-warming. No significant differences among AW, NGW,  
 470 GW, and NW were reported from ANOVA; data points are the mean  $\pm$  standard error.  
 471 One-way ANOVA results of  $Re$ ,  $CH_4$  uptake and  $N_2O$  emissions among the four  
 472 warming treatments were not significant, except that the  $CH_4$  uptake in the GS 2019  
 473 under the GW treatment was significantly higher than that of AW and NGW treatment  
 474 ( $P < 0.05$ ).



475  
 476 Figure 3 Response (presented by linear regression) of variation in ecosystem  
 477 respiration ( $Re$ ),  $CH_4$  uptake, and  $N_2O$  emission to changes in soil temperature under  
 478 AW, NGW and GW conditions in the alpine grassland, from 2016 to 2019. RR, the  
 479 natural logarithm of the response ratio of the mean value of the chosen variable in the  
 480 warming group to that in the control (NW) group.  $\Delta ST_{AW}$ , soil temperature of AW



481 minus that of NW;  $\Delta ST_{CW}$ , soil temperature of NGW minus that of NW;  $\Delta ST_{WW}$ , soil  
 482 temperature of GW minus that of NW; AW, warming throughout the year; NGW,  
 483 warming in the non-growing season only; GW, warming in the growing season only;  
 484 NW, non-warming.

	<i>Re</i>	CH <sub>4</sub> flux	N <sub>2</sub> O flux
NGW-NGS %	a 41.6    c 0.8    b -1.6	75.0   -4.1   0.8	43.8   -1.4   -1.9
NGW-GS %	6.4   6.3   9.0	-2.9   0.2   -2.7	1.3   4.0   -0.3
GW-NGS %	0.7   36.5   22.2	51.3   7.4   0.9	29.6   10.2   -2.0
GW-GS %	22.6   -12.4   23.4	-2.6   0.4   -2.4	3.8   0.9   <0.1
AW-AY %	9.5   22.3   10.1	15.3   6.2   -0.9	7.7   4.5   -1.9
NW-AY %	7.6   26.7   5.0	18.5   4.7   -0.9	21.5   -3.7   3.5

485

486 Figure 4 Influence of soil temperature and soil moisture on ecosystem respiration  
 487 (*Re*), CH<sub>4</sub> uptake, and N<sub>2</sub>O emission by variation-partitioning analysis under four  
 488 treatments in the growing season and non-growing season. a, Single effect of soil  
 489 temperature (%); b, single effect of soil moisture (%); c, joint effects of soil  
 490 temperature and moisture (%); NGW-NGS, greenhouse gas fluxes in non-growing  
 491 season under non-growing season warming treatment; NGW-GS, greenhouse gas  
 492 fluxes in growing season under non-growing season warming treatment; GW-NGS,  
 493 greenhouse gas fluxes in non-growing season under growing season warming  
 494 treatment; GW-GS, greenhouse gas fluxes in growing season under growing season

495 warming treatment; AW-AY, annual greenhouse gas fluxes under annual warming  
496 treatment; NW-AY, annual greenhouse gas fluxes without warming.