

Geographic variability in freshwater methane hydrogen isotope ratios and its implications for global isotopic source signatures

Peter M.J. Douglas¹, Emerald Stratigopoulos^{1*}, Jenny Park¹, Dawson Phan¹

¹Earth and Planetary Sciences, McGill University, Montreal, H3A 0E8, Canada

* Now at Department of Earth Sciences, University of Toronto, Toronto, M5S 3B1, Canada

Correspondence to: Peter M. J. Douglas (peter.douglas@mcgill.ca)

Abstract. There is growing interest in developing spatially resolved methane (CH₄) isotopic source signatures to aid in geographic source attribution of CH₄ emissions. CH₄ hydrogen isotope measurements ($\delta^2\text{H-CH}_4$) have the potential to be a powerful tool for geographic differentiation of CH₄ emissions from freshwater environments, as well as other microbial sources. This is because microbial $\delta^2\text{H-CH}_4$ values are partially dependent on the $\delta^2\text{H}$ of environmental water ($\delta^2\text{H-H}_2\text{O}$), which exhibits large and well-characterized spatial variability globally. We have refined the existing global relationship between $\delta^2\text{H-CH}_4$ and $\delta^2\text{H-H}_2\text{O}$ by compiling a more extensive global dataset of $\delta^2\text{H-CH}_4$ from freshwater environments, including wetlands, inland waters, and rice paddies, comprising a total of 129 different sites, and compared these with measurements and estimates of $\delta^2\text{H-H}_2\text{O}$, as well as $\delta^{13}\text{C-CH}_4$ and $\delta^{13}\text{C-CO}_2$ measurements. We found that estimates of $\delta^2\text{H-H}_2\text{O}$ explain approximately 42% of the observed variation in $\delta^2\text{H-CH}_4$, with a flatter slope than observed in previous studies. The inferred global $\delta^2\text{H-CH}_4$ vs. $\delta^2\text{H-H}_2\text{O}$ regression relationship is not sensitive to using either modelled precipitation $\delta^2\text{H}$ or measured $\delta^2\text{H-H}_2\text{O}$ as the predictor variable. The slope of the global freshwater relationship between $\delta^2\text{H-CH}_4$ and $\delta^2\text{H-H}_2\text{O}$ is similar to observations from incubation experiments, but is different from pure culture experiments. This result is consistent with previous suggestions that variation in the $\delta^2\text{H}$ of acetate, controlled by environmental $\delta^2\text{H-H}_2\text{O}$, is important in determining variation in $\delta^2\text{H-CH}_4$. The relationship between $\delta^2\text{H-CH}_4$ and $\delta^2\text{H-H}_2\text{O}$ leads to significant differences in the distribution of freshwater $\delta^2\text{H-CH}_4$ between the northern high latitudes (60–90°N), relative to other global regions. We estimate a flux-weighted global freshwater $\delta^2\text{H-CH}_4$ of $-310 \pm 15\text{‰}$, which is higher than most previous estimates. Comparison with $\delta^{13}\text{C}$ measurements of both CH₄ and CO₂ implies that residual $\delta^2\text{H-CH}_4$ variation is the result of complex interactions between CH₄ oxidation, variation in the dominant pathway of methanogenesis, and potentially other biogeochemical variables. We observe a significantly greater distribution of $\delta^2\text{H-CH}_4$ values, corrected for $\delta^2\text{H-H}_2\text{O}$, in inland waters relative to wetlands, and suggest this difference is caused by more prevalent CH₄ oxidation in inland waters. We used the expanded freshwater CH₄ isotopic dataset to calculate a bottom-up estimate of global source $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ that includes spatially resolved isotopic signatures for freshwater CH₄ sources. Our bottom-up global source $\delta^2\text{H-CH}_4$ estimate ($-278 \pm 15\text{‰}$) is higher than a previous estimate using a similar approach, as a result of the more enriched global freshwater $\delta^2\text{H-CH}_4$ signature derived from our dataset. However, it is

Peter Douglas 5/11/2021 1:05 PM

Deleted: D

Peter Douglas 5/11/2021 1:05 PM

Deleted: -

Peter Douglas 5/11/2021 1:05 PM

Deleted: $\delta\text{D-H}_2\text{O}$

Peter Douglas 5/11/2021 1:05 PM

Formatted: Subscript

Peter Douglas 5/17/2021 1:50 PM

Deleted:

Peter Douglas 5/11/2021 9:31 AM

Deleted: to

Peter Douglas 5/11/2021 9:33 AM

Deleted: from

Peter Douglas 5/11/2021 1:06 PM

Deleted: ,

Peter Douglas 5/11/2021 1:06 PM

Deleted: and

Peter Douglas 5/17/2021 1:51 PM

Deleted:

Peter Douglas 5/13/2021 12:16 PM

Deleted: of the residual variability in $\delta^2\text{H-CH}_4$

Peter Douglas 5/11/2021 9:37 AM

Deleted: does not support a dominant role for either differential isotopic fractionation related to methanogenesis pathways or methane oxidation in controlling variation in $\delta^2\text{H-CH}_4$, but instead suggests

Peter Douglas 5/11/2021 9:37 AM

Deleted: these and other biogeochemical variables

Peter Douglas 5/13/2021 12:17 PM

Deleted: higher

Peter Douglas 5/11/2021 1:08 PM

Deleted: CH₄

Peter Douglas 5/11/2021 1:08 PM

Deleted: sources

Peter Douglas 5/11/2021 1:09 PM

Deleted: The

in agreement with top-down estimates of global source $\delta^2\text{H-CH}_4$ based on atmospheric measurements and estimated atmospheric sink fractionations. In contrast our bottom-up global source $\delta^{13}\text{C-CH}_4$ estimate is lower than top-down estimates, partly as a result of a lack of $\delta^{13}\text{C-CH}_4$ data from C_4 plant dominated ecosystems. In general, we find there is a particular need for more data to constrain isotopic signatures for low-latitude microbial CH_4 sources.

1 Introduction

Methane (CH_4) is an important greenhouse gas that accounts for approximately 25% of current anthropogenic global warming, but we do not have a complete understanding of the current relative or absolute fluxes of different CH_4 sources to the atmosphere (Schwietzke et al., 2016; Saunio et al., 2019), nor is there consensus on the causes of recent decadal-scale changes in the rate of increase in atmospheric CH_4 (Kai et al., 2011; Pison et al., 2013; Rice et al., 2016; Schaefer et al., 2016; Worden et al., 2017; Thompson et al., 2018; Turner et al., 2019). Freshwater ecosystems are an integral component of the global CH_4 budget. They are one of the largest sources of atmospheric CH_4 and are unequivocally the largest natural, or non-anthropogenic, source (Bastviken et al., 2011; Saunio et al., 2019). At the same time the geographic distribution of freshwater CH_4 emissions, changes in the strength of this source through time, and the relative importance of wetland versus inland water CH_4 emissions all remain highly uncertain (Pison et al., 2013; Schaefer et al., 2016; Ganesan et al., 2018; Saunio et al., 2019; Turner et al., 2019). Gaining a better understanding of freshwater CH_4 emissions on a global scale is of great importance for understanding potential future climate feedbacks related to CH_4 emissions from these ecosystems (Bastviken et al., 2011; Koven et al., 2011; Yvon-Durocher et al., 2014; Zhang et al., 2017). It is also necessary in order to better constrain the quantity and rate of change of other CH_4 emissions sources, including anthropogenic sources from fossil fuels, agriculture, and waste (Kai et al., 2011; Pison et al., 2013; Schaefer et al., 2016).

Isotopic tracers, particularly $\delta^{13}\text{C}$, have proven to be very useful in constraining global CH_4 sources and sinks (Kai et al., 2011; Nisbet et al., 2016; Rice et al., 2016; Schaefer et al., 2016; Schwietzke et al., 2016; Nisbet et al., 2019). However, $\delta^{13}\text{C}$ source signatures cannot fully differentiate CH_4 sources, leaving residual ambiguity in source apportionment (Schaefer et al., 2016; Schwietzke et al., 2016; Worden et al., 2017; Turner et al., 2019). Applying additional isotopic tracers to atmospheric CH_4 monitoring has the potential to greatly improve our understanding of CH_4 sources and sinks (Turner et al., 2019; Saunio et al., 2020). Recently developed laser-based methods, including cavity ringdown spectroscopy, quantum cascade laser absorption spectroscopy, and tunable infrared laser direct absorption spectroscopy, in addition to continued application of isotope ratio mass spectrometry (Chen et al., 2016; Röckmann et al., 2016; Yacovitch et al., 2020) could greatly enhance the practicality of atmospheric $\delta^2\text{H-CH}_4$ measurements at greater spatial and temporal resolution, similar to recent developments for $\delta^{13}\text{C-CH}_4$ measurements (Zazzeri et al., 2015; Miles et al., 2018). $\delta^2\text{H-CH}_4$ measurements have proven useful in understanding past CH_4 sources in ice-core records (Whiticar

Peter Douglas 5/13/2021 12:20 PM

Deleted: Saunio et al., 2019;

and Schaefer, 2007; Mischler et al., 2009; Bock et al., 2010; Bock et al., 2017), but have seen only limited use in modern atmospheric CH₄ budgets (Kai et al., 2011; Rice et al., 2016), in part because of loosely constrained source terms, as well as relatively sparse atmospheric measurements. Atmospheric inversion models have shown that increased spatial and temporal resolution of $\delta^2\text{H-CH}_4$ measurements could provide substantial improvements in precision for global and regional methane budgets (Rigby et al., 2012).

$\delta^2\text{H-CH}_4$ measurements could prove especially useful in understanding freshwater CH₄ emissions. Freshwater $\delta^2\text{H-CH}_4$ is thought to be highly dependent on $\delta^2\text{H-H}_2\text{O}$ (Waldron et al., 1999a; Whiticar, 1999; Chanton et al., 2006). Since $\delta^2\text{H-H}_2\text{O}$ exhibits large geographic variation as a function of temperature and fractional precipitation (Rozanski et al., 1993; Bowen and Revenaugh, 2003), $\delta^2\text{H-CH}_4$ measurements have the potential to differentiate freshwater CH₄ sources by latitude. This approach has been applied in some ice core studies (Whiticar and Schaefer, 2007; Bock et al., 2010), but geographic source signals remain poorly constrained, in part because of small datasets and because of incompletely understood relationships between $\delta^2\text{H-H}_2\text{O}$ and $\delta^2\text{H-CH}_4$. In contrast, recent studies of modern atmospheric $\delta^2\text{H-CH}_4$ have typically not accounted for geographic variation in freshwater CH₄ sources (Kai et al., 2011; Rice et al., 2016). Relatedly, other studies have found an important role for variation in $\delta^2\text{H-H}_2\text{O}$ in controlling $\delta^2\text{H-CH}_4$ from biomass burning (Umezawa et al., 2011) and from plants irradiated by UV light (Vigano et al., 2010), as well as the $\delta^2\text{H}$ of H₂ produced by wood combustion (Röckmann et al., 2016).

In addition to variance caused by $\delta^2\text{H-H}_2\text{O}$, a number of additional biogeochemical variables have been proposed to influence $\delta^2\text{H-CH}_4$ in freshwater environments. These include differences in the predominant biochemical pathway of methanogenesis (Whiticar et al., 1986; Whiticar, 1999; Chanton et al., 2006), the extent of methane oxidation (Happell et al., 1994; Waldron et al., 1999a; Whiticar, 1999; Cadieux et al., 2016), isotopic fractionation resulting from diffusive gas transport (Waldron et al., 1999a; Chanton, 2005), and differences in the thermodynamic favorability or enzymatic reversibility of methanogenesis (Valentine et al., 2004b; Stolper et al., 2015; Douglas et al., 2016). These influences on $\delta^2\text{H-CH}_4$ have the potential to complicate geographic signals, but also provide the potential to differentiate ecosystem sources, if specific ecosystems are characterized by differing rates and pathways of methanogenesis, rates of CH₄ oxidation, or gas transport processes. A recent study proposed that freshwater $\delta^{13}\text{C-CH}_4$ could be differentiated geographically based on ecosystem differences in the prevalence of different methanogenic pathways and in the predominance of C₄ plants, in addition to the geographic distribution of wetland ecosystems (Ganesan et al., 2018). $\delta^2\text{H-CH}_4$ measurements have the potential to complement this approach by providing an additional isotopic parameter for differentiating ecosystem and geographic CH₄ source signatures.

In order to use $\delta^2\text{H-CH}_4$ as an indicator of freshwater ecosystem contributions to global and regional CH₄ emissions budgets, a clearer understanding of freshwater $\delta^2\text{H}$ source signals, and how they vary by geographic location, ecosystem type, and other variables is needed. In order to address this need we have assembled and analyzed a dataset of 897 $\delta^2\text{H-CH}_4$ measurements from 129 individual ecosystems, or sites, derived from 40 publications (Schoell, 1983;

117 Woltemate et al., 1984; Burke Jr and Sackett, 1986; Whiticar et al., 1986; Burke Jr et al., 1988; Burke Jr, 1992; Burke Jr
 118 et al., 1992; Lansdown et al., 1992; Lansdown, 1992; Martens et al., 1992; Wassmann et al., 1992; Happell et al., 1993;
 119 Levin et al., 1993; Happell et al., 1994; Wahlen, 1994; Bergamaschi, 1997; Chanton et al., 1997; Hornibrook et al., 1997;
 120 Tyler et al., 1997; Zimov et al., 1997; Bellisario et al., 1999; Popp et al., 1999; Waldron et al., 1999b; Chasar et al., 2000;
 121 Marik et al., 2002; Nakagawa et al., 2002b; Nakagawa et al., 2002a; Chanton et al., 2006; Walter et al., 2006; Walter et
 122 al., 2008; Alstad and Whiticar, 2011; Brosius et al., 2012; Sakagami et al., 2012; Bouchard et al., 2015; Stolper et al.,
 123 2015; Wang et al., 2015; Cadieux et al., 2016; Douglas et al., 2016; Thompson et al., 2016; Lecher et al., 2017). We have
 124 advanced existing datasets of freshwater $\delta^2\text{H-CH}_4$ (Whiticar et al., 1986; Waldron et al., 1999a; Sherwood et al., 2017) in
 125 the following key attributes: 1) compiling a significantly larger dataset than was previously available; 2) compiling
 126 paired $\delta^{13}\text{C-CH}_4$ data for all sites, $\delta^{13}\text{C-CO}_2$ data for 50% of sites, and $\delta^2\text{H-H}_2\text{O}$ data for 47% of sites; 3) compiling
 127 geographic coordinates for all sites, providing the ability to perform spatial analyses and compare with gridded datasets
 128 of precipitation isotopic composition; and 4) classifying all sites by ecosystem and sample type (dissolved vs. gas
 129 samples), allowing for a clearer differentiation of how these variables influence $\delta^2\text{H-CH}_4$.

Peter Douglas 5/11/2021 9:42 AM
 Formatted: Subscript

130 Using this data set we applied statistical analyses to address key questions surrounding the global distribution of
 131 freshwater $\delta^2\text{H-CH}_4$, the variables that control this distribution, and its implications for atmospheric $\delta^2\text{H-CH}_4$.
 132 Specifically, we investigated the nature of the global dependence of $\delta^2\text{H-CH}_4$ on $\delta^2\text{H-H}_2\text{O}$, and whether this relationship
 133 results in significant differences in freshwater $\delta^2\text{H-CH}_4$ by latitude. We also assessed whether variability in $\delta^{13}\text{C-CH}_4$,
 134 $\delta^{13}\text{C-CO}_2$, and α_{C} was correlated with $\delta^2\text{H-CH}_4$, and whether there are significant differences in $\delta^2\text{H-CH}_4$ between
 135 different ecosystem and sample types. Finally, we used our dataset, combined with other isotopic datasets (Sherwood et
 136 al., 2017) and flux estimates (Saunois et al., 2020), to estimate the global $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ of global emissions
 137 sources, and compared this with previous estimates based on atmospheric measurements or isotopic datasets (Whiticar
 138 and Schaefer, 2007; Rice et al., 2016; Sherwood et al., 2017),.

Peter Douglas 5/17/2021 2:00 PM
 Deleted: ,

139 2 Methods

140 2.1 Isotope Nomenclature

141 Hydrogen and carbon isotope ratios are primarily discussed as delta values, using the generalized formula (Coplen,
 142 2011):

$$143 \delta = \frac{(R_{\text{sample}} - R_{\text{standard}})}{R_{\text{standard}}} \quad (1)$$

144 where R is the ratio of the heavy isotope to the light isotope, and the standard is Vienna Standard Mean Ocean Water
 145 (VSMOW) for $\delta^2\text{H}$ and Vienna Pee Dee Belemnite (VPDB) for $\delta^{13}\text{C}$. δ values are expressed in per mil (‰) notation.

Peter Douglas 5/11/2021 9:43 AM
 Deleted: The isotope notation used in this study is
 briefly introduced here.

We also refer to the isotopic fractionation factor between two phases, or α , which is defined as:

$$\alpha_{a-b} = \frac{R_a}{R_b} = \frac{\delta_a + 1}{\delta_b + 1} \quad (2)$$

Specifically, we discuss the carbon isotope fractionation factor between CO₂ and CH₄ (α_C) and the hydrogen isotope fractionation factor between H₂O and CH₄ (α_H).

2.2 Dataset Compilation

2.2.1 Literature Survey

To identify datasets we used a set of search terms (methane OR CH₄ AND freshwater OR wetland OR peatland OR swamp OR marsh OR lake OR pond OR 'inland water' AND 'hydrogen isotope' OR 'δD' OR 'δ²H') in Google Scholar to find published papers that discussed this measurement. We also identified original publications using previously compiled datasets (Waldron et al., 1999a; Sherwood et al., 2017). Data for 90% of sites were from peer-reviewed publications. Data from 13 sites were from a Ph.D. dissertation (Lansdown, 1992).

2.2.2 Dataset structure

Most samples were associated with geographic coordinates in data tables or text documentation, or with specific geographic locations such as the name of a town or city. In a few cases we identified approximate geographic locations based on text descriptions of sampling sites, with the aid of Google Earth software. Sampling sites were defined as individual water bodies or wetlands as identified in the relevant study. In [one study](#) where a number of small ponds were sampled from the same location, we grouped ponds of a given type as a single site (Bouchard et al., 2015). We divided sampling sites into six ecosystem categories: 1) lakes and ponds (hereafter lakes), 2) rivers and floodplains (hereafter rivers), 3) bogs, 4) fens, 5) swamps and marshes, and 6) rice paddies. Most data (7 of 8 sites) in the rivers category are from floodplain lake or delta environments. Swamps and marshes were combined as one category because of a small number of sites, and because there is no clear indication of biogeochemical differences between these ecosystems. To make these categorizations we relied on site descriptions in the data source publications. We also analyzed data in two larger environment types, inland waters (lakes and rivers) and wetlands (bogs, fens, swamps and marshes, and rice paddies), which correspond to two flux categories (freshwaters and natural wetlands) documented by Saunio et al. (2020). While rice paddies are an anthropogenic ecosystem, they are wetlands where microbial methanogenesis occurs under generally similar conditions to natural wetlands, and therefore we included them as wetlands in our analysis. In some cases the type of wetland was not specified. We did not differentiate between ombrotrophic and minerotrophic peatlands since most publications did not specify this difference, although it has been inferred to be important for δ¹³C-

Peter Douglas 5/11/2021 1:46 PM

Deleted: some cases

CH₄ distributions (Hornibrook, 2009). For studies of bogs and fens that sampled by soil depth we have only included sample measurements from the upper 50 cm. This is based on the observation of large-scale isotopic variability with soil depth in these ecosystems (Hornibrook et al., 1997; Waldron et al., 1999b), and the observation that shallow peat is typically the dominant source of atmospheric emissions (Waldron et al., 1999b; Bowes and Hornibrook, 2006; Shoemaker et al., 2012), which is our primary focus in this study. Other wetland ecosystems were not sampled by soil depth.

We also categorized samples by the form in which CH₄ was sampled, differentiating between dissolved CH₄ and CH₄ emitted through diffusive fluxes, which we group as dissolved CH₄, and gas-phase samples, including bubbles sampled either by disturbing sediments or by collecting natural ebullition fluxes. In some cases the sampling method or type of sample was not specified, or samples were a mix of both categories, which we did not attempt to differentiate.

Where possible (78% of sites), $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ values, as well as $\delta^{13}\text{C-CO}_2$ and $\delta^2\text{H-H}_2\text{O}$, were gathered from data files or published tables. In a number of publications, representing 22% of sites, data were only available graphically. For these studies we used Webplot Digitizer (<https://automeris.io/WebPlotDigitizer/>) software to extract data for these parameters. Previous studies have shown that user errors from Webplot Digitizer are typically small, with 90% of user generated data within 1% of the actual value (Drevon et al., 2017). Based on this, we estimate a typical error for $\delta^2\text{H-CH}_4$ data of less than 3‰. Studies where data were derived from graphs are identified in Supplementary Table S1 (Douglas et al., 2020).

2.2.3 Estimates of $\delta^2\text{H-H}_2\text{O}$ and its effects on $\delta^2\text{H-CH}_4$

To estimate $\delta^2\text{H-H}_2\text{O}$ for sites where it was not measured we relied on estimates of the isotopic composition of precipitation ($\delta^2\text{H}_p$), derived from the Online Isotopes in Precipitation Calculator v.3.1 (OIPC3.1; www.waterisotopes.org; Bowen and Wilkinson, 2002; Bowen and Revenaugh, 2003; Bowen et al., 2005). Inputs for $\delta^2\text{H}_p$ estimates are latitude, longitude, and elevation. We estimated elevation for each site surface elevation at the site's geographic coordinates reported by Google Earth. We tabulated estimates of both annual and growing season precipitation-amount weighted $\delta^2\text{H}_p$, where the growing season is defined as months with a mean temperature greater than 0 °C. We then analysed whether annual or growing season $\delta^2\text{H}_p$ is a better estimate of environmental $\delta^2\text{H-H}_2\text{O}$ for both wetlands and inland waters by comparing these values with measured $\delta^2\text{H-H}_2\text{O}$ for sites with measurements (See Sect. 3.2). Based on this analysis, we then identified a 'best-estimate' $\delta^2\text{H-H}_2\text{O}$ value for each site, using an approach similar to that of Waldron et al. (1999a). Namely, we apply measured $\delta^2\text{H-H}_2\text{O}$ where available, and estimates based on the regression analyses detailed in Section 3.2 for sites without measurements.

To account for the effects of $\delta^2\text{H-H}_2\text{O}$ on $\delta^2\text{H-CH}_4$, we introduce the term $\delta^2\text{H-CH}_{4,w0}$, which is the estimated $\delta^2\text{H-CH}_4$ of a sample if it had formed in an environment where $\delta^2\text{H-H}_2\text{O} = 0\text{‰}$. This is defined by the equation:

$$\delta^2\text{H-CH}_{4,w0} = \delta^2\text{H-CH}_4 - (b \times \delta^2\text{H}_2\text{O}) \quad (3)$$

Peter Douglas 5/11/2021 1:51 PM

Deleted: , and growing season precipitation-amount weighted $\delta^2\text{H}_p$.

where $\delta^2\text{H-H}_2\text{O}$ is the ‘best-estimate’ value for each site described above, b is the slope of the regression relationship of $\delta^2\text{H-H}_2\text{O}$ vs. $\delta^2\text{H-CH}_4$ for the entire dataset, as reported in Sect. 3.3. We also performed the same calculation separately for the subset of sites with measured $\delta^2\text{H-H}_2\text{O}$. We analyze $\delta^2\text{H-CH}_{4,\text{w0}}$ instead of α_{H} because, as discussed in Sect. 3.3.1, the global relationship between $\delta^2\text{H}_\text{p}$ vs. $\delta^2\text{H-CH}_4$ does not correspond to a constant value of α_{H} , and therefore deviations from the global empirical relationship are more clearly expressed as a residual as opposed to a fractionation factor.

2.3 Statistical analyses

For all statistical analyses we use site-level mean isotopic values. This avoids biasing our analyses towards sites with a large number of measurements, since there are large differences in the number of samples analyzed per site (n ranges from 66 to 1). To calculate α_{C} we used average $\delta^{13}\text{C-CH}_4$ and $\delta^{13}\text{C-CO}_2$ at a given site. This approach entails some additional uncertainty in this variable, but was necessary because at many sites these measurements were not made on the same samples.

We perform a set of linear regression analyses $\delta^2\text{H-CH}_4$ against other isotopic variables, in addition to latitude. All statistical analyses were performed in Matlab. We considered $p < 0.05$ to be the threshold for identifying significant regression relationships. We chose to perform unweighted regression, as opposed to weighted regression based on the standard error of sample measurements, for two reasons. First, a small number of sites with a large number of measurements, and therefore small standard error, had a disproportionate effect on weighted regression results. Second, in environmental research unweighted regression is frequently less biased than weighted regression (Fletcher and Dixon, 2012). Based on a test proposed by Fletcher and Dixon (2012), unweighted regression is appropriate for this dataset. We used analysis of covariance to test for significant differences between regression relationships.

To compare isotopic data ($\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$) between groups (i.e. latitudinal bands, ecosystem types, sample types) we used non-parametric statistical tests to test whether the groups were from different distributions. We used non-parametric tests because some sample groups were not normally distributed, as determined by a Shapiro-Wilk test (Shapiro and Wilk, 1965). For comparing differences between the distributions of two groups we used the Mann-Whitney U-test (Mann and Whitney, 1947), whereas when comparing differences between the distributions of more than two groups we used the Kruskal-Wallis H-test (Kruskal and Wallis, 1952), combined with Dunn’s test to compare specific sample group pairs (Dunn, 1964). We considered $p < 0.05$ to be the threshold for identifying groups with significantly different distributions.

When comparing $\delta^{13}\text{C-CH}_4$ by latitude and ecosystem we combined the data from this study with additional data from Sherwood et al. (2017) (32 additional sites) where $\delta^2\text{H-CH}_4$ was not measured to make our dataset as representative as possible. To our knowledge this combined dataset is the largest available compiled dataset of freshwater $\delta^{13}\text{C-CH}_4$, although there are many more $\delta^{13}\text{C-CH}_4$ measurements that have not yet been aggregated. We did not include these

Peter Douglas 5/11/2021 1:55 PM

Deleted: (p < 0.05)

244 additional data when analysing differences by sample type, as sample type was not specified in the dataset of Sherwood
245 et al. (2017).

246 2.4 Estimation of global atmospheric CH₄ δ²H and δ¹³C source values

247 To better understand how latitudinal differences in wetland isotopic source signatures influence atmospheric δ²H-CH₄
248 and δ¹³C-CH₄, we calculated a ‘bottom-up’ mixing model of δ²H-CH₄ and δ¹³C-CH₄. For this calculation we ascribed all
249 CH₄ sources a flux (derived from Saunio et al., 2020; see details below) and a δ²H and δ¹³C value, and calculated the
250 global atmospheric source value using an isotopic mixing model. Because of non-linearity when calculating mixtures
251 using δ²H values, we performed the mixing equation using isotopic ratios (see Sect. 2.1). The mixing equation is as
252 follows:

$$253 R_{mix} = f_1 R_1 + f_2 R_2 + \dots + f_n R_n \quad (4)$$

254 where f_n is the fractional flux for each source term (i.e. the ratio of the source flux to total flux), and R_n is the isotope
255 ratio for each source term.

256 Values for the flux, δ²H, and δ¹³C applied for each source term are shown in Table 1. We used bottom-up source
257 fluxes from Saunio et al. (2020) for the period 2008-2017. For categories other than wetlands, inland waters, and rice
258 paddies, we used global fluxes and isotope values, since geographically resolved isotopic source signature estimates are
259 not available. For these sources we used δ²H and δ¹³C values published by Sherwood et al. (2017), using the mean value
260 for each source term. For wetlands, inland waters, and rice paddies, we used geographically resolved (60-90 °N; 30-60
261 °N, 90° S-30°N) fluxes derived from Saunio et al. (2019) for the period 2008-2017, and mean δ²H-CH₄ for these
262 latitudinal bands from this study.

263 To calculate mean δ¹³C-CH₄ from wetlands, inland waters, and rice paddies for different latitudinal bands we
264 combined the data from this study along with additional data from Sherwood et al. (2017) (32 additional sites) to make
265 our estimated source signatures as representative as possible. To our knowledge this combined dataset is the largest
266 available compiled dataset of freshwater δ¹³C-CH₄ (See Sect. 2.3). Sites dominated by C₄ plants are notably
267 underrepresented in this combined dataset. In addition, the biomass burning dataset of Sherwood et al. (2017) contains
268 very few data from C₄ plant combustion. We performed a separate estimate of global source δ¹³C-CH₄ that attempted to
269 correct for these likely biases by making two adjustments: 1) using the estimated low-latitude wetland δ¹³C-CH₄
270 signature of Ganesan et al., (2018) (-56.7‰), which takes into account the predicted spatial distribution of C₄ plant
271 dominated wetlands; and 2) using the biomass burning δ¹³C-CH₄ signature of Schwietzke et al., (2016) (-22.3‰), which
272 is weighted by the predicted contribution from C₄ plant combustion. We did not attempt to take into account δ¹³C-CH₄
273 from ruminants feeding on C₄ plants. For the C₄-plant corrected δ¹³C-CH₄ estimate we applied the same uncertainties that
274 are reported in Table 1.

Peter Douglas 5/11/2021 1:58 PM

Deleted: se

Peter Douglas 5/11/2021 1:58 PM

Deleted:

Peter Douglas 5/11/2021 1:58 PM

Deleted: ions

Table 1: Estimates of source-specific fluxes, $\delta^2\text{H}$ -CH₄, and $\delta^{13}\text{C}$ -CH₄, and their uncertainties, used in mixing models and Monte Carlo analyses

| Category | Flux | $\delta^2\text{H}$ signature | | $\delta^{13}\text{C}$ signature | |
|--|---------|------------------------------|------------|---------------------------------|-----------|
| | (Tg/Yr) | Uncertainty | (‰, VSMOW) | Uncertainty | (‰, VPDB) |
| Wetlands (<30N) | 115 | 37.5 | -301 | 15 | -64.4 |
| Wetlands (30-60N) | 25 | 16.5 | -324 | 14 | -61.8 |
| Wetlands (>60N) | 9 | 8.0 | -374 | 10 | -62.7 |
| Inland Waters (<30N) | 80 | 39.4 | -301 | 12 | -57.1 |
| Inland Waters (30-60N) | 64 | 31.9 | -308 | 18 | -62.0 |
| Inland Waters (>60N) | 16 | 7.5 | -347 | 9 | -65.0 |
| Geological (onshore) ^a | 38 | 13.0 | -189 | 44 | -43.8 |
| Wild animals ^b | 2 | 2.0 | -316 | 28 | -65.4 |
| Termites ^c | 9 | 6.0 | -343 | 50 | -63.4 |
| Permafrost soils (direct) ^d | 1 | 0.5 | -374 | 15 | -64.4 |
| Geological (offshore) ^a | 7 | 7.0 | -189 | 44 | -43.8 |
| Biogenic open and coastal ^e | 6 | 3.0 | -200 | 50 | -80.0 |
| Enteric fermentation and manure | 111 | 5.0 | -308 | 28 | -65.4 |
| Landfills and waste | 65 | 4.5 | -297 | 6 | -56.0 |
| Rice cultivation (<30N) | 19 | 1.2 | -324 | 8 | -55.0 |
| Rice cultivation (30-60N) | 12 | 0.5 | -325 | 8 | -62.3 |
| Coal mining | 42 | 15.5 | -232 | 5 | -49.5 |
| Oil and gas | 79 | 13.0 | -189 | 2 | -43.8 |
| Industry ^f | 3 | 3.0 | -189 | 2 | -43.8 |
| Transport ^f | 4 | 4.0 | -189 | 2 | -43.8 |
| Biomass burning ^g | 17 | 6.0 | -211 | 15 | -26.2 |
| Biofuel burning ^g | 12 | 2.0 | -211 | 15 | -26.2 |

a-No specific isotopic measurements in the database (Sherwood et al., 2017). We applied the mean isotopic values for oil and gas, and applied the standard deviation of for oil and gas as the uncertainty

b-No specific isotopic measurements in database (Sherwood et al., 2017). We used the isotopic values and uncertainties from livestock

c- Only one $\delta^2\text{H}$ measurement in database (Sherwood et al., 2017). We applied 50‰ as a conservative uncertainty estimate.

d- No specific isotopic measurement in database (Sherwood et al., 2017). We used the isotopic values and uncertainties for high-latitude wetlands

e- No specific isotopic measurements in database (Sherwood et al., 2017). We applied approximate isotopic values based on Whiticar, (1999), and conservatively large uncertainty estimates.

f-No specific isotopic measurements in database (Sherwood et al., 2017). We used the isotopic values and uncertainties for oil and gas.

g-We applied all isotopic measurements of biomass burning to both the biomass burning and biofuel burning categories. We did not correct for the relative proportion of C₃ and C₄ plant combustion sources (See Sect. 2.4)

279 Since fluxes from *other natural sources* are not differentiated for the period 2008-2017, we calculated the
280 proportional contribution of each category of other natural sources for the period 2000-2009 (Saunois et al., 2020), and
281 applied this to the total flux from other natural sources for 2008-2017. Inland waters and rice paddies do not have
282 geographically resolved fluxes reported in Saunois et al. (2020). Therefore, we calculated the proportion of *other natural*
283 *sources* attributed to inland waters from 2000-2009 (71%), and applied this proportion to the geographically resolved
284 fluxes of *other natural sources*. Similarly, we calculated the proportion of *agricultural and waste sources* attributed to
285 rice agriculture from 2008-2017 (15%), and applied this to the geographically resolved fluxes of *agricultural and waste*
286 *fluxes*.

287 To estimate uncertainty in the modelled total source $\delta^2\text{H}$ and $\delta^{13}\text{C}$ values we conducted Monte Carlo analyses
288 (Thompson et al., 1992). We first estimated the uncertainty for each flux, $\delta^2\text{H}$, and $\delta^{13}\text{C}$ term. Flux uncertainties were
289 defined as one half of the range of estimates provided by Saunois et al., (2020). For sources where fluxes were calculated
290 as a proportion of a larger flux, we applied the same proportional calculation to uncertainty estimates. In cases where one
291 half of the range of reported studies was larger than the flux estimate, we set the uncertainty to be equal to the flux
292 estimate to avoid negative fluxes in the mixing model. Isotopic source signal uncertainties were defined as the 95%
293 confidence interval of the mean value for a given source category. For some sources there is insufficient data to calculate
294 a 95% confidence interval, and we applied a conservative estimate of uncertainty for these sources, as detailed in Table 1.
295 Confidence intervals for fossil fuel isotopic source signatures do not take into account variation in emissions fluxes and
296 isotopic values between regions or resource types (i.e. conventional vs. unconventional reservoirs). This variation is
297 difficult to quantify with available datasets, but could imply additional uncertainty in global source signatures. We
298 recalculated the $\delta^2\text{H}$ and $\delta^{13}\text{C}$ mixing models 10,000 times, each time sampling inputs from the uncertainty distribution
299 for each variable. We assumed all uncertainties were normally distributed. We interpret the 2-sigma standard deviation of
300 the resulting Monte Carlo distributions as an estimate of the uncertainty of our total atmospheric CH_4 source isotopic
301 values. To examine how the Monte Carlo analyses were specifically influenced by uncertainty in isotopic source
302 signatures vs. flux estimates, we conducted sensitivity tests where we set the uncertainty in either isotopic source
303 signatures or flux estimates to zero. We also used the mixing model and Monte Carlo method to estimate the mean flux-
304 weighted freshwater $\delta^2\text{H}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CH}_4$, using only the inputs for freshwater environments (Wetlands, Inland
305 Waters, and Rice Cultivation) from Table 1 (See Sect. 3.5)

306 **3 Results and Discussion**


307 **3.1 Dataset distribution**

308 The dataset is primarily concentrated in the northern hemisphere (Fig. 1A), but is distributed across a wide range
309 of latitudes between 3°S to 73°N (Fig. 1C). The majority of sampled sites are from North America (Fig. 1B), but there are

Peter Douglas 5/11/2021 10:00 AM
Deleted:
Peter Douglas 5/11/2021 10:03 AM
Deleted: then

Peter Douglas 5/11/2021 2:05 PM
Deleted:
Peter Douglas 5/11/2021 2:05 PM
Deleted:

314 numerous sites from Eurasia. A much smaller number of sites are from South America and Africa. We define three
315 latitudinal bands for describing geographic trends: low latitudes (3°S to 30°N); mid-latitudes (30° to 60°N); and high-
316 latitudes; (60° to 90°N). This definition was used primarily because it corresponds with a commonly applied geographic
317 classification of CH₄ fluxes (Saunois et al., 2020).



318 **Figure 1: Distribution of sites shown; A) on a global map, with site mean CH₄- $\delta^2\text{H}$ values indicated in relation to a color bar.**
319 **Sites with and without measured $\delta^2\text{H}-\text{H}_2\text{O}$ are differentiated; B) on a map of North America; and C) as a histogram of sites by**
320 **latitude, differentiated between wetlands and inland waters. Dashed lines in (C) indicate divisions between low-latitude, mid-**
321 **latitude, and high-latitude sites.**
322

323 74 of 129 sites are classified as inland waters, primarily lakes (n = 66), with a smaller number from rivers (n =
324 8). To our knowledge, all of the inland water sites are natural ecosystems and do not include reservoirs. 55 sites are

Peter Douglas 5/11/2021 2:05 PM
Deleted:
Peter Douglas 5/11/2021 2:05 PM
Deleted:
Peter Douglas 5/11/2021 2:05 PM
Deleted:
Peter Douglas 5/11/2021 2:05 PM
Deleted:
Peter Douglas 5/11/2021 2:05 PM
Deleted:

classified as wetlands, including 16 bogs, 14 swamps and marshes, 12 fens, and 8 rice paddies. For the majority of sites (n = 84) gas samples were measured, whereas studies at 36 sites measured dissolved CH₄ or diffusive fluxes.

3.2 Use of $\delta^2\text{H}_p$ as an estimator for freshwater $\delta^2\text{H-H}_2\text{O}$

As discussed in Sect. 2.2.3, we regressed annual and growing season $\delta^2\text{H}_p$ against measured $\delta^2\text{H-H}_2\text{O}$ to determine which is a better estimator for sites where $\delta^2\text{H-H}_2\text{O}$ is not measured. We performed this analysis separately for wetland and inland water environments because these broad environmental categories have distinct hydrological characteristics. For all regression analyses we found strong correlations, with R^2 values between 0.82 to 0.88 (Fig. 2). For wetlands, regression using annual $\delta^2\text{H}_p$ produces a slightly better fit, and also produces a slope within error of 1 (Fig 2A), suggesting that variation in annual $\delta^2\text{H}_p$ scales proportionately with variation in measured $\delta^2\text{H-H}_2\text{O}$. However, the intercept of this relationship was significantly greater than 0 (19 ± 9 ‰). We interpret this intercept as indicating that evaporative isotopic enrichment is generally important in controlling $\delta^2\text{H-H}_2\text{O}$ in wetlands. A slope slightly greater than 1 is also consistent with evaporative enrichment, since greater evaporation rates would be expected in low-latitude environments with higher $\delta^2\text{H-H}_2\text{O}$. These results are consistent with detailed studies of wetland isotope hydrology that indicate a major contribution from groundwater, with highly dampened seasonal variability relative to precipitation, but also indicate evaporative enrichment of water isotopes in shallow soil water (Sprenger et al., 2017; David et al., 2018).

For inland waters, regression with growing season $\delta^2\text{H}_p$ produces a relationship that is within error of the 1:1 line (Fig. 2C), in contrast to annual $\delta^2\text{H}_p$, which produces a flatter slope (Fig. 2D). We infer that seasonal differences in $\delta^2\text{H}_p$ are important in determining $\delta^2\text{H-H}_2\text{O}$ in the inland water environments analyzed, especially at high latitudes, implying that these environments generally have water residence times on subannual timescales. This finding is generally consistent with evidence for seasonal variation in lake water isotopic compositions that is dependent on lake water residence times (Tyler et al., 2007; Jonsson et al., 2009). Lake water residence times vary widely, primarily as a function of lake size, but isotopic data implies that small lakes have water residence times of less than a year (Brooks et al., 2014), resulting in seasonal isotopic variability (Jonsson et al., 2009). Isotopic enrichment of lake water is highly variable, but is typically minor in humid and high-latitude regions (Jonsson et al., 2009; Brooks et al., 2014), which characterizes most of our study sites.

Based on these results we combine measured and estimated $\delta^2\text{H-H}_2\text{O}$ to determine a ‘best-estimate’ value for each site, an approach similar to that of Waldron et al. (1999a). For sites with measured $\delta^2\text{H-H}_2\text{O}$ values we use that value. For inland water sites without measured $\delta^2\text{H-H}_2\text{O}$ we use modeled growing season $\delta^2\text{H}_p$ since the regression of this against measured $\delta^2\text{H-H}_2\text{O}$ is indistinguishable from the 1:1 line (Fig. 2D). For wetland sites without measured $\delta^2\text{H-H}_2\text{O}$ we estimate $\delta^2\text{H-H}_2\text{O}$ using the regression relationship with annual precipitation $\delta^2\text{H-H}_2\text{O}$ shown in Fig. 2A. The root mean square errors (RMSE) of these relationships (16‰ for wetlands, 22‰ for inland waters) provide an estimate of

Peter Douglas 5/17/2021 3:18 PM

Deleted: compared modelled

Peter Douglas 5/13/2021 12:32 PM

Deleted: with

Peter Douglas 5/13/2021 12:31 PM

Deleted: comparisons

Peter Douglas 5/11/2021 10:53 AM

Deleted: }

Peter Douglas 5/11/2021 2:08 PM

Deleted: the measured

the uncertainty associated with estimating $\delta^2\text{H-H}_2\text{O}$ using $\delta^2\text{H}_p$. Given the uncertainty associated with estimating $\delta^2\text{H-H}_2\text{O}$ using $\delta^2\text{H}_p$, for all analyses presented below that depend on $\delta^2\text{H-H}_2\text{O}$ values we also analyse the dataset only including sites with measured $\delta^2\text{H-H}_2\text{O}$.




Figure 2: Scatter plots of annual or growing season $\delta^2\text{H}_p$ vs. measured $\delta^2\text{H-H}_2\text{O}$ for wetland (A,B) and inland water (C,D) sites. The red lines indicates the best fit, with a 95% confidence interval (gray envelopes), and the dashed black lines are the 1:1 relationship.

376 We carried out regression analyses of $\delta^2\text{H-H}_2\text{O}$ vs. $\delta^2\text{H-CH}_4$, both using ‘best-estimate’ $\delta^2\text{H-H}_2\text{O}$ as described in sect. 3.2
 377 (Fig. 3A), and only including sites with measured $\delta^2\text{H-H}_2\text{O}$ (Fig. 3B). In addition we analysed the relationship for all
 378 sites using annual (Fig. 3C) and growing season (Fig. 3D) $\delta^2\text{H}_p$. Identifying the relationship between modelled $\delta^2\text{H}_p$ and
 379 $\delta^2\text{H-CH}_4$ is of value because this could be used to create gridded global predictions of $\delta^2\text{H-CH}_4$ based on gridded datasets
 380 of $\delta^2\text{H}_p$ (Bowen and Revenaugh, 2003), as well as to predict the distribution of $\delta^2\text{H-CH}_4$ under past and future global
 381 climates using isotope enabled Earth system models (Zhu et al., 2017).

382 $\delta^2\text{H-CH}_4$ is significantly positively correlated with $\delta^2\text{H-H}_2\text{O}$ when using all four methods of estimating $\delta^2\text{H-}$
 383 H_2O (Fig. 3, Supplemental Table 2). This is the case when analysing all sites together, as well as when analysing
 384 wetlands and inland waters separately (Supplemental Table 2, Fig. 4). There is no significant difference in regression
 385 relationships, based on analysis of covariance, when $\delta^2\text{H-CH}_4$ is regressed against best-estimate $\delta^2\text{H-H}_2\text{O}$, measured $\delta^2\text{H-}$
 386 H_2O , or modelled $\delta^2\text{H}_p$, nor is there a major difference in R^2 values or RMSE (Supplemental Table S2). Regression with
 387 wetland sites consistently results in a higher R^2 values and lower RMSE than regression with inland water sites.

388 Given the similar results when regressing with estimated or measured $\delta^2\text{H-H}_2\text{O}$, we infer that using either the
 389 ‘best-estimate’ $\delta^2\text{H-H}_2\text{O}$ or modelled $\delta^2\text{H}_p$, instead of measured $\delta^2\text{H-H}_2\text{O}$, to predict $\delta^2\text{H-CH}_4$ does not result in
 390 substantial additional error. This implies that isotope-enabled Earth Systems models (ESMs) could be used to predict the
 391 distribution of freshwater $\delta^2\text{H-CH}_4$ under past and future climates based on modeled $\delta^2\text{H}_p$, although the substantial
 392 scatter in Figures 3C and D should be taken into account. The southern hemisphere is highly underrepresented in the
 393 $\delta^2\text{H-CH}_4$ dataset. However, the mechanisms linking $\delta^2\text{H-CH}_4$ with $\delta^2\text{H-H}_2\text{O}$ should not differ in the southern
 394 hemisphere, and we argue that the relationships observed in this study are suitable to predict southern hemisphere
 395 freshwater $\delta^2\text{H-CH}_4$. The choice of predicting $\delta^2\text{H-CH}_4$ using growing-season vs. annual precipitation $\delta^2\text{H}_p$ could be
 396 important, with steeper slopes overall when regressing against growing season $\delta^2\text{H}_p$. Based on our analysis in sect. 3.2,
 397 we suggest that annual $\delta^2\text{H}_p$ may be more appropriate for estimating wetland $\delta^2\text{H-CH}_4$, while growing season $\delta^2\text{H}_p$ may
 398 be more appropriate for estimating inland water $\delta^2\text{H-CH}_4$. Forthcoming research will combine gridded datasets of
 399 wetland distribution (Ganesan et al., 2018), modeled annual $\delta^2\text{H}_p$ (Bowen and Revenaugh, 2003), and the regression
 400 relationships from this study to predict spatially-resolved wetland $\delta^2\text{H-CH}_4$ at a global scale (Stell et al., in press).

Peter Douglas 5/13/2021 12:34 PM

Deleted: -

... (1)

Peter Douglas 5/11/2021 2:10 PM

Deleted: Wetland sites consistently have a steeper regression slope than inland water sites (Supplemental Table S2), but this difference is not significant.

Peter Douglas 5/11/2021 2:10 PM

Deleted: also

Peter Douglas 5/13/2021 12:35 PM

Deleted: available

Peter Douglas 5/17/2021 3:25 PM

Deleted: H₂O-

Peter Douglas 5/11/2021 10:56 AM

Deleted: Future




Figure 3: Scatter plots of $\delta^2\text{H-CH}_4$ vs. (A) best-estimate $\delta^2\text{H-H}_2\text{O}$; (B) measured $\delta^2\text{H-H}_2\text{O}$; (C) annual $\delta^2\text{H}_p$; and (D) growing season $\delta^2\text{H}_p$. Sites that were included in the analysis of Waldron et al., (1999a) are indicated. The regression relationship for the total dataset in each plot is shown by the red line, with its 95% confidence interval (grey envelope). The regression relationship and confidence interval for the dataset of Waldron et al., (1999a) is shown in blue. Uncertainties for reported regression relationships are standard errors.

Overall, our results are broadly consistent with those of Waldron et al., (1999a), and confirm the finding of that study that $\delta^2\text{H-H}_2\text{O}$ is the predominant predictor of global variation in $\delta^2\text{H-CH}_4$. All of the regression slopes produced using our dataset are flatter than the regression relationship found by Waldron et al. (1999a) using a smaller dataset (0.68 ± 0.1), although the slopes are not significantly different based on analysis of covariance. Based on this result we

infer that the true global relationship is likely flatter than that estimated by Waldron et al. (1999a). The difference between the regression relationships reported here and that of Waldron et al. (1999a) is largely a result of a much greater number of samples from the high latitudes (Fig. 1C), where $\delta^2\text{H-H}_2\text{O}$ values are typically lower. The small number of high-latitude sites sampled by Waldron et al. (1999a) are skewed towards the low end of the high-latitude $\delta^2\text{H-CH}_4$ data from this study (Fig. 3). A similarly flatter slope (0.54 ± 0.05) was found by Chanton et al. (2006) when combining a dataset of $\delta^2\text{H-CH}_4$ from Alaskan wetlands, which are included in this study, with the dataset of Waldron et al. (1999a). Based on the range of R^2 values shown in Figure 3, we estimate that $\delta^2\text{H-H}_2\text{O}$ explains approximately 42% of variability in $\delta^2\text{H-CH}_4$, implying substantial residual variability, with greater residual variability for inland water sites than for wetlands (Supplemental Table 2).

Given that $\delta^2\text{H-H}_2\text{O}$ is strongly influenced by latitude, we examined whether $\delta^2\text{H-CH}_4$ is also significantly correlated with latitude. There is indeed a significant, negative relationship between latitude and $\delta^2\text{H-CH}_4$, indicating an approximate decrease of 0.9‰ latitude (Fig. 4). The slope is significantly flatter than that for latitude vs. $\delta^2\text{H-H}_2\text{O}$ in this dataset (-2‰ latitude), which is consistent with the inferred slope for $\delta^2\text{H-H}_2\text{O}$ vs. $\delta^2\text{H-CH}_4$ (0.44 to 0.5). There is greater scatter in $\delta^2\text{H-CH}_4$ at higher latitudes, especially for inland waters, but it is unclear if this is simply a result of a larger sample set or of differences in the underlying processes controlling $\delta^2\text{H-CH}_4$. We discuss latitudinal differences in $\delta^2\text{H-CH}_4$ in further detail in Sect. 3.5




Figure 4: Scatter plots of $\delta^2\text{H-CH}_4$ and best-estimate $\delta^2\text{H-H}_2\text{O}$, vs. latitude ($^{\circ}\text{N}$). Sites with measured $\delta^2\text{H-H}_2\text{O}$ are indicated. Envelopes indicate 95% confidence intervals for regression lines.

Peter Douglas 5/11/2021 10:58 AM

Deleted: , but more data will be needed to further constrain this relationship

Peter Douglas 5/13/2021 12:37 PM

Deleted: As discussed below in sect. 3.3.1, our regression relationship slopes are very similar to that of the 'in-vitro' line of Waldron et al. (1999a).

Peter Douglas 5/17/2021 3:26 PM

Deleted: in

Peter Douglas 5/11/2021 10:59 AM

Deleted: , although it is also influenced by other geographic and climatic variables,

Peter Douglas 5/13/2021 12:42 PM

Deleted:

Peter Douglas 5/11/2021 2:14 PM

Deleted:

3.3.1 Comparison of $\delta^2\text{H-H}_2\text{O}$ vs $\delta^2\text{H-CH}_4$ relationships between environmental and experimental studies

To further understand the processes controlling the observed [freshwater](#) $\delta^2\text{H-H}_2\text{O}$ vs. $\delta^2\text{H-CH}_4$ relationships we compared them to results from pure culture and incubation experiments across a wide range of $\delta^2\text{H-H}_2\text{O}$ values (Fig. 5), focusing on regression against best-estimate $\delta^2\text{H-H}_2\text{O}$. The regression slopes for both wetlands and inland waters (0.5 and 0.42) are within error of the ‘in-vitro’ relationship compiled by Waldron et al. (1999a) (0.44), based on laboratory incubations from three separate studies (Schoell, 1980; Sugimoto and Wada, 1995; Waldron et al., 1998). The intercept for the wetland and inland water regressions is higher than that for the ‘in-vitro’ relationship, although only the difference with inland waters is significant. In contrast, the regression slope for pure-culture acetoclastic methanogenesis experiments is much flatter (0.18 to 0.2) (Valentine et al., 2004b; Gruen et al., 2018), consistent with the prediction that one hydrogen atom is exchanged between water and the acetate methyl group during CH_4 formation (Pine and Barker, 1956; Whiticar, 1999). The large difference in intercept between the two acetate pure culture datasets is likely a function of differences in the $\delta^2\text{H}$ of acetate [between the experiments](#), but could also be influenced by differences in kinetic isotope effects (Valentine et al., 2004b).

Pure culture hydrogenotrophic methanogenesis experiments (Gruen et al., 2018) yield a regression slope that is consistent with a constant α_{H} value, although α_{H} [can](#) clearly [vary](#) depending on experimental or environmental conditions (Valentine et al., 2004b; Stolper et al., 2015; Douglas et al., 2016). The wetland, inland water, and ‘in-vitro’ regression relationships are not consistent with a constant value of α_{H} (Fig. 5). Our comparison supports previous inferences that the in-vitro line of Waldron et al., (1999a) provides a good estimate of the slope of environmental $\delta^2\text{H-H}_2\text{O}$ vs. $\delta^2\text{H-CH}_4$ relationships. This slope is likely controlled by the relative proportion of acetoclastic and hydrogenotrophic methanogenesis, the net kinetic isotope effect associated with these two methanogenic pathways, and variance in $\delta^2\text{H}$ of acetate (Waldron et al., 1998; Waldron et al., 1999a; Valentine et al., 2004a), but the relative importance of these variables remains uncertain.

In particular, the $\delta^2\text{H}$ of acetate methyl hydrogen is probably influenced by environmental $\delta^2\text{H-H}_2\text{O}$, and therefore likely varies geographically as a function of $\delta^2\text{H}_\text{p}$, as originally hypothesized by Waldron et al. (1999a). To our knowledge there are no measurements of acetate or acetate-methyl $\delta^2\text{H}$ from natural environments with which to test this hypothesis. In general, variability in the $\delta^2\text{H}$ of environmental organic molecules in lake sediments and wetlands, including fatty acids and cellulose, is largely controlled by $\delta^2\text{H-H}_2\text{O}$ (Huang et al., 2002; Sachse et al., 2012; Mora and Zanazzi, 2017), albeit with widely varying fractionation factors. The $\delta^2\text{H}$ of methoxyl groups in plants has also been shown to vary as a function of $\delta^2\text{H-H}_2\text{O}$ (Vigano et al., 2010). Furthermore, culture experiments with acetogenic bacteria imply that there is rapid isotopic exchange between H_2 and H_2O during chemoautotrophic acetogenesis (Valentine et al., 2004a), implying that the $\delta^2\text{H}$ of [chemoautotrophic acetate](#) is also partially controlled by environmental $\delta^2\text{H-H}_2\text{O}$. Incubation experiments, such as those included in the ‘in-vitro line’ (Schoell, 1980; Sugimoto and Wada, 1995; Waldron

Peter Douglas 5/11/2021 11:44 AM

Deleted: varies

Peter Douglas 5/11/2021 2:15 PM

Deleted: }

Peter Douglas 5/11/2021 2:15 PM

Deleted:

et al., 1998), probably contain acetate- $\delta^2\text{H}$ that varies as a function of ambient $\delta^2\text{H-H}_2\text{O}$, given that the acetate in these incubation experiments was actively produced by fermentation and/or acetogenesis during the course of the experiment. This differs from pure cultures of methanogens, where acetate is provided in the culture medium and therefore does not vary in its $\delta^2\text{H}$ value (Valentine et al., 2004b; Gruen et al., 2018).

Peter Douglas 5/11/2021 11:45 AM
Deleted: would




Figure 5: Scatter plots of $\delta^2\text{H-CH}_4$ vs. $\delta^2\text{H-H}_2\text{O}$ for wetlands, inland waters, landfills, and cow rumen, compared with incubation and pure-culture experiments. Regression lines and confidence intervals corresponding to each dataset (except landfills and cow rumen) are shown. Dashed gray lines indicate constant values of α_{H} . Regression line statistics are listed in Supplemental Table S2. Plotted $\delta^2\text{H-H}_2\text{O}$ values are ‘best-estimate’ values for wetlands and inland waters, measured values for culture experiments, and a combination of measured values and annual $\delta^2\text{H}_\text{p}$ for landfills and cow rumen (See supplemental Table S3 for more details).

3.4 Relationship of $\delta^2\text{H-CH}_4$ with $\delta^{13}\text{C-CH}_4$, $\delta^{13}\text{C-CO}_2$, and α_{C}

As shown in Fig. 3, there is a large amount of residual variability in $\delta^2\text{H-CH}_4$ that is not explained by $\delta^2\text{H-H}_2\text{O}$. Several biogeochemical variables have been proposed to influence freshwater $\delta^2\text{H-CH}_4$ independently of $\delta^2\text{H-H}_2\text{O}$, including the predominant biochemical pathway of methanogenesis (Whiticar et al., 1986;Whiticar, 1999;Chanton et al., 2006), the extent of methane oxidation (Happell et al., 1994;Waldron et al., 1999a;Whiticar, 1999;Cadieux et al., 2016), isotopic fractionation resulting from diffusive gas transport (Waldron et al., 1999a;Chanton, 2005), and differences in the thermodynamic favorability or reversibility of methanogenesis (Valentine et al., 2004b;Stolper et al., 2015;Douglas et al.,

2016). These variables are also predicted to cause differences in $\delta^{13}\text{C-CH}_4$, $\delta^{13}\text{C-CO}_2$, and α_c . Therefore, we analysed co-variation between $\delta^2\text{H-CH}_{4,\text{wO}}$ (see definition in Sect. 2.2.3) and $\delta^{13}\text{C-CH}_4$, $\delta^{13}\text{C-CO}_2$, and α_c to see if it could partially explain the residual variability in $\delta^2\text{H-CH}_4$ (Fig. 6).

In order to facilitate interpretation of isotopic co-variation, we estimated approximate vectors of predicted isotopic co-variation for the four variables being considered (Fig. 6). We emphasize that these vectors are uncertain, and while they can be considered indicators for the sign of the slope of co-variation and the relative magnitude of expected isotopic variability, they are not precise representations of the slope or intercept of isotopic co-variation. In reality, isotopic co-variance associated with these processes likely varies depending on specific environmental conditions, although the sign of co-variance should be consistent. The starting point for the vectors is arbitrarily set to typical isotopic values for inferred acetoclastic methanogenesis in freshwater systems (Whiticar, 1999). We based the vectors for differences in the dominant methanogenic pathway and methane oxidation on Figures 8, 5, and 10 in Whiticar (1999). These figures are widely applied to interpret environmental isotopic data related to CH_4 cycling. However, we note that both environmental and experimental research has questioned whether differences in the dominant methanogenic pathway has an influence on $\delta^2\text{H-CH}_4$ (Waldron et al., 1998; Waldron et al., 1999a). Differences in $\delta^2\text{H-CH}_4$ between hydrogenotrophic and acetoclastic methanogenesis are likely highly dependent on both the $\delta^2\text{H}$ of acetate and the carbon and hydrogen kinetic isotope effects for both methanogenic pathways, both of which are poorly constrained in natural environments and are likely to vary between sites (see Sect. 3.3.1). We did not differentiate between anaerobic and aerobic methane oxidation, and the vectors shown are similar to experimental results for aerobic methane oxidation (Wang et al., 2016).

The vector for isotopic fractionation related to gas-phase diffusion is based on the calculations of Chanton (2005), and indicates isotopic change for residual gas following a diffusive loss. Gas-liquid diffusion is predicted to have a much smaller isotopic effect (Chanton, 2005). The vector for differences in enzymatic reversibility are based on experiments where CH_4 and CO_2 isotopic compositions were measured together with changes in methane production rate or Gibbs free energy (Valentine et al., 2004b; Penning et al., 2005). We note that these studies did not measure $\delta^2\text{H-CH}_4$ in the same experiments as $\delta^{13}\text{C-CH}_4$ or $\delta^{13}\text{C-CO}_2$, implying large uncertainty in the co-variance vectors. More detail on the estimated vectors is provided in the Supplementary Text.

Peter Douglas 5/13/2021 12:44 PM

Deleted: -

Peter Douglas 5/13/2021 12:44 PM

Deleted: -

Peter Douglas 5/13/2021 12:44 PM

Formatted: Indent: First line: 0.5"




Figure 6: Scatter plots of $\delta^2\text{H}-\text{CH}_{4,\text{w0}}$ vs. (A) $\delta^{13}\text{C}-\text{CH}_4$, (B) α_c , and (C) $\delta^{13}\text{C}-\text{CO}_2$. Approximate vectors for isotopic covariation related to four biogeochemical variables are shown. See details in Sect. 3.4 and the supplemental text. Regression relationships are shown for wetland and inland water sites, with envelopes indicating 95% confidence intervals. Regression statistics are shown here for relationships with significant correlations ($p < 0.05$). All regression statistics are detailed in Supplemental Table S4.

539 We observe significant positive correlations between $\delta^2\text{H-CH}_{4,\text{w0}}$, calculated using best estimate $\delta^2\text{H-H}_2\text{O}$, and
 540 both $\delta^{13}\text{C-CO}_2$, and α_c for wetland sites (Fig. 6B,C; Supplemental Table S4). We do not observe a significant correlation
 541 between these variables for inland water sites or for the dataset as a whole. We also observe a weak but significant
 542 negative correlation between $\delta^2\text{H-CH}_{4,\text{w0}}$ and $\delta^{13}\text{C-CH}_4$ for all sites, but not for data disaggregated into wetlands and
 543 inland water categories (Fig. 6A). The correlations shown in Figure 6 should be interpreted with caution, since repeating
 544 this analysis only using sites with measured $\delta^2\text{H-H}_2\text{O}$ does not result in any significant correlations (Supplemental Table
 545 S4). It is unclear whether this different result when using best-estimate or measured $\delta^2\text{H-H}_2\text{O}$ represents a bias related to
 546 estimating $\delta^2\text{H-H}_2\text{O}$ using $\delta^2\text{H}_p$, or is an effect of the much smaller sample size for sites with $\delta^2\text{H-H}_2\text{O}$ measurements. If
 547 accurate, the observed significant positive correlations in Figures 6B and C suggest that residual variability in $\delta^2\text{H-CH}_4$
 548 in wetlands is more strongly controlled by biogeochemical variables related to methanogenesis, namely differences in
 549 methanogenic pathway or thermodynamic favorability, than post-production processes such as diffusive transport and
 550 CH_4 oxidation. For inland water sites our analysis suggests that no single biogeochemical variable has clear effect in
 551 controlling residual variability in $\delta^2\text{H-CH}_4$.

552 Overall, our results are not consistent with arguments that residual variability in freshwater $\delta^2\text{H-CH}_4$ is
 553 dominantly controlled by either differences in methanogenic pathway (Chanton et al., 2006), or post-production
 554 processes (Waldron et al., 1999a). Instead they highlight the combined influence of a complex set of variables and
 555 processes that are difficult to disentangle on an inter-site basis using $\delta^{13}\text{C}$ measurements alone. It is also important to
 556 note the likely importance of variables that could influence $\delta^{13}\text{C-CH}_4$ or $\delta^{13}\text{C-CO}_2$ but not necessarily affect $\delta^2\text{H-CH}_4$,
 557 including variance in the $\delta^{13}\text{C}$ of soil or sediment organic matter (Conrad et al., 2011; Ganesan et al., 2018), diverse
 558 metabolic and environmental sources and sinks of CO_2 in aquatic environments, and Rayleigh fractionation associated
 559 with CH_4 carbon substrate depletion (Whiticar, 1999). Finally, the possible role of other carbon substrates, such as
 560 methanol, in CH_4 production could be important in controlling isotopic co-variation. Culture experiments suggest that
 561 CH_4 produced from methanol has low $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values relative to other pathways (Krzycki et al., 1987; Penger et al.,
 562 2012; Gruen et al., 2018), although the importance of this difference in environmental CH_4 is unclear.

563 Further research examining intra-site isotopic co-variation, which largely avoids complications associated with
 564 estimating $\delta^2\text{H-H}_2\text{O}$, would help to more clearly resolve the relative importance of these processes, and how they vary
 565 between environments. Expanded research using methyl fluoride to inhibit acetoclastic methanogenesis (Penning et al.,
 566 2005; Penning and Conrad, 2007; Conrad et al., 2011), with a particular focus on $\delta^2\text{H-CH}_4$ measurements, would also help
 567 to clarify the importance of methanogenic pathway on isotopic co-variation. Finally, an expanded application of
 568 measurements of clumped isotopes, which have distinctive patterns of variation related to these processes (Douglas et al.,

Peter Douglas 5/13/2021 1:32 PM
 Deleted: very
 Peter Douglas 5/13/2021 1:32 PM
 Deleted: ,
 Peter Douglas 5/13/2021 1:32 PM
 Deleted: ,
 Peter Douglas 5/13/2021 1:32 PM
 Deleted: ,
 Peter Douglas 5/11/2021 11:51 AM
 Deleted: significant

Peter Douglas 5/11/2021 11:51 AM
 Deleted: However, the residual variability in $\delta^2\text{H-CH}_4$ explained by $\delta^{13}\text{C-CO}_2$ and α_c in wetlands is relatively small, specifically between 19 to 28% based on the R^2 values in Figures 6B and C.
 Peter Douglas 5/11/2021 10:33 AM
 Deleted: It is intriguing that we observe the strongest correlation in wetlands between $\delta^2\text{H-CH}_{4,\text{w0}}$ and $\delta^{13}\text{C-CO}_2$, since it is probable that a wide range of biotic and abiotic processes unrelated to methane cycling influence $\delta^{13}\text{C-CO}_2$. This suggests that measurements of $\delta^{13}\text{C-CO}_2$ are important for future research on environmental variables controlling wetland $\delta^2\text{H-CH}_4$.

Peter Douglas 5/17/2021 3:32 PM
 Deleted: e
 Peter Douglas 5/17/2021 3:32 PM
 Deleted: variability

2016; Douglas et al., 2017; Young et al., 2017; Douglas et al., 2020), would also be of value in determining their relative importance in controlling $\delta^2\text{H-CH}_4$ values in freshwater environments.

3.5 Differences in $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ by latitude

When analysing all sites together we found a significant difference in the distribution of $\delta^2\text{H-CH}_4$ between high-latitude sites (median: -351‰) and both low (median: -298‰) and mid-latitude sites (median: -320‰) (Fig. 7A). However, we did not find a significant difference in the distribution of low- and mid-latitude sites. Similar differences were found when the data were disaggregated into wetland and inland water sites. We also found that the distribution of $\delta^{13}\text{C-CH}_4$ for low latitude sites (median: -61.6‰) was significantly higher than for high latitude sites (median: -63.0‰), but that mid-latitude sites (median: -60.3‰) were not significantly different from the other two latitudinal zones (Fig. 7B). The observed difference by latitudinal zone in $\delta^{13}\text{C-CH}_4$ appears to be driven primarily by latitudinal differences between inland water sites, where a similar pattern is found. In wetland sites we found no significant differences in the distribution of $\delta^{13}\text{C-CH}_4$ by latitude.




Figure 7: Boxplots of (A) $\delta^2\text{H-CH}_4$ and (B) $\delta^{13}\text{C-CH}_4$ for sites differentiated by latitude, for all data, wetlands, and inland waters. Numbers in parentheses indicate the number of sites for each category. Red lines indicate medians, boxes indicate 25th and 75th percentiles, whiskers indicate 95th and 5th percentiles, and outliers are shown as red crosses. Notches indicate the 95% confidence intervals of the median value; where notches overlap the edges of the box this indicates the median confidence interval exceeds the 75th or 25th percentile. Black points and error bars indicate the category mean and 95% confidence interval of the mean. Gray lines indicate the estimated flux-weighted mean values for global freshwater CH_4 , and dashed lines indicate the 95% confidence interval of this value. Asterisks in (A) indicate that high-latitude sites have significantly different distributions from other latitudinal bands. Asterisks in (B) indicate groups that have significantly different distributions from one another, within a specific environmental category. Two extremely low outliers ($<-80\text{‰}$; high latitude wetland and inland water) are not shown in (B).

Estimates of flux-weighted mean freshwater $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$, calculated using the Monte Carlo approach described in Sect. 2.4, are $-310\pm 15\text{‰}$ (Fig. 7A) and $-61.5\pm 3\text{‰}$ (Fig. 7B) respectively. Flux weighted mean values for natural wetlands (not including inland waters or rice paddies) are $-310\pm 25\text{‰}$ for $\delta^2\text{H-CH}_4$ and $-63.9\pm 3.3\text{‰}$ for $\delta^{13}\text{C-CH}_4$.

615 Flux weighted mean values for inland waters are $-309 \pm 31\text{‰}$ for $\delta^2\text{H-CH}_4$ and $-60 \pm 5.7\text{‰}$ for $\delta^{13}\text{C-CH}_4$. As discussed in
616 Sect. 2.4 there are limited data in our dataset or that of Sherwood et al., (2017) from C_4 plant dominated wetlands, and
617 therefore our low-latitude and flux-weighted mean $\delta^{13}\text{C-CH}_4$ values for wetlands are probably biased towards low values.

618 Differences in $\delta^2\text{H-CH}_4$ by latitude has the potential to aid in geographic discrimination of freshwater methane
619 sources, both because it is based on a clear mechanistic linkage with $\delta^2\text{H-H}_2\text{O}$ (Figs. 3 and 4), and because geographic
620 variation in $\delta^2\text{H-H}_2\text{O}$ is relatively well understood (Bowen and Revenaugh, 2003; Bowen et al., 2005). However, recent
621 studies of atmospheric $\delta^2\text{H-CH}_4$ variation have typically not accounted for geographic variation in source signals. As an
622 example, Rice et al., (2016) apply a constant $\delta^2\text{H-CH}_4$ of -322‰ for both low-latitude ($0\text{--}30^\circ\text{ N}$) and high latitude (30--
623 90° N) wetland emissions. Based on our dataset this estimate is an inaccurate representation of wetland $\delta^2\text{H-CH}_4$ for
624 either $0\text{--}30^\circ\text{ N}$ (mean: $-305 \pm 13\text{‰}$) or $30\text{--}90^\circ\text{ N}$ (mean: $-345 \pm 11\text{‰}$). Studies of ice core measurements have more
625 frequently differentiated freshwater $\delta^2\text{H-CH}_4$ values as a function of latitude. For example, Bock et al., (2010)
626 differentiated $\delta^2\text{H-CH}_4$ between tropical (-320‰) and boreal (-370‰) wetlands. This tropical wetland signature is
627 significantly lower than our estimate of low-latitude wetland $\delta^2\text{H-CH}_4$, although the boreal wetland signature is similar to
628 our mean value for high-latitude wetlands ($-374 \pm 10\text{‰}$). Overall, our results imply that accounting for latitudinal variation
629 in freshwater $\delta^2\text{H-CH}_4$, along with accurate latitudinal flux estimates, is important for developing accurate estimates of
630 global freshwater $\delta^2\text{H-CH}_4$ source signatures.

631 Our analysis indicates significant differences in the distribution of freshwater $\delta^{13}\text{C-CH}_4$ between the low- and
632 high-latitudes, but mid-latitude sites cannot be differentiated. Furthermore our results do not indicate significant
633 latitudinal differences in $\delta^{13}\text{C-CH}_4$ for wetland sites in particular. This is in contrast to previous studies that have
634 inferred significant differences in wetland $\delta^{13}\text{C-CH}_4$ by latitude (Bock et al., 2010; Rice et al., 2016; Ganesan et al., 2018).
635 An important caveat is that we have not analyzed a comprehensive dataset of freshwater $\delta^{13}\text{C-CH}_4$, for which there are
636 much more published data than for $\delta^2\text{H-CH}_4$, although our analysis does comprise the largest dataset of freshwater $\delta^{13}\text{C-}$
637 CH_4 compiled to date (See Sect. 2.3). In addition, our analysis does not take into account the geographic distribution of
638 different ecosystem categories, although we do not find significant differences in $\delta^{13}\text{C-CH}_4$ between ecosystem
639 categories (Fig. 8; Sect. 3.6). Low-latitude ecosystems dominated by C_4 plants are underrepresented both in our dataset
640 and that of Sherwood et al., (2017), and accounting for this would likely lead to a more enriched low-latitude wetland
641 $\delta^{13}\text{C-CH}_4$. In contrast, high-latitude ecosystems, including bogs, are relatively well represented in these datasets (Fig. 8),
642 and we suggest that inferences of especially low $\delta^{13}\text{C-CH}_4$ in high-latitude wetlands (Bock et al., 2010; Rice et al.,
643 2016; Ganesan et al., 2018) are not consistent with the compiled dataset of in-situ measurements. However, we note that
644 atmospheric estimates of high-latitude wetland $\delta^{13}\text{C-CH}_4$ ($\sim -68 \pm 4\text{‰}$; Fisher et al., 2011) are lower than the median or
645 mean value shown in Figure 7B, and are in close agreement with the relatively low values predicted by (Ganesan et al.,
646 2018). Ombrotrophic and minerotrophic peatlands have distinctive $\delta^{13}\text{C-CH}_4$ signatures (Bellisario et al., 1999; Bowes

Peter Douglas 5/11/2021 11:55 AM

Deleted: higher

Peter Douglas 5/11/2021 2:30 PM

Deleted: , whereas we do observe significant differences between the low- and high-latitudes for inland water sites

Peter Douglas 5/11/2021 11:56 AM

Deleted: . However,

Peter Douglas 5/11/2021 11:56 AM

Deleted: especially

653 and Hornibrook, 2006;Hornibrook, 2009), with lower signatures in ombrotrophic peatlands. We did not differentiate
654 peatlands by trophic status, and it is possible that the dataset of high-latitude wetland in-situ measurements is biased
655 towards minerotrophic peatlands with relatively high $\delta^{13}\text{C-CH}_4$.

656 Latitudinal differences in $\delta^{13}\text{C-CH}_4$ inferred by Ganesan et al. (2018) were based on two key mechanisms: (1)
657 differences in methanogenic pathway between different types of wetlands, especially between minerotrophic fens and
658 ombrotrophic bogs; and (2) differential inputs of organic matter from C_3 and C_4 plants. Because inferred latitudinal
659 differences in $\delta^{13}\text{C-CH}_4$ and $\delta^2\text{H-CH}_4$ are caused by different mechanisms, they could be highly complementary in
660 validating estimates of freshwater emissions by latitude. It is also important to note that previous assessments of
661 latitudinal differences in $\delta^{13}\text{C-CH}_4$ did not include inland water environments. Our analysis suggests that latitudinal
662 variation in $\delta^{13}\text{C-CH}_4$ in inland waters may be more pronounced than in wetlands, although the mechanisms causing this
663 difference will need to be elucidated with further study. A benefit of geographic discrimination based on $\delta^2\text{H-CH}_4$ is that
664 the same causal mechanism applies to all freshwater emissions, including both wetlands and inland waters.

665 3.5.1 Potential for geographic discrimination of other microbial methane sources based on $\delta^2\text{H-CH}_4$

666 We speculate that latitudinal differences in $\delta^2\text{H-CH}_4$ should also be observed in other fluxes of microbial methane from
667 terrestrial environments, including enteric fermentation in livestock and wild animals, manure ponds, landfills, and
668 termites. This is because microbial methanogenesis in all of these environments will incorporate hydrogen from
669 environmental water, and therefore will be influenced by variation in precipitation $\delta^2\text{H}$. There are limited data currently
670 available to test this prediction, but $\delta^2\text{H-CH}_4$ data from cow rumen and landfills are available with either specified
671 locations or $\delta^2\text{H-H}_2\text{O}$ (Burke Jr, 1993;Levin et al., 1993;Liptay et al., 1998;Bilek et al., 2001;Wang et al., 2015;Teasdale
672 et al., 2019). These data plot in a range that is consistent with the $\delta^2\text{H-CH}_4$ vs. $\delta^2\text{H-H}_2\text{O}$ relationships for freshwater CH_4
673 (Fig. 5). Landfill data are only available for a very small range of estimated $\delta^2\text{H-H}_2\text{O}$, making it impossible to assess for
674 geographic variation currently. $\delta^2\text{H-CH}_4$ data from cow rumen span a much wider range, and express substantial variation
675 that is independent of $\delta^2\text{H-H}_2\text{O}$, but largely overlap measurements from freshwater environments. Based on these limited
676 data, variation observed in incubation studies that simulate landfill conditions (Schoell, 1980;Waldron et al., 1998), and
677 our understanding of the influence of $\delta^2\text{H-H}_2\text{O}$ on microbial $\delta^2\text{H-CH}_4$ (Fig. 6), we suggest that both landfill and cow
678 rumen $\delta^2\text{H-CH}_4$ likely vary geographically as a function of $\delta^2\text{H-H}_2\text{O}$. If validated, this variation could also be used to
679 distinguish these CH_4 sources geographically. More data are clearly needed to test this conjecture, and it will also be
680 important to evaluate how closely annual or seasonal $\delta^2\text{H}_p$ corresponds to environmental $\delta^2\text{H-H}_2\text{O}$ in both landfills and
681 cow rumen. Relatedly, the $\delta^2\text{H}$ of CH_4 emitted by biomass burning or directly by plants has also been shown to vary as a
682 function of $\delta^2\text{H-H}_2\text{O}$ (Vigano et al., 2010;Umezawa et al., 2011).

Peter Douglas 5/13/2021 1:38 PM

Deleted: . However, the cow rumen data

Peter Douglas 5/13/2021 1:38 PM

Deleted: span a range that is similar to that
observed in

687 When comparing ecosystems, we analyze $\delta^2\text{H-CH}_{4,\text{w0}}$ values to account for variability related to differences in $\delta^2\text{H-H}_2\text{O}$.
 688 Ecosystem types are not evenly distributed by latitude, and therefore have different distributions of $\delta^2\text{H-H}_2\text{O}$ values. Our
 689 analysis does not find a significant difference in the distribution of $\delta^2\text{H-CH}_{4,\text{w0}}$ between ecosystems, which could be
 690 partly a result of small sample sizes for most ecosystem categories (Fig. 8). Comparing the broader categories of inland
 691 waters and wetlands, we do observe a significant difference in $\delta^2\text{H-CH}_{4,\text{w0}}$ distributions, with inland waters shifted
 692 towards higher values (median: -296‰) than wetlands (median: -311‰). We repeated this analysis only including sites
 693 with measured $\delta^2\text{H-H}_2\text{O}$ and found the same results in terms of category differences (Supplemental Figure S1). We did
 694 not observe any significant differences in $\delta^{13}\text{C-CH}_4$ distributions between ecosystems, nor was there a significant
 695 difference in $\delta^{13}\text{C-CH}_4$ distributions between inland waters and wetlands.

696 The significant difference in the distribution of $\delta^2\text{H-CH}_{4,\text{w0}}$ between inland waters and wetlands is primarily a
 697 result of the difference in $\delta^2\text{H-CH}_4$ between these environments in the high latitudes (Figs. 3, 4, and 7). We suggest this
 698 difference could be related to a greater overall prevalence of CH_4 oxidation in inland waters. As shown in Figure 6, the
 699 lack of positive co-variation between $\delta^2\text{H-CH}_{4,\text{w0}}$ and $\delta^{13}\text{C-CO}_2$, and α_c could be interpreted to support a greater role for
 700 CH_4 oxidation to control $\delta^2\text{H-CH}_{4,\text{w0}}$ in inland waters relative to wetlands, although this result requires further validation.
 701 In lakes that undergo seasonal overturning and water column oxygenation there may be a greater overall effect of CH_4
 702 oxidation than there are in wetlands typically. The absence of significant differences between ecosystems in terms of
 703 $\delta^{13}\text{C-CH}_4$ (Fig. 8B) is in contrast to previous studies that have suggested that fens and bogs in particular have distinctive
 704 $\delta^{13}\text{C-CH}_4$ (Ganesan et al., 2018). Bogs in particular have a very wide distribution of $\delta^{13}\text{C-CH}_4$ that could represent
 705 differences between minerotrophic and ombrotrophic bogs (Hornibrook, 2009), which we did not differentiate in our
 706 dataset. This result should be interpreted with caution given that our dataset is not a comprehensive compilation of
 707 published $\delta^{13}\text{C-CH}_4$ data, although it is the largest compiled dataset available (Sect. 2.3). We argue that inferred
 708 differences in $\delta^{13}\text{C-CH}_4$ between wetland ecosystem categories should be further verified with more comprehensive data
 709 assimilation and additional measurements.

Peter Douglas 5/11/2021 11:59 AM

Deleted: There are differences in the median values by ecosystem, with rivers (-283‰) exhibiting relatively enriched median $\delta^2\text{H-CH}_{4,\text{w0}}$, and fens (-310‰) and rice paddies (-314‰) exhibiting relatively low median values. However, given the small sample sizes and large variance in most of these categories, o

Peter Douglas 5/11/2021 11:59 AM

Deleted: infer

Peter Douglas 5/11/2021 9:33 AM

Deleted: with a

Peter Douglas 5/11/2021 9:34 AM

Deleted: find

Peter Douglas 5/11/2021 9:34 AM

Deleted: ,

Peter Douglas 5/11/2021 12:00 PM

Deleted: The median $\delta^{13}\text{C-CH}_4$ value for bogs was relatively low (-66‰), while median values for fens (-60.3‰) and rice paddies (-60.3‰) were relatively high, but there was a large range in values for all of these ecosystems.

Peter Douglas 5/13/2021 1:39 PM

Deleted: the overarching categories of

Peter Douglas 5/11/2021 12:07 PM

Deleted: are unsure of the mechanism causing this difference, though it

Peter Douglas 5/11/2021 12:07 PM

Deleted: is likely

Peter Douglas 5/11/2021 12:10 PM

Deleted: The absence of significant differences in $\delta^2\text{H-CH}_{4,\text{w0}}$ distributions between specific ecosystem categories could be the result of small samples sizes for most ecosystems. Further study could be targeted towards verifying and testing the apparent differences shown in Figure 8A. Generally lower $\delta^2\text{H-CH}_{4,\text{w0}}$ in rice paddies and fens could reflect a greater proportion of acetoclastic methanogenesis inferred for these ecosystems (Conrad and Klose, 1999; Hornibrook, 2009; Ganesan et al., 2018), or possibly more thermodynamically favorable methanogenesis related to high carbon substrate, H_2 , or nutrient concentrations. Both of these explanations would be consistent with the relatively high median $\delta^{13}\text{C-CH}_4$ values in these ecosystems (Fig. 7B, see also Fig. 6A). High median values in river ecosystems, in contrast, may be a function of generally greater rates of oxidation, given that these environments are also characterized by relatively high $\delta^{13}\text{C-CH}_4$ (Fig. 8B), and the potential for greater water-column oxygenation in fluvial environments with turbulent flow (Devol et al., 1987). However, our river dataset is highly biased [2]

Peter Douglas 5/11/2021 2:33 PM

Deleted:




Figure 8: Boxplots of (A) $\delta^2\text{H-CH}_{4,w0}$ and (B) $\delta^{13}\text{C-CH}_4$ for sites differentiated by ecosystem type. Numbers in parentheses indicate the number of sites for each category. Boxplot parameters are as in Fig. 7. Black points and error bars indicate the category mean and 95% confidence interval of the mean. Gray lines indicate the mean values across all categories and the dashed lines indicate the 95% confidence interval of this value. Two extremely low outliers (<-80‰; lake and fen) are not shown in (B). IW- Inland Waters; WL- Wetlands; S/M- Swamps and marshes. Asterisks in A indicate that inland waters and wetlands have significantly different distributions.

3.7 Differences in $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ by sample type

As with comparing ecosystems, when comparing sample types we analyze $\delta^2\text{H-CH}_{4,w0}$ values to normalize for variability related to differences in $\delta^2\text{H-H}_2\text{O}$, since sample types are not distributed evenly by latitude. When comparing sample types, dissolved CH_4 samples do not have a significantly different $\delta^2\text{H-CH}_{4,w0}$ distribution for the dataset as a whole, nor is there a significant difference between these groups in wetland sites (Fig. 9A). There is, however, a significant difference in inland water sites, with dissolved CH_4 samples having a more enriched distribution (median: -270‰) vs. gas samples (median: -302‰). We repeated this analysis only including sites with measured $\delta^2\text{H-H}_2\text{O}$ and found the same results in terms of category differences (Supplemental Figure S2). We did not observe a significant difference in the distribution of $\delta^{13}\text{C}$ between dissolved and gas-phase CH_4 samples, either for the dataset as a whole or when the dataset was disaggregated into wetlands and inland waters (Fig. 9B).

We suggest that the higher $\delta^2\text{H-CH}_{4,w0}$ in dissolved vs. gas samples for inland waters could be a result of generally greater oxidation of dissolved CH_4 in inland water environments, potentially as a result of longer exposure to aerobic conditions in lake or river water columns. This is in contrast to wetlands, where aerobic conditions are generally limited to the uppermost layers of wetlands proximate to the water table. However, our dataset for inland water dissolved

Peter Douglas 5/11/2021 10:35 AM

Deleted: 5

Peter Douglas 5/11/2021 2:37 PM

Deleted: s

Peter Douglas 5/11/2021 12:12 PM

Formatted: Indent: First line: 0.5"

CH₄ is quite small (n=9), and more data are needed to test this hypothesis. Furthermore, it is unclear why oxidation in inland water dissolved CH₄ would be more strongly expressed in terms of δ²H-CH_{4,w0} (Fig. 9A) than δ¹³C (Fig. 9B). Overall, our data imply that isotopic differences between dissolved and gas phase methane are relatively minor on a global basis, especially in wetlands. This result could imply that the relative balance of diffusive vs. ebullition gas fluxes does not have a large effect on the isotopic composition of freshwater CH₄ emissions. However, our study does not specifically account for isotopic fractionation occurring during diffusive or plant-mediated transport (Hornibrook, 2009), and most of our dissolved sample data are of *in-situ* dissolved CH₄ and not diffusive fluxes. More isotopic data specifically focused on diffusive methane emissions, for example using measurements of gas sampled from chambers, would help to resolve this question, as would more comparisons of the isotopic composition of diffusive and ebullition CH₄ emissions from the same ecosystem.




Figure 9: Boxplots of (A) δ²H-CH_{4,w0} and (B) δ¹³C-CH₄ for sites differentiated by sample type. Numbers in parentheses indicate the number of sites for each category. Boxplot parameters are as in Fig. 7. Black points and error bars indicate the category mean and 95% confidence interval of the mean. Gray lines indicate the mean values across all categories and the dashed lines indicate the 95% confidence interval of this value. Two extremely low outliers (<-80‰; dissolved wetland and gas inland water) are not shown in (B). Asterisks in A indicate that dissolved and gas-phase CH₄ samples from inland water sites have significantly different distributions.

3.8 Estimates of global emissions source δ²H-CH₄ and δ¹³C-CH₄

Our mixing model estimates a global source δ²H-CH₄ of -278±15‰, and a global source δ¹³C-CH₄ of -56.4±2.6‰ (Fig. 10). Monte Carlo sensitivity tests that only included uncertainty in either isotopic source signatures or flux estimates suggest that larger uncertainty is associated with isotopic source signatures (12‰ for δ²H; 2.2‰ for δ¹³C) than with flux estimates (8‰ for δ²H; 1.4‰ for δ¹³C). When correcting for wetland and biomass burning emissions from C₄ plant ecosystems, as described in Section 2.4, our estimate of global source δ¹³C-CH₄ increases to -55.2±2.6‰. Our estimate of


Peter Douglas 5/11/2021 12:13 PM
Deleted: n

Peter Douglas 5/11/2021 2:37 PM
Deleted: s

Peter Douglas 5/17/2021 3:48 PM
Deleted: and Monte Carlo analyses

Peter Douglas 5/13/2021 1:43 PM
Deleted: -

832 global source $\delta^2\text{H-CH}_4$ is substantially higher than a previous bottom-up estimate using a similar approach (-295‰; Fig.
 833 10) (Whiticar and Schaefer, 2007). This difference can be largely attributed to the application of more depleted $\delta^2\text{H-CH}_4$
 834 source signatures for tropical wetlands (-360 ‰), and to a lesser extent boreal wetlands (-380 ‰), by Whiticar and
 835 Schaefer (2007).



836
 837 **Figure 10: Comparison of estimates of dual-isotope global source $\delta^2\text{H-CH}_4$ and $\delta^{13}\text{C-CH}_4$ from this and previous studies. Error**
 838 **bars from this study indicate the 2σ standard deviation from Monte Carlo analysis. Gray dot and error bars indicate an**
 839 **estimate corrected for the lack of data from wetlands and biomass burning in C_4 plant environments, as described in Sect. 2.4.**
 840 **Error bars for Rice et al., (2016), indicate the range of values estimated in that study between 1977-2005. Error bars for**
 841 **Sherwood et al., (2017) reflect the combined measurement uncertainty and uncertainty in sink fractionations reported in that**
 842 **study. Whiticar and Schaefer (2007) did not provide uncertainties for their estimates.**

843 Our bottom-up estimate of global source $\delta^2\text{H-CH}_4$ substantially overlaps the range of top-down estimates (-258
 844 to -289‰) based on atmospheric $\delta^2\text{H-CH}_4$ measurements from 1977-2005 and a box model of sink fluxes and kinetic
 845 isotope effects (Rice et al., 2016) (Fig. 10). It is also within error of simpler top-down estimates calculated based on
 846 mean atmospheric measurements and estimates of a constant sink fractionation factor (Whiticar and Schaefer,
 847 2007; Sherwood et al., 2017). Sherwood et al., (2017) estimate a very wide range of possible global source $\delta^2\text{H-CH}_4$
 848 values based on a relatively large atmospheric sink fractionation with large uncertainty (-235±80‰). This range overlaps
 849 with our bottom up estimate, although its mid-point is substantially lower than our estimate. We argue that the box-model

Peter Douglas 5/11/2021 2:38 PM

Deleted: by

Peter Douglas 5/17/2021 3:50 PM

Deleted: Another key difference is their inclusion of a relatively large flux and enriched $\delta^2\text{H-CH}_4$ signature from aerobic methane production from plants by Whiticar and Schaefer (2007), which is not included as a CH_4 source in our calculations.

856 method used to account for sink fractionations applied by Rice et al. (2016) probably provides a more accurate
857 representation of global-source isotopic composition than the other top-down estimates shown in Figure 10. The
858 estimates of Rice et al. (2016) are also supported by the results of a global inversion model. Overall, the overlap between
859 our bottom-up estimate of global source $\delta^2\text{H-CH}_4$ with top-down estimates is encouraging, and suggests that the
860 estimates of emission source $\delta^2\text{H-CH}_4$ signatures applied in this study are reasonably accurate. However, as discussed
861 below, there is still substantial [scope](#) to further constrain these estimates and to reduce uncertainty.

862 Our bottom-up estimate of global source $\delta^{13}\text{C-CH}_4$ is lower than the other top-down and bottom-up estimates
863 shown in Figure 10. As discussed above, there is likely a bias in our freshwater CH_4 isotopic database in that it includes
864 very few wetland sites from C_4 -plant dominated ecosystems. When correcting for this, as well as [for](#) CH_4 emissions from
865 [combustion of \$\text{C}_4\$ plants](#) (Fig. 10), our estimate shifts to a more enriched value that is within uncertainty of other
866 estimates. Clearly, accounting for the effect of C_4 plants in wetland and biomass burning CH_4 emissions, and potentially
867 also in enteric fermentation emissions, is important for accurate estimates of global source $\delta^{13}\text{C-CH}_4$. As discussed
868 below, other sources of error in both isotopic source signatures and inventory-based flux estimates could also partially
869 account for our relatively low global source $\delta^{13}\text{C-CH}_4$ estimate. [For example, variation in fossil fuel isotopic signatures](#)
870 [between regions and resource types is potentially an additional source of uncertainty that is not accounted for in this](#)
871 [estimate](#).

872 Previous studies have argued, on the basis of comparing atmospheric measurements and emissions source $\delta^{13}\text{C-}$
873 CH_4 signatures, that there are biases in global emissions inventories, specifically that fossil fuel emissions estimates are
874 too low, and that either microbial emissions estimates are too high (Schwietzke et al., 2016), or that biomass burning
875 estimates are too high (Worden et al., 2017). We argue that greater analysis of $\delta^2\text{H-CH}_4$ measurements could be valuable
876 for evaluating these and other emissions scenarios, as has been suggested previously (Rigby et al., 2012). This is
877 especially true for determining the relative proportion of fossil fuel and microbial emissions, since these sources have
878 widely differing $\delta^2\text{H-CH}_4$ signatures (Table 1). Currently, atmospheric $\delta^2\text{H-CH}_4$ measurements are not a routine
879 component of CH_4 monitoring programs, but we argue that based on both their value in constraining emissions sources
880 and sinks (Rigby et al., 2012), and the increasing practicality of high-frequency measurements (Chen et al.,
881 2016; Röckmann et al., 2016; Yacovitch et al., 2020), that there should be a renewed focus on these measurements.

882 The uncertainty in our bottom-up estimates, the overall greater uncertainty associated with isotopic source
883 signatures in our Monte Carlo calculations, and the apparent discrepancies for $\delta^{13}\text{C-CH}_4$ shown in Figure 10, also imply
884 that isotopic source signatures for specific sources could be greatly improved. As noted by Rigby et al. (2012), the impact
885 of improved isotopic source signatures increases as measurement precision improves. We have discussed above the
886 importance of increased data assimilation and measurements from tropical wetlands, with a particular focus on C_4 plant
887 dominated ecosystems. Using the isotopic source signal uncertainties and emissions fluxes shown in Table 1, we
888 identified the sources with the greatest flux-weighted uncertainty in isotopic signatures. Based on this analysis, the

Peter Douglas 5/13/2021 1:44 PM

Deleted: (Whiticar and Schaefer, 2007; Sherwood et al., 2017)

Peter Douglas 5/13/2021 1:45 PM

Deleted: room

Peter Douglas 5/11/2021 2:42 PM

Deleted: biomass burning

greatest uncertainty for global source $\delta^2\text{H-CH}_4$ estimates comes from source signatures for enteric fermentation and manure, low-latitude wetlands, onshore geological emissions, low-latitude and mid-latitude inland waters, termites, and landfills. We identified the same source categories as having the greatest flux-weighted uncertainty for $\delta^{13}\text{C-CH}_4$, with the exception of termites, but repeat the caveat that the underlying dataset is less comprehensive for $\delta^{13}\text{C-CH}_4$. We argue that these source categories should be considered priorities for future emissions source isotopic characterization through data assimilation and additional measurements. As discussed in Sect. 3.5.1, evaluation of possible latitudinal variation in enteric fermentation and landfill $\delta^2\text{H-CH}_4$ is particularly promising.

Peter Douglas 5/11/2021 12:39 PM

Deleted: In particular, a

5 Conclusions

Our analysis of an expanded isotopic dataset for freshwater CH_4 confirms the previous finding that $\delta^2\text{H-H}_2\text{O}$ is the primary determinant of $\delta^2\text{H-CH}_4$ on a global scale (Waldron et al., 1999a), but also finds that the slope of this relationship is probably flatter than was inferred previously (Fig. 3). This flatter slope is primarily the result of the inclusion of a much larger number of high-latitude sites with low $\delta^2\text{H-H}_2\text{O}$ in our dataset. We find that the inferred relationship between $\delta^2\text{H-CH}_4$ and $\delta^2\text{H-H}_2\text{O}$ is not highly sensitive to whether measured $\delta^2\text{H-H}_2\text{O}$, modeled $\delta^2\text{H}_p$, or a combination of the two (i.e. a best-estimate) is used to estimate $\delta^2\text{H-H}_2\text{O}$. This implies that gridded datasets of $\delta^2\text{H}_p$ or isotope-enabled climate models could be used to predict the distribution of $\delta^2\text{H-CH}_4$ in the present, as well as under past and future climates. Our analysis also suggests that annual $\delta^2\text{H}_p$ may be a better predictor for wetland $\delta^2\text{H-CH}_4$, while seasonal $\delta^2\text{H}_p$ may be a better predictor of inland water $\delta^2\text{H-CH}_4$. The slope of $\delta^2\text{H-CH}_4$ vs. $\delta^2\text{H-H}_2\text{O}$ in both wetlands and inland waters agrees well with that found in incubation experiments (Schoell, 1980; Sugimoto and Wada, 1995; Waldron et al., 1998; Waldron et al., 1999a), and we concur with previous inferences that this slope is partly controlled by variation in the $\delta^2\text{H}$ of acetate as a function of $\delta^2\text{H-H}_2\text{O}$ (Waldron et al., 1999a). Analysis of co-variation of $\delta^2\text{H-CH}_{4,w0}$ with $\delta^{13}\text{C-CH}_4$, $\delta^{13}\text{C-CO}_2$, and α_c suggest that residual variation in $\delta^2\text{H-CH}_4$ is influenced by a complex set of biogeochemical variables, including both variable isotopic fractionation related to methanogenesis, and post-production isotopic fractionation related to CH_4 oxidation and diffusive gas transport. A significant positive correlation between $\delta^2\text{H-CH}_{4,w0}$ and both $\delta^{13}\text{C-CO}_2$, and α_c in wetlands suggests that [hydrogen isotope](#) fractionation related to methanogenesis pathway [or enzymatic reversibility](#) may be more important in these environments, but this result is dependent on the method used to estimate $\delta^2\text{H-H}_2\text{O}$ and requires further validation.

Peter Douglas 5/11/2021 2:44 PM

Deleted: variable

Peter Douglas 5/13/2021 1:48 PM

Deleted: and

Peter Douglas 5/13/2021 1:49 PM

Deleted: thermodynamics

The dependence of $\delta^2\text{H-CH}_4$ on $\delta^2\text{H-H}_2\text{O}$ leads to clear latitudinal differences in $\delta^2\text{H-CH}_4$, with particularly low values from high latitude sites (Fig. 4; Fig. 7A). The mechanism for latitudinal differences in $\delta^2\text{H-CH}_4$ is distinct from proposed mechanisms for latitudinal differences in $\delta^{13}\text{C-CH}_4$ (Ganesan et al., 2018), implying that these two isotopic tracers are complementary in differentiating geographic emissions sources. We estimate a global flux-weighted $\delta^2\text{H-CH}_4$ signature from freshwater environments of $-310 \pm 15\text{‰}$, which is enriched relative to values used in previous source

Peter Douglas 5/11/2021 2:45 PM

Deleted:

929 apportionment studies (Rice et al., 2016; Bock et al., 2017). We observe a significantly higher $\delta^2\text{H-CH}_{4,\text{w0}}$ distribution in
930 inland waters relative to wetlands (Fig. 8A), which we suggest is a result of greater rates of CH_4 oxidation in inland
931 waters. We do not find significant differences between more specific ecosystem categories, nor do we find significant
932 differences in $\delta^2\text{H-CH}_{4,\text{w0}}$ between sample types (Fig. 9A), with the exception of higher values in dissolved CH_4 relative
933 to gas-phase CH_4 in inland water environments.

934 Our bottom-up estimate of the global $\delta^2\text{H-CH}_4$ source signature, $-278 \pm 15\text{‰}$, is higher than previous bottom-up
935 estimates (Whiticar and Schaefer, 2007), but is within the range of top-down estimates based on atmospheric
936 measurements and modeled sink fractionations (Rice et al., 2016). In contrast, our bottom-up estimate of global $\delta^{13}\text{C-}$
937 CH_4 , $-56.4 \pm 2.6\text{‰}$, is low relative to top-down estimates, which is partially explained by a lack of data from C_4 plant-
938 dominated ecosystems in the freshwater CH_4 isotopic dataset. The agreement between bottom-up and top-down global
939 $\delta^2\text{H-CH}_4$ estimates suggests that our current understanding of $\delta^2\text{H-CH}_4$ source signatures, when combined with
940 inventory-based flux estimates (Saunio et al., 2020), is consistent with atmospheric measurements. This supports the
941 argument that increased measurements and modeling of atmospheric $\delta^2\text{H-CH}_4$ could help to constrain global CH_4
942 budgets (Rigby et al., 2012). However, there is clearly a need to better constrain source signatures for both $\delta^2\text{H-CH}_4$ and
943 $\delta^{13}\text{C-CH}_4$, especially from low-latitude microbial sources.

944
945 **Data Availability:** The datasets used in this paper (Supplementary Tables 1-4) are publicly available: Douglas, Peter;
946 Stratigopoulos, Emerald; Park, Jenny; Phan, Dawson (2020): Data for geographic variability in freshwater methane
947 hydrogen isotope ratios and its implications for emissions source apportionment and microbial biogeochemistry. figshare.
948 Dataset. <https://doi.org/10.6084/m9.figshare.13194833.v2>

949
950 **Author Contribution:** PMJD designed the project, assisted with compiling the data, analyzed the data, and wrote the
951 manuscript; ES and JP compiled the data, and assisted with analyzing the data and editing the manuscript; DP developed
952 code for mixing model and Monte Carlo calculations, and assisted with analyzing the data and editing the manuscript.

953
954 **Competing Interests:** The authors declare they have no competing interests.

955
956 **Acknowledgments:** We thank all of the researchers whose published data made this analysis possible (See Supplemental
957 Table 1). We also thank Susan Waldron, Edward Hornibrook, and an anonymous reviewer for constructive feedback.
958 This research was partially funded by McGill Science Undergraduate Research Awards to ES and JP and by NSERC
959 Discovery Grant 2017-03902 to PMJD.

960
961

Peter Douglas 5/11/2021 12:40 PM

Deleted: , but there are apparent differences between some wetland ecosystems that could be verified with larger datasets

Peter Douglas 5/11/2021 2:46 PM

Deleted: . We also do not

Peter Douglas 5/13/2021 1:50 PM

Deleted: ,

- 968
- 969 Alstad, K. P., and Whiticar, M. J.: Carbon and hydrogen isotope ratio characterization of methane dynamics for Fluxnet
 970 Peatland Ecosystems, *Org Geochem*, 42, 548-558, 2011.
- 971 Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prast, A.: Freshwater methane emissions offset the
 972 continental carbon sink, *Science*, 331, 50-50, 2011.
- 973 Bellisario, L., Bubier, J., Moore, T., and Chanton, J.: Controls on CH₄ emissions from a northern peatland, *Global*
 974 *Biogeochem Cy*, 13, 81-91, 1999.
- 975 Bergamaschi, P.: Seasonal variations of stable hydrogen and carbon isotope ratios in methane from a Chinese rice paddy,
 976 *Journal of Geophysical Research: Atmospheres*, 102, 25383-25393, 1997.
- 977 Bilek, R., Tyler, S., Kurihara, M., and Yagi, K.: Investigation of cattle methane production and emission over a 24-hour
 978 period using measurements of $\delta^{13}\text{C}$ and δD of emitted CH₄ and rumen water, *Journal of Geophysical Research:*
 979 *Atmospheres*, 106, 15405-15413, 2001.
- 980 Bock, M., Schmitt, J., Möller, L., Spahn, R., Blunier, T., and Fischer, H.: Hydrogen isotopes preclude marine hydrate
 981 CH₄ emissions at the onset of Dansgaard-Oeschger events, *Science*, 328, 1686-1689, 2010.
- 982 Bock, M., Schmitt, J., Beck, J., Seth, B., Chappellaz, J., and Fischer, H.: Glacial/interglacial wetland, biomass burning,
 983 and geologic methane emissions constrained by dual stable isotopic CH₄ ice core records, *Proceedings of the National*
 984 *Academy of Sciences*, 114, E5778-E5786, 2017.
- 985 Bouchard, F., Laurion, I., Preskienis, V., Fortier, D., Xu, X., and Whiticar, M.: Modern to millennium-old greenhouse
 986 gases emitted from ponds and lakes of the Eastern Canadian Arctic (Bylot Island, Nunavut), *Biogeosciences*, 12, 7279-
 987 7298, 2015.
- 988 Bowen, G. J., and Wilkinson, B.: Spatial distribution of $\delta^{18}\text{O}$ in meteoric precipitation, *Geology*, 30, 315-318, 2002.
- 989 Bowen, G. J., and Revenaugh, J.: Interpolating the isotopic composition of modern meteoric precipitation, *Water*
 990 *Resources Research*, 39, 2003.
- 991 Bowen, G. J., Wassenaar, L. L., and Hobson, K. A.: Global application of stable hydrogen and oxygen isotopes to wildlife
 992 forensics, *Oecologia*, 143, 337-348, 2005.
- 993 Bowes, H. L., and Hornibrook, E. R.: Emission of highly ¹³C-depleted methane from an upland blanket mire, *Geophys*
 994 *Res Lett*, 33, 2006.
- 995 Brooks, J. R., Gibson, J. J., Birks, S. J., Weber, M. H., Rodecap, K. D., and Stoddard, J. L.: Stable isotope estimates of
 996 evaporation: inflow and water residence time for lakes across the United States as a tool for national lake water quality
 997 assessments, *Limnol Oceanogr*, 59, 2150-2165, 2014.
- 998 Brosius, L., Walter Anthony, K., Grosse, G., Chanton, J., Farquharson, L., Overduin, P. P., and Meyer, H.: Using the
 999 deuterium isotope composition of permafrost meltwater to constrain thermokarst lake contributions to atmospheric CH₄
 1000 during the last deglaciation, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 117, 2012.
- 1001 Burke Jr, R.: SHALLOW AQUATIC SEDIMENTS, *Bacterial Gas. Conference*, Mila, September 25-26, 1989, 1992, 47,
- 1002 Burke Jr, R. A., and Sackett, W. M.: Stable hydrogen and carbon isotopic compositions of biogenic methanes from
 1003 several shallow aquatic environments, in, *ACS Publications*, 1986.
- 1004 Burke Jr, R. A., Barber, T. R., and Sackett, W. M.: Methane flux and stable hydrogen and carbon isotope composition of
 1005 sedimentary methane from the Florida Everglades, *Global Biogeochem Cy*, 2, 329-340, 1988.
- 1006 Burke Jr, R. A., Barber, T. R., and Sackett, W. M.: Seasonal variations of stable hydrogen and carbon isotope ratios of
 1007 methane in subtropical freshwater sediments, *Global Biogeochem Cy*, 6, 125-138, 1992.
- 1008 Burke Jr, R. A.: Possible influence of hydrogen concentration on microbial methane stable hydrogen isotopic
 1009 composition, *Chemosphere*, 26, 55-67, 1993.
- 1010 Cadieux, S. B., White, J. R., Sauer, P. E., Peng, Y., Goldman, A. E., and Pratt, L. M.: Large fractionations of C and H
 1011 isotopes related to methane oxidation in Arctic lakes, *Geochim Cosmochim Acta*, 187, 141-155, 2016.
- 1012 Chanton, J., Whiting, G., Blair, N., Lindau, C., and Bollich, P.: Methane emission from rice: Stable isotopes, diurnal
 1013 variations, and CO₂ exchange, *Global Biogeochem Cy*, 11, 15-27, 1997.
- 1014 Chanton, J. P.: The effect of gas transport on the isotope signature of methane in wetlands, *Org Geochem*, 36, 753-768,
 1015 2005.

Peter Douglas 5/13/2021 1:50 PM

Deleted: -

Peter Douglas 5/17/2021 3:58 PM

Deleted: , -, Artn 1299 -

... [4]

Chanton, J. P., Fields, D., and Hines, M. E.: Controls on the hydrogen isotopic composition of biogenic methane from high-latitude terrestrial wetlands, *Journal of Geophysical Research: Biogeosciences* (2005–2012), 111, 2006.

Chasar, L., Chanton, J., Glaser, P. H., and Siegel, D.: Methane concentration and stable isotope distribution as evidence of rhizospheric processes: Comparison of a fen and bog in the Glacial Lake Agassiz Peatland complex, *Annals of Botany*, 86, 655–663, 2000.

Chen, Y., Lehmann, K. K., Peng, Y., Pratt, L., White, J., Cadieux, S., Sherwood Lollar, B., Lacrampe-Couloume, G., and Onstott, T. C.: Hydrogen Isotopic Composition of Arctic and Atmospheric CH₄ Determined by a Portable Near-Infrared Cavity Ring-Down Spectrometer with a Cryogenic Pre-Concentrator, *Astrobiology*, 16, 787–797, 2016.

Conrad, R., Noll, M., Claus, P., Klose, M., Bastos, W., and Enrich-Prast, A.: Stable carbon isotope discrimination and microbiology of methane formation in tropical anoxic lake sediments, *Biogeosciences*, 8, 795, 2011.

Coplen, T. B.: Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results, *Rapid Commun Mass Sp*, 25, 2538–2560, 2011.

David, K., Timms, W., Hughes, C. E., Crawford, J., and McGeeney, D.: Application of the pore water stable isotope method and hydrogeological approaches to characterise a wetland system, *Hydrology and Earth System Sciences*, 22, 6023–6041, 2018.

Douglas, P. M. J., Stolper, D., Smith, D., Walter Anthony, K., Paull, C., Dallimore, S., Wik, M., Crill, P., Winterdahl, M., Eiler, J., and Sessions, A. L.: Diverse origins of Arctic and Subarctic methane point source emissions identified with multiply-substituted isotopologues, *Geochim Cosmochim Acta*, 188, 163–188, 2016.

Douglas, P. M. J., Stolper, D. A., Eiler, J. M., Sessions, A. L., Lawson, M., Shuai, Y., Bishop, A., Podlaha, O. G., Ferreira, A. A., and Neto, E. V. S.: Methane clumped isotopes: progress and potential for a new isotopic tracer, *Org Geochem*, 113, 262–282, 2017.

Douglas, P. M. J., Gonzalez Moguel, R., Walter Anthony, K. M., Wik, M., Crill, P. M., Dawson, K. S., Smith, D. A., Yanay, E., Lloyd, M. K., and Stolper, D. A.: Clumped isotopes link older carbon substrates with slower rates of methanogenesis in northern lakes, *Geophys Res Lett*, 47, e2019GL086756, 2020.

Douglas, P. M. J.; Stratigopoulos, E.; Park, J.; Phan, D.: [Data for geographic variability in freshwater methane hydrogen isotope ratios and its implications for emissions source apportionment and microbial biogeochemistry. figshare.](https://doi.org/10.6084/m9.figshare.13194833.v2)

Drevon, D., Fursa, S. R., and Malcolm, A. L.: Intercoder reliability and validity of WebPlotDigitizer in extracting graphed data, *Behavior modification*, 41, 323–339, 2017.

Dunn, O. J.: Multiple comparisons using rank sums, *Technometrics*, 6, 241–252, 1964.

Fisher, R. E., Sriskantharajah, S., Lowry, D., Lanoisellé, M., Fowler, C., James, R., Hermansen, O., Lund Myhre, C., Stohl, A., Greinert, J., and Nisbet, E. G.: Arctic methane sources: Isotopic evidence for atmospheric inputs, *Geophys Res Lett*, 38, L21803, 2011.

Fletcher, D., and Dixon, P. M.: Modelling data from different sites, times or studies: weighted vs. unweighted regression, *Methods in Ecology and Evolution*, 3, 168–176, 2012.

Ganesan, A., Stell, A., Gedney, N., Comyn-Platt, E., Hayman, G., Rigby, M., Poulter, B., and Hornibrook, E.: Spatially resolved isotopic source signatures of wetland methane emissions, *Geophys Res Lett*, 45, 3737–3745, 2018.

Gruen, D. S., Wang, D. T., Könneke, M., Topçuoğlu, B. D., Stewart, L. C., Goldhammer, T., Holden, J. F., Hinrichs, K.-U., and Ono, S.: Experimental investigation on the controls of clumped isotopologue and hydrogen isotope ratios in microbial methane, *Geochim Cosmochim Acta*, 237, 339–356, 2018.

Happell, J. D., Chanton, J. P., Whiting, G. J., and Showers, W. J.: Stable isotopes as tracers of methane dynamics in Everglades marshes with and without active populations of methane oxidizing bacteria, *Journal of Geophysical Research: Atmospheres*, 98, 14771–14782, 1993.

Happell, J. D., Chanton, J. P., and Showers, W. S.: The influence of methane oxidation on the stable isotopic composition of methane emitted from Florida swamp forests, *Geochim Cosmochim Acta*, 58, 4377–4388, 1994.

Hornibrook, E. R., Longstaffe, F. J., and Fyfe, W. S.: Spatial distribution of microbial methane production pathways in temperate zone wetland soils: stable carbon and hydrogen isotope evidence, *Geochim Cosmochim Acta*, 61, 745–753, 1997.

Peter Douglas 5/11/2021 12:42 PM

Deleted: Conrad, R., and Klose, M.: How specific is the inhibition by methyl fluoride of acetoclastic methanogenesis in anoxic rice field soil?, *FEMS Microbiology Ecology*, 30, 47–56, 1999. .

Peter Douglas 5/11/2021 12:42 PM

Deleted: Devol, A. H., Quay, P. D., Richey, J. E., and Martinelli, L. A.: The role of gas exchange in the inorganic carbon, oxygen, and ²²²Rn budgets of the Amazon River I, *Limnol Oceanogr*, 32, 235–248, 1987. .

Peter Douglas 5/17/2021 4:00 PM

Deleted:

Peter Douglas 5/17/2021 4:00 PM

Deleted:

Peter Douglas 5/17/2021 3:59 PM

Formatted: Font:10 pt

1078 Hornibrook, E. R.: The stable carbon isotope composition of methane produced and emitted from northern peatlands,
 1079 Baird, A., Belyea, L., Comas, X., Reeve, A., and Slater, L., American Geophysical Union, Geophysical Monograph
 1080 Series, 184, 187-203, 2009.
 1081 Huang, Y., Shuman, B., Wang, Y., and Webb III, T.: Hydrogen isotope ratios of palmitic acid in lacustrine sediments
 1082 record late Quaternary climate variations, *Geology*, 30, 1103-1106, 2002.
 1083 Jonsson, C. E., Leng, M. J., Rosqvist, G. C., Seibert, J., and Arrowsmith, C.: Stable oxygen and hydrogen isotopes in
 1084 sub-Arctic lake waters from northern Sweden, *J Hydrol*, 376, 143-151, 2009.
 1085 Kai, F. M., Tyler, S. C., Randerson, J. T., and Blake, D. R.: Reduced methane growth rate explained by decreased
 1086 Northern Hemisphere microbial sources, *Nature*, 476, 194-197, 2011.
 1087 Kruskal, W. H., and Wallis, W. A.: Use of ranks in one-criterion variance analysis, *Journal of the American statistical*
 1088 *Association*, 47, 583-621, 1952.
 1089 Krzycki, J., Kenealy, W., DeNiro, M., and Zeikus, J.: Stable carbon isotope fractionation by *Methanosarcina barkeri*
 1090 during methanogenesis from acetate, methanol, or carbon dioxide-hydrogen, *Applied and Environmental Microbiology*,
 1091 53, 2597-2599, 1987.
 1092 Lansdown, J., Quay, P., and King, S.: CH₄ production via CO₂ reduction in a temperate bog: A source of ¹³C-depleted
 1093 CH₄, *Geochim Cosmochim Acta*, 56, 3493-3503, 1992.
 1094 Lansdown, J. M.: The carbon and hydrogen stable isotope composition of methane released from natural wetlands and
 1095 ruminants, PhD Dissertation, University of Washington, 1992.
 1096 Lecher, A. L., Chuang, P. C., Singleton, M., and Paytan, A.: Sources of methane to an Arctic lake in Alaska: An isotopic
 1097 investigation, *Journal of Geophysical Research: Biogeosciences*, 122, 753-766, 2017.
 1098 Levin, I., Bergamaschi, P., Dörr, H., and Trapp, D.: Stable isotopic signature of methane from major sources in Germany,
 1099 *Chemosphere*, 26, 161-177, 1993.
 1100 Liptay, K., Chanton, J., Czepiel, P., and Mosher, B.: Use of stable isotopes to determine methane oxidation in landfill
 1101 cover soils, *Journal of Geophysical Research: Atmospheres*, 103, 8243-8250, 1998.
 1102 Mann, H. B., and Whitney, D. R.: On a test of whether one of two random variables is stochastically larger than the
 1103 other, *The annals of mathematical statistics*, 50-60, 1947.
 1104 Marik, T., Fischer, H., Conen, F., and Smith, K.: Seasonal variations in stable carbon and hydrogen isotope ratios in
 1105 methane from rice fields, *Global Biogeochem Cy*, 16, 41-41-41-11, 2002.
 1106 Martens, C. S., Kelley, C. A., Chanton, J. P., and Showers, W. J.: Carbon and hydrogen isotopic characterization of
 1107 methane from wetlands and lakes of the Yukon-Kuskokwim delta, western Alaska, *Journal of Geophysical Research:*
 1108 *Atmospheres* (1984–2012), 97, 16689-16701, 1992.
 1109 Miles, N. L., Martins, D. K., Richardson, S. J., Rella, C. W., Arata, C., Lauvaux, T., Davis, K. J., Barkley, Z. R.,
 1110 McKain, K., and Sweeney, C.: Calibration and field testing of cavity ring-down laser spectrometers measuring CH₄, CO
 1111 2, and $\delta^{13}\text{C}$ CH₄ deployed on towers in the Marcellus Shale region, *Atmospheric Measurement Techniques*, 11, 1273-
 1112 1295, 2018.
 1113 Mischler, J. A., Sowers, T. A., Alley, R. B., Battle, M., McConnell, J., Mitchell, L., Popp, T., Sofen, E., and Spencer, M.:
 1114 Carbon and hydrogen isotopic composition of methane over the last 1000 years, *Global Biogeochem Cy*, 23, 2009.
 1115 Mora, G., and Zanazzi, A.: Hydrogen isotope ratios of moss cellulose and source water in wetlands of Lake Superior,
 1116 United States reveal their potential for quantitative paleoclimatic reconstructions, *Chem Geol*, 468, 75-83, 2017.
 1117 Nakagawa, F., Yoshida, N., Sugimoto, A., Wada, E., Yoshioka, T., Ueda, S., and Vijarnsorn, P.: Stable isotope and
 1118 radiocarbon compositions of methane emitted from tropical rice paddies and swamps in Southern Thailand,
 1119 *Biogeochemistry*, 61, 1-19, 2002a.
 1120 Nakagawa, F., Yoshida, N., Nojiri, Y., and Makarov, V.: Production of methane from alasses in eastern Siberia:
 1121 Implications from its ¹⁴C and stable isotopic compositions, *Global Biogeochem Cy*, 16, 14-11-14-15, 2002b.
 1122 Nisbet, E., Dlugokencky, E., Manning, M., Lowry, D., Fisher, R., France, J., Michel, S., Miller, J., White, J., and
 1123 Vaughn, B.: Rising atmospheric methane: 2007–2014 growth and isotopic shift, *Global Biogeochem Cy*, 30, 1356-1370,
 1124 2016.
 1125 Nisbet, E. G., Manning, M., Dlugokencky, E., Fisher, R., Lowry, D., Michel, S., Myhre, C. L., Platt, S. M., Allen, G.,
 1126 and Bousquet, P.: Very strong atmospheric methane growth in the 4 years 2014–2017: Implications for the Paris
 1127 Agreement, *Global Biogeochem Cy*, 33, 318-342, 2019.

1128 Penger, J., Conrad, R., and Blaser, M.: Stable carbon isotope fractionation by methylophilic methanogenic archaea,
 1129 Applied and environmental microbiology, 78, 7596-7602, 2012.
 1130 Penning, H., Plugge, C. M., Galand, P. E., and Conrad, R.: Variation of carbon isotope fractionation in hydrogenotrophic
 1131 methanogenic microbial cultures and environmental samples at different energy status, Global Change Biol, 11, 2103-
 1132 2113, 2005.
 1133 Penning, H., and Conrad, R.: Quantification of carbon flow from stable isotope fractionation in rice field soils with
 1134 different organic matter content, Org Geochem, 38, 2058-2069, 2007.
 1135 Pine, M. J., and Barker, H.: Studies on the methane fermentation XII.: The pathway of hydrogen in the acetate
 1136 fermentation, Journal of Bacteriology, 71, 644-648, 1956.
 1137 Pison, I., Ringeval, B., Bousquet, P., Prigent, C., and Papa, F.: Stable atmospheric methane in the 2000s: key-role of
 1138 emissions from natural wetlands, 2013.
 1139 Popp, T. J., Chanton, J. P., Whiting, G. J., and Grant, N.: Methane stable isotope distribution at a Carex dominated fen in
 1140 north central Alberta, Global Biogeochem Cy, 13, 1063-1077, 1999.
 1141 Rice, A. L., Butenhoff, C. L., Teama, D. G., Röger, F. H., Khalil, M. A. K., and Rasmussen, R. A.: Atmospheric methane
 1142 isotopic record favors fossil sources flat in 1980s and 1990s with recent increase, Proceedings of the National Academy
 1143 of Sciences, 113, 10791-10796, 2016.
 1144 Rigby, M., Manning, A., and Prinn, R.: The value of high-frequency, high-precision methane isotopologue
 1145 measurements for source and sink estimation, Journal of Geophysical Research: Atmospheres, 117, 2012.
 1146 Röckmann, T., Eyer, S., Van Der Veen, C., Popp, M. E., Tuzson, B., Monteil, G., Houweling, S., Harris, E., Brunner, D.,
 1147 and Fischer, H.: In situ observations of the isotopic composition of methane at the Cabauw tall tower site, Atmos. Chem.
 1148 Phys., 16, 10469-10487, 2016.
 1149 Rozanski, K., Araguás-Araguás, L., and Gonfiantini, R.: Isotopic patterns in modern global precipitation, Geophysical
 1150 Monograph Series, 78, 1-36, 1993.
 1151 Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R.,
 1152 McInerney, F. A., and van der Meer, M. T. J.: Molecular paleohydrology: Interpreting the hydrogen-isotopic composition
 1153 of lipid biomarkers from photosynthesizing organisms, Annual Review of Earth and Planetary Sciences, 40, 221-249,
 1154 2012.
 1155 Sakagami, H., Takahashi, N., Hachikubo, A., Minami, H., Yamashita, S., Shoji, H., Khlystov, O., Kalmychikov, G.,
 1156 Grachev, M., and De Batist, M.: Molecular and isotopic composition of hydrate-bound and dissolved gases in the
 1157 southern basin of Lake Baikal, based on an improved headspace gas method, Geo-Mar Lett, 32, 465-472, 2012.
 1158 Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E.
 1159 J., Houweling, S., and Patra, P. K.: The global methane budget 2000–2017, Earth System Science Data, 12, 1561-1623,
 1160 2020.
 1161 Schaefer, H., Fletcher, S. E. M., Veidt, C., Lassey, K. R., Brailsford, G. W., Bromley, T. M., Dlugokencky, E. J., Michel,
 1162 S. E., Miller, J. B., and Levin, I.: A 21st-century shift from fossil-fuel to biogenic methane emissions indicated by
 1163 $\delta^{13}\text{CH}_4$, Science, 352, 80-84, 2016.
 1164 Schoell, M.: The hydrogen and carbon isotopic composition of methane from natural gases of various origins, Geochim
 1165 Cosmochim Acta, 44, 649-661, 1980.
 1166 Schoell, M.: Genetic characterization of natural gases, AAPG bulletin, 67, 2225-2238, 1983.
 1167 Schwietzke, S., Sherwood, O. A., Bruhwiler, L. M., Miller, J. B., Etiope, G., Dlugokencky, E. J., Michel, S. E., Arling,
 1168 V. A., Vaughn, B. H., and White, J. W.: Upward revision of global fossil fuel methane emissions based on isotope
 1169 database, Nature, 538, 88-91, 2016.
 1170 Shapiro, S. S., and Wilk, M. B.: An analysis of variance test for normality (complete samples), Biometrika, 52, 591-611,
 1171 1965.
 1172 Sherwood, O. A., Schwietzke, S., Arling, V. A., and Etiope, G.: Global inventory of gas geochemistry data from fossil
 1173 fuel, microbial and burning sources, version 2017, Earth System Science Data, 9, 2017.
 1174 Shoemaker, J., Varner, R., and Schrag, D.: Characterization of subsurface methane production and release over 3 years at
 1175 a New Hampshire wetland, Geochim Cosmochim Acta, 91, 120-139, 2012.
 1176 Sprenger, M., Tetzlaff, D., Tunaley, C., Dick, J., and Soulsby, C.: Evaporation fractionation in a peatland drainage
 1177 network affects stream water isotope composition, Water Resources Research, 53, 851-866, 2017.

1178 [Stell, A. C., Douglas, P.M.J., Rigby, M., and Ganesan, A.L.: The impact of spatially varying wetland source signatures](#)
1179 [on the atmospheric variability of \$\delta\text{D-CH}_4\$, Philosophical Transactions of the Royal Society A. Accepted article in press,](#)
1180 [2021.](#)

1181 Stolper, D., Martini, A., Clog, M., Douglas, P., Shusta, S., Valentine, D., Sessions, A., and Eiler, J.: Distinguishing and
1182 understanding thermogenic and biogenic sources of methane using multiply substituted isotopologues, *Geochim*
1183 *Cosmochim Acta*, 161, 219-247, 2015.

1184 Sugimoto, A., and Wada, E.: Hydrogen isotopic composition of bacterial methane: CO_2/H_2 reduction and acetate
1185 fermentation, *Geochim Cosmochim Acta*, 59, 1329-1337, 1995.

1186 Teasdale, C. J., Hall, J. A., Martin, J. P., and Manning, D. A.: Discriminating methane sources in ground gas emissions in
1187 NW England, *Quarterly Journal of Engineering Geology and Hydrogeology*, 52, 110-122, 2019.

1188 Thompson, H. A., White, J. R., Pratt, L. M., and Sauer, P. E.: Spatial variation in flux, $\delta^{13}\text{C}$ and $\delta^2\text{H}$ of methane in a
1189 small Arctic lake with fringing wetland in western Greenland, *Biogeochemistry*, 131, 17-33, 2016.

1190 Thompson, K. M., Burmaster, D. E., and Croucher, E. A.: Monte Carlo techniques for quantitative uncertainty analysis in
1191 public health risk assessments, *Risk Analysis*, 12, 53-63, 1992.

1192 Thompson, R. L., Nisbet, E., Pisso, I., Stohl, A., Blake, D., Dlugokencky, E., Helmig, D., and White, J.: Variability in
1193 atmospheric methane from fossil fuel and microbial sources over the last three decades, *Geophys Res Lett*, 45, 11,499-
1194 411,508, 2018.

1195 Turner, A. J., Frankenberg, C., and Kort, E. A.: Interpreting contemporary trends in atmospheric methane, *Proceedings of*
1196 *the National Academy of Sciences*, 116, 2805-2813, 2019.

1197 Tyler, J. J., Leng, M. J., and Arrowsmith, C.: Seasonality and the isotope hydrology of Lochnagar, a Scottish mountain
1198 lake: implications for palaeoclimate research, *The Holocene*, 17, 717-727, 2007.

1199 Tyler, S., Bilek, R., Sass, R., and Fisher, F.: Methane oxidation and pathways of production in a Texas paddy field
1200 deduced from measurements of flux, $\delta^{13}\text{C}$, and δD of CH_4 , *Global Biogeochem Cy*, 11, 323-348, 1997.

1201 Umezawa, T., Aoki, S., Kim, Y., Morimoto, S., and Nakazawa, T.: Carbon and hydrogen stable isotopic ratios of
1202 methane emitted from wetlands and wildfires in Alaska: Aircraft observations and bonfire experiments, *Journal of*
1203 *Geophysical Research: Atmospheres*, 116, 2011.

1204 Valentine, D., Sessions, A., Tyler, S., and Chidthaisong, A.: Hydrogen isotope fractionation during H_2/CO_2
1205 acetogenesis: hydrogen utilization efficiency and the origin of lipid-bound hydrogen, *Geobiology*, 2, 179-188, 2004a.

1206 Valentine, D. L., Chidthaisong, A., Rice, A., Reeburgh, W. S., and Tyler, S. C.: Carbon and hydrogen isotope
1207 fractionation by moderately thermophilic methanogens, *Geochim Cosmochim Acta*, 68, 1571-1590, 2004b.

1208 Vigano, I., Holzinger, R., Keppler, F., Greule, M., Brand, W. A., Geilmann, H., Van Weelden, H., and Röckmann, T.:
1209 Water drives the deuterium content of the methane emitted from plants, *Geochim Cosmochim Acta*, 74, 3865-3873, 2010.

1210 Wahlen, M.: Carbon dioxide, carbon monoxide and methane in the atmosphere: abundance and isotopic composition,
1211 *Stable isotopes in ecology and environmental science*, 93-113, 1994.

1212 Waldron, S., Watson-Craig, I. A., Hall, A. J., and Fallick, A. E.: The carbon and hydrogen stable isotope composition of
1213 bacteriogenic methane: a laboratory study using a landfill inoculum, *Geomicrobiology Journal*, 15, 157-169, 1998.

1214 Waldron, S., Lansdown, J., Scott, E., Fallick, A., and Hall, A.: The global influence of the hydrogen isotope composition
1215 of water on that of bacteriogenic methane from shallow freshwater environments, *Geochim Cosmochim Acta*, 63, 2237-
1216 2245, 1999a.

1217 Waldron, S., Hall, A. J., and Fallick, A. E.: Enigmatic stable isotope dynamics of deep peat methane, *Global Biogeochem*
1218 *Cy*, 13, 93-100, 1999b.

1219 Walter, K., Chanton, J. P., Chapin, F. S., Schuur, E. A. G., and Zimov, S. A.: Methane production and bubble emissions
1220 from arctic lakes: Isotopic implications for source pathways and ages, *J Geophys Res-Biogeophys*, 113, [2008](#).

1221 Walter, K. M., Zimov, S., Chanton, J. P., Verbyla, D., and Chapin, F. S.: Methane bubbling from Siberian thaw lakes as a
1222 positive feedback to climate warming, *Nature*, 443, 71-75, 2006.

1223 Wang, D. T., Gruen, D. S., Sherwood Lollar, B., Hinrichs, K.-U., Stewart, L. C., Holden, J. F., Hristov, A. N., Pohlman,
1224 J. W., Morrill, P. L., Könneke, M., Delwiche, K. B., Reeves, E. P., Sutcliffe, C. N., Ritter, D. J., Seewald, J. S.,
1225 McIntosh, J. C., Hemond, H. F., Kubo, M. D., Cardace, D., Hoehler, T. M., and Ono, S.: Nonequilibrium clumped
1226 isotope signals in microbial methane, *Science*, 348, 428-431, 2015.

1227 Wang, D. T., Welander, P. V., and Ono, S.: Fractionation of the methane isotopologues $^{13}\text{CH}_4$, $^{12}\text{CH}_3\text{D}$, and $^{13}\text{CH}_3\text{D}$
 1228 during aerobic oxidation of methane by *Methylococcus capsulatus* (Bath), *Geochim Cosmochim Acta*, 192, 186-202, 2016.
 1229 Wassmann, R., Thein, U., Whiticar, M., Rennenburg, H., Seiler, W., and Junk, W.: Methane emissions from the Amazon
 1230 floodplain: characterization of production and transport, *Global Biogeochem Cy*, 6, 3-13, 1992.
 1231 Whiticar, M., and Schaefer, H.: Constraining past global tropospheric methane budgets with carbon and hydrogen isotope
 1232 ratios in ice, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365,
 1233 1793-1828, 2007.
 1234 Whiticar, M. J., Faber, E., and Schoell, M.: Biogenic methane formation in marine and freshwater environments: CO_2
 1235 reduction vs. acetate fermentation—Isotope evidence, *Geochim Cosmochim Acta*, 50, 693-709, 1986.
 1236 Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem Geol*,
 1237 161, 291-314, 1999.
 1238 Woltemate, I., Whiticar, M., and Schoell, M.: Carbon and hydrogen isotopic composition of bacterial methane in a
 1239 shallow freshwater lake, *Limnol Oceanogr*, 29, 985-992, 1984.
 1240 Worden, J. R., Bloom, A. A., Pandey, S., Jiang, Z., Worden, H. M., Walker, T. W., Houweling, S., and Röckmann, T.:
 1241 Reduced biomass burning emissions reconcile conflicting estimates of the post-2006 atmospheric methane budget,
 1242 *Nature communications*, 8, 1-11, 2017.
 1243 Yacovitch, T. I., Daube, C., and Herndon, S. C.: Methane emissions from offshore oil and gas platforms in the Gulf of
 1244 Mexico, *Environmental Science & Technology*, 54, 3530-3538, 2020.
 1245 Young, E., Kohl, I., Sherwood Lollar, B., Etiope, G., Rumble, D., Li, S., Haghnegahdar, M., Schauble, E., McCain, K.,
 1246 Foustoukos, D., Sutcliffe, C., Warr, O., and Ballentine, C.: The relative abundances of resolved CH_2D_2 and $^{13}\text{CH}_3\text{D}$ and
 1247 mechanisms controlling isotopic bond ordering in abiotic and biotic methane gases, *Geochim Cosmochim Acta*, 203, 2017.
 1248 Yvon-Durocher, G., Allen, A. P., Bastviken, D., Conrad, R., Gudas, C., St-Pierre, A., Thanh-Duc, N., and Del Giorgio,
 1249 P. A.: Methane fluxes show consistent temperature dependence across microbial to ecosystem scales, *Nature*, 507, 488-
 1250 491, 2014.
 1251 Zazzeri, G., Lowry, D., Fisher, R., France, J., Lanoisellé, M., and Nisbet, E.: Plume mapping and isotopic
 1252 characterisation of anthropogenic methane sources, *Atmos Environ*, 110, 151-162, 2015.
 1253 Zhu, J., Liu, Z., Brady, E., Otto-Bliesner, B., Zhang, J., Noone, D., Tomas, R., Nusbaumer, J., Wong, T., and Jahn, A.:
 1254 Reduced ENSO variability at the LGM revealed by an isotope-enabled Earth system model, *Geophys Res Lett*, 44, 6984-
 1255 6992, 2017.
 1256 Zimov, S., Voropaev, Y. V., Semiletov, I., Davidov, S., Prosiannikov, S., Chapin, F. S., Chapin, M., Trumbore, S., and
 1257 Tyler, S.: North Siberian lakes: a methane source fueled by Pleistocene carbon, *Science*, 277, 800-802, 1997.
 1258