

Spatiotemporal patterns of N₂ fixation in coastal waters derived from rate measurements and remote sensing

5

Mindaugas Zilius^{1*}, Irma Vybernaite-Lubiene¹, Diana Vaiciute¹, Donata Overlingė¹, Evelina Griniene¹,
Anastasija Zaiko^{2,3}, Stefano Bonaglia^{1,4}, Iris Liskow⁵, Maren Voss⁵, Agneta Andersson⁶, Sonia Brugel⁶,
10 Tobia Politi¹, and Paul A. Bukaveckas^{7*}

¹Marine Research Institute, Klaipeda University, Klaipeda, 92294, Lithuania

²Coastal and Freshwater Group, Cawthron Institute, Nelson, 7042, New Zealand

15 ³Zealand Institute of Marine Science, University of Auckland, Auckland, Private Bag 92019, New Zealand

⁴Department of Marine Sciences, University of Gothenburg, Box 461, Gothenburg, 40530, Sweden

⁵Department of Biological Oceanography, Leibniz Institute for Baltic Sea Research, Rostock, 18119, Germany


⁶Department of Ecology and Environmental Sciences, Umeå University, Umeå, 90187, Sweden

⁷Center for Environmental Studies, Virginia Commonwealth University, Richmond, VA 23284, USA

20

25

30



40 **Figure S1:** Nitrogen fluxes associated with riverine input to the Curonian Lagoon (total nitrogen, grey bars) and N₂ fixation in water column based on remote sensing of Chlorophyll-a (green bars), and daily Nemunas River discharge (grey line) in 2018. N₂ fixation rates are mean and standard error (some error bars not visible). The Lithuanian Hydro-Meteorological Service provides discharge data at the gauging section.