

# Carbon dioxide fluxes and carbon balance of an agricultural grassland in southern Finland

Laura Heimsch<sup>1</sup>, Annalea Lohila<sup>1,2</sup>, Juha-Pekka Tuovinen<sup>1</sup>, Henriikka Vekuri<sup>1</sup>, Jussi Heinonsalo<sup>1,3</sup>, Olli Nevalainen<sup>1</sup>, Mika Korkiakoski<sup>1</sup>, Jari Liski<sup>1</sup>, Tuomas Laurila<sup>1</sup>, and Liisa Kulmala<sup>1,3</sup>

<sup>1</sup>Finnish Meteorological Institute, P.O. Box 503, 00101 Helsinki, Finland

<sup>2</sup>Institute for Atmospheric and Earth System Research, Physics, P.O. Box 64, 00014 University of Helsinki, Finland

<sup>3</sup>Institute for Atmospheric and Earth System Research, Forest Sciences, P.O. Box 27, 00014 University of Helsinki, Finland

**Correspondence:** Laura Heimsch (laura.heimsch@fmi.fi)

**Abstract.** A significant proportion of the global carbon emissions to the atmosphere originates from agriculture. Therefore, continuous long-term monitoring of CO<sub>2</sub> fluxes is essential to understand the carbon dynamics and balances of different agricultural sites. Here we present results from a new eddy covariance flux measurement site located in southern Finland. We measured CO<sub>2</sub> and H<sub>2</sub>O fluxes at this agricultural grassland site for two years, from May 2018 to May 2020. Especially the first summer experienced prolonged dry periods, which affected the CO<sub>2</sub> fluxes, and substantially larger fluxes were observed in the second summer. During the dry summer, leaf area index (LAI) was notably lower than in the second summer. Water use efficiency increased with LAI in a similar manner in both years, but photosynthetic capacity per leaf area was lower during the dry summer. The annual carbon balance was calculated based on the CO<sub>2</sub> fluxes and management measures, which included input of carbon as organic fertilisers and output as yield. The carbon balance of the field was ~~-50~~-57 ± ~~68~~10 g C m<sup>-2</sup> yr<sup>-1</sup> and ~~-118~~-86 ± ~~24~~12 g C m<sup>-2</sup> yr<sup>-1</sup> in the first and second study year, respectively. We estimated that on average the grassland ~~exceeded~~reached the global "4 per 1000" goal to increase the soil carbon content.

## 1 Introduction

Conventional and intensive agricultural practices cause significant carbon emissions while diminishing the soil organic matter (SOM) content. This leads to a reduction of soil quality and health (e.g. Houghton and Nassikas, 2017; Le Quéré et al., 2009, 2017; Lal, 2016; Paustian et al., 2000; Smith, 2008). Currently, agriculture is responsible for more than 10% of the global anthropogenic greenhouse gas (GHG) emissions to the atmosphere (Le Quéré et al., 2017). Soil type and properties, vegetation, climate and weather conditions as well as management practices all have a considerable effect on the carbon fluxes and balances of agroecosystems (Bolinder et al., 2010; Gomez-Casanovas et al., 2012; Jensen et al., 2017; Lorenz and Lal, 2018; Singh et al., 2018). Frequent ploughing, monocropping and intensive use of agrochemicals are the main contributors to the loss of SOM and the resulting carbon dioxide (CO<sub>2</sub>) emissions from land use (Ceschia et al., 2010; Reinsch et al., 2018; Yang et al., 2019). A change from conventional and intensive agricultural practices to regenerative and holistic farm management provides a substantial climate change mitigation potential (Lal, 2016). Increasing the amount of SOM in agroecosystems by applying enhanced management practices, such as lighter tillage, continuous plant cover, rotational grazing, agroforestry, increased

biodiversity and cover cropping, would not only help to mitigate climate change but also to restore soil quality and fertility. Especially, managed grasslands as part of agricultural systems have a high potential for substantial soil carbon sequestration (Soussana et al., 2010; Gilmanov et al., 2010; Yang et al., 2019). The importance of increasing soil organic carbon (SOC) content of agricultural soils has recently attained more attention, and the "4 per mille Soils for Food Security and Climate" initiative was launched at the 21<sup>st</sup> Conference of the Parties to the United Nations Framework Convention on Climate Change in Paris in 2015 (Minasny et al., 2017). The aim of this initiative is to increase the soil carbon stock on all land surfaces in the upper 2 metres by 0.4% annually. This would be enough to sequester carbon from the atmosphere by an amount equivalent to the annual anthropogenic GHG emissions. However, the initiative states that the most potential SOC increases can be achieved on managed agricultural lands. In that case, the "4 per 1000" means increasing of SOC at the top 1-m layer of agricultural soils by 0.4% annually. That would effectively offset approximately 20–35% of the global GHG emissions.

Agricultural ecosystems are highly prone to impacts of climate change, which induces a risk for food production. One of the possible impacts of climate change on agricultural ecosystems is associated with the changes in seasonal weather conditions and the resulting alteration in the carbon and water balance of these ecosystems (Ciais et al., 2014; Donnelly et al., 2017; Harrison et al., 2019). Severe drought events and storms causing considerable damage to agriculture have already been observed across Europe (Ciais et al., 2005; Wolf et al., 2013; Bastos et al., 2020). Moreover, adverse climatic impacts may be amplified by current and prior land use practices if they have not supported ecosystem resilience (Brunsell et al., 2014). For instance, a deeper root system is likely to buffer the negative impacts of climate variability. Also, high plant species diversity, compared to monocultures, favours the efficiency of plant water consumption and resilience to drought (De Boeck et al., 2006). As gross primary production (GPP) is closely related to ecosystem evapotranspiration (ET) via stomatal functions (Fricker and Willmer, 2012), changes in terrestrial water balance are potentially reflected in GPP and thus in the carbon balance of agricultural grasslands. The effect of water stress can be studied, for instance, by analysing ecosystem water use efficiency (WUE), i.e. the amount of carbon assimilated per unit of water lost by ET (Steduto, 1996). Generally, the productivity of a grassland ecosystem correlates with WUE, and thus ecosystems with a high productivity usually also have a high WUE (Hu et al., 2008). Environmental factors ~~are mainly regulating WUE via the effects on~~ regulate WUE via effects on stomatal conductance and GPP, and during prolonged drought periods, for example, temperature-induced downregulation of GPP may reduce WUE of grasslands in particular (Gharun et al., 2020). Furthermore, the WUE response depends on the intensity of the drought (Xu et al., 2019). However, the drought effects are also strongly related to season, as Wolf et al. (2013) reported that the WUE of Swiss grassland ecosystems did not respond to a spring drought and Bastos et al. (2020) concluded that the spring weather may either amplify or dampen the carbon and water dynamics during the following summer.

Better understanding of climatic impacts of agriculture and the effects of improved practices from the perspective of soil health and vitality is needed in order to develop tools for better environmental management of these ecosystems. Continuous long-term measurements of the atmosphere-ecosystem fluxes are needed to identify the key factors affecting carbon dynamics of different ecosystems, to quantify the resulting carbon balance and its components, and to verify soil carbon and ecosystem models. Moreover, ~~the~~ is-are high-quality GHG flux data ~~is-are~~ is-are needed for a reliable, global monitoring and verification system of agricultural carbon fluxes and soil carbon sequestration and stability (Smith et al., 2020).

The eddy covariance (EC) method is widely used for measuring CO<sub>2</sub> and energy fluxes in different ecosystems and climatic conditions (Aubinet et al., 2012). The high-frequency measurements provided by EC allow a direct quantification and analysis of gas exchange between the ecosystem and atmosphere. The carbon balance calculated from EC data, combined with the additional carbon fluxes caused by management, serves as an important measure for determining the climatic impact of agricultural ecosystems (e.g. Baldocchi, 2003; Baldocchi et al., 2018). However, continuous GHG flux measurements on agricultural sites, especially on mineral soils and grasslands, are still scarce in the Nordic countries (Shurpali et al., 2009; Lind et al., 2020; Jensen et al., 2017).

The aim of this study is to ~~determine the magnitude and seasonal dynamics of the carbon balance of~~ investigate, based on EC measurements, CO<sub>2</sub> exchange between the atmosphere and a managed forage grassland in southern Finland. In particular, we had three specific research questions:

1. ~~How does the CO<sub>2</sub> exchange and carbon balances vary between the study years~~ What is the magnitude of the annual carbon balance and its components?
2. Does the grass photosynthesis indicate occasional drought-related responses?
3. How does the possible carbon sink relate to the carbon sequestration objective of the “4 per 1000” initiative?


For the purposes of this study, we collected field data on the net exchange of CO<sub>2</sub> and H<sub>2</sub>O, soil and vegetation properties and meteorological variables on an agricultural grassland in southern Finland during two years, from May 2018 to May 2020.

## 2 Material and methods

### 2.1 Site description

The flux measurements were conducted at the Qvidja farm in southern Finland (60.29550°N, 22.39281°E; elevation 5 m) from May 2018 to May 2020 (Fig. 1). The site belongs to the hemiboreal climate zone. From 1981 to 2010, the mean annual air temperature and precipitation at the Kaarina Yltöinen weather station, located 13 km northeast of Qvidja, were 5.4°C and 679 mm, respectively (Pirinen et al., 2012). The experimental field in Qvidja has mineral soil (clay loam) and it covers 16.25 ha. It was cultivated as forage grassland during the study years. From 2008 to 2016, the field was managed intensively with conventional practices, and it was in annual crop rotation. In 2017, the field management practices were converted towards more sustainable and environmentally friendly farming by increasing the use of organic fertilisers and perennials, restricting the use of pesticides and increasing plant species biodiversity. The current grass and clover mixture was sown as an undergrown species with broad bean in spring 2017. The predominant ~~grass~~-species were timothy (*Phleum pratense*), meadow fescue (*Festuca pratensis*) and white clover (*Trifolium repens*).

Grass was harvested for silage for the first time on 12 June 2018. As the grass cover was fairly sparse later in the summer due to drought, repair seeding was done on 3 September 2018 to restore the drought-induced damage. The seed mixture included 35% of timothy, 30% of rye-grasses (*Lolium spp.*), 20% of common meadow-grass (*Poa pratensis*) and 15% of red fescue



**Figure 1.** Experimental field with the sectors representing the target area ~~the area that~~ covers 3.9 ha. Eddy covariance tower is located in the centre of the sectors. ~~Wind-EC data from wind~~ directions from 30 to 140° were ~~filtered out~~ ~~discarded~~ due to another experimental plot locating in that part of the field. (Orthophoto from National Land Survey of Finland)

90 (*Festuca rubra*). Timothy, meadow fescue and clover remained as the predominant species also in 2019 and early 2020. On 21 August 2018, the grass was cut at approximately 15 cm, but the yield was left in the field. The second harvest of 2018 occurred on 23 September. In 2019, the grass was harvested on 11 June and 20 August. In June 2018, a conventional cutting height of 6 cm was used, whereas in the other harvests the grass was cut at 15 cm.

In 2018, the field was fertilised twice, on 16 July and 24 August, with 2800 kg ha<sup>-1</sup> and 1800 kg ha<sup>-1</sup> of NK-molasses, 95 respectively (Table 1). NK-molasses was a byproduct of the sugar industry. It contained 67% of organic matter (OM) and 4.4% of nitrogen and had the C:N ratio of 9. According to the product information, the molasses included 205 g kg<sup>-1</sup> of organic carbon. In addition, it contained potassium and small proportions of sulphur, magnesium, calcium and sodium.

In May 2019, the field was fertilised with a mixture of side products from industries of starch potato processing, biowaste processing and ethanol production out of sawdust. This fertilisation mixture contained ~~65% of~~ 70% (of dry weight) of OM, 100 1.3% of nitrogen, 0.2% of phosphorus, 3% of potassium and 0.4% of sulphur, as well as small amounts of calcium, magnesium, zinc, copper and manganese. Approximately 4600 kg ha<sup>-1</sup> was applied on the field on 8 May (Table 1). On 26 June after the first harvest, 220 kg ha<sup>-1</sup> of mineral fertiliser was applied. This fertiliser contained 23% of nitrogen, 10% of phosphorus and 8% of potassium.



## 2.2 Measurement setup

105 The CO<sub>2</sub> and H<sub>2</sub>O fluxes were measured with the micrometeorological EC method. The flux measurements started on 3 May 2018, and here we analysed data collected from 4 May 2018 to 3 May 2020. From this point on, the periods of 4 May 2018 – 3 May 2019 and 4 May 2019 – 3 May 2020 are referred to as the first and second EC measurement year, respectively.

The EC instrumentation consisted of an enclosed infrared CO<sub>2</sub>/H<sub>2</sub>O gas analyser (LI-7200, LI-COR Biosciences, NE, USA), which detects the CO<sub>2</sub> and H<sub>2</sub>O mixing ratios, and a three-dimensional sonic anemometer (uSonic-3 Scientific, METEK  
110 GmbH, Elmshorn, Germany) to measure wind speed and air temperature. The data were recorded at 10-Hz frequency. The measurement height was 2.3 m. The flow rate was about 12 l min<sup>-1</sup>, and the length of the 4-mm stainless steel inlet tube with 2 µm Swagelok sinter was 0.8 m. ~~The gas analyser was calibrated with a zero. The~~ CO<sub>2</sub> ~~concentration air as a reference~~  
~~gas in May 2018 and March 2020. measurements were regularly checked with zero and span gases, and the LI-7200 was~~  
~~recalibrated when necessary. The H<sub>2</sub>O measurements were compared with the data obtained from a dedicated humidity sensor;~~  
115 ~~no recalibration was necessary.~~

The micrometeorological sign convention is used throughout the paper, with a negative value indicating the flux from the atmosphere to the ecosystem (net uptake) and a positive value indicating the flux from the ecosystem to the atmosphere (net emission).

Auxiliary meteorological measurements were conducted next to the flux tower. These included soil moisture observations  
120 at the depth of 0.1 m (ML3 ThetaProbe sensor, Delta-T Devices Ltd., Cambridge, UK) and soil temperature profile at the depths of 5, 10 and 30 cm (Pt100 IKES sensors, Nokeval Oy, Nokia, Finland). The soil temperature data were collected with a Vaisala QML201C datalogger (Vaisala Oyj, Vantaa, Finland). Photosynthetically active radiation (PQS PAR sensor, Kipp & Zonen B.V., Delft, The Netherlands), global and reflected solar radiation (CMP3 radiometer, Kipp & Zonen), and air temperature and relative humidity (Humicap HMP155, Vaisala Oyj) were measured at the height of 1.8 m. In addition, precipitation  
125 was measured with ~~Pluvio2~~ (a weighing rain gauge (Pluvio2, OTT HydroMet GmbH, Kempten, Germany)). Meteorological measurements started on 8 May 2018, and the data were recorded as 30-min averages, excluding the precipitation which was recorded as 1-min values. Snow ~~cover~~ depth was recorded at the weather station of Kaarina Yltöinen.

The leaf area index (LAI) data were obtained from the Sentinel-2 satellite as daily values on the clear-sky days. LAI was calculated from the Sentinel-2 bottom-of-atmosphere products (L2A) using the Google Earth Engine (GEE) and a Python  
130 implementation of the Biophysical Processor toolbox (Weiss and Baret, 2016) available in Sentinel Application Platform (SNAP) software. The cloudy, cloud-shadowed and snowy data were filtered out using the scene classification band available in the L2A products.


## 2.3 Eddy covariance data processing

The turbulent fluxes were determined as the covariance between the variations of vertical wind component and gas mixing ratio  
135 recorded at 10 Hz. They were calculated as 30-min block averages applying standard procedures, including double coordinate rotation and lag determination based on cross-correlation analysis (Rebmann et al., 2012). The systematic flux loss due to

the incomplete frequency response of the measurement system was corrected according to the empirical method described by Laurila et al. (2005).

The EC data from 5 January to 28 March 2019 were affected by technical issues with an inlet filter, which resulted in an erroneous reading of the internal analyser pressure. For this period, the 10-Hz mixing ratios were recalculated from the recorded absorbance data using the instrument-specific calibration functions. The mean CO<sub>2</sub> mixing ratio was set to 410 ppm in these calculations.

The following acceptance criteria were applied to screen the 30-min averaged CO<sub>2</sub> flux data: number of spikes in the raw data < 150 of 18,000, relative stationarity of CO<sub>2</sub> flux (Foken et al., 2012) < 50%, mean CO<sub>2</sub> mixing ratio > 380 ppm, variance of CO<sub>2</sub> mixing ratio < 15 ppm<sup>2</sup> between April and September and < 5 ppm<sup>2</sup> between October and March, and wind direction within 0–30° or 140–360°. Furthermore, the data were discarded during the periods of weak turbulence and when the flux footprint was not sufficiently representative of the target grassland, as estimated with the footprint model of Kormann and Meixner (2001). For these, we applied a friction velocity limit of 0.06 m s<sup>-1</sup> and a cumulative footprint limit of 0.7. The further screening applied to H<sub>2</sub>O fluxes included: H<sub>2</sub>O flux > 0, relative stationarity of H<sub>2</sub>O flux < 50% and variance of H<sub>2</sub>O mixing ratio < 1 (mmol mol<sup>-1</sup>)<sup>2</sup>. After applying these filtering criteria, the coverage of CO<sub>2</sub> and H<sub>2</sub>O flux data accepted for further analysis was 44% and 30% of all the 30-min periods during the two measurement years, respectively ([for CO<sub>2</sub>, day/night 55%/33%, April–September/October–March 48%/38%; for H<sub>2</sub>O, day/night 49%/11%, April–September/October–March 41%/16%](#)). Most of the accepted CO<sub>2</sub> and H<sub>2</sub>O flux data were collected when the wind direction was in the south-southwest sector (Fig. 2).



**Figure 2.** Number of accepted flux measurements within 20° sectors around the flux tower during the first and second year. Data from 30° to 140° were discarded.

## 2.4 Soil temperature model

The soil temperature sensor at the depth of 5 cm malfunctioned during the first measurement year, and these data were replaced with values derived from air temperature using the model presented by Rankinen et al. (2004). This model also takes into account the effect of possible snow cover on soil temperature. The following equation was used to obtain 30-min soil temperatures at 5 cm from 8 May 2018 to 3 May 2019:-

$$160 \quad T_z^{t+1} = T_z^t + \left[ \frac{\Delta t K_T}{C_A (2Z_s)^2} (T_{air}^t - T_z^t) \right] e^{-f_s D_s}$$

where  $T_z^{t+1}$  is the soil temperature at the depth of  $Z_s$  on the following day,  $T_z^t$  is the soil temperature of the current day,  $\Delta t$  is the length of the timestep,  $K_T$  is soil thermal conductivity, which was set to  $1 \text{ W m}^{-1} \text{ K}^{-1}$ ,  $C_A$  is the apparent heat capacity which is the sum of specific heat capacity of the soil  $C_s = 0.5 \times 10^{-6} \text{ J m}^{-3} \text{ K}^{-1}$  and specific heat capacity due to freezing and thawing  $C_{ice} = 4 \times 10^{-6} \text{ J m}^{-3} \text{ K}^{-1}$ , and  $T_{air}^t$  is the measured air temperature. The impact of snow cover was taken into account in the last term of the equation where  $f_s$  is an empirical snow parameter, which was set to  $10 \text{ m}^{-1}$ , and  $D_s$  is the measured snow depth. The model predictions were compared to measurements at the experimental field between June 2019 and May 2020. During summertime, the changes in soil temperature were fairly well captured by the model, whereas in the wintertime, the model tended to create larger changes in temperature than the actual measurements showed (Fig. ??)-

Comparison of measured and modelled soil temperature at 5 cm depth from June 2019 to May 2020.

## 170 2.4 Flux partitioning and gap-filling

To calculate  $\text{CO}_2$  balances and to ~~conduct further seasonal analysis of the flux components the~~ analyse the components of the net exchange between the field and the atmosphere, the measured  $\text{CO}_2$  flux data (i.e. net ecosystem exchange, NEE) were partitioned to GPP and total ecosystem respiration ( $R_{eco}$ ) and gap-filled based on this partitioning:

$$NEE = GPP + R_{eco} \quad (1)$$

175 The gap-filled GPP and  $R_{eco}$  were calculated with empirical response functions by first fitting these functions to the flux data.  $R_{eco}$  was expressed as a function of temperature (Lloyd and Taylor, 1994):

$$R_{eco} = R_0 e^{\frac{E_0 \left( \frac{1}{T_1} - \frac{1}{T_s - T_0} \right)}{}} e^{\frac{E_0 \left( \frac{1}{T_1} - \frac{1}{T_a - T_0} \right)}{}} \quad (2)$$

where  $R_0$  is the respiration rate ( $\text{mg m}^{-2} \text{ s}^{-1}$ ) at the reference ~~soil~~ temperature of 283.15 K,  $T_0 = 227.13 \text{ K}$ ,  $T_1 = 56.02 \text{ K}$ , ~~and  $E_0 = 308 \text{ K}$  is the long-term is~~ ecosystem sensitivity coefficient (Lloyd and Taylor, 1994) that describes the temperature response of soil respiration, and  $T_{sa}$  is the ~~soil temperature at the depth of 5 cm~~ air temperature.

GPP was modelled as a function of photosynthetically active radiation (PAR,  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) ~~and daily effective phytomass index (PI) as:~~ as:

$$GPP = \frac{PI \times \alpha \times PAR \times GP_{max}}{\alpha \times PAR + GP_{max}} \quad (3)$$

185 ~~where PI is an empirically-determined variable introduced to describe the seasonal changes in the photosynthetically active vegetation (Aurela et al., 2001), where~~  $\alpha$  is the apparent quantum yield ( $\text{mg } \mu\text{mol}^{-1}$ ), and  $GP_{max}$  denotes the asymptotic  $\text{CO}_2$  uptake rate in optimal light conditions ( $\text{mg m}^{-2} \text{s}^{-1}$ ). Further details on the ~~PI determination and~~ gap-filling procedure ~~are provided in the Appendix ?? and A, respectively~~ is provided in Appendix A. Energy fluxes were gap-filled following the description in ~~the~~ Appendix B.

190 To study the differences in photosynthetic capacity of the grass field between the two growing seasons, daily  $GP_{1200}$  values were calculated with the estimated  $\alpha$  and  $GP_{max}$  values, i.e. GPP was normalised to  $PAR = 1200 \mu\text{mol m}^{-2} \text{s}^{-1}$ .

## 2.5 ~~Carbon~~ Net ecosystem carbon balance

~~The~~ In this study, the system boundaries include the main components of the carbon balance of the ~~agricultural ecosystem field ecosystem studied. The carbon balance~~ was calculated by adding up the 30-min NEE fluxes, the imported carbon in the form  
195 of organic fertilisers and the ~~removal of carbon~~ carbon removed as harvested biomass:

$$C_{balance} = C_H + C_F + \sum_{i=1}^{mn} NEE_i \quad (4)$$

where  $C_H$  is the amount of carbon in harvested biomass,  $C_F$  is the amount of carbon in imported fertilisation and ~~mn~~ is the total number of timesteps in the period for which the balance was calculated. Thus, the carbon balance indicates the net ecosystem carbon balance as defined by Chapin et al. (2006) without the contribution of carbon monoxide, methane, volatile organic and  
200 particulate compounds or leaching. This balance is commonly called the net biome production (Kutsch et al., 2010). Biomass was converted to carbon by multiplying the dry weight by 0.42 (Lohila et al., 2004). The following sign convention was used: the carbon imported into the ecosystem corresponds to a negative flux and the carbon removed from the system corresponds to a positive flux.

## 2.6 Uncertainty analysis

205 The  $\text{CO}_2$  balance between the field and the atmosphere, which is calculated based on the EC measurements, includes multiple potential error sources. Uncertainties are associated, for example, with the stochastic nature of turbulence and incomplete sampling of large eddies, the performance of instruments and the flux variation caused by the limited area of the target ecosystem

(Aubinet et al., 2012). Some of these errors were compensated for in the data processing and screening .~~Here we included in the uncertainty estimate the~~ and the most relevant random error sources, i.e. the statistical measurement error ( $E_{meas}$ ) and the error caused by gap-filling ( $E_{gap}$ ) ~~Aurela et al. (2002)~~ (Aurela et al., 2002), were included in the uncertainty estimate:

$$E_{meas} = \sqrt{\sum_{i=1}^n (NEE_{meas,i} - NEE_{mod,i})^2} \quad (5)$$

where  $NEE_{meas}$  is the filtered 30-min flux,  $NEE_{mod}$  is the corresponding modelled NEE (Eqs. 1–3), and  $n$  is the number of measured data.

$$E_{gap} = \sqrt{\sum_{i=1}^N (E_{GPP,i}^2 + E_{Reco,i}^2)} \quad (6)$$

where  $E_{GPP}$  and  $E_{Reco}$  are the errors of modelled GPP and  $R_{eco}$ , respectively.  $N$  is the number of gaps in the data.

The standard error propagation principle was used in estimating the total uncertainty ( $E_{tot}$ ) of the annual carbon balance:

$$E_{tot} = \sqrt{E_{meas}^2 + E_{gap}^2} \quad (7)$$

## 2.7 Water use efficiency

The ecosystem WUE was defined as the ratio of GPP to ET, i.e. H<sub>2</sub>O flux:

$$WUE = \frac{GPP}{ET} \quad (8)$$

where daily means of GPP and ET were used. The ET data corresponding to a latent heat flux lower than 30 W m<sup>-2</sup> were discarded (Abraha et al., 2016). Furthermore, days with precipitation were eliminated.

## 2.8 Soil carbon content

Soil carbon content was determined from 1-m-deep core samples taken within the flux source area. The samples were taken in October 2018 using a hydraulic corer installed to a tractor ~~in October 2018.~~ The diameter of the sample cylinder was 151 mm.



Subsamples were taken along the 1-m core at 16 points, and soil organic carbon (SOC,  $\text{kg m}^{-2}$ ) content in each subsample  
230 was analysed using a VarioMax CN analyser (Elementar Analysensysteme GmbH, Germany).

### 3 Results

#### 3.1 Meteorological conditions

The annual mean air temperature at the study site was  $7.6^{\circ}\text{C}$  and  $7.7^{\circ}\text{C}$  in the first and second measurement year, respectively. Both years were warm compared to the long-term (1981–2010) average of  $5.4^{\circ}\text{C}$  measured at a nearby weather station (Pirinen  
235 et al., 2012). The annual precipitation sum was lower in the first year (473 mm) and higher in the second year (855 mm) than the long-time average (679 mm).

The thermal growing season, defined here as the period when the daily mean temperature exceeded permanently  $5^{\circ}\text{C}$ , started on 14 April in 2018, i.e. before the EC measurements started. In 2019 and 2020, the thermal growing season began on 16 April and 18 April, respectively. The thermal growing season ended on 17 November and 26 October in 2018 and 2019, respectively.  
240 Thus, the thermal growing season length was 218 days in 2018 and 194 days in 2019. Meteorological conditions during the main growing season between May and September varied substantially between the two years. The mean air temperature during these months was  $16.7^{\circ}\text{C}$  and  $14.5^{\circ}\text{C}$  in 2018 and 2019, respectively. ~~The mean growing season soil temperatures were similar to the air temperatures with  $16.4^{\circ}\text{C}$  (modelled) in 2018 and  $14.5^{\circ}\text{C}$  in 2019.~~ During the same period, the mean daily PAR was about 12% higher in 2018 than in 2019 ( $460$  vs.  $410 \mu\text{mol m}^{-2} \text{s}^{-1}$ ), while the precipitation sum was 32%  
245 lower (212 vs. 312 mm).


During winter 2018–2019, permanent snow cover was recorded from 17 December [2018](#) to 26 March 2019. In the following winter (2019–2020), there were only two short snow-cover periods: 5–8 February and 30–31 March 2020. The maximum snow depth in the first winter was 33 cm, whereas in the second winter it was 3 cm. The mean wintertime (November–March) air temperature was  $-0.2^{\circ}\text{C}$  in 2018–2019 and  $2.2^{\circ}\text{C}$  in 2019–2020. ~~The warmer winter in the second measurement year was also  
250 observed in the mean soil temperature ( $-0.9$  vs.  $1.6^{\circ}\text{C}$ ).~~

Soil moisture content at the depth of 10 cm varied between  $0.16$  and  $0.55 \text{ m}^3 \text{ m}^{-3}$  during the study period. In several occasions, the daily mean soil moisture dropped to about  $0.2 \text{ m}^3 \text{ m}^{-3}$ . During the growing seasons, such low values indicate substantial drought, while in the winter, rapid data drops were likely related to soil freezing. The average soil moisture during the growing season in 2019 was higher than in 2018 ( $0.30$  vs.  $0.26 \text{ m}^3 \text{ m}^{-3}$ ). As a result of the higher precipitation in 2019,  
255 soil moisture occasionally increased up to  $0.4 \text{ m}^3 \text{ m}^{-3}$ , i.e. close to the saturated values observed in winter.

#### 3.2 Fluxes

At the beginning of the measurements, the net  $\text{CO}_2$  fluxes were negative (Fig. 3), and the air ~~and soil temperatures were~~  
[temperature was](#) already well above  $10^{\circ}\text{C}$  (Fig. 4). Net uptake was observed until the first harvest around mid-June 2018. This harvest and the following management events during ~~the~~ that growing season induced large short-term variations in the

260 CO<sub>2</sub> fluxes. Similarly, in the second study year, large impacts on CO<sub>2</sub> fluxes were observed after the management events. During the growing season, the mean NEE was  $-0.13$  and  $-0.21$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup> in 2018 and 2019, respectively. During the wintertime, no significant CO<sub>2</sub> uptake occurred, and the positive fluxes were small compared to the nocturnal fluxes in summer. The mean measured NEE between December 2018 and February 2019 was  $0.03$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, and during the same period in 2019–2020 it was  $0.04$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>.



**Figure 3.** Accepted 30-min a) net ecosystem exchange (NEE) and b) H<sub>2</sub>O flux measurements from May 2018 to May 2020. Vertical lines with H and F indicate harvest and fertilisation, respectively.

265 Seasonal patterns were observed also in the H<sub>2</sub>O fluxes (Fig. 3). In the spring, the ecosystem ET started to increase reaching the highest levels between June and August, after which it gradually decreased to wintertime values, i.e. close to zero. The mean growing season H<sub>2</sub>O flux was  $34.7$  mg H<sub>2</sub>O m<sup>-2</sup> s<sup>-1</sup> in 2018 and  $35.5$  mg H<sub>2</sub>O m<sup>-2</sup> s<sup>-1</sup> in 2019. The wintertime (December–February) mean H<sub>2</sub>O flux was  $3.6$  mg H<sub>2</sub>O m<sup>-2</sup> and  $3.7$  mg H<sub>2</sub>O m<sup>-2</sup> in 2018–2019 and 2019–2020, respectively.

The experimental field was harvested and fertilised twice during each of the studied growing seasons (Table 1). The effect  
 270 of management was investigated by comparing the mean fluxes 5-five days before and after the harvest dates (Table A1). The harvest in June 2018 changed the mean CO<sub>2</sub> flux from a net sink of  $-0.28$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup> to a source of  $0.03$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, i.e. increased the net efflux by  $0.31$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>. The first harvest of 2019 increased NEE by  $0.47$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, but as the pre-harvest mean NEE was  $-0.50$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, the field remained as a net sink. As a result of the second harvest on 23 September 2018, the mean sink reduced from  $-0.10$  to  $-0.02$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, while the harvest on 20 August  
 275 2019 caused the sink to change from  $-0.25$  to  $-0.02$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>. Thus, after all the harvests with a cutting height of 15 cm, the mean sink rate was diminished to  $-0.02$  or  $-0.03$  mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>.

In the first growing season, the first and second fertilisation events with organic substances increased NEE by 0.27 and 0.08 mg CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>, respectively, i.e. diminished the CO<sub>2</sub> sink (Fig. 3, Table A1). During the ~~5~~five days after the harvest in May 2019, the field acted as a CO<sub>2</sub> source. A similar trend was not observed in June 2019, as mineral fertiliser was used and thus  
 280 no organic substances were added to the soil. Each of the fertilisation events were followed by rain within the next ~~5~~five days. However, the mean soil moisture ~~remained either at the depth of 10 cm either~~remained the same or decreased slightly (Fig. 4, Table A1). Furthermore, the mean air temperature increased after the fertilisations in July 2018 and May 2019, potentially affecting CO<sub>2</sub> fluxes. After the fertilisation events with organic substances in July 2018, August 2018 and May 2019, the mean PAR was 7%, 29% and 12% lower, respectively, than the 5-day mean before the fertilisation, complicating the interpretation of  
 285 fertilisation impacts on the CO<sub>2</sub> fluxes. The effect of management on H<sub>2</sub>O fluxes could not be disentangled from the present data (Fig. 3b).

~~The PI calculated from the flux data was consistent with the seasonal changes in the~~The LAI derived from Sentinel-2 images (Fig. 4d) ~~varied greatly between the years~~. The higher LAI in 2019 indicated that there was more photosynthesising green biomass before the first and second harvest compared to 2018. The effect of larger leaf area was also observed in the  
 290 differences in the photosynthetic capacity (GP<sub>1200</sub>) of the grassland between the study years (Fig. 5a). The years differed significantly (p < 0.05) in terms of GP<sub>1200</sub> at all levels of LAI (>1). Larger LAI values were observed throughout 2019, indicating that grass was growing better than in 2018. Furthermore, the grassland was photosynthesising more efficiently with the same leaf area in 2019 than in the previous year (Fig. 5a).


### 3.3 Water use efficiency

295 The ecosystem WUE estimate showed different seasonal variation during the studied growing seasons (Fig. 6). Generally, WUE was higher in 2019 than in 2018 throughout the growing season. WUE increased before the first harvest around mid-June in both years, indicating more efficient CO<sub>2</sub> uptake in terms of water use than during the spring. The 5-day mean WUE before the first harvest was ~~2.6 and 2.9~~2.8 and 3.0 g CO<sub>2</sub> (kg H<sub>2</sub>O)<sup>-1</sup> in 2018 and 2019, respectively. Due to the harvest, it dropped to ~~0.8-0.9~~ g CO<sub>2</sub> (kg H<sub>2</sub>O)<sup>-1</sup> in 2018 and to ~~2.2-2.6~~ g CO<sub>2</sub> (kg H<sub>2</sub>O)<sup>-1</sup> in 2019. During the latter growing season,  
 300 WUE increased steadily towards 4 g CO<sub>2</sub> (kg H<sub>2</sub>O)<sup>-1</sup> until the second harvest in August, whereas ~~WUE in 2018 it~~ remained predominantly below 2 g CO<sub>2</sub> (kg H<sub>2</sub>O)<sup>-1</sup> during the same period~~in 2018~~. In the end of August and early September, WUE was at the same level in both years.


The LAI derived from the Sentinel-2 data was compared to the daily WUE values (Fig. 5b) to further ~~interpret the relation~~cast light on the relationship between vegetation status and ~~ecosystem~~-WUE. While WUE was on average lower in 2018 than  
 305 2019, the difference at a given LAI was not significant (p > 0.05). However, in both years the daily WUE increased in a similarly linear manner in relation to LAI.

### 3.4 Carbon balance and soil carbon content


The carbon balance of the studied grass field was ~~-50-57 ± 68-10~~ g C m<sup>-2</sup> yr<sup>-1</sup> ~~, i.e. not different from zero,~~ in the first year, ~~while and~~ the balance of the second year was ~~negative, -118-86 ± 24-12~~ g C m<sup>-2</sup> yr<sup>-1</sup>, i.e. the field acted as a net carbon



**Figure 4.** Daily mean a) air and soil (depth = 0.05 m) temperature, b) photosynthetically active radiation (PAR), c) precipitation and soil moisture (depth = 0.1 m), d) ~~phytomass (PI)~~ and ~~leaf area indices index~~ (LAI), and e) daily mean NEE, GPP,  $R_{eco}$  and cumulative carbon flux from May 2018 to May 2020.



**Figure 5.** a) Daily photosynthetic capacity ( $GP_{1200}$ ) and b) water use efficiency (WUE) as a function of leaf area index (LAI) during the two growing seasons. Grey areas represent the uncertainty estimation bands.



**Figure 6.** Daily water use efficiency (WUE) during two growing seasons.



**Table 1.** Different management events and their C inputs (fertilisation) and C outputs (harvest). During the cutting in August 2018, the grass was not collected and thus did not result ~~to~~in any C flux allocated to management.

Date	Management	Output (dry weight kg ha <sup>-1</sup> )	Input (kg ha <sup>-1</sup> )	Carbon (g m <sup>-2</sup> )
12 Jun 2018	Harvest	1985		83
16 Jul 2018	Fertilisation		-2800	-57
21 Aug 2018	Cutting	-	-	-
24 Aug 2018	Fertilisation		-1755	-36
23 Sep 2018	Harvest	348		15
8 May 2019	Fertilisation		-4606	<del>-93</del> <u>-43</u>
11 Jun 2019	Harvest	3107		130
20 Jun 2019	Fertilisation (mineral)	-	-	-
20 Aug 2019	Harvest	1029		43

**Table 2.** ~~Annual~~The annual carbon balances and their components (g C m<sup>-2</sup> yr<sup>-1</sup>) for the two measurement years. Negative values indicate C input into the ecosystem, whereas positive values indicate C loss. Management (M) is the sum of the C fluxes due to harvest (positive) and fertilisation (negative) events (Table 1). The values after  $\pm$  represent the uncertainty in NEE.

	NEE	GPP	R <sub>eco</sub>	M	Total balance
First year	<del>-55</del> <u>-62</u>	<del>-1034</del> <u>-1121</u>	<del>972</del> <u>1053</u>	5	<del>-50</del> <u>-57</u> $\pm$ <del>68</del> <u>10</u>
Second year	<del>-198</del> <u>-216</u>	<del>-1480</del> <u>-1583</u>	<del>1291</del> <u>1362</u>	<del>80</del> <u>130</u>	<del>-118</del> <u>-86</u> $\pm$ <del>24</del> <u>12</u>


310 sink in both years (Table 2). ~~All the~~The magnitude of all components of the carbon balance were smaller in the first year than in the second one, GPP by ~~30~~29%, R<sub>eco</sub> by ~~25~~23% and management by ~~94~~-%.

~~There was a major difference in the~~ 96%. The components in the mean annual CO<sub>2</sub> ~~balancees between the~~ fluxes between the field and the atmosphere indicated major differences also between the growing seasons (Table 3). In 2019, the ~~growing season net uptake magnitude of the growing season NEE~~ was 78%, GPP 49% and R<sub>eco</sub> 42% higher than in 2018.

315 The average soil carbon content in the 1-m layer was 16.59  $\pm$  2.25 kg m<sup>-2</sup> (average  $\pm$  standard deviation), with the highest SOC found in the top 30-cm layer (Fig. 7). The carbon balance of 2018 was 0.3% of the average SOC, and in 2019 this ratio was ~~0.7~~0.5%. On average, the annual carbon input to the soil accounted for ~~0.5~~0.4% of the SOC.

**Table 3.** ~~Growing season (from 4 May to 30 September)~~Net ecosystem exchange of CO<sub>2</sub> ~~balancees (NEE, g CO<sub>2</sub> m<sup>-2</sup> m<sup>-2</sup>), its components gross primary production (GPP) and total ET ecosystem respiration (R<sub>eco</sub>), and evapotranspiration (ET, mm) during the growing season (4 May to 30 September), in 2018 and 2019.~~

<u>Year</u>	NEE	GPP	R <sub>eco</sub>	ET
2018	<del>-650</del> <u>-601</u>	<del>-3190</del> <u>-3330</u>	<del>2510</del> <u>2715</u>	297
2019	<del>-1160</del> <u>-1176</u>	<del>-4740</del> <u>-4955</u>	<del>3560</del> <u>3771</u>	283



**Figure 7.** a) Soil organic carbon (SOC) content at different depths in the 1-m deep soil samples, and b) the total SOC in the samples. Numbers from 1 to 5 indicate sample numbers.

## 4 Discussion

### 4.1 Fluxes and carbon balance

320 There is an urgent need to find evidence-based climate-friendly practices in agriculture also in the boreal region, where the growing season is short and varieties differ from those cultivated in the temperate region. The carbon fluxes ~~on an~~ we measured on the agricultural grassland at the Qvidja farm in southern Finland ~~were clearly different between the two study years~~ clearly indicated that this site was a sink of atmospheric carbon. The annual NEE was ~~-55~~ -62 g C m<sup>-2</sup> yr<sup>-1</sup> in the first study year (4 May 2018 – 3 May 2019) and ~~-198~~ -216 g C m<sup>-2</sup> yr<sup>-1</sup> in the second year (4 May 2019 – 3 May 2020). The GPP showed

325 notable variation between the study years as the annual GPP was ~~-1034 and -1480~~ -1121 and -1583 g C m<sup>-2</sup> yr<sup>-1</sup> in the first and second year, respectively. Gilmanov et al. (2010) ~~have reported the GPP of European managed grasslands to be in the reported a~~ range of -2107 to -1410 g C m<sup>-2</sup> yr<sup>-1</sup> for the GPP of European managed grasslands. Our results fall ~~in to the lower range or below these GPP values~~ below or in the lower end of this range. The annual R<sub>eco</sub> in Qvidja ~~was also varying also varied~~ between the study years (~~972 and 1291~~ 1053 and 1362 g C m<sup>-2</sup> yr<sup>-1</sup>). Globally, the annual R<sub>eco</sub> of managed grasslands

330 is reported to vary within a wide range from 31 to 2150 g C m<sup>-2</sup> yr<sup>-1</sup>. The average R<sub>eco</sub> was 1445 and 647 g C m<sup>-2</sup> yr<sup>-1</sup> on the intensively and extensively managed grasslands, respectively (Gilmanov et al., 2010), and the ~~The~~ annual R<sub>eco</sub> in Qvidja ~~falls between these values in both study years. Regarding only the European grasslands, the annual R<sub>eco</sub> is reported to vary between 494 and 1623 g C m<sup>-2</sup> yr<sup>-1</sup> (Gilmanov et al., 2007).~~ Our observations are thus also within this range.

The carbon balance of the grass field ecosystem in Qvidja was close to neutral (-50 ~~To answer our first research question, we~~ concluded that the carbon balance was negative in both study years (-57 ± 68 ~~10~~ g C m<sup>-2</sup> yr<sup>-1</sup> ) in the first study year (4 May 2018 – 3 May 2019), and in the second year (4 May 2019 – 3 May 2020) the field was a moderate carbon sink (-118 and -86 ~~± 24~~ 12 g C m<sup>-2</sup> yr<sup>-1</sup>). ~~, and thus the field acted as a net carbon sink during the study period.~~ Carbon balances, including the

carbon equivalent of  $\text{N}_2\text{O}$ ,  $\text{CH}_4$  and management-related carbon fluxes, ~~have been widely studied in other~~ were studied in nine European agricultural grassland sites ~~between 2002–2004 (Soussana et al., 2007). These nine grassland~~ (Soussana et al., 2007). All sites acted mainly as net carbon sinks in 2002–2004, the annual net carbon balance ranging from  $-446$  to  $251 \text{ g CO}_2\text{-C eq. m}^{-2} \text{ yr}^{-1}$ , where 13 of the 17 measured annual balances were negative. Our ~~study-site carbon balance, which excludes  $\text{N}_2\text{O}$  and  $\text{CH}_4$  fluxes,~~ falls into this range. In comparison, the Finnish agricultural ~~sites-fields~~ measured so far ~~are-were~~ generally carbon sources when ecosystem-atmosphere  $\text{CO}_2$  fluxes, harvests and the carbon supplied to the system as fertilisers were considered (Heikkinen et al., 2013; Shurpali et al., 2009; Lind et al., 2016; Lohila et al., 2004). Lind et al. (2016) reported ~~slightly higher annual net uptake of atmospheric  $\text{CO}_2$ -a slightly more negative annual NEE~~ (two-year average NEE  $-259 \text{ g C m}^{-2} \text{ yr}^{-1}$ ) for a grassland site on mineral soil than we observed in Qvidja. However, by considering the total carbon balance of the system by taking into account the carbon fluxes ~~caused-by-management~~ related to biomass removal as grass yield, it was concluded that their site acted as a net carbon source. Mineral fertilisers were used during their study, and thereby no carbon was imported to the field to compensate for the biomass removal from the system as harvests. Similar management-related carbon flux patterns ~~related to management were reported by Eichelmann et al. (2016). The annual NEE of the agricultural grassland in their study in Canada was more negative (average NEE were observed by Eichelmann et al. (2016), who reported a more negative NEE (average  $-405 \text{ g C m}^{-2} \text{ yr}^{-1}$ ) for an agricultural grassland in Canada~~ than the NEE in Qvidja. ~~However;~~ however, the two-year mean annual carbon balance ~~of the Canadian field~~ was positive when biomass removal was taken into account, i.e. the Canadian field was a net source of carbon. It is noteworthy that the yield in Qvidja was substantially smaller than at the other two study sites (Lind et al., 2016; Eichelmann et al., 2016), at which the total balance became positive when the management, i.e. harvests and fertilisation, was taken into account. ~~However, as the harvested grass was used as feed for farm animals, there was no need for a higher yield at Qvidja in either of the study years~~ The total carbon balance of the field depends greatly both on the amount of organic matter imported to the system as fertilisers and on the harvest yields, which are affected, for instance, by the applied cutting height.

Analysis of the weather variables in Qvidja indicated that temperature and moisture conditions were associated with the differences in ~~fluxes~~  $\text{CO}_2$  flux dynamics and carbon balance between the study years. The growing season was warmer and drier in 2018 than 2019, with 13% lower mean soil moisture, 32% lower precipitation,  $2.2^\circ\text{C}$  higher mean air temperature and 12% higher mean radiation during the growing season, and substantially smaller fluxes were observed in the first year. This is in accordance with Shurpali et al. (2009) who observed a positive correlation between the uptake of atmospheric  $\text{CO}_2$  (GPP) and both soil moisture and air temperature on another Finnish agricultural grassland. According to their conclusions, moderate temperature with high soil moisture ~~favoured~~ enhanced  $\text{CO}_2$  uptake. Furthermore, Flanagan et al. (2002) and Kurc and Small (2007) concluded that ~~photosynthesis of grassland favours~~ rather wet summer conditions ~~-favoured photosynthetic activity in grasslands~~. These findings would support the conclusion that low soil moisture and high temperatures were the main factors limiting  $\text{CO}_2$  uptake at our study site in the summer 2018.

~~To answer our first research question, we conclude that there were notable year-to-year differences in the carbon balances, but the reason behind this variation~~ However, this question remains partly open, as weather conditions, grass age and grass leaf area all showed different dynamics between the study years. In Finland, it is typical to grow grasslands for 3–4 years before ~~the~~

grass renewal. In Qvidja, the grass was not renewed between the study years, which may have led to the larger fluxes observed in the second year when the grass root system, for instance, was ~~possibly more developed~~enhancing carbon uptake likely to be more developed, enhancing water and nutrient availability and thus reducing the effect of drought stress. Furthermore, the leaf area was larger, and other capabilities, such as microbial symbioses (e.g. de Vries et al., 2020; Harman and Uphoff, 2019; Moreau et al., 2019), of the more developed grass may have increased carbon uptake. The lower leaf area during the first year was most probably also ~~connected due~~ to the dry summer, as shortage of water is a growth-limiting factor. Besides the leaf area, the photosynthetic potential per leaf area was lower in the first year, indicating either drought stress or shortage of nutrients, as temperature, a widely limiting factor in northern latitudes, was high enough during both summers not to restrict photosynthesis. In any case, a more specific analysis of the driving and inhibiting environmental factors will require a longer measurement period.

Our second research question concerned the drought-related restrictions of photosynthesis. It has been widely recognised that in dry conditions plants are able to reduce transpiration by stomatal regulation (~~Pirasteh-Anosheh et al., 2016~~)(Willmer and Fricker, 1996). However, grasses seem to limit stomatal functions only in severe, prolonged drought conditions (Wolf et al., 2013; Xu et al., 2019), and thus occasional or seasonal drought events may not be observed in the ecosystem WUE of grasslands. In our study, WUE values were predominantly lower in 2018 than in 2019. This was most probably explained by the differences in LAI, as the relationship between WUE and LAI was similar ~~in during~~ both growing seasons (Fig. 5b). Furthermore, the drier conditions with high temperatures in the summer 2018 may have resulted in a decoupling of assimilation and transpiration and in temperature-induced downregulation of GPP (Gharun et al., 2020), as ET was similar in both years (Table 3). Therefore, the clearly lower leaf-area-based photosynthetic capacity ( $GP_{1200}$ ) in 2018 compared to 2019 probably indicates ~~drought-related stress in drought-related stress on~~ photosynthetic processes despite the similar leaf-area-based WUE (Fig. 5). It is noteworthy that the WUE analysis was performed by means of the total ecosystem ET rather than plant transpiration, which would have enabled a more direct determination of the actual plant WUE and thus a simpler interpretation of plant processes and their relation to LAI. Nevertheless, days with even slight precipitation were eliminated from the analysis, and therefore we can assume that during the growing season most of the water flux arises from transpiration. In general, WUE at our study site varied mainly between 0 and ~~5.4~~ 4 g C (kg H<sub>2</sub>O)<sup>-1</sup>. This is consistent with the WUEs observed ~~for on~~ northern grasslands (0–7 g C (kg H<sub>2</sub>O)<sup>-1</sup>) (Tang et al., 2014).

The different management practices, such as fertilisation and the choice of grass cutting height, were slightly different in the first and second year, which probably had an impact on the carbon balances. In June 2018, a conventional cutting height of 6 cm was used, whereas in the other harvests the grass was cut at 15 cm. The higher cutting height may have enhanced the regrowth of ~~the grasses~~grass, especially in the more favourable weather conditions in 2019, and with a larger leaf area higher CO<sub>2</sub> uptake was observed right after the harvest. Only after the 6-cm harvest, the field turned to a net source of CO<sub>2</sub>. With a low cutting height, it was more likely that the grass was cut below the growing point, particularly in dry conditions, which affects the stand longevity and stress tolerance (Jones and Tracy, 2018). As the weather was warm and dry during the harvest events in June in both years, a higher cutting height may have served as a vital management improvement.

The field was mainly fertilised with organic substances, and thus carbon was imported to the system, affecting the net carbon balance. After each of the fertilisation events with organic material, the respiration of the field ~~seemed to increase~~increased, whereas mineral fertilisation was not observed to have an immediate effect on CO<sub>2</sub> fluxes. Increased respiration was likely to occur due to microbial activity of the organic fertilisers. Gilmanov et al. (2007) observed on a Danish agricultural grassland that, although the application of manure increased respiration, ~~also~~ the plant uptake of CO<sub>2</sub> was notably higher than at the other sites studied. Fornara et al. (2016) also concluded, based on their 43-yr study, that manure fertilisation substantially increased soil carbon sequestration of a grassland ecosystem in Northern Ireland. Although the type of the organic fertiliser possibly plays a crucial role, the application of carbon to the system has a direct effect on the carbon balance, but there is also an indirect effect on its components R<sub>eco</sub> and GPP via soil and plant functions.

Concerning our final research question on the relation of possible carbon sink to the international “4 per 1000” carbon sequestration initiative (Minasny et al., 2017), our results show that ~~, on average, the the~~ the Qvidja field acted as a net annual carbon sink by increasing the soil carbon content by ~~0.50~~0.4% annually over the studied period. Thus the site fulfilled the goal of the “4 per 1000” initiative and contributed to the short-term climate change mitigation. Furthermore, the annual carbon balance of our second study year (~~-118~~-86 g C m<sup>-2</sup> yr<sup>-1</sup>) ~~is with improved management practices fits~~ in the upper range of annual carbon sequestration potential (80–120 g C m<sup>-2</sup> yr<sup>-1</sup>) that is evaluated to be attainable with improved management practices (Lal, 2016). Thus, this study demonstrates the potential for a positive impact of northern agricultural grasslands in terms of climate change mitigation.

## 4.2 Errors and uncertainties

Uncertainties ~~with the data in the results~~ are mainly related to the gaps in the measurement data, which required gap-filling of ~~those periods missing measurements~~ with modelled data. The length of a gap increases the related uncertainty, but in our data there were only three longer gaps (4, 8 and 9 days), which all occurred during the first winter, when temperatures were low and only minor fluxes could have been observed. All the other gaps were shorter than 3 days. However, each gap contributed to the uncertainty and were included in the carbon balance calculations. Further uncertainties, which were not included in the error estimates, were ~~caused by the the soil temperature modelling for the first study year and the management flux estimates. involved in the yield measurements and fertilisation input estimates.~~

Carbon balance was calculated based on the ecosystem-atmosphere CO<sub>2</sub> fluxes and the inputs and outputs of harvest and fertilisation. Thus, no other gaseous carbon compounds, such as methane, were considered. Regina et al. (2007) reported that the annual methane ~~exchange balances~~ of a Finnish clay soil ~~varied between during two years were~~ -0.009 and 0.034 g CH<sub>4</sub> m<sup>-2</sup> yr<sup>-1</sup> ~~during two years in 2000–2002. Thus, based.~~ Based on this estimate, the possible carbon emission ~~from as~~ methane accounts for less than 1% of our annual carbon balance.

Leaching of dissolved carbon and emissions of volatile organic compounds may have had an effect on the annual carbon balance. Leaching of carbon from the agricultural soils is mainly driven by ~~the~~ meteorological and hydrological conditions (Manninen et al., 2018), but it is also affected by soil properties (Don and Schulze, 2008). Large variations in soil moisture and temperature and precipitation may increase the solubility of SOM. Generally, however, clay soils retain carbon better than



other soil types. Furthermore, ploughing increases leaching as mineralisation of SOM is enhanced. Depending on precipitation and hydrological and chemical properties of the soil, carbon leaching on grasslands may equal approximately ~~to~~ 25% of the annual carbon balance calculated based on NEE, harvest and fertilisation (Kindler et al., 2011). At our study site, the effect of leaching on the annual carbon balance could be assumed to be fairly small in both summers due to low soil moisture and low precipitation. ~~On the other hand, during wet periods, the leaching may have had a small effect on carbon balance, even though there may have been a temporary contribution during wet periods.~~ However, a more ~~precise~~ accurate carbon balance estimate would require further measurements, including leaching and other carbon-containing gases.

## 5 Conclusions

The agricultural grassland site located at Qvidja in southern Finland acted as a net carbon sink during the two years studied. The carbon balance of the first study year was ~~-50~~ ~~-57~~  $\pm$  ~~68~~ ~~10~~ g C m<sup>-2</sup> yr<sup>-1</sup> and in the second year it was ~~-118~~ ~~-86~~  $\pm$  ~~24~~ ~~12~~ g C m<sup>-2</sup> yr<sup>-1</sup>. We estimated that on average the grassland ~~exceeded~~ reached the goal of the "4 per 1000" initiative intending to increase soil carbon content. The data and results presented here act as a basis for the future studies ~~at this site~~ that focus on the conversion of this farm from intensive agricultural practices towards more sustainable agricultural management ~~and its impacts~~, especially on the impacts of such a conversion on the GHG fluxes occurring on mineral soils in northern conditions. ~~Further~~ Even though we could quantify the sink capacity of the field, further research with longer-term measurements ~~would be~~ is needed to evaluate the persistence of carbon sequestration and storage. Longer time series ~~would be~~ are also essential to study more closely the causes of the interannual variation of GHG fluxes and carbon and water balances at this site, for which the present study provides a baseline.

*Data availability.* The flux and meteorological data as well as the SOC measurements and LAI data are available at Zenodo (<https://doi.org/10.5281/zenodo.4647078>, Heimsch et al. 2020).

## Appendix A: ~~Effective phytomass index~~

~~The PI was used to refine the gap-filling of GPP, especially in the case of long gaps in the nighttime data, based on which GPP was parameterised. PI reflects the development of LAI but, being derived from the daytime NEE measurements, is more dynamic than LAI and thus describes more precisely the course of the photosynthetic activity of plants (Aurela et al., 2001). PI was derived from the net ecosystem CO<sub>2</sub> exchange data by selecting fluxes at high PAR levels. The PAR limit was set to 700  $\mu\text{mol m}^{-2} \text{s}^{-1}$  from March to September and 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$  from October to February. The assumed respiration, i.e. fluxes when  $\text{PAR} < 20 \mu\text{mol m}^{-2} \text{s}^{-1}$ , was subtracted from from the NEE data. This was followed by averaging NEE and  $R_{\text{eco}}$  within a moving window, which was set to 3 days and increased to 5 or 7 days if necessary. Averaging was limited to the harvest dates by decreasing the window size step-by-step to 1.5 days, and similarly increasing it after the harvest. An average GPP was then calculated by subtracting  $R_{\text{eco}}$  from NEE and normalising to unity to obtain PI. Daily PI values were used for~~

calculating the GPP fluxes. Due to the scarcity of respiration data in July in both years and of the daytime data in winter, linear interpolation was applied to cover the missing daily PI values.

## Appendix A: Gap-filling of CO<sub>2</sub> fluxes

The flux data set was separated into sections at the harvest dates, and gap-filling was done separately for these sections by  
475 first parameterising and calculating  $R_{eco}$  and then GPP. The parameter  $R_0$  was determined for each day from the nighttime data ( $PAR < 20 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) with a 7-day moving window.  $E_0$  was determined within the same moving window as  $R_0$ . If there were less than 24 measurements within the time window, its length was increased by 1 day both in-at the beginning and at-the end until enough data were obtained.  $R_0$  was allowed to vary between 0.001 and 1  $\text{mg m}^{-2} \text{s}^{-1}$ . ~~Similarly, the~~ The same minimum number of observations and-within a 3-day moving window was used for determining  $\alpha$  and  $GP_{max}$  from the  
480 observed NEE from which the estimated  $R_{eco}$  had been subtracted.  $\alpha$  and  $GP_{max}$  were allowed to vary between ~~-0.1 and 0.5~~ -0.1 and 0.5 and ~~-0.00001~~ -0.00001  $\text{mg } \mu\text{mol}^{-1}$ , and ~~-5.0 and 0~~ -5.0 and 0  $\text{mg m}^{-2} \text{s}^{-1}$ , respectively. ~~From 5 December 2018 to 26 March 2019 and from 26 November 2019 to 15 March 2020, with no significant CO<sub>2</sub> uptake, a 5-day moving average was used to fill the gaps in NEE.~~

## Appendix B: Gap-filling of energy fluxes

485 The gaps in the net radiation ( $R_n$ ) time series were filled with the monthly mean diurnal cycles. Soil heat flux ( $G$ ) was not measured at our site, so it was estimated from the energy balance closure during the periods when the other energy fluxes were known. Gap-filling of  $G$  was done by assuming a constant ratio between  $G$  and  $R_n$  (Liebethal and Foken, 2007). The ratio of 0.24 was calculated with linear regression from the daytime data (between 10:00–15:00). The sensible and latent heat fluxes ( $Q_H$  and  $Q_E$ , respectively) were gap-filled based on the procedure described by Kowalski et al. (2003). The gaps in the  
490 daytime  $Q_H$  ( $R_n > 0$ ) were filled with monthly linear regression with  $R_n$ . The nighttime gaps in  $Q_H$  ( $R_n < 0$ ) were filled with the corresponding  $R_n$  values. The gaps in the daytime  $Q_E$  were filled in such a way that the monthly mean energy balance closure was achieved. The nighttime gaps in  $Q_E$  were set to 0.

## Appendix C: Management effect on fluxes

The immediate effect of management on the measured NEE and WUE ~~were-was~~ investigated by comparing the mean values  
495 of five days before and after the management day (Table A1).

*Author contributions.* JL and TL planned the flux measurements and TL was responsible for the setup. JPT made the post-processing data corrections and calculated the flux footprint. HV and MK developed the gap-filling code. LH filtered the data and carried out the data analysis.

**Table A1.** Mean flux and meteorological conditions 5 days before and after management. The management day is not included.

	NEE (mg CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )		WUE (g C kg <sup>-1</sup> H <sub>2</sub> O)		PAR (μmol m <sup>-2</sup> s <sup>-1</sup> )		Air T (°C)		Precipitation (mm)		Soil moisture (m <sup>3</sup> m <sup>-3</sup> )	
	Before	After	Before	After	Before	After	Before	After	Before	After	Before	After
Harvest 12 Jun 2018	-0.28	0.03	<del>2.6</del> <del>2.8</del>	<del>0.8</del> <del>0.9</del>	563	646	12.5	16.4	0	0	0.24	0.23
Fertilisation 16 Jul 2018	-0.27	0	<del>1.9</del> <del>2.4</del>	<del>2.0</del> <del>2.5</del>	516	480	19.9	22.4	0	8.3	0.23	0.21
Cutting 21 Aug 2018	na	-0.02	na	<del>1.8</del> <del>1.9</del>	na	290	na	17.2	0	1.2	na	0.23
Fertilisation 24 Aug 2018	-0.10	-0.02	2.0	<del>1.3</del> <del>1.5</del>	382	273	15.7	14.5	6.7	10.4	0.24	0.24
Harvest 23 Sep 2018	-0.10	-0.02	<del>2.4</del> <del>2.5</del>	<del>1.3</del> <del>1.4</del>	183	226	15.1	8.6	0.7	8.6	0.36	0.34
Fertilisation 8 May 2019	-0.17	0.17	2.4	<del>1.8</del> <del>2.0</del>	367	324	3.2	9.6	1.5	3.7	0.37	0.29
Harvest 11 Jun 2019	-0.50	-0.03	<del>2.9</del> <del>3.0</del>	<del>2.2</del> <del>2.6</del>	627	412	21.2	15.8	0.4	0.7	0.23	0.20
Fertilisation 20 Jun 2019	-0.08	-0.08	<del>1.8</del> <del>2.3</del>	<del>1.8</del> <del>2.1</del>	601	622	17.5	17.0	0	2.5	0.20	0.20
Harvest 20 Aug 2019	-0.25	-0.02	<del>3.1</del> <del>3.0</del>	1.4	268	354	16.0	15.9	12.6	9.9	0.35	0.36

JH provided the soil carbon data and ON processed the Sentinel-2 LAI data. LH, AL, JPT and LK prepared the manuscript with contributions from all co-authors.

500 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This study was supported by SITRA, Business Finland [grant 6905/31/2018] and The Strategic Research Council at the Academy of Finland [grant no 327214]. We acknowledge MSc student Niina Ruoho for skillful technical assistance. Qvidja farm owners and staff, especially Pekka Heikkinen and Jonathan Nylund are greatly acknowledged for diverse practical assistance and management of the field.

## 505 References

- Abraha, M., Gelfand, I., Hamilton, S. K., Shao, C., Su, Y.-J., Robertson, G. P., and Chen, J.: Ecosystem water-use efficiency of annual corn and perennial grasslands: contributions from land-use history and species composition, *Ecosystems*, 19, 1001–1012, <https://doi.org/10.1007/s10021-016-9981-2>, 2016.
- Aubinet, M., Vesala, T., and Papale, D.: Eddy covariance: a practical guide to measurement and data analysis, Springer Science & Business Media, <https://doi.org/10.1007/978-94-007-2351-1>, 2012.
- Aurela, M., Laurila, T., and Tuovinen, J.-P.: Seasonal CO<sub>2</sub> balances of a subarctic mire, *Journal of Geophysical Research: Atmospheres*, 106, 1623–1637, <https://doi.org/10.1029/2000JD900481>, 2001.
- Aurela, M., Laurila, T., and Tuovinen, J.-P.: Annual CO<sub>2</sub> balance of a subarctic fen in northern Europe: importance of the wintertime efflux, *Journal of Geophysical Research: Atmospheres*, 107, ACH–17, <https://doi.org/10.1029/2002JD002055>, 2002.
- 515 Baldocchi, D., Chu, H., and Reichstein, M.: Inter-annual variability of net and gross ecosystem carbon fluxes: A review, *Agricultural and Forest Meteorology*, 249, 520–533, <https://doi.org/10.1016/j.agrformet.2017.05.015>, 2018.
- Baldocchi, D. D.: Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future, *Global Change Biology*, 9, 479–492, <https://doi.org/10.1046/j.1365-2486.2003.00629.x>, 2003.
- Bastos, A., Ciais, P., Friedlingstein, P., Sitch, S., Pongratz, J., Fan, L., Wigneron, J., Weber, U., Reichstein, M., Fu, Z., et al.: Direct and seasonal legacy effects of the 2018 heat wave and drought on European ecosystem productivity, *Science Advances*, 6, eaba2724, <https://doi.org/10.1126/sciadv.aba2724>, 2020.
- 520 Bolinder, M., Kätterer, T., Andrén, O., Ericson, L., Parent, L.-E., and Kirchmann, H.: Long-term soil organic carbon and nitrogen dynamics in forage-based crop rotations in Northern Sweden (63–64 N), *Agriculture, Ecosystems & Environment*, 138, 335–342, <https://doi.org/10.1016/j.agee.2010.06.009>, 2010.
- 525 Brunsell, N., Nippert, J., and Buck, T.: Impacts of seasonality and surface heterogeneity on water-use efficiency in mesic grasslands, *Ecohydrology*, 7, 1223–1233, <https://doi.org/10.1002/eco.1455>, 2014.
- Ceschia, E., Béziat, P., Dejoux, J.-F., Aubinet, M., Bernhofer, C., Bodson, B., Buchmann, N., Carrara, A., Cellier, P., Di Tommasi, P., et al.: Management effects on net ecosystem carbon and GHG budgets at European crop sites, *Agriculture, Ecosystems & Environment*, 139, 363–383, <https://doi.org/10.1016/j.agee.2010.09.020>, 2010.
- 530 Chapin, F. S., Woodwell, G. M., Randerson, J. T., Rastetter, E. B., Lovett, G. M., Baldocchi, D. D., Clark, D. A., Harmon, M. E., Schimel, D. S., Valentini, R., et al.: Reconciling carbon-cycle concepts, terminology, and methods, *Ecosystems*, 9, 1041–1050, 2006.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., et al.: Europe-wide reduction in primary productivity caused by the heat and drought in 2003, *Nature*, 437, 529–533, <https://doi.org/10.1038/nature03972>, 2005.
- 535 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., et al.: Carbon and other biogeochemical cycles, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 465–570, Cambridge University Press, [https://pure.mpg.de/rest/items/item\\_2058766/component/file\\_2058769/content](https://pure.mpg.de/rest/items/item_2058766/component/file_2058769/content), 2014.
- De Boeck, H. J., Lemmens, C. M., Bossuyt, H., Malchair, S., Carnol, M., Merckx, R., Nijs, I., and Ceulemans, R.: How do climate warming and plant species richness affect water use in experimental grasslands?, *Plant and Soil*, 288, 249–261, <https://doi.org/10.1007/s11104-006-9112-5>, 2006.
- 540



- de Vries, F. T., Griffiths, R. I., Knight, C. G., Nicolitch, O., and Williams, A.: Harnessing rhizosphere microbiomes for drought-resilient crop production, *Science*, 368, 270–274, <https://doi.org/10.1126/science.aaz5192>, 2020.
- Don, A. and Schulze, E.-D.: Controls on fluxes and export of dissolved organic carbon in grasslands with contrasting soil types, *Biogeochemistry*, 91, 117–131, <https://doi.org/10.1007/s10533-008-9263-y>, 2008.
- Donnelly, C., Greuell, W., Andersson, J., Gerten, D., Pisacane, G., Roudier, P., and Ludwig, F.: Impacts of climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level, *Climatic Change*, 143, 13–26, <https://doi.org/10.1007/s10584-017-1971-7>, 2017.
- Eichelmann, E., Wagner-Riddle, C., Warland, J., Deen, B., and Voroney, P.: Carbon dioxide exchange dynamics over a mature switchgrass stand, *GCB Bioenergy*, 8, 428–442, <https://doi.org/10.1111/gcbb.12259>, 2016.
- Flanagan, L. B., Wever, L. A., and Carlson, P. J.: Seasonal and interannual variation in carbon dioxide exchange and carbon balance in a northern temperate grassland, *Global Change Biology*, 8, 599–615, <https://doi.org/10.1046/j.1365-2486.2002.00491.x>, 2002.
- Foken, T., Leuning, R., Oncley, S. R., Mauder, M., and Aubinet, M.: Corrections and data quality control, in: *Eddy covariance*, pp. 85–131, Springer, [https://doi.org/10.1007/978-94-007-2351-1\\_4](https://doi.org/10.1007/978-94-007-2351-1_4), 2012.
- Fornara, D. A., Wasson, E.-A., Christie, P., and Watson, C. J.: Long-term nutrient fertilization and the carbon balance of permanent grassland: any evidence for sustainable intensification?, *Biogeosciences*, 13, 4975, <https://doi.org/10.5194/bg-13-4975-2016>, 2016.
- Fricker, M. and Willmer, C.: *Stomata*, Springer Science & Business Media, <https://doi.org/10.1007/978-94-011-0579-8>, 2012.
- Gharun, M., Hörtnagl, L., Paul-Limoges, E., Ghiasi, S., Feigenwinter, I., Burri, S., Marquardt, K., Etzold, S., Zweifel, R., Eugster, W., et al.: Physiological response of Swiss ecosystems to 2018 drought across plant types and elevation, *Philosophical Transactions of the Royal Society B*, 375, 20190 521, <https://doi.org/10.1098/rstb.2019.0521>, 2020.
- Gilmanov, T., Soussana, J.-F., Aires, L., Allard, V., Ammann, C., Balzarolo, M., Barcza, Z., Bernhofer, C., Campbell, C., Cernusca, A., et al.: Partitioning European grassland net ecosystem CO<sub>2</sub> exchange into gross primary productivity and ecosystem respiration using light response function analysis, *Agriculture, Ecosystems & Environment*, 121, 93–120, <https://doi.org/10.1016/j.agee.2006.12.008>, 2007.
- Gilmanov, T. G., Aires, L., Barcza, Z., Baron, V. S., Belelli, L., Beringer, J., Billesbach, D., Bonal, D., Bradford, J., Ceschia, E., et al.: Productivity, respiration, and light-response parameters of world grassland and agroecosystems derived from flux-tower measurements, *Rangeland Ecology & Management*, 63, 16–39, <https://doi.org/10.2111/REM-D-09-00072.1>, 2010.
- Gomez-Casanovas, N., Matamala, R., Cook, D. R., and Gonzalez-Meler, M. A.: Net ecosystem exchange modifies the relationship between the autotrophic and heterotrophic components of soil respiration with abiotic factors in prairie grasslands, *Global Change Biology*, 18, 2532–2545, <https://doi.org/10.1111/j.1365-2486.2012.02721.x>, 2012.
- Harman, G. E. and Uphoff, N.: Symbiotic root-endophytic soil microbes improve crop productivity and provide environmental benefits, *Scientifica*, 2019, <https://doi.org/10.1155/2019/9106395>, 2019.
- Harrison, P. A., Dunford, R. W., Holman, I. P., Cojocaru, G., Madsen, M. S., Chen, P.-Y., Pedde, S., and Sandars, D.: Differences between low-end and high-end climate change impacts in Europe across multiple sectors, *Regional environmental change*, 19, 695–709, <https://doi.org/10.1007/s10113-018-1352-4>, 2019.
- Heikkinen, J., Ketoja, E., Nuutinen, V., and Regina, K.: Declining trend of carbon in Finnish cropland soils in 1974–2009, *Global Change Biology*, 19, 1456–1469, <https://doi.org/10.1111/gcb.12137>, 2013.
- Houghton, R. A. and Nassikas, A. A.: Global and regional fluxes of carbon from land use and land cover change 1850–2015, *Global Biogeochemical Cycles*, 31, 456–472, <https://doi.org/10.1002/2016GB005546>, 2017.

- Hu, Z., Yu, G., Fu, Y., Sun, X., Li, Y., Shi, P., Wang, Y., and Zheng, Z.: Effects of vegetation control on ecosystem water use efficiency within and among four grassland ecosystems in China, *Global Change Biology*, 14, 1609–1619, <https://doi.org/10.1111/j.1365-2486.2008.01582.x>, 2008.
- Jensen, R., Herbst, M., and Friborg, T.: Direct and indirect controls of the interannual variability in atmospheric CO<sub>2</sub> exchange of three contrasting ecosystems in Denmark, *Agricultural and Forest Meteorology*, 233, 12–31, <https://doi.org/10.1016/j.agrformet.2016.10.023>, 2017.
- Jones, G. and Tracy, B.: Persistence and productivity of orchardgrass and orchardgrass/alfalfa mixtures as affected by cutting height, *Grass and Forage Science*, 73, 544–552, <https://doi.org/10.1111/gfs.12309>, 2018.
- Kindler, R., Siemens, J., Kaiser, K., Walmsley, D. C., Bernhofer, C., Buchmann, N., Cellier, P., Eugster, W., Gleixner, G., GRÜNWALD, T., et al.: Dissolved carbon leaching from soil is a crucial component of the net ecosystem carbon balance, *Global Change Biology*, 17, 1167–1185, <https://doi.org/10.1111/j.1365-2486.2010.02282.x>, 2011.
- Kormann, R. and Meixner, F. X.: An analytical footprint model for non-neutral stratification, *Boundary-Layer Meteorology*, 99, 207–224, <https://doi.org/10.1023/A:1018991015119>, 2001.
- Kowalski, S., Sartore, M., Burrell, R., Berbigier, P., and Loustau, D.: The annual carbon budget of a French pine forest (*Pinus pinaster*) following harvest, *Global Change Biology*, 9, 1051–1065, <https://doi.org/10.1046/j.1365-2486.2003.00627.x>, 2003.
- Kurc, S. A. and Small, E. E.: Soil moisture variations and ecosystem-scale fluxes of water and carbon in semiarid grassland and shrubland, *Water Resources Research*, 43, <https://doi.org/10.1029/2006WR005011>, 2007.
- Kutsch, W., Aubinet, M., Buchmann, N., Smith, P., Osborne, B., Eugster, W., Wattenbach, M., Schrumpf, M., Schulze, E., Tomelleri, E., et al.: The net biome production of full crop rotations in Europe, *Agriculture, ecosystems & environment*, 139, 336–345, 2010.
- Lal, R.: Beyond COP 21: potential and challenges of the “4 per Thousand” initiative, *Journal of Soil and Water Conservation*, 71, 20A–25A, <https://doi.org/10.2489/jswc.71.1.20A>, 2016.
- Laurila, T., Tuovinen, J.-P., Lohila, A., Hatakka, J., Aurela, M., Thum, T., Pihlatie, M., Rinne, J., and Vesala, T.: Measuring methane emissions from a landfill using a cost-effective micrometeorological method, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005GL023462>, 2005.
- Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R. A., Foster, P., et al.: Trends in the sources and sinks of carbon dioxide, *Nature Geoscience*, 2, 831–836, <https://doi.org/10.1038/ngeo689>, 2009.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Pongratz, J., Manning, A. C., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Jackson, R. B., et al.: Global carbon budget 2017, *Earth System Science Data Discussions*, pp. 1–79, <https://doi.org/10.5194/essd-10-405-2018>, 2017.
- Liebethal, C. and Foken, T.: Evaluation of six parameterization approaches for the ground heat flux, *Theoretical and Applied Climatology*, 88, 43–56, <https://doi.org/10.1007/s00704-005-0234-0>, 2007.
- Lind, S. E., Shurpali, N. J., Peltola, O., Mammarella, I., Hyvönen, N., Maljanen, M., Rätty, M., Virkajärvi, P., and Martikainen, P. J.: Carbon dioxide exchange of a perennial bioenergy crop cultivation on a mineral soil, *Biogeosciences*, 13, 1255–1268, <https://doi.org/10.5194/bg-13-1255-2016>, 2016.
- Lind, S. E., Virkajärvi, P., Hyvönen, N. P., Maljanen, M., Kivimäenpää, M., Jokinen, S., Antikainen, S., Latva, M., Rätty, M., Martikainen, P. J., et al.: Carbon dioxide and methane exchange of a perennial grassland on a boreal mineral soil, *Boreal Environmental Research*, 25, 1–17, <http://urn.fi/URN:NBN:fi-fe202002135420>, 2020.

- Lloyd, J. and Taylor, J.: On the temperature dependence of soil respiration, *Functional Ecology*, pp. 315–323, <https://doi.org/10.2307/2389824>, 1994.
- Lohila, A., Aurela, M., Tuovinen, J.-P., and Laurila, T.: Annual CO<sub>2</sub> exchange of a peat field growing spring barley or perennial forage grass, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004715>, 2004.
- 620 Lorenz, K. and Lal, R.: Soil Carbon Stock, in: *Carbon Sequestration in Agricultural Ecosystems*, pp. 39–136, Springer, [https://doi.org/10.1007/978-3-319-92318-5\\_2](https://doi.org/10.1007/978-3-319-92318-5_2), 2018.
- Manninen, N., Soinne, H., Lemola, R., Hoikkala, L., and Turtola, E.: Effects of agricultural land use on dissolved organic carbon and nitrogen in surface runoff and subsurface drainage, *Science of the Total Environment*, 618, 1519–1528, <https://doi.org/10.1016/j.scitotenv.2017.09.319>, 2018.
- 625 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., et al.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- Moreau, D., Bardgett, R. D., Finlay, R. D., Jones, D. L., and Philippot, L.: A plant perspective on nitrogen cycling in the rhizosphere, *Functional Ecology*, 33, 540–552, <https://doi.org/10.1111/1365-2435.13303>, 2019.
- Paustian, K., Six, J., Elliott, E., and Hunt, H.: Management options for reducing CO<sub>2</sub> emissions from agricultural soils, *Biogeochemistry*, 48, 147–163, <https://doi.org/10.1023/A:1006271331703>, 2000.
- 630 Pirasteh-Anosheh, H., Saed-Moucheshi, A., Pakniyat, H., and Pessarakli, M.: Stomatal responses to drought stress, *Water Stress and Crop Plants* (Chichester, UK: John Wiley & Sons, Ltd, pp. 24–40, <https://doi.org/10.1002/9781119054450>, 2016.
- Pirinen, P., Simola, H., Aalto, J., Kaukoranta, J.-P., Karlsson, P., and Ruuhela, R.: Tilastoja Suomen ilmastosta 1981–2010, *Ilmatieteen laitos*, 2012.
- 635 Rankinen, K., Karvonen, T., and Butterfield, D.: A simple model for predicting soil temperature in snow-covered and seasonally frozen soil: model description and testing, *Hydrology and Earth System Sciences Discussions*, 8, 706–716, <https://hal.archives-ouvertes.fr/hal-00304956>, 2004.
- Rebmann, C., Kolle, O., Heinesch, B., Queck, R., Ibrom, A., and Aubinet, M.: Data acquisition and flux calculations, in: *Eddy covariance*, pp. 59–83, Springer, [https://doi.org/10.1007/978-94-007-2351-1\\_3](https://doi.org/10.1007/978-94-007-2351-1_3), 2012.
- 640 Regina, K., Pihlatie, M., Esala, M., and Alakukku, L.: Methane fluxes on boreal arable soils, *Agriculture, Ecosystems & Environment*, 119, 346–352, <https://doi.org/10.1016/j.agee.2006.08.002>, 2007.
- Reinsch, T., Loges, R., Kluß, C., and Taube, F.: Effect of grassland ploughing and reseeded on CO<sub>2</sub> emissions and soil carbon stocks, *Agriculture, ecosystems & environment*, 265, 374–383, <https://doi.org/10.1016/j.agee.2018.06.020>, 2018.
- Shurpali, N. J., Hyvönen, N. P., Huttunen, J. T., Clement, R. J., Reichstein, M., Nykänen, H., Biasi, C., and Martikainen, P. J.: Cultivation of a perennial grass for bioenergy on a boreal organic soil—carbon sink or source?, *GCB Bioenergy*, 1, 35–50, <https://doi.org/10.1111/j.1757-1707.2009.01003.x>, 2009.
- 645 Singh, M., Sarkar, B., Sarkar, S., Churchman, J., Bolan, N., Mandal, S., Menon, M., Purakayastha, T. J., and Beerling, D. J.: Stabilization of soil organic carbon as influenced by clay mineralogy, in: *Advances in Agronomy*, vol. 148, pp. 33–84, Elsevier, <https://doi.org/10.1016/bs.agron.2017.11.001>, 2018.
- 650 Smith, P.: Land use change and soil organic carbon dynamics, *Nutrient Cycling in Agroecosystems*, 81, 169–178, <https://doi.org/10.1007/s10705-007-9138-y>, 2008.

- Smith, P., Soussana, J.-F., Angers, D., Schipper, L., Chenu, C., Rasse, D. P., Batjes, N. H., van Egmond, F., McNeill, S., Kuhnert, M., et al.: How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal, *Global Change Biology*, 26, 219–241, <https://doi.org/10.1111/gcb.14815>, 2020.
- 655 Soussana, J.-F., Allard, V., Pilegaard, K., Ambus, P., Amman, C., Campbell, C., Ceschia, E., Clifton-Brown, J., Czóbel, S., Domingues, R., et al.: Full accounting of the greenhouse gas (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>) budget of nine European grassland sites, *Agriculture, Ecosystems & Environment*, 121, 121–134, <https://doi.org/10.1016/j.agee.2006.12.022>, 2007.
- Soussana, J.-F., Tallec, T., and Blanfort, V.: Mitigating the greenhouse gas balance of ruminant production systems through carbon sequestration in grasslands, *Animal: The International Journal of Animal Biosciences*, 4, 334, <https://doi.org/10.1017/S175173110999078>, 2010.
- 660 Steduto, P.: Water use efficiency, in: *Sustainability of Irrigated Agriculture*, pp. 193–209, Springer, [https://doi.org/10.1007/978-94-015-8700-6\\_12](https://doi.org/10.1007/978-94-015-8700-6_12), 1996.
- Tang, X., Li, H., Desai, A. R., Nagy, Z., Luo, J., Kolb, T. E., Oliosio, A., Xu, X., Yao, L., Kutsch, W., et al.: How is water-use efficiency of terrestrial ecosystems distributed and changing on Earth?, *Scientific reports*, 4, 7483, <https://doi.org/10.1038/srep07483>, 2014.
- Weiss, M. and Baret, F.: S2ToolBox Level 2 products: LAI, FAPAR, FCOVER, Available online: [step.esa.int/docs/extra/ATBD\\_S2ToolBox\\_L2B\\_V1.1.pdf](https://step.esa.int/docs/extra/ATBD_S2ToolBox_L2B_V1.1.pdf), 2016.
- 665 Willmer, C. and Fricker, M.: *Stomata*, vol. 2, Springer Science & Business Media, 1996.
- Wolf, S., Eugster, W., Ammann, C., Häni, M., Zielis, S., Hiller, R., Stieger, J., Imer, D., Merbold, L., and Buchmann, N.: Contrasting response of grassland versus forest carbon and water fluxes to spring drought in Switzerland, *Environmental Research Letters*, 8, 035 007, <https://doi.org/10.1088/1748-9326/8/3/035007>, 2013.
- 670 Xu, H.-j., Wang, X.-p., Zhao, C.-y., and Zhang, X.-x.: Responses of ecosystem water use efficiency to meteorological drought under different biomes and drought magnitudes in northern China, *Agricultural and Forest Meteorology*, 278, 107 660, <https://doi.org/10.1016/j.agrformet.2019.107660>, 2019.
- Yang, Y., Tilman, D., Furey, G., and Lehman, C.: Soil carbon sequestration accelerated by restoration of grassland biodiversity, *Nature communications*, 10, 1–7, <https://doi.org/10.1038/s41467-019-08636-w>, 2019.