

# Multi-compartment kinetic-allometric model of radionuclide bioaccumulation in marine fish

Roman Bezhenar<sup>1</sup>, Kyeong Ok Kim<sup>2</sup>, Vladimir Maderich<sup>1</sup>, Govert de With<sup>3</sup>, and Kyung Tae Jung<sup>4</sup>

<sup>1</sup>Institute of Mathematical Machine and System Problems, Glushkov av., 42, Kyiv 03187, Ukraine

<sup>2</sup>Korea Institute of Ocean Science and Technology, Metropolitan city Busan, Republic of Korea

<sup>3</sup>NRG, Utrechtseweg 310, 6800 ES Arnhem, the Netherlands


<sup>4</sup>Oceanic Consulting & Trading, 403 Munhwa-building, 90 Yangpyong-ro, Seoul, Republic of Korea

**Correspondence:** Vladimir Maderich (vladmad@gmail.com)

**Abstract.** A model of the radionuclide accumulation in fish taking into account the contribution of different tissues and allometry is presented. The basic model assumptions are as follows: (i) A fish organism is represented by several compartments in which radionuclides are homogeneously distributed; (ii) The compartments correspond to three groups of organs/tissues: muscle, bones and organs (kidney, liver, gonads, etc.) differing in metabolic function; (iii) Two input compartments include gills absorbing contamination from water and digestive tract through which contaminated food is absorbed; (iv) The absorbed radionuclide is redistributed between organs/tissues according to their metabolic functions; (v) The elimination of assimilated elements from each group of organs/tissues differs, reflecting differences in specific tissues/organs in which elements were accumulated; and (vi) The food and water uptake rates, elimination rate and growth rate depend on the metabolic rate, which is scaled by fish mass to the 3/4 power. The analytical solutions of the system of model equations describing dynamics of the assimilation and elimination of <sup>134</sup>Cs, <sup>57</sup>Co, <sup>60</sup>Co, <sup>54</sup>Mn and <sup>65</sup>Zn, which are preferably accumulated in different tissues, exhibited good agreement with the laboratory experiments. The developed multi-compartment kinetic-allometric model was embedded into the compartment-box model POSEIDON-R, which describes transport of radionuclides in water, accumulation in the sediment, and transfer of radionuclides through the pelagic and benthic food webs. The POSEIDON-R model was applied for the simulation of the transport and fate of <sup>60</sup>Co and <sup>54</sup>Mn routinely released from Forsmark NPP located on the Baltic Sea coast of Sweden and for calculation of <sup>90</sup>Sr concentration in fish after the accident at Fukushima Dai-ichi NPP. ~~Predicted~~ Computed concentrations of radionuclides in fish agree with the measurements much better than ~~predicted-calculated~~ using standard whole-body model and target tissue model. The model with the defined generic parameters could be used in different marine environments without calibration based on *a posteriori* information, which is important for emergency decision support systems.

## 1 Introduction

Accumulation of radionuclides in marine organisms is a complicated process that is governed by uptake of radionuclides from water, sediment and food and by depuration. In turn, these processes depend on the chemical properties of elements, their roles in metabolic processes, the positions of organisms in the food web and marine environmental parameters. In the case of chronic



**Figure 1.** Distribution of accumulated <sup>137</sup>Cs activities of isotopes Cs, <sup>90</sup>Sr and <sup>60</sup>Co in muscle, bone, liver and kidneys according to previously reported data (Yankovich, 2003; Yankovich et al., 2010).

exposure, the radiological assessment models often assumed an equilibrium approach (Carvalho, 2018), in which concentration in the organism relates to the concentration in water using a biological accumulation factor (*BAF*). However, to describe highly time dependent transfer processes resulting from accidental releases, dynamic models for the uptake and retention of activity in marine organisms are necessary (Vives i Batlle et al., 2016). According to Takata et al. (2019), the effective ~~recession times~~ half-lives of post-Fukushima Dai-ichi Nuclear Power Plant (FDNPP) accident disequilibrium of <sup>137</sup>Cs in biota ranged from 100 to 1,100 days. The most commonly used bioaccumulation models are the whole-body models, where the organism is represented as a single box in which contamination is evenly distributed (e.g. Fowler and Fisher, 2004; ~~Tateda et al., 2013;~~ Vives i Batlle et al., 2016). However, the distribution of radionuclides in organisms, and in particular in fish, is non-uniform. For example, the highest concentration of radiocaesium in fish is observed in the muscle, while the highest concentrations of the actinides, plutonium and americium, are measured in specific organs (Coughtrey and Thorne, 1983). Moreover, Vives i Batlle (2012) noted that elimination of activity from organisms occurred with different rates that can be interpreted as elimination from different tissues/organs with different metabolism. In a first approximation, this is used in the “target tissue” approach (Heling et al., 2002; ~~Tateda et al., 2013;~~ Maderich et al., 2014a,b; Bezhenar et al., 2016), where radionuclides are grouped into several classes depending on the type of tissues in which a specific radionuclide accumulates preferentially (target tissue). However, the contribution of other tissues with greater mass than the mass of the target tissue can be commensurate with the contribution of the target tissue to the amount of radioactivity in the body. ~~This is observed in Fig. 1, which is built from~~ Distribution of accumulated activities of isotopes Cs, Sr and Co in muscle, bone and liver estimated from previously reported data (Yankovich, 2003; Yankovich et al., 2010) where are shown in Fig. 1. The accumulated activity in a given tissue was calculated as a ratio of tissue mass fraction (%) (Yankovich, 2003) to body-to-tissue concentration ratio (Yankovich et al., 2010). As seen in Fig. 1, the accumulated activity of <sup>90</sup>Sr in muscle is not negligible in comparison with accumulated activity in bones, whereas the accumulated activity of <sup>60</sup>Co is redistributed between muscle, bone and ~~organs~~ liver.

A more general approach to the description of the radionuclide accumulation in the tissues of fish is using the physiologically based pharmacokinetic (PBPK) models (Barron et al., 1990; Thomann et al., 1997; Garnier-Laplace et al., 2000; Otero-Muras


et al., 2010; [Grech et al. 2019](#)). In the PBPK models, the fish organism is represented as three groups of compartments: absorption compartments simulating uptake of contaminants, distribution compartments simulating tissues and organs, and excretion compartments. The exchange of contaminants between compartments is limited by blood flux perfusing compartments. However, these models require a significant number of parameters depending on elements, fish species and marine environments. They must be determined from the laboratory experiments (Thomann et al., 1997) or by the optimization procedures (Otero-Muras et al., 2010). ~~Notice that~~[Note that, with the exception of model \(Grech et al. 2019\),](#) PBPK fish models do not ~~yet~~ include scaling (allometric) relationships between metabolic rates and organism mass (West et al., 1997; Higley and Bytwerk, 2007; Vives i Batlle et al., 2007; Beresford et al., 2016). Therefore, there is a need to develop a generic model of intermediate complexity between the one-compartment model and the PBPK model taking into account (i) the heterogeneity of the distribution of contamination in fish tissues and (ii) the allometric relationships between metabolic rates and organism mass. Such a model can be used for accidental release simulations without local calibration, which is a complicated task in the circumstances of the accident.

In this paper, a new approach for predicting radionuclide accumulation in fish taking into account the contributions of different tissues and allometry is presented. The [developed multi-compartment kinetic-allometric \(MCKA\) model is embedded into the box model POSEIDON-R \(Lepicard et al., 2004; Maderich et al., 2014a,b; 2018b; Bezhenar et al., 2016\) which describes transport of radionuclides in water, accumulation in the sediment, and transfer of radionuclides through the pelagic and benthic food webs.](#) The paper is organized as follows. The [MCKA](#) model is described in Section 2. The comparison with laboratory experiments is given in Section 3. The results of simulation of several radionuclides in the marine environment for regular and accidental releases are described in Section 4. The conclusions are presented in Section 5.

## 2 Model

### 2.1 Model equations

Here, a simple multi-compartmental model to simulate kinetics of radionuclides in the fish is described. The basic assumptions are as follows: (i) a fish organism is represented by several compartments in which radionuclides are homogeneously distributed; (ii) the compartments correspond to three groups of organs/tissues differing in metabolic function: ~~flesh~~[muscle](#), bones and organs (kidney, liver, gonads, etc.); (iii) two input compartments include gills which absorb contamination from water and digestive tract through which contaminated food is absorbed; (iv) the absorbed radionuclide is redistributed between organs/tissues according to their metabolic functions; (v) the elimination of assimilated elements from each group of organs/tissues differs, reflecting differences in the specific tissues/organs in which elements were accumulated; (vi) the food and water uptake rates, elimination rate and growth rate depend on the metabolic rate, which is scaled by fish mass to the 3/4 power following general theory (West et al., 1997)~~describing transport of essential materials through space-filling fractal networks of branching tubes in organism.~~ [A schematic representation of the model is shown in Fig. 2.](#)



**Figure 2.** Schematic of the multi-compartment kinetic-allometric model

The equation for concentration of radionuclide  $C_1$  [Bq kg<sup>-1</sup> wet weight (WW)] in the gill compartment ( $i = 1$ ) is written as

$$80 \quad \frac{dm_1 C_1}{dt} = K_w m C_w - k_1 m_1 C_1 - \lambda_1 m_1 C_1. \quad (1)$$

The equation for concentration of unabsorbed radionuclide  $C_2$  [Bq kg<sup>-1</sup> WW] in the digestive tract compartment ( $i = 2$ ) is

$$\frac{dm_2 C_2}{dt} = K_f m C_f - k_2 m_2 C_2 - \lambda_2 m_2 C_2. \quad (2)$$

The equations for concentrations of radionuclide  $C_i$  [Bq kg<sup>-1</sup> WW] in the muscle ( $i = 3$ ), bones ( $i = 4$ ) and organs ( $i = 5$ ) are

$$85 \quad \frac{dm_i C_i}{dt} = k_{2i} m_2 C_2 + k_{1i} m_1 C_1 - \lambda_i m_i C_i. \quad (3)$$

Here,  $t$  is time;  $m_i$  is the mass of the  $i$ -th tissues;  $m$  [kg] is the total mass of fish;  $C_w$  [Bq m<sup>-3</sup>] is the concentration of radionuclide in water;  $C_f$  [Bq kg<sup>-1</sup> WW] is the concentration of radionuclide in food;  $K_w$  [(m<sup>3</sup>(kg d)<sup>-1</sup>] is a water uptake rate per unit fresh weight of fish and  $K_f$  [(kg(kg d)<sup>-1</sup>] is a food uptake rate per unit fresh weight of fish;  $\lambda_1$  is a loss rate from gills to water [d<sup>-1</sup>];  $k_1$ ,  $k_{1i}$ ,  $k_2$ , and  $k_{2i}$  [d<sup>-1</sup>] are transfer rates between tissues,  $\lambda_2$  [d<sup>-1</sup>] is the egestion rate from the digestive tract;  $\lambda_i$  [d<sup>-1</sup>] is the absorbed radionuclide elimination rate from different tissues/organs ( $i = \overline{3, 5}$ ).

The activity concentration in the food  $C_f$  is expressed by the following equation, summing for a total of  $n$  prey types

$$C_f = \sum_{j=0}^n C_{prey,j} P_j \frac{drw_{pred}}{drw_{prey,j}}, \quad (4)$$

where  $C_{prey,j}$  is the activity concentration in prey of type  $0 \leq j \leq n$ ,  $P_j$  is preference for prey of type  $j$ ,  $drw_{pred}$  is the dry weight fraction of fish, and  $drw_{prey,j}$  is the dry weight fraction of prey of type  $j$ . The mean whole-body concentration of

95 activity in the organism  $C_{wb}$  and whole-body activity  $A_{wb}$  [Bq] are calculated as

$$A_{wb} = mC_{wb} = \sum_{i=1}^5 m_i C_i = m \sum_{i=1}^5 \mu_i C_i, \quad (5)$$

where  $\mu_i$  are weighting factors, ( $\sum_{i=1}^5 \mu_i = 1$ ).

Transfer rates  $k_1$  and  $k_2$  are related with tissue transfer rates  $k_{1i}$  and  $k_{2i}$  as

$$k_1 = \sum_{i=3}^5 k_{1i}, \quad k_2 = \sum_{i=3}^5 k_{2i}. \quad (6)$$

100 Summing eqns. (1) to (3) yields the equation for total concentration of activity in fish  $C_{wb}$  as

$$\frac{dC_{wb}}{dt} = K_w C_w + K_f C_f - \lambda_1 \mu_1 C_1 - \lambda_2 \mu_2 C_2 - \sum_{i=3}^5 \lambda_i \mu_i C_i - \lambda_g C_{wb}, \quad (7)$$

where  $\lambda_g$  is the organism growth rate, defined as

$$\lambda_g = \frac{1}{m} \frac{dm}{dt}. \quad (8)$$

The growth dilution can be ignored in the model calculations when  $\lambda_g \ll \lambda_i$ . For short-lived radionuclides,  $\lambda_i$  should be  
 105 corrected taking into account the physical decay. The assimilation efficiencies of elements from water  $AE_w$  and food  $AE_f$   
 (Pouil et al., 2018) can be introduced, assuming that uptake from water and food is equilibrated by loss to the water from gills  
 and through the egestion. The corresponding relations are

$$AE_w = \frac{K_w C_w - \lambda_1 \mu_1 C_1}{K_w C_w}, \quad (9)$$

$$AE_f = \frac{K_f C_f - \lambda_2 \mu_2 C_2}{K_f C_f}. \quad (10)$$

110 Taking into account the relations (9)-(10) for constant  $AE_w$  and  $AE_f$ , the equation (7) will be similar to the standard whole-body  
 single compartment equation

$$\frac{dC_{wb}}{dt} = AE_w K_w C_w + AE_f K_f C_f - (\lambda_{wb} + \lambda_g) C_{wb}, \quad (11)$$

if elimination terms in (7) are replaced by a single term  $\lambda_{wb} C_{wb}$ , assuming that  $\lambda_{wb}$  is a single whole-body elimination rate  
 (Fowler and Fisher, 2004).

115 The food and water uptake rates, elimination rate and growth rate depend on the metabolic rate, which in turn is known to  
 scale by the organism mass. Here, we employed quarter-power scaling for uptake, elimination and growth rates derived from  
 general theory (West et al., 1997). [This theory predict for all organisms the 3/4 power law for metabolic rates. It describes  
 transport of essential materials through space-filling fractal networks of branching tubes assuming that the energy dissipation  
 is minimized and that the terminal branch of the network is a size-invariant. The scaling relations are](#)

$$120 \quad K_w(m) = \alpha_w m^{-1/4}, \quad K_f(m) = \alpha_f m^{-1/4}, \quad \lambda_g(m) = \alpha_g m^{-1/4}, \quad \lambda_i(m) = \alpha_i m^{-1/4}, \quad (12)$$

**Table 1.** Parameters in allometric relations, standard deviation *STD* of parameters and number of measurements *N*.

Constant	Value	<i>N</i>	<i>STD</i>	Data source
$\alpha_w$	0.08	1	-	Mathews et al. (2008)
$\alpha_f$	0.012	7	0.005	Alava and Gobas (2016)
$\alpha_g$	0.0012	7	0.0003	Alava and Gobas (2016)
$\alpha_1$	<del>0.8</del> 800	-	-	This study
$\alpha_2$	0.75	7	0.18	Andersen (1984); Pouil et al. (2017)
$\alpha_3$	0.007	8	0.0017	Jeffree et al. (2006); Mathews and Fisher (2008); Mathews et al. (2008)
$\alpha_4$	0.001	-	-	Heling et al. (2002)
$\alpha_5$	0.0275	1	-	Rouleau et al. (1995)

where  $\alpha_w, \alpha_f, \alpha_g, \alpha_i$  ( $i = \overline{1,5}$ ) are constants. These parameters can also depend on temperature, salinity and fish age (e.g. Belharet et al., 2019; Heling and Bezhenar, 2009). Notice that a number of laboratory experiments (Thomas and Fisher, 2010) showed that temperature exerts no major influence on uptake and elimination, whereas the effect of salinity varies for elements (Heling and Bezhenar, 2009; Jeffree et al, 2017). Here, we did not analyze these factors requiring separate consideration.

125 The values of constants  $\alpha_w, \alpha_f, \alpha_g, \alpha_i$  ( $i = \overline{1,5}$ ) estimated from laboratory experiments and marine data are provided in Table 1, whereas values of weighting factors  $\mu_1=0.01, \mu_2=0.01, \mu_3=0.78, \mu_4=0.12, \mu_5=0.08$ , were chosen using estimates from Yankovich ~~et al. (2010). The model parameters for characteristic values of the fish mass are given in Table S1 (Supplementary Material).~~ (2003).

## 2.2 Kinetics in equilibrium state

130 The model parameters can be estimated using measurement data and applying the kinetic equations under equilibrium conditions. Equations (1) - (3) rewritten for radionuclide concentrations in the equilibrium state are

$$\mu_1 C_1 = \frac{K_w C_w}{k_1 + \lambda_1 + \lambda_g}, \quad (13)$$

$$\mu_2 C_2 = \frac{K_f C_f}{k_2 + \lambda_2 + \lambda_g}, \quad (14)$$

$$k_{2i} \frac{K_f C_f}{k_2 + \lambda_2 + \lambda_g} + k_{1i} \frac{K_w C_w}{k_1 + \lambda_1 + \lambda_g} = \lambda_i \mu_i C_i + \lambda_g \mu_i C_i. \quad (15)$$

135 Then, using (13)-(14), the assimilation efficiencies of elements from water  $AE_w$  and food  $AE_f$  are rewritten as

$$AE_w = \frac{k_1 + \lambda_g}{k_1 + \lambda_1 + \lambda_g}, \quad AE_f = \frac{k_2 + \lambda_g}{k_2 + \lambda_2 + \lambda_g} \quad (16)$$

When  $\lambda_1 \gg \lambda_g, \lambda_2 \gg \lambda_g, \lambda_1 \gg \lambda_g, \lambda_2 \gg \lambda_g$ , we determine from (16) and (12), approximately,

$$k_1 = \frac{AE_w \lambda_1}{1 - AE_w} = \frac{AE_w \alpha_1}{(1 - AE_w) m^{1/4}}, \quad k_2 = \frac{AE_f \lambda_2}{1 - AE_f} = \frac{AE_f \alpha_2}{(1 - AE_f) m^{1/4}}. \quad (17)$$

~~The equations under~~ This assumption also imposes requirement on the modelling of radionuclides with decay constant  $\lambda \ll \lambda_i$ .

140 The equations (17) ~~is-are~~ used to relate kinetic coefficients of the model with experimentally determined parameters  $AE_w$ ,  $AE_f$ ,  $\lambda_1$  and  $\lambda_2$ . The ~~equations under~~ equation (15) can be rewritten as

$$AE_{fi}\alpha_f BAF_{food} + AE_{wi}\alpha_w = \mu_i(\alpha_i + \alpha_g) \frac{BAF_{wb}}{CR_i}, \quad (18)$$

where  $AE_{wi}$  and  $AE_{fi}$  are assimilation efficiencies for tissue  $i$ ,

$$\sum_{i=3}^5 \mu_i AE_{wi} \approx AE_w, \quad \sum_{i=3}^5 \mu_i AE_{fi} \approx AE_f. \quad (19)$$

145 The assimilation efficiencies are expressed through kinetic coefficients as

$$\mu_i AE_{wi} = \frac{k_{1i}}{k_1} AE_w, \quad \mu_i AE_{fi} = \frac{k_{2i}}{k_2} AE_f. \quad (20)$$

~~The bioaccumulation factors for food  $BAF_{food}$ , for~~ We define bioconcentration factor ( $BCF$ ) as ratio of whole-body of fish to water concentrations with no dietary intake, bioaccumulation factor ( $BAF$ ) as ratio of whole-body of fish  ~~$BAF_{wb}$  and to water concentrations with dietary intake,~~ body-to-tissue concentration ratio ( $CR_i$ ) as ratio of whole-body to  $i$ -th tissue concentrations,

150 ~~whereas ratio of food to water concentrations is indicated as  $BAF_{food}$ . The parameters  $BAF$ ,  $CR_i$  and  $BAF_{food}$  are described as~~

$$BAF_{food} = \frac{C_{food}}{C_w}, \quad BAF_{wb} = \frac{C_{wb}}{C_w}, \quad CR_i = \frac{C_{wb}}{C_i}, \quad BAF_{food} = \frac{C_{food}}{C_w}. \quad (21)$$

Values of  $BAF$  for different radionuclides are ~~available in IAEA (expressed as  $CF$  in (IAEA, 2004).~~ Yankovich et al. (2010) provide  $CR_i$  based on aggregate experimental data for marine fish. Assume that the kinetics of assimilation in fish tissues are

155 similar for radionuclides absorbed from water and food, i.e.

$$\frac{AE_{wi}}{AE_w} = \frac{AE_{fi}}{AE_f}. \quad (22)$$

Notice that assimilation for some elements can be considered as route dependent (Reinfelder et al., 1999), and so (22) is only a first approximation. Inserting (22) into (18) yields

$$\frac{AE_{fi}}{AE_f} (AE_f \alpha_f BAF_{food} + AE_w \alpha_w) = BAF_{wb} \frac{\mu_i(\alpha_i + \alpha_g)}{CR_i}, \quad (23)$$

160 Summing (23) for  $i = 3, 4, 5$  yields

$$AE_f \alpha_f BAF_{food} + AE_w \alpha_w = BAF_{wb} \sum_{i=3}^5 \frac{\mu_i(\alpha_i + \alpha_g)}{CR_i}. \quad (24)$$

Using (23) and (24), ratio  $AE_{fi}/AE_f$  can be written as

$$\frac{AE_{fi}}{AE_f} = \mu_i(\alpha_i + \alpha_g) / \left( CR_i \sum_{i=3}^5 \frac{\mu_i(\alpha_i + \alpha_g)}{CR_i} \right). \quad (25)$$

**Table 2.** The food assimilation efficiency  $AE_f$  (Pouil et al., 2018), assimilation efficiency of elements from water  $AE_w$ , tissue assimilation efficiencies  $AE_{fi}$  and  $TTF$  for several elements.

Element	$AE_f$	<u><math>AE_w</math></u>	$AE_{fi}$
Cs	0.76	<u>0.001<sup>1</sup></u>	0
Sr	0.29 <sup>*2</sup>	<u>0.00003<sup>3</sup></u>	0
Co	0.081	<u>0.0025<sup>1</sup></u>	0
Mn	0.24	<u>0.0045<sup>1</sup></u>	0
Zn	0.22	<u>0.0065<sup>1</sup></u>	0
<u>Ag-0.033-0.016-0.007-0.24-0.03-Cu-0.2-0.07-0.03-0.96-0.16-Cd-0.2-0.15-0.06-0.95-0.29-Cr-0.023-0.01-0.009-0.17-0.03</u> -height			

<sup>1</sup>The values were estimated from experiments by Mathews and Fisher (2008)

<sup>2</sup>The value was estimated using  $BAF$  from IAEA (2004).

<sup>3</sup>Heling and Bezhenar (2009) for seawater environment.

The values of kinetic coefficients  $k_1$ ,  $k_2$  and assimilation efficiencies for tissues  $AE_{fi}$  were calculated from (17) and (25) using  
165 assimilation efficiencies from experimental data (Pouil et al., 2018). The values of  $AE_{fi}$  for several radionuclides are given in  
Table 2. Equation (24) can be rearranged to express the ratio of  $BAF_{wb}$  to  $BAF_{food}$  taking into account dominance of dietary  
intake over water intake (Mathews and Fisher, 2009). This ratio is the trophic transfer factor ( $TTF$ ), written as

$$TTF = AE_f \alpha_f / \sum_{i=3}^5 \frac{\mu_i (\alpha_i + \alpha_g)}{CR_i}. \quad (26)$$

A  $TTF > 1$  indicates possible biomagnification, and  $TTF < 1$  indicates that biodiminution is likely. As follows from (26),  
170 the  $TTF$  value does not depend on the mass of fish. The  $TTF$  values calculated by using (26) are given in Table 2. Among  
the considered elements, only caesium ( $TTF > 1$ ) may be biomagnified in the food chain, in agreement with Kasamatsu and  
Ishikawa (1997), where it was found that the  $BAF$  of  $^{137}\text{Cs}$  increased with increasing trophic level. The concentrations of  
other radionuclides in Table 2 decreased with the increase of trophic level ( $TTF < 1$ ), which is consistent with the findings  
presented by Cardwell et al. (2013), where an inverse relationship was obtained between trophic levels and the concentration  
175 of inorganic metals in water chains.

Notice that values of  $AE_f$ ,  $AE_w$  and  $BAF$  do not depend on the fish mass. The literature data reveal diverse relationships  
between fish mass and both radionuclide bioconcentration ( $BCF$ ) and bioaccumulation ( $BAF$ ) factors. In particular, data of  
laboratory experiments (Mathews et al., 2008) showed that there is no significant relationship between bioconcentration factor  
 $BCF$  and fish size for most studied aqueous metals. The  $BAF$  in larger and older fish of the same species can differ from  
180 smaller and younger fish due to the change of habitat and diet with age (e.g. Kasamatsu and Ishikawa, 1997; Ishikawa et al.,  
1999; Kim et al., 2019) that does not contradict our findings, however, in this study we did not consider the change of prey  
preference along the fish growth.



### 3 Comparison with laboratory experiments

#### 3.1 Depuration of radionuclides after pulse-like feeding

185 Retention of absorbed elements in fish after single feeding was often used to estimate  $AE_f$  and depuration rate (e.g. Jeffree et al., 2006; Mathews and Fisher, 2008; Mathews et al., 2008; Pouil et al., 2017). According to Goldstein and Elwood (1971), single feeding can be approximated by a delta function  $\delta(t)$  at  $t = 0$  as

$$K_f m C_f = A_f \delta(t), \quad (27)$$

where  $A_f$  is the total amount of ingested activity. The solutions of equations (1)-(3) for activities  $A_2 = m_2 C_2$  and  $A_i = m_i C_i$  and for initial conditions  $A_2 = A_i = 0$  are


$$\frac{A_2}{A_f} = \frac{m_2 C_2}{A_f} = \exp(-(k_2 + \lambda_2)t), \quad (28)$$

$$\frac{A_i}{A_f} = \frac{m_i C_i}{A_f} = \frac{k_{2i}}{k_2 + \lambda_2 - \lambda_i} [\exp(-\lambda_i t) - \exp(-(k_2 + \lambda_2)t)]. \quad (29)$$

As follows from solutions (28)-(29), the decay of activity in the fish organisms includes a fast component with decay constant representing transfer of activity to the fish body and unabsorbed element egestion from the digestive tract, along with a slow component which is governed by elimination constants for  $i$  tissues. These solutions are generalized solutions of the equations of the sequentially linked two-compartment model by Goldstein and Elwood (1971), whereas in the D-DAT model (Vives i Batlle et al. 2008), an organism was represented by two boxes with parallel kinetics, also describing "slow" and "fast" exchange processes.

The solutions (28)-(29) can be compared with laboratory experiments in which depuration of metals from the fish after single feeding was studied. In the experiment by Mathews and Fisher (2008), the retention of several radioisotopes in juvenile sea bream (*Sparus auratus*) was considered. The average wet weight of the fish was 0.0001 kg. These fish were fed radiolabelled *Artemia salina nauplii*. The fish were allowed to feed for one hour, after which metal retention was observed in clean water over a 15-day period. The solutions with parameters corresponding to the fish mass and metal  $AE$  (Pouil et al., 2018) were compared with experimental data for  $^{134}\text{Cs}$ ,  $^{60}\text{Co}$ ,  $^{54}\text{Mn}$ , and  $^{65}\text{Zn}$  in Fig. 2-3. As seen in Fig. 23, both model and experiments showed two phases (fast and slow) of radionuclide elimination. Most of the activity is contained in muscle; however, the first 5 days after feeding the concentrations of  $^{60}\text{Co}$ ,  $^{54}\text{Mn}$ , and  $^{65}\text{Zn}$  in organs are much greater than in the muscle.

The solutions were also compared with laboratory experiments for predator fish (Mathews et al., 2008). In these experiments, the retention of several radioisotopes in sea bream (*Sparus auratus*), turbot (*Psetta maxima*) and spotted dogfish (*Scyliorhinus canicula*) was studied. Immature *S. auratus* (wet weight 0.012 kg), *P. maxima* (wet weight 0.027 kg), and *S. canicula* (wet weight 0.008 kg) were fed radiolabelled prey fish (juvenile *S. auratus*). After this feeding, fish were fed unlabelled prey fish for three weeks. In Fig. 34, the solutions (28)-(29) were compared with the laboratory experiments in which prey fish were labelled by  $^{134}\text{Cs}$ ,  $^{57}\text{Co}$ ,  $^{54}\text{Mn}$ , and  $^{65}\text{Zn}$ . The solution (28)-(29) and experiment agreed, demonstrating general dependence of the depuration process on fish mass. Differences between experimental data for different species may be due to differences in anatomy and physiology, as discussed by Jeffree et al. (2006) for *P. maxima* and *S. canicula*. The model, unlike the situation



**Figure 3.** Retention of radionuclides in whole body and tissues of juvenile sea bream (*Sparus auratus*). The simulations are compared with whole body measurements by Mathews and Fisher (2008).

215 for prey fish (Fig. 23), underestimates the total concentrations of  $^{57}\text{Co}$  and  $^{54}\text{Mn}$  in comparison with experiments, which is probably due to the neglect of other factors, except body weight, for the bioaccumulation kinetics. [Parameters of MCKA model for fish from experiments \(Mathews and Fisher, 2008; Mathews et al., 2008\) are given in Table S1, whereas Tables S2-S4 show dependence on radionuclides of the transfer rates  \$k\_{2,i}\$  in different fishes.](#)


### 3.2 Bioconcentration of dissolved radionuclides from sea water

220 Uptake and absorption in fish of elements from water were studied in several laboratory experiments (e.g. Jeffree et al., 2006; Mathews and Fisher, 2008; Mathews et al., 2008). The modelling of the absorption of elements can be used to estimate an assimilation efficiency  $AE_w$ . An analytical solution of equations (1) and (3) with initial conditions  $C_i = 0$  at  $t = 0$  is written as

$$\frac{\mu_1 C_1}{C_w} = \frac{K_w}{k_1 + \lambda_1} [1 - \exp(-(k_1 + \lambda_1)t)], \quad (30)$$

$$225 \frac{\mu_i C_i}{C_w} = \frac{k_{1i} K_w}{k_1 + \lambda_1 - \lambda_i} \frac{k_{1i} K_w}{(k_1 + \lambda_1) \lambda_i} \left[ 1 - \frac{k_1 + \lambda_1}{k_1 + \lambda_1 - \lambda_i} \exp(-\lambda_i t) + \frac{\lambda_i}{k_1 + \lambda_1 - \lambda_i} \exp(-(k_1 + \lambda_1)t) \right], \quad (i = \overline{3, 5}) \quad (31)$$

These solutions were compared with laboratory experiments for prey fish (Mathews and Fisher et al., 2008) and for predator fish (Jeffree et al., 2006). In the experiment by Mathews and Fisher et al. (2008), the uptake of several radioisotopes by juvenile *S. auratus* (wet weight 0.0002 kg) was studied during 25 days of exposure, whereas in experiments by Jeffree et al. (2006),



**Figure 4.** Retention of radionuclides in whole bodies of predator fish: sea bream (*Sparus auratus*), turbot (*Psetta maxima*) and spotted dogfish (*Scyliorhinus canicula*). The simulations are compared with whole body measurements by Mathews et al. (2008).

immature *P. maxima* (wet weight 0.0061 kg) and *S. canicula* (wet weight 0.0067 kg) were used for study during a 15-day period.


230 Parameters of MCKA model for these fishes are given in Table S5, whereas Tables S6-S8 show dependence on radionuclides of the transfer rates  $k_{1,i}$  in different fishes. The comparison of the analytical solution (30)-(31) with experimental data with respect to bioconcentration factor ( $BCF = C_{wb}/C_w$  [ $\text{l kg}^{-1}$ ]) is presented in Fig. 4-5. The values of  $AE_w$  were selected to approximate the experiment for small prey fish (Mathews, Fisher, et al., 2008). They differ for different metals. For  $^{134}\text{Cs}$ , the value of  $AE_w$  was 0.001, whereas for  $^{57}\text{Co}$ ,  $^{60}\text{Co}$ ,  $^{54}\text{Mn}$ , and  $^{65}\text{Zn}$ , these values were 0.0050, 0.0025, 0.0045, 0.0065,

235 respectively. This contrasted with  $AE_f$ , which is larger for  $^{134}\text{Cs}$  than for  $^{57}\text{Co}$ ,  $^{60}\text{Co}$ ,  $^{54}\text{Mn}$ , and  $^{65}\text{Zn}$  (Pouil et al., 2017). The estimated values of  $AE_w$  are presented in Table 2. The abovementioned values of  $AE_w$  were used to calculate  $BCF$  for larger predator fish in experiments by Jeffree et al. (2006). As seen in Fig. 45, values of  $AE_w$  for considered elements are of the order  $10^{-3}$ , which is in agreement with most models. However, comparison with larger fish highlighted some differences between species of fish, as discussed by Jeffree et al. (2006), and differences between model and experiment for a constant

240 value of  $AE_w$ . At the same time, it is known that dietary intake of metals dominates over water intake (Mathews and Fisher, 2009). Therefore, deviations in values of  $AE_w$  would not be significantly affected by the full uptake of elements from the marine environment.

The parameter  $\alpha_1$  can be estimated from the relations (13) and (16) according to the experimental data for equilibrium conditions. An average value of gill  $BCF_1$  is approximately  $10 \text{ l kg}^{-1}$  for  $^{58}\text{Co}$ ,  $^{54}\text{Mn}$ ,  $^{134}\text{Cs}$  and  $^{65}\text{Zn}$  (Jefferies and Hewitt, 1971;

245 Pentreath, 1973). Then for  $AE_w=0.001$  we obtained  $\alpha_1=$ 0.8-800. With the selected value of  $\alpha_1$ , the process of adaptation of



**Figure 5.** Simulated  $BCF$  in marine fish during the period of exposition in water. The simulations are compared with isotope measurements by Mathews and Fisher et al. (2008) in juvenile sea bream (*Sparus auratus*) and measurements by Jeffree et al. (2006) in turbot (*Psetta maxima*) and spotted dogfish (*Scyliorhinus canicula*).

the gill tissue to changes in the concentration of radioactivity in water is much faster than for other tissues of the fish. In addition, as follows from the solution (30)-(31) at  $\lambda_1 \gg \lambda_i$ , the contribution of gill contamination to the whole-body contamination is small.

### 3.3 Simplification of the model based on the results of analytical solutions and laboratory experiments

250 Comparison of the model with against laboratory experiments on the retention of absorbed elements in fish after single feeding has shown the importance of including in the model demonstrated the need to include the kinetic characteristics of the digestive tract compartment describing in the model when highly non-equilibrium transfer dynamics are expected. However, for modelling of food uptake in marine environment with multiple feedings the simple equilibrium assumption (14) can be used. At the same time, the analytical solution describing the bioconcentration due to the uptake and absorption in fish of elements from water, as well as the results of the laboratory experiment, show that the contribution to the gills (13) is negligible. Therefore, for modelling of uptake from water the equilibrium assumption can also be used as it shown above. The corresponding simplified equations for muscle, bone and organs can be rewritten as

$$\mu_i \frac{dC_i}{dt} = AE_{fi} K_f C_f + AE_{wi} K_w C_w - \lambda_i \mu_i C_i - \lambda_g \mu_i C_i. \quad (32)$$

The uncertainty in calculations using equations (32) arises due to (i) limited experimental data to define allometry constants  $\alpha_f$ ,  $\alpha_w$ ,  $\alpha_3$ ,  $\alpha_4$ ,  $\alpha_5$ ; (ii) large intervals in the values of known  $AE_f$  coefficients; (iii) unknown  $AE_f$  values for many radionuclides;

(iv) lack of experimental data about  $AE_w$ ; (v) limited experimental data for whole-body to tissue concentration ratios  $CR_i$  in the marine fish. Therefore, a sensitivity analysis is necessary to estimate uncertainty of the simulation results. We estimated the effects of variations in above parameters in the equation (24) on the value of  $BAF_{wb}$   $BAF$  in equilibrium state. The simple local sensitivity analysis and One-At-a-Time method (Pianosi et al., 2016) was used. The sensitivity of model output  
265 was estimated using a sensitivity index ( $SI$ ) calculated following Hamby (1994) as

$$SI = \frac{D_{max} - D_{min}}{D_{max}}, \quad (33)$$

where  $D_{max}$  and  $D_{min}$  are the outputs corresponded to maximal and minimal input parameter values, respectively. Similarly to Bezhenar et al. (2016), the range for every parameter was defined as follows: minimum value was set to half the reference value and maximum value was set to twice the reference value. Calculated  $SI$  for three radionuclides, which are preferably  
270 accumulated in different tissues:  $^{137}\text{Cs}$ ,  $^{90}\text{Sr}$ , and  $^{60}\text{Co}$ , are given in [Table-S2](#) [Fig. S1](#).


The results of sensitivity study suggest that model results are most sensitive to variations of  $AE_f$  and  $\alpha_f$  for  $^{137}\text{Cs}$  and  $^{60}\text{Co}$ , whereas for  $^{60}\text{Co}$  they are almost not sensitive to variations of  $AE_w$  and  $\alpha_w$ . Note that model results are also sensitive to the variations of parameters related to tissues where radionuclide is mainly accumulated:  $\alpha_3$  and  $CR_3$  for  $^{137}\text{Cs}$ ,  $\alpha_5$  and  $CR_5$  for  $^{60}\text{Co}$ . For  $^{90}\text{Sr}$  the model results are moderately sensitive to the variations of most above considered parameters.

## 275 4 Model applications

### 4.1 Modified ~~compartment~~ POSEIDON-R box model

In order to predict the accumulation of radionuclides in fish in the marine environment using the ~~multi-compartment kinetic allometric~~ ~~(MCKA)~~ MCKA model described above, it is necessary to calculate changes in concentration in water and in bottom sediments and to calculate the transport of radionuclides through food chains. The POSEIDON-R ~~compartment~~ box model (Lepicard et al., 2004; Maderich et al., 2014a,b; 2018b; Bezhenar et al., 2016) can be used to simulate the marine environment as a system  
280 of 3D ~~compartments~~ boxes for the water column, bottom sediment, and food web. The water column ~~compartment~~ box is vertically subdivided into layers. The suspended matter settles in the water column. The bottom sediment ~~compartment~~ box is divided into three layers (Fig. [S1](#) [S2](#)). The downward burial processes operate in all three sediment layers. Maderich et al. (2018b) described the POSEIDON-R ~~compartment~~ model in detail.

285 A food web model that includes pelagic and benthic food chains is implemented within the POSEIDON-R ~~compartment~~ box model (Bezhenar et al., 2016). In the food web model, marine organisms are grouped into classes according to trophic level and species type (Fig. [S2](#) [S3](#)). The food chains differ between the pelagic zone and the benthic zone. Pelagic organisms comprise primary producers (phytoplankton) and consumers (zooplankton, non-piscivorous (forage) fish, and piscivorous fish). In the benthic food chain, radionuclides are transferred from algae and contaminated bottom sediments to deposit-feeding inverte-  
290 brates, demersal fish, and benthic predators. Bottom sediments include both organic and inorganic components. Radioactivity is assumed to be assimilated by benthic organisms from the organic components of the bottom deposits. Other food web components are crustaceans (detritus feeders), molluscs (filter feeders), and coastal predators, which feed throughout the water




**Figure 6.** Compartment Box system around the Forsmark NPP. The numbers denote regional boxes in the box system of the Baltic Sea (Bezhenar et al., 2016). An additional “Inner box” is separated from regional box 68 by the blue line. The coastal box (red rectangle) surrounds the area, where cooling water from NPP is discharged.

column in shallow coastal waters. All organisms take in radionuclides both via the food web and directly from the water. [Table S9 in the Supplementary Material contains food preferences for organisms in the food web which are used in the model.](#) Details of the transfer of radiocaesium through the marine food web are presented by Bezhenar et al. (2016) and Maderich et al. (2018b).

[The POSEIDON-R model can handle different types of radioactive releases: including atmospheric fallout and point sources associated with routine releases from nuclear facilities located directly on the coast or point sources associated with accidental releases \(Lepicard et al., 2004\). For coastal discharges occurring in the large \('regional'\) boxes, 'coastal' release boxes are nested into the 'regional' box system. The intermediary boxes between 'coastal' and 'regional' boxes are called 'inner' boxes.](#)

#### 4.2 Release of radionuclides during normal operation of the Forsmark nuclear power plant

This section presents the simulation results of  $^{60}\text{Co}$  and  $^{54}\text{Mn}$  routine release into the marine environment from Forsmark NPP, located on the Baltic Sea coast of Sweden. The compartment POSEIDON-R model with embedded food web MCKA model, one-compartment fish model and bioaccumulation factor (equilibrium) model was customized for the Baltic Sea, as described by Bezhenar et al. (2016) and Maderich et al. (2018a). ~~A set of nested boxes~~ [The nested boxes \('inner' and 'coastal' boxes\)](#) inside the regional box no. 68 in the Baltic Sea box system ~~was-were~~ added to resolve the ~~near field of radionuclide~~



**Figure 7.** Release rates of <sup>60</sup>Co and <sup>54</sup>Mn according to measurements (Forsmark, 2014).

concentration-radionuclide concentration in the near field (Fig. 56). Parameters of the "innerbox" and "coastalbox" 'inner' and 'coastal' boxes are based on data from (SKBAquilonius, 2010). The main parameters of boxes with zooming in to the NPP are presented in Table S3-S10 in the Supplement. The simulation results for <sup>60</sup>Co and <sup>54</sup>Mn were compared with measurements for the smallest coastal box, where measurement data for bottom sediments and fish were available (Forsmark, 2014). The measurement data were compared with simulations for two species of fish: herring (*Clupea harengus membras*) as a non-piscivorous fish and pike (*Esox lucius*) as a coastal predator. There is no information on the mass of fish caught in the vicinity of Forsmark NPP. Therefore, we used estimates of the masses typical for prey and predatory fish, which are given in Table S4-S11. In the one-compartment model, two parameters must be prescribed: assimilation efficiency and biological half-life  $T_{0.5} = \ln 2 \lambda_{wb}^{-1} T_{0.5} = \ln(2) \lambda_{wb}^{-1}$ . Assimilation efficiency (see Table 2) was obtained from Pouil et al. (2018). Baudin et al. (1997) used  $T_{0.5} = 21$  d for <sup>60</sup>Co in the one-compartment model. The average value for  $T_{0.5}$  in predatory marine fish is 40 d (Beresford et al., 2015). Therefore, for <sup>60</sup>Co, we used  $T_{0.5} = 20$  d for prey fish and  $T_{0.5} = 40$  d for predatory fish. There are very limited data for  $T_{0.5}$  values in marine fish for <sup>54</sup>Mn. According to Beresford et al. (2015),  $T_{0.5}$  is in the range between 20 and 40 days. Therefore, we used the same values of  $T_{0.5}$  for <sup>54</sup>Mn, as for <sup>60</sup>Co.

The release rates According to Jeffrey et al. (2017), the uptake and depuration kinetics of <sup>60</sup>Co and <sup>54</sup>Mn for fish species in marine, brackish and freshwater environments are very similar. Therefore, we can apply the model parameters defined for marine environment (see Table S11) to reconstruct the herring and pike contamination by the above-mentioned radionuclides in the area near the Forsmark NPP with low salinity (3-5 PSU).

The release rate of <sup>60</sup>Co from the Forsmark NPP (Forsmark, 2014) are plotted in Fig. 6-7. As seen in Fig. 7a8a, the results of simulation for the concentration of <sup>60</sup>Co in the bottom sediments are in good agreement with the measurements

(Forsmark, 2014) in the coastal box for the wide range of employed values of sediment distribution coefficient  $K_d$ : from  $K_d = 3 \cdot 10^5 \div 2 \cdot 10^6 \text{ L kg}^{-1}$  for margin seas to  $K_d = 5 \cdot 10^7 \text{ L kg}^{-1}$  for open ocean (IAEA, 2004). The benthic food web (Bezhenar et al., 2016a), which describes the transfer of radioactivity from bottom sediments to deposit-feeding invertebrates and finally to fish, is quite important in this range of  $K_d$  values. The results from the POSEIDON-R calculations obtained with the MCKA model are compared (Fig. 7b8b,c) with measurements and results of calculations obtained with a one-compartment model and with an equilibrium approach using a standard BAF value (IAEA, 2004). Whereas one-compartment and equilibrium models underestimated the concentration of  $^{60}\text{Co}$  in fish, the MCKA model using generic parameters yields better agreement with measurements for both non-piscivorous fish (Fig. 7b8b) and coastal predator feeding by pelagic and benthic organisms in the coastal area (Fig. 7e8c).

Similarly, the behaviour of  $^{54}\text{Mn}$  in the marine environment near the Forsmark NPP is modelled. The release rate of  $^{54}\text{Mn}$  from the NPP is also plotted in Fig. 6-7 using data from Forsmark (2014). Comparison of the simulated concentration of  $^{54}\text{Mn}$  in bottom sediments with measurements (Forsmark, 2014) for the Forsmark coastal box is given in Fig. 8a8d. Good agreement was obtained when a standard value of  $K_d = 2 \cdot 10^6 \text{ L kg}^{-1}$  for  $^{54}\text{Mn}$  in margin seas (IAEA, 2004) was used. This means that, as in the case of  $^{60}\text{Co}$ , a significant fraction of radionuclide is deposited at the bottom, and the benthic food web should be considered. Similarly to the  $^{60}\text{Co}$  case, obtained results of simulation are also compared with results obtained using the one-compartment model and equilibrium approach. Again, the MCKA model yields the best agreement with measurements for both non-piscivorous fish (Fig. 8b8e) and coastal predators (Fig. 8e8f); however, this agreement is slightly worse than in the  $^{60}\text{Co}$  case. Notice that the BAF in the equilibrium approach can be locally estimated using *a posteriori* data. However, the MCKA model provided good agreement with measurements using only *a priori* information, which is important in the case of accidents, as considered in the next section.

### 4.3 Accumulation of $^{90}\text{Sr}$ in the fish after the Fukushima Dai-ichi accident


Following caesium,  $^{90}\text{Sr}$  is the second most important radiologically long-lived radionuclide released as a result of the FDNPP accident. It is highly soluble in water and exhibits relatively high ability for assimilation by marine organisms due to similar chemical properties with calcium. The atmospheric deposition of  $^{90}\text{Sr}$  is usually not taken into account due to its low volatility. Most of the  $^{90}\text{Sr}$  released from the FDNPP was directly released to the ocean, with estimates of total inventory in the range from 0.04 to 1.0 PBq (Buesseler et al., 2017). Here, we extended the simulation by Maderich et al. (2014b) of transfer and fate of  $^{90}\text{Sr}$  resulting from the FDNPP accident using the POSEIDON-R model complemented by the food web model (Bezhenar et al., 2016) and MCKA fish model.

The POSEIDON-R model was customized for the Northwestern Pacific and adjacent seas (the East China Sea, the Yellow Sea and the East/Japan Sea) as in (Maderich et al., 2018a). A total of 188 ~~compartments~~ boxes covered this region. The ~~compartments~~ boxes around the FDNPP are shown on Fig. 9 with an additional 4x4 km coastal box in the vicinity of FDNPP. The coastal box was included for the model validation with  $^{90}\text{Sr}$  measurements conducted in this specific area. Details of customization are given in Maderich et al. (2014a,b; 2018a).



Comparison between calculated and measured  $^{60}\text{Co}$  (Forsmark, 2014) concentrations in bottom sediment (a), non-piscivorous fish (herring)


(b) and coastal predator fish (pike) for the coastal box.



**Figure 8.** Comparison between calculated and measured  $^{54}\text{Mn}$ - $^{60}\text{Co}$  (Forsmark, 2014) concentrations in bottom sediment (a), non-piscivorous fish (herring) (b) and coastal predator fish (pike) (c) and  $^{54}\text{Mn}$  concentrations in bottom sediment (d), non-piscivorous fish (herring) (e) and coastal predator fish (pike) (f) for the coastal box.

The historical contamination due to global atmospheric deposition in the period from 1945-2010 was simulated according to (Maderich et al., 2014b) with data from the MARiS database (2020). The value of the accidental release was estimated as 160 TBq (16 TBq day<sup>-1</sup> during 10 days), that was consistent with the range reported by Buesseler et al. (2017). In the post-accidental period, the continuous leakage of  $^{90}\text{Sr}$  due to groundwater transport of radioactivity from the NPP site was monitored (Castrillejo et al., 2016). Therefore, in the simulation the conservative scenario was used for release of  $^{90}\text{Sr}$  in the post-accidental period (Fig. 9); the release of  $^{90}\text{Sr}$  was assumed equal to  $^{137}\text{Cs}$  release (Maderich et al. (2018a). Comparison between calculated and measured  $^{90}\text{Sr}$  concentrations in water (a), bottom sediment (b), and piscivorous fish (c) for the coastal box. Comparison between calculated and measured  $^{90}\text{Sr}$  concentrations in water (a), bottom sediment (b), and piscivorous fish (c) for box no. 173.


Comparison between calculated and measured  $^{90}\text{Sr}$  concentrations in water, bottom sediment, and piscivorous fish for the coastal box and box no. 173 are shown in Figs. 10 and 11, respectively. Fig. 10. Measured concentrations of  $^{90}\text{Sr}$  in the water and bottom sediments before the accident were obtained from the MEXT database (MEXT, 2020). Concentrations of  $^{90}\text{Sr}$  after the accident at TEPCO (Tokyo Electric Power Company) sampling points near the discharging canals (T1 and T2) and



**Figure 9.** (a) Boxes along the eastern coast of Japan with fine resolution in the area around the FDNPP (Maderich et al., 2018a). Deep boxes with 3 vertical layers in the water column are coloured by blue, boxes with 2 vertical layers are marked by vertical stripes, and shallow one-layer boxes are white. (b) Release rates of  $^{90}\text{Sr}$  in the accidental (bold line) and post-accidental (dashed line) periods.

at different distances offshore (TD-5 inside the coastal box area, T7, TD-1 and TD-9 for outer box no. 173) are available in the NRA (Nuclear Regulation Authority) database (NRA, 2020). The results of simulation show that the concentration of  $^{90}\text{Sr}$  in the seawater reaches the maximum just after the accidental release (Figs. 10a, [Had](#)). Notice that the box model gives the average concentrations for each box, which means that local concentrations may be larger or smaller than the average (Fig. 10).  
 375 Especially large differences can occur during the accidental release under strongly non-equilibrium conditions. Further large dispersion of measured concentrations shows that non-equilibrium conditions remained for a long time in the area close to NPP (Fig. 10a). The agreement between calculated and measured concentrations (Figs. 10a, [Had](#)) could be a confirmation of the correctness of estimation of the source term. This is also confirmed by agreement of measured and simulated concentrations  
 380 of  $^{90}\text{Sr}$  in the bottom sediment (Fig. 10b, [e](#)). Notice that there is no measurement data for bottom sediment in box no. 173 (Fig. [Hb](#))-[10e](#)).

The calculated concentration of  $^{90}\text{Sr}$  in piscivorous fish was compared with measurement data for fat greenling (*Hexagrammos otakii*) before the accident (JCAC, 2020) and with data from (NRA, 2020) and Miki et al. (2017) after the accident. The NRA data for  $^{90}\text{Sr}$  are very limited. Therefore, different species of piscivorous fish were considered, such as sharks (*Triakis scyllium*, *Squatina japonica*), rockfish (*Sebastes cheni*) and seabass (*Lateolabrax japonicus*). [Parameters of MCKA model for](#)




**Figure 10.** Comparison between calculated and measured  $^{90}\text{Sr}$  concentrations in water (a), bottom sediment (b), piscivorous fish (c) for the coastal box and in water (d), bottom sediment (e), and piscivorous fish (f) for box no. 173.

these fishes are given in Table S11. The simulation results with the MCKA model agree well with the experimental observations (Fig. 10c, Hef), while the target tissue (TT) approach underestimates the concentration of  $^{90}\text{Sr}$  in the fish.

390 Comparison of the equations (1)-(3) and (7) of the MCKA model and the equation (11) of the standard whole-body model demonstrates that the main difference is found in the description of the whole-body elimination rate  $\lambda_{wb}$ . Whereas in the whole-body model the value of  $\lambda_{wb}$  is constant, in the MCKA model it is the ratio of activity weighted tissue elimination rates to whole-body activity

$$\lambda_{wb} = \frac{\sum_{i=3}^5 \mu_i \lambda_i C_i}{C_{wb}}. \quad (34)$$

395 Therefore, in the MCKA model, the value of  $\lambda_{wb}$  can vary over time, depending on the uptake of radionuclide and the tissue elimination rates. The time variation of  $\lambda_{wb}$  computed from (34) is shown in Fig. 11 for three different cases: (a) pulse-like feeding experiment (Mathews and Fisher, 2008), (b) release of  $^{60}\text{Co}$  during normal operation of the Forsmark NPP and (c) accumulation of  $^{90}\text{Sr}$  in the fish due to the FDNPP accident. As seen in the plots,  $\lambda_{wb}$  varies considerably when there is non-equilibrium, such as in the case of a pulse-like feeding or an accident. Even in case of a routine release,  $\lambda_{wb}$  follows any changes in the release rate (Fig. 11b). In case of the FDNPP accident, the calculated  $\lambda_{wb}$  shows some tendency towards an equilibrium value, but after a pulse-like release of  $^{90}\text{Sr}$  in 2011  $\lambda_{wb}$  doubled following the release of activity and then slowly



**Figure 11.** The calculated  $\lambda_{wb}$  for three scenarios: (a) the pulse-like feeding experiment (Mathews, Fisher (2008)), (b) the release of  $^{60}\text{Co}$  during normal operation of the Forsmark NPP and (c) the accumulation of  $^{90}\text{Sr}$  in the fish due to the FDNPP accident.

400 converged to the quasi-equilibrium state governed by the global deposition. Notice that in this case we extended simulation  
 period to 2040 extrapolating deposition data and FDNPP release data in Fig. 9b. Therefore, whole-body model with a constant  
 $\lambda_{wb}$ , that is calibrated using observational data, cannot to correctly describe such transient processes in the organism. This is  
 confirmed in the Fig. 10 by comparing the results from the MCKA model and the one-compartment model. Here the  $AE$  value  
 in both two models are the same, whereas the equilibrium value  $\lambda_{wb}$  was calculated in the MCKA model using value before  
 405 2011 for piscivorous fish ( $\lambda_{wb} = 0.0027 \text{ d}^{-1}$ ). The one-compartment model simulation results using parameters from MCKA  
 model are close to the MCKA model results at initial stage of accidental release. However, over time, the concentration  $^{90}\text{Sr}$   
 in fish tends to equilibrium faster than the MCKA model predicts, which is explained by the time-dependent behavior of  $\lambda_{wb}$   
 in the MCKA model. These results suggest also that the MCKA model could be effectively used to find equivalent parameters  
 for the one-compartment model as was done here.

## 410 5 Conclusions

A new approach to predicting the accumulation of radionuclides in fish taking into account heterogeneity of distribution of con-  
 tamination in the organism and dependence of metabolic process rates on the fish mass was developed. The fish organism was  
 represented by compartments for three groups of tissues/organs (muscle, bone, organs) and two input compartments represent-  
 ing gills and digestive tract. The absorbed elements are redistributed between organs/tissues and then eliminated according to  
 415 their metabolic function. The food and water uptake rates, elimination rate and growth rate depend on the metabolic rate, which  
 is scaled by the fish mass to the 3/4 power, but do not depend on the radionuclide. At the same time, the activity is distributed  
 between the different tissues and organs according to the tissue assimilation efficiencies, which differ per radionuclide (Table  
 2), but that do not depend on fish mass. Therefore, the transfer rates can be associated with specific radionuclide and fish mass  
 as shown e.g. in Tables S2-S4 and S6-S8. The position of the fish species in the trophic level also affects the concentration of  
 420 activity in the organism.

This model is of intermediate complexity and provides an alternative for the basic/simplistic whole-body models and the highly advanced PBPK models. The main difference between MCKA and whole-body models was found in the description of the whole-body elimination rate  $\lambda_{wb}$ . Whereas in the whole-body model the value of  $\lambda_{wb}$  is constant, in the MCKA model it is the ratio of activity weighted tissue elimination rates to whole-body activity as described (34). The elimination rate  $\lambda_{wb}$  varies considerably in non-equilibrium state of fish, such as in the case of a pulse-like feeding or an accident.

The trophic transfer factors ( $TTF$ ) were calculated for 9-5 elements using assimilation efficiencies  $AE_f$  obtained from laboratory data. Among considered elements, only caesium (Cs) ~~may biologically magnify when transferring upwards into the level may elevate in the predator fish of the~~ food chain ( $TTF > 1$ ). This is in agreement with measurements. The concentrations of other elements (Sr, Co, Mn, Zn, ~~Ag, Cu, Cd, and Cr~~) decrease with the increase of trophic level ( $TTF < 1$ ). The kinetics of the assimilation and elimination of  $^{134}\text{Cs}$ ,  $^{57}\text{Co}$ ,  $^{60}\text{Co}$ ,  $^{54}\text{Mn}$  and  $^{65}\text{Zn}$ , which are preferably accumulated in different tissues, were analyzed using the analytical solutions of a system of model equations. These solutions exhibited good agreement with the laboratory experiments for the depuration process after single feeding of fish with radiolabelled prey and with respect to uptake of activity from water. Notice that for relatively slow processes in the marine environment, transfer processes in the gills and digestive tract can be close to equilibrium, which allows consideration of only a three-compartment (muscle, bone, organs) version of the model.

The developed MCKA model was embedded into the ~~compartment-box~~ model POSEIDON-R, which describes the transfer of radionuclides through the pelagic and benthic food webs. The POSEIDON-R model was applied for the simulation of the transport and fate of  $^{60}\text{Co}$  and  $^{54}\text{Mn}$  routinely released from Forsmark NPP located on the Baltic Sea coast of Sweden and for calculation of  $^{90}\text{Sr}$  concentration in fish after the accident at Fukushima Dai-ichi NPP. Predicted concentrations of radionuclides in fish agreed well with the measurements in both case studies. It is shown that the MCKA model with the defined generic parameters could be used in different marine environments without calibration based on *a posteriori* information, which is important for emergency decision support systems (Periáñez et al., 2019).

*Author contributions.* RB, VM and KJ conducted the literature review and designed the study. RB, VM and KJ developed the model and performed modelling of laboratory experiments. GdV and KK collected data for case studies. RB, VM and KK performed the simulations for case studies. RB, VM, KJ and GdV analysed results of simulations and wrote the initial manuscript. All authors edited and approved the final manuscript text

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This work was partially supported by KIOST major project PE99812, National Research Foundation of Ukraine projects nos. 2020.02/0048 and 2020.01/0421, and IAEA Coordinated Research Project K41017. Authors are grateful to three anonymous reviewers for useful suggestions that helped to improve the manuscript.

## References

- Alava, J. and Gobas, F.: Modeling  $^{137}\text{Cs}$  bioaccumulation in the salmon-resident killer whale food web of the Northeastern Pacific following the Fukushima Nuclear Accident, *Science of the Total Environment*, 544, 56-67, 2016.
- Andersen, N. G.: Depletion rates of gastrointestinal content in common goby (*Pomatoschistus microps* (Kr.)). Effects of temperature and fish size, *Dana*, 3, 31-42, 1984.
- [Aquiloni, K.: The marine ecosystems at Forsmark and Laxemar-Simpevarp. SR-Site Biosphere. Sweden. SKB, 495. pp., 2010.](#)
- Barron, M. G., Stehly, G. R. and Hayton, W.L.: Pharmacokinetic modeling in aquatic animals. Models and concepts, *Aquatic Toxicology* 17, 187-212, 1990.
- Baudin, J. P., Veran, M. P., Adam, C., and Garnier-Laplace, J.: Co-60 transfer from water to the rainbow trout (*Oncorhynchus mykiss* Walbaum). *Archives of Environmental Contamination and Toxicology* 33, 230-237, 1997.
- Belharet, M., Charmasson, S., Tsumune, D., Arnaud, M. and Estournel, C.: Numerical modeling of  $^{137}\text{Cs}$  content in the pelagic species of the Japanese Pacific coast following the Fukushima Dai-ichi Nuclear Power Plant accident using a size structured food-web model. *PLoS ONE* 14(3), e0212616, 2019. <https://doi.org/10.1371/journal.pone.0212616>
- Beresford, N.A., Beaugelin-Seiller, K., Burgos, J., Cujic, M., Fesenko, S., Kryshev, A., Pachal, N., Real, A., Su, B.S., Tagami, K., Vives i Batlle, J., Vives-Lynch, S., Wells, C. and Wood M.D.: Radionuclide biological half-life values for terrestrial and aquatic wildlife. *J. Environ. Radioactivity* 150, 270-276, 2015.
- Beresford, N.A., Wood, M.D., Vives i Batlle, J., Yankovich, T.L., Bradshaw, C. and Willey, N.: Making the most of what we have: application of extrapolation approaches in radioecological wildlife transfer models. *Journal of Environmental Radioactivity* 151, 373-386, 2016.
- Bezhenar, R., Jung, K.T., Maderich, V., Willemsen, S., de With, G. and Qiao, F.: Transfer of radiocaesium from contaminated bottom sediments to marine organisms through benthic food chain in post-Fukushima and post-Chernobyl periods. *Biogeosciences* 13, 3021-3034, 2016.
- Buesseler, K., Dai, M., Aoyama, M., Benitez-Nelson, C., Charmasson, S., Higley, K., Maderich, V., Masque, P., Morris, P. J., Oughton, D. and Smith, J. N.: Fukushima Daiichi-derived radionuclides in the Ocean: transport, fate, and impacts. *Annual Reviews of Marine Sciences*, 9, 173-203, 2017. <https://doi.org/10.1146/annurev-marine-010816-060733>.
- Cardwell, R.D., DeForest, D.K., Brix, K.V. and Adams, W.J.: Do Cd, Cu, Ni, Pb and Zn biomagnify in aquatic ecosystems? *Reviews of Environmental Contamination and Toxicology* 226, 101-122, 2004.
- Carvalho, F.P.: Radionuclide concentration processes in marine organisms: A comprehensive review, *J. Environ. Radioactivity*, 186, 124-130, 2018.
- Castrillejo, M., Casacuberta, N., Breier, C.F., Pike, S.M., Masque, P. and Buesseler, K.O.: Reassessment of  $^{90}\text{Sr}$ ,  $^{137}\text{Cs}$  and  $^{134}\text{Cs}$  in the coast off Japan derived from the Fukushima Dai-ichi Nuclear Accident. *Environmental Science and Technology*, 50, 173-180, 2016.
- Coughtrey, P.J. and Thorne, M.C.: Radionuclide distribution and transport in terrestrial and aquatic ecosystems: A critical review of data, vol 2, A. A. Balkema, Rotterdam, 1983.
- Forsmark: Forsmark omgivningsdata utdrag ur databas 20140113.xlsx, 2014.
- Fowler, S.W. and Fisher, N.S. Radionuclides in the biosphere. In: H. D. Livingston (Ed.) Ch. 6. Marine Radioactivity. *Radioactivity in environment*, v.6. Elsevier. 167-203, 2004.
- Garnier-Laplace, J., Adam, C., Lathuilliere, T., Baudin, J. and Clabaut, M.: A simple fish physiological model for radioecologists exemplified for  $^{54}\text{Mn}$  direct transfer and rainbow trout (*Oncorhynchus mykiss* W.). *J. Environ. Radioactivity*, 49, 35-53, 2006.

- Goldstein, R. A. and Elwood, J.W.: A Two-compartment, three-parameter model for the absorption and retention of ingested elements by animals. *Ecology*, 52, 935-939, 1971.
- 490 [Grech, A., Tebb, C., Brochot, C., Bois, F.Y., Bado-Nilles, A., Dorne, J.L., Quignot, N., Beaudouin, R.: Generic physiologically-based toxicokinetic modelling for fish: Integration of environmental factors and species variability. \*Science of The Total Environment\*, 651, 516-531, 2019.](#)
- Hamby, D. M.: A review of techniques for parameter sensitivity analysis of environmental models, *Environ. Monit. Assess.*, 32, 135-154, 1994.
- 495 Higley, K.A. and Bytwerk, D.P.: Generic approaches to transfer. *Journal of Environmental Radioactivity*, 98, 4-23, 2007.
- Heling, R., Koziy, L. and Bulgakov, V.: On the dynamical uptake model developed for the uptake of radionuclides in marine organisms for the POSEIDON-R model system. *Radioprotection* 37, C1, 833-838, 2002.
- Heling, R., and Bezhenar, R.: Modification of the dynamic radionuclide uptake model BURN by salinity driven transfer parameters for the marine food web and its integration in POSEIDON-R. *Radioprotection*, 44, 741-746, 2009.
- 500 IAEA (International Atomic Energy Agency): Sediment distribution coefficients and concentration factors for biota in the marine environment, Technical Report Series No 422, IAEA, Vienna, Austria, 2004.
- Ishikawa, Y., Yamada, K., Nonaka, N., Marumo, K. and Ueda, T.: Size-dependent concentrations of radiocesium and stable elements in muscles of flathead flounder *Hippoglossoides dubius*. *Fish Sci (Tokyo)*, 61:981-985, ~~1999~~-1995.
- Jefferies, D. F. and Hewitt, C. J.: The accumulation and excretion of radioactive caesium by the plaice and thornback ray. *J. mar. biol. Ass. UK*, 51, 411-422, 1971.
- 505 Jeffrey, R.A., Warnau, M., Teyssie, J.-L. and Markich, S.J.: Comparison of the bioaccumulation from seawater and depuration of heavy metals and radionuclides in the spotted dogfish *Scyliorhinus canicula* (Chondrichthys) and the turbot *Psetta maxima* (Actinopterygii: Teleostei). *Sci. Total Environ.* 368, 839-852, 2006.
- Jeffrey, R. A., Markich, S., Oberhansli, F. and Teyssie, J.L.: Radionuclide biokinetics in the Russian sturgeon and phylogenetic consistencies with cartilaginous and bony marine fishes. *J. Environ. Radioactivity*, 202, 25-31, 2017.
- 510 JCAC (Japan Chemical Analysis Center): Environmental Radioactivity Database. Available at <https://www.jcac.or.jp/>, last accessed: 26 October 2020.
- Kim, S.H., Lee, H., Lee, S.H. and Kim, I.: Distribution and accumulation of artificial radionuclides in marine products around Korean Peninsula. *Marine Pollution Bulletin*, 146, 521-531, 2019.
- 515 Kasamatsu, F. and Ishikawa, Y.: Natural variation of radionuclide <sup>137</sup>Cs concentration in marine organisms with special reference to the effect of food habits and trophic level. *Marine Ecology Progress Series*. 160,109-120, 1997.
- Lepicard, S., Heling, R. and Maderich, V.: POSEIDON-R/RODOS models for radiological assessment of marine environment after accidental releases: application to coastal areas of the Baltic, Black and North Seas, *J. Environ. Radioactivity*, 72, 153-161, 2004.
- Maderich, V., Bezhenar, R., Heling, R., De With, G., Jung, K.T., Myoung, J.G., Cho, Y.-K., Qiao, F. and Robertson, L.: Regional long-term model of radioactivity dispersion and fate in the Northwestern Pacific and adjacent seas: application to the Fukushima Dai-ichi accident. *J. Environ. Radioactivity*, 131, 4-18, 2014a.
- 520 Maderich, V., Jung, K.T., Bezhenar, R., de With, G., Qiao, F., Casacuberta, N., Masque, P. and Kim, Y.H.: Dispersion and fate of <sup>90</sup>Sr in the Northwestern Pacific and adjacent seas: global fallout and the Fukushima Dai-ichi accident, *Sci. Total Environ.*, 494-495, 261-271, 2014b.

- 525 Maderich, V., Bezhenar, R., Tateda, Y., Aoyama, M. and Tsumune, D.: Similarities and differences of  $^{137}\text{Cs}$  distributions in the marine environments of the Baltic and Black seas and off the Fukushima Dai-ichi nuclear power plant in model assessments, *Marine Pollution Bulletin*, 135, 895-906, 2018a.
- Maderich, V., Bezhenar, R., Tateda, Y., Aoyama, M., Tsumune, D., Jung, K. T. and de With, G.: The POSEIDON-R compartment model for the prediction of transport and fate of radionuclides in the marine environment. *MethodsX*, 5, 1251-1266, 2018b.
- MARIS (Marine Information System) (2020): Data available at <http://maris.iaea.org/>, last accessed: 26 October 2020.
- 530 Mathews, T. and Fisher, N.S.: Trophic transfer of seven trace metals in a four-step marine food chain. *Marine Ecology Progress Series* 367, 23-33, 2008.
- Mathews, T., Fisher, N.S., Jeffree, R.A. and Teysse, J.-L.: Assimilation and retention of metals in teleost and elasmobranch fishes following dietary exposure. *Marine Ecology Progress Series* 360, 1-12, 2008.
- Mathews, T. and Fisher, N.S.: Dominance of dietary intake of metals in marine elasmobranch and teleost fish. *Sci. Total Environ.* 407, 5156-5161, 2009.
- 535 MEXT (Japanese Ministry of Education, Culture, Sports, Science and Technology): Environmental radiation database. Available at <http://search.kankyo-hoshano.go.jp/servlet/search.top>, last accessed: 26 October 2020.
- Miki, S., Fujimoto, K., Shigenobu, Y., Ambe, D., Kaeriyama, H., Takagi, K., Ono, T., Watanabe, T., Sugisaki, H. and Morita, T.: Concentrations of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}/^{90}\text{Sr}$  activity ratios in marine fishes after the Fukushima Dai-ichi Nuclear Power Plant accident. *Fisheries Oceanography*, 26, 221-233, 2017.
- 540 NRA (Monitoring information of environmental radioactivity level, Japan): Monitoring information of environmental radioactivity level. <http://radioactivity.nsr.go.jp/en/list/205/list-1.html>, last accessed: 26 October 2020.
- Otero-Muras, I., Franco-Uria, A., Alonso, A.A. and Balsa-Canto, E.: Dynamic multi-compartmental modelling of metal bioaccumulation in fish: Identifiability implications. *Environmental Modelling and Software*, 25, 344-353, 2010.
- 545 Pentreath, R. J.: The accumulation from sea water of  $^{65}\text{Zn}$ ,  $^{54}\text{Mn}$ ,  $^{58}\text{Co}$ , and  $^{59}\text{Fe}$ , by the thornback ray, *Raja clavata l.* *J. exp. mar. Biol. Ecol.*, 12, 327-334, 1973.
- Periáñez, R., Bezhenar, R., Brovchenko, I., Duffa, C., Iosjpe, M., Jung, K.T., Kim, K.O., Kobayashi, T., Liptak, L., Little, A., Maderich, V., McGinnity, P., Min, B.I., Nies, H., Osvath, I., Suh, K.S. and de With, G.: Marine radionuclide transport modelling: Recent developments, problems and challenges. *Environmental Modelling and Software*, 122, 104523, 2019. <https://doi.org/10.1016/j.envsoft.2019.104523>.
- 550 Pianosi, F., Beven, K., Freer, J., Hall, J. W., Rougier, J., Stephenson, D. B. and Wagener, T.: Sensitivity analysis of environmental models: A systematic review with practical workflow. *Environmental Modelling & Software*, 79, 214-232, 2016.
- Pouil, S., Warnau, M., Oberhansli, F., Teysse, J.-L., Bustamante, P. and Metian, M.: Comparing single-feeding and multi-feeding approaches for experimentally assessing trophic transfer of metals in fish. *Environmental Toxicology and Chemistry*, 36, 1227-1234, 2017.
- Pouil, S., Bustamante, P., Warnau, M. and Metian, M.: Overview of trace element trophic transfer in fish through the concept of assimilation efficiency. *Marine Ecology Progress Series* 588, 243-254, 2018.
- 555 Reinfelder, J. R., Fisher, N. S., Luoma, S. N., Nichols, J. W. and Wang, W.-X.: Trace element trophic transfer in aquatic organisms: a critique of the kinetic model approach. *Sci Total Environ* 219:117-135, 1998.
- Rouleau, C., Tjalve, H., Gottofrey, J. and Pelletier, E.: Uptake, distribution and elimination of  $^{54}\text{Mn(II)}$  in the brown trout (*Salmo Trutta*). *Environmental Toxicology and Chemistry*. 14, 483-490, 1995.
- 560 ~~SKB (Swedish Nuclear Fuel and Waste Management Company): The marine ecosystems at Forsmark and Laxemar-Simpevarp. Report TR-10-03, SKB, 495 p., 2010.~~



- Takata, H., Johansen, M.P., Kusakabe, M., Ikenoue, T., Yokota, M. and Takaku, H.: A 30-year record reveals re-equilibration rates of  $^{137}\text{Cs}$  in marine biota after the Fukushima Dai-ichi nuclear power plant accident: Concentration ratios in pre-and post-event conditions. *Science of the Total Environment*. 675, 694-704, 2019.
- 565 Tateda, Y., Tsumune, D. and Tsubono, T.: Simulation of radioactive cesium transfer in the southern Fukushima coastal biota using a dynamic food chain transfer model. *J. Environ. Radioactivity* 124, 1-12, 2013.
- Thomann, R.V., Shkreli, F. and Harrison, S.: A pharmacokinetic model of cadmium in rainbow trout. *Environmental Toxicology and Chemistry*. 16, 2268-2274, 1997.
- Thomas, D. M. and Fisher, N. S.: Evaluation of body size and temperature on  $^{137}\text{Cs}$  uptake in marine animals. *J. Environ. Radioactivity* 202, 570 25-31, 2019.
- Vives i Batlle, J. , Wilson, R.C. and McDonald, P.: Allometric methodology for the calculation of biokinetic parameters for marine biota, *Sci. Total Environ.*, 388, 256-269, 2007.
- Vives Batlle, J., Wilson, R.C., Watts, S.J., Jones, S.R., McDonald, P. and Vives-Lynch, S.: Dynamic model for the assessment of radiological exposure to marine biota. *J. Environ. Radioactivity* 99, 1711-1730, 2008.
- 575 Vives i Batlle, J.: Radioactivity in the Marine Environment. *Encyclopedia of Sustainability Science and Technology*. R.A. Meyers (eds). Springer, New York, 8387-8425, 2012.
- Vives i Batlle, J., Beresford, N.A., Beaugelin-Seiller, K., Bezhenar, R., Brown, J., Cheng, J.-J., Čujić, M., Dragović, S., Duffa, C., Fiévet, B., Hosseini, A., Jung, K.T., Kamboj, S., Keum, D.-K., Kobayashi, T., Kryshev, A., LePoire, D., Maderich, V., Min, B.-I., Perriñez, R., Sazykina, T., Suh, K.-S., Yu, C., Wang, C. and Heling, R.: Inter-comparison of dynamic models for radionuclide transfer to marine biota 580 in a Fukushima accident scenario. *J. Environ. Radioactivity*, 153, 31-50, 2016.
- West, G.B., Brown, J.H. and Enquist, B.J.: A general model for the origin of allometric scaling laws in biology. *Science* 276, 122-126, 1997.
- [Yankovich T.L.: Towards an improved ability to estimate internal dose to non-human biota: development of conceptual models for reference non-human biota. Proceedings of third international symposium on the protection of the environment from ionising radiation. IAEA, Vienna, Austria, 365-373, 2003.](#)
- 585 Yankovich, T., Beresford, N., Wood, M., Aono, T., Anderson, P., Barnett, C.L., Bennett, P., Brown, J.E., Fesenko, S., Fesenko, J., Hosseini, A., Howard, B.J., Johansen, P., Phaneuf, M.M., Tagami, K., Takata, H., Twining, J.R. and Uchida, S.: Whole-body to tissue-specific concentration ratios for use in biota dose assessments for animals, *Radiation Environ. Biophysics*, 49, 549-565, 2010.