

# Recent extreme drought events in the Amazon rainforest: Assessment of different precipitation and evapotranspiration datasets, and drought indicators

Phillip Papastefanou<sup>1</sup>, Christian S. Zang<sup>2</sup>, Zlatan Angelov<sup>1</sup>, Aline Anderson de Castro<sup>3</sup>, Juan Carlos Jimenez<sup>4</sup>, Luiz Felipe Campos De Rezende<sup>3</sup>, Romina Ruscica<sup>3,6,7</sup>, Boris Sakschewski<sup>8</sup>, Anna Sörensson<sup>5,6,7</sup>, Kirsten Thonicke<sup>8</sup>, Carolina Vera<sup>5,6,7</sup>, Nicolas Viovy<sup>9</sup>, Celso Von Randow<sup>3</sup> and Anja Rammig<sup>1</sup>

<sup>1</sup> Technical University of Munich, TUM School of Life Sciences Weihenstephan, Freising, Germany

<sup>2</sup> University of Applied Sciences Weihenstephan-Triesdorf, Department of Forestry, Freising, Germany

<sup>3</sup> Earth System Sciences Centre, National Institute for Spatial Research, São José dos Campos, São Paulo, Brazil

<sup>4</sup> GCU/IPL, University of Valencia, Valencia, Spain.

<sup>5</sup> Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Ciencias de la Atmósfera y los Océanos. Buenos Aires, Argentina.

<sup>6</sup> CONICET – Universidad de Buenos Aires. Centro de Investigaciones del Mar y la Atmósfera (CIMA). Buenos Aires, Argentina.

<sup>7</sup> CNRS – IRD – CONICET – UBA. Instituto Franco-Argentino para el Estudio del Clima y sus Impactos (UMI 3351 IFAECI). Centro de Investigaciones del Mar y la Atmósfera (CIMA). Buenos Aires, Argentina.

<sup>8</sup> Potsdam Institute for Climate Impact Research (PIK), Telegraphenberg A31, Potsdam, 14473, Germany

<sup>9</sup> LSCE, CEA-CNRS-Univ Paris-Saclay, Saclay, France

Correspondence to: Phillip Papastefanou (papa@tum.de)

Style Definition	... [22]
Style Definition	... [21]
Style Definition	... [20]
Style Definition	... [19]
Style Definition	... [18]
Style Definition	... [17]
Style Definition	... [16]
Style Definition	... [15]
Style Definition	... [14]
Style Definition	... [13]
Style Definition	... [12]
Style Definition	... [11]
Style Definition	... [10]
Style Definition	... [9]
Style Definition	... [8]
Style Definition	... [7]
Style Definition	... [6]
Style Definition	... [5]
Style Definition	... [4]
Style Definition	... [3]
Style Definition	... [2]
Style Definition	... [1]
Deleted: Quantifying the spatial extent and intensity of recent	
Formatted	... [24]
Formatted	... [23]
Formatted	... [25]
Deleted: their impacts on the carbon cycle	
Formatted	... [26]
Deleted: Zang <sup>1</sup> ...ang <sup>2</sup> , Zlatan Angelov <sup>1</sup> , Aline Anderson d	... [27]
Formatted	... [28]
Deleted: Sakschewski <sup>7</sup> ...akschewski <sup>8</sup> , Anna Sörensson <sup>4,5</sup>	... [29]
Formatted	... [30]
Deleted: Thonicke <sup>7</sup> ...honicke <sup>8</sup> , Carolina Vera <sup>4,5</sup>	... [31]
Formatted	... [32]
Deleted: Viovy <sup>8</sup> ...iovy <sup>9</sup> , Celso Von Randow <sup>2</sup>	... [33]
Deleted: <sup>2</sup> Earth	
Deleted: <sup>3</sup> GCU	
Deleted: <sup>4</sup> Universidad	
Deleted: <sup>5</sup> CONICET	
Deleted: <sup>6</sup> CNRS	
Deleted: <sup>7</sup> Potsdam	
Deleted: <sup>8</sup> LSCE	

## Abstract

Over the last decades, the Amazon rainforest was hit by multiple severe drought events. Here, we assess the severity and spatial extent of the extreme drought years 2005, 2010, and 2015/2016 in the Amazon region and their impacts on the regional carbon cycle. As an indicator of drought stress in the Amazon rainforest, we use the widely applied maximum cumulative water deficit (MCWD). Evaluating nine state-of-the-art precipitation datasets for the Amazon region, we find that the spatial extent of the drought in 2005 ranges from 2.2 to 3.0 (mean = 2.7) million km<sup>2</sup> (37 – 51% of the Amazon basin, mean = 45%) where MCWD indicates at least moderate drought conditions (relative MCWD anomaly < -0.5). In 2010, the affected area was about 16% larger, ranging from 3.0 up to 4.4 (mean = 3.6) million km<sup>2</sup> (51 – 74%, mean = 61%). In 2016, the mean area affected by drought stress was between 2005 and 2010 (mean = 3.2 million km<sup>2</sup>; 55% of the Amazon basin), but the general disagreement between datasets was larger, ranging from 2.4 up to 4.1 million km<sup>2</sup> (40–69%). In addition, we compare differences and similarities among datasets using the self-calibrating Palmer Drought Severity Index (scPDSI) and a rainfall anomaly index (RAI). We find that scPDSI shows a stronger, and RAI a much weaker drought impact in terms of extent and severity for the year 2016 compared to MCWD. We further investigate the impact of varying evapotranspiration on the drought indicators using two state-of-the-art evapotranspiration datasets. Generally, the variability in drought stress is most dependent on the drought indicator (60%), followed by the choice of precipitation dataset (20%) and the evapotranspiration dataset (20%). Using a fixed, constant evapotranspiration rate instead of variable evapotranspiration can lead to an overestimation of drought stress in the parts of Amazon basins that have a more pronounced dry season (for example in 2010). We conclude that for investigating impacts of droughts on the Amazon rainforest and e.g. its carbon cycle, multiple datasets for precipitation and also variable evapotranspiration input should be considered. Furthermore, considering different drought indices can help to understand the complex characteristics that drought events in the Amazon have.

**Deleted:** .

**Deleted:** .

**Deleted:**

**Deleted:** an ensemble of

**Deleted:** empirical MCWD-mortality relationship, we calculate biomass losses for the 2005 drought event. We show

**Deleted:** the majority of the datasets agree on biomass losses of about 1.2 Petagram carbon (PgC), but the overall range is between 0.7 and 1.6 PgC. Disagreement across datasets increased, (1) when comparing the total area of severe and extreme drought signals and (2) when comparing spatial drought location across datasets.

**Deleted:** deriving

**Deleted:** Basin based on precipitation

**Deleted:** This is especially relevant when assessing the impact of drought on the Amazon rainforest and its carbon cycle.

## 1 Introduction

The severe drought events occurring in 2005, 2010, and 2015/16 in the Amazon basin are reasons for concern regarding their frequency and severity, and their impacts on the Amazon rainforest. Different large-scale atmospheric processes related to increased sea surface temperature (SST) in the Pacific and the Atlantic Ocean seem to be responsible for such repeated mega-drought events (Coelho et al., 2012): While the drought 2015/16 was driven by a record-level El Niño event enhanced by the strong underlying global warming trend (Jimenez et al., 2018), the 2010 drought was a combination of a moderate El Niño event and anomalously warm SSTs in the tropical North Atlantic (Marengo et al., 2011; Marengo & Espinoza, 2016). Similarly, the 2005 drought was attributed to anomalies of warm SSTs in the North Atlantic (Marengo, Nobre, Tomasella, Oyama, et al., 2008; Zeng et al., 2008). In consequence, such events differ in their strength, their timing, and spatial patterns, and thus, impacted regions differ. While drought events related to El Niño events show a Southwest to Northeast gradient with dry conditions over the NE Amazon region (Malhi et al., 2008), drought events caused by anomalously warm North Atlantic SSTs show a North-South gradient with dry conditions in the southern Amazon region (Lewis et al., 2011; Marengo et al., 2008). Even in the case of El Niño events, SSTs anomalies over the Eastern Pacific (EP) or the Central Pacific (CP) can lead to different impacts and spatial patterns of drought (Jimenez et al., 2019). In addition to their influence on temperature, recent El Niño events also showed amplified atmospheric vapor pressure deficit anomalies (Barkhordarian et al., 2019; Rifai et al., 2019). The impacts of such drought events on humid tropical forests, which are often not adapted to longer-lasting dryness, are severe. Increased forest mortality connected to drought events was observed in central and southern Amazonia (Feldpausch et al., 2016; Lewis et al., 2011; Phillips et al., 2009), as well as shifts in tree species composition (Esquivel-Muelbert et al., 2019). Droughts are assumed to be one of the main drivers for the observed decline in the Amazon carbon sink, indicating that more carbon is lost to the atmosphere than taken up by the forest (Hubau et al., 2020). Thus, such extreme drought events are altering the carbon cycle of the Amazon forest (Feldpausch et al., 2016; Gloor et al., 2015; Hubau et al., 2020; Phillips et al., 2009).

Losing tropical forests in the Amazon region through increased mortality under drought also has implications for regional and continental scale water cycling (Ruiz-Vásquez et al., 2020). The rainforest transpires enormous amounts of water which is transported by winds to remote regions far beyond the borders of the rainforest (e.g. Dirmeyer et al., 2009; van der Ent et al., 2010; Zemp et al., 2014, 2017). In addition, the ongoing deforestation in the Amazon rainforest further decreases forest cover and thus, transpiration rates, leading to a rainfall decline and enhanced drought conditions in a positive feedback loop (Miralles et al., 2019; Zemp et al., 2017). It can be expected that ongoing climate change most likely will cause stronger and more frequent drought events in the Amazon (Cai et al., 2015; Jiang et al., 2020; Marengo & Espinoza, 2016).

For assessing the severity, the spatial extent, and, in particular, the impacts of such drought events on existing ecosystems, different gridded precipitation datasets are available which in some cases differ strongly in magnitude and spatio-temporal distribution of precipitation amounts (Golian et al., 2019). Typical problems of precipitation data for South America encompass the underestimation of extreme rainfall events in both dry or wet seasons (Blacutt et al., 2015; Giles et al., 2020). Therefore,

**Deleted:** (Coelho et al., 2012): While the drought 2015/16 was driven by a record-level El Niño event enhanced by the strong underlying global warming trend (Jimenez et al., 2018), the 2010 drought was a combination of a moderate El Niño event and anomalously warm SSTs in the tropical North Atlantic (Marengo & Espinoza, 2016; Marengo et al., 2011). Similarly, the 2005 drought was attributed to anomalies of warm SSTs in the North Atlantic (Marengo et al., 2008; Zeng et al., 2008). In consequence, such events differ in their strength, their timing, and spatial patterns, and thus, impacted regions differ. While drought events related to El Niño events show a Southwest to Northeast gradient with dry conditions over the NE Amazon region (Malhi et al., 2008), drought events caused by anomalously warm North Atlantic SSTs show a North-South gradient with dry conditions in the southern Amazon region (Lewis et al., 2011; Marengo et al., 2008). Even in the case of El Niño events, SSTs anomalies over the Eastern Pacific (EP) or the Central Pacific (CP) can lead to different impacts and spatial patterns of drought (Jimenez et al., 2019). In addition to their influence on temperature, recent El Niño events also showed amplified atmospheric vapor pressure deficit anomalies (Barkhordarian et al., 2019; Rifai et al., 2019). The impacts of such drought events on humid tropical forests, which are often not adapted to longer-lasting dryness, are severe. Increased forest mortality connected to drought events was observed in central and southern Amazonia (Feldpausch et al., 2016; Lewis et al., 2011; Phillips et al., 2009), as well as shifts in tree species composition (Esquivel-Muelbert et al., 2019). Droughts are assumed to be one of the main drivers for the observed decline in the Amazon carbon sink, indicating that more carbon is lost to the atmosphere than taken up by the forest (Hubau et al., 2020). Thus, such extreme drought events are altering the carbon cycle of the Amazon forest (Feldpausch et al., 2016; Gloor et al., 2015; Hubau et al., 2020; Phillips et al., 2009).

**Formatted:** German

**Deleted:** Losing tropical forests in the Amazon region through increased mortality under drought also has implications for regional and continental scale water cycling (Ruiz-Vásquez et al., 2020). The rainforest transpires enormous amounts of water which is transported by winds to remote regions far beyond the borders of the rainforest (e.g. Dirmeyer et al., 2009; van der Ent et al., 2010; Zemp et al., 2014; Zemp et al., 2017). In addition, the ongoing deforestation in the Amazon rainforest further decreases forest cover and thus, transpiration rates, leading to a rainfall decline and enhanced drought conditions in a positive feedback loop (Miralles et al., 2019; Zemp et al., 2017). It can be expected that ongoing climate change most likely will cause stronger and more frequent drought events in the Amazon (Cai et al., 2015; Jia et al., 2019; Marengo & Espinoza, 2016).<sup>4</sup> For assessing the severity, the spatial extent, and, in particular, the impacts of such drought events on existing ecosystems, different gridded precipitation datasets are available which in some cases differ strongly in magnitude and spatio-temporal distribution of precipitation amounts (Golian et al., 2019). Typical problems of precipitation data for South America encompass the underestimation of extreme rainfall events in both dry or wet seasons (Blacutt et al., 2015; Giles et al., 2020). Therefore, while for the Amazon region, the recent drought events have been assessed in terms of severity (Jiménez-Muñoz et al., 2016; Jimenez et al., 2018) and impacts (Phillips et al. 2009, Lewis et al. 2011, Feldpausch et al. 2016) based on single precipitation data sets, a systematic analysis of how the most frequent used precipitation datasets differ regarding the spatial extent, location and severity of recent extreme drought events, is currently missing.<sup>4</sup>

... [34]

while for the Amazon region, the recent drought events have been assessed in terms of severity (Jimenez et al., 2018; Jiménez-Muñoz et al., 2016) and impacts (Feldpausch et al., 2016; Lewis et al., 2011; Phillips et al., 2009) based on single precipitation data sets, a systematic analysis of how the most frequent used precipitation datasets differ regarding the spatial extent, location and severity of recent extreme drought events, is currently missing.

For our study, we selected precipitation from nine different datasets: (1, 2) Data from the Tropical Rainfall Measurement Mission (TRMM) version 6 and 7 (Huffman et al., 2007) which have been frequently used, e.g. to estimate drought impacts on the carbon balance (Lewis et al., 2011; Malhi et al., 2009) and are assumed to represent precipitation patterns in the Amazon region best since they are derived from radar measurements (Huffman et al., 2007). (3) CHIRPS (Climate Hazards group Infrared Precipitation with Stations, Espinoza et al., 2019), which has been used to study regional hydro-climatic and environmental changes in the Amazon Basin. These two datasets only provide precipitation and no information about other climatic variables such as temperature or radiation. In addition, we selected five datasets that are often used as drivers for ecosystem models (e.g. in Forkel et al., 2019; Yang et al., 2015) and – in contrast to the other datasets – provide information for more climate variables: Data from the Climate Research Unit (CRU) with a joint project reanalysis (NCEP, National Centers for Environmental Prediction) applied, (4) the CRUNCEP (version 8, Viovy, 2018), (5) the WATCH-WFDEI (WATCH: Water and Global Change, Weedon et al., 2011). WFDEI: WATCH Forcing Data methodology applied to ERA-Interim, Weedon et al., 2014) dataset, originally derived from global sub-daily observations merged with integrations from a general circulation model, (6) the GSWP3 (Global Soil Wetness phase 3, Kim et al. in prep) dataset which is closely related to WATCH-WFDEI, relying on a similar forcing but with a different bias-correction method applied, (7) the newer GLDAS (Global Land Data Assimilation System) 2.1, which is derived from various geostationary infrared satellite measurements and microwave observations (Rodell et al., 2004), (8) the latest ECMWF atmospheric reanalysis dataset, ERA5, which is the successor of ERA-Interim, providing higher spatial and temporal resolutions and a more recent model and data assimilation system than the previous ERA-Interim reanalysis (Albergel et al., 2018), and, finally, (9) the GPCC (named after the Global Precipitation Climatology Centre) dataset (Schneider et al., 2018), which is based on globally available land stations (rain gauges) combined with an empirical interpolation method (Willmott et al., 1985). The data sets were chosen because they are often used to force Dynamic Global Vegetation and hydrological simulation models in climate impacts studies. A more detailed description of the datasets is given in the methods section.

We evaluate the precipitation datasets based on the Maximum Cumulative Water Deficit (MCWD; Aragão et al., 2007), a well-established drought index that is particularly suitable for estimating drought stress in the Amazon region (e.g. Esquivel-Muelbert et al., 2019; Lewis et al., 2011; Malhi et al., 2009; Phillips et al., 2009; Zang et al., 2020). In addition, we included two other measures to complement our analysis: A rainfall anomaly index (RAI), which does account for the mean deviation (in units of standard deviation) of precipitation during the driest months of the year, and the scPDSI (self-calibrating Palmer Drought Index, Wells et al., 2004). The scPDSI index has a more complex formulation compared to RAI and MCWD and takes available soil water content into account. Both RAI and scPDSI have been used in studies describing the recent Amazonian drought events (e.g. Jiménez-Muñoz et al., 2016; Lewis et al., 2011). Many studies (e.g. Flack-Prain et al., 2019;

[Hubau et al., 2020](#)) currently still use a fixed evapotranspiration rate for the calculation of MCWD instead of using evapotranspiration datasets as input. To assess the robustness of a fixed evapotranspiration rate, we include two evapotranspiration datasets GLEAM (Martens et al., 2017) and DOLCE (Hobeichi et al., 2018) for the calculation of MCWD and scPDSI. The goals of our study are (1) to analyse and quantify the uncertainty in strength, extent, and location of three recent Amazon droughts in the years 2005, 2010, and 2015/2016 in precipitation from nine state-of-the-art precipitation or climate datasets based on MCWD; (2) to examine differences among these drought events by taking two additional drought indicators RAI and scPDSI and two evapotranspiration datasets into account.

## 2 Methods

### 2.1 Study area

[Our study covers the Amazon river basin as delineated by Döll & Lehner \(2002\)](#), see black contour in Fig. 1). Using 0.5° spatial resolution in longitude and latitude results in 1946 grid cells of interest for this study area. Note that differences in the comparison of our results with [Lewis et al., \(2011\)](#) arise because of differences in the delineation of the Amazon region, i.e. the area used in our study is 0.6 million km<sup>2</sup> larger.

**Deleted:** Our study covers the Amazon river basin as delineated by Döll & Lehner (2002),

**Deleted:** Lewis et al. (2011)

### 2.2 Data sources

In the following, we briefly describe the nine precipitation datasets applied in our study (see also Table 1): The Tropical Rainfall Measuring Mission (TRMM v7) product ([Huffman et al., 2007](#)) is a precipitation-only dataset based on multiple microwave-infrared satellite data developed as a joint product between NASA and the Japan Aerospace Exploration Agency (JAXA). We also included the predecessor v6 for comparison in our study, because it has been frequently and prominently used to derive drought impacts to the Amazon Basin (e.g. Lewis et al., 2011; Phillips et al., 2009) and shows significantly lower precipitation throughout the basin compared to v7 (Seto et al., 2011). CHIRPS (Climate Hazards group Infrared Precipitation with Station) is a novel dataset (Funk et al., 2015) which is a quasi-global (full longitude, but only 50°S – 50°N latitude extent) precipitation-only merged product, based on multi-satellite estimates (similar to TRMM 6 and TRMM 7) and approx. 2,000 in-situ observations per month in South America. TRMM 6, TRMM 7 and CHIRPS share the quasi-global spatial extent, however, in comparison to TRMM 6, TRMM 7 with a resolution of 0.25° x 0.25°, CHIRPS has a much higher spatial resolution of 0.05° x 0.05°. ERA5 (Muñoz-Sabater et al., 2018) shows improvements in, e.g., land evapotranspiration, surface soil moisture and turbulent heat fluxes over its predecessor ERA-Interim (Albergel et al., 2018), which we decided not to include in our study as it showed higher systematic errors over tropical areas (Nogueira, 2020). Similarly, CRUNCEP (Viovy, 2018) is generated based on a reanalysis from the national centers for environmental prediction (NCEP) and the National Center for Atmospheric Research (NCAR), corrected with the CRU TS3.2 (Harris et al., 2014) dataset. GPCC is mainly based on data from rain-gauge land stations. Similar to CRUNCEP, it is also based on the NCEP reanalysis dataset and has been used in global drought studies (Ziese et al., 2014). Both GPCC and CRUNCEP cover the longest periods of all selected datasets in this study with time spans from 1891 until 2016 and from 1901 until 2016, respectively. WATCH-WFDEI (Weedon et al., 2011; 2014) is based on the reanalysis ERA-Interim corrected with GPCC precipitation. GSWP3 (Kim et al. in prep.) is based on the atmospheric reanalysis method “20CR” (20th Century Reanalysis version 2, Compo et al., 2013), which has been dynamically downscaled to 0.5° x 0.5° resolution. Corrections with observational data have not only been applied to precipitation but also to short/longwave radiation, air temperature and the daily temperature range. Both WATCH-WFDEI and GSWP end in the year 2010. The GLDAS 2.1 dataset is built by using the ‘Noah Land surface model’ forced by the Goddard Earth Observing System (GEOS) Data Assimilation System with corrected precipitation and radiation (Rodell et al., 2004; Sheffield et al., 2006). Starting in January 2000 (Version 2.1), it is the dataset with the latest time onset and hence

**Deleted:** (Huffman et al., 2007) is a precipitation-only dataset based on multiple microwave-infrared satellite data developed as a joint product between NASA and the Japan Aerospace Exploration Agency (JAXA). We also included the predecessor v6 for comparison in our study, because it has been frequently and prominently used to derive drought impacts to the Amazon Basin (e.g. Lewis et al., 2011; Phillips et al., 2009) and shows significantly lower precipitation throughout the basin compared to v7 (Seto et al., 2011). CHIRPS (Climate Hazards group Infrared Precipitation with Station) is a novel dataset (Funk et al., 2015) which is a quasi-global (full longitude, but only 50°S – 50°N latitude extent) precipitation-only merged product, based on multi-satellite estimates (similar to TRMM 6 and TRMM 7) and approx. 2,000 in-situ observations per month in South America. TRMM 6, TRMM 7 and CHIRPS share the quasi-global spatial extent, however, in comparison to TRMM 6, TRMM 7 with a resolution of 0.25° x 0.25°, CHIRPS has a much higher spatial resolution of 0.05° x 0.05°. ERA5 (Muñoz-Sabater et al., 2018) shows improvements in, e.g., land evapotranspiration, surface soil moisture and turbulent heat fluxes over its predecessor ERA-Interim (Albergel et al., 2018), which we decided not to include in our study as it showed higher systematic errors over tropical areas (Nogueira, 2020). Similarly, CRUNCEP (Viovy, 2018) is generated based on a reanalysis from the national centers for environmental prediction (NCEP) and the National Center for Atmospheric Research (NCAR), corrected with the CRU TS3.2 (Harris et al., 2014) dataset. GPCC is mainly based on data from rain-gauge land stations. Similar to CRUNCEP, it is also based on the NCEP reanalysis dataset and has been used in global drought studies (Ziese et al., 2014). Both GPCC and CRUNCEP cover the longest periods of all selected datasets in this study with time spans from 1891 until 2016 and from 1901 until 2016, respectively. WATCH-WFDEI (Weedon et al., 2011; 2014) is based on the reanalysis ERA-Interim corrected with GPCC precipitation. GSWP3 (Kim et al. in prep.) is based on the atmospheric reanalysis method “20CR” (20th Century Reanalysis version 2, Compo et al., 2013), which has been dynamically downscaled to 0.5° x 0.5° resolution. Corrections with observational data have not only been applied to precipitation but also to short/longwave radiation, air temperature and the daily temperature range. Both WATCH-WFDEI and GSWP end in the year 2010. The GLDAS 2.1 dataset is built by using the ‘Noah Land surface model’ forced by the Goddard Earth Observing System (GEOS) Data Assimilation System with corrected precipitation and radiation (Rodell et al., 2004; Sheffield et al., 2006).



anomaly ~~every~~ analysis conducted in our study. We also estimated seasonal patterns of cumulative water deficit (CWD), by defining  $rCWD$  similar to  $rMCWD$  as the relative anomaly of each month's CWD in units of standard deviation.

**Deleted:** for the majority of the  
**Deleted:** except for the impact of drought on aboveground biomass (section 2.4 and Fig. 4), where we use the  $aMCWD$  anomaly

### 2.3.2. Calculation of rainfall anomaly index (RAI)

For the rainfall anomaly index, dry season rainfall was taken as the mean precipitation from July-September following Lewis et al. (2011). Like for the MCWD estimation, we calculated the mean dry season rainfall from a baseline period 2000-2010 to investigate the drought impacts of 2005 and 2010, and for 2016 we selected a baseline period from 2000 to 2016 excluding 2005, 2010, and 2016. The relative rainfall anomaly index ( $rRAI$ ) was estimated as 'standardized anomaly' from the baseline period similarly to the  $rMCWD$  anomaly calculation. As  $rRAI$  only reflects the precipitation anomaly during July and September, it can also be described as a dry season anomaly.

### 2.3.3. Calculation of the self-calibrating Palmer Drought Severity Index (scPDSI)

The self-calibrating Palmer Drought Severity Index (scPDSI, Wells et al., 2004) has in recent studies been used to assess the impacts of droughts on the Amazon basin (e.g. Jiménez-Muñoz et al., 2016). It improves the original PDSI by using a self-calibrating procedure based on historical climate data, eliminating the empirically derived climatic characteristics. Next to precipitation, it also takes monthly ~~evapotranspiration~~ ET into account. In our study, we use ET data generated ~~from DOLCE and GLEAM (section 2.4)~~. Additionally, the scPDSI takes soil water capacity as input, which we assumed here as a constant value of 100 mm. scPDSI was estimated using the R package *scPDSI* (Ruida et al., 2018).

**Deleted:** potential  
**Deleted:** by the ERA5 reanalysis.  
**Deleted:** (Ruida et al., 2018).

To enable cross-comparison with the  $rMCWD$  and  $rRAI$  anomalies, we selected identical baseline periods from 2000 to 2010 for the 2005 and 2010 events, and from 2000 to 2016 for the 2016 drought event. Again, we used the relative deviation  ~~$rscPDSI$~~ , defined as 'standardized anomaly' from the baseline period of monthly scPDSI values as drought indicator.

**Deleted:**  $rscPDSI$

## 2.4. Evapotranspiration datasets

**Deleted:** 2.3.

In addition to assuming a constant evapotranspiration  $ET(t) = ET_{fixed} = 100$  mm for the calculation of MCWD, and for the calculation of scPDSI we use the two *ET* datasets GLEAM and DOLCE. The Global Land Evaporation Amsterdam Model (GLEAM) v3a dataset (Martens et al., 2017) is derived from a set of algorithms incorporating satellite-observed soil moisture, vegetation optical depth, reanalysis air temperature and radiation, and multiple precipitation datasets. The Derived Optimal Linear Combination Evapotranspiration (DOLCE, Hobeichi et al., 2018) dataset is derived by combining and weighting multiple other evapotranspiration datasets, also including GLEAM.

## 2.5. Calculation of drought area and extent

Each grid cell's area was approximated as a trapezoid to its boundary coordinates (in  $0.5^\circ \times 0.5^\circ$  resolution), resulting in an area between 2900 and 3090 km<sup>2</sup> per grid cell. Accumulating the associated areas over all grid cells resulted in a total area of 5.94 million km<sup>2</sup> representing the Amazon Basin. Note that for comparison of our results with Lewis et al. (2011) differences in absolute areas arise because of differences in study area size (5.94 vs. 5.3 million km<sup>2</sup>, respectively). For the calculation of the drought-affected area, we summed up the area of grid cells that matched the respective drought classification (e.g.  $rMCWD$  anomaly  $< -2.5$  for extreme drought stress). The spatial agreement of drought location among datasets was estimated by selecting the grid cells matching the drought classification per dataset and subsequently counting the number of datasets per grid cells showing the respective drought classification.

### **Deleted: 2.4. Estimating carbon losses during drought events**

To estimate carbon loss during drought events, we used a simple linear relation between  $\Delta MCWD$  anomaly and  $\Delta AGB$ , the change in aboveground biomass, i.e. biomass carbon losses in the Amazon basin derived from plot measurements (Lewis et al., 2011):

$$\Delta AGB = 0.3778 - 0.052 * \Delta MCWD \quad (2)$$

The equation was derived from Amazon plot inventory data measured across the RAINFOR network to estimate the impact of the 2005 drought event (Lewis et al. 2011). To calculate the  $\Delta AGB$  anomaly in Eq. 2, we calculated the  $\Delta MCWD$  anomaly of each gridcell in 2005 for each of the precipitation datasets in our study. The total biomass carbon loss (in PgC) across the Amazon basin is then calculated by summing up  $\Delta AGB$  anomaly for all gridcells weighted by each gridcell's size.

### 3. Results

All areas in the following section are expressed as percentage [with respect to the entire Amazon basin](#) according to our delineation (5.94 million km<sup>2</sup>). For an overview of the areas affected in million km<sup>2</sup>, see [Tab. S1](#) and [S2](#).

Deleted: of

Deleted: Table S2

Deleted: S3

#### 3.1 Comparison of total drought area based on relative MCWD anomaly

We first evaluate differences in  $rMCWD$  for 2016 across the datasets (Fig. 1). Here, we find that the spatial patterns of the  $rMCWD$  anomaly generally match across the available datasets, showing severe and extreme drought stress mainly in the northern Amazon basin. Only GLDAS diverges, showing extreme drought stress in the Central and Western part of Amazonia (Fig. 1d) where none of the other datasets [shows](#) any drought stress during the same year. The other datasets mostly differ regarding the intensity of the drought stress. While ERA5 and TRMM7 show values  $rMCWD < -2.5$  in the Columbian part of the basin, CRUNCEP and GPCC do show such a strong drought impact only in Northern Brazil. The absolute areas of drought stress across different severity levels are similar across most datasets with only GLDAS showing a significantly larger area affected by extreme drought stress of  $rMCWD < -2.5$ .

Deleted: show

Across all precipitation datasets, in 2005, an area ranging from 37 to 51% (mean 45%) of the whole Amazon basin, was moderately affected (Table [S1](#), Fig. 2a). ERA5 displayed the smallest area affected by moderate drought (2.2 million km<sup>2</sup>, Tab. 1, Fig. 2), while CHIRPS and CRUNCEP showed a vast affected area (3.0 million km<sup>2</sup>), an area about 36% larger than displayed by ERA5. For severe and extreme drought conditions, ERA5 shows the smallest affected area with 3% and 1% of the basin affected. For severe drought conditions, CRUNCEP suggests that an area approximately 3 times larger was affected compared to ERA5 (0.2 million km<sup>2</sup> vs. 0.6 million km<sup>2</sup>). CRUNCEP and GLDAS also encompass the largest area of extreme drought stress (0.2 million km<sup>2</sup>; 3% of the basin less than  $rMCWD < -2.5$ , Fig. 2a).

Deleted: S2

During the 2010 drought, a larger area ranging between a minimum of 52% (GPCC) and a maximum of 74% (TRMM 6) was affected by moderate drought stress, which is about 36% larger than during the 2005 drought (3.6 million km<sup>2</sup> vs. 2.7 million km<sup>2</sup>, Table [S1](#), Fig. 2). In addition, the area under severe drought stress was on average 25% larger compared to 2005 and the area affected by extreme drought was double the size of the 2005 drought event. Particularly, GLDAS and TRMM 6 showed the largest area affected throughout the three drought classifications (Fig. 2b).

Deleted: S2

For 2016, two datasets (CHIRPS and CRUNCEP) showed with 38% a considerably smaller area that was moderately affected by drought stress compared to GLDAS with 63% of the area affected, respectively (datasets ranging between 2.2 and 3.7 million km<sup>2</sup>). Generally, in 2016, the size of the area affected by moderate drought was in between the size of the area affected in 2005 and 2010, but the extent of severely and extremely drought-affected areas was larger. Here, particularly GLDAS followed by GPCC showed the largest affected area, with 21% severely affected and 6% extremely affected ([Tab. S2](#)).

Deleted: Table S3

### 3.2 Spatial agreement of rainfall datasets using the $r$ MCDW anomaly

While the agreement of the total area affected by drought is relatively high (see 3.1) the data sets only partly agree on the spatial extent and location of extreme drought conditions, particularly during the 2010 and 2016 events (Fig. 3). For 2005, all datasets agree on the drought epicentre being in Central Amazonia. Datasets agree that an area of about 15 % of the Amazon Basin was at least moderately affected (Fig. 3a). Only a small overlap was found for the area affected by severe and extreme drought stress (Fig. 3b, c). Here, only half of the datasets agreed on 4% of central Amazonia being severely and 1.5% extremely affected.

**Deleted:** epicenter

**Deleted:** located

For 2010, all datasets agreed on an affected area of 21% in the Amazon basin, and half of the datasets agreed on an area of 60% of the Amazon Basin being moderately affected by drought stress (Fig. 3d). The 2010 drought displayed no central hotspot, but three most affected areas in the Eastern, Southern and central parts of Amazonia on which most of the datasets agreed (Fig. 3d). Severe drought stress in 2010 was in the southern part of Amazonia, where four datasets agreed (Fig. 3e), while for extreme drought stress almost no overlap between datasets was found (Fig. 3f).

**Deleted:** located

For 2016, all datasets agreed on an area of about 7% of moderate drought stress and half of the datasets agreed on 51% of the basin being affected (Fig. 3g). Agreement for severe and extreme drought stress was lower compared to the other drought years (Fig. 3h, i). Most of the datasets located the epicentre of the drought in the north-western Amazon basin. Some datasets also showed the South-Central part of the basin being severely affected (Fig. 3i).

**Deleted:** epicenter

**Deleted:** western

We could not find any pronounced biases between the precipitation datasets (Fig. S3 - S5), but a generally higher correlation of the  $r$ MCWD anomalies for 2005 compared to 2010 and 2016. Only ERA5 and GLDAS showed some spikes in the  $r$ MCWD anomalies that are located within the high latitude regions of the Andes.

**Deleted:** 3.3 Estimating the variation of carbon losses during drought events<sup>1</sup>

For the different precipitation datasets and based on the linear relation between  $a$ MCWD and  $a$ AGB anomaly, we derive carbon losses for 2005 to be in the range of 0.7-1.6 PgC with ERA5 showing the smallest and CRUNCEP the strongest impact regarding carbon losses (Fig. 4). The mean biomass loss overall datasets was 1.2 PgC with five of the nine estimated carbon losses being close to that mean (difference of  $a$ AGB anomaly less than 0.1 PgC to the mean value). Because no relationship between the anomalies of  $a$ MCWD and  $a$ AGB could be verified for 2010 (Feldpausch et al., 2016) we did not estimate the impacts on AGB for the other drought years 2010 and 2016....

### 3.3 Constant versus varying evapotranspiration rates: Effects on drought severity and extent estimates

We find that assuming a constant ET rate of 100 mm month<sup>-1</sup> is only realistic in the Northern part of the Amazon basin and only when comparing to the DOLCE dataset (Fig. 4a, b), which shows ET rates of about 100mm month<sup>-1</sup> during both the wettest (as averaged between June to August) and the driest months (as averaged between January and March). Using GLEAM, average ET rates are between 30 and 50% higher than 100mm month<sup>-1</sup> during the wettest months (Fig. 4c) and remain higher than 100mm month<sup>-1</sup> also in the Northern part during the dry season (Fig. 4f). Evapotranspiration rates can be as low as 50mm month<sup>-1</sup> on average throughout the driest months for both ET datasets in the South of the basin (Fig. 4b. and 4f).

This spatial heterogeneity in evapotranspiration rates has implications for the extent and severity of drought stress expressed as  $r$ MCWD anomaly when compared to using constant evapotranspiration. Using the two evapotranspiration datasets we find lower drought impacts across most parts of the Amazon basin for the two years 2005 and 2010 (Fig. 4c, d, g, h). In 2005 the mean area of moderate drought stress is lower when using variable ET: 44% of the basin for GLEAM and 39% for DOLCE.

compared 46% for a constant ET. Interestingly, these differences were not particularly located in the epicentre of drought during that year (see Fig. 3a, b, c), but rather in the South and the high latitude regions toward the Andes (Fig. 4c, g). The total area of severe drought stress did only slightly decrease from 9% (constant ET) to 8% (GLEAM) and 7% (DOLCE). In 2010, we find stronger differences between variable and constant ET. The area of moderate drought stress is 52% for GLEAM and 49% for DOLCE which is significantly lower than the 60% when using constant evapotranspiration. For this year the areas of these differences (Fig. 4d, h) strongly overlap with the epicentres of the drought (see Fig. 3d, e, f). Consequently, also the areas of severe drought stress are lower (7% for GLEAM, 8% for DOLCE) compared to using constant evapotranspiration (12%). We find similar patterns for 2016 (not shown) where the mean area of severe drought stress is approximately 11% for both GLEAM and DOLCE, which is lower compared to using constant ET (15%).

### 3.4 Comparison of drought indices: $r$ MCDW, $r$ scPDSI and $r$ RAI anomalies

Similar to  $r$ MCWD, there is variable agreement among datasets when evaluating the other two drought metrics,  $r$ RAI and  $r$ scPDSI (Fig. 5). The largest dry season anomaly ( $r$ RAI) in 2005 was displayed by GPCC with 6.5% (0.4 million km<sup>2</sup>, Table 2), followed by TRMM 7 with 5.7% of the Amazon basin being severely affected. ERA 5 showed with 3% the smallest area affected. In 2005, spatial patterns of  $r$ RAI matched with  $r$ MCWD anomalies despite  $r$ MCWD anomalies showing a larger area affected by severe drought stress (Fig. 5a, d).  $r$ scPDSI displayed the smallest area affected by drought stress in 2005 with also only GPCC and TRMM 7 showing with 5.5% and 3.1% the largest severely affected area, respectively. All other datasets showed less than 1% of severe drought-affected areas in 2005. The small spatial area of  $r$ scPDSI differed compared to the other two drought indicators (Fig 5a, d, g): Some areas showed a strong disagreement between drought indices, e.g. Central Amazonia was hit by severe drought stress according to  $r$ MCWD and  $r$ RAI (with 3-4 climate datasets in agreement) while, in contrast,  $r$ scPDSI did not indicate abnormally dry conditions there.

In 2010, the differences of drought-affected areas were even more pronounced between the three indices (Fig. 5b, e, h). Here, ERA5 and TRMM7 showed the largest areas affected by severe drought stress based on the dry season  $r$ RAI anomaly with 7% and 5%, respectively. Using  $r$ scPDSI all datasets showed an area between 1% and 2.5% severely affected. Interestingly, the area affected based on  $r$ MCWD roughly encompasses the area affected by  $r$ RAI, but additionally shows a large area in the South-Eastern part of the basin being affected by severe drought stress (Fig. 5b, e).

In 2016,  $r$ scPDSI shows the largest area affected by drought stress with GLDAS showing 39% (followed by TRMM7, 16%) of the basin being severely affected. Four datasets agreed on the affected area in the northeastern part of the basin (Fig. 5i). Only one dataset (GLDAS) showed severe drought stress in 2016 when calculating dry season rainfall anomalies ( $r$ RAI, Fig

Deleted:  $r$ scPDSI

Deleted: S2

Deleted:  $r$ scPDSI

Deleted:  $r$ scPDSI

5c), indicating no pronounced anomalies in dry season rainfall according to all other datasets.  $r_{MCWD}$  and  $r_{scPDSI}$  roughly agreed on the northern part of the basin being severely affected (Fig. 5f, i).

Average seasonal patterns are quite consistent across datasets but differ depending on the choice of drought index and drought event (Fig. 6). The strongest (most negative) rainfall anomaly was visible from May to July during the 2005 drought event (Fig. 6a). Accumulating such low rainfall estimates resulted in very low values of  $r_{CWD}$  during that period (Fig. 6d) in 2005.  $r_{scPDSI}$  values were also low, but more constant throughout the year (Fig. 6g).

The 2010 drought followed similar patterns regarding  $r_{RAI}$  with a lower absolute impact during May to July compared to 2005 (Fig 6b). Interestingly, the wet season months March to May showed a strong anomaly during 2010 compared to the 2005 event. Subsequently,  $r_{CWD}$  was also already lower during the wet season in 2010 compared to 2005 (Fig. 6e).  $r_{scPDSI}$  anomalies values were similar for 2010 compared to 2005 with a slightly downward trend towards the end of the year (Fig. 6g, h).

To investigate the seasonal patterns of 2016 we also considered the drought indices of 2015 since both years were El Niño years. We found a strong rainfall anomaly already starting during September 2015 continuing until April 2016 (Fig. 6c). Consequently, also  $r_{CWD}$  values were very low during that period (Fig. 6f). While  $r_{MCWD}$  was applied as the maximum value from October to September, drought stress before October of the previous year cannot be accounted for when using  $r_{MCWD}$ . The two-year drought impact was also visible using  $scPDSI$  (Fig. 6i) showing a steady decline from 2015 to 2016.

### **3.5 Overall variability: precipitation datasets vs. drought indices vs. evapotranspiration datasets**

When assessing the variability of drought severity and extent across the nine different precipitation datasets, the two drought indices ( $r_{MCWD}$  and  $r_{scPDSI}$ ) and the two evapotranspiration datasets (DOLCE and GLEAM), we find that across all drought events the choice of drought index accounts for roughly 60% of the variability while both the precipitation dataset and the evapotranspiration dataset account for 20%, each (Tab. 2).

#### 4. Discussion

We assessed the severity and spatial extent of the extreme drought years 2005, 2010, and 2015/2016 in the Amazon region by computing different drought indices using a range of precipitation datasets. When analyzing how drought conditions are captured in nine different precipitation datasets for the Amazon basin, we find that while the datasets mostly agree on the extent of the drought area, they differ in their location of drought.

##### Critical aspects regarding the detection of drought events in the Amazon basin

###### *Drought indices*

The idea of defining water deficit based on evapotranspiration rates goes back to Stephenson (1998) and the MCWD is now one of the most widely used indicators to assess drought stress in tropical forests (e.g. Lewis et al., 2011, Phillips et al., 2009, Esquivel-Muelbert et al., 2019). In its simplest form, the calculation of MCWD only requires precipitation data and assumes a constant evapotranspiration (ET) rate of 100 mm month<sup>-1</sup> (Aragão et al., 2007). Although the simplicity of *r*MCWD and  $\alpha$ MCWD is a main advantage, a fixed ET (which we also used in our study) is inappropriate for regions other than the lowland tropics, where the lower supply of energy may result in lower ET values. Most importantly, an approximated ET does not account for either seasonal variation (driven mainly by radiation, temperature, and phenology) or spatial variation in ET related to soil and root properties (Malhi et al., 2009). Hence, changes in *r*MCWD are purely accounting for changes in rainfall (Phillips et al., 2009). In contrast, *sc*PDSI is driven with spatially and temporally resolved evapotranspiration data. However, currently available evapotranspiration products for the Amazon rainforest show significant differences in areas and extent of evapotranspiration (Sörensson & Ruscica, 2018), hence introducing another source of uncertainty when using them for the calculation of drought indices. In the last decade, better products of spatially and temporally resolved evapotranspiration data (e.g. ERA5) have been developed and an increasing number of studies are now estimating MCWD based on such data (e.g. Staal et al., 2020). However, using a constant evapotranspiration (ET) rate of 100 mm month<sup>-1</sup> across the Amazon rainforest is still very common (e.g. Flack-Prain et al., 2019; Koch et al., 2021). Using variable evapotranspiration consistently reduced the moderate drought-affected area by 5-10% per drought event (Fig. 4). Extending the baseline period of the MCWD calculation to include also years before 2001 leads to overall lower MCWD values and, hence, an increased intensity of the three drought events. This finding highlights the drought anomaly that the recent decade from 2001 to 2016 has compared to the years before that period.

The key difference between the three drought indices applied in our study is the temporal resolution: RAI is only calculated for the three driest months (July-September) and thus, for example, a rainy season with deficient rainfall is not captured. MCWD, in contrast, accumulates over 12 months and is reset to zero at the end of the hydrological year. In this way, drought events caused by low precipitation in both dry- and rainy seasons are captured, however, drought events lasting for more than a year are not detected. *sc*PDSI captures multi-year drought events and is not reset to zero at the end of the hydrological year.

**Deleted:** We found a wide range between 0.7 and 1.7 PgC of potential biomass losses during 2005 with most datasets showing an impact of about 1.2 PgC.

**Deleted:** ,

**Deleted:**

**Deleted:** (here: ERA5).

**Deleted:** (Sörensson & Ruscica, 2018),

**Deleted:** . Staal et al., 2020).

**Deleted:** Flack-Prain et al., 2019; Koch et al., 2021).

**Formatted:** English (US)

**Deleted:** We investigated the effect of choosing variable evapotranspiration and a longer baseline in our MCWD calculation (Fig. S3).

**Deleted:** -20

**Deleted:** S3a, b, c). It also affected the intensity of the drought stress, e.g. areas previously classified as extreme drought stress were now classified as areas under severe drought stress. This reduction is expected as ERA5 takes the above-mentioned lower ET values in the highland tropics into account which overall leads to higher MCWD values in this region. Because of the strength and consistency of this effect we recommend testing the MCWD calculation regarding its sensitivity to variable ET in the tropical rainforest in future studies. In contrast, extending

**Deleted:** (Fig. S3d, e, f).

**Deleted:**

**Formatted:** English (UK)

These differences between the drought indicators can be seen for the three drought events analysed in this study. In 2005,  $rRAI$  and  $rMCWD$  values roughly match in location of the epicenter indicating a particularly strong anomaly during the dry season (Fig. 5a, d). This does not apply to the 2010 drought event, where despite some dry season anomaly an even stronger anomaly during the wet season is visible (Fig. 6b, e). [The 2015/2016 drought event is classified as a severe multi-year drought according to Yang et al. \(2018\), which is also displayed in our analysis when using  \$rscPDSI\$  \(Fig 6i\).](#)  $rMCWD$  and  $rRAI$ , however, do not agree on a spatially and temporally extensive drought event in 2016 (Fig. 5c, f, i), but instead display distinct regions of severe drought stress. Seasonal patterns of the three drought indices support this assumption (Fig. 6): Resetting  $rMCWD$  once per year neglects any influences of drought events of the preceding year (Fig. 6c). While the drought indices used in this study showed pronounced differences in spatial and temporal dynamics, including all of them can help better understanding the different characteristics that drought events can have in the Amazon basin.

A common drawback of all drought metrics used in our study is their incapability to explicitly represent the effect of increasing atmospheric vapor pressure deficit (VPD) on plant water stress. [A steady amplification of atmospheric vapor pressure deficit \(VPD\) has been detected over the Amazon basin \(Barkhordarian et al., 2019; Rifai et al., 2019\). Such stronger atmospheric water demand leads to additional water loss of plants during drought, subsequently increasing the severity of droughts. Hence, the role of VPD during a drought and as a driver for plant stress should not be underestimated \(Grossiord et al., 2020\). With increasing data availability and better estimates of VPD across the Amazon region, it should be included in future drought assessments \(e.g. Castro et al., 2020\). One possibility to account for the influences of VPD is choosing temporal and spatially resolved evapotranspiration instead of constant evapotranspiration in the calculation of MCWD. Future studies could further investigate the relationships between MCWD, ET, and VPD and the impacts on biomass.](#)

[Furthermore, in the last decade, new methods have been developed that assess impacts of drought on ecosystems, e.g. analyses based on solar-induced fluorescence \(SIF\) data show that tall forests are less sensitive to rainfall compared to short forests \(Giardina et al., 2018\). Also, vegetation optical depth \(VOD\) used as a proxy for water content in forests is a promising satellite-derived indicator for mortality and impacts of droughts on forests \(Rao et al., 2019\). However, conducting analyses over the Amazon rainforest based on VOD is difficult, because of the limited penetration depth of microwaves in dense tropical forests \(Chaparro et al., 2019\), and the influences of vegetation water status \(Xu et al., 2021\). So far, VOD data could only be applied with limited success across tropical rainforests \(Konings & Gentine, 2017\). Future studies should estimate the impacts of droughts based on multiple drought characteristics. For example, Toomey et al. \(2011\) show that considering both, heat stress and soil moisture stress greatly improves the explanatory power of drought impacts in the Amazon basin.](#)

#### Precipitation datasets

For the three drought events in 2005, 2010 and 2016, CHIRPS, GLDAS and ERA5 diverted the most from the other datasets regarding the spatial drought extent. ERA5 shows the smallest area of moderate drought stress during 2005 but one of the largest areas in 2010 (Fig. 2). We found no obvious bias between the precipitation datasets regarding distribution and frequency of monthly rainfall (SI Fig. 2) with only ERA5 showing higher rainfall more frequently. Although TRMM7 and CHIRPS are

**Deleted:** The 2015/2016 drought event is classified as a severe multi-year drought according to Yang et al. (2018), which is also displayed in our analysis when using  $rscPDSI$  (Fig 6i).

**Deleted:** A steady amplification of atmospheric vapor pressure deficit (VPD) has been detected over the Amazon basin (Barkhordarian et al., 2019; Rifai et al., 2019). Such stronger atmospheric water demand leads to additional water loss of plants during drought, subsequently increasing the severity of droughts. Hence, the role of VPD during a drought and as a driver for plant stress should not be underestimated (Grossiord et al., 2020). With increasing data availability and better estimates of VPD across the Amazon region, it should be included in future drought assessments (e.g. Castro et al., 2020). One possibility to account for the influences of VPD is choosing temporal and spatially

**Deleted:** Furthermore, in the last decade, new methods have been developed that assess impacts of drought on ecosystems, e.g. analyses based on solar-induced fluorescence (SIF) data show that tall forests are less sensitive to rainfall compared to short forests (Giardina et al., 2018). Also, vegetation optical depth (VOD) used as a proxy for water content in forests is a promising satellite-derived indicator for mortality and impacts of droughts on forests (Rao et al., 2019). However, conducting analyses over the Amazon rainforest based on VOD is difficult, because of the limited penetration depth of microwaves in dense tropical forests (Chaparro et al. 2019), and the influences of vegetation water status (Xu et al. 2021). So far, VOD data could only be applied with limited success across tropical rainforests (Konings & Gentine, 2017). Future studies should estimate the impacts of droughts based on multiple drought characteristics. For example, Toomey et al. (2011) show that considering both, heat stress and soil moisture stress greatly improves the explanatory power of drought impacts in the Amazon basin. ¶

#### ¶ Precipitation datasets¶

For the three drought events in 2005, 2010 and 2016, CHIRPS, GLDAS and ERA5 diverted the most from the other datasets regarding the spatial drought extent. ERA5 shows the smallest area of moderate drought stress during 2005 but one of the largest areas in 2010 (Fig. 2). We found no obvious bias between the precipitation datasets regarding distribution and frequency of monthly rainfall (Fig. S2...

based on the same satellite data as their input, they differ regarding the size of the drought area, especially during 2016 (Fig. 2). Lewis et al. (2011) estimated an area of 47% (2.5 million km<sup>2</sup>) of the Amazon basin moderately affected in 2005 using the TRMM6 dataset, which compares well with the size of the affected area for the majority of datasets analyzed in our study (considering our 0.6 million km<sup>2</sup> larger study area; see Methods). For 2010, Lewis et al. (2011) reported an area of 3.2 million km<sup>2</sup> being affected in comparison to 4.5 million km<sup>2</sup> in our analysis using TRMM6 with very similar spatial patterns. The newer product, TRMM7, however, shows less frequent rainfall but heavier rainfall than CHIRPS maintaining a similar total amount of precipitation (Giles et al., 2020). Also, both versions (TRMM6 and TRMM7) differ regarding the total area affected by drought stress in 2005 and in particular in 2010, where TRMM6 showed a 10% larger area of the Amazon basin affected. This can be explained by the generally higher precipitation rates detected in the TRMM7 dataset in comparison to TRMM6 (Seto et al., 2011) leading to lower absolute values of rMCWD. Spatially, this difference was most pronounced in the western and northern parts of Amazonia, in the *Acre* and *Roraima* states, and in Peru. Because of such higher precipitation rates in TRMM7 as compared to TRMM6, and subsequently the much stronger drought response according to our analysis, studies based on TRMM6 only might overstate the actual drought conditions and should be revisited. Precipitation datasets usually show remarkable differences in the representation of occurrence, frequency, intensity and location of events, mainly due to their nature of high spatial and temporal variability (Covey et al., 2016; Dirmeyer et al., 2012). Generally, the sparse network of observations in the Amazon rainforest may explain the differences across precipitation datasets and drought indices for datasets that rely on station data. Within the last decade, the number of observations increased, due to a new denser network of stations. [This may improve the reanalysis models that are used for several precipitation datasets applied here, however, it does not improve datasets that only rely on gauge observations. Bias-correction is also applied different across precipitation datasets. So doe CRUNCEP and WATCH\\_WFDEI use two different gridded bias corrections inputs, while the simulated precipitation fields of ERA5 are not using any bias corrections. Different datasets that are used for bias corrections can give very different results on regional scales \(Doblas-Reyes et al., 2021\)](#)

Jiménez-Muñoz et al. (2016) quantified drought extend using the scPDSI and found that 40%, 25% and 10% of the Amazon basin were affected by moderate, severe, and extreme drought stress, respectively, in March 2016. While we did not evaluate scPDSI directly but focused on  $r_{scPDSI}$  to allow for a better cross-comparison to the other drought indicators, we found similar patterns for moderate drought stress (47% of the basin affected), but different patterns under severe (11%) and extreme (1%) drought stress when evaluating  $r_{scPDSI}$  using the ERA5 dataset. Our estimation diverted from the results of Jiménez-Muñoz et al. (2016) mainly because of our different drought classification, but also due to a different reference area (see Methods).

In addition, Jiménez-Muñoz et al. (2016) used spatially resolved information on soil water capacity when calculating scPDSI and a longer baseline period (year onset is 1979 in their study vs. 2000 in our study). Furthermore, the choice of the precipitation dataset plays an important role. [Compared to the datasets considered in our study, ERA 5 showed the weakest drought impact during the 2016 drought event. GLDAS and TRMM7 showed a much stronger drought impact with over 70% of the area moderately and between 15% and 39% severely affected \(Tab. S2\). This is particularly interesting because recent](#)

**Deleted:** (Giles et al., 2020).

**Deleted:** (Covey et al., 2016; Dirmeyer et al., 2012).

**Deleted:**

**Deleted:** This may improve the reanalysis models that are used for several precipitation datasets applied here, however, it does not improve datasets that only rely on gauge observations.

**Formatted:** English (UK)

**Deleted:**  $r_{scPDSI}$

**Formatted:** English (UK)

**Deleted:**  $r_{scPDSI}$

**Deleted:** Compared to the datasets considered in our study, ERA 5 showed the weakest drought impact during the 2016 drought event. GLDAS and TRMM7 showed a much stronger drought impact with over 70% of the area moderately and between 15% and 39% severely affected (Table S3). This is particularly interesting because recent studies identify TRMM7, CHIRPS and ERA5 as the best precipitation datasets when comparing to gauge observations in South America (Albergel et al., 2018; Burton et al., 2018; Rifai et al., 2019)....

studies identify TRMM7, CHIRPS and ERA5 as the best precipitation datasets when comparing to gauge observations in South America (Albergel et al., 2018; Burton et al., 2018; Rifai et al., 2019). The higher scPDSI variability across the precipitation datasets can be explained with the more complex algorithm (including the self-calibrating mechanism) the index has compared to MCWD and RAI.

Formatted: English (UK)

Formatted: English (UK)

Formatted: Font: Not Bold

#### Evapotranspiration datasets

Using a variable ET dataset over constant ET of 100mm month<sup>-1</sup> leads to smaller areas affected by drought stress depending on the year and drought location (Fig. 4). According to our findings using a constant ET of 100mm month<sup>-1</sup> introduces not only a change in drought-affected areas, but rather a bias, as drought intensity and spatial extent are consistently higher for all drought years. The reason for this bias lies within the calculation of MCWD which computes stronger deficits for higher values of ET (e.g. 100 mm month<sup>-1</sup>) than for lower values (e.g. 50 mm month<sup>-1</sup>) during months with low precipitation. This bias can be rather small during drought events that are located in the northern, wetter parts of the basin (as in 2005), but it can also be quite strong for droughts that are located in the southern parts which have a more pronounced dry season (as in 2010).

#### Implications for drought impact analyses in the Amazon rainforest

Drought leads to increased tree mortality and carbon losses in tropical forests (Hubau et al., 2020; Lewis et al., 2011; Phillips et al., 2009). With the prospect of more severe and frequent droughts in a future climate, more precise estimates of how much carbon is lost from reductions in growth and drought-induced mortality are necessary. Currently, the Amazon rainforest is acting as a carbon sink, thereby removing CO<sub>2</sub> from the atmosphere, but with more frequent and severe drought events, this sink is already declining (Hubau et al. 2020). Lewis et al. (2011) estimated a total loss of biomass for the Amazon basin in 2005 of 1.6 Pg C and a 38% more severe impact of 2.2 PgC for 2010 based on TRMM6. [Using TRMM7 instead of TRMM6 and using variable ET would likely decrease the impact of the 2010 drought on vegetation carbon as calculated in Lewis et al. 2011.](#)

Deleted: estimating

Deleted: impacts on the carbon cycle of

Deleted: Later studies however found that the relationship between  $\alpha$ MCWD and  $\alpha$ AGB does not hold for the 2010 drought event (Feldpausch et al., 2016). When applied to the  $\alpha$ MCWD derived from the precipitation datasets in our study, we still can calculate the loss of biomass of the 2005 drought event to be in the range of 0.7-1.7 PgC (Fig. 4). This is in the range of the regional average annual carbon uptake (1-2 PgC) per year, and thus, has the potential to turn the carbon sink into a carbon source. We acknowledge that our estimates are based on a relatively simple, empirically derived relation that does not take the biomass variability across the whole Amazon basin and individual forest/tree responses to drought into account. It however gives a rough estimation of potential carbon losses during drought and an idea of how much it varies depending on the precipitation datasets applied in a study. In addition, we would like to note that the empirical biomass  $\alpha$ MCWD relation from Lewis et al. (2011) has been estimated with constant ET=100 mm. When using evapotranspiration data (from ERA5) for the calculation of  $\alpha$ MCWD, we find generally lower biomass losses (between 10-20% lower, Fig. S3), and thus, the use of MCWD should be carefully viewed via its sensitivity to ET. While previous studies found that the MCWD calculation can be quite robust, in our analysis, MCWD is sensitive to the evapotranspiration input and baseline period (Fig. S3).

Deleted: Furthermore, our totally

Deleted:

Deleted: .

Deleted: (Feldpausch et al., 2016).

The affected areas (Fig. 2) for the drought events might be underestimated as (1) the total duration of the 2016 drought was longer than 12 months (see above paragraph and Fig. 6) and can hence not be fully captured by the standard 12-month period of the  $\alpha$ MCWD and  $r$ MCWD calculation used in this study, (2) Potential lag effects due to delayed plant mortality within the subsequent years are not considered so far. We would recommend for future studies to investigate the relationship of biomass losses with other drought indices (such as scPDSI) in a similar manner as done in Lewis et al. (2011). As the biomass of the Amazon rainforest is heterogeneously distributed (e.g. Saatchi et al., 2011), large-scale drought-induced biomass losses which result from a severe  $\alpha$ MCWD anomaly should be interpreted carefully. Differences in the amount of biomass in different forest types, species composition, and critical hydraulic processes should be considered when estimating potential biomass losses under drought stress (Feldpausch et al., 2016). A step forward would be to use, for example, remotely sensed biomass maps to account for regional biomass distributions (e.g. Avitabile et al., 2016) or to simulate drought impacts with dynamic global

vegetation models (DGVMs). DGVMs simulate the carbon- and water cycle of the biosphere in a process-based way, accounting for the interplay of carbon uptake and water loss through stomatal opening, evapotranspiration (ET), carbon assimilation via photosynthesis, and carbon allocation to different plant compartments such as leaves, wood, and roots (e.g. Schaphoff et al., 2018; Smith et al., 2014). The simulated response of tropical forests in DGVMs is particularly sensitive to precipitation input under present and future climate change scenarios (e.g. Seiler et al., 2015). Therefore, we recommend using multiple climate forcing datasets to test for climate data uncertainty also under present climate conditions. Particularly, studies based on TRMM6 should possibly be revisited and complemented with more forcing datasets for their analysis.

## 6. Conclusions

We find substantial variation in the spatial extent, location, and timing of the extreme drought events in the years 2005, 2010 and 2016 in the Amazon basin. Depending on the precipitation dataset and drought index used, the area affected by severe (extreme) drought varied between 0% and 39% (0% and 13.7%) for the 2016 event. Especially the area under severe drought conditions changed from almost no severe drought stress (5 out of 6 datasets) when using  $rRAI$  to greater than 10% when using  $rMCWD$  and  $rscPDSI$  instead. The variation partly results from the application of different drought metrics ( $rMCWD$ ,  $rRAI$ ,  $rscPDSI$ ) and from differences in the underlying precipitation datasets. Such differences also propagate when quantifying the impacts of droughts on the carbon cycle of the Amazon rainforest and result in a large variability in biomass carbon losses for a particular drought year. The estimated intensity of droughts depends predominantly on the selected drought indicator and to a lesser extent on the choices of precipitation and evapotranspiration dataset.

We, therefore, recommend applying several drought metrics, climate (precipitation) datasets and, if available, evapotranspiration datasets to account for model uncertainty when assessing the spatial extent, duration, and location of droughts. We regard it as an important step when assessing drought impacts on tropical rainforests also under current climate conditions. Communicating the uncertainty in the estimation of drought events and their impacts on the Amazon rainforest is highly relevant and thus, multiple datasets should be applied by any large-scale study on drought impacts on vegetation.

## 7. Code availability

All scripts to reproduce analysis and figures are available at <https://github.com/PhillipPapastefanou/DroughtAnalysis>

## 8. Data availability

All datasets are available following the references in the method section.

## 9. Author contribution

P.P. and A.R. conceived the study and wrote the first draft of the manuscript. All authors contributed to the development of the analysis and the writing of the manuscript.


## 10. Competing interests

Formatted: English (UK)

Deleted: We found the biomass loss to vary between 0.7 and 1.6 PgC during the 2005 drought depending on the precipitation forcing.

Deleted: as well as drought metrics

The authors declare no competing interests.



## 11. References

- Albergel, C., Dutra, E., Munier, S., Calvet, J.-C., Muñoz-Sabater, J., de Rosnay, P., & Balsamo, G. (2018). ERA-5 and ERA-Interim driven ISBA land surface model simulations: Which one performs better? *Hydrology and Earth System Sciences*, 22(6), 3515–3532. <https://doi.org/10.5194/hess-22-3515-2018>
- Aragão, L. E. O. C., Malhi, Y., Roman-Cuesta, R. M., Saatchi, S., Anderson, L. O., & Shimabukuro, Y. E. (2007). Spatial patterns and fire response of recent Amazonian droughts. *Geophysical Research Letters*, 34(7), L07701. <https://doi.org/10.1029/2006GL028946>
- Barkhordarian, A., Saatchi, S. S., Behrangi, A., Loikith, P. C., & Mechoso, C. R. (2019). A Recent Systematic Increase in Vapor Pressure Deficit over Tropical South America. *Scientific Reports*, 9(1), 15331. <https://doi.org/10.1038/s41598-019-51857-8>
- Blacutt, L. A., Herdies, D. L., de Gonçalves, L. G. G., Vila, D. A., & Andrade, M. (2015). Precipitation comparison for the CFSR, MERRA, TRMM3B42 and Combined Scheme datasets in Bolivia. *Atmospheric Research*, 163, 117–131. <https://doi.org/10.1016/j.atmosres.2015.02.002>
- Burton, C., Rifai, S., & Malhi, Y. (2018). Inter-comparison and assessment of gridded climate products over tropical forests during the 2015/2016 El Niño. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170406. <https://doi.org/10.1098/rstb.2017.0406>
- Cai, W., Santoso, A., Wang, G., Yeh, S.-W., An, S.-I., Cobb, K. M., Collins, M., Guilyardi, E., Jin, F.-F., Kug, J.-S., Lengaigne, M., McPhaden, M. J., Takahashi, K., Timmermann, A., Vecchi, G., Watanabe, M., & Wu, L. (2015). ENSO and greenhouse warming. *Nature Climate Change*, 5(9), 849–859. <https://doi.org/10.1038/nclimate2743>
- Castro, A. O., Chen, J., Zang, C. S., Shekhar, A., Jimenez, J. C., Bhattacharjee, S., Kindu, M., Morales, V. H., & Rammig, A. (2020). OCO-2 Solar-Induced Chlorophyll Fluorescence Variability across Ecoregions of the Amazon Basin and the Extreme Drought Effects of El Niño (2015–2016). *Remote Sensing*, 12(7), 1202–1202. <https://doi.org/10.3390/rs12071202>

- Chaparro, D., Duveiller, G., Piles, M., Cescatti, A., Vall-Ilossera, M., Camps, A., & Entekhabi, D. (2019). Sensitivity of L-band vegetation optical depth to carbon stocks in tropical forests: A comparison to higher frequencies and optical indices. *Remote Sensing of Environment*, 232, 111303. <https://doi.org/10.1016/j.rse.2019.111303>
- Coelho, C. A. S., Cavalcanti, J. A. F., Costa, S. M. S., Freitas, S. R., Ito, E. R., Luz, G., Santos, A. F., Nobre, C. A., Marengo, J. A., & Pezza, A. B. (2012). Climate diagnostics of three major drought events in the Amazon and illustrations of their seasonal precipitation predictions. *Meteorological Applications*, 19(2), 237–255. <https://doi.org/10.1002/met.1324>
- Compo, G. P., Sardeshmukh, P. D., Whitaker, J. S., Brohan, P., Jones, P. D., & McColl, C. (2013). Independent confirmation of global land warming without the use of station temperatures. *Geophysical Research Letters*, 40(12), 3170–3174. <https://doi.org/10.1002/grl.50425>
- Covey, C., Gleckler, P. J., Doutriaux, C., Williams, D. N., Dai, A., Fasullo, J., Trenberth, K., & Berg, A. (2016). Metrics for the Diurnal Cycle of Precipitation: Toward Routine Benchmarks for Climate Models. *Journal of Climate*, 29(12), 4461–4471. <https://doi.org/10.1175/JCLI-D-15-0664.1>
- da Rocha, H. R., Goulden, M. L., Miller, S. D., Menton, M. C., Pinto, L. D. V. O., de Freitas, H. C., & e Silva Figueira, A. M. (2004). SEASONALITY OF WATER AND HEAT FLUXES OVER A TROPICAL FOREST IN EASTERN AMAZONIA. *Ecological Applications*, 14(sp4), 22–32. <https://doi.org/10.1890/02-6001>
- Dirmeyer, P. A., Cash, B. A., Kinter, J. L., Jung, T., Marx, L., Satoh, M., Stan, C., Tomita, H., Towers, P., Wedi, N., Achuthavarier, D., Adams, J. M., Altschuler, E. L., Huang, B., Jin, E. K., & Manganello, J. (2012). Simulating the diurnal cycle of rainfall in global climate models: Resolution versus parameterization. *Climate Dynamics*, 39(1–2), 399–418. <https://doi.org/10.1007/s00382-011-1127-9>
- Dirmeyer, P. A., Schlosser, C. A., & Brubaker, K. L. (2009). Precipitation, Recycling, and Land Memory: An Integrated Analysis. *Journal of Hydrometeorology*, 10(1), 278–288. <https://doi.org/10.1175/2008JHM1016.1>
- Doblas-Reyes, F. J., Sorensson, A. A., Almazroui, M., Dosio, A., Gutowski, W. J., Haarsma, R., Hamdi, R., Hewitson, B., Kwon, W.-T., Lamptey, B. L., Maraun, D., Stephenson, T. S., Takayabu, I., Terray, L., Turner, A., & Zuo, Z. (2021). *Linking global to regional climate change* (V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Pean, S. Berger,

- N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekci, R. Yu, & B. Zhou, Eds.). Cambridge University Press. <https://centaur.reading.ac.uk/99896/>
- Döll, P., & Lehner, B. (2002). Validation of a new global 30-min drainage direction map. *Journal of Hydrology*, 258(1–4), 214–231. [https://doi.org/10.1016/S0022-1694\(01\)00565-0](https://doi.org/10.1016/S0022-1694(01)00565-0)
- Espinoza, J. C., Sörensson, A. A., Ronchail, J., Molina-Carpio, J., Segura, H., Gutierrez-Cori, O., Ruscica, R., Condom, T., & Wongchuig-Correa, S. (2019). Regional hydro-climatic changes in the Southern Amazon Basin (Upper Madeira Basin) during the 1982–2017 period. *Journal of Hydrology: Regional Studies*, 26, 100637. <https://doi.org/10.1016/j.ejrh.2019.100637>
- Esquivel-Muelbert, A., Baker, T. R., Dexter, K. G., Lewis, S. L., Brienen, R. J. W., Feldpausch, T. R., Lloyd, J., Monteagudo-Mendoza, A., Arroyo, L., Álvarez-Dávila, E., Higuchi, N., Marimon, B. S., Marimon-Junior, B. H., Silveira, M., Vilanova, E., Gloor, E., Malhi, Y., Chave, J., Barlow, J., ... Phillips, O. L. (2019). Compositional response of Amazon forests to climate change. *Global Change Biology*, 25(1), 39–56. <https://doi.org/10.1111/gcb.14413>
- Feldpausch, T. R., Phillips, O. L., Brienen, R. J. W., Gloor, E., Lloyd, J., Lopez-Gonzalez, G., Monteagudo-Mendoza, A., Malhi, Y., Alarcón, A., Álvarez Dávila, E., Alvarez-Loayza, P., Andrade, A., Aragao, L. E. O. C., Arroyo, L., Aymard C., G. A., Baker, T. R., Baraloto, C., Barroso, J., Bonal, D., ... Vos, V. A. (2016). Amazon forest response to repeated droughts. *Global Biogeochemical Cycles*, 30(7), 964–982. <https://doi.org/10.1002/2015GB005133>
- Flack-Prain, S., Meir, P., Malhi, Y., Smallman, T. L., & Williams, M. (2019). The importance of physiological, structural and trait responses to drought stress in driving spatial and temporal variation in GPP across Amazon forests. *Biogeosciences*, 16(22), 4463–4484. <https://doi.org/10.5194/bg-16-4463-2019>
- Forkel, M., Drüke, M., Thurner, M., Dorigo, W., Schaphoff, S., Thonicke, K., von Bloh, W., & Carvalhais, N. (2019). Constraining modelled global vegetation dynamics and carbon turnover using multiple satellite observations. *Scientific Reports*, 9(1), 18757. <https://doi.org/10.1038/s41598-019-55187-7>
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., & Michaelsen, J. (2015). The climate hazards infrared precipitation with stations—A new environmental record for monitoring extremes. *Scientific Data*, 2(1), 150066. <https://doi.org/10.1038/sdata.2015.66>

- [Giardina, F., Konings, A. G., Kennedy, D., Alemohammad, S. H., Oliveira, R. S., Uriarte, M., & Gentine, P. \(2018\). Tall Amazonian forests are less sensitive to precipitation variability. \*Nature Geoscience\*, 11\(6\), 405–409. <https://doi.org/10.1038/s41561-018-0133-5>](#)
- [Giles, J. A., Ruscica, R. C., & Menéndez, C. G. \(2020\). The diurnal cycle of precipitation over South America represented by five gridded datasets. \*International Journal of Climatology\*, 40\(2\), 668–686. <https://doi.org/10.1002/joc.6229>](#)
- [Gloor, M., Barichivich, J., Ziv, G., Brienen, R., Schöngart, J., Peylin, P., Ladvocat Cintra, B. B., Feldpausch, T., Phillips, O., & Baker, J. \(2015\). Recent Amazon climate as background for possible ongoing and future changes of Amazon humid forests. \*Global Biogeochemical Cycles\*, 29\(9\), 1384–1399. <https://doi.org/10.1002/2014GB005080>](#)
- [Golian, S., Javadian, M., & Behrangi, A. \(2019\). On the use of satellite, gauge, and reanalysis precipitation products for drought studies. \*Environmental Research Letters\*, 14\(7\), 075005. <https://doi.org/10.1088/1748-9326/ab2203>](#)
- [Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., Sperry, J. S., & McDowell, N. G. \(2020\). Plant responses to rising vapor pressure deficit. \*New Phytologist\*, 226\(6\), 1550–1566. <https://doi.org/10.1111/nph.16485>](#)
- [Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. \(2014\). Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset: UPDATED HIGH-RESOLUTION GRIDS OF MONTHLY CLIMATIC OBSERVATIONS. \*International Journal of Climatology\*, 34\(3\), 623–642. <https://doi.org/10.1002/joc.3711>](#)
- [Hobeichi, S., Abramowitz, G., Evans, J., & Ukkola, A. \(2018\). Derived Optimal Linear Combination Evapotranspiration \(DOLCE\): A global gridded synthesis ET estimate. \*Hydrology and Earth System Sciences\*, 22\(2\), 1317–1336. <https://doi.org/10.5194/hess-22-1317-2018>](#)
- [Hubau, W., Lewis, S. L., Phillips, O. L., Affum-Baffoe, K., Beeckman, H., Cuni-Sanchez, A., Daniels, A. K., Ewango, C. E. N., Fauset, S., Mukinzi, J. M., Sheil, D., Sonké, B., Sullivan, M. J. P., Sunderland, T. C. H., Taedoumg, H., Thomas, S. C., White, L. J. T., Abernethy, K. A., Adu-Bredu, S., ... Zomagho, L. \(2020\). Asynchronous carbon sink saturation in African and Amazonian tropical forests. \*Nature\*, 579\(7797\), 80–87. <https://doi.org/10.1038/s41586-020-2035-0>](#)

- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P., & Stocker, E. F. (2007). The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales. *Journal of Hydrometeorology*, 8(1), 38–55. <https://doi.org/10.1175/JHM560.1>
- Jiang, M., Medlyn, B. E., Drake, J. E., Duursma, R. A., Anderson, I. C., Barton, C. V. M., Boer, M. M., Carrillo, Y., Castañeda-Gómez, L., Collins, L., Crous, K. Y., De Kauwe, M. G., dos Santos, B. M., Emmerson, K. M., Facey, S. L., Gherlenda, A. N., Gimeno, T. E., Hasegawa, S., Johnson, S. N., ... Ellsworth, D. S. (2020). The fate of carbon in a mature forest under carbon dioxide enrichment. *Nature*, 580(7802), 227–231. <https://doi.org/10.1038/s41586-020-2128-9>
- Jimenez, J. C., Barichivich, J., Mattar, C., Takahashi, K., Santamaría-Artigas, A., Sobrino, J. A., & Malhi, Y. (2018). Spatio-temporal patterns of thermal anomalies and drought over tropical forests driven by recent extreme climatic anomalies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170300. <https://doi.org/10.1098/rstb.2017.0300>
- Jimenez, J. C., Marengo, J. A., Alves, L. M., Sulca, J. C., Takahashi, K., Ferrett, S., & Collins, M. (2019). The role of ENSO flavours and TNA on recent droughts over Amazon forests and the Northeast Brazil region. *International Journal of Climatology*, 41(7), 3761–3780. <https://doi.org/10.1002/joc.6453>
- Jiménez-Muñoz, J. C., Mattar, C., Barichivich, J., Santamaría-Artigas, A., Takahashi, K., Malhi, Y., Sobrino, J. A., & Schrier, G. van der. (2016). Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–2016. *Scientific Reports*, 6(1), 33130. <https://doi.org/10.1038/srep33130>
- Koch, A., Hubau, W., & Lewis, S. L. (2021). Earth System Models Are Not Capturing Present-Day Tropical Forest Carbon Dynamics. *Earth's Future*, 9(5). <https://doi.org/10.1029/2020EF001874>
- Konings, A. G., & Gentine, P. (2017). Global variations in ecosystem-scale isohydricity. *Global Change Biology*, 23(2), 891–905. <https://doi.org/10.1111/gcb.13389>
- Lewis, S. L., Brando, P. M., Phillips, O. L., van der Heijden, G. M. F., & Nepstad, D. (2011). The 2010 Amazon Drought. *Science*, 331(6017), 554–554. <https://doi.org/10.1126/science.1200807>

- [Malhi, Y., Aragao, L. E. O. C., Galbraith, D., Huntingford, C., Fisher, R., Zelazowski, P., Sitch, S., McSweeney, C., & Meir, P. \(2009\). Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. \*Proceedings of the National Academy of Sciences\*, 106\(49\), 20610–20615. <https://doi.org/10.1073/pnas.0804619106>](#)
- [Malhi, Y., Roberts, J. T., Betts, R. A., Killeen, T. J., Li, W., & Nobre, C. A. \(2008\). Climate Change, Deforestation, and the Fate of the Amazon. \*Science\*, 319\(5860\), 169–172. <https://doi.org/10.1126/science.1146961>](#)
- [Marengo, J. A., & Espinoza, J. C. \(2016\). Extreme seasonal droughts and floods in Amazonia: Causes, trends and impacts: EXTREMES IN AMAZONIA. \*International Journal of Climatology\*, 36\(3\), 1033–1050. <https://doi.org/10.1002/joc.4420>](#)
- [Marengo, J. A., Nobre, C. A., Tomasella, J., Cardoso, M. F., & Oyama, M. D. \(2008\). Hydro-climatic and ecological behaviour of the drought of Amazonia in 2005. \*Philosophical Transactions of the Royal Society B: Biological Sciences\*, 363\(1498\), 1773–1778. <https://doi.org/10.1098/rstb.2007.0015>](#)
- [Marengo, J. A., Nobre, C. A., Tomasella, J., Oyama, M. D., Sampaio de Oliveira, G., de Oliveira, R., Camargo, H., Alves, L. M., & Brown, I. F. \(2008\). The Drought of Amazonia in 2005. \*Journal of Climate\*, 21\(3\), 495–516. <https://doi.org/10.1175/2007JCLI1600.1>](#)
- [Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R., & Rodriguez, D. A. \(2011\). The drought of 2010 in the context of historical droughts in the Amazon region: DROUGHT AMAZON 2010. \*Geophysical Research Letters\*, 38\(12\), n/a–n/a. <https://doi.org/10.1029/2011GL047436>](#)
- [Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., & Verhoest, N. E. C. \(2017\). GLEAM v3: Satellite-based land evaporation and root-zone soil moisture. \*Geoscientific Model Development\*, 10\(5\), 1903–1925. <https://doi.org/10.5194/gmd-10-1903-2017>](#)
- [Miralles, D. G., Gentile, P., Seneviratne, S. I., & Teuling, A. J. \(2019\). Land-atmospheric feedbacks during droughts and heatwaves: State of the science and current challenges: Land feedbacks during droughts and heatwaves. \*Annals of the New York Academy of Sciences\*, 1436\(1\), 19–35. <https://doi.org/10.1111/nyas.13912>](#)
- [Muñoz-Sabater, J., Dutra, E., Balsamo, G., Boussetta, S., Zsoter, E., Albergel, C., & Agusti-Panareda, A. \(2018\). ERA5-Land: An improved version of the ERA5 reanalysis land component.](#)

- Nogueira, M. (2020). Inter-comparison of ERA-5, ERA-interim and GPCP rainfall over the last 40 years: Process-based analysis of systematic and random differences. *Journal of Hydrology*, 583, 124632. <https://doi.org/10.1016/j.jhydrol.2020.124632>
- Phillips, O. L., Aragão, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., López-González, G., Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C. A., van der Heijden, G., Almeida, S., Amaral, I., Arroyo, L., Aymard, G., Baker, T. R., Bánki, O., Blanc, L., Bonal, D., ... Torres-Lezama, A. (2009). Drought Sensitivity of the Amazon Rainforest. *Science*, 323(5919), 1344–1347. <https://doi.org/10.1126/science.1164033>
- Rao, K., Anderegg, W. R. L., Sala, A., Martínez-Vilalta, J., & Konings, A. G. (2019). Satellite-based vegetation optical depth as an indicator of drought-driven tree mortality. *Remote Sensing of Environment*, 227, 125–136. <https://doi.org/10.1016/j.rse.2019.03.026>
- Rifai, S. W., Li, S., & Malhi, Y. (2019). Coupling of El Niño events and long-term warming leads to pervasive climate extremes in the terrestrial tropics. *Environmental Research Letters*, 14(10), 105002. <https://doi.org/10.1088/1748-9326/ab402f>
- Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., & Toll, D. (2004). The Global Land Data Assimilation System. *Bulletin of the American Meteorological Society*, 85(3), 381–394. <https://doi.org/10.1175/BAMS-85-3-381>
- Ruida, Z., Chen, X., Wang, Z., Lai, C., & Goddard, S. (2018). *Package scPDSI*. <https://github.com/Sibada/scPDSI>
- Schneider, U., Becker, A., Finger, P., Anja, M.-C., & Markus, Z. (2018). *GPCC Full Data Monthly Version 2018.0 at 0.5°: Monthly Land-Surface Precipitation from Rain-Gauges built on GTS-based and Historic Data*. [https://doi.org/10.5676/DWD\\_GPCC/FD\\_M\\_V2018\\_050](https://doi.org/10.5676/DWD_GPCC/FD_M_V2018_050)
- Seto, S., Iguchi, T., & Meneghini, R. (2011). Comparison of TRMM PR V6 and V7 focusing heavy rainfall. 2011 *IEEE International Geoscience and Remote Sensing Symposium*, 2582–2585. <https://doi.org/10.1109/IGARSS.2011.6049769>
- Sheffield, J., Goteti, G., & Wood, E. F. (2006). Development of a 50-Year High-Resolution Global Dataset of Meteorological Forcings for Land Surface Modeling. *Journal of Climate*, 19(13), 3088–3111. <https://doi.org/10.1175/JCLI3790.1>

- Sörensson, A. A., & Ruscica, R. C. (2018). Intercomparison and Uncertainty Assessment of Nine Evapotranspiration Estimates Over South America. *Water Resources Research*, 54(4), 2891–2908. <https://doi.org/10.1002/2017WR021682>
- Staal, A., Fetzer, I., Wang-Erlandsson, L., Bosmans, J. H. C., Dekker, S. C., van Nes, E. H., Rockström, J., & Tuinenburg, O. A. (2020). Hysteresis of tropical forests in the 21st century. *Nature Communications*, 11(1), 4978. <https://doi.org/10.1038/s41467-020-18728-7>
- Toomey, M., Roberts, D. A., Still, C., Goulden, M. L., & McFadden, J. P. (2011). Remotely sensed heat anomalies linked with Amazonian forest biomass declines: AMAZON DROUGHT THERMAL ANOMALIES. *Geophysical Research Letters*, 38(19), n/a-n/a. <https://doi.org/10.1029/2011GL049041>
- van der Ent, R. J., Savenije, H. H. G., Schaeffli, B., & Steele-Dunne, S. C. (2010). Origin and fate of atmospheric moisture over continents: ORIGIN AND FATE OF ATMOSPHERIC MOISTURE. *Water Resources Research*, 46(9). <https://doi.org/10.1029/2010WR009127>
- Viovy, N. (2018). *CRUNCEP Version 7—Atmospheric Forcing Data for the Community Land Model*. <http://rda.ucar.edu/datasets/ds314.3/0%22>
- von Randow, C., Manzi, A. O., Kruijt, B., de Oliveira, P. J., Zanchi, F. B., Silva, R. L., Hodnett, M. G., Gash, J. H. C., Elbers, J. A., Waterloo, M. J., Cardoso, F. L., & Kabat, P. (2004). Comparative measurements and seasonal variations in energy and carbon exchange over forest and pasture in South West Amazonia. *Theoretical and Applied Climatology*, 78(1–3). <https://doi.org/10.1007/s00704-004-0041-z>
- Weedon, G. P., Balsamo, G., Bellouin, N., Gomes, S., Best, M. J., & Viterbo, P. (2014). The WFDEI meteorological forcing data set: WATCH Forcing Data methodology applied to ERA-Interim reanalysis data. *Water Resources Research*, 50(9), 7505–7514. <https://doi.org/10.1002/2014WR015638>
- Weedon, G. P., Gomes, S., Viterbo, P., Shuttleworth, W. J., Blyth, E., Österle, H., Adam, J. C., Bellouin, N., Boucher, O., & Best, M. (2011). Creation of the WATCH Forcing Data and Its Use to Assess Global and Regional Reference Crop Evaporation over Land during the Twentieth Century. *Journal of Hydrometeorology*, 12(5), 823–848. <https://doi.org/10.1175/2011JHM1369.1>

- [Wells, N., Goddard, S., & Hayes, M. J. \(2004\). A Self-Calibrating Palmer Drought Severity Index. \*Journal of Climate\*, 17\(12\), 2335–2351. \[https://doi.org/10.1175/1520-0442\\(2004\\)017<2335:ASPDSI>2.0.CO;2\]\(https://doi.org/10.1175/1520-0442\(2004\)017<2335:ASPDSI>2.0.CO;2\)](#)
- [Willmott, C. J., Rowe, C. M., & Philpot, W. D. \(1985\). Small-Scale Climate Maps: A Sensitivity Analysis of Some Common Assumptions Associated with Grid-Point Interpolation and Contouring. \*The American Cartographer\*, 12\(1\), 5–16. <https://doi.org/10.1559/152304085783914686>](#)
- [Xu, X., Konings, A. G., Longo, M., Feldman, A., Xu, L., Saatchi, S., Wu, D., Wu, J., & Moorcroft, P. \(2021\). Leaf surface water, not plant water stress, drives diurnal variation in tropical forest canopy water content. \*New Phytologist\*, 231\(1\), 122–136. <https://doi.org/10.1111/nph.17254>](#)
- [Yang, H., Piao, S., Zeng, Z., Ciais, P., Yin, Y., Friedlingstein, P., Sitch, S., Ahlström, A., Guimberteau, M., Huntingford, C., Levis, S., Levy, P. E., Huang, M., Li, Y., Li, X., Lomas, M. R., Peylin, P., Poulter, B., Viovy, N., ... Wang, L. \(2015\). Multicriteria evaluation of discharge simulation in Dynamic Global Vegetation Models. \*Journal of Geophysical Research: Atmospheres\*, 120\(15\), 7488–7505. <https://doi.org/10.1002/2015JD023129>](#)
- [Yang, Y., Saatchi, S. S., Xu, L., Yu, Y., Choi, S., Phillips, N., Kennedy, R., Keller, M., Knyazikhin, Y., & Myneni, R. B. \(2018\). Post-drought decline of the Amazon carbon sink. \*Nature Communications\*, 9\(1\), 3172. <https://doi.org/10.1038/s41467-018-05668-6>](#)
- [Zang, C. S., Buras, A., Esquivel-Muelbert, A., Jump, A. S., Rigling, A., & Rammig, A. \(2020\). Standardized drought indices in ecological research: Why one size does not fit all. \*Global Change Biology\*, 26\(2\), 322–324. <https://doi.org/10.1111/gcb.14809>](#)
- [Zemp, D. C., Schleussner, C.-F., Barbosa, H. M. J., Hirota, M., Montade, V., Sampaio, G., Staal, A., Wang-Erlandsson, L., & Rammig, A. \(2017\). Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks. \*Nature Communications\*, 8\(1\), 14681. <https://doi.org/10.1038/ncomms14681>](#)
- [Zemp, D. C., Schleussner, C.-F., Barbosa, H. M. J., van der Ent, R. J., Donges, J. F., Heinke, J., Sampaio, G., & Rammig, A. \(2014\). On the importance of cascading moisture recycling in South America. \*Atmospheric Chemistry and Physics\*, 14\(23\), 13337–13359. <https://doi.org/10.5194/acp-14-13337-2014>](#)

Zeng, N., Yoon, J.-H., Marengo, J. A., Subramaniam, A., Nobre, C. A., Mariotti, A., & Neelin, J. D. (2008). Causes and impacts of the 2005 Amazon drought. *Environmental Research Letters*, 3(1), 014002. <https://doi.org/10.1088/1748-9326/3/1/014002>

Ziese, M., Schneider, U., Meyer-Christoffer, A., Schamm, K., Vido, J., Finger, P., Bissolli, P., Pietzsch, S., & Becker, A. (2014). The GPCP Drought Index – a new, combined and gridded global drought index. *Earth System Science Data*, 6(2), 285–295. <https://doi.org/10.5194/essd-6-285-2014>

Tables

Precipitation dataset	Abbreviation	Details	Resolutions	Derived from	References
Climate Hazards group Infrared Precipitation with Stations	CHIRPS	quasi-global (50°S-50°N)	high resolution (0.05°), daily, pentadal, and monthly	Remote sensing, in-situ observations	Funk et al., 2015
Tropical Rainfall Measurement Misson	TRMM v6 3b43	quasi-global (50°S-50°N)	Quarter degree resolution (0.25°) daily, pentadal, and monthly	Remote sensing	Huffman et al., 2007
Tropical Rainfall Measurement Misson	TRMM v7 3B43	quasi-global (50°S-50°N)	Quarter degree resolution (0.25°), daily, pentadal, and monthly	Remote sensing	Huffman et al., 2007
	CRU_NCEP V8	global	Half degree resolution (0.5°), daily, pentadal and monthly	<a href="#">Reanalysis corrected by CRU gridded observational dataset</a>	Viovy et al., 2017
ERA5		global	Quarter degree resolution (0.25°), sub-daily, daily, monthly	<a href="#">Reanalysis</a>	Albergel et al., 2018
Global Land Data Assimilation System	GLDAS 2.1	global	Quarter degree resolution (0.25°), daily, pentadal, and monthly	<a href="#">Geostationary satellite infrared cloud-top temperature</a>	Rodell et al., 2004

**Deleted:** Albergel, C., Dutra, E., Munier, S., Calvet, J.-C., Munoz-Sabater, J., de Rosnay, P., & Balsamo, G. (2018). ERA-5 and ERA-Interim driven ISBA land surface model simulations: which one performs better? *Hydrology and Earth System Sciences*, 22(6), 3515–3532. <https://doi.org/10.5194/hess-22-3515-2018>

Aragão, L. E. O. C., Malhi, Y., Roman-Cuesta, R. M., Saatchi, S., Anderson, L. O., & Shimabukuro, Y. E. (2007). Spatial patterns and fire response of recent Amazonian droughts. *Geophysical Research Letters*, 34(7). <https://doi.org/10.1029/2006GL028946>

Avitabile, V., Herold, M., Heuvelink, G. B. M., Lewis, S. L., Phillips, O. L., Asner, G. P., Armston, J., Ashton, P. S., Banin, L., Bayol, N., Berry, N. J., Boeckx, P., de Jong, B. H. J., DeVries, B., Girardin, C. A. J., Kearsley, E., Lindsell, J. A., Lopez-Gonzalez, G., Lucas, R., ... Willcock, S. (2016). An integrated pan-tropical biomass map using multiple reference datasets. *Global Change Biology*, 22(4), 1406–1420. <https://doi.org/10.1111/gcb.13139>

Barkhordarian, A., Saatchi, S. S., Behrang, A., Loikith, P. C., & Mechocho, C. R. (2019). A Recent Systematic Increase in Vapor Pressure Deficit over Tropical South America. *Scientific Reports*, 9(1), 15331. <https://doi.org/10.1038/s41598-019-51857-8>

Blacutt, L. A., Herdies, D. L., de Gonçalves, L. G. G., Vila, D. A., & Andrade, M. (2015). Precipitation comparison for the CFSR, MERRA, TRMM3B42 and Combined Scheme datasets in Bolivia. *Atmospheric Research*, 163, 117–131. <https://doi.org/10.1016/j.atmosres.2015.02.002>

Burton, C., Rifai, S., & Malhi, Y. (2018). Inter-comparison and assessment of gridded climate products over tropical forests during the 2015/2016 El Niño. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170406. <https://doi.org/10.1098/rstb.2017.0406>

Cai, W., Santos, A., Wang, G., Yeh, S.-W. W., An, S.-I. I., Cobb, K. M., Collins, M., Guilyardi, E., Jin, F.-F. F., Kug, J.-S. S., Lengaigne, M., McPhaden, M. J., Takahashi, K., Timmermann, A., Vecchi, G., Watanabe, M., & Wu, L. (2015). ENSO and greenhouse warming. *Nature Climate Change*, 5(9), 849–859. <https://doi.org/10.1038/nclimate2743>

Castro, A. O., Chen, J., Zang, C. S., Shekhar, A., Jimenez, J. C., Bhattacharjee, S., Kindu, M., Morales, V. H., & Rammig, A. (2020). OCO-2 Solar-Induced Chlorophyll Fluorescence Variability across Ecoregions of the Amazon Basin and the Extreme Drought Effects of El Niño (2015–2016). *Remote Sensing*, 12(7), 1202. <https://doi.org/10.3390/rs12071202>

Coelho, C. A. S., Cavalcanti, I. A. F., Costa, S. M. S., Freitas, S. R., Ito, E. R., Luz, G., Santos, A. F., Nobre, C. A., Marengo, J. A., & Pezza, A. B. (2012). Climate diagnostics of three major drought events in the Amazon and illustrations of their seasonal precipitation predictions. *Meteorological Applications*, 19(2), 237–255. <https://doi.org/10.1002/met.1324>

Compo, G. P., Sardeshmukh, P. D., Whitaker, J. S., Brohan, P., Jones, P. D., & McColl, C. (2013). Independent confirmation of global land warming without the use of station temperatures. *Geophysical Research Letters*, 40(12), 3170–3174. <https://doi.org/10.1002/grl.50425>

Covey, C., Gleckler, P. J., Doutriaux, C., Williams, D. N., Dai, A., Fasullo, J., Trenberth, K., & Berg, A. (2016). Metrics for the Diurnal Cycle of Precipitation: Toward Routine Benchmarking. [\[35\]](#)

**Formatted Table**

**Deleted:** Mainly in-situ observations

**Deleted:** Land surface models, remote sensing, in-situ observations

**Deleted:** Land surface models

				<a href="#">measurements and microwave observation techniques.</a>	
Global Precipitation Climatology Centre at Deutscher Wetterdienst	GPCC2018	global	Quarter degree resolution (0.25°), monthly	<a href="#">Gridded in-situ observations</a>	Schneider et al., 2018
Global Soil Wetness Project Phase 3	GSWP3	global	Half degree resolution (0.5°), daily, monthly	<a href="#">Reanalysis (20CR) corrected with gridded observation (GPCC)</a>	H. Kim et al. n.d.;
WATCH Forcing Data (WFD) + WATCH Forcing Data methodology applied to ERA-Interim data (WFDEI)	WATCH_W FDEI	global	Half degree resolution (0.5°), daily, monthly	<a href="#">Hydrological model applied to ERA Interim data</a>	Weedon et al., 2011, 2014

Table 1: Overview of the 10 precipitation datasets used in our study. Columns show the name of the dataset, the official abbreviation, the [short abbreviation used in here](#), the spatial and temporal resolution, and the references.

**Deleted:** Land surface models, remote sensing, in-situ observations...

**Formatted:** Swedish

**Deleted:** <http://hydro.iis.u-tokyo.ac.jp/GSWP3/index.html>

**Formatted:** English (UK)

**Formatted:** English (UK)

**Deleted:** Land surface models, remote sensing, in-situ observations...


**Formatted:** Pattern: Clear (Light Orange)

**Deleted:** , the inputs the precipitation datasets are derived from, ...

Drought event	Fraction of overall variability in rMCWD anomaly based on		
	precipitation datasets	drought indicators	evapotranspiration datasets
2005	0.21	0.6	0.19
2010	0.21	0.58	0.21
2016	0.22	0.59	0.19

**Table 2: Fraction of overall variability in rMCWD anomaly based on precipitation datasets, drought indicators, and evapotranspiration datasets.**


**Figures**



**Figure 1: Relative MCWD anomalies (from October to September) as an indicator for drought stress in the Amazon basin during the record-breaking drought event in 2016. Displayed are only the datasets that include the year 2016 in their temporal range. The baseline period of the MCWD calculation is 2001 to 2016.**

**Formatted: Font: Bold**

Formatted: Normal



Re  
 $x < -2.5$  -

Deleted:





Figure 2: Total area of the Amazon basin affected by drought stress (%) according to relative MCWD anomaly for each of the precipitation datasets. Displayed are the three drought events (a) 2005, (b) 2010 and (c) 2016. The total area representing the Amazon basin in our study is 5.94 million km<sup>2</sup>. For absolute area affected, see Tab. S2 and S3.



Deleted:

Formatted: Font: Not Bold








Figure 3: Agreement of precipitation datasets on drought area as identified by relative MCWD anomalies. In columns, different levels of drought severity are displayed and rows show the different drought years 2005 (a-c), 2010 (d-f) and 2016 (g-i). The colors indicate the number of datasets that agree on a specific drought level in a given pixel. Drought severity levels are defined as moderate ( $rMCWD < -0.5$ ), severe ( $rMCWD < -2.0$ ) and extreme ( $rMCWD < -2.5$ ). Orange pixels indicate areas where only one dataset shows the respective drought stress (No agreement = "None"). White pixels represent areas where no dataset shows any drought signal. Note that in a-f, TRMM 6 and GSWP3 were excluded, as they were either very similar to its successor (TRMM 7) or due to a similar reanalysis procedure (WATCH\_WFDEI). In g-i, only six datasets were included, which cover the full time period until 2016.





**Figure 4: Spatial pattern of ET for the dry and wet season for the DOLCE and GLEAM datasets (a, b, e, f) and the differences between using the two ET datasets to calculate the rMCWD anomaly and the rMCWD based on the constant ET=100mm per month assumption for 2005 (c, g) and 2010 (d, h). Wet and dry season ET is calculated as mean from June to August and January and March, respectively. Negative (positive) differences of the rMCWD anomalies indicate an overestimation (underestimation) of drought stress when using ET=100mm per month compared to the respective evapotranspiration dataset.**



**Figure 5: Agreement of precipitation datasets on drought area as identified by different drought metrics. Comparison of the Amazon drought events in 2005, 2010 and 2016 (columns) vs three different drought indexes (rows): *rMCWD* (a-c), *rscPDSI* (d-f) and *rRAI* (g-i). Only the area affected by severe drought stress is displayed, which is defined equally for each of the drought indices. Orange pixels indicate areas where only one dataset shows the respective drought stress (“None”). White pixels represent areas where no dataset shows any drought signal.**

Formatted: Font: Not Bold





Figure 6: Monthly development of the Amazon drought events in 2005, 2010 and 2016 (columns) as described by the three different drought indices (rows):  $rMCWD$  (a-c),  $rscPDSI$  (d-f) and relative rainfall anomaly ( $rRAI$ , g-i). Colored lines indicate the indices of the 10% quantile of all gridcells of each of the different precipitation datasets. The indices are estimated as relative deviation from a 2001 to 2016 baseline period for each month.



<b>Page 1: [1] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Correspondence

<b>Page 1: [2] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Footer

<b>Page 1: [3] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Equation

<b>Page 1: [4] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Affiliation

<b>Page 1: [5] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

List Paragraph

<b>Page 1: [6] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

MS title

<b>Page 1: [7] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Copernicus\_Word\_template

<b>Page 1: [8] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Name

<b>Page 1: [9] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
-------------------------------------	-----------------------------	----------------------------

Kontakt

<b>Page 1: [10] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Header

<b>Page 1: [11] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Bullets

<b>Page 1: [12] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Betreff

<b>Page 1: [13] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Comment Text

<b>Page 1: [14] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Subtitle

<b>Page 1: [15] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Title

<b>Page 1: [16] Style Definition</b>	<b>Phillip Papastefanou</b>	<b>16/03/2022 19:48:00</b>
--------------------------------------	-----------------------------	----------------------------

Heading 6

**Page 1: [19] Style Definition Phillip Papastefanou 16/03/2022 19:48:00**

Heading 3

**Page 1: [20] Style Definition Phillip Papastefanou 16/03/2022 19:48:00**

Heading 2

**Page 1: [21] Style Definition Phillip Papastefanou 16/03/2022 19:48:00**

Heading 1

**Page 1: [22] Style Definition Phillip Papastefanou 16/03/2022 19:48:00**

Normal

**Page 1: [23] Formatted Phillip Papastefanou 16/03/2022 19:48:00**

Title, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

**Page 1: [24] Formatted Phillip Papastefanou 16/03/2022 19:48:00**

Section start: New page, Numbering: Restart each page

**Page 1: [25] Formatted Phillip Papastefanou 16/03/2022 19:48:00**

Font: 16 pt, Font colour: Custom Colour (RGB(60,64,67)), Highlight

**Page 1: [25] Formatted Phillip Papastefanou 16/03/2022 19:48:00**

Font: 16 pt, Font colour: Custom Colour (RGB(60,64,67)), Highlight

**Page 1: [26] Formatted Phillip Papastefanou 16/03/2022 19:48:00**

Font: 16 pt, Font colour: Auto

**Page 1: [27] Deleted Phillip Papastefanou 16/03/2022 19:48:00**

▼.....

**Page 1: [27] Deleted Phillip Papastefanou 16/03/2022 19:48:00**

▼.....

**Page 1: [27] Deleted Phillip Papastefanou 16/03/2022 19:48:00**

▼.....

**Page 1: [27] Deleted Phillip Papastefanou 16/03/2022 19:48:00**

▼.....

**Page 1: [27] Deleted Phillip Papastefanou 16/03/2022 19:48:00**

▼.....

▼

**Page 1: [29] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 1: [30] Formatted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

Superscript

**Page 1: [31] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 1: [31] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 1: [32] Formatted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

Superscript

**Page 1: [33] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 1: [33] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 3: [34] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 30: [35] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼

**Page 36: [36] Deleted**    **Phillip Papastefanou**    **16/03/2022 19:48:00**

▼