

Interactive comment on “Hypersaline tidal flats as important “Blue Carbon” systems: A case study from three ecosystems” by Dylan R. Brown et al.

Dylan R. Brown et al.

christian.sanders@scu.edu.au

Received and published: 9 February 2021

I consider the manuscript "Hypersaline tidal flats as important "Blue Carbon" systems: A case study from three ecosystems" to be relevant in the actual global environmental context. The study is addressing important issues and bringing new perspectives, that could be further considered in the approach and management of hypersaline tidal flats (HTFs). The HTFs potential capacity of long-term CO₂ storage and their inclusion in the blue carbon framework should be further addressed, as these ecosystems can have a potential beneficial effect in the reduction of greenhouse gases (GHG) from the atmosphere. Response: We appreciate these positive comments, below we respond to specific comments individually.

C1

At line 139-140, "were measured in a high-purity germanium (HPGe) planar or well gamma detectors", the use of the word "or" creates confusion whether the author refers to two different detectors or if is giving an alternative name for the same apparatus, this should be clarified for a better understanding. Response: We have reworded this sentence for clarity, Line 142; "Radionuclides from the uranium-238 (238U) decay series were measured in a high-purity germanium (HPGe) gamma detectors, a planar for the Gladstone and Guaratiba and a well detector for the Karumba samples."

At line 145, "detector efficiency determined from standard calibration" needs further explanation concerning the calibration of the detector, especially it is indicated to describe the used calibration standards (IAEA – name and type) or, if the calibration was made by Monte Carlo modelling, the software name that was used. On the other hand, the sediment dating models (CIC), that was used in this study, was proved to be, in many cases, an idealistic model, because it assumes constant sediment deposition (constant sedimentation). In this case, considering that the surface is occasionally flooded, the sedimentation rate could vary, especially in Guaratiba. In this point, the 210Pb distribution through the sediment column shows discrepancies from the theoretical exponential decrease that is expected (Figure 2). In Gladstone site 2, only CIC model can be applied, because the dating horizon is not reached, and the column does not have the full inventory, that is essential for the CRS model. Response: We have reworded this paragraph for clarity, Line 147; "...were calculated by multiplying the counts per minute by a correction factor that includes the gamma-ray intensity and detector efficiency determined from NIST Rocky Flat soils reference material." Also, we agree that the CRS model was not appropriate for this study, as such we opted for using the CIC model. Given the sedimentary excess Pb-210 profiles, we feel that the CIC was our best option.

At line 165, page 5, the assumption that the pressure in the chamber is 1 atm., in my opinion, in some cases, for example, if the chamber is exposed to sunlight, the pressure inside can increase, which could influence the CO₂ exhalation. Response:

C2

We now clarify this issue, as large changes in temperature are not expected over a short time interval of only 5-15 min. The text now read, Line 161: "Each short-term incubation lasted 5-15 min to achieve a linear change in CO₂ concentration within the chambers, and was associated with a maximum increased temperature ~ 2°C in relation to external conditions, indicating no bias due to warming and subsequent changes in the inner pressure and biological activity."

Overall, I consider that the present manuscript contains valuable scientific information, that needs to be available to the large audience. Response: Once again, we thank the reviewers for these reviews.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-426>, 2020.